

Intrinsic structural features of the human IRE1 α transmembrane domain sense membrane lipid saturation.

Hyunju Cho^{1,6,8,*}, Francesca Stanzione^{2,7,8}, Amrita Oak^{1,8}, Geun Hyang Kim³, Sindura Yerneni¹, Ling Qi^{3,4}, Amadeu K. Sum², Christina Chan^{1,5,9,*}

¹Department of Chemical Engineering and Materials Science, Michigan State University, East Lansing, MI 48824, USA

²Department of Chemical and Biological Engineering, Colorado School of Mines, Golden, CO 80401, USA

³Department of Molecular & Integrative Physiology, University of Michigan Medical School, Ann Arbor, MI 48105, USA

⁴ Division of Metabolism, Endocrinology & Diabetes, Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI 48105, USA.

⁵Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI 48824, USA

⁶Department of Chemistry, California Institute of Technology, Pasadena, CA 91125, USA

⁷Institute of Medical Science-University of Aberdeen, Aberdeen AB25 2ZD, UK.

⁸These authors contributed equally

⁹Lead Contact

*Correspondence: krischan@egr.msu.edu (C.C.), hjcho@caltech.edu (H.C.)

Summary

Activation of Inositol-Requiring Enzyme (IRE1 α) is an indispensable step in remedying the cellular stress associated with lipid perturbation in the endoplasmic reticulum (ER) membrane. IRE1 α is a single-spanning ER transmembrane protein possessing both kinase and endonuclease functions, and its activation can be fully achieved through the dimerization/oligomerization process. How IRE1 α senses membrane lipid saturation remains largely unresolved. Using both computational and experimental tools, we systematically investigated the dimerization process of the transmembrane domain (TMD) of IRE1 α and found that, with help of the serine 450 residue, the conserved tryptophan 457 residue buttresses the core dimerization interface of IRE1 α -TMD. The BiFC (Bimolecular Fluorescence Complementation) experiments revealed that mutation on these residues abolished the saturated fatty acid-induced dimerization in the ER membrane and subsequently inactivated IRE1 α activity *in vivo*. Therefore, our results suggest that the structural elements of IRE1 α -TMD serves as a key sensor that detects membrane aberrancy.

Keywords

IRE1 α /UPR/dimerization/membrane lipid saturation.

Introduction

The Endoplasmic Reticulum (ER) is a cellular compartment responsible for protein folding, lipid synthesis, and calcium storage. Physiological conditions such as elevated levels of free fatty acids and glucose, oxidative stress, and inflammatory cytokines are known to induce ER stress, leading to the accumulation of unfolded/misfolded proteins or toxic lipid species (Wang and Kaufman, 2012). Cellular adaptation to ER stress is achieved through activation of the unfolded protein response (UPR), an integrated signal transduction pathway mediated by three ER stress sensor proteins, IRE1 (Inositol-requiring enzyme 1), PERK (protein kinase R (PKR)-like ER kinase), and ATF6 (activating transcription factor 6) (Kaufman, 1999). UPR signaling coordinates the cellular responses by down-regulating protein translation, enhancing the expression of ER chaperone proteins that promote protein refolding, and activating proteases that are involved in the degradation of misfolded proteins. When these adaptive processes are insufficient to attenuate ER stress, the UPR triggers apoptosis (Hetz and Papa, 2018; Wang and Kaufman, 2014).

Among the three UPR sensor proteins, IRE1 α is the most conserved protein and is ubiquitously expressed in cells. IRE1 α is a type I transmembrane protein with an N-terminal luminal domain (LD) that acts as an ER stress sensor and a C-terminal KEN (kinase-extension nuclease) domain containing both Ser/Thr kinase and endoribonuclease activities (Korennykh and Walter, 2012). Under ER stress conditions, the LD of IRE1 α oligomerizes in the lumen, promoting self-association of the protein (Oikawa et al., 2009). The dimerization of the kinase domain (KD) facilitates trans-autophosphorylation, subsequently activating the RNase domain (RD) (Ali et al., 2011). The active form of IRE1 α splices mRNA of the X-Box binding protein-1 (XBP1), producing the spliced XBP1 transcription factor (XBP1^S) and thus resulting in the upregulation of genes involved in enhancing ER protein-folding capacity and degrading unfolded or misfolded proteins (Lee et al., 2003; Yoshida et al., 2001).

Dimerization of the LD or cytosolic domain is known to be important for IRE1 α activation, whereas dimerization of its transmembrane domain (TMD) in the activation process remains unclear. Hydrogen bonds and hydrophobic interactions stabilize the dimerization interface of the LD, and disruption of these interactions prevents the self-association of the protein, subsequently reducing both the autophosphorylation and the RNase activities of IRE1 α (Li et al., 2010; Zhou et al., 2006). In addition, mutations at the dimerization interface on the KD diminish both the kinase

and RNase activities (Ali et al., 2011). Interestingly, a previous study suggests that the IRE1 α molecule lacking the LD is able to sense the lipid perturbation in the membrane and contribute to IRE1 α activation through the TMD (Volmer et al., 2013). Using a reporter assay that detects its downstream factor, XBP1^S, their follow-up study concluded that generic sequence features of the IRE1 α -TMD are sufficient for sensing membrane aberrancy (Kono et al., 2017). However, due to the dynamic and complex nature of the UPR signaling network (Cho et al., 2013; Tsuru et al., 2016; Yoshida et al., 2001), direct observation of the IRE1 α -TMD conformation is required to establish which features of the IRE1 α -TMD are crucial for IRE1 α activation as well as lipid saturation sensing.

In this study, we investigated how human IRE1 α senses membrane lipid saturation. To directly monitor the conformation of IRE1 α -TMD, we developed an *in vivo* BiFC (Bimolecular Fluorescence Complementation) assay to confirm that lipid saturation induces the dimerization of IRE1 α -TMD in the ER membrane. Systematic approaches using the *in vivo* BiFC assay and MD simulations identified an essential residue, tryptophan 457, that supports the core dimerization interface of IRE1 α -TMD. Our data reveal a key regulatory mechanism in which unique structural features of the tryptophan residue modulate the saturated fatty acid-induced dimerization of IRE1 α -TMD, and thus promote an active conformation of IRE1 α .

Results

The isolated TMD of IRE1 α dimerizes in membranes.

To assess whether the TMD of IRE1 α dimerizes in the lipid bilayer, we used an established Förster resonance energy transfer (FRET) assay (Li et al., 2005) that measures the self-association of TMD peptides in 1-palmitoyl 2-oleoyl phosphatidylcholine (POPC) liposomes. When both donor (Fluorescein isothiocyanate (FITC)) and acceptor (Carboxytetramethylrhodamine (TAMRA)) labeled IRE1 α -TMD peptides were present in the POPC liposomes, we observed a decrease in the donor fluorescence and an increased in the acceptor fluorescence, indicating that FRET has occurred (Figure 1A). The observed FRET values at various acceptor to lipid ratios (Figure 1B) were comparable to the previously measured TMD dimerization of fibroblast growth factor receptor 3 (FGFR3) in the POPC liposomes using the same dye pair (Li et al., 2005). This indicates that the observed FRET in the IRE1 α -TMD containing liposomes is not entirely due to

random acceptor/donor colocalization and that IRE1 α -TMD has a propensity for self-association in liposomes.

Two additional experiments further support our conclusion that the IRE1 α -TMD dimerizes/oligomerizes in the lipid bilayer. First, SDS-PAGE analysis, which is commonly used to detect the oligomerization status of TMDs, showed that the IRE1 α -TMD peptide has several oligomeric states, as evident by the higher band (Figure 1C). The monomeric and dimeric states did not separate clearly, possibly reflecting the dynamic equilibrium between the two states during the electrophoresis, which is frequently observed in short TMD peptides (Ng and Deber, 2010). Second, the results of the TOXCAT assay, in which association between the TMDs in the inner membrane of *E. coli* drives expression of the reporter gene chloramphenicol acetyltransferase (CAT) (Russ and Engelman, 1999), suggested dimerization of the IRE1 α -TMD peptides in the cellular membrane. The CAT activity was normalized to a positive control (the TMD of glycophorin A (GpA)), which is known to dimerize strongly, while GpA-G83I served as a dimerization-impaired mutant (a negative control) (Lemmon et al., 1992). The CAT activity of IRE1 α -TMD was significantly higher than the negative control (Figure 1D), indicating that the IRE1 α -TMD dimerizes in a biological membrane. Furthermore, MalE complementation assay showed that the lack of a TMD did not allow *E. coli* to grow on maltose, while all other constructs exhibited robust growth. This indicates that the MBP domain of these fusion proteins was localized on the periplasm of the NT326 cells and the IRE1 α -TMD was correctly inserted in the cell membrane (Figure 1E). Therefore, we showed that the isolated TMD of the IRE1 α protein dimerizes/oligomerizes in lipid bilayers.

Both Trp457 and Ser450 support palmitate-activated dimerization of the IRE1 α - Δ LD *in vivo*.

Membrane lipid saturation induced by a saturated fatty acid, palmitate, is known to upregulate the IRE1 α activity (Kitai et al., 2013; Volmer et al., 2013). In contrast to the conventional UPR activation initiated by accumulation of unfolded protein in the ER lumen, membrane lipid saturation did not trigger the clustering of IRE1 α proteins via oligomerization of the LD, and thus palmitate is thought to use a different mechanism of activating the IRE1 α protein (Kitai et al., 2013). In addition, the IRE1 α protein lacking the LD (30-407 aa, denoted as IRE1 α - Δ LD) can be activated by palmitate (Volmer et al., 2013), suggesting that palmitate could directly regulate the conformation of IRE1 α -TMD. To directly monitor palmitate-induced dimerization *in*

in vivo, we employed a BiFC assay that has been successfully used to quantify dimerization of both the seven transmembrane domain (7TM) receptor and the receptor tyrosine kinases (RTKs) family proteins *in vivo* (Kosel et al., 2010; Tao and Maruyama, 2008). Therefore, to dissect the role of palmitate on the dimerization of IRE1 α -TMD, we used the IRE1 α - Δ LD construct devoid of the upstream signal (Volmer et al., 2013). The N-terminal Venus fragment (Myc-tagged VN) and the C-terminal Venus fragment (HA-tagged VC) were inserted downstream of the signal peptide (1-18 aa) on the IRE1 α - Δ LD (Figure 2A) (Shyu et al., 2006). Once both VN-IRE1 α - Δ LD and VC-IRE1 α - Δ LD are in-close proximity in the ER membrane, the Venus fragments will be able to reconstitute a native and fluorescent protein (Figure 2B).

We monitored the early response of palmitate-induced dimerization of IRE1 α - Δ LD in the IRE1 α ^{-/-} MEF cells expressing both VN-IRE1 α - Δ LD and VC-IRE1 α - Δ LD. IRE1 α ^{-/-} MEF cells have a negligible amount of endogenous IRE1 α proteins (Volmer et al., 2013) and thus allow the monitoring of the dimerization efficiency of exogenously expressing IRE1 variants *in vivo*. We first confirmed that the expression levels of IRE1 α - Δ LD in BSA- and palmitate-treated IRE1 α ^{-/-} MEF cells were comparable to the endogenous IRE1 α levels in its parental MEF-WT cells (Figure S1A and S1B). Consistent with this, the exogenous expression of IRE1 α - Δ LD did not alter the palmitate-induced activation of endogenous eIF2 α , a subsequent downstream factor of the other UPR sensor PERK (Figure S1C and S1D), demonstrating that the PERK pathway was not perturbed by exogenously expressing the IRE1 α - Δ LD proteins. In contrast, upon palmitate treatment at 2 h, the Venus fluorescence intensity increased significantly as compared to the BSA-treated control cells (Figure 2C and 2D), indicating the reconstitution of the Venus fluorescent protein and thus dimerization of the VN-IRE1 α - Δ LD-WT and VC-IRE1 α - Δ LD-WT proteins.

Several lines of evidence suggest that the observed dimerization of IRE1 α - Δ LD in the *in vivo* BiFC assay was specifically induced by palmitate. First, the IRE1 α - Δ LD proteins were expressed at similar levels in both the BSA- and palmitate-treated cells (Figure 2E and Figure S1A), suggesting that the enhanced dimer formation of IRE1 α - Δ LD by palmitate is not due to a difference in the protein expression levels between the two conditions. Second, the palmitate-induced dimerization of IRE1 α - Δ LD-WT was further corroborated by the enhanced XBP1 splicing efficiency (Figure 2F and 2G). Third, a monounsaturated fatty acid, oleate (OA), failed to induce the dimerization of IRE1 α - Δ LD (Figure 2C and 2D), suggesting that the induced dimerization of IRE1 α - Δ LD in the palmitate-treated cells is specifically due to lipid saturation. Furthermore,

consistent with the previous study (Volmer et al., 2013), tunicamycin, an ER stress inducer that perturbs protein folding in the ER lumen, was also unable to induce the dimerization of IRE1 α - Δ LD (Figure 2C and 2D). Therefore, the enhanced dimerization of IRE1 α - Δ LD *in vivo* is specific to the increased lipid saturation due to palmitate.

To systematically identify the key residues responsible for the dimerization of IRE1 α - Δ LD, we performed alanine-scanning mutagenesis on the IRE1 α -TMD using the BiFC assay. All residues, except for two native alanines, were mutated to alanine in the VN-IRE1 α - Δ LD and VC-IRE1 α - Δ LD constructs (Figure 2A). Among the tested mutations, both S450A and W457A were not able to reconstitute the fluorescence in the palmitate-treated cells, indicating an inability to dimerize in the ER membrane (Figure 2C and 2H). The loss of palmitate-induced dimerization of IRE1 α - Δ LD with the W457A and S450A mutants resulted in a catalytically inactive state associated with their failure to generate XBP1^S (Figure 2F and 2G). The dimerization and RNase defects of both W457A and S450A were not due to lower levels of protein expression than the WT or mislocalization of the mutant proteins under the palmitate-treated condition (Figure 2E and S2A and S2B, respectively). The obtained Pearson's correlation coefficients for W457A and S450A were not significantly different from the WT, suggesting that these mutants primarily co-localize with the ER membrane marker protein, calnexin, in the BSA and palmitate-treated Hep3B cells (Figure S2A and S2B). Therefore, our *in vivo* data demonstrate an essential role of both Trp457 and Ser450 in modulating the palmitate-induced dimerization of IRE1 α - Δ LD.

IRE1 α -TMD serves as a sensor for lipid membrane saturation in mammals.

Results of a previous study with the yeast Ire1 showed that the juxta-membrane amphipathic helix, but not its TMD, plays a crucial role in sensing aberrant physical membrane properties (Halbleib et al., 2017). Since the amphipathic helix is conserved on the human IRE1 α (Figure 3A) (Halbleib et al., 2017; Kono et al., 2017), we further assessed whether this is also important in regulating the dimerization of IRE1 α - Δ LD through activation by membrane lipid saturation. To this end, we mutated two conserved hydrophobic residues (V437 and M440) on the amphipathic helix to arginine and performed the BiFC assay. In contrast to the yeast Ire1 (Halbleib et al., 2017), the two amphipathic helix mutations did not prevent the dimerization of the IRE1 α - Δ LD in the ER membrane (Figure 3B and 3C). V437R and M440R helix mutant proteins had similar expression levels as the WT protein (Figure 3D), and they also co-localized with calnexin

in both the BSA and palmitate-treated cells (Figure S2A and S2B). Furthermore, the amphipathic helix mutant proteins exhibited WT-like XBP1 splicing activity (Figure 3E and 3F). Therefore, together with the previous study (Kono et al., 2017), our *in vivo* results strongly suggest that the human IRE1 α uses a different sensing mechanism for membrane lipid saturation than the yeast Ire1.

To exclude the possibility that the N-terminal LD affects the dimerization conformation of the IRE1 α -TMD, we used full-length IRE1 α BiFC constructs, VN-IRE1 α -FL and VC-IRE1 α -FL (Figure 4A) and further monitored their dimerization in IRE1 α ^{-/-} MEF cells. Palmitate induced the dimerization of full-length IRE1 α , as evident by an increase in the reconstituted green fluorescence (Figure 4B). Consistent with the observation of the IRE1 α - Δ LD (Figure 2C and 2H), both W457A and S450A mutations on the IRE1 α -FL dramatically reduced the dimerization in the palmitate-treated cells (Figure 4B and 4C). Since the active IRE1 α dimers further facilitate the formation of high-order-oligomers that re-localize into discrete foci *in vivo* (Li et al., 2010), we further calculated the IRE1 α foci-positive cells. In agreement with their dimerization capacity (Figure 4B and 4C), both the W457A and S450A mutants significantly reduced the IRE1 α foci-positive cells (Figure 4D), while the total protein levels of the IRE1 α mutants were unaltered (Figure 4E). In the BiFC experiments, the fluorescence only results upon the formation of the IRE1 α dimer. Thus, the decreased number of foci-positive cells in both mutants is due mainly to their reduced dimerization ability. In addition, since the expression levels of IRE1 α -FL variants in the IRE1 α ^{-/-} MEF cells are similar to the endogenous IRE1 α level in the MEF-WT cells (Figure S1A and S1B and 4E), the dimerization/oligomerization defect in the W457A and S450A mutants is unlikely due to non-physiologically high concentrations of the IRE1 α -FL variants. These full-length IRE1 α proteins co-localized with the ER membrane protein, calnexin, with similar Pearson's correlation coefficients (Figure S3A and S3B). Consistent with this, both the IRE1 α -FL-W457A and IRE1 α -FL-S450A mutants in the palmitate-treated cells showed significantly reduced XBP1^S fractions (Figure 4F and 4G). Therefore, these results suggest that both Trp457 and Ser450 play a crucial role in not only the initial dimerization process but also the IRE1 α oligomer formation that could activate XBP1 splicing.

To further test whether the reduction of XBP1 splicing efficiencies on the W457A and S450A mutants resulted from lipid perturbation, but not accumulation of unfolded proteins upon palmitate treatment, we used an established Y161A mutation that is defective in sensing unfolded

proteins (Kono et al., 2017). Similar to the results with the IRE1 α - Δ LD (Figure 2F and 2G), the XBP1 splicing activities of both IRE1 α -FL-Y161A-W457A and IRE1 α -FL-Y161A-S450A were significantly lower than IRE1 α -FL-Y161A (Figure 4H and 4I). Collectively, our *in vivo* results suggest that the intrinsic structural elements of the IRE1 α -TMD could be a key factor in sensing membrane lipid saturation in mammals.

MD simulations identified a key role of Trp457 in a potential dimerization process of the IRE1 α -TMD.

To further investigate the potential roles of both Trp457 and Ser450 on the dimerization process of IRE1 α -TMD, we performed atomistic MD simulations, where two IRE1 α -TMD peptides were inserted in an explicit POPC lipid bilayer. Due to the lack of structural information on the IRE1 α -TMD, three different initial conformations of IRE1 α -TMD were arbitrarily assigned (Model A, B, and C) (Figure S4A). We estimated the dimerization tendency of the TMD peptides in the POPC bilayer based on both the distance between the center of mass (CM) of the peptides and the inter-peptide interactions. In model A, the CM distance initially decreased from 11.8 Å (Figure S4B, X mark) to 9.9 ± 0.4 Å and then returned to the initial value (Figure S4B). In model B, the CM distance decreased from 11.8 Å to 10.3 ± 0.3 Å while, in model C, the CM distance increased to 14.0 Å (Figure S4B). Consistent with this, the total number of atomic contacts within 6 Å between the two peptides in model C was significantly lower than both models A and B (Figure S4C).

To analyze the nature of the inter-peptide interactions, we calculated the average conformation of the three models over the last 50 ns of the simulation. From the analysis, we found that both models A and B have common conformational features in the IRE1 α -TMD peptides. First, the TMD peptides had a slightly tilted conformation in the POPC bilayer with a tilt angle of $8.9 \pm 1.5^\circ$ and $5.0 \pm 2.0^\circ$ in models A and B, respectively (Figure S4D). In contrast, to escape from the hydrophobic acyl chains of the POPC to the more polar head groups, the hydrophilic TMD residues in model C rearranged drastically, thereby leading to a deformed and tilted conformation in the membrane (Figure S4D) (Killian, 1998). Second, Trp457 was observed in the dimerization interface of the TMD peptides. In the simulation with model A, Trp457 in TMD1 and Trp457 and Phe460 in TMD2 were clustered at the interface of the TMD peptides (Figure S5A). In this interface, Trp457 in TMD1 and Phe460 in TMD2 formed the T-shaped aromatic interaction which

is slightly greater (~ 6 Å) than an ideal T-shaped stacking value (~ 5 Å) (Janowski and Pulay, 2007) (Figure S5B). In the initial structure of model B, the two Trp457 residues were located on the opposite sides of the TMD peptides (CM distance = 14 Å, green in Figure 5A). However, they rapidly changed orientations to relocate to the center of the TMD peptide interface, forming a stable aromatic interaction (CM distance = 4-5 Å, orange in Figure 5A) (also see movie S1). Similar to model A (Figure S5A), the position of Trp457 at the interface was transiently stabilized by the aromatic interactions with the Phe460 residue during the simulation (Figure 5B). Given that F460A did not exhibit any significant defect in the dimerization of IRE1 α - Δ LD *in vivo* (Figure 2H, the observed Phe460-Trp457 interactions in models A and B could be intermediate states that are not essential for the IRE1 α -TMD dimerization. After the first 150 ns of simulation of model B, the distance between the benzene and indole centroids decreased (Figure 5C, left panel), creating an aromatic pocket with an ideal aromatic sandwich stacking configuration of the tryptophan residues (Figure 5C, right panel). Furthermore, the ring orientations of Trp457 allowed the formation of H-bonds between the NH of the indole groups and the carbonyl oxygen of Trp457, Leu454, and Leu453 (Figure S6A and S6B). In addition to Trp457, Ser450, located near the N-terminus of the TMD, transiently formed H-bonds with Ser450 and Thr446 on the other peptide (Figure S6C and S6D). Interestingly, inspection of the average conformation of model C revealed that the Trp457 was not located at the interface, suggesting that the lack of the Trp457 at the peptide interface could destabilize the dimer of IRE1 α -TMD, inducing a dramatically tilted conformation in the POPC bilayer (Figure S4D). Together with the *in vivo* experimental results (Figure 2 and 4), the MD simulation results from both models A and B suggest that Trp457 could play a key role in the IRE1 α -TMD dimerization process.

To provide broad insight on the alanine mutagenesis results (Figure 2H), we further calculated the atomic contact number (cutoff = 6 Å) for each residue in model B. The analysis revealed that Trp457 was the most frequently localized residue to the dimerization interface during the last 50 ns of simulation time (normalized to 1 in Figure 5D) and residues including Thr446, Ser450, Leu453, and Leu454 had about 30-40% of the contact numbers of Trp457 (Figure 5D). This is consistent with the *in vivo* data showing that W457A, S450A, and T446A have defects in the dimerization of IRE1 α - Δ LD (Figure 5D, red). By contrast, the alanine mutation of both Leu453 and Leu454 did not alter the dimerization efficiency of IRE1 α - Δ LD (Figure 5D, blue). In addition, both the I447A and I461A mutants exhibited less than 50% of the WT dimerization efficiency

(Figure 5D, green). It is notable that all the defective residues identified from the BiFC assay (Figure 2H) are localized in the dimerization interface obtained from model B (Figure 5E). Therefore, we speculate that model B could represent the initial dimerization stages of the IRE1 α -TMD.

Trp457 locates near the center of the POPC bilayer.

Tryptophan is found abundantly in membrane proteins. Due to its amphipathic nature, tryptophan is known to stabilize the membrane proteins by anchoring the interactions in the membrane-water interface (Yau et al., 1998). However, tryptophan often locates toward the center of the TMD helix to support the aromatic interaction, which mediates the TMD dimerization (Haeger et al., 2010; Ridder et al., 2005; Sal-Man et al., 2007). Using the MD simulation results, we further analyzed both the density profile of the Trp457 and POPC head group as a function of the distance from the bilayer center along the direction normal to the bilayer plane (Figure 5F). The preferred location of Trp457 is close to the center of the lipid bilayer, i.e., the hydrophobic core region of the POPC bilayer (Figure 5F and 5G). The calculated root mean square distance of Trp457 from the center of the bilayer ($\langle Z_{cf}^2 \rangle^{1/2}$) is 0.76 ± 0.12 Å, corroborating a density profile that locates the tryptophan residues in proximity of the center of the POPC lipid bilayer.

To experimentally determine the location of Trp457 in the POPC lipid bilayer, we used the parallax method that estimates the membrane penetration depth of the tryptophan residue (Chattopadhyay and London, 1987). The parallax method allows the determination of the apparent location of tryptophan by comparing the quenching of the tryptophan fluorescence by two different spin-labeled POPC lipids, 5-Doxyl PC (shallower quencher), and 16-Doxyl PC (deeper quencher). Compared to the fluorescence intensity in the absence of the quenchers (Figure 5H, black), Trp457 was quenched dramatically by the deeply located nitroxide lipid quencher ($F/F_0 = 0.099$) (Figure 5H, red), but much less efficiently by the shallow quencher ($F/F_0 = 0.71$) (Figure 5H, blue). We further calculated the distance of Trp457 from the center of the bilayer (Z_{cf}) as described in the Methods. The average depth of penetration of Trp457 from the center of the bilayer was estimated at 0.88 ± 0.35 Å, consistent with the computationally calculated value of 0.76 ± 0.12 Å. Therefore, both the computational and experimental results indicate that the tryptophan residue locates near the center of the bilayer.

Intrinsic structural features of Trp457 are important for the dimerization process of IRE1 α -TMD.

Of the aromatic residues, the side chain of tryptophan is unique with its amphipathic nature, enabling it to participate in both aromatic interactions and hydrogen bonding (NH- π and CH- π) (Samanta et al., 1999). This multi-functional side chain makes the tryptophan residue a key contributor, both to the process of protein folding and stabilization of the folded protein structures. To address whether the tryptophan residue is a unique feature in the 457 position for the IRE1 α -TMD dimerization, Trp457 was mutated to either alanine or phenylalanine. W457F preserves the aromatic ring but removes both the H-bond donor and dipole moment properties of the tryptophan whereas W457A removes both the aromatic interactions and hydrogen bonds. Both W457A and W457F mutants significantly reduced the dimerization ability of IRE1 α -TMD in the cellular membrane (Figure 6A). Therefore, the experimental results suggest that the aromatic characteristic alone is insufficient to promote self-association, and the H-bond or the dipole moment on the tryptophan residue could be important in regulating the TMD dimerization of the IRE1 α protein.

To elucidate the potential role of tryptophan on the TMD dimerization process, we performed further MD simulations of the W457 mutants (W457A and W457F) (Figure 6B and 6C) starting from the initial conformation of model B. The CM distances of W457A and W457F were not significantly different from the WT (Figure 6D). In the W457F model, in contrast to the WT (Figure 5A), the benzene ring only formed a T-shaped aromatic interaction during the first half of the simulation (CM distance = 6 Å, blue in Figure 6E) and then they moved apart, breaking the aromatic interaction in the last 100 ns (CM distance = 7.3 Å, red and orange in Figure 6E). Therefore, both the computational and experimental results suggest that the predominant effect of the W $\rightarrow$ F substitution results from the removal of the electrostatic interactions between the indole dipole and the polar groups, supporting that the aromatic characteristic alone is insufficient to drive the TMD dimerization.

To further understand the potential mechanisms by which Ser450 regulates the dimerization of IRE1 α -TMD, we performed MD simulations with the S450A peptides (Figure 6F). Serine substitution to alanine only changes the side chain of CH₂OH to CH₃, thus eliminating the capability of H-Bonds. The CM distance between the S450A peptides increased significantly as compared to WT (Figure 6G). As observed in the WT (Figure 5A), the CM distance between the

indole rings on Trp457 also rapidly decreased (14 Å to 7.5 Å) in S450A. Due to the loss of H-bonds, this distance increased back to 9.5 Å and remained constant over the simulation time except for a short period around 100 ns (Figure 6H). Therefore, the MD simulation results suggest that the primary effect of S450A could be a loss of the Trp457-Trp457 interactions. If this hypothesis is correct, we would not observe any synergetic effect in the S450A-W457A double mutant cells. Consistent with this notion, the dimerization efficiency of IRE1 α in the S450A-W457A cells is approximately 20% of the WT and thus is not significantly different from the single mutant (S450A or W457A) cells (Figure S7A and S7B vs. Figure 4B and 4C). Taken together, the H-bonds formed by Ser450 could contribute to the stability of the Trp457-Trp457 interaction during the IRE1 α -TMD dimerization process.

Discussion

Here we propose a mechanistic model of human IRE1 α activation by membrane lipid saturation. Under normal conditions, four functional domains (LD, TMD, KD, and RD) of the IRE1 α present in an inactive monomer conformation (Figure 7). In particular, the Trp457 residue could remain on the opposite side of the dimerization interface (Figure 7, back-to-back). Given that palmitate alters the physical properties of the membrane, e.g., fluidity and lipid packing (Covino et al., 2016; Leekumjorn et al., 2008), the tryptophan residues rapidly rotate and support the core dimerization interface of the IRE1 α -TMD (Figure 7, face-to-face). In addition, the Ser450 residue assists the dimerization stability by forming H-bonds with itself and with Thr446. The dimerization of the IRE1 α -TMD further brings the cytosolic domains together which could induce the face-to-face conformation of KD via a currently unknown mechanism, leading subsequently to an active, phosphorylated conformation (Figure 7) (Ali et al., 2011; Li et al., 2010). The activated IRE1 α enables the cleavage of the Xbp1 mRNA for downstream transcriptional regulation of UPR genes. Therefore, our results showed that the TMD of IRE1 α serves as an active signaling modulator of membrane lipid saturation.

In membrane proteins, it is well known that tryptophan preferentially localizes in the membrane interface, due to its overall amphipathic nature and the aromaticity of the indole moiety (Yau et al., 1998). Our bioinformatics analysis revealed that about 30% of the single-spanning transmembrane proteins (a total of 2365 human proteins) contain tryptophan residues. One-fifth

of these proteins have tryptophan near the center of the lipid bilayer (cutoff = $\pm 30\%$ from the center amino acid residue in the TM domain) (Table S1). Additionally, several previous studies suggest that tryptophan supports self-association of the helical TMD segments (Haeger et al., 2010; Ridder et al., 2005; Sal-Man et al., 2007). Our study further demonstrates that the conserved tryptophan (Figure S8A) is an essential amino acid residue for the dimerization process. Additionally, the chloride intracellular channel protein 1 (CLIC1), one of the positive hits in our tryptophan location analysis (Table S1), has been shown previously to stabilize the TM dimerization in the membrane by the cation (Lys37)- π interaction (Trp35) (Peter et al., 2014). The WxK motif is known to increase the strength of oligomerization of the TMD helices (Johnson et al., 2007) and it is conserved on the chloride intracellular channel protein family except for CLIC3 (Table S1). Remarkably, the tryptophan location analysis showed that another ER sensor protein, PERK, has two conserved tryptophan residues (Trp521 and Trp522) near the center of the TMD sequence (Table S1 and Figure S8B), suggesting dimerization of the PERK TM domain might be mediated by the cation (Lys523)- π interaction (Trp521). Nevertheless, whether the tryptophan residues play a crucial role in activating the PERK protein remains to be tested.

An intriguing feature we observed in the MD simulations is a rotameric switch of the tryptophan residue of IRE1 α -TMD (Figure 5A and Movie S1). In model B, two Trp457 residues initially pointed in opposite directions, and then they rapidly flipped towards the center of the TM dimerization interface to maximize their interactions (Figure 5A and Movie S1). The rotameric switch could optimize the IRE1 α -TMD interface by providing the driving force for other H-bonds to form (Figure S6A and S6B). The tryptophan rotameric switch model is most commonly observed in G-protein-coupled receptors (GPCRs). A highly conserved tryptophan residue (Trp^{6.48}) in TMD6 was rotated toward another conserved phenylalanine residue (Phe^{5.47}) in TMD5, and these two aromatic residues formed an edge-to-face aromatic interaction (Holst et al., 2010; Li et al., 2013; Lyman et al., 2009). This rotameric switch is known to be involved in the activation process of GPCRs (Holst et al., 2010; Li et al., 2013; Schwartz et al., 2006). Interestingly, a recent study showed that Trp1042 on the TM domain of the Mga2 protein senses the lipid saturation of the yeast ER membrane and subsequently triggers the TM helix rotation to induce an active Mga2 conformation for transcriptional upregulation of fatty acid desaturation (Covino et al., 2016). Consistent with this, our MD simulation and *in vivo* results suggest that the Trp457 rotameric switch of the IRE1 α -TMD induced by palmitate triggers the active conformation of the cytosolic

domain of the IRE1 α protein to induce XBP1 mRNA splicing. Taken together, the rotameric switch could be important for activation of membrane proteins.

In addition to the tryptophan residue, the MD simulations predicted that the residues at the interface of the IRE1 α -TM domain consisting of Ser450, Leu453, and Leu454 are involved in the H-bonds and hydrophobic interactions (Figure S6C and S6D). Our BiFC experiments showed that S450A dramatically reduces the dimerization of the IRE1 α (Figure 2 and 4). Interestingly, these residues are located as a leucine-zipper-like motif (SxxLxxx motif) (Figure S8A) which was also observed in the parallel TM dimerization of the mouse erythropoietin receptor (EpoR) protein (Ruan et al., 2004). In addition, both the tryptophan and SxxLxxx motif on IRE1 α are highly conserved in mammals, but not in yeast Ire1 (Figure S8A) (Kono et al., 2017). Due to the different ER membrane lipid composition between yeast and mammals (van Meer et al., 2008), the TMD sequences on the IRE1 α protein of mammals could have evolved to be particularly sensitive to lipids in the ER membrane while the yeast Ire1 relies on the juxta-membrane amphipathic helix sequences (Halbleib et al., 2017).

Why are the intrinsic structural features of the IRE1 α -TMD required for sensing the membrane lipid perturbation? Results of a previous study showed that replacing the IRE1 α -TMD sequence by its scrambled TMD sequence or calnexin TMD sequence did not completely abolish the XBP1 splicing activity of the IRE1 α protein (Volmer et al., 2013). However, the XBP1 splicing efficiencies for those two constructs and the WT under the same treatment conditions were significantly different in the IRE1 α -FL and IRE1 α - Δ LD expressing cells (Volmer et al., 2013), suggesting that the IRE1 α TMD sequence cannot be fully replaced. Given that the UPR signaling pathways are dynamically modulated by palmitate (Cho et al., 2013), the intrinsic structural feature of the IRE1 α -TMD could be crucial for fine-tuning the dynamic UPR signaling induced by membrane lipid saturation. The bulky tryptophan residue 457 could be accommodated in the loosely packed lipid bilayer whereas in a more densely packed bilayer, the formation of Trp457-Trp457 aromatic stacking interaction would be energetically more favorable (Covino et al., 2016). Based on our MD simulation results, this rotation is expected to be very fast (~ 10 ns) (Figure 5A) and thus would allow for rapid response to small lipid perturbations in the ER membrane. In contrast, due to lipid packing in the presence of palmitate, the TMD lacking the tryptophan residue could passively diffuse and eventually dimerize under prolonged palmitate exposure (Kono et al.,

2017). Although this hypothesis remains to be tested, we propose that the tryptophan residue in the IRE1 α -TMD enables a dynamic switch by rapid rotation upon membrane lipid saturation.

From the MD simulation results, we propose that the dimerization process of IRE1 α -TMD uses a “dual-lock” strategy. In this scenario, the Trp457 is a core lock that buttresses the dimerization interface and the Ser450 is a secondary lock that further secures the core interactions. The S450A mutant showed a partial rotation of Trp457 residues (Figure 6F and 6H), representing an intermediate state during the IRE1 α -TMD dimerization process. This intermediate state would sense the perturbation of the lipid composition in the ER membrane, however, it is insufficient to drive a stable dimerization of the IRE1 α -TMD and to transmit signals to the cytosolic domain of IRE1 α *in vivo* (Figure 2 and 4). Further mechanistic study is needed to confirm that S450A is an isolated mutant that decouples lipid saturation sensing from the dimerization of IRE1 α -TMD, nonetheless, we suggest that the full rotation of the Trp457 residues is required for an active conformation of IRE1 α .

Whereas the conventional UPR initiates by BiP release from IRE1 α -LD upon accumulation of unfolded proteins in the ER lumen and subsequent dimerization of IRE1 α -LD, our data with IRE1 α - Δ LD and IRE1 α -Y161A (Figure 2 and 4) demonstrate that palmitate directly activates IRE1 α through the dimerization of the TMD. This further suggests that, depending on the nature of the physiological stimuli, IRE1 α could use different sensing mechanisms. Given that IRE1 α dynamically assembles with numerous co-factors and regulators to activate its cytosolic domain that constitutes the UPRosome (Hetz and Glimcher, 2009), whether this non-canonical activation of IRE1 α by lipid perturbation requires specific factors in the UPRosome is currently unknown. Interestingly, a recent study showed that Bax/Bak, one of the co-factors that binds to the cytosolic domain of IRE1 α (Hetz et al., 2006), is required for palmitate-induced activation of ER stress (Zhang et al., 2018). Therefore, the mapping of the IRE1 α UPRosome that is induced by lipid perturbation remains to be explored. This, together with our results, would provide further insight into how IRE1 α could customize its sensing and regulatory domains for a specific stimulus.

Acknowledgements

We thank Dr. Heedeok Hong for valuable discussions on the parallax method. This study was supported in part by the National Science Foundation (CBET 1510895, CBET 1547518, and CBET 1802992) and the National Institutes of Health (R01GM079688, R01GM089866, and R21CA176854).

Author Contributions

Conceptualization, H.C., A.K.S, and C.C.; Experiments, H.C., A.O., G.H.K., and S.Y.; MD simulations, F.S.; Data Analysis, H.C., F.S., A.O., G.H.K., L.Q., and C.C.; Manuscript writing, H.C., F.S., and A.O.; Manuscript edition, H.C., A.K.S, C.C. with all authors' input; Funding Acquisition, C.C. and A.K.S.

Declaration of Interests

The authors declare no competing interests.

References

- Ali, M.M.U., Bagratuni, T., Davenport, E.L., Nowak, P.R., Silva-Santisteban, M.C., Hardcastle, A., McAndrews, C., Rowlands, M.G., Morgan, G.J., Aherne, W., *et al.* (2011). Structure of the Ire1 autophosphorylation complex and implications for the unfolded protein response. *Embo Journal* 30, 894-905.
- Berendsen, H., Postma, JPM, van Gunsteren, WF, Hermans, JJ (1981). Interaction models for water in relation to protein hydration (D. Reidel Publishing Company).
- Buchan, D.W., Ward, S.M., Lobley, A.E., Nugent, T.C., Bryson, K., and Jones, D.T. (2010). Protein annotation and modelling servers at University College London. *Nucleic Acids Res* 38, W563-568.
- Chattopadhyay, A., and London, E. (1987). Parallax method for direct measurement of membrane penetration depth utilizing fluorescence quenching by spin-labeled phospholipids. *Biochemistry* 26, 39-45.
- Cho, H., Lamarca, R., and Chan, C. (2012). Oligomerization of the transmembrane domain of IRE1 α in SDS micelles. *Biochem Biophys Res Commun* 427, 764-767.
- Cho, H., Wu, M., Zhang, L., Thompson, R., Nath, A., and Chan, C. (2013). Signaling dynamics of palmitate-induced ER stress responses mediated by ATF4 in HepG2 cells. *BMC Syst Biol* 7, 9.
- Covino, R., Ballweg, S., Stordeur, C., Michaelis, J.B., Puth, K., Wernig, F., Bahrami, A., Ernst, A.M., Hummer, G., and Ernst, R. (2016). A Eukaryotic Sensor for Membrane Lipid Saturation. *Mol Cell* 63, 49-59.
- De Lano, W. (2002). The PyMOL Molecular Graphics System (San Carlos, CA, USA.: DeLano Scientific).
- Gautier, R., Douguet, D., Antonny, B., and Drin, G. (2008). HELIQUEST: a web server to screen sequences with specific alpha-helical properties. *Bioinformatics* 24, 2101-2102.
- Haeger, S., Kuzmin, D., Detro-Dassen, S., Lang, N., Kilb, M., Tsetlin, V., Betz, H., Laube, B., and Schmalzing, G. (2010). An intramembrane aromatic network determines pentameric assembly of Cys-loop receptors. *Nat Struct Mol Biol* 17, 90-98.
- Halbleib, K., Pesek, K., Covino, R., Hofbauer, H.F., Wunnicke, D., Hänel, I., Hummer, G., and Ernst, R. (2017). Activation of the Unfolded Protein Response by Lipid Bilayer Stress. *Mol Cell* 67, 673-684.e678.
- Hess B, K.C., Van der Spoel D & Lindhal E (2008). GROMACS 4: Algorithms for Highly Efficient, Load-Balanced, and Scalable Molecular Simulation. *J Chem Theory Comput* 4, 435-447.
- Hetz, C., Bernasconi, P., Fisher, J., Lee, A.H., Bassik, M.C., Antonsson, B., Brandt, G.S., Iwakoshi, N.N., Schinzel, A., Glimcher, L.H., *et al.* (2006). Proapoptotic BAX and BAK modulate the unfolded protein response by a direct interaction with IRE1 α . *Science* 312, 572-576.
- Hetz, C., and Glimcher, L.H. (2009). Fine-tuning of the unfolded protein response: Assembling the IRE1 α interactome. *Mol Cell* 35, 551-561.
- Hetz, C., and Papa, F.R. (2018). The Unfolded Protein Response and Cell Fate Control. *Mol Cell* 69, 169-181.
- Holst, B., Nygaard, R., Valentin-Hansen, L., Bach, A., Engelstoft, M.S., Petersen, P.S., Frimurer, T.M., and Schwartz, T.W. (2010). A conserved aromatic lock for the tryptophan rotameric switch in TM-VI of seven-transmembrane receptors. *J Biol Chem* 285, 3973-3985.
- Humphrey, W., Dalke, A., and Schulten, K. (1996). VMD: Visual molecular dynamics. *Journal of Molecular Graphics & Modelling* 14, 33-38.
- Janowski, T., and Pulay, P. (2007). High accuracy benchmark calculations on the benzene dimer potential energy surface. *Chemical Physics Letters* 447, 27-32.
- Johnson, R.M., Hecht, K., and Deber, C.M. (2007). Aromatic and cation-pi interactions enhance helix-helix association in a membrane environment. *Biochemistry* 46, 9208-9214.

Kalli, A.C., Hall, B.A., Campbell, I.D., and Sansom, M.S. (2011). A helix heterodimer in a lipid bilayer: prediction of the structure of an integrin transmembrane domain via multiscale simulations. *Structure* **19**, 1477-1484.

Kaufman, R.J. (1999). Stress signaling from the lumen of the endoplasmic reticulum: coordination of gene transcriptional and translational controls (vol 13, pg 1211, 1999). *Genes Dev* **13**, 1898-1898.

Killian, J.A. (1998). Hydrophobic mismatch between proteins and lipids in membranes. *Biochim Biophys Acta* **1376**, 401-415.

Kitai, Y., Ariyama, H., Kono, N., Oikawa, D., Iwawaki, T., and Arai, H. (2013). Membrane lipid saturation activates IRE1 α without inducing clustering. *Genes Cells* **18**, 798-809.

Kono, N., Amin-Wetzel, N., and Ron, D. (2017). Generic membrane-spanning features endow IRE1 α with responsiveness to membrane aberrancy. *Mol Biol Cell* **28**, 2318-2332.

Korennykh, A., and Walter, P. (2012). Structural basis of the unfolded protein response. *Annu Rev Cell Dev Biol* **28**, 251-277.

Kosel, D., Heiker, J.T., Juhl, C., Wottawah, C.M., Blüher, M., Mörl, K., and Beck-Sickinger, A.G. (2010). Dimerization of adiponectin receptor 1 is inhibited by adiponectin. *J Cell Sci* **123**, 1320-1328.

Lee, A.H., Iwakoshi, N.N., and Glimcher, L.H. (2003). XBP-1 regulates a subset of endoplasmic reticulum resident chaperone genes in the unfolded protein response. *Molecular and Cellular Biology* **23**, 7448-7459.

Leekumjorn, S., and Sum, A.K. (2007). Molecular characterization of gel and liquid-crystalline structures of fully hydrated POPC and POPE bilayers. *J Phys Chem B* **111**, 6026-6033.

Leekumjorn, S., Wu, Y., Sum, A.K., and Chan, C. (2008). Experimental and computational studies investigating trehalose protection of HepG2 cells from palmitate-induced toxicity. *Biophys J* **94**, 2869-2883.

Lemmon, M.A., Flanagan, J.M., Hunt, J.F., Adair, B.D., Bormann, B.J., Dempsey, C.E., and Engelman, D.M. (1992). Glycophorin A dimerization is driven by specific interactions between transmembrane alpha-helices. *J Biol Chem* **267**, 7683-7689.

Li, E., You, M., and Hristova, K. (2005). Sodium dodecyl sulfate-polyacrylamide gel electrophoresis and forster resonance energy transfer suggest weak interactions between fibroblast growth factor receptor 3 (FGFR3) transmembrane domains in the absence of extracellular domains and ligands. *Biochemistry* **44**, 352-360.

Li, H., Korennykh, A.V., Behrman, S.L., and Walter, P. (2010). Mammalian endoplasmic reticulum stress sensor IRE1 signals by dynamic clustering. *Proc Natl Acad Sci U S A* **107**, 16113-16118.

Li, J., Jonsson, A.L., Beuming, T., Shelley, J.C., and Voth, G.A. (2013). Ligand-dependent activation and deactivation of the human adenosine A(2A) receptor. *J Am Chem Soc* **135**, 8749-8759.

Litman, B.J., Lewis, E.N., and Levin, I.W. (1991). Packing characteristics of highly unsaturated bilayer lipids: Raman spectroscopic studies of multilamellar phosphatidylcholine dispersions. *Biochemistry* **30**, 313-319.

Lyman, E., Higgs, C., Kim, B., Lupyan, D., Shelley, J.C., Farid, R., and Voth, G.A. (2009). A role for a specific cholesterol interaction in stabilizing the Apo configuration of the human A(2A) adenosine receptor. *Structure* **17**, 1660-1668.

Maiorano, J.N., and Davidson, W.S. (2000). The orientation of helix 4 in apolipoprotein A-I-containing reconstituted high density lipoproteins. *J Biol Chem* **275**, 17374-17380.

Marrink, S.J., Berger, O., Tieleman, P., and Jahnig, F. (1998). Adhesion forces of lipids in a phospholipid membrane studied by molecular dynamics simulations. *Biophys J* **74**, 931-943.

McCloy, R.A., Rogers, S., Caldon, C.E., Lorca, T., Castro, A., and Burgess, A. (2014). Partial inhibition of Cdk1 in G 2 phase overrides the SAC and decouples mitotic events. *Cell Cycle* **13**, 1400-1412.

Ng, D.P., and Deber, C.M. (2010). Modulation of the oligomerization of myelin proteolipid protein by transmembrane helix interaction motifs. *Biochemistry* **49**, 6896-6902.

Oikawa, D., Kimata, Y., Kohno, K., and Iwawaki, T. (2009). Activation of mammalian IRE1 α upon ER stress depends on dissociation of BiP rather than on direct interaction with unfolded proteins. *Exp Cell Res* **315**, 2496-2504.

Peter, B., Polyansky, A.A., Fanucchi, S., and Dirr, H.W. (2014). A Lys-Trp cation- π interaction mediates the dimerization and function of the chloride intracellular channel protein 1 transmembrane domain. *Biochemistry* 53, 57-67.

Ren, J., Lew, S., Wang, Z., and London, E. (1997). Transmembrane orientation of hydrophobic α -helices is regulated both by the relationship of helix length to bilayer thickness and by the cholesterol concentration. *Biochemistry* 36, 10213-10220.

Ridder, A., Skupjen, P., Unterreitmeier, S., and Langosch, D. (2005). Tryptophan supports interaction of transmembrane helices. *J Mol Biol* 354, 894-902.

Ruan, W., Becker, V., Klingmüller, U., and Langosch, D. (2004). The interface between self-assembling erythropoietin receptor transmembrane segments corresponds to a membrane-spanning leucine zipper. *J Biol Chem* 279, 3273-3279.

Russ, W.P., and Engelman, D.M. (1999). TOXCAT: a measure of transmembrane helix association in a biological membrane. *Proc Natl Acad Sci U S A* 96, 863-868.

Sal-Man, N., Gerber, D., Bloch, I., and Shai, Y. (2007). Specificity in transmembrane helix-helix interactions mediated by aromatic residues. *J Biol Chem* 282, 19753-19761.

Samanta, U., Pal, D., and Chakrabarti, P. (1999). Packing of aromatic rings against tryptophan residues in proteins. *Acta Crystallogr D Biol Crystallogr* 55, 1421-1427.

Schwartz, T.W., Frimurer, T.M., Holst, B., Rosenkilde, M.M., and Elling, C.E. (2006). Molecular mechanism of 7TM receptor activation--a global toggle switch model. *Annu Rev Pharmacol Toxicol* 46, 481-519.

Shyu, Y.J., Liu, H., Deng, X., and Hu, C.D. (2006). Identification of new fluorescent protein fragments for bimolecular fluorescence complementation analysis under physiological conditions. *Biotechniques* 40, 61-66.

Sirk, T.W., Brown, E.F., Sum, A.K., and Friedman, M. (2008). Molecular dynamics study on the biophysical interactions of seven green tea catechins with lipid bilayers of cell membranes. *Journal of agricultural and food chemistry* 56, 7750-7758.

Tao, R.H., and Maruyama, I.N. (2008). All EGF(ErbB) receptors have preformed homo- and heterodimeric structures in living cells. *J Cell Sci* 121, 3207-3217.

Tsuru, A., Imai, Y., Saito, M., and Kohno, K. (2016). Novel mechanism of enhancing IRE1 α -XBP1 signalling via the PERK-ATF4 pathway. *Sci Rep* 6, 24217.

Van der Spoel, D., Lindahl, E., Hess, B., Groenhof, G., Mark, A.E., and Berendsen, H.J.C. (2005). GROMACS: Fast, flexible, and free. *Journal of computational chemistry* 26, 1701-1718.

van Meer, G., Voelker, D.R., and Feigenson, G.W. (2008). Membrane lipids: where they are and how they behave. *Nat Rev Mol Cell Biol* 9, 112-124.

Volmer, R., van der Ploeg, K., and Ron, D. (2013). Membrane lipid saturation activates endoplasmic reticulum unfolded protein response transducers through their transmembrane domains. *Proceedings of the National Academy of Sciences of the United States of America* 110, 4628-4633.

Wang, M., and Kaufman, R.J. (2014). The impact of the endoplasmic reticulum protein-folding environment on cancer development. *Nat Rev Cancer* 14, 581-597.

Wang, S., and Kaufman, R.J. (2012). The impact of the unfolded protein response on human disease. *The Journal of cell biology* 197, 857-867.

Yau, W.M., Wimley, W.C., Gawrisch, K., and White, S.H. (1998). The preference of tryptophan for membrane interfaces. *Biochemistry* 37, 14713-14718.

Yoshida, H., Matsui, T., Yamamoto, A., Okada, T., and Mori, K. (2001). XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell* 107, 881-891.

Zhang, E., Lu, X., Yin, S., Yan, M., Lu, S., Fan, L., and Hu, H. (2018). The functional role of Bax/Bak in palmitate-induced lipoapoptosis. *Food Chem Toxicol* 123, 268-274.

Zhang, Y., Luo, Y., Deng, Y., Mu, Y., and Wei, G. (2012). Lipid interaction and membrane perturbation of human islet amyloid polypeptide monomer and dimer by molecular dynamics simulations. *PLoS One* 7, e38191.

Zhou, J., Liu, C.Y., Back, S.H., Clark, R.L., Peisach, D., Xu, Z., and Kaufman, R.J. (2006). The crystal structure of human IRE1 luminal domain reveals a conserved dimerization interface required for activation of the unfolded protein response. *Proceedings of the National Academy of Sciences of the United States of America* 103, 14343-14348.

Figure legends

Figure 1. The isolated IRE1 α -TMD dimerizes in both synthetic and cellular membranes.

(A) Fluorescence spectra of FITC/TAMRA-labeled TMD peptides inserted in POPC liposomes. 0.1 mol % of donor (FITC-TMD) or 0.1 mol % of acceptor (TAMRA-TMD) peptide was reconstituted in 400 μ M of lipid. The spectra (excitation at 439 nm) were measured for the samples containing the donor and acceptor peptides (red), as well as control samples containing only the donor (black) and only the acceptor (blue) peptides.

(B) Dependence of fluorescence energy transfer as a function of the acceptor/lipid molar ratio. The ratio of donor-labeled to acceptor-labeled peptides was kept at 1 while the ratio of total peptide to lipid was varied. The final concentration of the lipid was fixed at 400 μ M. The FRET efficiency was calculated as described in Methods. Values are mean $\pm$ SD, with $n = 3$.

(C) SDS-PAGE analysis of the IRE1 α -TMD peptide. The peptides dissolved in 1x lithium dodecyl sulfate (LDS) sample loading buffer were loaded onto 9% Bis-Tris gel. Bands were visualized using silver staining.

(D) (Upper) TOXCAT assays. CAT activities from cells expressing ToxR-TMD-MBP were quantified and normalized to GpA-WT. GpA-WT and its disruptive mutant (GpA-G83I) served as positive and negative controls, respectively. Values are mean $\pm$ SD, with $n = 6$. (Lower) Western Blotting analysis. The expression levels of ToxR-TMD-MBP proteins were measured using anti-MBP antibody. The expression levels were also normalized to correct the CAT activity.

(E) MalE complementation assay. The TOXCAT chimeric proteins (ToxR-TMD-MBP) were plated on M9 agar containing 0.4% maltose. MalE activity was used to show appropriate TMD topology.

Figure 2. W457A and S450A mutations abolish both dimerization and activity of the IRE1 α - Δ LD *in vivo*.

(A) Scheme of IRE1 α - Δ LD constructs used in the BiFC assay. Myc-tagged Venus N-terminus fragment and HA-tagged Venus C-terminus fragment were inserted between the signal peptide

(SP) and the truncated LD. The amino acid residue numbers of the human IRE1 α are shown in the boxes. The TMD sequence of IRE1 α is shown on top.

(B) Schematic diagram of the BiFC assay. Homodimerization of the IRE1 α - Δ LD results in a functional Venus fluorescent protein.

(C) Specific homodimerization of IRE1 α - Δ LD detected by BiFC. IRE1 $\alpha^{-/-}$ MEF cells co-expressing VN-IRE1 α - Δ LD and VC-IRE1 α - Δ LD or their W457A and S450A mutant constructs were treated with 2% BSA media, 0.4 mM palmitate (PA) or 0.4 mM oleate (OA) dissolved in 2% BSA media, or 5 μ g/mL tunicamycin (Tm) for 2 h. The Venus fluorescence was detected by confocal microscopy. Bars 42 μ m.

(D) Fluorescence quantification for (C). The corrected total cell fluorescence (CTCF) was calculated for each cell and then normalized to the PA-treated samples. 16-30 cells were quantified for each replicate. Values are reported as mean $\pm$ SD ($n = 2$), $**p < 0.001$ (Student's t-test).

(E) Western blot analysis of the total IRE1 α - Δ LD expression levels. As described in (C), cells were treated with BSA or 0.4 mM PA for 2 h. GAPDH was used as a loading control.

(F) RT-PCR of Xbp1 mRNA from cells transfected with VN-IRE1 α - Δ LD and VC-IRE1 α - Δ LD or their W457A and S450A mutant constructs. Cells were treated with 2% BSA or 0.4 mM PA for 2 h, and total mRNA isolated from the cells was subjected to RT-PCR analysis. The unspliced (U) and spliced (S) products and heteroduplex species (HD) are indicated. GAPDH was used as a loading control.

(G) Quantification of spliced XBP1 (XBP1^S). The fraction of spliced Xbp1 mRNA in each sample was quantified as described in the Methods. Data represent the means $\pm$ SD ($n = 3$), $**p < 0.001$ (Student's t-test).

(H) Alanine scanning mutagenesis of the IRE1 α -TMD. IRE1 $\alpha^{-/-}$ MEF cells co-expressing alanine mutant VN and VC plasmids were treated with BSA or 0.4 mM PA for 2 h. The fold-change of Venus fluorescence between the BSA- and PA-treated cells was quantified and normalized to WT. Values are reported as mean $\pm$ SE ($n = 2$).

Figure 3. Mutations on the juxtamembrane amphipathic helix do not significantly affect the dimerization and activity of IRE1 α - Δ LD in the cells.

(A) HeliQuest analysis of the amphipathic helix (IRE1 α 433-443 aa) (Gautier et al., 2008). Hydrophobic residues mutated to arginine (V437R and M440R) are shown in blue and the tryptophan residue is shown in red.

(B) IRE1 $\alpha^{-/-}$ MEF cells co-expressing VN-IRE1 α - Δ LD and VC-IRE1 α - Δ LD or their V437R and M440R mutants were treated with 2% BSA or 0.4 mM PA for 2 h. Venus fluorescence was detected with confocal microscopy. Bars 42 μ m.

(C) Quantification of Venus fluorescence in the presence of BSA or PA. The CTCF values were calculated for each treatment and mutation and the fold changes upon PA treatment were then normalized to the IRE1 α - Δ LD-WT sample. Values are reported as mean $\pm$ SD (n = 2), **p < 0.001 (Student's t-test).

(D) Western blot analysis of the total IRE1 α - Δ LD expression levels. GAPDH was used as a loading control.

(E) RT-PCR of Xbp1 mRNA from IRE1 $\alpha^{-/-}$ MEF cells co-transfected with VC-IRE1 α - Δ LD-WT and VN-IRE1 α - Δ LD-WT or their mutants, V437R and M440R. BSA and PA treatment conditions are the same as (B).

(F) Quantification of splicing activity of IRE1 α - Δ LD-WT, V437R and M440R mutants in the presence of BSA or PA. Values are reported as mean $\pm$ SD (n = 3), **p < 0.001 (Student's t-test).

Figure 4. The structural elements in the IRE1 α -TMD sense lipid aberrancy and activate the IRE1 α -XBP1 pathway

(A) Scheme of IRE1 α full-length (FL) constructs used in the BiFC assay. The Myc-tagged Venus N-terminus fragment and HA-tagged Venus C-terminus fragment were inserted between SP and LD.

(B) IRE1 $\alpha^{-/-}$ MEF cells co-expressing VN-IRE1 α -WT-FL and VC- IRE1 α -WT-FL, or their W457A and S450A mutants, were treated with 2% BSA or 0.4 mM PA for 2 h. Venus fluorescence was detected with confocal microscopy. Bars 42 μ m.

(C) Quantification of Venus fluorescence in (B). The CTCF values were calculated for each treatment and mutation. The fluorescence fold-change upon PA treatment was normalized to WT. Values are reported as mean $\pm$ SD (n = 3), **p < 0.001 (Student's t-test).

(D) Western blot of the WT and mutant samples with indicated antibodies. Total protein extracts were obtained from BSA- and PA-treated cells in (B).

(E) Quantification of the IRE1 α foci-positive cells in (B). Percentage of cells that contain at least one IRE1 α foci was calculated as described in the Methods. Values are reported as mean $\pm$ SD (n = 3), *p < 0.05 (Student's t-test).

(F) RT-PCR of Xbp1 mRNA from IRE1 $\alpha^{-/-}$ MEF cells co-transfected with VC-IRE1 α -FL-WT and VN-IRE1 α -FL-WT or their mutants W457A and S450A. Cells were treated with 0.4 mM PA for 0, 0.5, 1, 2 h. The unspliced (U) and spliced (S) products and heteroduplex species (HD) are indicated.

(G) Quantification of splicing activity of IRE1 α -FL-WT, W457A, and S450A mutants in (E). Values are reported as mean $\pm$ SD (n = 2), **p < 0.001 (Student's t-test).

(H) RT-PCR of Xbp1 mRNA from IRE1 $\alpha^{-/-}$ MEF cells transfected with pcDNA-hIRE1 α -FL-Y161A, pcDNA-hIRE1 α -FL-Y161A-W457A, and pcDNA-hIRE1 α -FL-Y161A-S450A. Cells were treated with BSA or 0.4 mM PA for 2 h.

(I) Quantification of splicing activity of IRE1 α -FL-Y161A, IRE1 α -FL-Y161A-W457A, and IRE1 α -FL-Y161A-S450A. Values are mean $\pm$ SD, with n = 2, **p < 0.001 (Student's t-test). Error bars may not be visible due to low values.

Figure 5. Trp457 residues form an aromatic stacking interaction in the center of the TMD dimerization interface.

(A) (Left) Distance between the center of mass (CM) of the two indole rings of Trp457 in model B. (Right) Representative structures at various time points. The colored indole rings correspond to the same colored arrows in the left panel.

(B) (Left) Distance between the CM of Trp457 and Phe460 for each TMD peptide. (Right) A representative structure of the aromatic pocket in the last 50 ns of the simulation is shown.

(C) (Left) Distance between two hydrogen atoms of the indole rings used to evaluate the alignment of the Trp457 in the simulation. Two hydrogen atoms at position 2 and 6 in the indole ring represent H1 (black) and H2 (red), respectively. (Right) Average structure obtained from the last 50 ns of the simulation time. H1 and H2 are highlighted in black and red, respectively. Nitrogen atom is colored in blue and other hydrogen atoms not considered for this analysis are colored in white.

(D) Residue contact numbers were calculated as the number of times each residue comes within 6 Å of the dimerization interface during the last 50 ns of the simulation. The contact numbers were further normalized to the contact number of the Trp457 residue. Dimerization deficiency was calculated as [1-Fluorescence] from the data in Figure 2H. Residues identified from both the MD simulations and alanine scanning experiments, residues identified only from the MD simulations, and residues identified only from the alanine scanning experiments are shown in red, blue, and green, respectively.

(E) Location of residues identified from the MD simulations or alanine scanning experiments are highlighted on the IRE1 α -TMD dimer structure obtained from model B. Colors are shown as in (D).

(F) Density profile in z dimension of the Trp457 (red) and POPC head group (black) as a function of distance from the center of the membrane. The density distributions were collected from 10,000 snapshots during the last 50 ns of the simulations. The root mean square distance of Trp457 from the center of the bilayer ($\langle Z_{\text{cf}}^2 \rangle^{1/2}$) was calculated over the last 50 ns of the simulation.

(G) Closer view of the density profile of (F).

(H) Nitroxide lipid quenching of IRE1 α -TMD peptide in small unilamellar vesicles (SUVs). The SUVs containing TMD peptide, POPC, and spin-labeled shallow quencher (5 Doxyl PC) or deep

quencher (16 Doxyl PC) were prepared as described in the Methods. The tryptophan fluorescence was excited at 280 nm and the emission spectra were obtained by scanning from 295nm to 450nm.

Figure 6. Intrinsic structural features of Trp457 are important for the dimerization process of IRE1 α -TMD and Ser450 stabilizes the Trp457-Trp457 interaction.

(A) TOXCAT assays of IRE1 α -TMD-WT and its W457A and W457F mutants. (Upper) CAT activities from cells expressing ToxR-TMD-MBP were quantified and normalized to the WT. (Lower) The protein expression levels for each sample were also normalized to the ToxR-TMD-MBP protein. Values are mean $\pm$ SD (n = 6), **p < 0.001 (Student's t-test).

(B) Initial (0 ns) and final (200 ns) conformation of the W457A mutant. The mutated alanine residues are colored in red.

(C) Initial and final conformations of the W457F mutant. The mutated phenylalanine residues are colored in red.

(D) Distance between the CM of the TMD peptides. The black, orange, and blue lines correspond to the WT, W457A, and W457F mutants, respectively.

(E) (Left) Distance between W457F residues as measured from the CM of the two benzene rings. (Right) Representative structures at various time points. The colors of the phenylalanine residues match the color of the arrows in the left panel.

(F) Initial and final conformations of the S450A mutant. Both mutated serine residues and Trp457 residues are colored in red and magenta, respectively.

(G) Distance between the CM of WT (black) and S450A (red) peptides.

(H) Distance between the indole rings of Trp457 residues in WT (black) and S450A (red) peptides.

Figure 7. The proposed model of IRE1 α activation by membrane lipid saturation.

Under normal conditions, IRE1 α presents as an inactive monomer. Upon palmitate treatment, the center-localized tryptophan residues (red) rapidly initiate the TMD dimerization process and

support the core interactions for the IRE1 α -TMD. The serine residue (green) assists in the stability of the TMD dimer during this process. The dimerization of IRE1 α -TMD further induces dimerization of LD, KD, and RD, subsequently resulting in an active conformation.

Figures

Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.





Figure 7.



STAR•METHODS

Experimental Model and Subject Details

Cell lines used:

IRE1 $\alpha^{-/-}$ MEF grown in DMEM containing 4500 mg/L glucose, L-glutamine, sodium pyruvate and sodium bicarbonate supplemented with 10% fetal bovine serum +10% FBS at 37°C in 5% CO₂

Hep3B grown in DMEM containing 4500 mg/L glucose, L-glutamine, sodium pyruvate and sodium bicarbonate supplemented with 10% fetal bovine serum +10% FBS at 37°C in 5% CO₂

Quantification and Statistical Analysis

Statistical details of experiments can be found in the figure legend. The value of n is specified for each figure in the legend as well as the p-value when statistical test was performed.

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-calnexin	Santa Cruz Biotech	Cat# sc-23954
Anti-HA	Abcam	Cat# ab9110
Anti-Myc	Abcam	Cat# ab9106
Goat Anti-Rabbit IgG- Alexa Fluor® 488	Abcam	Cat# 150077
Goat Anti-Mouse IgG- Alexa Fluor® 568	Abcam	Cat# 175701
Anti-IRE1 α	Cell Signaling Tech	Cat# 3294S
Anti-GAPDH	GeneTex	Cat# GTX100118
Anti-MBP	New England Biolabs	Cat# E8032
Bacterial and Virus Strains		
E coli Top10	ThermoFisher	Cat# C404010
NT326(malE deficient) E coli	Donald M. Engelman Lab	
Chemicals, Peptides, and Recombinant Proteins		
1-palmitoyl-2-stearoyl-(5-doxyl)-sn-glycero-3-phosphocholine	Avanti Lipids	Cat# 810601
1-palmitoyl-2-stearoyl-(16-doxyl)-sn-glycero-3-phosphocholine	Avanti Lipids	Cat# 810604
1-palmitoyl-2-oleoyl-glycero-3-phosphocholine	Avanti Lipids	Cat# 850457
⁴⁴⁴ KDMATIILSTFLLIGWVAFIITYPLS ⁴⁶⁷ -KK	Peptide 2.0	custom
FITC- ⁴⁴⁴ KDMATIILSTFLLIGWVAFIITYPLS ⁴⁶⁷ -KK	Peptide 2.0	custom
TAMRA- ⁴⁴⁴ KDMATIILSTFLLIGWVAFIITYPLS ⁴⁶⁷ -KK	Peptide 2.0	custom

Phos-tag™ Acrylamide 5 mM Aqueous Solution	FUJIFILM Wako Pure Chemical Corporation	Cat# 304-93526
Experimental Models: Cell Lines		
IRE1 $\alpha^{-/-}$ MEF	David Ron lab	
Hep3B	ATCC	
Oligonucleotides		
5'-GAACCAGGAGTTAAGAACACG-3'	mXBP1-f (for XBP1 splicing assay)	Eurofins Genomics
5'-AGGCAACAGTGTCTCAGAGTCC-3'	mXBP1-r (for XBP1 splicing assay)	Eurofins Genomics
5'-GGGGAATCGAGCTAGCATGGCTACCATCAT CCTGA-3'	IRE1 α -TM-f (IRE1 α - TM amplification for pcc-IRE1 α -WT)	Eurofins Genomics
5'-GGGTTGATCAGGATCCCAGGGGATAGGTG ATGATG-3'	IRE1 α -TM-r (IRE1 α - TM amplification for pcc-IRE1 α -WT)	Eurofins Genomics
5'-ATAAATCGGCTCCAACTTGGGGAATCGAG-3'	pcc-f (pccKAN vector amplification for pcc-IRE1 α -WT)	Eurofins Genomics
5'-GTTTAAAGCTGGATTGGCTTGGGTTGATCA GGATC-3'	pcc-r (pccKAN vector amplification for pcc-IRE1 α -WT)	Eurofins Genomics
Recombinant DNA		
pBiFC-VC155	Addgene	Plasmid# 22011
pBiFC-VN155(I152L)	Addgene	Plasmid# 27097
pcDNA-hIRE1 α	Addgene	Plasmid# 21892
pccKAN	Donald M. Engelman Lab	
pccGpA-WT	Donald M. Engelman Lab	
pccGpA-G83I	Donald M. Engelman Lab	

Contact for Reagent and Resource Sharing

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Professor Christina Chan (krischan@egr.msu.edu).

Method Details

Cell culture and transfection

IRE1 $\alpha^{-/-}$ mouse embryonic fibroblast (MEF) cells were maintained at 37°C at 5% CO₂ in Dulbecco's Modified Eagle Medium (DMEM, Thermo cat# 11995065) containing 4500 mg/L

glucose, L-glutamine, sodium pyruvate, and sodium bicarbonate supplemented with 10% fetal bovine serum. Cells were reverse-transfected with Lipofectamine® LTX reagent (Thermo Fisher Scientific) for 24 h. Afterward, the media was removed and replaced with complete media containing only 2% bovine serum albumin (BSA) or 0.4 mM palmitate (PA) in 2% BSA media.

Cloning of VC-IRE1 α and VN-IRE1 α plasmids

pBiFC-VC155 and pBiFC-VN155 (I152L) were a gift from Chang-Deng Hu (Addgene plasmid # 22011 and #27097) (Shyu et al., 2006). pcDNA-hIRE1 α (a gift originally from Randal J. Kaufman (University of Michigan, MI, USA)) was used to amplify the IRE1 α gene. An IRE1 α signal peptide sequence (1-18 aa) was added at the N-terminal of the coding sequence upstream of the HA- and Myc-tagged VC and VN fragments. The coding sequences of the truncated luminal domain (19-29aa, 408-443aa) followed by the transmembrane and cytosolic domains (444-977 aa) were inserted at the C-terminus of the VC and VN fragments (see Figure 2A). For the IRE1 α -full length BiFC constructs, the HA- and Myc-tagged VC and VN fragments were inserted between the signal peptide sequence and the C-terminus coding sequences of IRE1 α (18-977aa). Primers for the alanine point mutations of IRE1 α -TMD and the V437R, M440R, Y161A mutations were designed and performed according to guidelines from the QuikChange Lightning Site-Directed Mutagenesis kit (Agilent). The sequences of the mutations were confirmed by DNA sequencing services (Eurofins Genomics).

Immunofluorescence

Hep3B Cells were reverse transfected with pBiFC-VC155-IRE1 α - Δ LD and pBiFC-VN155-IRE1 α - Δ LD or pBiFC-VC155-IRE1 α -FL and pBiFC-VN155-IRE1 α -FL plasmids using Lipofectamine LTX for 24 h and then fixed, permeabilized, and co-stained with an anti-calnexin antibody (Santa Cruz Biotech, cat# sc-23954), an anti-HA antibody (Abcam, cat# ab91110), or an anti-Myc antibody (Abcam, cat# ab9106). Appropriate secondary antibodies conjugated with Alexa 488 (Abcam, cat# 150077) or Alexa 568 (Abcam cat# 175701) were further incubated. Representative images for cells expressing Myc-VN-IRE1 α - Δ LD protein are shown although the cells expressing HA-VC-IRE1 α - Δ LD protein had the same colocalization pattern.

Bimolecular fluorescence complementation (BiFC) assay

μ -plates (Ibidi) were used for confocal microscopy to detect Venus fluorescence. IRE1 $\alpha^{-/-}$ MEF cells were seeded at a density of 3×10^6 cells/mL in each well and co-transfected with VN155-IRE1 α - Δ LD and VC155-IRE1 α - Δ LD (WT or V437R/M440R/W457A/S450A) plasmids. Cells were transfected as described above and then treated with BSA or PA media for 2 h. The Olympus FluoView FV1000 imaging system, with an UPLFLN 20X NA 0.5 objective and a PLAPON 60X O NA 1.42 objective, was used to observe the formation of the Venus fluorophore at $\lambda_{\text{ex}} = 515\text{nm}$ and $\lambda_{\text{em}} = 535\text{nm}$. The BiFC assay for IRE1 α -FL was performed for the same conditions as IRE1 α - Δ LD constructs using the VN155-IRE1 α -FL and VC155-IRE1 α -FL (WT or W457A/S450A/W457A-S450A) plasmids. The corrected total cell fluorescence (CTCF) was calculated for each cell (McCloy et al., 2014) using the following equation: $\text{CTCF} = [\text{Integrated Density} - (\text{Area of selected cell} \times \text{mean fluorescence of background readings})]$. The number of cells that contain at least one foci was analyzed using ImageJ (Fiji) software with the following parameter settings, threshold 15-255, circularity 0-1, and size (pixel²) 100-10000.

XPB1 splicing assay

IRE1 $\alpha^{-/-}$ MEF cells were reverse transfected with the plasmids as described above. Total mRNA was collected and the one-step RT-PCR kit (Qiagen) was used to amplify XPB1 cDNA. PCR primers were designed to flank the 26-bp splicing sequence of XPB1. PCR reaction was performed using the following primers: mXPB1f 5'-GAACCAGGAGTTAAGAACACG-3' and mXPB1r 5'-AGGCAACAGTGTCTCAGAGTCC-3'. The PCR products were run on a 2.5% agarose gel made in 0.5X TBE. Gels were visualized and quantified using Molecular Imager ChemiDoc XRS System (Bio Rad). The splicing efficiency of XPB1 (% XPB1) was calculated with the following equation: $\text{XPB1 splicing (\%)} = [\text{Intensity (XPB1}^{\text{S}}) + 0.5 * \text{Intensity (XPB1}^{\text{HD}})] / [\text{Intensity (XPB1}^{\text{S}}) + \text{Intensity (XPB1}^{\text{HD}}) + \text{Intensity (XPB1}^{\text{U}})] * 100$

Förster resonance energy transfer (FRET) measurements

The lysine residues on the N and C-termini (K-⁴⁴⁴KDMATIILSTFLIGWVAFIITYPLS⁴⁶⁷-KK) were added to enhance the solubility and stability in the lipid bilayer. Unlabeled TM peptide and N-terminal labeled-TMD peptides with FITC (Fluorescein isothiocyanate) or Carboxytetramethylrhodamine (TAMRA) were synthesized from Peptide 2.0 Inc. Peptides dissolved in TFE (Trifluoroethanol) and POPC (1-

palmitoyl 2-oleoyl phosphatidylcholine) suspended in chloroform were mixed and then the organic solvents were removed using a nitrogen gas stream. The peptide/lipid mixture was reconstituted in 10 mM phosphate buffer (pH 7.0) containing 500 mM NaCl, freeze-thawed three times, and equilibrated at RT for 1 h before the fluorescence measurement.

FRET experiments were performed with FluoroMax-4 (Horiba), as previously described (Cho et al., 2012). For FITC/TAMRA as a donor/acceptor pair, the excitation wavelength was set at 439 nm and emission spectra were collected from 480 to 650 nm. FRET efficiency was calculated from measurements of donor fluorescence intensity at 515 nm in the absence and presence of the acceptor: $\text{FRET efficiency} = (ID - IDA)/ID$, where ID and IDA are the donor fluorescence intensities of the samples containing only donor-labeled peptides and the samples with both donor- and acceptor-labeled peptides, respectively. The contribution to the emission at 515 nm from the direct excitation of the acceptor was removed by subtracting the spectra of the samples containing only acceptor-labeled peptides. In addition, the background signal from the lipid was eliminated by subtracting the spectra of liposome samples without peptides.

SDS-PAGE for IRE1 α -TMD peptide

The IRE1 α -TMD at the final concentration 2 μ M was incubated in 1X lithium dodecyl sulfate (LDS) buffer (final concentration = 2% LDS) (ThermoFisher Scientific Inc., cat# NP0007) for 20 min at room temperature. 5 μ L of sample was loaded onto 9% Bis-Tris gel and ran in MES buffer. Spectra Multicolor Low Range Protein Ladder (Thermo Fisher Scientific Inc., cat# 26628) was used as a molecular weight marker. The IRE1 α -TMD peptides were visualized with Silver staining.

Tryptophan fluorescence measurements and tryptophan depth calculation

For nitroxide lipid quenching experiments, the small unilamellar vesicles (SUVs) were prepared by a modified method from a previous study (Maiorano and Davidson, 2000). The spin labels used, 5 doxyl PC (1-palmitoyl-2-stearoyl-(5-doxyl)-sn-glycero-3-phosphocholine) and 16 doxyl PC (1-palmitoyl-2-stearoyl-(16-doxyl)-sn-glycero-3-phosphocholine) were obtained from Avanti Lipids. The molar ratio of POPC: doxylPC: peptide was 68:12:1 for all the assays. The lipid and peptide mixture in chloroform was dried under nitrogen, vacuum dessicated for 8 h to remove all traces of chloroform. 10 μ L ethanol was further added to the lipid-peptide film. The

sample was then vortexed and incubated on ice for 10 min. After diluting the sample with 1 mL of 10 mM Tris-HCl buffer (pH 7.0), bath sonication was performed for 1 h. The spectra were recorded at room temperature on a Fluoromax-4 at λ_{ex} = 280nm. Emission spectra were collected from 295nm to 450nm. The fluorescence from POPC SUVs was subtracted from the samples containing spin label and peptide.

The distance of the Trp from the center of the bilayer (Z_{cf}) is calculated with the following equation (Ren et al., 1997):

$$Z_{cf} = L_{cd} - \left[\frac{\ln \left(\frac{\left(\frac{F_s}{F_0} \right)^2}{\frac{F_d}{F_0}} \right)}{\pi C} - 2L_{ds}^2 + 4L_{cd}^2 \right] / 4(L_{ds} + L_{cd})$$

where F_0 is the fluorescence intensity in the absence of quencher, F_s is the fluorescence intensity in the presence of the shallow quencher (5-doxyl PC), F_d is the fluorescence intensity in the presence of the deep quencher (16-doxyl PC), L_{cd} is the distance from the center of the bilayer to the deep quencher (2.25 Å), and L_{ds} is the distance between the shallow (12.15 Å) and deep quenchers.

TOXCAT chimera constructions and assay

The TOXCAT vectors, pccKAN (TM deficient negative control), pccGpA-WT (positive control), and pccGpA-G83I (negative control), along with NT326 (malE deficient) *E.coli* strain, were kindly provided by Dr. Donald M. Engelman (Yale University, CT, USA) (Russ and Engelman, 1999). DNA fragment coding for the IRE1 α -TMD residues (444-466 aa) was amplified from the plasmid pcDNA-hIRE1 α and then inserted in-frame into the NheI and BamHI sites of pccKAN to generate the ToxR-IRE1 α -TMD-MBP (maltose binding protein) fusion protein. Individual mutants of IRE1 α -TMD were created through mutation of the pcc-IRE1 α -WT plasmid using the QuikChange Lightning site-directed mutagenesis kit (Agilent). The TOXCAT chimeras were transformed into NT326 cells and grown at 37°C until A600 was 0.6. The harvested cells were lysed using the xTractor cell lysis buffer (Clontech) containing 0.2 mg/mL lysozyme. The CAT concentration was quantified with the CAT enzyme-linked immunosorbent

assay kit (Roche Applied Science) according to the product manual. Chimera protein expression was quantified with western blotting using anti-MBP antibody (New England Biolabs, cat# E8032) and used to normalize the CAT activity of the various constructs.

MalE complementation assay

To confirm correct membrane insertion, NT326 cells expressing the TOXCAT chimeras were grown overnight in LB media. The cells were inoculated in M9 minimal media and cultured for 8 h. The cells were streaked onto M9 minimal media agar plates containing 0.4% maltose as the only carbon source and incubated for 3 days at 37°C. The images of the agar plates were collected using the Molecular Imager ChemiDoc XRS System (Bio Rad).

Molecular dynamics (MD) protocol

MD simulations of the TM peptides in lipid bilayer were performed with the GRONINGEN Machine for Chemical Simulations (GROMACS) v4.5.4 package (Hess B, 2008), using the gromos53a6 force field, extended to improve the simulation of the lipid components consistent with our previous work (Leekumjorn and Sum, 2007; Leekumjorn et al., 2008; Marrink et al., 1998; Sirk et al., 2008). The lipid bilayer consisted of 288 1-palmitoyl 2-oleoyl phosphatidylcholine (POPC) lipids and 11500 water molecules (SPC model (Berendsen, 1981)) in the liquid-crystalline state (gel-to-liquid crystalline phase transition at 270 K (Litman et al., 1991)). This model was chosen since it gives an approximate representation of the phospholipid constituents in the ER (Buchan et al., 2010).

The topology file for the POPC molecule was consistent with the previous topology described in (Leekumjorn and Sum, 2007; Leekumjorn et al., 2008; Marrink et al., 1998; Sirk et al., 2008). An initial 100 ns MD simulation of a pure POPC bilayer was performed to equilibrate the membrane. The TM helix sequence and all mutants were built with the PyMol software (De Lano, 2002), assigning an α -helical conformation as predicted by PSIPRED: Protein Structure Prediction Server (<http://psipred.net/psiform.html>) (Kalli et al., 2011). We built three different dimer models of the TM peptides (models A, B, C). Starting from a dimer conformation (model A), the other two potential dimers were gained by systematically rotating each TM peptide 90° in model A around its backbone (z-axis) clockwise or counter-clockwise (Zhang et al., 2012). All

the TM chains in the three dimer models have the same initial conformation and the only difference for the three dimers is the helix-helix interface.

The TM segments were then inserted in the pre-equilibrated POPC bilayer at 1.2 nm from each other. The insertion required the removal of two lipid molecules per leaflet. Chloride ions were added to neutralize the charges in the simulation box. To relax the bond geometries, the potential energy of the system (lipid bilayer, peptides, water and ions) was minimized by using the steepest-descent method until convergence was reached. The solvent was then relaxed by 1 ns of MD at 300K, restraining the peptides' atomic positions with a harmonic potential of $1000 \text{ kJ mol}^{-1} \text{ nm}^{-2}$ and then for another 1 ns decreasing the harmonic potential to $500 \text{ kJ mol}^{-1} \text{ nm}^{-2}$. The production simulations were performed at 300 K and 1 bar in an isothermal-isobaric ensemble (NPT), using the Berendsen thermostat and barostat. Lipids, water, peptide, and counterions were separately coupled to the temperature bath by weak coupling with a coupling constant of 0.1 ps. The pressure was also weakly coupled (with a coupling constant of 1.0 ps and compressibility of 4.5×10^{-5} using a semi-isotropic scheme in which only the pressure in the x and y were coupled together, but the normal pressure (in the z -direction) was not, as recommended for membrane protein simulations.

A simulation time step of 2 fs was used with all bond lengths constrained via the LINCS algorithm. Long-range electrostatic interactions were calculated with the Particle-Mesh Ewald (PME) method. Short-range interactions were cut-off at 1 nm. The production run consisted of 200 ns, and only the last 50 ns were used for the calculation of the average structure. The configurations in these production runs were stored every 2500 steps (5 ps), ensuring negligible statistical correlation between successively stored configurations. The trajectories were analyzed using GROMACS (Van der Spoel et al., 2005) and Visual Molecular Dynamics (VMD) (Humphrey et al., 1996).