

Application of protoplast technology to CRISPR/Cas9 mutagenesis: from single-cell mutation detection to mutant plant regeneration

Choun-Sea Lin^{1,*} , Chen-Tran Hsu¹, Ling-Hung Yang¹, Lan-Ying Lee², Jin-Yuan Fu¹, Qiao-Wei Cheng¹, Fu-Hui Wu¹, Han C.-W. Hsiao³, Yesheng Zhang⁴, Ru Zhang⁴, Wan-Jung Chang^{1,†}, Chen-Ting Yu¹, Wen Wang⁴, Li-Jen Liao⁵, Stanton B. Gelvin^{2,*}  and Ming-Che Shih^{1,*}

¹Agricultural Biotechnology Research Center, Academia Sinica, Taipei, Taiwan

²Department of Biological Sciences, Purdue University, West Lafayette, IN, USA

³Department of Bioinformatics and Medical Engineering, Asia University, Taichung City, Taiwan

⁴State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, China

⁵Institute of Life Science, National Kaohsiung Normal University, Kaohsiung, Taiwan

Received 6 August 2017;

revised 22 November 2017;

accepted 3 December 2017.

*Correspondence (Tel +886-2-2787-2094;

fax +886-2-2651-5600; email

cslin99@gate.sinica.edu.tw (C.-S.L.) and

(Tel +1-765-494-4939; fax +1-765-496-

1496; email gelvin@purdue.edu (S.B.G.))

and; (Tel +886-2-2789-9403; fax +886-2-

2651-5600; email

mcshih@gate.sinica.edu.tw (M.-C.S.))

†Present address: Department of

Biochemistry, University of Pennsylvania,

Philadelphia, PA, 19104-6030, USA.

Summary

Plant protoplasts are useful for assessing the efficiency of clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated protein 9 (Cas9) mutagenesis. We improved the process of protoplast isolation and transfection of several plant species. We also developed a method to isolate and regenerate single mutagenized *Nicotiana tabacum* protoplasts into mature plants. Following transfection of protoplasts with constructs encoding Cas9 and sgRNAs, target gene DNA could be amplified for further analysis to determine mutagenesis efficiency. We investigated *N. tabacum* protoplasts and derived regenerated plants for targeted mutagenesis of the *phytoene desaturase* (*NtPDS*) gene. Genotyping of albino regenerants indicated that all four *NtPDS* alleles were mutated in amphidiploid tobacco, and no Cas9 DNA could be detected in most regenerated plants.

Keywords: protoplast isolation, CRISPR/Cas9, protoplast regeneration, single-cell analysis.

Introduction

Genome engineering is an important component of the relatively novel field of synthetic biology. Gene editing, the directed change of a specific DNA sequence, is an important element of genome engineering. The clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated protein 9 (Cas9) system is a convenient genome-editing tool that requires only two reagents: Cas9 protein and a single guide RNA (sgRNA) (Feng *et al.*, 2013; Gaj *et al.*, 2013; Li *et al.*, 2013; Nekrasov *et al.*, 2013; Shan *et al.*, 2013). Following CRISPR-mediated mutagenesis, integrated transgenes encoding gene editing reagents can often be removed from the genome through genetic segregation, mitigating public concerns regarding genetically modified organisms (Huang *et al.*, 2016). CRISPR/Cas9 mutagenesis is therefore becoming an important technology for basic plant science and agriculture.

Numerous published studies have described various vectors to improve CRISPR-mediated target mutagenesis. These studies describe parameters such as promoters, different versions of Cas9, and the use of multiple sgRNAs (Ali *et al.*, 2015, 2016; Belhaj *et al.*, 2013; Bortesi and Fischer, 2015; Butt *et al.*, 2017; Cermak *et al.*, 2017; Eid *et al.*, 2016; Kaya *et al.*, 2016; Ma *et al.*, 2015; Murovec *et al.*, 2017; Shimatani *et al.*, 2017; Wang *et al.*, 2015b; Yan *et al.*, 2015). Not only Cas9, but other

endonucleases such as Cpf1 can induce mutations (Endo *et al.*, 2016; Kim *et al.*, 2017; Mahfouz, 2017; Xu *et al.*, 2017). However, stable transformation to evaluate CRISPR mutagenesis efficacy can be time-consuming. Transient protoplast transfection is an alternative strategy to test multiple mutagenesis parameters rapidly. Protoplasts from at least five crop species (rice, wheat, maize, lettuce, and tomato), in addition to *Arabidopsis* and tobacco, have been used to evaluate gene editing reagents using CRISPR/Cas9-based systems (Cermak *et al.*, 2015; Liang *et al.*, 2014; Shan *et al.*, 2014; Woo *et al.*, 2015). Although several methods have been developed to obtain and transfect *Arabidopsis* protoplasts (Sheen, 2001; Wu *et al.*, 2009), protoplast isolation remains a bottleneck to testing genome-editing reagents in many crop species.

Previously, DNA from pooled mutagenized protoplasts was used to determine target site mutagenesis efficiency (Cermak *et al.*, 2015; Liang *et al.*, 2014; Shan *et al.*, 2014; Woo *et al.*, 2015). The target region is amplified by PCR, and the resulting amplicons are further evaluated using restriction fragment length polymorphism (RFLP) (Feng *et al.*, 2013; Nekrasov *et al.*, 2013; Shan *et al.*, 2013) or cleaved amplified polymorphic sequence analysis (Kaya *et al.*, 2016; Mikami *et al.*, 2015a,b; Shimatani *et al.*, 2017), T7 endonuclease I (T7E1) analysis (Kim *et al.*, 2017; Woo *et al.*, 2015) or next-generation sequencing (NGS) (Kim *et al.*, 2017; Woo *et al.*, 2015).

Because the PCR amplicons constitute a mixture of wild-type and mutated DNA, mutagenesis efficiency is determined by calculating the gel image density (RFLP and T7E1 assays) or the per cent of mutant sequences (by NGS). It is often difficult to detect low target site mutagenesis efficiency. False-positive results can occur because of incomplete restriction endonuclease digestion or PCR errors. Although NGS can resolve these problems, the process can be time-consuming and costly. Because there are only two potential target site alleles in a diploid cell, single-cell DNA analysis can rapidly determine mutagenesis efficiency. Several single-cell isolation methods have been published (Brennecke *et al.*, 2013; Efroni and Birnbaum, 2016; Efroni *et al.*, 2015; Gierahn *et al.*, 2017; Klein *et al.*, 2015; Macosko *et al.*, 2015; Yamamoto *et al.*, 2016). Single-cell analyses have been applied to transcriptome and metabolome studies (Efroni *et al.*, 2015; Yamamoto *et al.*, 2016). However, these studies required expensive facilities or technically demanding protocols, including flow cytometry (Gierahn *et al.*, 2017; Klein *et al.*, 2015; Macosko *et al.*, 2015) or microinjection (Yamamoto *et al.*, 2016). A convenient and reliable single-cell isolation protocol would greatly benefit plant scientists conducting gene editing experiments.

Protoplasts can be used to determine target site mutagenesis efficiency and can be regenerated into plants (Woo *et al.*, 2015). Furthermore, genome-editing reagents such as sgRNAs and Cas9 protein can be synthesized and assembled *in vitro* to form active ribonucleoprotein (RNP) complexes. These complexes can be delivered into protoplasts and mutagenize the target gene. Thus, target mutants can be obtained without the presence of exogenous DNA (Kim *et al.*, 2017; Liang *et al.*, 2017; Woo *et al.*, 2015). Such DNA-free genome editing avoids stable introduction of transgenes. Although geminiviruses can be used in whole plants to deliver donor DNA for homology-directed repair (HDR; Wang *et al.*, 2017), protoplast transfection is an alternative to deliver high amounts of DNA required for HDR. However, protoplast regeneration is difficult in most plant species.

In this report, we further develop protoplast isolation protocols for several crop and ornamental species, and the model plant *Arabidopsis*. We used these protoplasts to evaluate CRISPR/Cas9 mutagenesis efficiency. We describe a simple single-protoplast isolation protocol and use this protocol to edit the tobacco *NtPDS* gene. Multiple plants regenerated from single mutagenized tobacco protoplasts contain a variety of CRISPR-induced mutations.

Results

Improvement of protoplast isolation for transfection

We designed a tool to generate multiple longitudinal cuts in monocot seedlings. A razor blade was divided into four pieces, which were stacked in parallel on a scalpel handle (Figure 1a). In previous reports, seedlings were cut in cross section (Zhang *et al.*, 2011; Figure 1b). In this report, seedlings were sorted and arranged in parallel (or affixed to clear tape) for longitudinal cutting (Figure 1c). When a seedling is cut in cross section, only the cells near the edge are digested (Figure 1d); cutting in longitudinal sections increases the release of protoplasts (Figure 1e). The protoplast yield from rice seedlings subjected to longitudinal cutting was higher than that from seedlings subjected to cross-cutting [4.8×10^6 protoplasts/g fresh weight (FW) vs 2.2×10^6 protoplasts/g FW]. Longitudinal cutting permitted efficient cell wall digestion by cellulose R10 and macerozyme R10 enzymes (Table S1), which are less expensive than are cellulase RS and macerozyme RS enzymes used

previously (Zhang *et al.*, 2011). Longitudinal cutting was used successfully for five Poaceae species: rice, wheat, maize, millet and bamboo. Protoplasts derived from this protocol were transfected with a red fluorescence protein (RFP) gene by a PEG-mediated method. Total intact cell number was calculated, and the transfection efficiency was calculated as the percentage of intact protoplasts with RFP fluorescence/intact protoplasts. Transfection efficiencies of these protoplasts were >40% (Figure 1j and k, bamboo: 54%; millet: 51%; rice: 44%; maize: 47%; wheat: 41%; Figure S1).

We previously established an efficient 'Tape-*Arabidopsis* Sandwich' protoplast isolation protocol (Wu *et al.*, 2009). This protocol can be applied to several Brassicaceae species, including *Brassica oleracea*, *B. napus*, *Cleome spinosa*, *C. monophylla*, and *C. gynandra*. Using 3M Scotch Tape, we peeled off the epidermal layer of leaves or cotyledons, facilitating protoplast release. Our results indicated that 2- to 3-week-old *in vitro* cotyledons from *B. napus* were more suitable for protoplast isolation than were 1-week-old cotyledons. For *Cleome* species, mature leaves from plants grown in a greenhouse were suitable. *C. gynandra* is a C4 plant, so there are two types of protoplasts, from mesophyll and bundle sheath cells. Protoplasts from all six Brassicaceae species isolated by the Tape Sandwich method were transfected using a PEG-mediated method with efficiencies >40% (Figure S1; *Arabidopsis*: 67%; broccoli: 43%; rapeseed: 63%; *C. gynandra*: 46%; *C. spinosa*: 79%; *C. monophylla*: 83%).

It was difficult to isolate high-quality protoplasts from tomato leaves using the Tape Sandwich method (data not shown). Instead, a suspension cell line was developed from tomato hypocotyls using the tomato cultivar 'Micro-Tom' (Figure 1f–i). Tomato hypocotyls were cut, callus cultures were developed, and the callus used to establish a suspension cell culture (Figure 1g,h). The proliferation rate of this tomato suspension line is similar to that of tobacco BY-2 cells. Protoplasts were isolated by incubating the suspension cells in a cell wall digestion buffer. Similar to BY-2 protoplasts, the tomato protoplasts were transfected using a PEG-mediated transfection method with 63% efficiency (Figures 1l,m, and S1).

CRISPR/Cas9-mediated mutagenesis of protoplasts

To evaluate mutagenesis efficiency, we targeted in most species the phytoene desaturase (*PDS*) gene. For each *PDS* gene, we chose the sgRNA such that a restriction enzyme site upstream of the protospacer adjacent motif (PAM; Figure 2a) sequence may be lost if target mutations occurred. If there were no suitable sgRNA targeting site (e.g. lacking a PAM sequence) or unique sequence for *PDS*, a published sgRNA for another gene was used.

Protoplasts from each species were transfected with a VirD2-NLS-mRFP plasmid (Lee *et al.*, 2008) and pCambia1300-OsU3 (*AarI*)-Cas9 or pCambia1300-OsU6(*AarI*)-Cas9 carrying the species-specific sgRNA (Figures 2b and S2). After 24 or 48 h, DNA from total protoplasts was extracted and the target gene amplified by PCR. The PCR product was digested with a restriction enzyme whose site is adjacent to the PAM, and the products separated by electrophoresis through an agarose gel (Figure 2c). Undigested DNA appeared in the 48 h sample. This result suggests that some genomic DNA had been mutated, disrupting the restriction endonuclease site targeted by the sgRNA. The undigested PCR fragments were cloned into a T/A vector, transformed into *Escherichia coli*, and the insert subjected to colony PCR. The PCR fragment was digested by the appropriate restriction endonuclease, and PCR products not digested by the




Figure 1 Improved protoplast isolation. (a) A razor blade was divided into four pieces and the pieces assembled in parallel in a scalpel handle. (b) Seedlings of rice were cut in cross section and placed in digestion solution. Bar = 1 cm. (c) Seedlings of rice were cut longitudinally (in the same direction as the veins) and placed in digestion solution. Bar = 1 cm. (d) Microscopic image of rice seedlings cut in cross section (perpendicular to vascular bundles) after 3-h digestion. Bar = 10 μ m. (e) Microscopic image of rice seedlings cut in longitudinally (parallel to vascular bundles) after 3-h digestion. Bar = 10 μ m. (f) Tomato hypocotyl sections were incubated in medium supplemented with 10 mg/L NAA and photographed after 1 month. Bar = 1 cm. (g) Microscopic image of tomato suspension cells. Bar = 40 μ m. (h) Tomato suspension cells grown in 1 mg/L 2,4-D after 7 days. Bar = 1 cm. (i) Tomato suspension cells formed calli on solid medium supplemented with 1 mg/L 2,4-D. Bar = 1 cm. (j) mRFP-NLS plasmid DNA was delivered to rice protoplasts using a PEG-mediated method. Protoplasts were photographed after 24 h. Red colour indicates RFP epifluorescence. Bar = 50 μ m. (k) Overlay of epifluorescence and bright field images of transfected rice protoplasts. Bar = 50 μ m. (l) mRFP-NLS plasmid DNA was delivered into tomato Micro-Tom protoplasts using a PEG-mediated method. Protoplasts were photographed after 24 h. Red colour indicates RFP epifluorescence. Bar = 50 μ m. (m) Overlay of epifluorescence and bright field images of transfected tomato protoplasts. Bar = 50 μ m.

restriction enzyme were sequenced (Figure 2d–f). Protoplast transfection and CRISPR/Cas9 editing were evaluated for each of the nine species. The results are summarized in Table 1.

CRISPR editing of *Poaceae* protoplasts

Bamboo

For the bamboo *PDS* gene, a target sequence (sg2) that contains a *BsaI* restriction site was chosen. Results of the DNA amplifications and digestions after the transfection are shown in Figure S3. After 48-h transfection, there was a minor undigested band indicating a mutation efficiency of only 6.6%. {The intensity of the undigested band is 52.800 arbitrary units (au). The sum of the intensities of the digested bands is 1434.911 au. The transfection efficiency is 54%. Therefore, the mutation frequency is $[52.8 / (52.8 + 1434.911)] / 0.54 = 6.6\%$.}. The PCR products amplified by the first set of primers were cloned, and clones with putative mutated regions were sequenced. Five of 40 clones showed mutations (12.5%), which were either deletions (1–13 bp) or a 1-bp substitution (Figure S4a). Although base substitutions are

relatively rare, several studies reported such CRISPR/Cas9-induced mutations (Cermak *et al.*, 2015; Ikeda *et al.*, 2016; Li *et al.*, 2013; Liang *et al.*, 2016; Mikami *et al.*, 2015a,b; Schiml *et al.*, 2014; Wang *et al.*, 2016; Xu *et al.*, 2014; Zhang *et al.*, 2014). To reduce the cloning work and increase the sensitivity of mutant validation, we amplified DNA (using a second set of primers) from the restriction enzyme mixture and digested the amplicons with *BsaI* to detect mutations again. Meanwhile, the second PCR products were cloned into a T/A vector (Figure S4b). After enrichment of the mutated amplified DNA, 71% (5/7) of the clones carried mutations, all of which were deletions (3–8 bp).

Millet

The results in millet were similar to those of bamboo, with a more significant undigested band in protoplast DNA PCR products after 48 h (Figure S5). The scanned electrophoresis image indicated a mutation frequency of 10.2% after 48-h incubation (Figure S5a). Sequencing results of the second PCR product indicated that most of the mutant clones carried deletions before the PAM, with




Figure 2 Schematic showing mutation of target site sequences after genome-editing using a protoplast transfection system. (a) The genomic sequence of the target gene is used to design an sgRNA sequence and primers. Target sites containing a restriction enzyme site are chosen. Blue, Target site; red, protospacer adjacent motif (PAM). The single guide RNA (sgRNA) was cloned into pCambia1300-OsU3-Cas9 (or OsU6). Plasmid DNA is delivered into protoplasts by a PEG-mediated method. (b) Fluorescence image of transfected millet protoplasts. The plasmid mRFP-NLS is used as a marker to determine the transfection efficiency. Cells fluorescing red are successfully transfected. (c) DNA from nontransfected (WT) or transfected (OsU3-24 h or OsU3-48 h) protoplasts are isolated and used as a PCR template for target gene amplification (*PDS* in this case). PCR products with (+) or without (–) *Pst*I digestion. (d) Sanger method sequencing result of one aliquot pool of PCR product. The PCR product pool contains a mixture of wild-type and differentially mutagenized DNA [SNPs (single nucleotide polymorphisms), insertions and deletions]. (e) An aliquot of mutagenized PCR products is subcloned into a T/A cloning vector. The inserts of individual colonies are amplified by colony PCR and the mutation confirmed by digestion with *Pst*I. Clones containing a putative mutagenized target gene (indicated by star marks) are subjected to sequencing. (f) DNA sequencing results of mutated clones. First line, wild type; blue, target site; red, PAM; purple, insertion; –, deleted nucleotides.

one showing an insertion and one showing an insertion/deletion (line 7, the sequence with the grey colour code; Figure S5c).

Rice

The transfection of rice protoplasts with pCambia1300-OsU3 (*Aar*I)-OsPDS resulted in a minor undigested band after DNA amplification 24 h after transfection and a more visible band after 48 h (Figure S6). The scanned gel image indicated a mutation frequency of 7.3%. The second PCR product cloning and sequencing showed mutations at the predicted site (the fourth nucleotide before the PAM). Most mutations were deletions. There were two cases of insertions; one was a 1-bp insertion, and the other contained 175 nucleotides of vector sequence (Figure S6c).

Maize

The maize inositol phosphate kinase (*IPK*) gene was targeted using two sgRNA sequences (Liang *et al.*, 2014). These sgRNAs were cloned individually into pCambia1300-OsU3 (*Aar*I). *Sac*I and *Bam*HI were used to validate mutations effected by gRNA1 and gRNA2, respectively. After 48-h transfection, there were minor undigested bands in both gRNA1 and 2 treatments, indicating that the mutation efficiency was low. When individual gRNAs were used for transfection, gRNA2 was more efficient than was

gRNA1 48 h after transfection (Figure S7, mutation frequency, gRNA1: 0.2%; gRNA2: 1.1%). Sequencing revealed mutations consisting of either deletions or insertions (Figures S7c and d). When both guide RNAs were co-transfected, the first PCR did not reveal any truncated product. However, there was a ~300-bp smaller PCR product after the second PCR (Figure S8a). These results confirmed that the DNA region between sgRNA1 and 2 was deleted (Figure S8b).

CRISPR editing of *Brassicaceae* protoplasts

Arabidopsis

Arabidopsis protoplasts were transfected with pCambia1300-OsU3 (*Aar*I)-AtPDS3. Although the AtPDS3 sgRNA was driven by either a rice OsU3 or OsU6 promoter, CRISPR/Cas9-mediated mutations were detected. Using the construct carrying the OsU3 promoter, DNA extracted from protoplasts amplified with AtPDS3 primers and digested with *Nco*I showed faint undigested bands 24 h after transfection (Figure S9a). The results were similar when the OsU6 promoter was used (Figure S9b). After 48 h, more intense undigested bands were observed. The mutation frequency was 6.5% in the OsU6 treatment after 48-h incubation. The PCR products were cloned and sequencing indicated that mutations (deletions and insertions) initiated at or before the fourth nucleotide before the PAM (Figure S9d).

Table 1 Targeted mutations in protoplasts of nine plant species

					Sequenced mutations (bp)		
	Species	Transfection efficiency (%)	Target gene	Mutagenesis efficiency (%)	Insertions	Deletions	Other
Poaceae	<i>Bambusa oldhamii</i>	54	<i>PDS</i>	6.6	—	1, 3, 4, 6, 8, 11, 13	Replace 1
	<i>Setaria italica</i>	51	<i>PDS</i>	10.2	1	1, 3, 5, 6, 7, 8, 14, 31, 43	—
	<i>Oryza sativa</i>	44	<i>PDS</i>	7.3	1, 175	1, 9, 11	—
	<i>Zea mays</i>	47	<i>IPK</i>	0.2*	1	5	—
				1.1 [†]	1	1	—
Brassicaceae	<i>Arabidopsis thaliana</i>	67	<i>PDS</i>	6.5	1(A), 1(T), 2	−5	—
	<i>Brassica oleracea</i>	43	<i>GA4a</i>	75.2	1 (C), 1 (T)	2, 5, 6, 8, 10, 14	Delete 5, replace 5
	<i>Brassica napus</i>	63	<i>GA4a</i>	56.8	1 (A), 1 (T), 34	2, 5	—
Solanaceae	<i>Nicotiana tabacum</i>	41	<i>PDS</i>	15.0	14, 24	1, 3, 6, 8, 12	—
	<i>Solanum lycopersicum</i>	63	<i>PDS</i>	3.7	29	1, 2, 3, 8, 9	—

*sgRNA1.

†sgRNA2.

Broccoli

For broccoli, we targeted the sgRNA gene *BolGA4.a* at a *HphI* site (Lawrenson *et al.*, 2015). Both rice promoters, U3 and U6, were tested. Despite the use of a monocot promoter, genome editing in *B. oleracea* had a 75.2% mutation efficiency with OsU6 treatment after 48-h incubation (Figure S10). To confirm that the undigested PCR product is mutated by CRISPR/Cas9, the first PCR product was cloned and sequenced. The results indicated that 64.1% (25/39) of the clones were mutated at or before the fourth nucleotide preceding the PAM (Figure S11). These results are similar to those of the gel image. Using these two methods, both mutagenesis efficiency and the mutated sequences could be obtained.

Rapeseed

Because the rapeseed *GA4.a* gene is highly homologous to that of broccoli, the same construct was used for both species. The CRISPR/Cas9 editing efficiency was also high (in OsU3 treatment after 24-h incubation, 56.8%) in rapeseed protoplasts. The first PCR product was cloned and sequenced. Sequencing indicated both deletions and insertions (Figure S12).

CRISPR editing of *Solanaceae* protoplasts

To edit *Solanaceae* crop genomes, we used tobacco (BY-2) and tomato (Micro-Tom) suspension cells. The *PDS* homologs, which harbour a *MlyI* site, were targeted in both species. The OsU3 promoter was used to drive sgRNA expression in both cases. The results are shown in Figure S13 (for tobacco BY-2 cells, the mutation frequency was 15.0% after 48 h) and Figure S14 (for tomato Micro-Tom, the mutation frequency was 3.7% after 48 h).

Detecting CRISPR-mediated mutations in individual protoplasts

Transfection of protoplasts with CRISPR editing reagents usually results in a mixture of genotypes, including nonedited wild-type sequences and various differently edited genomes. Even when the protoplast transfection efficiency is high, frequently only a small percentage of cells contain edited genomes. It is often difficult to detect these rare edited genomes among a high background of nonedited genomes. We thus developed a protocol for analysing DNA from individual mutagenized protoplasts. Because the PCR template is from a single cell, it is easier to validate the electrophoresis results, including complete digestion of the amplicon DNA by restriction endonucleases. Mutations in the PCR product can be confirmed by Sanger sequencing.

We evaluated the parameters of transfection and single-cell analysis by targeting mutations in the tobacco *PDS* gene (*NtPDS*). Tobacco (*N. tabacum*) is an amphidiploid derived from *N. sylvestris* and *N. tomentosiformis* (Endo *et al.*, 2016; Sierro *et al.*, 2014). Specific primers were designed to amplify *PDS* genes from the *N. sylvestris* genome (S form) and the *N. tomentosiformis* genome (T form). We transfected *N. tabacum* protoplasts with various amounts of a mixture of a construct encoding the sgRNA which can target identical sequences in both genomes and the Cas9 protein (Kaya *et al.*, 2016) and isolated more than 20 individual protoplasts for each treatment.

Table 2 shows that with increasing amounts of DNA, a higher percentage of individual protoplasts showed *PDS* mutations. Using 20 µg of DNA, on average 55.3% of the protoplasts showed S form target mutations in three experiments. Because

the transfection efficiency in this experiment (Figure 3) was 43%, these results indicate that the S genome *PDS* target mutation efficiency was 83.7% ($0.36/0.43 = 83.7\%$, Data S1). Although the target sequence was identical, only 29.7% of the T form *PDS* genes showed mutations (Table 2). Similar S/T mutation ratios were seen when protoplasts were incubated for 2–4 days following transfection (Figure 4, Data S2). Triplicate experiments yielded similar results (Figure S15, Data S2).

To apply this single-protoplast protocol to other species, we investigated CRISPR-mediated mutagenesis of the maize *ZmIPK* gene. Poaceae protoplasts are smaller than those of tobacco and *Arabidopsis*, but we could isolate individual maize protoplasts after transfection. Similar to the experiments described above, we used two sgRNA to delete a fragment of *ZmIPK*. Using a protoplast mixture, deletions could not be detected during the first round of PCR analysis, but could be observed using a second round of PCR after a restriction enzyme was used to digest the amplicons resulting from the first PCR product (Figure S8). Because of these two rounds of PCR, it is difficult to calculate target site mutagenesis efficiency using the scanned gel image. However, using a single protoplast, we could detect deletions in 23% of the protoplasts (Figure 5). Because the transfection efficiency was 55%, the calculated mutagenesis efficiency was 41%. PCR products were cloned and sequenced to confirm the deletions. The sequence data are shown in Data S3.

According to these results, single-cell analysis is a convenient method to detect mutagenesis efficiency and to determine mutated sequences. Using pooled protoplast DNA as the template, the mutagenesis efficiency could be measured by gel image analysis. In the *ZmIPK* two sgRNA experiment (Figure 5), single-cell analysis was more sensitive because the low amount of the PCR product containing deletions was not observed in the scanned gel image (Figure S7). To obtain the mutated sequences, the pooled DNA PCR products had to be cloned for sequencing. Using single-cell analysis, the PCR product could be sequenced without cloning. Heterozygous PCR products were identified using bioinformatics tools. The scanned gel image analysis plus PCR product cloning/sequencing, and the single-cell analysis which we present in this report are alternative methods to determine mutagenesis efficiency and mutated sequences. The results from these two methodologies were consistent.

Table 2 Mutation of *NtPDS* genes from single transfected tobacco protoplasts

DNA (µg)	Gene*	0	5	10	20
Experiment 1	S	0	9	23	36
	T	0	9	9	9
Experiment 2	S	0	65	70	85
	T	0	15	40	50
Experiment 3	S	0	10	60	45
	T	0	5	25	30
% Target mutation (±SE)	S	0	28.0 ± 32.0	51.0 ± 24.8	55.3 ± 26.1
	T	0	9.7 ± 5.0	24.7 ± 15.5	29.7 ± 20.5

*S, *N. sylvestris* gene; T, *N. tomentosiformis* gene. Bold-face type indicates the average ± s.d. among the multiple experiments.




Figure 3 Single-protoplast analysis of the effect of plasmid dosage on *NtPDS* target mutagenesis. Different amounts of plasmid DNA (containing the expression cassette of *NtPDS* sgRNA and *SaCas9*; Kaya *et al.*, 2016) were transfected into tobacco mesophyll protoplasts. After three days, the target mutations were analysed by RFLP. S, *N. sylvestris* form; T, *N. tomentosiformis* form. Green box, wild-type RFLP control; red box, albino mutant RFLP control. Numbers above the gel lanes indicate protoplast number.

Regeneration of CRISPR-edited plants from single protoplasts

To show that CRISPR-mediated mutagenesis can be useful to generate genome-edited plants, we targeted the *NtPDS* gene of tobacco. The *PDS* gene is important for chlorophyll biogenesis (Qin *et al.*, 2007), and homozygous *PDS* mutants are albino. Because *N. tabacum* contains four *PDS* genes (two each from the *N. sylvestris* and *N. tomentosiformis* progenitors), mutation of all four genes is necessary to obtain albino plants. Following transfection, protoplasts were transferred to growth medium to obtain calli. We obtained both albino and green calli (Figure 6a), which we regenerated into shoots. Green calli grew more quickly than did albino calli, but we were able to obtain shoots from both types of calli (Figure 6b–d). From three biological repeats, albino calli represented 29.1% of all regenerated calli generated using 20 µg DNA, although the percentage of albino calli was lower using less DNA (Table 3). We analysed DNA of 20 shoots from albino and green calli (Figures 7 and S16, Table 4). As expected, all albino shoots contained targeted mutations in all of the *PDS* genes. DNA sequence analysis of the few amplicons from white calli that were digested by *Bst*NI indicated that this restriction endonuclease site remained, but that other frame-shift mutations occurred (Data S4). From the green tissue, 44.4% of the shoots contained edited S form *PDS* genes, but only 25.6% contained edited T form genes (Figures 7 and S16, Table 4). Sequencing of PCR *NtPDS* gene products from green shoots confirmed the

mutations (Data S4). For all shoots (green and albino) investigated, 58.8% contained at least one mutated *NtPDS* gene.

The percentage of shoots with the albino phenotype was DNA dosage dependent. Ten green shoot clusters for each repetition (three repeats) for each DNA amount were subjected to RFLP and sequencing analysis (Table 3, Figure S17, Data S5). No albino calli were observed without the use of Cas9/sgRNA-encoding DNA, and only ~4% of the shoot clusters obtained using 5 and 10 µg DNA/reaction were albino (Table 3). The reaction using 20 µg DNA/reaction resulted in the highest percentage of albino shoots.

We further investigated whether the regenerated shoots contained the *Cas9* gene. Thirty-one of 180 albino shoots (17.2%) contained *Cas9*. Only four of the 180 green regenerants (2.2%) contained *Cas9*. Thus, we could obtain regenerated shoots from *PDS*-edited plants that lack *Cas9* (Table 4).

Discussion

Improvement of protoplast isolation

CRISPR/Cas9 technology is a powerful tool for plant breeding and research. CRISPR technology is still evolving, including the use of novel Cas9 nucleases, rules for the design of gRNAs and algorithms to predict target and 'off-target' sequences. The performance of these new systems and sgRNA target efficiency still relies on empirical results. However, stable transformation to evaluate a construct is time-consuming and costly, with at least




Figure 4 Single-protoplast analysis of the effect of incubation times on *NtPDS* target mutagenesis. Tobacco protoplasts were transfected with 20 µg plasmid DNA (containing the expression cassette of *NtPDS* sgRNA and *SaCas9*; Kaya *et al.*, 2016) and incubated for various number of days. The target mutation was analysed by RFLP. S, *sylvestris* form; T, *tomentosiformis* form. Green box, wild-type RFLP control; red box, albino mutant RFLP control. Numbers above the gel lanes indicate protoplast number.

one report claiming that in maize >90 candidate target sites must be screened to find a suitable sgRNAs (Zhu *et al.*, 2016). Therefore, a CRISPR platform with a rapid and efficient evaluation protocol is needed. A protoplast transient transfection system with high transfection efficiency fulfils this need. The bottleneck of isolating high-quality protoplasts from different species has hindered this application.

For rice protoplast isolation, we made two modifications to previous methods (Chen *et al.*, 2006; Zhang *et al.*, 2011) to improve yield and reduce costs. First, instead of cutting the seedlings into cross section pieces, we cut the rice seedlings parallel to the veins. Early attempts to use random or cross section cutting resulted in fewer and lower quality protoplasts, likely because there are proportionally more cell walls in the same-sized cross-sectional area. Longitudinal cuts generated more damage to cells along the cut edge but allowed easier access of the maceration enzymes to the cells, resulting in more protoplasts released during digestion (Figure 1d,e). When seedlings were cut in cross section, more cells remained within the leaf sheath after digestion. We substituted the less expensive Cellulase R10 for the more expensive Cellulase RS (Zhang *et al.*, 2011). We could apply this protocol to other Poaceae species, including wheat, bamboo, millet and corn. We believe this convenient method will benefit Poaceae crop research.

The Tape Sandwich method can be used to isolate protoplasts from Brassicaceae species [*Arabidopsis* (Wu *et al.*, 2009), broccoli, *Cleome*, and rapeseed (Figure S1)] and from other family, such as

Euphorbiaceae (Poinsettia, Pitzschke and Persak, 2012). However, this protocol could not be applied to tomato leaves because the epidermal cells could easily be pulled off by tape.

Although tomato mesophyll protoplasts can be isolated by leaf cutting (Niedz *et al.*, 1985), this is often not convenient. Suspension cell lines are an alternative material for tomato protoplast isolation. These lines can be maintained in a controlled environment without variations caused by temperature or light (Nagata *et al.*, 1992). The tobacco suspension cell line BY-2 has been used for several purposes, such as subcellular protein localization, BiFC vector validation (Lee *et al.*, 2008) and CRISPR gene editing (Mercx *et al.*, 2016). In addition to a 'Micro-Tom' tomato cell line, we also established a line from a different variety (CL5915; World Vegetable Center, Tainan, Taiwan) following this protocol. Other advantages of suspension cells include the ease of transgenic material containment and the ability to maintain good manufacturing practices for production of pharmacological proteins or antibodies, which can be secreted to the extracellular medium for easy purification (Mercx *et al.*, 2016).

Validation of CRISPR/Cas9-mediated targeting mutagenesis in plant protoplasts

Bamboo

Bamboo is an economically and ecologically important vegetable and forest plant in Asia (Ma *et al.*, 2015). Bamboo grows quickly and is a sustainable and environmentally friendly crop. Bamboo




Figure 5 Targeted mutagenesis of the *ZmIPK* gene in protoplasts. (a) The gene structure of *ZmIPK*. Blue, gRNA positions; green, two pairs of primers for single-cell PCR analysis; red, size of PCR product from the wild-type gene; orange, size between two gRNAs; purple, theoretical size after precise deletion. (b) Maize protoplasts were transfected with two sgRNAs that targeted different regions of the *ZmIPK* gene. DNA from several single protoplasts was amplified by PCR 72 h after transfection. P, pooled DNA PCR product; –, no plasmid DNA transfected. Numbers above the gel lanes indicate protoplast number.




Figure 6 *NtPDS* mutant plants derived from CRISPR mutagenesis of transfected protoplasts. (a) Growing calli were embedded in shooting medium for 1.5 months. (a*) Magnified view of one portion of the plate shown in (a). Bar = 1 cm. (b) Green shoot clusters were subcultured in shooting medium for 3 weeks. (b*) Magnified view of one portion of the plate shown in (b). Bar = 1 cm. (c) Albino calli were subcultured in shooting medium for 3 weeks. (c*) Magnified view of one portion of the plate shown in (c). Bar = 1 cm. (d) Albino calli were subcultured in shooting medium after 7 weeks. (d*) Magnified view of one portion of the plate shown in (d). Bar = 1 cm.

Table 3 Analysis of regenerated green and albino shoots from CRISPR-mutagenized tobacco protoplasts

Number of green and albino shoots regenerated after transfection with various amounts of Cas9- and sgRNA-encoding plasmid DNA				
DNA (μg)	0	5	10	20
Green shoots (repeat 1)	113	127	62	139
Green shoots (repeat 2)	119	174	148	48
Green shoots (repeat 3)	136	129	167	123
Total green shoots	368	430	377	310
Albino shoots (repeat 1)	0	3	4	54
Albino shoots (repeat 2)	0	11	6	28
Albino shoots (repeat 3)	0	4	2	45
Total albino shoots	0	18	12	127
% albino shoots (±SE)	0	4.0 ± 1.2	3.1 ± 1.5	29.1 ± 6.9
Mutated <i>NtPDS</i> genes in green shoots (%)				
Repeat 1	0	10	20	80
Repeat 2	0	30	10	30
Repeat 3	0	30	30	40
Total mutant genes (±SE)	0	23.3 ± 11.5	20.0 ± 10.0	50.0 ± 26.5
Mutated <i>NtPDS</i> genes in all shoots (%)				
Repeat 1	0	12.1	24.8	85.6
Repeat 2	0	34.1	13.5	55.8
Repeat 3	0	32.1	30.8	56.1
Total mutant genes (±SE)	0	26.1 ± 12.2	23.1 ± 8.8	65.8 ± 17.1

Bold-face type indicates the average±s.d. among the multiple experiments.

protoplasts have been isolated from suspension cells and leaves (Hisamoto and Kobayashi, 2010; Huang *et al.*, 1989, 1990; Yeh *et al.*, 2011). Protoplasts derived from *in vitro* propagated shoots have been transfected for subcellular protein localization studies (Yeh *et al.*, 2011). In this report, the protoplast yield was

increased using longitudinal cutting from *in vitro* material and immature leaves of greenhouse-grown plants. We demonstrated that the bamboo genome can be mutated by the CRISPR/Cas9 system.

Millet

Millet is an important feed and food crop in arid regions (Pan *et al.*, 2015). Millet is a C4-photosynthesis plant. Because the genome size of millet is smaller than that of other grass crops (490 M), it has become a model plant for C4 photosynthesis (Doust *et al.*, 2009). Because transformation of millet is inefficient (~5.5%; Wang *et al.*, 2011), it is difficult to use a traditional transformation strategy to test sgRNAs for genome editing. Protoplasts can be used to screen for suitable sgRNA and Cas9 constructs for use in stable transformation. Xiang *et al.* (2004) developed a millet protoplast isolation protocol. Millet protoplasts have been used for protoplast transfection and for BiFC investigation (Liu *et al.*, 2016). Using protoplast transfection, we demonstrate that the millet genome can be mutated using the CRISPR/Cas9 system.

Rice

Since the first report of CRISPR/Cas9-mediated genome editing in rice, there have been more than 30 articles published about rice genome editing using CRISPR/Cas9 (e.g. Feng *et al.*, 2013; Mao *et al.*, 2013; Miao *et al.*, 2013; Shan *et al.*, 2013; Xie and Yang, 2013). Compared with other crop plants, CRISPR technology in rice is mature. Modifications reported include single sgRNA knockouts and deletion of a large DNA fragment (>200 kb; Zhou *et al.*, 2014). DNA-free genome editing has also been reported (Woo *et al.*, 2015). CRISPR/Cas9 has been used in rice breeding to improve phenotypes to change leaf morphology (Ikeda *et al.*, 2016), grain number, panicle architecture, grain size and plant architecture (Li *et al.*, 2016a). CRISPR/Cas9 has also been used to induce male sterility (Li *et al.*, 2016b), herbicide-tolerance (Sun *et al.*, 2016) and pathogen resistance (Wang *et al.*, 2016). Numerous studies used protoplasts to confirm the editing efficiency of the sgRNA and Cas9 protein (Jiang *et al.*, 2013; Li *et al.*, 2016b; Lowder *et al.*, 2015; Shan *et al.*, 2013, 2014; Wang *et al.*, 2016; Woo *et al.*, 2015; Xie and Yang, 2013; Xie *et al.*, 2015). These articles indicate that a convenient and high transient transfection protoplast protocol is useful to determine the efficiency of designed gRNAs. We improved rice protoplast isolation and showed that these protoplasts can be used for transient transfection. Our convenient rice protoplast isolation




Figure 7 Targeted mutagenesis of *NtPDS* genes from regenerated tobacco protoplasts. Albino (A) and green (G) plantlets from regenerated protoplasts derived from 20 μg plasmid DNA treatment. Twenty plantlets of each type were analysed for *NtPDS* mutagenesis. The target sequences were amplified and validated using RFLP. S, *N. sylvestris* form; T, *N. tomentosiformis* form. Green box, wild-type RFLP control; red box, albino mutant RFLP control.

Table 4 Analysis of regenerated green and albino shoots from CRISPR-mutagenized tobacco protoplasts which were transfected with 20 µg plasmid DNA

	Experiment 1	Experiment 2	Experiment 3	Average
Phenotypes of regenerated shoots				
Albino/Green	0.20 ± 0.07 (169/969)	0.52 ± 0.20 (205/379)	0.37 ± 0.11 (183/503)	0.36
Mutated <i>NtPDS</i> genes in green shoots (%)				
S	33.3 ± 2.9	46.7 ± 10.4	53.3 ± 2.9	44.4
T	15.0 ± 5.0	25.0 ± 10.0	36.7 ± 7.6	25.6
Total mutant genes (±SE)	35.0 ± 0.0	46.7 ± 10.4	53.3 ± 2.9	45.0
Mutated <i>NtPDS</i> genes in white shoots (%)				
S	100.0 ± 0.0	100.0 ± 0.0	100.0 ± 0.0	100
T	100.0 ± 0.0	100.0 ± 0.0	100.0 ± 0.0	100
Total mutant genes (±SE)	100.0 ± 0.0	100.0 ± 0.0	100.0 ± 0.0	100
Mutated <i>NtPDS</i> genes in all shoots (%)				
Total mutant genes (±SE)	45.6 ± 23.0	64.9 ± 6.0	65.9 ± 3.3	58.8
Shoots containing Cas9 gene (%)				
Green	1.7 ± 2.9	3.3 ± 5.8	1.7 ± 2.9	2.2
Albino	10.0 ± 13.2	15.0 ± 8.7	26.7 ± 15.3	17.2

Bold-face type indicates the average±s.d. among the multiple experiments.

protocol will be useful to reduce the time to evaluate the efficiency of nucleases and sgRNAs.

Maize

Protoplasts isolated from some maize genotypes can be regenerated into plants (Rhodes *et al.*, 1988; Sheen, 2001). There are several maize CRISPR/Cas9 studies that used protoplasts (e.g. Feng *et al.*, 2016; Liang *et al.*, 2014; Svitashv *et al.*, 2015; Xing *et al.*, 2014; Zhu *et al.*, 2016). In this report, we targeted the *IPK* gene with RFLP validation in protoplasts. Using two sgRNAs, we could generate a large (300 bp) deletion within this gene.

OsU3 and OsU6 application in dicots

Most current CRISPR/Cas9 studies have focused on developing only monocot or dicot vectors (Jiang *et al.*, 2013). In our study, we used the rice promoters OsU3 and OsU6 to express sgRNAs in a vector containing *Cas9* and *sgRNA* genes. The results indicate that monocot promoters can be used efficiently in both monocot and dicot protoplasts.

Broccoli

Broccoli plants can be regenerated from mesophyll-derived protoplasts (Robertson and Earle, 1986). These protoplasts can be transfected using a PEG-mediated method (Nugent *et al.*, 2006). We used the Tape Sandwich method to establish a broccoli protoplast transfection system using cotyledons as starting material. Lawrenson *et al.* (2015) established a broccoli CRISPR/Cas9 mutagenesis protocol. We used their sgRNA sequence in this study. In a broccoli *GA4a* CRISPR/Cas9 stable transformation study, 80% mutation frequency was achieved using the *Arabidopsis* U6 promoter to drive the same sgRNA (Lawrenson *et al.*, 2015). We conclude that our high mutagenesis results from the chosen target gene and the high efficiency of protoplast isolation and transfection. The sgRNA mutation efficiency has been positively correlated with GC content, and the *GA4a* gene has a high GC ratio (65%; Ren *et al.*, 2014). This speculation has to be proven by further sgRNA investigations.

Rapeseed

Rapeseed is cultivated mainly for its oil-rich seed, the third largest source of vegetable oil in the world (<http://apps.fas.usda.gov/psd/online/circulars/oilseeds.pdf>). Kartha *et al.* (1974) described a rapeseed protoplast isolation protocol, and plantlets could be regenerated from these protoplasts. A rapeseed protoplast transient transfection system has also been established (Wang *et al.*, 2015a). Our results indicate that the rapeseed protoplast genome can be mutated by CRISPR/Cas9.

Solanaceae

Tobacco is one of the first crops in which CRISPR/Cas9 genome editing was demonstrated (Jiang *et al.*, 2013; Li *et al.*, 2013; Nekrasov *et al.*, 2013). There are many articles demonstrating that the tomato genome can be edited by CRISPR/Cas9 (Brooks *et al.*, 2014; Cermak *et al.*, 2015). In previous studies, the sgRNA was driven by an *Arabidopsis* U6 promoter. We showed that the same vectors used in monocots (pCambia1300-OsU3(Aar1)-Cas9 vectors) can be used in Solanaceae (tobacco and tomato), even though the promoters are derived from rice.

Application of single-cell analysis to genome editing

Numerous studies have demonstrated the utility of protoplasts to evaluate the efficiency of various CRISPR systems to mutate specific target genes (Feng *et al.*, 2013; Li *et al.*, 2013; Shan *et al.*, 2013; Xie and Yang, 2013). Populations of protoplasts are transfected with DNA encoding Cas9 (or other editing proteins) and genes encoding sgRNAs. Alternatively, Cas9 protein/sgRNA complexes can be transfected into protoplasts (Woo *et al.*, 2015). Total DNA from the protoplast population is isolated and assayed for mutagenesis frequency using RFLP or other methods of analysis, such as the T7E1 surveyor assay (Woo *et al.*, 2015). When the mutation frequency is high, it is relatively easy to detect mutated molecules after electrophoresis through agarose gels. However, as we have shown above with the *IPK* gene of maize, the frequency of CRISPR-mediated mutagenesis can be low,

requiring a second round of PCR to amplify sequences from the small amount of nondigested DNA extracted from the gel.

Single-cell (protoplast) analysis is a sensitive and convenient method to evaluate the efficiency of various sgRNAs for CRISPR-mediated mutagenesis (Figure S18). Because only two copies of a given gene exist in a diploid cell (or four copies in an amphidiploid such as *N. tabacum*), edited copies of the gene will represent 50% of the gene molecules (for heterozygous cells) or 100% of the gene molecules (for homozygous or biallelic cells). PCR products of these genes can directly be sequenced without the need for prior cloning. Because of the small amount of DNA in a single protoplast, a gene of interest cannot be amplified by one step PCR directly, even if the protocol is increased to 50 cycles. Target sequences can be amplified in a second round of PCR (use 1 µL of the first step PCR product as a template). It is important to use nested primers for the second round of PCR. If the same primers are used for both PCR steps, only a smear of products is obtained. In a previous protocol, transfected protoplasts had to be cultured in W5 solution (Wu et al., 2009), then transferred to liquid callus medium for further regeneration. Our new single-protoplast analysis protocol allows faster analysis using much less plant material.

Homozygous transgene-free mutant plants can be obtained from protoplasts in the first generation

One of the advantages of using protoplasts is that DNA-free genome editing can be conducted using preassembled RNP complexes consisting of Cas9 protein and sgRNAs. These complexes were introduced directly into protoplasts using a polyethylene glycol-based method (Woo et al., 2015) to generate mutagenized lettuce. RNP complexes can also be delivered to calli by particle bombardment (Liang et al., 2017). Generation of homozygous or biallelic mutants lacking transgenes encoding genome-editing reagents in the first generation is important for those plant species that cannot easily be crossed to eliminate transgenes. These include vegetatively propagated species, such as potato and banana, and species with long generation times, such as woody trees.

In this report, we show that we can generate transgene-free genome-edited plants by transfecting protoplasts with DNA encoding sgRNAs and Cas9 protein (Figure S19). Most regenerated *N. tabacum* plants containing edited *PDS* genes lacked the Cas9-encoding transgene DNA (Table 4). Homozygous transgene-free mutant plants could be regenerated in the T1 generation. Similar mutagenesis results were reported for wheat (Zhang et al., 2016) and lettuce (Woo et al., 2015). Compared to RNP complexes, DNA encoding the genome-editing reagents is less expensive and more convenient to use. One advantage of RNP complexes is their higher target mutation efficiency (Woo et al., 2015). If the efficiency of a given sgRNA were high, we recommend using DNA encoding genome-editing reagents. As we have shown, protoplast transfection can be highly efficient and up to 60% of mutagenized regenerated tobacco plants contained at least one mutated *PDS* gene (Tables 3 and 4). In hexaploid wheat, only 0.6% (67/10448) of the plants transformed by CRISPR constructs were homozygous mutants (Zhang et al., 2016). We have shown that for allotetraploid tobacco, ~30% of the plants regenerated from mutagenized protoplasts were homozygous/tetra-allelic mutants. In theory, we could obtain 154 transgene-free albino mutants from one transfection

$\{[(169 + 205 + 183)/3] \times (1 - 0.172)\}$, a number much higher than can be obtained by particle bombardment (Liang et al., 2017; Zhang et al., 2016). Thus, CRISPR-based mutagenesis of protoplasts can be a useful technology for polyploid crops, especially horticultural crops that are propagated via asexual methods. Our studies suggest that this new technology can reduce breeding time significantly.

Protoplast regeneration remains a bottleneck for this platform. Especially for monocot species, high regeneration efficiency protocols are not available (Liang et al., 2017; Zhang et al., 2016). Improvements in protoplast regeneration protocols will be important for genome editing and the delivery of high editing efficiency reagents (RNP and donor DNA) to obtain transgene-free crops.

Materials and methods

Plant materials

Millet (*Setaria italica*), corn (*Zea mays*), rice (*Oryza sativa*), broccoli (*Brassica oleracea*), and rapeseed (*B. napus*) plants were grown on filter paper with drip irrigation in an environment-controlled chamber with a long photoperiod (16-h light/8-h dark) at 25 °C. Two-week-old plants were used for protoplast isolation. Bamboo (*Bambusa oldhamii*) shoots were incubated in Murashige and Skoog (MS) salts medium (Sigma-Aldrich, St. Louis, MO) supplemented with 0.1 mg/L TDZ (Lin et al., 2007). *Arabidopsis* seeds were grown in soil using a 16-h light/8-h dark cycle at 26 °C. *Cleome spinosa*, *C. monophylla*, and *C. gynandra* plants were grown in a greenhouse. Tobacco BY-2 cells were grown in MS salts supplemented with 1 mg/L thiamine-HCl, 370 mg/L KH_2PO_4 , 30 g/L sucrose, and 2 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D), pH 5.7 (Lee et al., 2008). A tomato Micro-Tom cell line was subcultured in MS medium supplemented with 1 mg/L 2,4-D, pH 5.7. Detailed information on this cell line is presented in Figure 1.

Protoplast isolation and PEG-mediated protoplast transfection

Protoplast isolation and transfection were performed following Wu et al. (2009) for Brassicaceae, Zhang et al. (2011) for Poaceae, and Lee et al. (2008) for Solanaceae, with modifications. Details are described in Table S1 for Poaceae, Table S2 for Brassicaceae, and Table S3 for Solanaceae. Following transfection, 1.5 mL W5 solution were added, the protoplasts were transferred into six-well plates (precoated with 1% BSA solution), and cultured in the dark at room temperature for 48 h.

Plasmid constructions

The primers OsU3-*HindIII*-f and OsU3-*HindIII*-r were used to amplify the gRNA cassette (OsU3 promoter, *AarI* restriction site and gRNA scaffold) from the plasmid pU3-gRNA which was ligated into *HindIII* digested pCAMBIA1300 to generate pCAMBIA1300-OsU3(*AarI*) (Figure S2). The Cas9 cassette fragment [35S promoter, human codon-optimized *SpCas9* (*hSpCas9*) gene and terminator] was obtained from 35S-Cas9-SK by digestion with *SaI* and *EcoRI* and ligated into *SaI/EcoRI* digested pCAMBIA1300-OsU3(*AarI*) to generate pCAMBIA1300-OsU3(*AarI*)-Cas9 (Accession no.: KX400856). The same strategy was used to generate pCAMBIA1300-OsU6(*AarI*)-Cas9 (Accession no.: KX388151). Detailed information is found in Supplemental Material and Methods.

Mutation validation in protoplasts

Genomic DNA was extracted from pooled protoplasts using a Mini GenoPlus Genomic DNA Extraction Kit (Viogene, New Taipei City, Taiwan). To amplify the genomic region targeted by the sgRNA, two pairs of primers were designed. The restriction enzyme and primer sequence information are shown in the Supplemental Figures. PCR conditions are 94 °C for 5 min, 35 cycles of 94 °C for 30 s, annealing (55–63 °C; detailed information is shown in the Supplemental Figures) for 30 s, polymerization at 72 °C for 30 s, followed by 72 °C for 3 min. The PCR product was digested by the appropriate restriction enzyme and the products subjected to electrophoresis. Electrophoresis gel images were analysed by ImageJ (Schneider *et al.*, 2012). The intensity of the undigested band was divided by the total intensity of all bands to generate the raw mutation frequency. The raw mutation frequency was divided by the transfection efficiency to obtain the final mutation frequency. Secondary PCR products were cloned into the T&A™ vector (FYC002-20P; Yeastern Biotech Co. LTD, New Taipei City, Taiwan). Colonies harbouring the edited DNA were screened by a PCR/restriction enzyme assay, and the insert DNA was sequenced. VirD2-NLS-mRFP plasmid DNA (Lee *et al.*, 2008) was transfected into protoplasts by PEG-mediated methods as a positive control to determine transfection efficiency.

Single-protoplast isolation and target mutagenesis validation

Transfected protoplasts were centrifuged (300× *g*, 3 min., 25 °C, Eppendorf Centrifuge 5804R; Eppendorf, Hamburg, Germany) and transferred to liquid callus medium [1/2 MS medium supplemented with 0.4 M mannitol, 30 g/L sucrose, 1 mg/L 1-naphthaleneacetic acid (NAA) and 0.3 mg/L kinetin]. Protoplast concentration was measured using a hemocytometer and adjusted to 1 cell/μL. A single protoplast was isolated using a pipette (SelectPette, Genomics, New Taipei City, Taiwan), transferred to a slide, and the condition checked by microscopy (Olympus, Tokyo, Japan). A single protoplast was transferred to 20 μL PCR mixture containing the first pair of primers for the initial DNA amplification. One microliter of the PCR product was used as a template for a second round of PCR using a second pair of primers. Amplicons resulting from the second PCR were digested by the appropriate restriction endonuclease and analysed by electrophoresis to validate mutation of the target sequence. A schematic for single-cell analysis is shown in Figure S18.

Tobacco protoplast regeneration

We used a protocol modified from Liao (1990, Figure S19). Transfected protoplasts were incubated in a 5-cm-diameter Petri dish containing liquid callus medium. After 2- to 3-week incubation in the dark, the protoplasts proliferated and formed dust-like calli. The calli were embedded in solidified callus medium in a 9-cm-diameter petri dish and incubated at 25 °C for 3–4 weeks in the dark. Calli larger than 3 mm were embedded in shoot medium for shoot induction. After 1 month at 25 °C in the light (light/dark: 16/8 h; 3000 lux), shoot clusters containing leaves >5 mm were transferred to fresh shoot medium for 2–3 weeks for further shoot proliferation. Shoot clusters with leaves were then transferred to solidified root medium.

Acknowledgements

We thank Caixia Gao of the Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, China, Jiankang Zhu of the Shanghai Center for Plant Stress Biology, Chinese Academy of Sciences, China, Seiichi Toki and Masaki Endo, National Institute of Agrobiological Sciences, Japan, for providing the vectors containing the gRNA cassette and the Cas9 cassette, respectively, used in this work.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by funds from the Agricultural Biotechnology Research Center, Academia Sinica of Taiwan (to CSL and MCS).

Author contributions

CSL, LYL, SBG, and MCS conceived and designed the experiments. YZ, RZ, WW, and WJC constructed vectors. CSL, LHY, CTH, and FHW performed targeted mutagenesis and protoplast-related experiments. CTH, CSL, and LJL conducted protoplast regeneration. CTH, QWC, FJY, CTY, LHY, and FHW analysed the data. CSL, LYL, LHY, CTH, SBG, and HCWH interpreted the data. CSL, LYL, SBG, and MCS wrote the manuscript.

References

- Ali, Z., Abulfaraj, A., Idris, A., Ali, S., Tashkandi, M. and Mahfouz, M.M. (2015) CRISPR/Cas9-mediated viral interference in plants. *Genome Biol.* **16**, 238.
- Ali, Z., Ali, S., Tashkandi, M., Zaidi, S.S. and Mahfouz, M.M. (2016) CRISPR/Cas9-mediated immunity to geminiviruses: differential interference and evasion. *Sci. Rep.* **6**, 26912.
- Belhaj, K., Chaparro-Garcia, A., Kamoun, S. and Nekrasov, V. (2013) Plant genome editing made easy: targeted mutagenesis in model and crop plants using the CRISPR/Cas system. *Plant Methods*, **9**, 39.
- Bortesi, L. and Fischer, R. (2015) The CRISPR/Cas9 system for plant genome editing and beyond. *Biotechnol. Adv.* **33**, 41–52.
- Brennecke, P., Anders, S., Kim, J.K., Kolodziejczyk, A.A., Zhang, X., Proserpio, V., Baying, B. *et al.* (2013) Accounting for technical noise in single-cell RNA-seq experiments. *Nat. Methods*, **10**, 1093–1095.
- Brooks, C., Nekrasov, V., Lippman, Z.B. and Van Eck, J. (2014) Efficient gene editing in tomato in the first generation using the clustered regularly interspaced short palindromic repeats/CRISPR-associated9 system. *Plant Physiol.* **166**, 1292–1297.
- Butt, H., Eid, A., Ali, Z., Atia, M.A.M., Mokhtar, M.M., Hassan, N., Lee, C.M. *et al.* (2017) Efficient CRISPR/Cas9-mediated genome editing using a chimeric single-guide RNA molecule. *Front. Plant Sci.* **8**, 1441.
- Cermak, T., Baltes, N.J., Cegan, R., Zhang, Y. and Voytas, D.F. (2015) High-frequency, precise modification of the tomato genome. *Genome Biol.* **16**, 232.
- Cermak, T., Curtin, S.J., Gil-Humanes, J., Cegan, R., Kono, T.J.Y., Konecna, E., Belanto, J.J. *et al.* (2017) A multipurpose toolkit to enable advanced genome engineering in plants. *Plant Cell*, **29**, 1196–1217.
- Chen, S., Tao, L., Zeng, L., Vega-Sanchez, M.E., Umemura, K. and Wang, G.L. (2006) A highly efficient transient protoplast system for analyzing defense gene expression and protein-protein interactions in rice. *Mol. Plant Pathol.* **7**, 417–427.
- Doust, A.N., Kellogg, E.A., Devos, K.M. and Bennetzen, J.L. (2009) Foxtail millet: a sequence-driven grass model system. *Plant Physiol.* **149**, 137–141.

- Efroni, I. and Birnbaum, K.D. (2016) The potential of single-cell profiling in plants. *Genome Biol.* **17**, 65.
- Efroni, I., Ip, P.L., Nawy, T., Mello, A. and Birnbaum, K.D. (2015) Quantification of cell identity from single-cell gene expression profiles. *Genome Biol.* **16**, 9.
- Eid, A., Ali, Z. and Mahfouz, M.M. (2016) High efficiency of targeted mutagenesis in arabidopsis via meiotic promoter-driven expression of Cas9 endonuclease. *Plant Cell Rep.* **35**, 1555–1558.
- Endo, A., Masafumi, M., Kaya, H. and Toki, S. (2016) Efficient targeted mutagenesis of rice and tobacco genomes using Cpf1 from *Francisella novicida*. *Sci. Rep.* **6**, 38169.
- Feng, Z.Y., Zhang, B.T., Ding, W.N., Liu, X.D., Yang, D.L., Wei, P.L., Cao, F.Q. et al. (2013) Efficient genome editing in plants using a CRISPR/Cas system. *Cell Res.* **23**, 1229–1232.
- Feng, C., Yuan, J., Wang, R., Liu, Y., Birchler, J.A. and Han, F. (2016) Efficient targeted genome modification in maize using CRISPR/Cas9 system. *J. Genet. Genomics*, **43**, 37–43.
- Gaj, T., Gersbach, C.A. and Barbas, C.F. (2013) ZFN, TALEN, and CRISPR/Cas-based methods for genome engineering. *Trends Biotechnol.* **31**, 397–405.
- Gierahn, T.M., Wadsworth, 2nd, M.H., Hughes, T.K., Bryson, B.D., Butler, A., Satija, R., Fortune, S. et al. (2017) Seq-Well: portable, low-cost RNA sequencing of single cells at high throughput. *Nat. Methods*, **14**, 395–398.
- Hisamoto, Y. and Kobayashi, M. (2010) Protoplast isolation from bamboo leaves. *Plant Biotechnol.* **27**, 353–373.
- Huang, L.C., Chen, W.L. and Huang, B.L. (1989) Tissue-culture investigations of bamboo. 3. A method for viable protoplast isolation from *Bambusa* cells of liquid suspension-culture. *Bot. Bull. Acad. Sinica*, **30**, 49–57.
- Huang, L.C., Huang, B.L. and Chen, W.L. (1990) Tissue-culture investigations of Bamboo. 5. Recovery of callus from protoplasts of suspension-cultured *Bambusa* cells. *Bot. Bull. Acad. Sinica*, **31**, 29–34.
- Huang, S., Weigel, D., Beachy, R.N. and Li, J. (2016) A proposed regulatory framework for genome-edited crops. *Nat. Genet.* **48**, 109–111.
- Ikedo, T., Tanaka, W., Mikami, M., Endo, M. and Hirano, H.Y. (2016) Generation of artificial drooping leaf mutants by CRISPR-Cas9 technology in rice. *Genes Genet. Syst.* **90**, 231–235.
- Jiang, W.Z., Zhou, H.B., Bi, H.H., Fromm, M., Yang, B. and Weeks, D.P. (2013) Demonstration of CRISPR/Cas9/sgRNA-mediated targeted gene modification in *Arabidopsis*, tobacco, sorghum and rice. *Nucleic Acids Res.* **41**, e188.
- Kartha, K.K., Michayluk, M.R., Kao, K.N., Gamborg, O.L. and Constabel, F. (1974) Callus formation and plant regeneration from mesophyll protoplasts of rape plants (*Brassica-napus* L cv Zephyr). *Plant Sci. Lett.* **3**, 265–271.
- Kaya, H., Mikami, M., Endo, A., Endo, M. and Toki, S. (2016) Highly specific targeted mutagenesis in plants using *Staphylococcus aureus* Cas9. *Sci. Rep.* **6**, 26871.
- Kim, H., Kim, S.T., Ryu, J., Kang, B.C., Kim, J.S. and Kim, S.G. (2017) CRISPR/Cpf1-mediated DNA-free plant genome editing. *Nat. Commun.* **8**, 14406.
- Klein, A.M., Mazutis, L., Akartuna, I., Tallapragada, N., Veres, A., Li, V. et al. (2015) Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells. *Cell*, **161**, 1187–1201.
- Lawrenson, T., Shorinola, O., Stacey, N., Li, C.D., Ostergaard, L., Patron, N., Uauy, C. et al. (2015) Induction of targeted, heritable mutations in barley and *Brassica oleracea* using RNA-guided Cas9 nuclease. *Genome Biol.* **16**, 258.
- Lee, L.Y., Fang, M.J., Kuang, L.Y. and Gelvin, S.B. (2008) Vectors for multi-color bimolecular fluorescence complementation to investigate protein–protein interactions in living plant cells. *Plant Methods*, **4**, 24.
- Li, J.F., Norville, J.E., Aach, J., McCormack, M., Zhang, D.D., Bush, J., Church, G.M. et al. (2013) Multiplex and homologous recombination-mediated genome editing in *Arabidopsis* and *Nicotiana benthamiana* using guide RNA and Cas9. *Nat. Biotechnol.* **31**, 688–691.
- Li, M., Li, X., Zhou, Z., Wu, P., Fang, M., Pan, X., Lin, Q. et al. (2016a) Reassessment of the four yield-related genes *Gn1a*, *DEP1*, *GS3*, and *IPA1* in rice using a CRISPR/Cas9 system. *Front. Plant Sci.* **7**, 377.
- Li, Q.L., Zhang, D.B., Chen, M.J., Liang, W.Q., Wei, J.J., Qi, Y.P. and Yuan, Z. (2016b) Development of japonica photo-sensitive genic male sterile rice lines by editing carbon starved anther using CRISPR/Cas9. *J. Genet. Genomics*, **43**, 415–419.
- Liang, Z., Zhang, K., Chen, K.L. and Gao, C.X. (2014) Targeted mutagenesis in *Zea mays* using TALENs and the CRISPR/Cas system. *J. Genet. Genomics*, **41**, 63–68.
- Liang, G., Zhang, H., Lou, D. and Yu, D. (2016) Selection of highly efficient sgRNAs for CRISPR/Cas9-based plant genome editing. *Sci. Rep.* **6**, 21451.
- Liang, Z., Chen, K., Li, T., Zhang, Y., Wang, Y., Zhao, Q., Liu, J. et al. (2017) Efficient DNA-free genome editing of bread wheat using CRISPR/Cas9 ribonucleoprotein complexes. *Nat. Commun.* **8**, 14261.
- Liao, L.J. (1990) *Studies on the characteristics of interspecific somatic hybrids between the N. sylvestris and N. tomentosiformis*. PhD Dissertation, National Taiwan University, Taipei, Taiwan.
- Lin, C.S., Kalpana, K., Chang, W.C. and Lin, N.S. (2007) Improving multiple shoot proliferation in bamboo mosaic virus-free *Bambusa oldhamii* Munro propagation by liquid culture. *HortScience*, **42**, 1243–1246.
- Liu, X.T., Tang, S., Jia, G.Q., Schnable, J.C., Su, H.X., Tang, C.J., Zhi, H. et al. (2016) The C-terminal motif of SiAGO1b is required for the regulation of growth, development and stress responses in foxtail millet (*Setaria italica* (L.) P. Beauv.). *J. Exp. Bot.* **67**, 3237–3249.
- Lowder, L.G., Zhang, D., Baites, N.J., Paul, 3rd, J.W., Tang, X., Zheng, X., Voytas, D.F. et al. (2015) A CRISPR/Cas9 toolbox for multiplexed plant genome editing and transcriptional regulation. *Plant Physiol.* **169**, 971–985.
- Ma, P.F., Zhang, Y.X., Guo, Z.H. and Li, D.Z. (2015) Evidence for horizontal transfer of mitochondrial DNA to the plastid genome in a bamboo genus. *Sci. Rep.* **5**, 11608.
- Macosko, E.Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M. et al. (2015) Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. *Cell*, **161**, 1202–1214.
- Mahfouz, M.M. (2017) Genome editing: the efficient tool CRISPR-Cpf1. *Nat. Plants*, **3**, 17028.
- Mao, Y., Zhang, H., Xu, N., Zhang, B., Gou, F. and Zhu, J.K. (2013) Application of the CRISPR-Cas system for efficient genome engineering in plants. *Mol. Plant*, **6**, 2008–2011.
- Mercx, S., Tollet, J., Magy, B., Navarre, C. and Boutry, M. (2016) Gene inactivation by CRISPR-Cas9 in *Nicotiana tabacum* BY-2 suspension cells. *Front. Plant Sci.* **7**, 40.
- Miao, J., Guo, D.S., Zhang, J.Z., Huang, Q.P., Qin, G.J., Zhang, X., Wan, J.M. et al. (2013) Targeted mutagenesis in rice using CRISPR-Cas system. *Cell Res.* **23**, 1233–1236.
- Mikami, M., Toki, S. and Endo, M. (2015a) Comparison of CRISPR/Cas9 expression constructs for efficient targeted mutagenesis in rice. *Plant Mol. Biol.* **88**, 561–572.
- Mikami, M., Toki, S. and Endo, M. (2015b) Parameters affecting frequency of CRISPR/Cas9 mediated targeted mutagenesis in rice. *Plant Cell Rep.* **34**, 1807–1815.
- Murovec, J., Pirc, Z. and Yang, B. (2017) New variants of CRISPR RNA-guided genome editing enzymes. *Plant Biotechnol. J.* **15**, 917–926.
- Nagata, T., Nemoto, Y. and Hasezawa, S. (1992) Tobacco BY-2 cell-line as the HeLa-cell in the cell biology of higher-plants. *Int. Rev. Cytol.* **132**, 1–30.
- Nekrasov, V., Staskawicz, B., Weigel, D., Jones, J.D.G. and Kamoun, S. (2013) Targeted mutagenesis in the model plant *Nicotiana benthamiana* using Cas9 RNA-guided endonuclease. *Nat. Biotechnol.* **31**, 691–693.
- Niedz, R.P., Rutter, S.M., Handley, L.W. and Sink, K.C. (1985) Plant-regeneration from leaf protoplasts of six tomato cultivars. *Plant Sci.* **39**, 199–204.
- Nugent, G.D., Coyne, S., Nguyen, T.T., Kavanagh, T.A. and Dix, P.J. (2006) Nuclear and plastid transformation of *Brassica oleracea* var. *botrytis* (cauliflower) using PEG-mediated uptake of DNA into protoplasts. *Plant Sci.* **170**, 135–142.
- Pan, Y.L., Ma, X., Liang, H.W., Zhao, Q., Zhu, D.Y. and Yu, J.J. (2015) Spatial and temporal activity of the foxtail millet (*Setaria italica*) seed-specific promoter pF128. *Planta*, **241**, 57–67.
- Pitzschke, A. and Persak, H. (2012) Poinsettia protoplasts—a simple, robust and efficient system for transient gene expression studies. *Plant Methods*, **8**, 14.
- Qin, G., Gu, H., Ma, L., Peng, Y., Deng, X.W., Chen, Z. and Qu, L.J. (2007) Disruption of phytoene desaturase gene results in albino and dwarf phenotypes in *Arabidopsis* by impairing chlorophyll, carotenoid, and gibberellin biosynthesis. *Cell Res.* **17**, 471–482.
- Ren, X., Yang, Z., Xu, J., Sun, J., Mao, D., Hu, Y., Yang, S.J. et al. (2014) Enhanced specificity and efficiency of the CRISPR/Cas9 system with optimized sgRNA parameters in *Drosophila*. *Cell Rep.* **9**, 1151–1162.
- Rhodes, C.A., Pierce, D.A., Mettler, I.J., Mascarenhas, D. and Detmer, J.J. (1988) Genetically transformed maize plants from protoplasts. *Science*, **240**, 204–207.

- Robertson, D. and Earle, E.D. (1986) Plant regeneration from leaf protoplasts of *Brassica oleracea* var. *italica* CV Green Comet broccoli. *Plant Cell Rep.* **5**, 61–64.
- Schimpl, S., Fauser, F. and Puchta, H. (2014) The CRISPR/Cas system can be used as nuclease for in planta gene targeting and as paired nickases for directed mutagenesis in *Arabidopsis* resulting in heritable progeny. *Plant J.* **80**, 1139–1150.
- Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods*, **9**, 671–675.
- Shan, Q.W., Wang, Y.P., Li, J., Zhang, Y., Chen, K.L., Liang, Z., Zhang, K. et al. (2013) Targeted genome modification of crop plants using a CRISPR-Cas system. *Nat. Biotechnol.* **31**, 686–688.
- Shan, Q.W., Wang, Y.P., Li, J. and Gao, C.X. (2014) Genome editing in rice and wheat using the CRISPR/Cas system. *Nat. Protoc.* **9**, 2395–2410.
- Sheen, J. (2001) Signal transduction in maize and *Arabidopsis* mesophyll protoplasts. *Plant Physiol.* **127**, 1466–1475.
- Shimatani, Z., Kashojiya, S., Takayama, M., Terada, R., Arazoe, T., Ishii, H., Teramura, H. et al. (2017) Targeted base editing in rice and tomato using a CRISPR-Cas9 cytidine deaminase fusion. *Nat. Biotechnol.* **35**, 441–443.
- Sierro, N., Battey, J.N., Ouadi, S., Bakaher, N., Bovet, L., Willig, A., Goepfert, S. et al. (2014) The tobacco genome sequence and its comparison with those of tomato and potato. *Nat. Commun.* **5**, 3833.
- Sun, Y.W., Zhang, X., Wu, C.Y., He, Y.B., Ma, Y.Z., Hou, H., Guo, X.P. et al. (2016) Engineering herbicide-resistant rice plants through CRISPR/Cas9-mediated homologous recombination of acetolactate synthase. *Mol. Plant*, **9**, 628–631.
- Svitashev, S., Young, J.K., Schwartz, C., Gao, H.R., Falco, S.C. and Cigan, A.M. (2015) Targeted mutagenesis, precise gene editing, and site-specific gene insertion in maize using Cas9 and guide RNA. *Plant Physiol.* **169**, 931–945.
- Wang, M.Z., Pan, Y.L., Li, C., Liu, C., Zhao, Q., Ao, G.M. and Yu, J.J. (2011) Culturing of immature inflorescences and *Agrobacterium*-mediated transformation of foxtail millet (*Setaria italica*). *Afr. J. Biotechnol.* **10**, 16466–16479.
- Wang, B., Guo, X., Wang, C., Ma, J., Niu, F., Zhang, H., Yang, B. et al. (2015a) Identification and characterization of plant-specific NAC gene family in canola (*Brassica napus* L.) reveal novel members involved in cell death. *Plant Mol. Biol.* **87**, 395–411.
- Wang, Z.P., Xing, H.L., Dong, L., Zhang, H.Y., Han, C.Y., Wang, X.C. and Chen, Q.J. (2015b) Egg cell-specific promoter-controlled CRISPR/Cas9 efficiently generates homozygous mutants for multiple target genes in *Arabidopsis* in a single generation. *Genome Biol.* **16**, 144.
- Wang, F., Wang, C., Liu, P., Lei, C., Hao, W., Gao, Y., Liu, Y.G. et al. (2016) Enhanced rice blast resistance by CRISPR/Cas9-targeted mutagenesis of the ERF transcription factor gene *OsERF922*. *PLoS ONE*, **11**, e0154027.
- Wang, M., Lu, Y., Botella, J.R., Mao, Y., Hua, K. and Zhu, J.K. (2017) Gene targeting by homology-directed repair in rice using a Geminivirus-based CRISPR/Cas9 system. *Mol. Plant*, **10**, 1007–1010. <https://doi.org/10.1016/j.molp.2017.03.002>.
- Woo, J.W., Kim, J., Kwon, S.I., Corvalan, C., Cho, S.W., Kim, H., Kim, S.G. et al. (2015) DNA-free genome editing in plants with preassembled CRISPR-Cas9 ribonucleoproteins. *Nat. Biotechnol.* **33**, 1162–1164.
- Wu, F.H., Shen, S.C., Lee, L.Y., Lee, S.H., Chan, M.T. and Lin, C.S. (2009) Tape-*Arabidopsis* Sandwich—a simpler *Arabidopsis* protoplast isolation method. *Plant Methods*, **5**, 16.
- Xiang, F.N., Xia, G.M., Zhi, D.Y., Wang, J., Nie, H. and Chen, H.M. (2004) Regeneration of somatic hybrids in relation to the nuclear and cytoplasmic genomes of wheat and *Setaria italica*. *Genome*, **47**, 680–688.
- Xie, K. and Yang, Y. (2013) RNA-guided genome editing in plants using a CRISPR-Cas system. *Mol. Plant*, **6**, 1975–1983.
- Xie, K., Minkenberg, B. and Yang, Y. (2015) Boosting CRISPR/Cas9 multiplex editing capability with the endogenous tRNA-processing system. *Proc. Natl Acad. Sci. USA*, **112**, 3570–3575.
- Xing, H.L., Dong, L., Wang, Z.P., Zhang, H.Y., Han, C.Y., Liu, B., Wang, X.C. et al. (2014) A CRISPR/Cas9 toolkit for multiplex genome editing in plants. *BMC Plant Biol.* **14**, 327.
- Xu, R., Li, H., Qin, R., Wang, L., Li, L., Wei, P. and Yang, J. (2014) Gene targeting using the *Agrobacterium tumefaciens*-mediated CRISPR-Cas system in rice. *Rice (N Y)*, **7**, 5.
- Xu, R., Qin, R., Li, H., Li, D., Li, L., Wei, P. and Yang, J. (2017) Generation of targeted mutant rice using a CRISPR-Cpf1 system. *Plant Biotechnol. J.* **15**, 713–717.
- Yamamoto, K., Takahashi, K., Mizuno, H., Anegawa, A., Ishizaki, K., Fukaki, H., Ohnishi, M. et al. (2016) Cell-specific localization of alkaloids in *Catharanthus roseus* stem tissue measured with Imaging MS and Single-cell MS. *Proc. Natl Acad. Sci. USA*, **113**, 3891–3896.
- Yan, L., Wei, S., Wu, Y., Hu, R., Li, H., Yang, W. and Xie, Q. (2015) High-efficiency genome editing in *Arabidopsis* using YAO promoter-driven CRISPR/Cas9 system. *Mol. Plant*, **8**, 1820–1823.
- Yeh, S.H., Lin, C.S., Wu, F.H. and Wang, A.Y. (2011) Analysis of the expression of *BohL1*, which encodes an LSD1-like zinc finger protein in *Bambusa oldhamii*. *Planta*, **234**, 1179–1189.
- Zhang, Y., Su, J., Duan, S., Ao, Y., Dai, J., Liu, J., Wang, P. et al. (2011) A highly efficient rice green tissue protoplast system for transient gene expression and studying light/chloroplast-related processes. *Plant Methods*, **7**, 30.
- Zhang, H., Zhang, J., Wei, P., Zhang, B., Gou, F., Feng, Z., Mao, Y. et al. (2014) The CRISPR/Cas9 system produces specific and homozygous targeted gene editing in rice in one generation. *Plant Biotechnol. J.* **12**, 797–807.
- Zhang, Y., Liang, Z., Zong, Y., Wang, Y., Liu, J., Chen, K., Qiu, J.L. et al. (2016) Efficient and transgene-free genome editing in wheat through transient expression of CRISPR/Cas9 DNA or RNA. *Nat. Commun.* **7**, 12617.
- Zhou, H., Liu, B., Weeks, D.P., Spalding, M.H. and Yang, B. (2014) Large chromosomal deletions and heritable small genetic changes induced by CRISPR/Cas9 in rice. *Nucleic Acids Res.* **42**, 10903–10914.
- Zhu, J.J., Song, N., Sun, S.L., Yang, W.L., Zhao, H.M., Song, W.B. and Lai, J.S. (2016) Efficiency and inheritance of targeted mutagenesis in maize using CRISPR-Cas9. *J. Genet. Genomics*, **43**, 25–36.

Supporting information

Additional Supporting Information may be found online in the supporting information tab for this article:

Figure S1 Protoplast transfection of various species.

Figure S2 Construction of pCambia1300-OsU3-Cas9.

Figure S3 Targeted mutagenesis of *Bambusa oldhamii* protoplasts.

Figure S4 The putative mutated *BoPDS* PCR products from *Bambusa oldhamii* protoplasts were cloned and validated by sequencing.

Figure S5 Targeted mutagenesis of *Setaria italica* protoplasts.

Figure S6 Targeted mutagenesis of *Oryza sativa* protoplasts.

Figure S7 Targeted mutagenesis of *Zea mays* protoplasts.

Figure S8 The PCR-RFLP-PCR results of *ZmIPK* mutagenesis.

Figure S9 Targeted mutagenesis of *Arabidopsis thaliana* protoplasts using constructs carrying the OsU3 or OsU6 monocot promoter.

Figure S10 Targeted mutagenesis of *Brassica oleracea* protoplasts using constructs carrying the OsU3 or OsU6 monocot promoter.

Figure S11 The putative mutated *BoGA4a* PCR products from *Brassica oleracea* protoplasts were cloned and validated by sequencing.

Figure S12 Targeted mutagenesis of *B. napus* protoplasts using constructs carrying the OsU3 or OsU6 monocot promoter.

Figure S13 Targeted mutagenesis of *Nicotiana tabacum* protoplasts.

Figure S14 Targeted mutagenesis of *Solanum lycopersicum* protoplasts.

Figure S15 Effect of incubation times on *NtPDS* target mutagenesis analysed in single protoplasts in Experiments 2 and 3.

Figure S16 Targeted mutagenesis of *NtPDS* in tobacco protoplast regenerants of Experiment 1.

Figure S17 Effect of plasmid dosage on *NtPDS* mutagenesis in

tobacco protoplast regenerants.

Figure S18 Schematic representation of single-cell isolation and validation of targeted mutagenesis.

Figure S19 Schematic of tobacco protoplast regeneration.

Table S1. Protocol for protoplast isolation and PEG transformation of different Poaceae species.

Table S2 Protocol for protoplast isolation and PEG transformation of different Brassicaceae species.

Table S3 Protocol for protoplast isolation and PEG transformation of different Solanaceae species.


Data S1 The *NtPDS* sequences of Figure 3 (Experiment 1) and Table 2.

Data S2 The *NtPDS* sequences of Figure 4 (Experiment 1) and Figure S15 (Experiment 2 and 3).

Data S3 The sequences of *Zea mays* single-cell *ZmIPK* genes in Figure 5.


Data S4 The *NtPDS* sequences of Figure 7 (Exp2R1) and Table 4.

Data S5 The *NtPDS* sequences of Figure S17.




Supplemental Figure 1. Protoplast transfection of various species.

Protoplasts from 13 species were isolated following the methods described in Materials and Methods section. Twenty μg mRFP-NLS plasmid DNA was separately delivered to protoplasts from each of the 13 species using a PEG-mediated method. Number after the species name: transfection efficiency. Protoplasts were photographed after 24 hrs. For each sample: left, epifluorescence; right, overlay of epifluorescence and bright field. Bars = 50 μm .




Supplemental Figure 2. Construction of pCambia1300-OsU3-Cas9.

Any target site fragment encoding the guide RNA with suitable adaptor sequences can be added to the sgRNA scaffold by insertion into the *AarI* site.



Supplemental Figure 3. Targeted mutagenesis of *Bambusa oldhamii* protoplasts.

- A. Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU3-*BoPDS* sgRNA1 and sgRNA2, and incubated for 48 hrs. Lanes 4 and 5, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU3-*BoPDS* sgRNA2, and incubated 48 hrs. Lanes 6 and 7, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plants.
- B. Information on the target gene, including the primer sequences used to amplify the target region after transfection and the restriction enzyme used in this experiment.




Supplemental Figure 4. The putative mutated *BoPDS* PCR products from *Bambusa oldhamii* protoplasts were cloned and validated by sequencing. DNA of *B. oldhamii* protoplasts 48 hrs after transfection with 35S-Cas9-OsU3-*BoPDS* sgRNA was isolated and subjected to PCR using primers targeting the *PDS* gene (A) or a second round of PCR (B). The products were subjected to TA cloning. Inserted DNA was amplified by PCR and sequenced by the Sanger method.

- A. The PCR products (from Supplemental Figure 3A) were TA-cloned, and colony PCR products were digested by *Bsa*I. Yellow star: putative *BoPDS* mutant clones. The putative mutant clones were subjected to Sanger sequencing. The sequence of the wild-type target gene (blue) and the PAM sequence (red) are highlighted.
- B. In this experiment, a second PCR step was added after PCR-RFLP (A). The mixture was subjected to restriction digestion after PCR amplification. The PCR product was TA-cloned and subjected to colony PCR. The colony PCR products were digested by *Bsa*I. Yellow star: putative mutant *PDS* clones. These clones were subjected to Sanger sequencing. The sequence of the wild-type target gene (blue) and the PAM sequence (red) are highlighted.




Supplemental Figure 5. Targeted mutagenesis of *Setaria italica* protoplasts.

- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plant. PCR products of genomic DNA from protoplasts treated with 35S-Cas9-OsU3-*SiPDS* sgRNA incubated 24 hrs (Lanes 4 and 5) and 48 hrs (Lanes 6 and 7); -, undigested control.
- Information on the target gene, including the primer sequences used to amplify the target region after transfection and the restriction enzyme used in this experiment.
- Sequence-based detection of mutations induced by Cas9-OsU3-*SiPDS* sgRNA. Wild-type sequence of the target gene (blue), PAM sequence (red), and insertion sequences (purple) are highlighted. The light grey color indicates inserted DNA within the deletion region.




Supplemental Figure 6. Targeted mutagenesis of *Oryza sativa* protoplasts.

- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control. PCR products of genomic DNA from protoplasts exposed to 35S-Cas9-OsU3- *OsPDS* sgRNA incubated 24 hrs (Lanes 4 and 5) and 48 hrs (Lanes 6 and 7) -, undigested control.
- Information on the target gene, including the primer sequences used to amplify the target region after transfection and the restriction enzyme used in this experiment.
- Sequence-based detection of mutations induced by Cas9-OsU3-*OsPDS* sgRNA. Sequence of the wild-type target gene (blue). The protospacer-adjacent motif (PAM, red) sequence and inserted sequences (purple) are highlighted. The number in parentheses indicates the size of the insert.



Supplemental Figure 7. Targeted mutagenesis of *Zea mays* protoplasts.

- Maize protoplasts were transfected with different sgRNAs: 1, 35S-Cas9-OsU3-*ZmIPK* sgRNA1; 2, 35S-Cas9-OsU3-*ZmIPK* sgRNA2; or 1+2, co-transfected with both sgRNAs 1 and 2 that target different regions of the *IPK* gene. DNA from the protoplast pool was amplified by PCR 48 hrs after transfection. The PCR products were digested by restriction enzymes (sgRNA1, *SacI*; sgRNA2, *BamHI*) according to the targeted sequence.
- Information on the target gene regions, including the primer sequences used to amplify the target regions and the restriction enzymes used in this experiment.
- Sequence-based detection of mutations induced by Cas9-OsU3-*ZmIPK* sgRNA1. Wild-type sequence of the target gene (blue), the PAM sequence (red), and the mutated nucleotide (green) are highlighted.
- Sequence-based detection of mutations induced by Cas9-OsU3-*ZmIPK* sgRNA2. Wild-type sequence of the target gene (blue) and the PAM sequence (red) are highlighted.




Supplemental Figure 8. The PCR-RFLP-PCR results of *ZmIPK* mutagenesis.

- A. PCR-amplified and restriction enzyme-digested (PCR-RFLP) DNA from Supplemental Figure 7A was re-amplified by PCR from the restriction enzyme mixture. Lane 1, DNA markers; Lane 2, PCR-amplified DNA from protoplasts transfected with only *ZmIPK* sgRNA1 was digested with *SacI* and subjected to a second PCR (PCR-RFLP-PCR); Lane 3, PCR-amplified DNA from protoplasts transfected with only *ZmIPK* sgRNA2 was digested with *Bam*HI and subjected to a second PCR (PCR-RFLP-PCR); Lanes 4-6, PCR-amplified DNA from protoplasts co-transfected with both *ZmIPK* sgRNA1 and sgRNA2 was digested with *SacI* (Lane 4), *Bam*HI (Lane 5), or *SacI* and *Bam*HI (Lane 6) and then subjected to a second round of PCR (PCR-RFLP-PCR); Lanes 7-9, PCR-amplified DNA from wild-type protoplasts was digested with *SacI* (Lane 7), *Bam*HI (Lane 8), or *SacI* and *Bam*HI (Lane 9) and then subjected to a second round of PCR.
- B. Sequence-based detection of deletions induced by co-transfection of maize protoplasts with Cas9-OsU3-*ZmIPK* sgRNA1 and sgRNA2. Sequence of the wild-type target gene (blue) and the PAM sequence (red) are highlighted. The number in parentheses indicates the size of the deletion.




Supplemental Figure 9. Targeted mutagenesis of *Arabidopsis thaliana* protoplasts using constructs carrying the OsU3 or OsU6 monocot promoter.

- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plants. PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU3-*AtPDS* sgRNA after incubation for 24 hrs (Lanes 4 and 5) or 48 hrs (Lanes 6 and 7).
- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plants. PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU6-*AtPDS* sgRNA after incubation for 24 hrs (Lanes 4 and 5) or 48 hrs (Lanes 6 and 7).
- Information on the target gene, including the primer sequences used to amplify the target region and the restriction enzyme used in this experiment.
- Sequence-based detection of mutations induced by Cas9-OsU6-*AtPDS* sgRNA. Sequence of the wild-type target gene (blue), the PAM sequence (red), and inserted sequences (purple) are highlighted.



Supplemental Figure 10. Targeted mutagenesis of *Brassica oleracea* protoplasts using constructs carrying the OsU3 or OsU6 monocot promoter.

- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plants. PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU3-*BolGA4a* sgRNA after incubation for 24 hrs (Lanes 4 and 5) or 48 hrs (Lanes 6 and 7).
- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plants. PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU6-*BolGA4a* sgRNA after incubation for 24 hrs (Lanes 4 and 5) or 48 hrs (Lanes 6 and 7).
- Information on the target genes, including the primer sequences which were used to amplify the target region, and the restriction enzyme which was used in this experiment.




Supplemental Figure 11. The putative mutated *BolGA4a* PCR products from *Brassica oleracea* protoplasts were cloned and validated by sequencing. DNA from *B. oleracea* protoplasts transfected with 35S-Cas9-OsU3-*BolGA4a* sgRNA after 24 hrs of incubation was isolated, subjected to PCR, and T/A-cloned.

- The colony PCR products were digested with *HphI*. +, *HphI* digested; -, control. Symbols colored in white, putative mutated clones; Gray, wild-type.
- Sequence-based detection of mutations induced by Cas9-OsU3-*BolGA4a* sgRNA. Sequence of the wild-type target gene (blue), the PAM sequence (red), and inserted sequences (purple) are highlighted.



Supplemental Figure 12. Targeted mutagenesis of *B. napus* protoplasts using constructs carrying the OsU3 or OsU6 monocot promoter.

- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plants. PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU3- *BnGA4a* sgRNA after incubation for 24 hrs (Lanes 4 and 5) or 48 hrs (Lanes 6 and 7).
- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plants. PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU6- *BnGA4a* sgRNA after incubation for 24 hrs (Lanes 4 and 5) or 48 hrs (Lanes 6 and 7).
- Information on the target gene, including the primer sequences used to amplify the target region and the restriction enzyme used in this experiment.
- Sequence-based detection of mutations induced by Cas9-OsU3-*BnGA4a* sgRNA. Sequence of the wild-type target gene (blue), the PAM sequence (red), and inserted sequences (purple) are highlighted.



Supplemental Figure 13. Targeted mutagenesis of *Nicotiana tabacum* protoplasts.


- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plants. PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU3-*NtPDS* sgRNA after incubation for 24 hrs (Lanes 4 and 5) or 48 hrs (Lanes 6 and 7).
- Information on the target gene, including the primer sequences used to amplify the target region and the restriction enzyme used in this experiment.
- Sequence-based detection of mutations induced by Cas9-OsU3-*NtPDS* sgRNA. Sequence of the wild-type target gene (blue), the PAM sequence (red), and inserted sequences (purple) are highlighted. The numbers in parentheses indicate the size of the insertion.




Supplemental Figure 14. Targeted mutagenesis of *Solanum lycopersicum* protoplasts.

- Detection of mutations using PCR-RFLP. Lane 1, DNA marker; Lanes 2 and 3, PCR products of digested (+) and undigested (-) genomic DNA from protoplasts of wild-type control plants. PCR products of digested (+) and undigested (-) genomic DNA from protoplasts treated with 35S-Cas9-OsU3-*SIPDS* sgRNA after incubation for 24 hrs (Lanes 4 and 5) or 48 hrs (Lanes 6 and 7).
- Information on the target gene, including the primer sequences used to amplify the target region and the restriction enzyme used in this experiment.
- Sequence-based detection of mutations induced by Cas9-OsU3-*SIPDS* sgRNA. Sequence of the wild-type target gene (blue), the PAM sequence (red) and inserted sequences (purple) are highlighted. The number in parentheses indicates the size of the insertion.

Exp. 2



Exp. 3



Supplemental Figure 15. Effect of incubation times on *NtPDS* target mutagenesis analyzed in single protoplasts in Experiments 2 and 3.

Tobacco protoplasts were transfected with 20 µg plasmid DNA containing the expression cassette of *NtPDS* sgRNA and SaCas9 (Kaya et al., 2016) and incubated for various number of days. Target mutations were analyzed by RFLP. S, *N. sylvestris* form; T, *N. tomentosiformis* form. Green frame, wild-type RFLP control; red frame, albino mutant RFLP control.

Exp. 1

Repeat 1

Albino S 100

T 100

Green S 35

T 15

Repeat 2

Albino S 100

T 100

Green S 35

T 20


Repeat 3

Albino S 100

T 100


Green S 30

T 10




Supplemental Figure 16. Targeted mutagenesis of *NtPDS* in tobacco protoplast regenerants of Experiment 1.

Protoplast regenerants derived from 20 µg plasmid DNA treatments were divided into albino (A) and green (G) plantlets. Twenty plantlets in each type were analyzed for targeted mutagenesis. The target sequences were amplified and validated using RFLP. S, *N. sylvestris* form; T, *N. tomentosiformis* form. Green frame, wild-type RFLP control; red frame, albino mutant RFLP control.




Supplemental Figure 17. Effect of plasmid dosage on *NtPDS* mutagenesis in tobacco protoplast regenerants.

Tobacco protoplasts were transfected with 0-20 µg plasmid DNA containing the expression cassettes *NtPDS* sgRNA and SaCas9 (Kaya et al., 2016) and regenerated into shoots. Targeted mutagenesis of 10 green regenerants from each treatment were analyzed by RFLP. S, *N. sylvestris* form; T, *N. tomentosiformis* form. Green frame, wild-type RFLP control; red frame, albino mutant RFLP control.



Supplemental Figure 18. Schematic representation of single cell isolation and validation of targeted mutagenesis.

Protoplasts are isolated and transfected (detailed information is given in Supplemental Table 1). The protoplasts are centrifuged and transferred to liquid callus medium (1/2 MS medium supplemented with 0.4 M mannitol, 30 g/L sucrose, 1 mg/L NAA, and 0.3 mg/L kinetin). The protoplast concentration is measured using a haemocytometer and adjusted to 1 cell/μl using the same medium. A single protoplast is isolated using a 2 μl laboratory pipette. The protoplast is transferred to a slide and the condition is checked by microscopy (Olympus, Tokyo, Japan). The protoplast is transferred to PCR mixture containing the first pair of primers for primary DNA amplification. One μl of product is used as the template for a second round of PCR using a second pair of primers. The second PCR product is digested by a restriction enzyme and analyzed by electrophoresis to validate mutagenesis of the target sequences.



Supplemental Figure 19. Schematic of tobacco protoplast regeneration.

The protocol is modified from Liao (1990). The transfected protoplasts are incubated in a 5 cm diameter petri dish containing liquid callus medium (1/2MS medium supplemented with 0.4 M mannitol, 30 g/L sucrose, 1 mg/L NAA and 3 mg/L kinetin). After 2-3 weeks, the proliferating protoplasts form dust-like calli. The calli are embedded in solidified callus medium in a 9 cm diameter petri dish for 3-4 weeks. In the callus stage, the explants are incubated in the dark (gray background). Calli larger than 3 mm can be embedded in shooting medium (MS medium supplemented with 2 mg/L kinetin, 0.3 mg/L IAA, 0.4 M mannitol, and 30 g/L sucrose) for shoot induction. After one month, the multiple shoots which contain leaves or are of a size larger than 5 mm are transferred to fresh shooting medium for 2-3 weeks for shoot proliferation. Multiple shoots with leaves are transferred to solidified rooting medium (MS medium supplemented with 0.1 mg/L IAA, and 30 g/L sucrose). After shooting, all explants are incubated in light/dark cycling conditions (light/dark: 16/8 hrs). KIN: kinetin.

Supplemental Material and Methods

The pCAMBIA1300-OsU3-Cas9 and pCAMBIA1300-OsU6-Cas9 plasmids can be used for both protoplast transfection and *Agrobacterium*-mediated transformation. The nucleotide “A” was included immediately upstream of the 5’ end of the target sequence, and a “G” was included immediately upstream of the 5’ end of the target sequence, when cloned into the plasmids pOsU3-sgRNA or pOsU6-sgRNA, respectively. According to the target sequence, we synthesized a pair of complementary oligonucleotides with 5’ adapters (pOsU3-sgRNA: Fw 5'-GGCA-20 bp-3', Rv: 5'-AAAC-20 bp-3'; pOsU6-sgRNA: Fw 5'-CTTG-20 bp-3', Rv: 5'-AAAC-20 bp-3'). The pair of complementary oligonucleotides with their corresponding adapters was annealed, phosphorylated, and ligated to the linearized plasmids digested with *AarI* (Fermentas/Thermo Scientific Inc., Waltham, MA) using T4 DNA ligase (New England Biolabs Inc., Ipswich, MA). The ligation mixture was transformed into *E. coli* DH5 α , and at least eight clones were picked, subjected to PCR, and the PCR products of candidate clones were analyzed by Sanger DNA sequencing. A plasmid containing the correct target sequence was later used for transfection into protoplasts.

Supplemental Table 1. Protocol for protoplast isolation and PEG transformation of different Poaceae species

Species	Digestion Buffer	Digestion condition	Wash Condition	Centrifugation Speed	Cell number	MMG Buffer	PEG solution	PEG Transfection time
<i>Setaria italica</i>	1.5% (wt/vol) Cellulase R10, 0.75% (wt/vol) macerozyme R10, 10 mM MES (pH5.7), 0.6 M mannitol, 55 °C for 10 min, cool to RT, 10 mM CaCl ₂ • 2H ₂ O, and 0.1% (wt/vol) BSA sterilize buffer with a 0.45 µm filter	3 hrs/50 rpm (25°C)	W5 solution: 154 mM NaCl, 125 mM CaCl ₂ • 2H ₂ O, 5 mM KCl, 2 mM MES (pH5.7), 5 mM glucose, centrifugation for 2 min Wash twice Keep cell on ice for 30 min	100g	1.2×10^7 /g FW	0.6M mannitol, 15 mM MgCl ₂ , 4 mM MES (pH 5.7)	40% (wt/vol) PEG 4000, 0.6 M mannitol , Warm the solution at 60°C until dissolved and cool down to 40°C 0.1 M CaCl ₂ • 2H ₂ O,	20 min
<i>Oryza sativa</i>				200g	4.2×10^6 /g FW			
<i>Triticum aestivum</i>				100g	4.1×10^6 /g FW			
<i>Zea mays</i>				150g	2.3×10^6 /g FW			

Supplemental Table 2. Protocol for protoplast isolation and PEG transformation of different Brassicaceae species

Species	Digestion Buffer	Digestion condition	Wash Condition	Centrifugation Speed	Cell number	MMG Buffer	PEG solution	PEG Transfection time
<i>Arabidopsis thaliana</i>	1% (wt/vol) Cellulase R10, 0.25% (wt/vol) macerozyme R10, 20 mM MES (pH5.7), 20 mM KCl, 0.4 M mannitol, 55 °C for 10 min, cool to RT 10 mM CaCl ₂ • 2H ₂ O and 0.1% BSA sterilize buffer with a 0.45 µm filter	1.5hrs/50 rpm (25°C)	W5 solution: 154 mM NaCl 125 mM CaCl ₂ • 2H ₂ O 5 mM KCl 2 mM MES (pH5.7) 5 mM glucose Centrifugation for 2 min Wash twice Keep cell on ice for 30 min	100g	3.0×10^7 /g FW	0.4 M mannitol 15 mM MgCl ₂ 4 mM MES (pH 5.7)	40% (wt/vol) PEG 4000 0.2 M mannitol 0.1 M CaCl ₂ • 2H ₂ O	6 min
<i>Brassica oleracea</i>	1.5% (wt/vol) Cellulase R10,	3hrs/50 rpm (25°C)			4.2×10^6 /g FW			10 min
<i>Brassica napus</i>	0.75% (wt/vol) macerozyme R10,				2.2×10^6 /g FW	0.6 M mannitol 15 mM MgCl ₂	40% (wt/vol) PEG 4000,	6 min
<i>Cleome</i> spp.	10mM MES (pH5.7), 0.6M mannitol, 55 °C for 10 min, cool to RT 10 mM CaCl ₂ • 2H ₂ O and 0.1% (wt/vol) BSA sterilize buffer with a 0.45-µm filter	1.5hrs/50 rpm (25°C)			3.5×10^6 /g FW	4 mM MES (pH 5.7)	0.6 M mannitol, Warm the solution at 60 °C until dissolved and cool down to 40°C 0.1 M CaCl ₂ • 2H ₂ O	10 min

Supplemental Table 3. Protocol for protoplast isolation and PEG transformation of different Solanaceae species

Species	Digestion Buffer	Digestion condition	Wash Condition	Centrifugation Speed	Cell number	MMG Buffer	PEG solution	PEG Transfection time
<i>Solanum tuberosum</i>	1.5% (wt/vol) Cellulase R10, 0.75% (wt/vol) macerozyme R10, 10 mM MES (pH5.7), 0.6 M mannitol, 55 °C for 10 min, cool to RT 1 mM CaCl ₂ • 2H ₂ O and 0.1% (wt/vol) BSA sterilize buffer with a 0.45 µm filter	3 hrs/50 rpm (25°C)	W5 solution: 154 mM NaCl, 125 mM CaCl ₂ • 2H ₂ O, 5 mM KCl, 2 mM MES (pH5.7), 5 mM glucose, centrifugation for 2 min Wash twice Keep cell on ice for 30 min	100g	6.4×10^6 /g FW	0.6M mannitol, 15 mM MgCl ₂ , 4 mM MES (pH 5.7)	40% (wt/vol) PEG 4000, 0.6 M mannitol , Warm the solution at 60 °C until dissolved and cool down to 40°C 0.1 M CaCl ₂ • 2H ₂ O	20 min
<i>Solanum lycopersicum</i> (cv microtom) suspension cells	1.5% (wt/vol) Cellulase R10, 0.75% (wt/vol) macerozyme R10, 10 mM MES (pH5.7), 0.6 M mannitol, 10 mM sodium acetate 55 °C for 10 min, cool to RT, 10 mM CaCl ₂ • 2H ₂ O, and 0.1% (wt/vol) BSA sterilize buffer with a 0.45 µm filter	4 hrs/50 rpm (35°C)		250g	7.2×10^6 /g FW			
<i>Nicotiana tabacum</i> BY-2 suspension cells				250g	1.4×10^7 /g FW			

Supplemental Data 1. The *NtPDS* sequences of Figure 3 (Experiment 1) and Table 2.

Red box in Sample no. column indicates all four copies of *NtPDS* have the same mutations. S, *N. sylvestris* form; T, *N. tomentosiformis* form. Orange box-Homo, homozygous in this form; Light blue box-hetero, heterozygous in this form; W, wild-type; I, insertion; D, deletion; Letter in red, the mutated nucleotide.

Experiment 1

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
R1 5 µg-12	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R1 5 µg-18	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
R1 5 µg-21	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
R1 5 µg-22	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R1 10 µg-3	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
R1 10 µg-4	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			D	GATGCCTAACAAGC --G GGGGAG	
R1 10 µg-8	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
R1 10 µg-14	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R1 10 µg-16	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
R1 10 µg-19	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R1 20 µg-1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	

			I	GATGCCTAACAAGCTCAGGGGAG	
R1 20 µg-6	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC- (+12 bps)	
R1 20 µg-8	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R1 20 µg-9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R1 20 µg-13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R1 20 µg-16	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R1 20 µg-18	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
		Hetero	I	GATGCCTAACAAGCACAGGGGAG	
R1 20 µg-19	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
R1 20 µg-21	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	

Experiment 2

Sample no.	Genome type	genotype	Mutation type	NtPDS sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
R2 5 µg-1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R2 5 µg-5	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R2 5 µg-6	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
R2 5 µg-8	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R2 5 µg-9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	

			I	GATGCCTAACAAGC A CAGGGGAG	
R2 5 µg-10	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 5 µg-12	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 5 µg-13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 5 µg-15	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			D	GATGCCTAACAAGC- (-8 bps)	
R2 5 µg-16	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 5 µg-17	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 5 µg-18	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 5 µg-19	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 5 µg-20	S	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 10 µg-2	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 10 µg-3	S	Homo	D	GATGCCTAACAAGC -- AGGGGAA	
			D	GATGCCTAACAAGC -- AGGGGAG	
R2 10 µg-4	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
		Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	D	GATGCCTAACAAGC -- AGGGGAA	
			D	GATGCCTAACAAGC -- AGGGGAA	
R2 10 µg-5	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 10 µg-6	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 10 µg-7	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	

			I	GATGCCTAACAAGC T CAGGGGAA	
R2 10 µg-8	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 10 µg-9	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 10 µg-10	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCC T AGGGGAG	
R2 10 µg-11	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			D	GATGCCTAACAAGC--AGGGGAG	
R2 10 µg-12	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCC T AGGGGAA	
R2 10 µg-13	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC TT CAGGGGA	
R2 10 µg-14	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 10 µg-16	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 10 µg-18	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
		Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 10 µg-20	S	Hetero	D	GATGCCTAACAA-C-CAGGGGAG	
		Hetero	D	GATGCCTAACAT G ---AGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 20 µg-1	S	Homo	I	GATGCCTAACAAGC (+106 bps)	
			I	GATGCCTAACAAGC (+106 bps)	
	T	Homo	D	GATGCCTAACAAGC--AGGGGAA	
			D	GATGCCTAACAAGC--AGGGGAA	
R2 20 µg-2	S	Hetero	D	GATGCCTAACAA---CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	D	GATGCCTAACAA---CAGGGGAG	

			D	GATGCCTAACAA---CAGGGGAG	
R2 20 µg-3	S	Hetero	W	GATGCCTAACAAAGC-CAGGGGAG	
			I	GATGCCTAACAAAGCTCAGGGGAG	
R2 20 µg-4	S	Hetero	I	GATGCCTAACAAAGCTCAGGGGAG	
			I	GATGCCTAACAAAGCGCAGGGGAG	
R2 20 µg-5	T	Homo	I	GATGCCTAACAAAGCTCAGGGGAA	
			I	GATGCCTAACAAAGCTCAGGGGAA	
R2 20 µg-6	S	Homo	I	GATGCCTAACAAAGCTCAGGGGAG	
			I	GATGCCTAACAAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAAGC-CAGGGGAA	
			I	GATGCCTAACAAAGCTCAGGGGAA	
R2 20 µg-7	S	Hetero	W	GATGCCTAACAAAGC-CAGGGGAG	
			I	GATGCCTAACAAAGCA CAGGGGAG	
R2 20 µg-8	S	Hetero	W	GATGCCTAACAAAGC-CAGGGGAG	
			I	GATGCCTAACAAAGCTCAGGGGAG	
R2 20 µg-9	S	Hetero	W	GATGCCTAACAAAGC-CAGGGGAG	
			I	GATGCCTAACAAAGCA CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAAGC-CAGGGGAA	
			I	GATGCCTAACAAAGCA CAGGGGAA	
R2 20 µg-10	S	Hetero	W	GATGCCTAACAAAGC-CAGGGGAG	
			I	GATGCCTAACAAAGCA CAGGGGAG	
	T	Homo	I	GATGCCTAACAAAGCTCAGGGGAA	
			I	GATGCCTAACAAAGCTCAGGGGAA	
R2 20 µg-11	S	Hetero	I	GATGCCTAACAAAGCA CAGGGGAG	
			I	GATGCCTAACAAAGCTCAGGGGAG	
R2 20 µg-12	S	Hetero	I	GATGCCTAACAAAGCTCAGGGGAG	
			I	GATGCCTAACAAAGCGCATGGGAG	
	T	Hetero	I	GATGCCTAACAAAGCA CAGGGGAA	
			I	GATGCCTAACAAAGCTCAGGGGAA	
R2 20 µg-13	S	Hetero	I	GATGCCTAACAAAGCTCAGGGGAG	
			D	GATGCCTAACAAAGC--AGGGGAG	
	T	Hetero	D	GATGCCTAACAAAGC--AGGGGAA	
			I	GATGCCTAACAAAGCTCAGGGGAA	
R2 20 µg-14	S	Homo	I	GATGCCTAACAAAGCTCAGGGGAG	
			I	GATGCCTAACAAAGCTCAGGGGAG	
R2 20 µg-16	T	Hetero	D	GATGCCTAACAAAGC--AGGGGAA	
			I	GATGCCTAACAAAGCA CAGGGGAA	

R2 20 µg-17	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 20 µg-18	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 20 µg-19	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC- (+11 bps)	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 20 µg-20	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	

Experiment 3

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
R3 5 µg-2	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R3 5 µg-10	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R3 5 µg-14	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R3 10 µg-1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R3 10 µg-4	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R3 10 µg-5	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R3 10 µg-8	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R3 10 µg-9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R3 10 µg-10	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	

			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
R3 10 µg-11	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
R3 10 µg-15	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
R3 10 µg-16	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
R3 10 µg-17	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			D	GATGCCTAACAAGC- (-11 bps)	
R3 10 µg-19	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R3 10 µg-20	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R3 20 µg-1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R3 20 µg-7	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
R3 20 µg-8	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R3 20 µg-9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R3 20 µg-11	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
R3 20 µg-13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	

			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
R3 20 µg-14	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R3 20 µg-17	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R3 20 µg-18	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
R3 20 µg-20	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	

Supplemental Data 2. The *NtPDS* sequences of Figure 4 (Experiment 1) and Supplemental Figure 15 (Experiment 2 and 3). Red box in Sample no. column indicates four copies of *NtPDS* have same mutations. S: *sylvestris* form. T: *tomentosiformis* form. Orange box-Homo: homozygous in this form. Light blue box-hetero: heterozygous in this form. W: wild type. I: insertion. D: deletion. Letter in red: the mutated nucleotide.

Experiment 1

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
R1 0 day-5	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R1 0 day-14	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
R1 1 day-9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R1 1 day-17	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R1 2 day-5	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R1 2 day-6	T	Homo	D	GATGCCT (-17 bps)	
			D	GATGCCT (-17 bps)	
R1 2 day-9	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R1 2 day-11	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R1 2 day-12	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
R1 2 day-15	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	

			I	GATGCCTAACAAGCACAGGGGAG	
R1 2 day-16	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R1 2 day-17	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R1 2 day-19	S	Homo	I	GATGCCTAACAAGATCAGGGGAG	
			I	GATGCCTAACAAGATCAGGGGAG	
R1 2 day-20	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGATCAGGGGAG	
R1 3 day-1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R1 3 day-6	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R1 3 day-8	S	Homo	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
R1 3 day-11	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R1 3 day-13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
R1 3 day-14	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R1 3 day-15	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
R1 4 day-3	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R1 4 day-4	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R1 4 day-6	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R1 4 day-9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R1 4 day-11	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	

R1 4 day-15	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC C AGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC C AAGGGGAA	
R1 4 day-16	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC C AAGGGGAA	

Experiment 2

Sample no.	Genome type	genotype	Mutation type	NtPDS sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
R2 0 day-9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 0 day-17	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 1 day-2	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC CT CAGGGGA	
	T	Homo	I	GATGCCTAACAAGC AC AGGGGAA	
			I	GATGCCTAACAAGC AC AGGGGAA	
R2 1 day-9	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 1 day-17	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC C AGGGGAA	
R2 1 day-19	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAG A -CAGGGGAG	
R2 2 day-1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC AC AGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 2 day-3	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 2 day-4	S	Homo	I	GATGCCTAACAAGC AC AGGGGAG	
			I	GATGCCTAACAAGC AC AGGGGAG	

	T	Homo	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
R2 2 day-7	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 2 day-9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 2 day-11	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
		Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
R2 2 day-12	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
		Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
R2 2 day-13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
		Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
R2 2 day-16	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 2 day-20	S	Hetero	W	GATGCCTAACAAGA-CAGGGGAG	
			I	GATGCCTAACAAGA T CAGGGGAG	
R2 3 day-1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 3 day-2	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
R2 3 day-3	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
R2 3 day-4	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 3 day-5	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 3 day-6	S	Hetero	D	GATGCCTAACAAGC --T GGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			D	GATGCCTAACAAGC --A GGGGAA	
R2 3 day-7	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 3 day-8	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	

R2 3 day-9	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
R2 3 day-10	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 3 day-11	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 3 day-12	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 3 day-13	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
R2 3 day-14	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 3 day-15	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
R2 3 day-16	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 3 day-17	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
R2 3 day-18	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
R2 3 day-19	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 3 day-20	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R2 4 day-6	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R2 4 day-10	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 4 day-11	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	

R2 4 day-13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 4 day-14	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 4 day-15	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R2 4 day-16	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
R2 4 day-18	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			D	GATGCCTAACAAGC-(-17 bps)	
R2 4 day-20	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	

Experiment 3

Sample no.	Genome type	genotype	Mutation type	NtPDS sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
R3 0 day-19	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAG A -CAGGGGAG	
R3 1 day-1	T	Homo	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
R3 1 day-2	S	Homo	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
R3 1 day-6	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R3 1 day-8	T	Hetero	D	GATGCCTAACAAGC--AGGGGAA	
			D	GATGCCTAACAAG A --AGGGGAA	
R3 1 day-15	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	

R3 2 day-2	S	Hetero	D	GATGCCTAACA---TCAGGGGAG	
		Hetero	D	GATGCCTAACA---TCAGGGGAG	
R3 2 day-3	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
R3 2 day-5	T	Homo	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
R3 2 day-7	S	Hetero	D	GATGCCTAACAAGC--CAGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	D	GATGCCTAACAAGC--AGGGGAA	
			D	GATGCCTAACAAGC--AGGGGAA	
R3 2 day-11	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R3 2 day-15	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R3 2 day-16	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R3 2 day-18	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
R3 3 day-1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R3 3 day-3	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			D	GATGCCTAACAAGC-(-71 bps)	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			D	GATGCCTAACA- (11 bps)	
R3 3 day-4	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
R3 3 day-5	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
R3 3 day-7	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
R3 3 day-8	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	

R3 3 day-9	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	D	GATGCC----- (-26 bps)	
			D	GATGCC----- (-26 bps)	
R3 3 day-12	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
R3 3 day-13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC-CAGGGGAA	
R3 3 day-15	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R3 3 day-16	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R3 3 day-18	S	Homo	D	GATGCCTAA-----CAGGGGAG	
			D	GATGCCTAA-----CAGGGGAG	
R3 4 day-1	S	Homo	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
R3 4 day-3	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
R3 4 day-6	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
R3 4 day-7	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC-CAG (+5 bps)	
R3 4 day-8	S	Hetero	D	GATGCCTAACA-----GGGGAG	
			I	GATGCCTAACAAGCTCGGGGAAG	
R3 4 day-16	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
R3 4 day-17	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	

Supplemental Data 3. The sequences of *Zea mays* single cell *ZmIPK* genes in Figure 5.

>no._1

CCGGGGCCTCGTGGCGTAGCCCGGGAGTAGTTGATGTCAATGACGAGATAGCGGTCGCCGGCGCGGACGTCG
CGGATCATGTGGAAGTTGAAGAGTTGCAGGCCAGCGCGCGGCGGAGGCCGCCGCGATCTGGTTGATGAAT
GCGGCGGGCGGCACGACGGCGTCCTCGAGACTCTTTTCGCCGTAGTACTCTCCGCCGTGCGCTCAGTGGGGA
GGTTGGAGACCTGGGAGAAGGAGACGGATGCTCGGAGACGACCTGGAGCATGGAGATGCGGTTGTGGAGGC
GGTCGATGGCGTGGGGCGGGTCGACGATGGGGACGGCCGGGTGGCGCGCGGCGAAGGCCACGAGCTGGGC
GCGCCAGTCGTCTCCGTAGAGCTTGTGA

>no._4

CGGTCTCTTGGCGTAGCCCGGGAGTAGTTGATGTCAATGACGAGATAGCGGTCGCCGGCGCGGACGTCGCGG
ATCATGTGGAAGTTGAAGAGTTGCAGGCCAGCGCGCGGCGGAGGCCGCCGCGATCTGGTTGATGAATGCG
GCGGGCGGCACGACGGCGTCCTCGAGACTCTTTTCGCCGTAGTACTCTCCGCCGTGCGCTCAGTGGGGAGGT
TGGAGACCTGGGAGAAGGAGACGGATGCTCGGAGACGACCTGGAGCATGGAGATGCGGTTGTGGAGGCGGT
CGATGGCGTGGGGCGGGTCGACGATGGGGACGGCCGGGTGGCGCGCGGCGAAGGCCACGAGCTGGGCGCG
CCAGTCGTCTCCGAGG

>no._13

ACGTAATCTTGTGGGACTTGGCGGTGCCGTGCGGCGACGAGGGGCTTGGCGATGAGCGGGAAGCGGAGCGCG
GCAAGGAGTCCGAAGTCGGCGAGCGCGGCAGCGTCGTAGACGACGACCTGGCTGGGGATACCGAAAGTGCT
GTCCTGGTCGGCGGCGTGGTCGAGCTCGGAGACGACCTGGAGCATGGAGATGCGGTTGTGGAGGCGGTGAT
GGCGTGGGGCGGGTCGACGATGGGGACGGCCGGGTGGCGCGCGGCGAAGGCCACGAGCTGGGCGCGCCAG
TCGTCTCCGTAGAGCTTGTGA

>no._16

TCGGGGCTCTTGGCGTAGCCCGGGAGTAGTTGATGTCAATGACGAGATAGCGGTCGCCGGCGCGGACGTCGC
GGATCATGTGGAAGTTGAAGAGTTGCAGGCCAGCGCGCGGCGGAGGCCGCCGCGATCTGGTTGATGAATG
CGGCGGACGGCACGACGGCGTCCTCGAGACTCTTTTCGCCGTAGTACTCTCCGCCGTGCGCTCAGTGGGGAG
GTTGGAGACCTGGGAGAAGGAGACGGATAGCTCGGAGACGACCTGGAGCATGGAGATGCGGTTGTGGAGGC
GGTCGATGGCGTGGGGCGGGTCGACGATGGGGACGGCCGGGTGGCGCGCGGCGAAGGCCACGAGCTGGGC
GCGCCAGTCGTCTCCGTGAAGGCTTGTGA

>no._18

CCGGTCTCTTGGCGTAGCCCGGGAGTAGTTGATGTCAATGACGAGATAGCGGTCGCCGGCGCGGACGTCGCG
GATCATGTGGAAGTTGAAGAGTTGCAGGCCAGCGCGCGGCGGAGGCCGCCGCGATCTGGTTGATGAATGC
GGCGGGCGGCACGACGGCGTCCTCGAGACTCTTTTCGCCGTAGTACTCTCCGCCGTGCGCTCAGTGGGGAGG
TTGGAGACCTGGGAGAAGGAGACGGATGCTCGGAGACGACCTGGAGCATGGAGATGCGGTTGTGGAGGCGG
TCGATGGCGTGGGGCGGGTCGACGATGGGGACGGCCGGGTGGCGCGCGGCGAAGGCCACGAGCTGGGCGC
GCCAGTCGTCTCCGTGAAGCTTGTGA

Supplemental Data 4. The *NtPDS* sequences of Figure 7 (Exp2R1) and Table 4. Red box in Sample no. column indicates all four copies of the *NtPDS* gene have the same mutations. S, *N. sylvestris* form; T, *N. tomentosiformis* form. Orange box-Homo, homozygous in this form; Light blue box-hetero, heterozygous in this form. W, wild-type; I, insertion; D, deletion; Letter in red, the mutated nucleotide.

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S			GATGCCTAACAAGC-CAGGGGAG	
	T			GATGCCTAACAAGC-CAGGGGAA	
Exp1R1A1	S	Hetero	D	GATGCCTAACAAGC-(-14 bps)	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCAAGGGGAA	
Exp1R1A2	S	Hetero	I	GATGCCTAACAAGCAAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R1A3	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp1R1A4	S	Homo	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp1R1A5	S	Homo	I	GATGCCTAACAAGCAAGGGGAG	
			I	GATGCCTAACAAGCAAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R1A6	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
Exp1R1A7	S	Hetero	I	GATGCCTAACAAGCAAGGGAAA	
			I	GATGCCTAACAAGCTCAGGGAAA	

	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R1A8	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp1R1A9	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp1R1A10	S	Homo	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R1A11	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp1R1A12	S	Hetero	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R1A13	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp1R1A14	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp1R1A15	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			D	GATGCCTAACAAGC--AGGGGAG	
	T	Homo	D	GATGCCTAACAAGC--AGGGGAA	
			D	GATGCCTAACAAGC--AGGGGAA	
Exp1R1A16	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCGCAGGGGAA	

			I	GATGCCTAACAAGC A CAGGGGAA	
Exp1R1A17	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC G CTGGGGAA	
			I	GATGCCTAACAAGC G CTGGGGAA	
Exp1R1A18	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC G CTGGGGAA	
			I	GATGCCTAACAAGC G CTGGGGAA	
Exp1R1A19	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			D	GATGCCTAACCAG-----GGGAG	
	T	Hetero	I	GATGCCTAACAAGC C CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R1A20	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	D	GATGCCTAACAAGC-- C AGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp1R1G1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAA	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G2	S	Homo	I	GATGCCTAACCAGG T CAGGGGAG	
			I	GATGCCTAACCAGG T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			D	GATGCCTAACAAGC-- A GGGGAA	
Exp1R1G3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G4	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	

			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G10	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G11	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAA	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G12	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
	T	Homo	D	GATGCCTAACA-----CAGGGGAA	
			D	GATGCCTAACA-----CAGGGGAA	
Exp1R1G13	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G14	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Exp1R1G15	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G16	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G17	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G18	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R1G19	S	Homo	W	GATGCCTAACAAGC G CAGGGGAG	
			W	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
		Hetero	I	GATGCCTAACAAGC G CAGGGGAA	
Exp1R1G20	S		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2A1	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G G C AGGGGA	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R2A2	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			D	GATGCCTAACAAGC -- AGGGGAG	

	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2A3	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Homo	D	GATGCCTAACAA---CAGGGGAA	
			D	GATGCCTAACAA---CAGGGGAA	
Exp1R2A4	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCAAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2A5	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2A6	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2A7	S	Hetero	I	GATGCCTAACAAGCAAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCAAGGGGAA	
			D	GATGCCTAACAAGC--AGGGGAA	
Exp1R2A8	S	Hetero	I	GATGCCTAACAAGCAAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2A9	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp1R2A10	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2A11	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	

			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2A12	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp1R2A13	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2A14	S	Homo	I	GATGCCTAACAAGCAAGGGGAG	
			I	GATGCCTAACAAGCAAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCAAGGGGAA	
			I	GATGCCTAACAAGCAAGGGGAA	
Exp1R2A15	S	Hetero	I	GATGCCTAACAAGCAAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	D	GATGCCTAACAAG---AGGGGAA	
			I	GATGCCTAACAAGCAAGGGGAA	
Exp1R2A16	S	Hetero	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCGGCAGGGGA	
	T	Hetero	I	GATGCCTAACAAGCAAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAG	
Exp1R2A17	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp1R2A18	S	Homo	I	GATGCCTAACAAGCAAGGGGAG	
			I	GATGCCTAACAAGCAAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp1R2A19	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2A20	S	Hetero	I	GATGCCTAACAAGCAAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCAAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	

Exp1R2G1	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGC (+33 bps)	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R2G2	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G5	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G7	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp1R2G8	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	

Exp1R2G10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
		Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R2G11	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G12	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G13	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G14	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G15	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G16	S	Homo	I	GATGCCTAACAAGC A CAGGGGA	
			I	GATGCCTAACAAGC A CAGGGGA	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G17	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G18	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G19	S		W	GATGCCTAACAAGC-CAGGGGAG	

			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R2G20	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Sample no.	Genome type	genotype	Mutation type	NtPDS sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3A1	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC C CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R3A2	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC C CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R3A3	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAG AG CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC C CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R3A4	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp1R3A5	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp1R3A6	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC C CAGGGGAA	
			D	GATGCCTAACAAGC (-25 bps)	

Exp1R3A7	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			D	GATGCCTAACAAGC --- GGGGAG	
	T	Homo	I	GATGCCTAACAAGC C CAGGGGAA	
			I	GATGCCTAACAAGC C CAGGGGAA	
Exp1R3A8	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC (+10 bps)	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R3A9	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp1R3A10	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC (+592 bps)	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R3A11	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R3A12	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC (+73 bps)	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp1R3A13	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp1R3A14	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp1R3A15	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC C CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp1R3A16	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	

			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp1R3A17	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R3A18	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp1R3A19	S	Hetero	I	GATGCCTAACAAGCCAGGGGAG	
			D	GATGCCTAACAAG-----AG	
	T	Hetero	I	GATGCCTAACAAGCAAGGGGAA	
			D	GATGCCTAACAAGC--AGGGGAA	
Exp1R3A20	S	Homo	D	GATGCCTAACAAG--CAGGGGAG	
			D	GATGCCTAACAAG--CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp1R3G1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp1R3G2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G5	S		W	GATGCCTAACAAGC-CAGGGGAG	

			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G6	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G10	S	Hetero	I	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC-CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G11	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G12	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			D	GATGCCTAACAAGC---GGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G14	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC-CAGGGGAG	

	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G15	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G16	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G17	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G18	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G19	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp1R3G20	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Regenerate no.	Genome type	genotype	Mutation type	NtPDS sequences	
Wild type	S			GATGCCTAACAAGC-CAGGGGAG	
	T			GATGCCTAACAAGC-CAGGGGAA	
Exp2R1A1	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	

			I	GATGCCTAACAAGCTCAGGGGAA	
Exp2R1A2	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp2R1A3	S	Homo	D	GATGCCT-----CAGGGGAG	
			D	GATGCCT-----CAGGGGAG	
	T	Hetero	D	GATGCCT-----CAGGGGAA	
			I	GATGCCTAACAAGC (+172 bps)	
Exp2R1A4	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp2R1A5	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			D	GATGCCTAACAAGCACAGGGGAG	
	T	Homo	D	GATGCCTAACAAGC--AGGGGAA	
			D	GATGCCTAACAAGC--AGGGGAA	
Exp2R1A6	S	Homo	D	GATGCCTAAC-----AGGGGAG	
			D	GATGCCTAAC-----AGGGGAG	
		Homo	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp2R1A7	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp2R1A8	S		D	GATGCCTAACAAGC--AGGGGAG	
			D	GATGCCTAACAAGC--AGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp2R1A9	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp2R1A10	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCGGGGGAA	
			D	GATGCCTAACAAGC--AGGGGAA	

Exp2R1A11	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R1A12	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R1A13	S	Homo	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp2R1A14	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R1A15	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R1A16	S	Hetero	I	GATGCCTAACAAGC (+477 bps)	
			I	GATGCCTAACAAGC (+475 bps)	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp2R1A17	S	Hetero	I	GATGCCTAACAAGC T TTGGTAGG	
			I	GATGCCTAACAAGC (+477 bps)	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R1A18	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp2R1A19	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R1A20	S	Homo	D	GATGCCTAACAAG -- CAGGGGAG	

			D	GATGCCTAACAAG--CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp2R1G1	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G2	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G3	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G4	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G5	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G6	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp2R1G7	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G8	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	

			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G10	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G11	S	Hetero	R	GATGCCTAACAAGC- T CAGGGGAG	
			D	GATGCCTAACAAGC- (-17 bps)	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G12	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G13	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G14	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G15	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G16	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G17	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G18	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	

	T	Hetero	D	GATGCCTAA-----AGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G19	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R1G20	S	Homo	W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Homo	W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type			W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2A1	S	Homo	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
	T	Homo	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp2R2A2	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R2A3	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp2R2A4	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp2R2A5	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	

Exp2R2A6	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCA CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp2R2A7	S	Homo	I	GATGCCTAACAAGCG CAGGGGAG	
			I	GATGCCTAACAAGCG CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCA CAGGGGAA	
Exp2R2A8	S	Homo	I	GATGCCTAACAAGCA CAGGGGAG	
			I	GATGCCTAACAAGCA CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp2R2A9	S	Hetero	I	GATGCCTAACAAGCA CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCA CAGGGGAA	
Exp2R2A10	S	Homo	I	GATGCCTAACAAGCA CAGGGGAG	
			I	GATGCCTAACAAGCA CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp2R2A11	S	Homo	D	GATGCCTAACAAGCA--GGGGAG	
			D	GATGCCTAACAAGCA--GGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCA CAGGGGAA	
Exp2R2A12	S	Homo	I	GATGCCTAACAAGCA CAGGGGAG	
			I	GATGCCTAACAAGCA CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCA CAGGGGAA	
			I	GATGCCTAACAAGCA CAGGGGAA	
Exp2R2A13	S	Homo	I	GATGCCTAACAAGCA CAGGGGAG	
			I	GATGCCTAACAAGCA CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCC CAGGGGAA	
			I	GATGCCTAACAAGCC CAGGGGAA	
Exp2R2A14	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCA CAGGGGAA	
			I	GATGCCTAACAAGCA CAGGGGAA	
Exp2R2A15	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	

			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			D	GATGCCTAACAAG -- CAGGGGAA	
Exp2R2A16	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp2R2A17	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp2R2A18	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp2R2A19	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			D	GATGCCTAACAAGC -- AGGGGAA	
Exp2R2A20	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			D	GATGCCTAACAAGC ---- GGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R2G1	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC C CAGGGGAA	
Exp2R2G2	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp2R2G3	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G4	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	

			I	GATGCCTAACAAGC A CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G5	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			D	GATGCCTAACA-----CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp2R2G6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G11	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G12	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R2G13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	

	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G14	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G15	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R2G16	S	Hetero	D	GATGCCTAACAAGC- C -GGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp2R2G17	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G18	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G19	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R2G20	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3A1	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	

			I	GATGCCTAACAAGC C AGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC C AGGGGAA	
			I	GATGCCTAACAAGC T AGGGGAA	
Exp2R3A2	S	Homo	D	G-----(-31 bps)	
			D	G-----(-31 bps)	
	T	Hetero	I	GATGCCTAACAAGC T AGGGGAA	
			D	GATGCCTAACAAGC--AGGGGAA	
Exp2R3A3	S	Homo	I	GATGCCTAACAAGC A AGGGGAG	
			I	GATGCCTAACAAGC A AGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T AGGGGAA	
			I	GATGCCTAACAAGC A AGGGGAA	
Exp2R3A4	S	Hetero	I	GATGCCTAACAAGC A AGGGGAG	
			I	GATGCCTAACAAGC G AGGGGAG	
	T	Homo	I	GATGCCTAACAAGC A AGGGGAA	
			I	GATGCCTAACAAGC A AGGGGAA	
Exp2R3A5	S	Hetero	I	GATGCCTAACAAGC T AGGGGAG	
			D	GATGCCTAACAAGC--AGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T AGGGGAA	
			I	GATGCCTAACAAGC A AGGGGAA	
Exp2R3A6	S	Homo	I	GATGCCTAACAAGC A AGGGGAG	
			I	GATGCCTAACAAGC A AGGGGAG	
	T	Homo	D	GATATTTGCGATGC--AGGGGAA	
			D	GATGCCTAACAAGC--AGGGGAA	
Exp2R3A7	S	Homo	I	GATGCCTAACAAGC T AGGGGAG	
			I	GATGCCTAACAAGC T AGGGGAG	
	T	Homo	I	GATGCCTAACAAGC A AGGGGAA	
			I	GATGCCTAACAAGC A AGGGGAA	
Exp2R3A8	S	Hetero	I	GATGCCTAACAAGC T AGGGGAG	
			D	GATGCCTAACAAGC--AGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T AGGGGAA	
			I	GATGCCTAACAAGC T AGGGGAA	
Exp2R3A9	S	Homo	I	GATGCCTAACAAGC T AGGGGAG	
			I	GATGCCTAACAAGC T AGGGGAG	
	T	Homo	I	GATGCCTAACAAGC A AGGGGAA	
			I	GATGCCTAACAAGC A AGGGGAA	
Exp2R3A10	S	Homo	I	GATGCCTAACAAGC A AGGGGAG	
			I	GATGCCTAACAAGC A AGGGGAG	

	T	Homo	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp2R3A11	S	Hetero	I	GATGCCTAACAAGC (+641 bps)	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	D	GATGCCTAACAAGC -- AGGGGAA	
			D	GATGCCTAACAAGC -- AGGGGAA	
Exp2R3A12	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R3A13	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	D	GATGCCTAACAAGC -- AGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R3A14	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R3A15	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	D	GATGCCTAACAAGC -- AGGGGAA	
			D	GATGCCTAACAAGC -- AGGGGAA	
Exp2R3A16	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
Exp2R3A17	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R3A18	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp2R3A19	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC C CAGGGGAA	

			I	GATGCCTAACAAGCTCAGGGGAA	
Exp2R3A20	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp2R3G1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G3	S	Hetero	I	GATGCCTAACAAGCCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G5	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G7	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	

			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G11	S	Hetero	W	GATGCCTAACAAGC--AGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G12	S	Homo	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R3G13	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G14	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp2R3G15	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G16	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp2R3G17	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			D	GATGCCTAACAAGC-(-11 bps)	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	

Exp2R3G18	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC C CAGGGGAA	
			I	GATGCCTAACAAGC C CAGGGGAA	
Exp2R3G19	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp2R3G20	S		W	GATGCCTAACAAGC-CAGGGGAG	
		Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1A1	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	D	GATGCCTAACAAGC -- AGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R1A2	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp3R1A3	S	Hetero	D	GATGCCTAACAAGC - (-7 bps)	
			I	GATGCCTAACAAGC C CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			D	GATGCCTAACAAGC -- AGGGGAA	
Exp3R1A4	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC C CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp3R1A5	S	Hetero	I	GATGCCTAACAAGC (+160 bps)	
			I	GATGCCTAACAAGC G CAGGGGAG	

	T	Hetero	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGC (+43 bps)	
Exp3R1A6	S	Hetero	I	GATGCCTAACAAGCGCAGGGGAG	
			D	GATGCCTAACAAGC--AGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1A7	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1A8	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp3R1A9	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1A10	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1A11	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp3R1A12	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp3R1A13	S	Hetero	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1A14	S	Hetero	I	GATGCCTAACAAGC (+82 bps)	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCGCAGGGGAA	

			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1A15	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp3R1A16	S	Hetero	D	GATGCCTAACAAGC--AGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1A17	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1A18	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp3R1A19	S	Hetero	D	GATGCCTAACAAGC--AGGGGAG	
			D	GATGCCTAACAAGC---GGGGAG	
	T	Hetero	I	GATGCCTAACAAGCCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1A20	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCGCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1G1	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1G2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	

			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G4	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			D	GATGCCTAACAAGC (-11 bps)	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1G5	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R1G6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G8	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G10	S	Hetero	I	GATGCCTAACAAGCCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G11	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G12	S	Hetero	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	

Exp3R1G13	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G14	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R1G15	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G16	S	Hetero	I	GATGCCTAACAAGC C CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R1G17	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G18	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G19	S	Homo	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R1G20	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Sample no.	Genome type	genotype	Mutation type	NtPDS sequences	
------------	-------------	----------	---------------	-----------------	--

Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2A1	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R2A2	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R2A3	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R2A4	S	Hetero	D	GATGCCTAACAAGC---GGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp3R2A5	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	D	GATGCCTAACAA---CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R2A6	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCGCAGGGGAA	
Exp3R2A7	S	Hetero	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R2A8	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R2A9	S	Hetero	I	GATGCCTAACAAGCCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	

	T	Hetero	I	GATGCCTAACAAGC C CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R2A10	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R2A11	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp3R2A12	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R2A13	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp3R2A14	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC GTGGGCCGA	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R2A15	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC (+213 bps)	
Exp3R2A16	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R2A17	S	Homo	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R2A18	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	

			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R2A19	S	Homo	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R2A20	S	Homo	D	GATGCCTAACAAGC--AGGGGAG	
			D	GATGCCTAACAAGC--AGGGGAG	
	T	Homo	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R2G1	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R2G2	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R2G3	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R2G4	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G5	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			D	GATGCCTAACA-----GGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R2G6	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R2G7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	

			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G8	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G11	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R2G12	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC C CAGGGGAA	
Exp3R2G13	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G14	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G15	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G16	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Exp3R2G17	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G18	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G19	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R2G20	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			D	GATGCCTAACA ---- CAGGGGAA	

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3A1	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC C CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
Exp3R3A2	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp3R3A3	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R3A4	S	Homo	I	GATGCCTAACAAGC A CAGGGGAG	

			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC G CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R3A5	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp3R3A6	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R3A7	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp3R3A8	S	Hetero	I	GATGCCTAACAAGC C CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp3R3A9	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Homo	I	GATGCCTAACAAGC T CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
Exp3R3A10	S	Hetero	I	GATGCCTAACAAGC C CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp3R3A11	S	Hetero	D	GATGCCTAACAAGC -- AGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGA	
Exp3R3A12	S	Hetero	I	GATGCCTAACAAGC C CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
Exp3R3A13	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	

	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R3A14	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R3A15	S	Hetero	I	GATGCCTAACAAGCACAGGGGAG	
			D	GATGCCTAACAA-----GGGGAG	
	T	Homo	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R3A16	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R3A17	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R3A18	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Homo	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R3A19	S	Homo	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R3A20	S	Homo	I	GATGCCTAACAAGCACAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGCACAGGGGAA	
			I	GATGCCTAACAAGCAAGGGGAA	
Exp3R3G1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	

	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G3	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGCTCAGGGGAA	
Exp3R3G4	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R3G5	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R3G6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G7	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			D	GATGCCTAACAAGC--AGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G8	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
Exp3R3G9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G10	S	Hetero	W	GATGCCTAACAAGC-GCAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R3G11	S	Hetero	I	GATGCCTAACAAGCGCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	

			I	GATGCCTAACAAGCACAGGGGAA	
Exp3R3G12	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G13	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G14	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G15	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G16	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G17	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G18	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCACAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G19	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
Exp3R3G20	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			D	GATGCCTAACAAGC-(-17 bps)	
	T	Hetero	I	GATGCCTAACAAGCTCAGGGGAA	
			I	GATGCCTAACAAGCACAGGGGAA	

Supplemental Data 5. The *NtPDS* sequences of Supplemental Figure 17. Red box in Sample no. column indicates four copies of *NtPDS* have same mutations. S: *sylvestris* form. T: *tomentosiformis* form. Orange box-Homo: homozygous in this form. Light blue box-hetero: heterozygous in this form. W: wild type. I: insertion. D: deletion. Letter in red: the mutated nucleotide.

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	

	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R1-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R1-1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R1-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R1-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R1-4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R1-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R1-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	

			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R1-7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R1-8	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
5 µg R1-9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R1-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R1-1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R1-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R1-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R1-4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R1-5	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

10 µg R1-6	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCAAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
10 µg R1-7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R1-8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R1-9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R1-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R1-1	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			D	GATGCCTAACAAGC---GGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
20 µg R1-2	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			W	GATGCCTAACAAGC--AGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCTCAGGGGAA	
20 µg R1-3	S	Hetero	I	GATGCCTAACAAGCCAGGGGAG	
			I	GATGCCTAACAAGCTCAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R1-4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R1-5	S	Hetero	I	GATGCCTAACAAGCAAGGGGAG	

			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R1-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Hetero	I	GATGCCTAACAAGC A CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R1-7	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R1-8	S	Hetero	D	GATGCCTAA-----CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
20 µg R1-9	S	Hetero	I	GATGCCTAACAAGC A CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			D	GATGCCTAACAAGC T CAGGGGAA	
20 µg R1-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Sample no.	Genome type	genotype	Mutation type	<i>NtPDS</i> sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	

			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R2-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R2-1	S	Homo	I	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

5 µg R2-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R2-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R2-4	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC (+153 bps)	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC A CAGGGGAA	
5 µg R2-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R2-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R2-7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R2-8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R2-9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R2-10	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC G CAGGGGAA	
10 µg R2-1	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	

			I	GATGCCTAACAAGCACAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R2-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R2-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R2-4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R2-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R2-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R2-7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R2-8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R2-9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R2-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	

	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R2-1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R2-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R2-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R2-4	S	Hetero	I	GATGCCTAACAAGC T CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			D	GATGCCTAACAAGC GTC (-7 bps)	
20 µg R2-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R2-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R2-7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R2-8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R2-9	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	

			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R2-10	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAA	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	

Sample no.	Genome type	genotype	Mutation type	NtPDS sequences	
Wild type	S		W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-7	S		W	GATGCCTAACAAGC-CAGGGGAG	

			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
0 µg R3-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R3-1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCAGGGGAA	
5 µg R3-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R3-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R3-4	S	Hetero	I	GATGCCTAACAAGCTCAGGGGAG	
			I	GATGCCTAACAAGCGCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCAGGGGAA	
5 µg R3-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R3-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	

	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R3-7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC-T CAGGGGAA	
5 µg R3-8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R3-9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			D	GATGCCTAACAAGC--AGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
5 µg R3-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R3-1	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R3-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R3-3	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R3-4	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R3-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	

			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R3-6	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC G CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R3-7	S	Homo	I	GATGCCTAACAAGC T CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	
10 µg R3-8	S	Hetero	W	GATGCCTAACAAGC-CAGGGGAG	
			I	GATGCCTAACAAGC A CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R3-9	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
10 µg R3-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R3-1	S	Hetero	I	GATGCCTAACAAGC C CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R3-2	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R3-3	S	Hetero	I	GATGCCTAACAAGC C CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R3-4	S	Hetero	I	GATGCCTAACAAGC G CAGGGGAG	
			I	GATGCCTAACAAGC T CAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGC T CAGGGGAA	

20 µg R3-5	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R3-6	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R3-7	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R3-8	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	
20 µg R3-9	S	Hetero	I	GATGCCTAACAAGCAGGGGAG	
			I	GATGCCTAACAAGCCAGGGGAG	
	T	Hetero	W	GATGCCTAACAAGC-CAGGGGAA	
			I	GATGCCTAACAAGCAGGGGAA	
20 µg R3-10	S		W	GATGCCTAACAAGC-CAGGGGAG	
			W	GATGCCTAACAAGC-CAGGGGAG	
	T		W	GATGCCTAACAAGC-CAGGGGAA	
			W	GATGCCTAACAAGC-CAGGGGAA	