



Historical baselines of coral cover on tropical reefs as estimated by expert opinion

Tyler D. Eddy¹, William W.L. Cheung¹ and John F. Bruno²

¹ Nippon Foundation—Nereus Program, Institute for the Oceans & Fisheries, University of British Columbia, Vancouver, British Columbia, Canada

² Department of Biological Sciences, University of North Carolina at Chapel Hill, Chapel Hill, NC, United States of America

ABSTRACT

Coral reefs are important habitats that represent global marine biodiversity hotspots and provide important benefits to people in many tropical regions. However, coral reefs are becoming increasingly threatened by climate change, overfishing, habitat destruction, and pollution. Historical baselines of coral cover are important to understand how much coral cover has been lost, e.g., to avoid the ‘shifting baseline syndrome’. There are few quantitative observations of coral reef cover prior to the industrial revolution, and therefore baselines of coral reef cover are difficult to estimate. Here, we use expert and ocean-user opinion surveys to estimate baselines of global coral reef cover. The overall mean estimated baseline coral cover was 59% ($\pm 19\%$ standard deviation), compared to an average of 58% ($\pm 18\%$ standard deviation) estimated by professional scientists. We did not find evidence of the shifting baseline syndrome, whereby respondents who first observed coral reefs more recently report lower estimates of baseline coral cover. These estimates of historical coral reef baseline cover are important for scientists, policy makers, and managers to understand the extent to which coral reefs have become depleted and to set appropriate recovery targets.

Subjects Conservation Biology, Ecology, Environmental Sciences, Marine Biology, Climate Change Biology

Keywords Historical ecology, Marine environmental history, Qualitative survey, Expert opinion survey, Historical baselines, Shifting baseline syndrome, Climate change

INTRODUCTION

Since the publication of the seminal article, ‘Anecdotes and the shifting baseline syndrome’ (*Pauly, 1995*), it has been recognized that it is important to understand what was pristine prior to large, human-induced changes to the natural environment (*Jackson et al., 2001; Lotze et al., 2006*). Historical baselines are important for scientists, managers, policy makers, and the general public to understand how much a population or habitat has been impacted by humans through time (*Johannes, 1981; Rosenberg et al., 2005; Eddy, Gardner & Pérez-Matus, 2010; McClenachan, Ferretti & Baum, 2012; Thurstan et al., 2015*). Baselines can also provide realistic and meaningful targets for restoration (*Pitcher, 2005*). Yet determining a natural baseline is often difficult, in part because there is little data on which to base estimates, and also because different approaches can produce very

Submitted 11 September 2017

Accepted 10 January 2018

Published 24 January 2018

Corresponding author

Tyler D. Eddy, t.eddy@oceans.ubc.ca

Academic editor

Richard Taylor

Additional Information and
Declarations can be found on
page 11

DOI 10.7717/peerj.4308

© Copyright
2018 Eddy et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

different estimates ([Bruno et al., 2014](#)). Without them, the perception of what was pristine or natural has been shown to change with younger generations, as the reference of their earliest memories is different than that of their parents or grandparents ([Sáenz-Arroyo et al., 2005](#); [McClenachan, 2009](#)).

Like many marine habitats, coral reefs have been drastically altered by human activities such as habitat destruction, nutrient and/or sediment runoff, overfishing, and increased frequency of bleaching events due to climate change. These impacts have resulted in significant declines in coral cover around the world ([Bruno et al., 2009](#)), including in the Caribbean Sea ([Gardner & Co, 2003](#); [Schutte, Selig & Bruno, 2010](#)), and the Pacific Ocean ([Gomez, Alcala & San Diego, 1981](#); [Bellwood et al., 2004](#); [Bruno & Selig, 2007](#)). While it has been suggested that the number of people living within proximity of coral reefs explains the degree of coral degradation ([Smith et al., 2016](#)), recent research suggests that global scale actions are required to prevent further loss ([Bruno & Valdivia, 2016](#)). The loss of coral reefs has implications for tropical and subtropical coastal societies globally, as they provide habitat for species that are important for subsistence and indigenous fisheries ([Cisneros-Montemayor et al., 2016](#)).

Coral cover, the percentage of the seafloor occupied by living scleractinian corals, is an important metric of reef state or “health”. Coral cover is related to habitat volume and complexity and is a good predictor of fish and invertebrate diversity and abundance ([Jones et al., 2004](#); [Idjadi & Edmunds, 2006](#); [Alvarez-Filip et al., 2009](#); [Dustan, Doherty & Pardede, 2013](#)). Coral cover also influences reef accretion; accretion rates generally decline with coral cover and on low cover reefs a greater portion of the carbonate reef structure is exposed to bioerosion (e.g., via urchin and parrotfish grazing) accelerating the loss of the reef framework ([Perry et al., 2013](#); [Kuffner & Toth, 2016](#)).

Syntheses of quantitative benthic reef surveys (i.e., based on *in situ* assessments, video or still images) suggest coral cover began to decline in most regions in the late 1970s—early 1980s ([Gardner & Co, 2003](#); [Bruno & Selig, 2007](#); [Schutte, Selig & Bruno, 2010](#)). Unfortunately, using published estimations of coral coverage to extrapolate global trends may suffer from reporting biases. Particularly, reef monitoring only became common after noticeable coral decline had already begun: even at regional scales, the number of sites surveyed annually was low (<10) until relatively recently, e.g., until the mid 1980s for the Great Barrier Reef and the late 1990s for the Greater Caribbean ([De’ath et al., 2012](#); [Jackson et al., 2014](#); [Schutte, Selig & Bruno, 2010](#); [Gardner & Co, 2003](#); [Bellwood et al., 2004](#)). Although there are early quantitative estimates of coral cover from the late 1960s and early 1970s, only a very small number of quantitative surveys were performed globally in any given year ([De’ath et al., 2012](#); [Bruno & Selig, 2007](#); [Jackson et al., 2014](#); [Schutte, Selig & Bruno, 2010](#); [Gardner & Co, 2003](#); [Bellwood et al., 2004](#)). Additionally, bias in site selection during this period could have influenced study outcomes. For example, scientists could have chosen reefs with extraordinary cover (or other characteristics) for aesthetic reasons. On the other hand, disturbance ecology was a popular research theme in the 1970s and the purpose of many of the quantitative surveys from the era was to document coral loss or low cover due to storms and predator outbreaks (and thus may have been biased against surveying high cover reefs). Overall, the inferences from long-term monitoring

studies about coral cover and other characteristics of reefs in the absence of anthropogenic disturbances are unfortunately limited.

In the absence of systematic quantitative surveys predating human impacts on coral reefs, there have been a number of attempts to understand historical baselines of coral reefs. Although paleoecology has been applied for decades to measure changes in coral composition, e.g., in response to natural and anthropogenic disturbances ([Aronson & Precht, 2001](#); [Pandolfi et al., 2003](#); [Wapnick, Precht & Aronson, 2004](#); [Pandolfi & Jackson, 2006](#)), this approach does not provide estimates of living coral cover. There have also been attempts to use photographic images to quantify coral reef percent cover prior to quantitative coral surveys ([Clark et al., 2016](#)); however, without a systematic survey, it is difficult to draw accurate conclusions about baselines of coral reef cover. An integrative approach combined archaeological deposits, ethnohistoric and anecdotal descriptions, and modern ecological and fishery data to evaluate the health of Hawaiian coral reefs over 700 years to document periods of reef decline and recovery ([Kittinger et al., 2011](#)). Interestingly, a recent study used navigational charts going back over 240 years to document a 52% decrease in patch reef cover in lagoons of the Florida Keys ([McClenachan et al., 2017](#)).

In the absence of quantitative data, drawing experience from coral reef experts through questionnaire surveys can help qualitatively estimate baselines. The “Reef Reminiscences” project gathered the memories of reef scientists of their work in the field half a century ago, painting a powerful picture of the state of reefs before most of their colleagues were born ([Sale & Szman, 2012](#)). There have been approaches in the fisheries literature to reconstruct baselines of fish stocks using historical resources to document declines that were much greater than previously understood ([Rosenberg et al., 2005](#)). Additionally, fishers’ ecological knowledge has been used to estimate the baselines of various fish stocks and fisheries (e.g., [Sadovy & Cheung, 2003](#); [Teh et al., 2007](#); [Eddy, Gardner & Pérez-Matus, 2010](#)). [Kleypas & Eakin \(2007\)](#) surveyed reef scientists to measure their opinions about the relative importance of different threats to coral reef ecosystems. Also, expert opinion has been used to quantify the relative importance of different threats to different components of marine environment where information is limited, such as for the deep sea ([MacDiarmid et al., 2012](#); [Eddy, 2014](#)), and to evaluate the effectiveness of fisheries management globally ([Mora et al., 2009](#)).

The purpose of this study was to estimate the coral cover baseline by assessing the opinions of coral reef scientists. We asked each participant what they believed “baseline coral cover” is. Our intent was that this value would be based on their own observations (which at the least would likely subconsciously bias their answer), their reading and interpretation of the literature, their discussions with colleagues, etc. We defined “baseline coral cover” as the global average pre-human impact, in shallow (1–15 m depth) fore reef environments. We also asked each participant questions related to their expertise and profession, level of experience, and the timing of their earliest field observations.

METHODS

To solicit as many as possible expert and ocean-user opinions about baselines of coral reefs, we designed a short, online survey that included four questions ([Table 1](#)). The survey

Table 1 Questions asked in online survey. See ‘Methods’ for further details that were provided to participants.

	Question
1	What is your position?
2	What was the first year that you observed a coral reef?
3	What is the highest percent cover of a coral reef that you have observed? Where and when was this?
4	What is your baseline estimate of coral reef cover?

was distributed via Twitter and the NOAA coral-list and remained open for three months (March 2–May 5, 2017). In both avenues, coral reef scientists, students studying coral reefs, managers, policy makers or NGO employees working in areas with coral reefs, and recreational divers who had dove on coral reefs were asked to fill out a quick survey. Due to the online nature of the survey, respondents were biased to those having access to email or the internet and are participants of the NOAA email coral list or the coral reef Twitter community or heard about it via word of mouth from someone who is. The Office of Human Research Ethics at the University of North Carolina at Chapel Hill determined that this submission did not constitute human subjects research as defined under federal regulations (45 CFR 46.102 (d or f) and 21 CFR 56.102(c)(e)(l)) and did not require IRB approval as communicated to JFB via email on March 1st, 2017 (Study # 17-0479).

The first question asked the respondent about their job type in order to differentiate professional scientists, managers, policy makers, and non-governmental organization (NGO) employees, students, and recreational divers. The second question asked what was the first year that the respondent observed a coral reef (to estimate the level of experience and the personal baseline reference). The third question asked what was the highest coral cover observed and where and when this occurred.

The fourth question asked for an estimation of baseline percent coral cover. In this question, we stated: *“What would the average coral cover be in the absence of any human impacts in shallow (1–15 m depth) fore reef environments?”* We specified that *this is not the same as the maximum achievable coverage; even before humans, reefs were disturbed by storms, disease, etc. and thus some proportion of reefs would have been in various stages of recovery and not at peak coral cover.* We indicated that *“this value surely varies among regions, habitats, depths, etc; but still, we’d like your help in developing a consensus about what the historical average was. And by “human impacts” we mean both local and global impacts including fishing, pollution, and ocean warming. Finally, we are assuming that you will integrate your own observations with your knowledge of the literature, discussions with colleagues, etc.”*

We then summarized the responses by type of participant and grouped the location of the highest coral cover observed by region using joy or ridge line plots in R. The ggjoy package version 0.3.0 was used to compute and draw a kernel density estimate (<https://github.com/clauswilke/ggjoy>). We also plotted estimates of historical baseline coral cover and highest coral cover observed as a function of time to test for evidence




Figure 1 First year that a coral reef was observed by survey respondent position. Distribution of responses for first year that a coral reef was observed by survey respondent position. Colors are used to visually differentiate among positions. Numbers of respondents by position: professional scientists = 133, students = 45, managers, policy makers or NGO employees = 12, recreational divers (rec divers) = 5.

Full-size [DOI: 10.7717/peerj.4308/fig-1](https://doi.org/10.7717/peerj.4308/fig-1)

of shifting baselines. Trend lines were calculated using the loess non-parametric local regression smoothing method in R.

To situate our baseline coral cover estimate alongside coral cover reported in quantitative surveys, we extracted published coral cover survey data for the earliest years from each study and plotted alongside our baseline estimate. Survey data sources were: (*De'ath et al., 2012*; *Bruno & Selig, 2007*; *Jackson et al., 2014*; *Schutte, Selig & Bruno, 2010*; *Gardner & Co, 2003*; *Bellwood et al., 2004*). We have not plotted quantitative baselines for the other regions reported in our qualitative survey—Atlantic Ocean, Persian Gulf, and Red Sea; although there are some surveys for locations within each region (ex. Brazil in the Atlantic, Egypt in the Red Sea, and Bahrain in the Persian Gulf), it is unclear if these individual locations are representative of the wider region.

RESULTS

Types of respondents

We received a total of 195 responses from 133 professional scientists (68%), 45 students (23%), 12 managers, policy makers or NGO employees (6%), and five recreational divers (3%).

First year coral reef observed

The earliest year that a coral reef was observed was listed at 1960, the average year was 1992, while the latest year listed was 2016 ([Fig. 1](#)). For scientists, the earliest year was also listed as 1960, the average was 1988, and the latest year was 2013.




Figure 2 Highest coral cover observed. Distribution of responses for highest coral cover observed by (A) survey respondent position and (B) region. Colors are used to visually differentiate among positions. Numbers of respondents by position: professional scientists = 133, students = 45, managers, policy makers or NGO employees = 12, recreational divers (rec divers) = 5. Number of respondents by region: Atlantic Ocean = 5, Caribbean = 54, Indian Ocean = 13, Pacific Ocean = 109, Persian Gulf = 4, Red Sea = 8.

Full-size [DOI: 10.7717/peerj.4308/fig-2](https://doi.org/10.7717/peerj.4308/fig-2)

Highest coral cover observed

The average highest percent cover reported was $79\% \pm 21\%$ SD (Fig. 2A). The earliest year that the highest percent cover observed by a respondent was 1962, the latest year reported was 2017. For scientists, the average highest coral percent cover was $82\% \pm 18\%$ SD (Fig. 2A). The earliest year that the highest percent cover was observed was 1970, the latest year was also 2017, and the average highest percent cover observed in 2001.

Five respondents listed having observed the highest coral cover at locations in the Atlantic Ocean, 54 in the Caribbean, 13 in the Indian Ocean, 109 in the Pacific Ocean, four in the Persian Gulf, and eight in the Red Sea. Two respondents did not list a location. The highest average coral cover was observed at locations in the Pacific Ocean at 87%, followed by the Indian Ocean at 79%, 76% in the Red Sea, 75% in the Persian Gulf, 63% in the Caribbean Sea, and 45% in the Atlantic Ocean (Fig. 2B).

Baseline estimates

The mean estimate of baseline coral cover was $59\% \pm 19\%$ SD, while for scientists it was $58\% \pm 18\%$ SD (Fig. 3A). Interestingly, there were differences in baseline coral cover estimations according to the location that respondents had observed the highest coral cover. Respondents who indicated a location in the Red Sea as being the place where they had observed the highest percent cover estimated the highest baseline cover, with a mean of 77% (Fig. 3B). This was followed by an average estimate of 62% coral cover for the Indian Ocean, 61% for the Pacific Ocean, 53% for the Caribbean Sea, 50% for the Persian Gulf, and 48% for the Atlantic Ocean (Fig. 3B).

Shifting baselines

We found no evidence of shifting baselines of coral reef cover as there is not a decline in the estimates of coral baselines with more recent first year of coral observations, and there appears to be a slight increase in baseline estimates over time (Fig. 4A). There was, however a decreasing trend in the highest coral cover observed as a function of time—which was represented by the first year of coral observation (Fig. 4B). These results indicate general decrease in highest observed coral coverage, but not the baseline.

Qualitative vs. quantitative baselines

Comparing the results from our baseline coral cover survey to published quantitative coral cover surveys dating back to the earliest years, we found that all quantitative surveys had lower values of coral cover (Fig. 5). The highest reported quantitative baseline value of 50% coral cover was reported from the Caribbean in 1977, while the lowest value was 28% from the Great Barrier reef in 1985 (Fig. 5).

DISCUSSION

Our expert opinion survey estimated global baseline coral cover at 59%, or 58% if only considering responses from non-student scientists. Baseline estimates varied depending on the region that respondents had observed the highest coral cover, such that estimates were the highest if a respondent had observed the highest coral cover in the Pacific, and lowest if observed in the Atlantic (Fig. 2B). These observations are in line with the ecology of the regions, as plating acroporid corals which can produce reefs with very high cover are not found in the Caribbean and some other regions. The global average baseline estimate from the survey appears to be biased by the proportion of respondents who reported to have observed corals in the Pacific who estimated a larger decline in coral cover relative to those




Figure 3 Estimates of baseline coral reef cover. Distribution of responses for expert opinion baseline estimates of coral reef cover by (A) survey respondent position and (B) according to the region where the respondent reported their highest observed coral cover. Colors are used to visually differentiate among positions. Numbers of respondents by position: professional scientists = 133, students = 45, managers, policy makers or NGO employees = 12, recreational divers (rec divers) = 5. Number of respondents by region: Atlantic Ocean = 5, Caribbean = 54, Indian Ocean = 13, Pacific Ocean = 109, Persian Gulf = 4, Red Sea = 8.

Full-size [DOI: 10.7717/peerj.4308/fig-3](https://doi.org/10.7717/peerj.4308/fig-3)

reported to have observed corals in other regions. This is consistent with a previous survey that asked respondents to estimate baselines for the Indo-Pacific and the Caribbean in 2013, that also reported differences between the regions (Bruno, 2013). For the Caribbean, 40.4% of respondents estimated baseline coral percent cover between 25–50%, 38.4% of respondents estimated 50–75% cover, 17.7% estimated <25% cover, and 4.9% estimated >75% cover (Bruno, 2013). For the Indo-Pacific, 43.5% of respondents estimated coral




Figure 4 Baseline estimates and highest coral cover observed over time. (A) Expert opinion estimates of baseline coral cover and (B) highest coral cover observed as a function of the first year that a respondent observed a coral reef. Box plots depict mean with standard error and tails show the minimum and maximum values. Red trend line indicates smoothed conditional mean with confidence interval in blue.

Full-size [DOI: 10.7717/peerj.4308/fig-4](https://doi.org/10.7717/peerj.4308/fig-4)

cover baseline between 50–75%, 29% estimated >75% cover, 26.1% estimated between 25–50%, while 3.4% estimated <25% cover (Bruno, 2013).

An additional source of variation among regions are the natural disturbances that can affect coral cover such as predator outbreaks and storms, hurricanes, and bleaching events. The presence and relative frequency and magnitude of these and other natural disturbances varies among reefs and regions. For example, cyclonic storms are absent within ~5% latitude of the equator and crown-of-thorns starfish (*Acanthaster planci*) are




Figure 5 Qualitative vs. quantitative baselines. Comparison of the expert opinion estimated global coral cover baseline from this study to the earliest coral cover surveys in different ocean regions. References for quantitative studies are found in 'Methods'. GBR, Great Barrier Reef.

Full-size DOI: [10.7717/peerj.4308/fig-5](https://doi.org/10.7717/peerj.4308/fig-5)

absent from the Caribbean. The degradation of some reefs appears to have begun centuries ago and it is very possible that human activities were already measurably affecting coral cover prior to the earliest field observations of our most experienced colleagues ([Wing & Wing, 2001](#); [Pandolfi et al., 2003](#)), therefore we could still be underestimating the baseline.

The mean expert opinion baseline estimate is lower than the average highest percent cover reported and also lower than some surveys from the 1970s (e.g., [Hughes, 1994](#); [Sheppard et al., 2017](#)). However, this seems reasonable because coral cover is reduced by natural disturbances including predators, storms, and diseases. Therefore, the baseline mean of a seascape or region would be substantially lower than the highest observed value, at least when integrated over time. Overestimating the cumulative impact (across larger scales of space and time than localized effects) of natural disturbances on coral cover could lead to an underestimate of the true baseline. On the other hand, it is possible that in the early years of reef science coral cover was atypically high, which could lead to an overestimate of the baseline. For example, [Woodley \(1992\)](#) argued that the high cover thickets of *Acropora* that dominated the reefs of Jamaica in the 1960s and 1970s were atypical and caused by an especially long lag in the return of large storms.

The lack of evidence for shifting baselines in survey respondents even with decrease in perceived coral coverage may indicate that spread of information/communication has generated consensus about historical baseline amongst the respondents. This may also suggest that the perceptions may be less independent among respondents. Either way, it appears that there is a general understanding among coral scientists and other surveyed groups that the baseline of coral reef percent cover was much greater than is generally reported today.

Expert opinion is a qualitative information source and will never replace quantitative data, however, in the absence of quantitative survey data dating to pre-industrial times, expert opinion is an invaluable resource. Recording the opinions of the relatively small number of scientists that worked on reefs when degradation was much less severe and widespread—from the 1950s to the early 1970s—should be considered a time sensitive priority ([Sale & Szmant, 2012](#)). There are of course, limitations to using expert opinion surveys, some of which we have addressed above related to geographical biases, as well as biases associated with recalled information from memory or personal experience which may be substantial ([Daw, 2010](#)). Additionally, we are unable to quantify the uncertainty associated with this historical baseline estimate of coral reef cover on tropical reefs. On the other hand, not only can expert opinion provide insight during time periods prior to quantitative data collection, they can also provide insight into parts of the ocean that have yet to be studied in depth, such as the deep sea ([MacDiarmid et al., 2012](#); [Eddy, 2014](#)).

CONCLUSIONS

Overall, we have provided an expert opinion estimate of the global coral percent cover baseline. While coral cover is presently declining at rapid rates around the world ([Selig, Casey & Bruno, 2012](#); [Bruno & Valdivia, 2016](#); [Hughes et al., 2017a](#); [Hughes et al., 2017b](#); [Hughes et al., 2018](#)), if we are capable of reversing these trends, it is important to understand what we should aim for as a target.

ACKNOWLEDGEMENTS

We thank everyone who participated in our survey and shared their knowledge, without which, this study would not have been possible. We also thank two anonymous reviewers and the editor for their comments.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the Nippon Foundation—The University of British Columbia Nereus Program. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Nippon Foundation—The University of British Columbia Nereus Program.

Competing Interests

John F. Bruno is an Academic Editor for PeerJ.

Author Contributions

- Tyler D. Eddy conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.

- William WL Cheung and John F. Bruno conceived and designed the experiments, performed the experiments, wrote the paper, reviewed drafts of the paper.

Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The Office of Human Research Ethics at the University of North Carolina at Chapel Hill, USA determined that this submission does not constitute human subjects research as defined under federal regulations [45 CFR 46.102 (d or f) and 21 CFR 56.102(c)(e)(l)] and does not require IRB approval (Study #: 17-0479).

Data Availability

The following information was supplied regarding data availability:

The raw survey data has been provided as a [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4308#supplemental-information>.

REFERENCES

- Alvarez-Filip L, Dulvy NK, Gill JA, Côté IM, Watkinson AR. 2009.** Flattening of Caribbean coral reefs: region-wide declines in architectural complexity. *Proceedings of the Royal Society B: Biological Sciences* **276**:3019–3025 DOI [10.1098/rspb.2009.0339](https://doi.org/10.1098/rspb.2009.0339).
- Aronson RB, Precht WF. 2001.** White-band disease and the changing face of Caribbean coral reefs. *Hydrobiologia* **460**:25–38 DOI [10.1023/A:1013103928980](https://doi.org/10.1023/A:1013103928980).
- Bellwood DR, Hughes TP, Folke C, Nystro M. 2004.** Confronting the coral reef crisis. *Nature* **429**(6994):827–833.
- Bruno JF. 2013.** Coral reef baselines survey: what do we think is natural? Available at https://figshare.com/articles/Coral_reef_baselines_survey_what_do_we_think_is_natural_/774627.
- Bruno JF, Precht WF, Vroom PS, Aronson RB. 2014.** Coral reef baselines: how much macroalgae is natural? *Marine Pollution Bulletin* **80**:24–29 DOI [10.1016/j.marpolbul.2014.01.010](https://doi.org/10.1016/j.marpolbul.2014.01.010).
- Bruno JF, Selig ER. 2007.** Regional decline of coral cover in the Indo-Pacific: timing, extent, and subregional comparisons. *PLOS ONE* **2**:e711 DOI [10.1371/journal.pone.0000711](https://doi.org/10.1371/journal.pone.0000711).
- Bruno JF, Sweatman H, Precht WF, Selig ER, Schutte VGW. 2009.** Assessing evidence of phase shifts from coral to macroalgal dominance on coral reefs. *Ecology* **90**:1478–1484 DOI [10.1890/08-1781.1](https://doi.org/10.1890/08-1781.1).
- Bruno JF, Valdivia A. 2016.** Coral reef degradation is not correlated with local human population density. *Scientific Reports* **6**:29778 DOI [10.1038/srep29778](https://doi.org/10.1038/srep29778).
- Cisneros-Montemayor AM, Pauly D, Weatherdon LV, Ota Y. 2016.** A global estimate of seafood consumption by coastal indigenous peoples. *PLOS ONE* **11**:e0166681 DOI [10.1371/journal.pone.0166681](https://doi.org/10.1371/journal.pone.0166681).

- Clark TR, Leonard ND, Zhao J, Brodie J, McCook LJ, Wachenfeld DR, Nguyen AD, Markham HL, Pandolfi JM. 2016. Historical photographs revisited: a case study for dating and characterizing recent loss of coral cover on the inshore Great Barrier Reef. *Scientific Reports* 6:19285 DOI 10.1038/srep19285.
- Daw TM. 2010. Shifting baselines and memory illusions: what should we worry about when inferring trends from resource user interviews?. *Animal Conservation* 13:534–535 DOI 10.1111/j.1469-1795.2010.00418.x.
- De’ath G, Fabricius KE, Sweatman H, Puotinen M. 2012. The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proceedings of the National Academy of Sciences of the United States of America* 109:17995–17999 DOI 10.1073/pnas.1208909109.
- Dustan P, Doherty O, Pardede S. 2013. Digital reef rugosity estimates coral reef habitat complexity. *PLOS ONE* 8:1–10 DOI 10.1371/journal.pone.0057386.
- Eddy TD. 2014. One hundred-fold difference between perceived and actual levels of marine protection in New Zealand. *Marine Policy* 46:61–67 DOI 10.1016/j.marpol.2014.01.004.
- Eddy TD, Gardner JPA, Pérez-Matus A. 2010. Applying fishers’ ecological knowledge to construct past and future lobster stocks in the Juan Fernández Archipelago, Chile. *PLOS ONE* 5:e13670 DOI 10.1371/journal.pone.0013670.
- Gardner TA, Co IM. 2003. Long-term region-wide declines in Caribbean corals. *Science* 301:958–960 DOI 10.1126/science.1086050.
- Gomez ED, Alcala AC, San Diego AC. 1981. Status of Philippine coral reefs—1981. *Proceedings of the Fourth International Coral Reef Symposium* 1:275–282.
- Hughes TP. 1994. Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef. *Science* 265:1547–1551 DOI 10.1126/science.265.5178.1547.
- Hughes TP, Anderson KD, Connolly SR, Heron SE, Kerry JT, Lough JM, Baird AH, Baum JK, Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H, Hobbs JPA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Pandolfi JM, Pratchett M, Schoepf V, Torda G, Wilson SK. 2018. Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science* 359(6371):80–83 DOI 10.1126/science.aan8048.
- Hughes TP, Barnes ML, Bellwood DR, Cinner JE, Cumming GS, Jackson JBC, Kleypas J, Van de Leemput IA, Lough JM, Morrison TH, Palumbi SR, Van Nes EH, Scheffer M. 2017b. Coral reefs in the Anthropocene. *Nature* 546:82–90 DOI 10.1038/nature22901.
- Hughes TP, Kerry J, Álvarez-Noriega M, Álvarez-Romero J, Anderson K, Baird A, Babcock R, Beger M, Bellwood D, Berkelmans R, Bridge T, Butler I, Byrne M, Cantin N, Comeau S, Connolly S, Cumming G, Dalton S, Diaz-Pulido G, Eakin CM, Figueira W, Gilmour J, Harrison H, Heron S, Hoey AS, Hobbs J-P, Hoogenboom M, Kennedy E, Kuo C-Y, Lough J, Lowe R, Liu G, Malcolm McCulloch HM, McWilliam M, Pandolfi J, Pears R, Pratchett M, Schoepf V, Simpson T, Skirving W, Sommer B, Torda G, Wachenfeld D, Willis B, Wilson S.

- 2017a. Global warming and recurrent mass bleaching of corals. *Nature* **543**:373–377 DOI [10.1038/nature21707](https://doi.org/10.1038/nature21707).
- Idjadi JA, Edmunds PJ. 2006. Scleractinian corals as facilitators for other invertebrates on a Caribbean reef. *Marine Ecology Progress Series* **319**:117–127 DOI [10.3354/meps319117](https://doi.org/10.3354/meps319117).
- Jackson JBC, Donovan MK, Cramer KL, Lam VV. 2014. Status and trends of Caribbean coral reefs: 1970–2012. In: *Global coral reef monitoring network*. Gland: IUCN.
- Jackson JBC, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, Bradbury RH, Cooke R, Erlandson J, Estes JA, Hughes TP, Kidwell S, Lange CB, Lenihan HS, Pandolfi JM, Peterson CH, Steneck RS, Tegner MJ, Warner RR. 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* **293**:629–638 DOI [10.1126/science.1059199](https://doi.org/10.1126/science.1059199).
- Johannes JE. 1981. *Words of the lagoon: fishing and marine lore in the Palau district of Micronesia*. Berkeley: University of California Press, 207 pp.
- Jones GP, McCormick MI, Srinivasan M, Eagle JV. 2004. Coral decline threatens fish biodiversity in marine reserves. *Proceedings of the National Academy of Sciences of the United States of America* **101**:8251–8253 DOI [10.1073/pnas.0401277101](https://doi.org/10.1073/pnas.0401277101).
- Kittinger JN, Pandolfi JM, Blodgett JH, Hunt TL, Jiang H, Maly K, McClenachan LE, Schultz JK, Wilcox BA. 2011. Historical reconstruction reveals recovery in Hawaiian coral reefs. *PLOS ONE* **6**(10):e25460 DOI [10.1371/journal.pone.0025460](https://doi.org/10.1371/journal.pone.0025460).
- Kleypas JA, Eakin CM. 2007. Scientists' perceptions of threats to coral reefs: results of a survey of coral reef researchers. *Bulletin of Marine Science* **80**:419–436 DOI [10.3354/esr00167](https://doi.org/10.3354/esr00167).
- Kuffner IB, Toth LT. 2016. A geological perspective on the degradation and conservation of western Atlantic coral reefs. *Conservation Biology* **30**:706–715 DOI [10.1111/cobi.12725](https://doi.org/10.1111/cobi.12725).
- Lotze HK, Lenihan HS, Bourque BJ, Bradbury RH, Cooke RG, Kay MC, Kidwell SM, Kirby MX, Peterson CH, Jackson JBC. 2006. Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science* **312**:1806–1809 DOI [10.1126/science.1128035](https://doi.org/10.1126/science.1128035).
- MacDiarmid A, McKenzie A, Sturman J, Beaumont J, Mikaloff-Fletcher S, Dunne J. 2012. Assessment of anthropogenic threats to New Zealand marine habitats. New Zealand aquatic environment and biodiversity report no. 93.
- McClenachan L. 2009. Historical declines of goliath grouper populations in South Florida, USA. *Marine Ecology Progress Series* **7**:175–181.
- McClenachan L, Ferretti F, Baum JK. 2012. From archives to conservation: why historical data are needed to set baselines for marine animals and ecosystems. *Conservation Letters* **5**:349–359 DOI [10.1111/j.1755-263X.2012.00253.x](https://doi.org/10.1111/j.1755-263X.2012.00253.x).
- McClenachan L, O'Connor G, Neal BP, Pandolfi JM, Jackson JBC. 2017. Ghost reefs: nautical charts document large spatial scale of coral reef loss over 240 years. *Science Advances* **3**:e1603155 DOI [10.1126/sciadv.1603155](https://doi.org/10.1126/sciadv.1603155).

- Mora C, Myers RA, Coll M, Libralato S, Pitcher TJ, Sumaila RU, Zeller D, Watson R, Gaston KJ, Worm B. 2009. Management effectiveness of the world's marine fisheries. *PLOS Biology* 7(6):e1000131 DOI 10.1371/journal.pbio.1000131.
- Pandolfi JM, Bradbury RH, Sala E, Hughes TP, Bjorndal KA, Cooke RG, McArdle D, McClenachan L, Newman MJH, Paredes G, Warner RR, Jackson JBC. 2003. Global trajectories of the long-term decline of coral reef. *Science* 301:955–958 DOI 10.1126/science.1085706.
- Pandolfi JM, Jackson JBC. 2006. Ecological persistence interrupted in Caribbean coral reefs. *Ecology Letters* 9:818–826 DOI 10.1111/j.1461-0248.2006.00933.x.
- Pauly D. 1995. Anecdotes and the shifting baseline syndrome of fisheries. *Trends in Ecology & Evolution* 10:430 DOI 10.1016/S0169-5347(00)89171-5.
- Perry CT, Murphy GN, Kench PS, Smithers SG, Edinger EN, Steneck RS, Mumby PJ. 2013. Caribbean-wide decline in carbonate production threatens coral reef growth. *Nature Communications* 4:Article 1402 DOI 10.1038/ncomms2409.
- Pitcher TJ. 2005. Back-to-the-future: a fresh policy initiative for fisheries and a restoration ecology for ocean ecosystems. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 360:107–121 DOI 10.1098/rstb.2004.1575.
- Rosenberg A, Bolster WJ, Alexander KE, Leavenworth WB, Cooper AB, McKenzie MG. 2005. The history of ocean resources: modeling cod biomass using historical records. *Frontiers in Ecology and the Environment* 2:83–90 DOI 10.1890/1540-9295(2005)003[0078:THOORM]2.0.CO;2.
- Sadovy YJ, Cheung WL. 2003. Near extinction of a highly fecund fish: the one that nearly got away. *Fish & Fisheries* 4:86–99 DOI 10.1046/j.1467-2979.2003.00104.x.
- Sáenz-Arroyo A, Roberts CM, Torre J, Cariño-Olvera M, Enríquez-Andrade RR. 2005. Rapidly shifting environmental baselines among fishers of the Gulf of California. *Proceedings of the Royal Society B*: 272:1957–1962 DOI 10.1098/rspb.2005.3175.
- Sale PF, Szmant AM. 2012. *Reef reminiscences: ratcheting back the shifted baselines concerning what reefs used to be*. Hamilton: United Nations University Institute for Water, Environment and Health, 35 pp.
- Schutte V, Selig E, Bruno J. 2010. Regional spatio-temporal trends in Caribbean coral reef benthic communities. *Marine Ecology Progress Series* 402:115–122 DOI 10.3354/meps08438.
- Selig ER, Casey KS, Bruno JF. 2012. Temperature-driven coral decline: the role of marine protected areas. *Global Change Biology* 18:1561–1570 DOI 10.1111/j.1365-2486.2012.02658.x.
- Sheppard C, Sheppard A, Mogg A, Bayley D, Dempsey AC, Roche R, Turner J, Purkis S. 2017. Coral bleaching and mortality in the Chagos Archipelago to 2017. *Atoll Research Bulletin* 613:1–25.
- Smith JE, Brainard R, Carter A, Dugas S, Edwards C, Harris J, Lewis L, Obura D, Rohwer F, Sala E, Vroom PS, Sandin S. 2016. Re-evaluating the health of coral reef communities: baselines and evidence for human impacts across the central

- Pacific. *Proceedings of the Royal Society B: Biological Sciences* **283**:Article 20151985
DOI [10.1098/rspb.2015.1985](https://doi.org/10.1098/rspb.2015.1985).
- Teh LSL, Zeller D, Cabanban A, Teh LCL, Sumaila UR. 2007.** Seasonality and historic trends in the reef fisheries of Pulau Banggi, Sabah, Malaysia. *Coral Reefs* **26**:251–263
DOI [10.1007/s00338-006-0182-x](https://doi.org/10.1007/s00338-006-0182-x).
- Thurstan RH, Buckley SM, Ortiz JC, Pandolfi JM. 2015.** Setting the record straight: assessing the reliability of retrospective accounts of change. *Conservation Letters* **9**:98–105 DOI [10.1111/conl.12184](https://doi.org/10.1111/conl.12184).
- Wapnick CM, Precht WF, Aronson RB. 2004.** Millennial-scale dynamics of staghorn coral in Discovery Bay, Jamaica. *Ecology Letters* **7**:354–361
DOI [10.1111/j.1461-0248.2004.00586.x](https://doi.org/10.1111/j.1461-0248.2004.00586.x).
- Wing S, Wing E. 2001.** Prehistoric fisheries in the Caribbean. *Coral Reefs* **20**:1–8
DOI [10.1007/s003380100142](https://doi.org/10.1007/s003380100142).
- Woodley JD. 1992.** The incidence of hurricanes on the north coast of Jamaica since 1870: are the classic reef descriptions a typical? *Hydrobiologia* **247**:133–138
DOI [10.1007/BF00008212](https://doi.org/10.1007/BF00008212).