

Spin-label scanning reveals conformational sensitivity of the bound helical interfaces of IA₃

Katie M. Dunleavy^{†1}, Eugene Milshteyn^{†1a}, Zachary Sorrentino^{†1b}, Natasha L. Pirman^{1c}, Zhanglong Liu^{1d}, Matthew B. Chandler^{1e}, Peter W. D'Amore^{1f}, and Gail E. Fanucci^{*1}

[†] Contributed equally

¹Department of Chemistry, PO BOX 117200, University of Florida, Gainesville, FL, 32611-7200 USA

Current addresses:

^a Department of Radiology and Biomedical Imaging, University of California, San Francisco, CA, USA, ^b College of Medicine, University of Florida, Gainesville, FL, USA, ^c MilliporeSigma, Burlington, MA, USA, ^d CVS Health, Woonsocket, RI, USA ^e Expertus Laboratories, Inc. Tampa, FL, USA, ^f Department of Orthopedic Surgery, LSU Health Science Center, New Orleans LA, USA.

*Contributing Author: Gail E. Fanucci

Address: University of Florida
Department of Chemistry
P.O. Box 117200
Gainesville, FL, 32611
Email: fanucci@chem.ufl.edu
Phone: (352)-392-0872

Abstract

IA₃ is an intrinsically disordered protein (IDP) that becomes helical when bound to yeast proteinase A (YPRA) or in the presence of the secondary stabilizer 2, 2, 2, trifluoroethanol (TFE). Here, site-directed spin-labeling (SDSL) continuous wave electron paramagnetic resonance (CW-EPR) spectroscopy and circular dichroism (CD) are used to characterize the TFE-induced helical conformation of IA₃ for a series of spin-labeled cysteine scanning constructs and varied amino acid substitutions. Results demonstrate that the N-terminal concave helical surface of IA₃, which is the buried interface when bound to YPRA, can be destabilized by the spin-label or bulky amino acid substitutions. In contrast, the helical tendency of IA₃ is enhanced when spin-labels are incorporated into the convex, i.e., solvent exposed, surface of IA₃. No impact of the spin-label within the C-terminal region was observed. This work further demonstrates the utility and sensitivity of SDSL CW-EPR for studies of IDPs. In general, care must be taken to ensure the

spin-label does not interfere with native helical tendencies and these studies provide us with knowledge of where to incorporate spin-labels for future SDSL investigations of IA₃.

Key Words

site-directed spin-labeling, intrinsically disordered protein, IA₃, nitroxide label, electron paramagnetic resonance, circular dichroism

Abbreviations

Intrinsically disordered protein (IDP); Intrinsically disordered region (IDR); Yeast proteinase A (YPRA); 2, 2, 2-trifluoroethanol (TFE); Circular dichroism (CD); Site-directed spin labeling (SDSL); Continuous wave electron paramagnetic resonance (CW-EPR); 3-(2-Iodoacetamido)-PROXYL (IAP); (1-oxyl-2,2,5,5-tetramethyl- Δ 3-pyrroline-3-methyl) methanethiosulfonate (MTSL); 4-maleimido-TEMPO (MSL); Dithiothreitol (DTT).

Introduction

Proteins or protein regions of 50 or more residues, which do not have stable secondary or tertiary structure under physiological conditions, are characterized as intrinsically disordered proteins (IDPs) or intrinsically disordered regions (IDRs) [1-7]. Although the structure-function paradigm suggests that 3D structure is necessary for function, the discovered importance of intrinsic disorder within proteins or protein regions of higher eukaryotic systems shows how function could arise from various unstructured states [3, 4, 8, 9]. IDPs have important roles in cellular signal transduction, translation, and transcription, influencing the study of disordered proteins by focusing on both the functions provided by IDPs, as well as the conformational changes associated when they bind to their target [4, 7, 10].

IA₃, an IDP found in *Saccharomyces cerevisiae*, is composed of 68 amino acid residues and acts as an inhibitor of yeast proteinase A (YPRA). Previous studies have shown IA₃ to adopt an α -helical conformation when bound in the active site of YPRA and when exposed to the secondary structure stabilizer 2,2,2-trifluoroethanol (TFE); with helical tendencies of IA₃ differing between the N-terminal residues (2-34) and C-terminal residues (35-68) [11-15]. X-ray crystallographic models of IA₃ bound to YPRA indicate α -helical structure in the N-terminal region (residues 2-34). In contrast, the C-terminal region (residues 35-68) has unresolved electron density, suggesting

disorder (**Figure 1**) [11]. Biophysical methods of circular dichroism (CD) spectroscopy, 2D ^1H ^{15}N nuclear magnetic resonance spectroscopy, molecular dynamics simulations, laser temperature-jump fluorescence spectroscopy, fluorescence resonance energy transfer spectroscopy, and site directed spin labeling (SDSL) provide data supporting a two-state transition for IA₃, where the C-terminus undergoes a helical transition but is less pronounced than the N-terminus [13-18].

SDSL, in combination with continuous wave electron paramagnetic resonance (CW-EPR) spectroscopy is biophysical tool for probing structure, dynamics and conformational changes in macromolecules. [19-23]. For proteins, typically a CW-EPR active chemical groups, such as a nitroxide spin label, is incorporated at desired locations. by chemical modification of a substituted cysteine, which provides a chemically reactive side group for covalent bond spin labeling. Recent progress in spin labeling methodology has also demonstrated the ability to incorporate non-natural amino acids for subsequent chemical modification, as well as novel schemes for specifically labeling tyrosine residues; a potentially very useful strategy for spin labeling IDPs [24-27].

The resultant CW-EPR spectral line shape reflects motional averaging of the nitroxide spin label [20, 21, 28]. Typically, the effects are described by three main types of motion, occurring in the 0.1-50 ns time scale [29, 30]. The corresponding correlation times are referred to as the following: τ_R , the overall tumbling of the protein, τ_I , the movement of the spin label about the bonds connecting it to the protein, and τ_B , the motion of the spin labeled protein backbone. SDSL CW-EPR has been applied to study structure-unstructured transitions in proteins as well as to IDP systems [5, 17, 22, 31, 32]. For structured proteins $> \sim 15$ kDa, line shape parameters such as the second moment and scaled mobility readily reveal conformational changes without the need to increase solution viscosity [29, 33]. Even ordered to disordered transitions in structured proteins are readily characterized by these “traditional” line shape parameter analyses [34, 35]. However, in IDPs, the degree of disorder and the dynamics of the system often times result in “isotropic-like” spectra, that fall into the fast motional averaging limit. In these cases, for relatively fast motional averaging, the ratio of the intensities of the low field ($h_{(+1)}$) and central field ($h_{(0)}$) transitions provide an alternative line shape analysis parameter, $h_{(1)}/h_{(0)}$, that is useful to describe conformational changes [5, 17, 22, 36].

Here, extending upon previous work, we performed a spin-labeled cysteine scanning profile of the N- and C-termini of IA₃, where a series of fifteen singly spin-labeled IA₃ constructs consisting of ten and five cysteine substitutions in the N- and C-terminus were generated, respectively (**Figure**

1). A comparison is provided between the termini on a site by site basis revealing variations in the degree of transition of the two termini and sensitivity of the helical content of the N-terminus to the spin-labeled cysteine substitution and select amino acid substitutions. The degree of site-specific helical propensity of the N-terminus, however, as these results show, is modulated by the introduction of a chemically modified cysteine residue. By further probing site V8 of IA₃, the specific effects on secondary structure due to site-specific amino acid substitution and spin labeling are analyzed. The size of an introduced residue or chemical modification has marked effects on the helical propensity within the N-terminal region of IA₃, whether found on the buried (concave) or solvent exposed (convex) side of the helix when bound to YPRA. By comparing SDSL and CD results, we are able to determine sites within IA₃ that can tolerate spin-label incorporation and retain WT helical tendencies; thus, we label those our “wild-type mimics”.

Materials and methods

Materials

BL21(DE3) pLysS cells were purchased from Invitrogen (Carlsbad, CA). DNA taq polymerase and DpnI were purchased from New England Biolabs (Ipswich, MA). A 5-mL chelating column was purchased from GE Healthcare (Wauwatosa, WI). A HiPrep 26/10 desalting column was purchased from Amersham (Pittsburgh, PA). 16.5% Tris-Tricene gels were purchased from Bio-Rad (Hercules, CA). 3-(2-Iodoacetamido)-PROXYL (IAP), (1-oxyl-2,2,5,5-tetramethyl-Δ³-pyrroline-3-methyl) methanethiosulfonate (MTSL), and 4-maleimido-TEMPO (MSL) spin label was purchased from Sigma Aldrich (St. Louis, MO). 0.60 i.d. x 0.84 o.d. capillary tubes were purchased from Fiber Optic Center (New Bradford, MA). Unless otherwise stated, all other reagents and products were purchased from Fisher Scientific (Pittsburgh, PA) and used as received.

Protein expression and purification of IA₃ wild type and mutants

E. coli codon-optimized DNA for the IA₃ gene from *S. cerevisiae*, and DNA primers used for site-directed mutagenesis were purchased from DNA2.0. The optimized IA₃ gene was cloned into the pET-22b⁺ vector containing a C-terminal 6xHis-tag, adding the sequence LEHHHHHH to the C-terminus as described previously [17]. Site directed mutagenesis via polymerase chain reaction was used to introduce non-native cysteines by designing mutagenic primers. The substituted sequence in each resultant plasmid was confirmed via Sanger DNA sequencing. WT and cysteine

substituted constructs were purified as described previously [17]. Following affinity chromatography, residual nickel from the column was removed by addition of EDTA to a concentration of 100 mM and cysteine reduction was ensured by addition of dithiothreitol (DTT) to a concentration of 0.1 mM. Protein purity was assessed using a 16.5% tris-tricine SDS-PAGE gel.

Spin labeling of IA₃ mutants

WT and cysteine substituted proteins were buffer exchanged into 50 mM sodium phosphate, 300 mM sodium chloride, 10% (v/v) glycerol, pH 7.4 using a desalting column. This step also removed residual nickel, EDTA, and DTT that were added as described above. After desalting, IAP, MTSL, or MSL nitroxide spin label, dissolved in ethanol, was added in excess (5-10X molar ratio) and allowed to react with cysteine substituted constructs for 10-12 h at room temperature in the dark. Spin labeling schemes are represented in **Figure 2A**. Excess spin label was removed by repeating the desalting buffer exchange process as described above.

Sample preparation for CW-EPR experiments

For each spin labeled construct, nine separate samples were prepared with increasing 5% (v/v) increments of 2, 2, 2-trifluoroethanol (TFE) that spanned 0-40%. Each individual sample was prepared at a final volume of 500 μ L: 300 μ L of 50 mM sodium phosphate, 300 mM sodium chloride, pH 7.4 buffer, (200-x) μ L of TFE, where x ranged from 0-200 μ L, and x μ L of appropriately prepared solution of 5X stock buffer diluted with water to keep all samples at equal ionic strength and pH. These larger 500 μ L sample volumes were prepared to help mitigate pipetting errors of small volumes. Samples of ~10 μ L were drawn into 0.60 i.d. x 0.84 o.d. capillary tubes, which were then flame sealed on both ends ensuring minimal to no TFE evaporation.

Sample preparation for CD experiments

For each spin labeled construct, samples were prepared with varying percentages of increasing TFE concentrations. For IAP labeled cysteine scanning and the V8C construct with varying spin label attachments, three samples were prepared with increasing 15% (v/v) increments of TFE from 0-30%. For amino acid scanning at the 8th position of the IA₃ sequence and cysteine or alanine substitution at the 11th position, six or nine samples were prepared with increasing 7% (v/v)

increments of TFE from 0-35% or increasing 5% (v/v) increments of TFE from 0-40%, respectively. Samples were prepared with 80-120 μ L of protein sample in 50 mM sodium phosphate, 300 mM sodium chloride, pH 7.4 to reach a final concentration of 10 μ M protein in 1000 μ L. The final volume was adjusted using 10 mM sodium phosphate, pH 7.0 buffer or TFE to reach appropriate % (v/v) TFE. For the V8C un-labeled construct, 0.1 mM DTT was present to prevent disulfide bonding of free cysteine residues.

CW X-band EPR data collection and analysis

A Bruker ER200 spectrometer with an ER023 M signal channel, an ER032 M field control unit, and a loop gap resonator was used to collect CW X-band EPR spectra. For all experiments the temperature was kept at a constant $27 \pm 0.2^\circ\text{C}$ using a nitrogen gas passed through a copper coil submersed into a water bath, while monitored with a temperature probe and thermometer from OMEGA Engineering Inc (Norwalk, CT). Prior to collecting spectra for each construct, the modulation amplitude and time constant were optimized for maximum signal-to-noise ratio without any line broadening. All spectra are reported as an average of four scans. Spectra were normalized using labVIEW software allowing for baseline correction and double integral area normalization. A representative nitroxide EPR spectrum with the typical three-line transition pattern is shown and labeled in **Figure 2B**. Spectral line shapes were analyzed by calculating the ratio of the intensities of the low field and center field transitions, $h_{(+1)}/h_{(0)}$ as a function of TFE percentage. Although the low field and center field transitions individually are not as sensitive to motion as the high field transition ($h_{(-1)}$), this ratio has been previously shown to change considerably as the unstructured-to-structured transition occurs and is not sensitive to errors in baseline correction or double integration [17, 36]. We have used this analysis previously to monitor the unstructured to helical transition in IA3 by TFE. Plots of $h_{(+1)}/h_{(0)}$ values versus the % TFE are sigmoidal in shape and are well fit by a two-state Boltzmann transition, given by Eq. (1).

$$y = \frac{(A_1 - A_2)}{1 + e^{(x - x_0)/dx}} + A_2 \quad (1)$$

A_1 is the initial value of the curve, A_2 is the final value of the curve and x_0 is the midpoint of the curve – which corresponds well to the midpoint of folding transitions observed by CD and NMR. Data were fit using Origin 8.5 software.

CD data collection and analysis

CD measurements were collected using an AVIV model 202 CD spectrometer set at 27°C. To collect the spectra, 400 μ L of each sample was loaded into a 1mm path length quartz cuvette, between runs, the cuvette was cleaned with nanopure H₂O and ethanol. Typically, four scans of each buffer blank and sample were collected between wavelengths of 200-250nm with 1 nm increments and a four second averaging time. Each set of CD scans for each construct was averaged, and the averaged buffer baseline was subtracted from each individual construct scan average. Units were converted to $\Delta\epsilon$ (molar circular dichroism, M⁻¹cm⁻¹) from the instruments output unit of Θ (ellipticity, mdeg) using Eq. 2.

$$\Delta\epsilon (M^{-1}cm^{-1}) = \frac{\Theta (mdeg) \times C \times L}{100 \times M} \quad (2)$$

Θ is the output ellipticity value from the CD measurement, C is the concentration of sample in g/L, L is the path length in cm, and M is the average molecular weight in g/mol.

Results and Discussion

SDSL CW-EPR spectroscopy reveals sites where spin-label incorporation alters helicity.

It is well known that the labeling of a naturally occurring or substituted cysteine may impact the native function of a protein. To understand if spin-label and cysteine substitution within IA₃ impacted folding or folding propensity, cysteine scanning was performed across the N- and C-term of IA₃. Sites K7, V8, S9, E10, I11, F12, Q13, S14, S15, and S27 were chosen within the N-terminus and span two helical turns containing both solvent exposed and buried residues in the α -helix when bound to YPRA. Sites Y57, N58, K59, L60, and K61, were selected in the C-terminus (**Figure 1**). CW X-band EPR spectra were collected for the fifteen IAP spin labeled-IA₃ variants (referred to within as P1-IA₃) as a function of TFE concentration that ranged from 0-40% (v/v) TFE. **Figure 3** shows representative CW-EPR spectra for 0%, 15%, and 30% (v/v) TFE for each P1-IA₃ variant. Spectra are plotted with normalized area to allow for easy visualization of changes in mobility. As mobility (defined as both the rate of motion and order of motion) decreases, the line shape broadens and becomes less intense. A decrease in mobility upon increasing TFE percentage is observed for nearly all sites and indicates a conformational change upon increasing TFE concentration (earlier work has ruled out the impact of increased solution viscosity) [17]. Inspection of the line shapes shows that the change in mobility differs among these 15 P1-IA₃ variants, indicating that some

spin-labeled cysteine substitutions affect the conformation of each state, altering the TFE-induced helical propensity of IA₃.

A more quantitative way to analyze the TFE-induced conformational changes for each variant is to plot $h_{(+1)}/h_{(0)}$ values as a function of TFE % (**Figure 4**). For isotropic motion, the $h_{(+1)}/h_{(0)}$ value is expected to be 1. As the mobility decreases as the protein becomes helical, lower values of $h_{(+1)}/h_{(0)}$ are expected. In the unfolded state (absence of TFE), all P1-IA₃ variants have similar CW-EPR spectra indicative of similar high nitroxide mobility (indicated by values of $h_{(+1)}/h_{(0)} \sim 0.9$). As a function of TFE concentration, the shape of each $h_{(+1)}/h_{(0)}$ plot can be well fit by Eq. 1, indicative of a 2-state change.

Several conclusions can be drawn from the data in **Figure 4**. Firstly, the sigmoidal behavior of the $h_{(+1)}/h_{(0)}$ parameter for N-terminal P1-IA₃ variants (gray shade, solid diamonds) varies both in sharpness and extent of transition, which is defined by the limiting $h_{(+1)}/h_{(0)}$ value. In contrast C-terminal P1-IA₃ variants (no shade, open circles) exhibit similar behavior, both in the sharpness and extent of transition. Secondly, there are site specific variations in the behavior of the $h_{(+1)}/h_{(0)}$ parameter in the N-terminus that follow a helical trend that maps to the surface of IA₃ in the YRPA bound site. Specifically, sites V8P1, I11P1, and S15P1 which are located on the concave buried face of the peptide when bound to YRPA, have small changes (< 0.05 over the 0-40% (v/v) TFE range) in their $h_{(+1)}/h_{(0)}$ values. In contrast, sites S9P1, Q13P1 and S27P1, which reside on the convex solvent exposed face when bound to YRPA, poses large changes in their $h_{(+1)}/h_{(0)}$ values (> 0.3 over the 0-40% TFE range). **Figure 5** shows the location of these residues in the IA₃:YRPA complex crystal structure. These results indicate a sensitivity of the N-terminal helicity to amino acid substitution.

To further explore these observations, CD spectroscopy was performed on each of the N-terminal P1-IA₃ variants with 0%, 15%, and 30% (v/v) TFE. Data are shown in **Figure 6**, which also overlays results from WT IA₃ (gray solid line) in 30% (v/v) TFE for comparison. For wild type IA₃, CD data reveal a predominantly random coil conformation for 0% (v/v) TFE, whereas with 30% (v/v) TFE the spectra reflect a mostly helical conformation with features at 208 nm and 222 nm indicative of helix formation [37, 38]. For all N-terminal P1-IA₃ variants, CD data confirm that in the absence of TFE, all constructs adopt a random coil conformation and that the addition of TFE induces a helical conformation. As was observed with the CW-EPR data, the degree of the helical transition, as reflected in the values of $\Delta\epsilon_{222\text{nm}}$ in CD spectra, varies when compared to WT

for the P1 variants. It can be seen that I11P1 and S14P1 possess CD spectra that very closely match that of WT (*i.e.*, represent SL wild-type mimics). V8P1 is found to be less helical than WT, whereas all other variants have greater helicity than WT. To directly compare the SDSL CW-EPR characterized unstructured-to-structured transition within the N-terminus to the degree of α -helicity from CD spectroscopy, we plotted the values of $\Delta h_{(+1)}/h_{(0)}$, or degree of transition ($h_{(+1)}/h_{(0)}$ (40% (v/v) TFE) - $h_{(+1)}/h_{(0)}$ (0% (v/v) TFE)) against the $\Delta\epsilon_{222\text{nm}}$ (30% (v/v) TFE) (Eq. 2) values from CD spectra for each P1-IA₃ variant (**Figure 7**). The $\Delta\epsilon_{222\text{nm}}$ value for WT IA₃ is displayed in **Figure 7** as a gray line to represent how the helical conformation of each variant compares to WT behavior. As can be seen, there is remarkable agreement in overall trend between these two parameters, confirming that the helical propensity of the N-terminal region in IA₃ is particularly sensitive to amino acid substitution/spin-labeling. It is striking that CW-EPR line shape analysis readily reflects this sensitivity to amino acid substitution/modification and in most cases the incorporation of the spin-label is found to enhance the helicity of IA₃.

Helical propensity at site V8 in IA₃ is modulated by amino acid substitution and size

Because the P1 substitution at site V8 revealed less helical character than WT, we proceeded to generate a series of substitutions at this site to probe how amino acid size and type impacted helical propensity. The following constructs were chosen: V8A and V8I (smaller and larger than V; respectively), V8P (typical helical disrupter), V8D (charged), and V8C (our cysteine mutation without spin-label). CD spectra of each construct in 30% (v/v) TFE are given in **Figure 8A**. It is found that TFE does induce a helical transition in each of these constructs. We assessed the impact of each substitution on helical content by calculating the difference in $\Delta\epsilon_{222\text{nm}}$ of the variant to WT. **Figure 8B** plots values of $\Delta(\Delta\epsilon_{222\text{nm}})$, showing that that as expected the proline substitution at site 8 is less helical than WT. Also as expected, the alanine substitution induces the greatest increase in helical content, whereas other substitutions resulted in near WT behavior.

Choice of spin-label at site V8C alters helical behavior of IA₃

The result that V8C contained WT helicity was surprising, which prompted us to examine if and how spin-label structure influenced helicity when incorporated at this site. Interestingly, we find that the P1 substitution (CYS labeled with IAP) disrupted helicity compared to WT, MSL-CYS behaved like WT and R1 (MTSL-CYS) slightly increased helical content (**Figures 8C and 8D**).

Perhaps the structure of the P1 label, which contains a pseudo-peptide bond may interfere with TFE induced folding. MSL is a bulkier label than P1, and R1 is oftentimes considered a compact label because of potential sulfur interactions with the protein backbone [39].

Nevertheless, results appear to indicate a structural bulkiness basis for disruption or enhancement of helical propensity. Indeed, the structure of IA₃ bound to YPRA shows a slight concave bend on the buried interface, which would make substitution to larger residues disrupt helical behavior and conversely then a smaller residue would enhance helicity. To test this idea, we made two additional constructs, V8P1-I11A and V8A-I11P1, which would place the smaller alanine residue next to the P1 spin-label. V8P1-I11A shows nearly WT helical behavior and although V8A-I11P1 is less helical than WT, it is more helical than V8P1 (**Figure 8E and 8F**).

Conclusions

It was determined through SDSL CW-EPR that the extent of helical order within the N-terminal region of IA₃ is easily perturbed by amino acid substitution and IAP labeling. The C-terminal region did not have similar susceptibility, where changes in labeled amino acid substitution did not reveal drastic changes in the unstructured to structured transition and as we showed previously, retained WT behavior [17]. CD results confirmed that the helical order within the N-terminus varied due to the location of the altered residue in relation to the buried, concave side, or the solvent-exposed, convex side of the protein. The concave face of IA₃, is negatively affected by cysteine mutation and labeling, whereas the convex surface increases in helical order. It is proposed that the change in helical propensity on either side of the protein are due to spatial availability when incorporating a cysteine residue and nitroxide probe. Residue V8, located on the concave side of the N-terminal region showed the greatest negative perturbation due to cysteine substitution, and spin labeling. Mutations to the V8 residue, including substitutions in residue charge and size did not drastically affect the helical propensity, but rather the addition of the IAP nitroxide probe had the greatest effect on helical formation.

These experiments demonstrate the care that must be taken when spin-labeling IDPs. Proper checks-and-balances are needed to ensure that a labeled site retains WT helical propensity (in addition to any other functional assays). Overall, the results provide us with known sites for spin-label incorporation that mimic WT helical tendencies that can be used in future studies of IA₃ conformational sampling studies.

Acknowledgements

This work was supported by NSF MCB 1329467 & MCB 1715384. The authors would like to thank Stephen J. Hagen and Robert McKenna for access to the Aviv CD spectrometer, and Christian Altenbach for Labview software to analyze the EPR data.

Conflict of interest

All authors declare no conflict of interest in this paper.

References

1. Dunker, A.K., et al., *Intrinsically disordered protein*. J Mol Graph Model, 2001. **19**(1): p. 26-59.
2. Tompa, P., *Intrinsically unstructured proteins*. Trends Biochem Sci, 2002. **27**(10): p. 527-33.
3. Wright, P.E. and H.J. Dyson, *Intrinsically unstructured proteins: re-assessing the protein structure-function paradigm*. J Mol Biol, 1999. **293**(2): p. 321-31.
4. Dyson, H.J. and P.E. Wright, *Intrinsically unstructured proteins and their functions*. Nat Rev Mol Cell Biol, 2005. **6**(3): p. 197-208.
5. Morin, B., et al., *Assessing induced folding of an intrinsically disordered protein by site-directed spin-labeling electron paramagnetic resonance spectroscopy*. J Phys Chem B, 2006. **110**(41): p. 20596-608.
6. Knott, M. and R.B. Best, *A preformed binding interface in the unbound ensemble of an intrinsically disordered protein: evidence from molecular simulations*. PLoS Comput Biol, 2012. **8**(7): p. e1002605.
7. Dunker, A.K., et al., *The unfoldomics decade: an update on intrinsically disordered proteins*. BMC Genomics, 2008. **9 Suppl 2**: p. S1.
8. Dyson, H.J. and P.E. Wright, *Coupling of folding and binding for unstructured proteins*. Curr Opin Struct Biol, 2002. **12**(1): p. 54-60.
9. Fuxreiter, M., et al., *Preformed structural elements feature in partner recognition by intrinsically unstructured proteins*. J Mol Biol, 2004. **338**(5): p. 1015-26.
10. Pancsa, R. and P. Tompa, *Structural disorder in eukaryotes*. PLoS One, 2012. **7**(4): p. e34687.
11. Li, M., et al., *The aspartic proteinase from Saccharomyces cerevisiae folds its own inhibitor into a helix*. Nat Struct Biol, 2000. **7**(2): p. 113-7.
12. Philip, L.H., et al., *The potency and specificity of the interaction between the IA3 inhibitor and its target aspartic proteinase from Saccharomyces cerevisiae*. J Biol Chem, 2001. **276**(3): p. 2023-30.
13. Green, T.B., et al., *IA3, an aspartic proteinase inhibitor from Saccharomyces cerevisiae, is intrinsically unstructured in solution*. Biochemistry, 2004. **43**(14): p. 4071-81.

14. Ganesh, O.K., et al., *Characterizing the residue level folding of the intrinsically unstructured IA3*. Biochemistry, 2006. **45**(45): p. 13585-96.
15. Narayanan, R., et al., *Kinetics of folding and binding of an intrinsically disordered protein: the inhibitor of yeast aspartic proteinase YPrA*. J Am Chem Soc, 2008. **130**(34): p. 11477-85.
16. Wang, J., et al., *Multi-scaled explorations of binding-induced folding of intrinsically disordered protein inhibitor IA3 to its target enzyme*. PLoS Comput Biol, 2011. **7**(4): p. e1001118.
17. Pirman, N.L., et al., *Characterization of the disordered-to-alpha-helical transition of IA(3) by SDSL-EPR spectroscopy*. Protein Sci, 2011. **20**(1): p. 150-9.
18. Le Breton, N., et al., *Diversification of EPR signatures in Site Directed Spin Labeling using a beta-phosphorylated nitroxide*. Phys Chem Chem Phys, 2014. **16**(9): p. 4202-9.
19. Hubbell, W.L., D.S. Cafiso, and C. Altenbach, *Identifying conformational changes with site-directed spin labeling*. Nat Struct Biol, 2000. **7**(9): p. 735-9.
20. Hubbell, W.L., et al., *Watching proteins move using site-directed spin labeling*. Structure, 1996. **4**(7): p. 779-83.
21. Fanucci, G.E. and D.S. Cafiso, *Recent advances and applications of site-directed spin labeling*. Curr Opin Struct Biol, 2006. **16**(5): p. 644-53.
22. Longhi, S., et al., *Probing structural transitions in both structured and disordered proteins using site-directed spin-labeling EPR spectroscopy*. J Pept Sci, 2011. **17**(5): p. 315-28.
23. Cafiso, D.S., *Identifying and quantitating conformational exchange in membrane proteins using site-directed spin labeling*. Acc Chem Res, 2014. **47**(10): p. 3102-9.
24. Fleissner, M.R., et al., *Site-directed spin labeling of a genetically encoded unnatural amino acid*. Proc Natl Acad Sci U S A, 2009. **106**(51): p. 21637-42.
25. Lorenzi, M., et al., *Tyrosine-targeted spin labeling and EPR spectroscopy: an alternative strategy for studying structural transitions in proteins*. Angew Chem Int Ed Engl, 2011. **50**(39): p. 9108-11.
26. Mileo, E., et al., *Enlarging the panoply of site-directed spin labeling electron paramagnetic resonance (SDSL-EPR): sensitive and selective spin-labeling of tyrosine using an isoindoline-based nitroxide*. Bioconjug Chem, 2013. **24**(6): p. 1110-7.
27. Mileo, E., et al., *Dynamics of the intrinsically disordered protein CP12 in its association with GAPDH in the green alga Chlamydomonas reinhardtii: a fuzzy complex*. Mol Biosyst, 2013. **9**(11): p. 2869-76.
28. Columbus, L. and W.L. Hubbell, *A new spin on protein dynamics*. Trends Biochem Sci, 2002. **27**(6): p. 288-95.
29. McHaourab, H.S., et al., *Motion of spin-labeled side chains in T4 lysozyme. Correlation with protein structure and dynamics*. Biochemistry, 1996. **35**(24): p. 7692-704.
30. Qin, P.Z., J. Iseri, and A. Oki, *A model system for investigating lineshape/structure correlations in RNA site-directed spin labeling*. Biochem Biophys Res Commun, 2006. **343**(1): p. 117-24.
31. Bartelli, N.L. and G.L. Hazelbauer, *Direct evidence that the carboxyl-terminal sequence of a bacterial chemoreceptor is an unstructured linker and enzyme tether*. Protein Sci, 2011. **20**(11): p. 1856-66.

32. Belle, V., et al., *Mapping alpha-helical induced folding within the intrinsically disordered C-terminal domain of the measles virus nucleoprotein by site-directed spin-labeling EPR spectroscopy*. Proteins, 2008. **73**(4): p. 973-88.
33. Frazier, A.A., et al., *Membrane orientation and position of the C2 domain from cPLA2 by site-directed spin labeling*. Biochemistry, 2002. **41**(20): p. 6282-92.
34. Fanucci, G.E., et al., *Substrate-induced conformational changes of the periplasmic N-terminus of an outer-membrane transporter by site-directed spin labeling*. Biochemistry, 2003. **42**(6): p. 1391-400.
35. Kim, M., G.E. Fanucci, and D.S. Cafiso, *Substrate-dependent transmembrane signaling in TonB-dependent transporters is not conserved*. Proc Natl Acad Sci U S A, 2007. **104**(29): p. 11975-80.
36. Esquiaqui, J.M., et al., *Conformational Flexibility and Dynamics of the Internal Loop and Helical Regions of the Kink-Turn Motif in the Glycine Riboswitch by Site-Directed Spin-Labeling*. Biochemistry, 2016. **55**(31): p. 4295-305.
37. Hirst, J.D. and C.L. Brooks, 3rd, *Helicity, circular dichroism and molecular dynamics of proteins*. J Mol Biol, 1994. **243**(2): p. 173-8.
38. Greenfield, N.J., *Using circular dichroism spectra to estimate protein secondary structure*. Nat Protoc, 2006. **1**(6): p. 2876-90.
39. Columbus, L. and W.L. Hubbell, *Mapping backbone dynamics in solution with site-directed spin labeling: GCN4-58 bZip free and bound to DNA*. Biochemistry, 2004. **43**(23): p. 7273-87.

Figures




Figure 1: Ribbon diagram of IA₃ bound to yeast proteinase A (YPRA). Structurally resolved residues (3-31) are rendered as a light orange α -helix, with the unresolved C-terminal residues represented as a sequence chain of the single letter amino acids. Sites chosen for spin label scanning within the resolved α -helix are represented as red spheres located at the alpha carbon positions. Sites that were spin-labeled within the C-terminal region are annotated by blue letters. (PDB: 1DPJ)




Figure 2: (A) Reaction schemes for chemical modification of a cysteine residue with various nitroxide probes. From top to bottom: IAP (3-(2-Iodoacetamido)-PROXYL) represented as P1, MTSL (2,2,5,5-tetramethyl-1-oxyl-3-methyl methanethiosulfonate) represented as R1, and MSL (3-maleimido-PROXYL). (B) A representative 100 G X-band CW-EPR spectrum with the three transition line pattern individually labeled: low field, $h_{(+1)}$, center field, $h_{(0)}$, and high field, $h_{(-1)}$. These three lines arise from the hyperfine interaction between the unpaired electron and the nitrogen nucleus of the spin label.




Figure 3: Stack plots of absorption area normalized 100 G X-band CW-EPR spectra of P1-IA₃ variants in the presence of 0%, 15%, and 30% (v/v) TFE, at $27 \pm 0.2^\circ\text{C}$. Spectra are vertically offset for visual clarity.




Figure 4: Plots of $h_{(+1)}/h_{(0)}$ values as a function of % (v/v) TFE for N-terminus (gray shade, solid diamonds) and C-terminus (no shade, open circles) P1-IA₃ variants. Solid lines through points are best fits of a two-state Boltzmann function (Eq. 1), consistent with a two-state transition of IA₃ from unstructured to α -helical; as described previously [17]. The size of each data point is larger than the standard deviation of triplicate measurements of the same sample.




Figure 5: (top) Ribbon diagram of IA₃ (dark gray) bound to YPRA (light gray), with chosen P1-labeled variants of solvent exposed and YPRA facing, buried residues highlighted. Solvent exposed highlighted residues are shown in orange, and buried residues are rendered in green. **(bottom)** Bound IA₃ structure, with YPRA removed. Buried residues located on concave side of structure shown in green, and solvent exposed residues located on convex side of structure shown in orange. (PDB: 1DPJ)




Figure 6: Circular dichroism spectra for N-terminal P1-IA₃ variants with WT spectra overlay at 30% (v/v) TFE concentration (solid gray line). P1-IA₃ variant spectra were collected with increasing % (v/v) TFE concentrations at 27°C as an average of four scans; 0% (v/v) TFE (dotted black line), 15% (v/v) TFE (dashed black line), and 30% (v/v) TFE (solid black line).




Figure 7: Comparison of helical propensity $\Delta(h_{(+1)}/h_{(0)})$ (black squares) to the $\Delta\epsilon \text{ (M}^{-1}\text{cm}^{-1}\text{)}$ at 222 nm (black circles) at 30% (v/v) TFE. Where error bars are not visible, the size of data point is larger than the standard deviation of triplicate measurement on the same sample.




Figure 8: CD data and analysis of constructs compared to WT. **(A)** CD spectra of V8 variants (V8P, V8D, V8I, V8C, and V8A) **(B)** $\Delta\Delta\epsilon$ of V8 variants compared to WT, **(C)** CD spectra of V8C-spin label variants (V8P1, V8MSL, and V8MR1), **(D)** $\Delta\Delta\epsilon$ for V8C-spin labeled variants, **(E)** CD spectra for V8-I11 variants (V8P1, V8A-I11P1, and V8P1-I11A), and **(F)** $\Delta\Delta\epsilon$ for V8-I11 variants. Red CD spectra of WT are added in each plot and where $\Delta\Delta\epsilon$ ($M^{-1}cm^{-1}$) is calculated as $\Delta\epsilon_{222nm}(WT) - \Delta\epsilon_{222nm}(variant)$ so that positive values represent more helical character and negative values indicate a decreased helicity.