



# Decavanadate Inhibits Mycobacterial Growth More Potently Than Other Oxovanadates

Nuttaporn Samart<sup>1,2</sup>, Zeyad Arhouma<sup>1,3</sup>, Santosh Kumar<sup>4</sup>, Heide A. Murakami<sup>1</sup>,  
Dean C. Crick<sup>3,4</sup> and Debbie C. Crans<sup>1,3\*</sup>

<sup>1</sup> Department of Chemistry, Colorado State University, Fort Collins, CO, United States, <sup>2</sup> Department of Chemistry, Rajabhat Rajanagarindra University, Chachoengsao, Thailand, <sup>3</sup> Cell and Molecular Biology Program, Colorado State University, Fort Collins, CO, United States, <sup>4</sup> Department of Microbiology, Immunology and Pathology, Colorado State University, Fort Collins, CO, United States

## OPEN ACCESS

### Edited by:

Christian Hartinger,  
University of Auckland, New Zealand

### Reviewed by:

Eugenio Garribba,  
University of Sassari, Italy  
Sanja Grgric-Sipka,  
University of Belgrade, Serbia

### \*Correspondence:

Debbie C. Crans  
debbie.crans@colostate.edu

### Specialty section:

This article was submitted to  
Inorganic Chemistry,  
a section of the journal  
Frontiers in Chemistry

Received: 24 June 2018

Accepted: 09 October 2018

Published: 20 November 2018

### Citation:

Samart N, Arhouma Z, Kumar S, Murakami HA, Crick DC and Crans DC (2018) Decavanadate Inhibits Mycobacterial Growth More Potently Than Other Oxovanadates. *Front. Chem.* 6:519.

doi: 10.3389/fchem.2018.00519

<sup>51</sup>V NMR spectroscopy is used to document, using speciation analysis, that one oxometalate is a more potent growth inhibitor of two Mycobacterial strains than other oxovanadates, thus demonstrating selectivity in its interaction with cells. Historically, oxometalates have had many applications in biological and medical studies, including study of the phase-problem in X-ray crystallography of the ribosome. The effect of different vanadate salts on the growth of *Mycobacterium smegmatis* (*M. smeg*) and *Mycobacterium tuberculosis* (*M. tb*) was investigated, and speciation was found to be critical for the observed growth inhibition. Specifically, the large orange-colored sodium decavanadate (V<sub>10</sub>O<sub>28</sub><sup>6-</sup>) anion was found to be a stronger inhibitor of growth of two mycobacterial species than the colorless oxovanadate prepared from sodium metavanadate. The vanadium(V) speciation in the growth media and conversion among species under growth conditions was monitored using <sup>51</sup>V NMR spectroscopy and speciation calculations. The findings presented in this work is particularly important in considering the many applications of polyoxometalates in biological and medical studies, such as the investigation of the phase-problem in X-ray crystallography for the ribosome. The findings presented in this work investigate the interactions of oxometalates with other biological systems.

**Keywords:** vanadate, speciation, sodium metavanadate, decavanadate, growth inhibition, tuberculosis

## INTRODUCTION


Polyoxometalates (POMs) are a class of compounds that have been investigated in a range of biological and biomedical systems, as their effects in cell culture studies and *in vivo* suggest that these compounds have potential for use as therapeutic agents (Moskovitz and Group, 1988; Hill et al., 1990; Rhule et al., 1998; Aureliano and Crans, 2009; Fraqueza et al., 2012; Aureliano et al., 2013, 2016; Wang et al., 2013; Aureliano and Ohlin, 2014; Leon et al., 2014; Kioseoglou et al., 2015; Shah et al., 2015; Sun et al., 2016; Fu et al., 2018; Gumerova et al., 2018). Decavanadate is a homopolyoxometalate anion, and one of the POMs that has been reported to have known biological effects, as documented by studies reported with cells as well as with

isolated enzyme systems. (Pluskey et al., 1996; Aureliano and Crans, 2009; Zhai et al., 2009; Fraqueza et al., 2012; Turner et al., 2012; Aureliano et al., 2013; Kioseoglou et al., 2013; Aureliano, 2014, 2016; Aureliano and Ohlin, 2014). Protein crystal structures have been reported for some protein-POM complexes such as those reported between the ribosome and a Dawson oxometallate (Weinstein et al., 1999; Auerbach-Nevo et al., 2005; Bashan and Yonath, 2008; Noeske et al., 2015). Other protein-POM complexes include protein complexes with smaller oxometalates such as decavanadate (Winkler et al., 2017). The limited stability of decavanadate at neutral pH would suggest that hydrolysis intermediates may form and generate stable complexes with proteins or cellular components. Speciation studies are important in this regard, and different species and possibilities must be considered when investigating the mode of action of systems that are not thermodynamically stable (Aureliano and Crans, 2009; Levina et al., 2017a). Even if the speciation is characterized, the active species and mode of action of these complex systems can be non-trivial to interpret (Willsky et al., 1984b, 1985, 2011; Delgado et al., 2005; Crans et al., 2011; Postal et al., 2016; Jakusch and Kiss, 2017). However, a wide range of activities have been reported depending on the protein, biological system or specific vanadium species (Crans, 2000; Crans et al., 2013; Correia et al., 2015; Postal et al., 2016). Recently, it has become clear that compound uptake is critical to the mode of action because many vanadium compounds are modified during the uptake process (Pessoa and Tomaz, 2010; Crans et al., 2011; Le et al., 2017; Levina et al., 2017a). In the case of a large anion such as decavanadate, the question is simply whether the species is too large to enter through protein channels and thus must be taken up through endocytosis or passive transport mechanisms. The alternative possibility is that the uptake is of the smaller vanadium oxovanadates, such as monomeric vanadate, which then oligomerizes to form decavanadate inside the cell. The formation of decavanadate has been demonstrated in yeast (*S. cerevisiae*) and thus makes this anion a desirable system to understand in greater detail (Willsky et al., 1984b, 1985).

Vanadium is a first-row transition metal ion and is in the group of transition metals that can form POMs (Baes, 1976; Chasteen, 1983; Vilas Boas and Costa Pessoa, 1987; Pope and Müller, 1991; Rehder, 1991; Crans et al., 2004, 2017). Vanadium is particularly prone to forming homopolyoxometalate ions as well (Baes, 1976; Aureliano and Crans, 2009). Indeed, pure crystalline metavanadate and orthovanadate upon dissolution will form several oxovanadate species containing vanadate monomer,  $V_1$ , vanadate dimer,  $V_2$ , vanadate tetramer,  $V_4$ , vanadate pentamer,  $V_5$  and decavanadate,  $V_{10}$  (Pettersson et al., 1983, 1985; Crans et al., 1990), **Figure 1**. Some of these species have been characterized using X-ray crystallography and have been found to interconvert in aqueous solution (Evans, 1966; Crans et al., 1990). However, the specifics of the reactions and their conditions vary. For example, vanadate and oligomeric species containing 2, 4, and 5 vanadium atoms are colorless and rapidly convert at neutral pH, **Figure 1A** (Crans et al., 1990). In contrast, decavanadate will form rapidly at acidic pH, but is only kinetically stable at neutral pH, **Figure 1B** (Baes, 1976; Pope and Müller, 1991; Aureliano and

Crans, 2009; Crans et al., 2017). Indeed, the kinetic studies have shown that decamer formation is a rapid process and much faster than the  $V_{10}$  decomposition in both neutral and basic solution (Clare et al., 1973a,b; Druskovich and Kepert, 1975; Comba and Helm, 1988; Kustin, 2015; Crans et al., 2017). The decomposition pathways investigated follow several different mechanisms and are dependent on the concentrations of  $H^+$  and  $OH^-$  and the other counter ions present in solution (Clare et al., 1973a,b; Druskovich and Kepert, 1975; Comba and Helm, 1988; Kustin, 2015). Information is needed describing how decavanadate interacts with membrane interfaces and cellular systems, including how decavanadates biological activities compare to monomeric vanadate. Specifically, an attractive alternative mode of action to simple decavanadate binding would be the direct delivery of a vanadium atom from decavanadate to a biomolecule resulting in the dissolution of the decavanadate cluster. The X-ray characterization of decavanadate shows that its dimensions are  $5.4 \text{ \AA} \times 7.7 \text{ \AA} \times 8.3 \text{ \AA}$ , **Figure 1** (Evans, 1966), a large size that cannot be accommodated by many biological transport channels. Therefore, any uptake of decavanadate is likely to be through endocytosis or a passive mechanism. An attractive alternative mode of uptake would involve dissolution of the cluster by direct delivery of vanadium atoms, for example, into a system such as a protein. However, such a mechanism is more difficult to investigate and will require more information regarding the potential interactions of the anion with ligands and interfaces. We have been addressing related questions for some time (Crans et al., 2017) and exploring the interactions of  $V_{10}$  with interfaces (Baruah et al., 2006; Crans et al., 2006, 2017; Samart et al., 2014; Sanchez-Lombardo et al., 2016). Indeed, more information is needed to be able to characterize the process of how the decavanadate converts to the vanadate monomer and smaller clusters, but such processes undoubtedly involve the molecular association of structures that form in solution. The work presented in this manuscript compares the effect of decavanadate and oxovanadates on two mycobacterial species. The results of this comparison are related to the questions of uptake of vanadate and will form the background information on which it will be possible to begin to address issues regarding potential delivery of V-atoms as well as uptake of decavanadate compared to monomeric and oligomeric vanadate.

Mycobacteria are particularly resistant to hydrophilic drug penetration, and thus studies with these organisms are of general interest. There have been several studies documenting the possibility that vanadium may affect growth of the bacteria, whether in a stimulatory or inhibitory manner; however, these studies did not consider the speciation of the studied compounds in growth media and in the biological context of the cells. For example, previous studies have been carried out investigating the effects of the simple salt vanadate on the cell growth of mycobacteria in the presence and absence of Tween-80 (Costello and Hedgecock, 1959). These studies investigated the possibility that vanadium may be stimulatory to growth because it would be able to replace  $Fe^{2+}$ ,  $Fe^{3+}$ , or biologically active Mo-atoms (Turian, 1951). Dose response curves reported in cultures in the presence of added metavanadate gave complete growth inhibition



**FIGURE 1 |** The schematic structures of (A) monomeric vanadate, monomeric and polymeric vanadate series (B) X-ray structure of  $\text{V}_{10}$  and the three different types of vanadium atoms in this complex ion with indicated dimensions (Evans, 1966). This figure was adapted with permission from Aureliano and Ohlin (2014), Crans et al. (2017).

observed at 5.0  $\mu\text{g}/\text{mL}$  V-atom for *M. tb* (H37Rv), at 10  $\mu\text{g}/\text{mL}$  V-atom for *Mycobacterium butyricum*, at 50  $\mu\text{g}/\text{mL}$  V-atom for *Mycobacterium phlei*, but even at 200  $\mu\text{g}/\text{mL}$  V-atom *M. smeg* was not completely inhibited (Costello and Hedgecock, 1959).

Additionally, studies have been carried out on vanadium coordination complexes and their effect on the growth of mycobacteria (Maiti and Ghosh, 1989; David et al., 2005; Correia et al., 2014; Gajera et al., 2015). The studies explore the possibility that a known drug, or drug derivative complexes binding to vanadium, may exert a greater synergistic activity (Correia et al., 2014; Gajera et al., 2015). For example, in a series of hydroxyquinoline (Correia et al., 2014), fluoroquinoline (Gajera et al., 2015) and acid hydrazides (Martins et al., 2015) vanadium(V) complex studies, the complexes were found with micromolar inhibitory potency against *M. tb*. In other systems the vanadium complex was found to have little if any effect on the growth of the bacterium (Maiti and Ghosh, 1989; Gajera et al., 2015). Additionally, a series of vanadium

thiosemicarbazone and hydroquinoline complexes tested against *M. tb* were found to have micromolar inhibition constant against growth, but the ligand was more potent than the complex (Correia et al., 2014; Gajera et al., 2015). In these cases, complexation to the vanadium protected the bacterium. Combined these studies demonstrate that vanadium compounds may be inhibitory or protective, depending on the environment and ligands that may be complexed to it, and thus it is of interest to investigate the effects of simple vanadium complexes prior to formation of a coordination complex.

In this work, we determine the effects of decavanadate and the rapidly exchanging oxovanadates on the growth of two mycobacterial species. These studies directly compare the effects of the two vanadium salt species and in doing so, will allow researchers to address the issue of whether such oxometalate species exert different effects, possibly due to uptake or conversion of the salt under physiological conditions. In our studies with *M. smeg* and *M. tb*, it was discovered

that decavanadate undergoes hydrolysis forming monomeric vanadate in the presence of the cells, suggesting some type of interaction with the bacterial cells or with the material excreted by the bacterial cells. Comparison of metavanadate and decavanadate demonstrates that these two oxovanadate species have very different effects on the growth of these mycobacterial species and depending on species, the potency varies.

## MATERIALS AND METHODS

### Materials

Sodium metavanadate (99.9% NaVO<sub>3</sub>) was purchased from Sigma-Aldrich, hydrochloric acid (36.5–38%, HCl) and citric acid (99% C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>) were purchased from Fischer. Deuterium oxide (D<sub>2</sub>O, 99.9%) was purchased from Cambridge Isotope Laboratories, Inc. All reagents and other chemicals were purchased from Sigma-Aldrich. Vanadium complexes were prepared according to literature (Samart et al., 2014; Sanchez-Lombardo et al., 2016).

## Preparation of Stock Solutions for NMR Spectroscopy Evaluation

### Vanadate

Colorless stock solutions of sodium metavanadate (NaVO<sub>3</sub>, 100 mM) were prepared in deuterium oxide (D<sub>2</sub>O). The suspension was heated to dissolve the solid, the solution was cooled to room temperature and then the pH was measured (~ pH 7–8) (Crans et al., 1990).

### Decavanadate

An orange-red solution of sodium decavanadate (Na<sub>6</sub>V<sub>10</sub>O<sub>28</sub>, 100 mM) was prepared from NaVO<sub>3</sub> (1.00 M, 1.22 g). Sodium metavanadate solutions were prepared directly by addition of solid NaVO<sub>3</sub> to D<sub>2</sub>O and heating the solution to dissolve the solid. Once the solution was clear, the pH was adjusted to 4.5–5.8 using 6 M DCl (DCl was prepared from 12 M HCl by the addition of D<sub>2</sub>O) resulting in an orange colored solution. The V<sub>10</sub> content in this solution was verified using <sup>51</sup>V NMR spectroscopy and assuming the vanadium was all present as V(V) the concentration of each oxovanadate could be calculated from the integration of the spectra (Crans et al., 1990).

### Citric Acid and V-Citrate Complex Solutions

Stock solutions of citric acid (100 mM, pH 2.33) were prepared in deuterium oxide (D<sub>2</sub>O). The vanadium-citrate complexes were prepared by adding equimolar amounts of citric acid and vanadium. The solution was diluted to form solutions with 75, 50, 10, and 5.0 mM vanadium(V).

### Preparation of Stock Solution to Cell Culture Studies

Stock solutions of vanadate and vanadium complexes were prepared in double distilled water (DI H<sub>2</sub>O), 7H9 media or in 100% DMSO depending on the solubility of these complexes. Specifically, metavanadate, orthovanadate or decavanadate were prepared in double distilled water. Isoniazid (INH) was dissolved in sterile distilled water and used as a positive control.




### Nuclear Magnetic Resonance (NMR) Measurements

All <sup>51</sup>V NMR spectroscopy measurements were taken on a Bruker spectrometer at 78.9 MHz for <sup>51</sup>V (400 MHz for <sup>1</sup>H). The chemical shifts were obtained using external reference for <sup>51</sup>V NMR of Na<sub>3</sub>VO<sub>4</sub> (100 mM, pH 12.9, signals at –535 (V<sub>1</sub>) and –559 ppm (V<sub>2</sub>)). The samples were prepared fresh to form vanadium-citrate complexes in media and the composition was investigated as a function of time with experiments carried out at time points: 0, 1, 5, and 24 h.

Speciation of vanadium was calculated using the integration of the vanadium peak(s) within the <sup>51</sup>V NMR spectra. The concentration of each species was determined using the known added concentration [assuming all vanadium is in the form of V(V)] (Crans et al., 1990), the integration of the vanadium peak(s) in the spectrum and by using the mole fractions for each signal, the concentration of each species could be calculated as shown in Table 1.

### Speciation Analysis

The interpretation of the <sup>51</sup>V NMR spectral data was supplemented by speciation calculations based on constants measured previously (Pettersson et al., 1983; Ehde et al., 1989; Selling et al., 1994). The species distribution diagrams were calculated by using HySS 2009 software (Alderighi et al., 1999; Carsella et al., 2017) and known speciation constants of the system at hand (Pettersson et al., 1983; Ehde et al., 1989; Selling et al., 1994). The citrate and the phosphate concentrations found in the Middlebrook 7H9 broth medium supplemented with 5% BSA, 2% dextrose, 5% catalase (ADC) enrichment, glycerol (0.2%, v/v), and Tween 80 (0.05%, v/v) were 0.48 and 24 mM, respectively (Bbl<sup>tm</sup>, 2018). The vanadium concentrations that were investigated were concentrations of 5, 3.3, and 10 mM. The speciation diagrams were constructed using the following equilibrium reactions for the binary H<sup>+</sup>-H<sub>2</sub>VO<sub>4</sub><sup>2-</sup> system [see equation (1)] (Pettersson et al., 1983), and two ternary system, H<sup>+</sup>-H<sub>2</sub>VO<sub>4</sub><sup>2-</sup>-C<sub>6</sub>H<sub>5</sub>O<sub>7</sub><sup>3-</sup> and H<sup>+</sup>-H<sub>2</sub>VO<sub>4</sub><sup>2-</sup>-H<sub>2</sub>PO<sub>4</sub><sup>-</sup> [see equation (2) (Ehde et al., 1989) and (3) (Selling et al., 1994)]. The alternate formulas were provided in the format of the equations provided for comparison.



As described previously, the equations above describe the nature of the complexes that form. For example, HVO<sub>4</sub><sup>2-</sup> is described as H<sub>2</sub>VO<sub>4</sub><sup>-</sup> minus H<sup>+</sup> and thus the species is described as (–1,1) where the p being –1 (Pettersson et al., 1983; Ehde et al., 1989).

**TABLE 1** | Formation constants of Vanadate species in a 0.6M NaCl system.

(p, q)	log $\beta$	Formula	Extended formula
<b>BINARY SYSTEM (H<sup>+</sup>-H<sub>2</sub>VO<sub>4</sub><sup>2-</sup>) REF (2)</b>			
-1, 1	-7.92	HVO <sub>4</sub> <sup>2-</sup>	(H <sup>+</sup> ) <sub>-1</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> )
-2, 2	-15.17	V <sub>2</sub> O <sub>7</sub> <sup>4-</sup>	(H <sup>+</sup> ) <sub>-2</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub>
-1, 2	-5.25	HV <sub>2</sub> O <sub>7</sub> <sup>3-</sup>	(H <sup>+</sup> ) <sub>-1</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub>
0, 2	2.77	H <sub>2</sub> V <sub>2</sub> O <sub>7</sub> <sup>2-</sup>	(H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub>
-2, 4	-8.88	V <sub>4</sub> O <sub>13</sub> <sup>6-</sup>	(H <sup>+</sup> ) <sub>-2</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>4</sub>
-1, 4	0.22	HV <sub>4</sub> O <sub>13</sub> <sup>5-</sup>	(H <sup>+</sup> ) <sub>-1</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>4</sub>
0, 4	10.0	V <sub>4</sub> O <sub>12</sub> <sup>4-</sup>	(H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>4</sub>
0, 5	12.4	V <sub>5</sub> O <sub>15</sub> <sup>5-</sup>	(H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>5</sub>
4, 10	52.1	V <sub>10</sub> O <sub>28</sub> <sup>6-</sup>	(H <sup>+</sup> ) <sub>4</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>10</sub>
5, 10	58.1	HV <sub>10</sub> O <sub>28</sub> <sup>5-</sup>	(H <sup>+</sup> ) <sub>5</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>10</sub>
6, 10	61.9	H <sub>2</sub> V <sub>10</sub> O <sub>28</sub> <sup>4-</sup>	(H <sup>+</sup> ) <sub>6</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>10</sub>
7, 10	63.5	H <sub>3</sub> V <sub>10</sub> O <sub>28</sub> <sup>3-</sup>	(H <sup>+</sup> ) <sub>7</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>10</sub>
2, 1	6.96	VO <sub>2</sub> <sup>+</sup>	(H <sup>+</sup> ) <sub>2</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>10</sub>
<b>TERNARY SYSTEM (H<sup>+</sup>-H<sub>2</sub>VO<sub>4</sub><sup>2-</sup>-C<sub>6</sub>H<sub>5</sub>O<sub>7</sub><sup>3-</sup>) REF (3)</b>			
1, 0, 1	5.217	Cit <sup>2-</sup>	(H <sup>+</sup> )(Cit <sup>3-</sup> )
2, 0, 1	9.298	Cit <sup>-</sup>	(H <sup>+</sup> ) <sub>2</sub> (Cit <sup>3-</sup> )
3, 0, 1	12.067	Cit	(H <sup>+</sup> ) <sub>3</sub> (Cit <sup>3-</sup> )
1, 2, 1	12.84	(H <sup>+</sup> )(H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub> (Cit <sup>3-</sup> )	
2, 2, 1	19.68	(H <sup>+</sup> ) <sub>2</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub> (Cit <sup>3-</sup> )	
3, 2, 1	24.12	(H <sup>+</sup> ) <sub>3</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub> (Cit <sup>3-</sup> )	
3, 1, 1	18.35	(H <sup>+</sup> ) <sub>3</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub> (Cit <sup>3-</sup> )	
2, 1, 1	14.1	(H <sup>+</sup> ) <sub>2</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub> (Cit <sup>3-</sup> )	
4, 2, 2	31.3	(H <sup>+</sup> ) <sub>4</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub> (Cit <sup>3-</sup> ) <sub>2</sub>	
5, 2, 2	35.3	(H <sup>+</sup> ) <sub>5</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub> (Cit <sup>3-</sup> ) <sub>2</sub>	
6, 2, 2	39.2	(H <sup>+</sup> ) <sub>6</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>2</sub> (Cit <sup>3-</sup> ) <sub>2</sub>	
<b>TERNARY SYSTEM (H<sup>+</sup>-H<sub>2</sub>VO<sub>4</sub><sup>2-</sup>-H<sub>2</sub>PO<sub>4</sub><sup>-</sup>) REF (4)</b>			
-2, 0, 1	-17.650	PO <sub>4</sub> <sup>3-</sup>	[(H <sup>+</sup> ) <sub>-2</sub> (H <sub>2</sub> PO <sub>4</sub> <sup>-</sup> )] <sup>3-</sup>
-1, 0, 1	-6.418	HPO <sub>4</sub> <sup>2-</sup>	[(H <sup>+</sup> ) <sub>-1</sub> (H <sub>2</sub> PO <sub>4</sub> <sup>-</sup> )] <sup>2-</sup>
1, 0, 1	1.772	H <sub>3</sub> PO <sub>4</sub>	[(H <sup>+</sup> )(H <sub>2</sub> PO <sub>4</sub> <sup>-</sup> )]
9, 14, 1	90.7	H <sub>3</sub> PV <sub>14</sub> O <sub>42</sub> <sup>6-</sup>	[(H <sup>+</sup> ) <sub>9</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>14</sub> (H <sub>2</sub> PO <sub>4</sub> <sup>-</sup> )] <sup>6-</sup>
10, 14, 1	94.84	H <sub>4</sub> PV <sub>14</sub> O <sub>42</sub> <sup>5-</sup>	[(H <sup>+</sup> ) <sub>10</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>14</sub> (H <sub>2</sub> PO <sub>4</sub> <sup>-</sup> )] <sup>5-</sup>
11, 14, 1	96.41	H <sub>5</sub> PV <sub>14</sub> O <sub>42</sub> <sup>4-</sup>	[(H <sup>+</sup> ) <sub>11</sub> (H <sub>2</sub> VO <sub>4</sub> <sup>2-</sup> ) <sub>14</sub> (H <sub>2</sub> PO <sub>4</sub> <sup>-</sup> )] <sup>4-</sup>

### Bacterial Strains and Culture Conditions

*Mycobacterium smegmatis* mc<sup>2</sup>155 was grown in 7H9 Middlebrook Medium supplemented with 0.2% (v/v) glycerol, 10% ADC, 0.05% Tween-80, and incubated at 37°C with shaking for 24 h. The growth of bacteria was monitored to mid-logarithmic growth phase using a spectrophotometer at 600 nm to an optical density of 0.6 (OD<sub>600nm</sub>) (Bbl<sup>tm</sup>, 2018). Metavanadate (40 mM, pH 8.6) and decavanadate [100 mM V<sub>10</sub> (1.0 M V-atoms), pH 3.8] stock solutions and the media with bacteria were added to a 96 well plate in a 3-fold dilution experiment monitored at times 0, 1, 5, and 24 h. The pH of these samples ranged from 5.5 to 6.3 before and after treatment with

V-compounds and growth; see spectra for in supplemental for details (Upadhyay et al., 2015).

Culture supernatant after cell growth generated for <sup>51</sup>V NMR analysis was prepared by running separate growth experiments for the *M. smeg* mc<sup>2</sup> 155 experiments with a total volume of 5 mL. At each time point, a 1 mL aliquot was removed for NMR analysis. <sup>51</sup>V NMR spectra were run at the time points 0, 1, 5, and 24 h without lock.

*M. tb* mc<sup>2</sup> 6230 is a nonpathogenic deletion mutant (ΔRD1 ΔpanCD) strain H37Rv that can safely be used in a Biosafety Level 2 conditions (Sambandamurthy et al., 2006). The bacteria were grown in 7H9 Middlebrook medium with the addition of D-pantothenate (24 mg/L) at 37°C to an optical density at 600 nm (OD<sub>600nm</sub>) of 0.6–0.8. The cultures were diluted and treated with the appropriate metavanadate and decavanadate stock solutions in a 96 well plate in a 3-fold dilution experiment, and incubated at 37°C for the duration of the experiment (5–7 days). The pH values of these samples ranged from 5.6 to 6.8 after growth for 5 days; see spectra for details.


<sup>51</sup>V NMR spectra were acquired on culture supernatant without lock (no D<sub>2</sub>O in sample) at the beginning and end of the bacterial growth experiment (0 and 5 days) for *M. tb* mc<sup>2</sup> 6230 growth samples.

## RESULTS


Investigating the effects of both vanadate and decavanadate on mycobacterial growth requires information on the speciation of vanadium under the conditions of the growth assay studies. *M. tb* and *M. smeg* were grown in supplemented Middlebrook 7H9 medium. Some of the components of the medium may form vanadium complexes. The major candidates for complex formation based on literature formation constants are phosphate (Gresser et al., 1986), citrate (Pettersson et al., 1985; Levina et al., 2017a), and amino acids (Crans, 2000; Rehder et al., 2002; Esbak et al., 2009). In the following, we investigated whether these complexes formed in the media, **Figure 1**.

### Reaction of Vanadate and Decavanadate With Components in Growth Media

<sup>51</sup>V NMR studies were carried out with vanadate and decavanadate solutions and Middlebrook 7H9 broth medium supplemented with 10% ADC enrichment (5% BSA, 2% dextrose, 5% catalase), glycerol (0.2%, v/v) and Tween 80 (0.05%, v/v). The mycobacteria grow well at pH 6.8, and decavanadate is known to form between pH 3 to 6.5 (Pope and Müller, 1991; Crans et al., 2004; Baruah et al., 2006; Aureliano and Crans, 2009). V<sub>10</sub> is formed at low pH values and the V<sub>10</sub> hydrolyzes at neutral pH. When V<sub>10</sub> is added to the growth media the pH was increased from pH 7.6 to 6.5/7 by the buffering capacity of the growth media and there is a potential for hydrolysis of V<sub>10</sub>. Importantly, little hydrolysis is observed in growth media in the absence of mycobacteria. NMR spectra were recorded at a range of pH values because the reactions of vanadate are very sensitive to pH (see **Supplemental Material**); however, the spectra shown in **Figures 2** and **3** are those of direct relevance to observation



**FIGURE 2 |**  $^{51}\text{V}$  NMR (78.9 MHz) spectra are shown of solution of decavanadate (10 mM  $\text{V}_{10}$ , 100 mM V-atoms). The samples are from the bottom up diluted  $\text{V}_{10}$  stock solution (100 mM V-atom) at pH 3.1; 10 mM  $\text{V}_{10}$  in the presence of 0.48 mM and 0.97 mM citrate at pH 2.8 and 2.2, respectively; 10 mM  $\text{V}_{10}$  in the presence of 24 mM  $\text{P}_i$  at pH 6.9; and finally 10 mM  $\text{V}_{10}$  in the presence of Middlebrook 7H9 broth medium supplemented with 10% ADC enrichment (5% BSA, 2% dextrose, 5% catalase), glycerol (0.2%, v/v) and Tween 80 (0.05%, v/v) recorded both in the absence and the presence of a capillary reference of 100 mM  $\text{Na}_3\text{VO}_4$ .




**FIGURE 3 |**  $^{51}\text{V}$  NMR (78.9 MHz) spectra are shown of solution of colorless oxovanadate (40 mM  $\text{V}_1$ , 40 mM V-atoms). The samples are from the bottom up diluted  $\text{V}_1$  stock solution (40 mM V-atom) at pH 8.3; 10 mM  $\text{V}_{10}$  in the presence of 24 mM  $\text{P}_i$  at pH 8.1; 10 mM  $\text{V}_{10}$  in the presence of 0.48 mM citrate at pH 6.3, and finally 10 mM  $\text{V}_{10}$  in the presence of Middlebrook 7H9 broth medium supplemented with 10% ADC enrichment (5% BSA, 2% dextrose, 5% catalase), glycerol (0.2%, v/v) and Tween 80 (0.05%, v/v) recorded both in the absence and the presence of a capillary reference of 100 mM  $\text{Na}_3\text{VO}_4$ . The spectrum labeled Reference is of the capillary reference alone (top spectrum). The key to the signals: V-oligomers,  $\text{V}_1$  monomer;  $\text{V}_2$ , dimer;  $\text{V}_4$ , tetramer;  $\text{V}_5$ , pentamer; VCit, V-citrate complex; PV, vanadate-phosphate complex.

and assignment of the species that form in the growth media (see below **Figure 4**). Importantly, all the spectra are referenced against a reference sample (labeled Ref. in **Figure 3**, containing 100 mM  $\text{Na}_3\text{VO}_4$ ).

In **Figure 2**, we show the spectra of decavanadate under various conditions. First, we show the spectrum of the stock solution at pH 3.1 with the three characteristic signals for the three different V-atoms in the  $\text{V}_{10}$  molecule. Next, we show two spectra with added citrate at two concentrations at pH 2.8 and 2.2 where the decavanadate will form VCit complex (two signals at  $-548$  and  $-552$  ppm). The next three spectra at higher pH values are shown first in the presence of phosphate ( $\text{P}_i$ ), demonstrating that the signals for  $\text{V}_{10}$  shifts as the pH increases from 3 to 6.9. As shown in the final two spectra in **Figure 2**,  $\text{V}_{10}$  in the presence of media that contains both citrate and  $\text{P}_i$  did not form any other V-products when all V-atoms are in the form of  $\text{V}_{10}$ . The spectrum of  $\text{V}_{10}$  in media is shown both in the presence and absence of a capillary tube filled with  $\text{Na}_3\text{VO}_4$  and used to reference the spectra (**Figure 2**). All the NMR spectra were referenced in this manner and the other NMR spectra are provided in the **Supplemental Material**. In summary, these spectra shown that when vanadium is present in the form of  $\text{V}_{10}$ , other vanadium species do not form in the cell media.

The spectra shown in **Figure 2** demonstrated that the  $\text{V}_{10}$  is stable in the growth media, which is consistent with studies in yeast media reported previously (Willsky et al., 1984a, 1985). The data shown in **Figure 2** demonstrated that at pH 2, the  $\text{V}_{10}$  forms complexes with citrate but as the pH increased to near neutral pH,  $\text{V}_{10}$  did not form such products with growth media components (Ehde et al., 1989; Zhou et al., 1999; Crans, 2000; Kaliva et al., 2002, 2003; Chen et al., 2007, 2014; Lodyga-Chruscinska et al., 2008). In contrast, when  $\text{V}_{10}$  is converted to the oxovanadates or  $\text{V}_1$  is the starting form at vanadium, the VCit and PV complexes are formed (this is generally observed in solutions at neutral or more basic pH values, **Figure 3**). However, if the pH of the growth media decreases (that is the acidity increase and pH value decrease),  $\text{V}_{10}$  hydrolyzes and forms oxometalates that are able to react with media components (see **Figures 2, 3**).

The spectra of solutions from metavanadate ( $\text{NaVO}_3$ ) containing vanadate monomer and other oxovanadates prepared in aqueous solution and in media are shown in **Figure 3**. Since vanadate solutions contain rapidly converting species (Crans et al., 1990) and the composition depends on pH and concentration, the species present is dependent on the solution composition. The first spectrum shows a solution containing the interconverting oxovanadates,  $\text{V}_1$ ,  $\text{V}_2$ ,  $\text{V}_4$ , and  $\text{V}_5$ . In the presence of  $\text{P}_i$  and at slightly lower pH value, the  $\text{V}_1$  signal shifted and the linewidth increased, which is indicative of the formation of a vanadate-phosphate complex (abbreviated PV on the spectra in **Figure 2**) (Gresser et al., 1986; Andersson et al., 2005). Although there are no X-ray structures for these species, the Gresser-Tracey team proposed that there is both a vanadium species that is four or five-coordinate and octahedral based on the chemistry shifts). The formation of the PV species (Gresser et al., 1986; Andersson et al., 2005) is rapid on the NMR time scale, and thus results in shifting of the vanadate monomer ( $\text{V}_1$ ) signal in place of observing two separate signals one for  $\text{V}_1$  and one for PV. As the pH decreased to 6.3,  $\text{V}_1$  protonates and the monomer signal shifts to about  $-560$  ppm. At these higher pH values, the



**FIGURE 4** | The growth curve is shown for treatment with vanadate (left, prepared from a 40 mM colorless metavanadate solution) and decavanadate [right, prepared from a 100 mM orange decavanadate solution (1.0 M V-atoms) on *M. tb*].

**TABLE 2** | The EC<sub>50</sub> values for V<sub>1</sub> and V<sub>10</sub> treated *Mycobacterium tuberculosis* and *smegmatis*.

EC <sub>50</sub> ( <i>M. tb</i> ) (mM)	Stand. error	EC <sub>50</sub> ( <i>M. tb</i> ) (mM V-atoms)	Stand. error	EC <sub>50</sub> ( <i>M. smeg</i> ) (mM)	Stand. error	EC <sub>50</sub> ( <i>M. smeg</i> ) (mM V-atoms)	Stand. error	
V <sub>1</sub>	2.0	0.43	2.0	0.43	0.19	0.071	0.19	0.071
V <sub>10</sub>	0.029	0.005	0.29	0.05	0.0037	0.0004	0.037	0.004

VCit complex is found to form in solutions containing both V<sub>1</sub> and citrate. Many different VCit complexes are known, however, the broad doublet signal observed for the VCit could be several structures as described in the literature (Ehde et al., 1989; Zhou et al., 1999; Crans, 2000; Kaliva et al., 2002, 2003; Chen et al., 2007, 2014; Lodyga-Chruscinska et al., 2008). The final spectra show the addition of the vanadate stock solution to media, which at pH 6.9 results in a solution that contains a signal of V<sub>1</sub> and PV, V<sub>2</sub>, and V<sub>4</sub> and a trace of V<sub>5</sub>. The amount of VCit formed is low and it is difficult to see the signal from the normal growth media spectrum, so we used an increased amplification to show the VCit signal and part of such spectrum is shown in the spectral insert.

In summary, if the growth of the mycobacteria is conducted near pH 6.8, the speciation of vanadium (V) will readily reach thermodynamic control. Solutions prepared from NaVO<sub>3</sub> will contain little to no V<sub>10</sub> and solutions containing V<sub>10</sub> will be relatively stable. Therefore, growth studies will be carried out with either V<sub>1</sub> or V<sub>10</sub> species, and it will be possible to determine if there is a difference in the growth effects on mycobacteria by these two species. That is the growth effects of the V<sub>1</sub> with the oligomeric oxovanadates and V<sub>10</sub> are measured in media with known vanadium speciation and under conditions where it is possible to observe the effect of each species.

## Growth Inhibition Experiments of Vanadate Monomer and Decavanadate

Growth inhibition experiments were designed to measure the effects of V<sub>10</sub> and V<sub>1</sub> each on *M. tb* mc<sup>2</sup> 6230 and *M. smeg* mc<sup>2</sup> 155. The measurements monitor growth using absorbance at 600 nm and the concentration of the vanadium compound

was changing with 3-fold dilution experiments. In this manner, the growth of the bacteria was measured over a 2,000-fold range of concentrations of vanadium compound. The results with the *M. tb* are shown in Figure 4, and the EC<sub>50</sub> (concentration of compound where growth is reduced by half) value calculated for the V<sub>1</sub> experiment was 2.0 mM (2.0 mM V-atoms) and for the V<sub>10</sub> experiment 29 μM (0.29 mM V-atoms, Table 2).

These results show that V<sub>10</sub> is a more potent inhibitor than V<sub>1</sub> by a factor of 100 in terms of concentration and a factor of 10 in terms of concentration of V-atoms. This result demonstrates that the large V<sub>10</sub> anion is more inhibitory than the V<sub>1</sub> atom even when counting for the fact that there are 10 V-atoms in each V<sub>10</sub> species. As discussed below, <sup>51</sup>V NMR spectroscopy was used to examine the speciation in the solutions and verify that the V<sub>10</sub> solution indeed contained V<sub>10</sub> (see below for details).

The growth effects were also measured for *M. smeg* mc<sup>2</sup> 155 for both V<sub>10</sub> and V<sub>1</sub> (data not shown). Similarly, the EC<sub>50</sub> values were calculated for the V<sub>1</sub> experiment to be 190 μM (0.190 mM V-atoms) and for the V<sub>10</sub> experiment to be 3.7 μM (0.037 mM V-atoms). These results show that the V<sub>10</sub> is a more potent inhibitor than V<sub>1</sub>, by a factor of 100 in terms of concentration and a factor of 10 in terms of concentration of V-atoms as we also observed for *M. tb*.

These results present evidence that the large compact V<sub>10</sub>-anion is a stronger inhibitor than V<sub>1</sub> of cell growth of the two mycobacterial strains. The more potent inhibition by V<sub>10</sub> compared to V<sub>1</sub> shows that the V<sub>10</sub> is an inhibitor, and that it is not V<sub>1</sub> formed from hydrolyzed V<sub>10</sub>. However, it should be mentioned that it is solutions of V<sub>10</sub> that are causing the inhibitory growth effect, and we cannot rule out the possibility that the effect is caused by several species including the hydrolysis

products, or some mechanism in which  $V_{10}$  delivers the V-atom to the cells. Regardless, these studies demonstrate the effects of simple oxovanadates and the large decavanadate. The fact that  $V_{10}$  has a greater effect than  $V_1$  (or 10  $V_1$  molecules) supports the interpretation that the growth effects observed are not due to  $V_1$  formed from hydrolyzed  $V_{10}$ . To support this interpretation, we acquired the  $^{51}\text{V}$  NMR spectra of the growth media solutions at various times during the growth experiment.

## Speciation Studies in Media of Mycobacteria Treated With Vanadate and Decavanadate

### $^{51}\text{V}$ NMR Spectra of Media That Have Grown Mycobacteria and Have Been Treated With Metavanadate


$^{51}\text{V}$  NMR spectra were recorded at several concentration and time-points in the growth media in which *M. smeg* had grown because as mentioned above the bacteria were subjected to a 2000-fold concentration range. In **Figure 5**, we show the  $^{51}\text{V}$  NMR spectra recorded at 3.3 mM and 10 mM oxometalates,

respectively. Both these series show the presence of the different oxometalates ( $V_1$ ,  $V_2$  and  $V_4$ ). In addition, neither series show any evidence for change as the experiment proceeded. This is somewhat surprising, because experiments with other cellular systems, such as yeast, fungi and red blood cells, all showed significant signs of signal reduction (Willsky et al., 1984b, 1985; Crans et al., 2002; Delgado et al., 2005; Jakusch et al., 2014). Since the  $V_1$  or oxovanadate is not a strong inhibitor of growth, it is possible that the simple vanadate system is not getting into the *M. smeg* cells very effectively.


### $^{51}\text{V}$ NMR Spectra of Media in Which Mycobacteria Have Been Treated With Decavanadate

$^{51}\text{V}$  NMR spectra were recorded at several concentrations and time-points in the growth media in which *M. smeg* had grown. In **Figure 6**, we show the  $^{51}\text{V}$  NMR spectra recorded at 3.3 and 10 mM, respectively. Both series show the presence of  $V_{10}$ . Although the stock solutions and growth media without bacterial cells contains 100%  $V_{10}$  (**Figure 2**), the addition of growth media containing cells immediately caused some decomposition of  $V_{10}$  (**Figure 6**). NMR spectra were recorded multiple times of the 3.3 mM V and resulted in approximately a 1:1 ratio of  $V_{10}$  to oxovanadates as determined by integration of the spectra. As shown in **Figure 6**, after 1 h the composition of the sample was very similar to the content at the beginning of treatment, whereas at the 5 h time point the  $V_{10}$  had decreased significantly and at 24 h very little was left. This demonstrates that the  $V_{10}$  is not stable in the media in the presence of the *M. smeg* cells at 33 mM total V-atom concentration even though the pH is between 5.8 and 6.8 which is a pH range where there should not have been any hydrolysis of  $V_{10}$  according to the known solution chemistry of  $V_{10}$ . However,  $V_{10}$  treatment may cause signal reduction or the  $V(\text{V})$  may interact with large structural components in the mycobacterial cells (or their excretion products) and the  $^{51}\text{V}$  NMR signal has broadened beyond detection. Precedent for these possibilities has been reported previously (Willsky et al., 1984b, 1985; Crans et al., 2002). In the series of spectra at 10 mM  $V_{10}$  in the media, there is less evidence for  $V_{10}$  removal is apparent at the different time points. However, when we integrated the signals a slow decrease in the  $V_{10}$  signals compared to the oxovanadate signals was clear. The ratio at time zero was 1:0.71, at time 1 h it was 1:1.08; at time 5 h it was 1:1.18 and at 24 h it was 1:1.24. Thus, providing evidence for a slow decrease in  $V_{10}$  content as the experiment progressed. This data is consistent with the possibility that there is some reduction of the vanadium from the POM resulting in hydrolysis and formation of the oxovanadates.

To further investigate whether there was any reduction taking place upon vanadium treatment, we also measured the color change in these samples, **Table 3**. More data and the pH measurements on all the samples shown are provided in the **Supplemental Material**, and the pH values are all in the range from 6.1 to 7.4. Vanadium(V) on reduction is known to change from colorless ( $V_1$  oxovanadates) or yellow/orange ( $V_{10}$ ) to a green color (Crans et al., 2004, 2010, 2013; Pessoa and Tomaz, 2010; Jakusch et al., 2014). The 7H9 media alone did not cause any reduction as evidenced by the media sample maintained the color prior to treatment, **Table 3**. All the samples containing



**FIGURE 5** | The  $^{51}\text{V}$  NMR spectra of media in which *M. smeg* had grown at 0, 5, and 24 h time points. Data for two concentrations are shown in (A) 3.3 mM  $V_1$  treatment (3.3 mM V-atoms) and (B) 10 mM  $V_1$  treatment (10 mM V-atoms). See **Figure 3** caption for key to labeling.



**FIGURE 6 |** The  $^{51}\text{V}$  NMR spectra of media in which *M. smeg* had grown at 0, 1, 5, and 24 hr time points. Data for two concentrations are shown in (A) 3.3 mM  $\text{V}_{10}$  treatment (33 mM V-atoms) and (B) 10 mM  $\text{V}_{10}$  treatment (100 mM V-atoms).

media and *M. smegmatis* treated with  $\text{V}_{10}$  developed a greenish tint as the treatment progressed beyond 5 h time points, whereas the samples treated with  $\text{V}_1$  remained colorless, see Table 3. This is consistent with reduction of some of the  $\text{V}(\text{V})$  in the  $\text{V}_{10}$  salt. Interestingly, similar reduction was not observed for the samples treated with the  $\text{V}_1$  solutions. These observations are consistent with the selective reduction of the  $\text{V}(\text{V})$  and that the reduction was more prevalent after longer treatments, higher concentrations of vanadium and  $\text{V}_{10}$  compared to  $\text{V}_1$ . Since the reduction only took place in the presence of  $\text{V}_{10}$ , this anion induced a response that was not induced by  $\text{V}_1$ . Furthermore, the supernatant of a batch of grown *M. smeg* bacteria and centrifuging down the bacteria and removing them was collected to investigate if the bacteria excrete a component responsible for the observed reduction. The  $\text{V}_{10}$  and  $\text{V}_1$  samples were added to this supernatant and the data are shown in column 4 in Table 3. This supernatant also generates the color change from orange to green upon incubation. This experiment showed that the reduction is something that was induced by a material excreted from the *M. smeg* cells during growth. This suggests that the reduction is likely to come from the excretion

of siderophores (Hider and Kong, 2010) or proteins such as the reported protein tyrosine phosphatases from mycobacteria (Beresford et al., 2009; Dutta et al., 2016). Since we also carried out such studies of supernatant that had been heated (see **Supplemental Material**), such a product is either a very heat stable protein or a siderophore-type of material.

Combined these studies show that there is selectivity in the effects of interactions of vanadium with mycobacteria, and that for these systems the most effective form is not the  $\text{V}_1$ , which is a potent phosphatase inhibitor and a common active form of vanadium. These studies show that the  $\text{V}_{10}$ , a POM is more potent inhibitor of growth of mycobacteria than the well-known phosphatase inhibitor,  $\text{V}_1$ . The studies also demonstrate that there is some redox chemistry involved in this process and that the mycobacteria excrete a component that interacts with  $\text{V}_{10}$  and cause the redox chemistry and hydrolysis of the  $\text{V}_{10}$ . Eventually the  $\text{V}_1$  samples would also show the color change consistent with reduction, but this required growth periods beyond several weeks and this process may be different than the reduction, of  $\text{V}_{10}$ .

## Modeling Speciation Vanadium species Distribution in Mycobacteria Media Added Metavanadate

To quantify the  $\text{V}$ -species that are present in the growth assay media, we carried out speciation analyses modeling experimentally the distribution of vanadium species in the media. This analysis is based on analyzing solutions containing  $\text{V}$ -species that are governed by thermodynamics and can be calculated the HySS program (Alderighi et al., 1999). This model can analyze the experimental conditions found in solution prepared by the addition of metavanadate and exposed to the conditions of the cell growth assay. We used the reactions reported previously with regard to the exchange of labile oxovanadates (Pettersson et al., 1983, 1985; Gresser et al., 1986; Ehde et al., 1989; Crans et al., 1990, 2000; Crans, 2000, 2005; Andersson et al., 2005; Baruah et al., 2006; Aureliano and Crans, 2009), the formation of the  $\text{Vcit}$  complex (Ehde et al., 1989; Crans, 2000), the formation of the  $\text{PV}$  complex (Gresser et al., 1986; Andersson et al., 2005), and the thermodynamics relating to the decavanadate deprotonation reactions (Crans, 2005; Baruah et al., 2006; Aureliano and Crans, 2009). The known speciation parameters used for vanadate at various pH values were measured in the presence of 0.6 M NaCl to keep the salt concentration constant for all the components in the system. As shown in Figure 5, the species composition for the conditions of three different concentrations of vanadium (0.0050, 3.3, and 10 mM) are illustrated. As discussed above, the vanadium samples prepared with decavanadate will not contain this speciation distribution because all  $\text{V}(\text{V})$  was converted to  $\text{V}_{10}$ . The addition of this solution to the media will only slowly hydrolyze the  $\text{V}_{10}$  species to the equilibrium oxovanadate mixture if above or near pH 7.

The speciation analysis of the conditions observed in the media with freely exchangeable oxovanadate shown in Figure 3 demonstrate that an observable amount of  $\text{VCit}$  complex forms. Depending on the concentration of the vanadium, the contribution of  $\text{VCit}$  is larger percentage-wise at the lower

**TABLE 3** | Samples treated with V<sub>10</sub> and V<sub>1</sub> in 7H9 media, *M. smeg* cultures and supernatant after *M. smeg*.

Time	media7H9	<i>M. smeg</i> in 7H9 media	Supernatant after <i>M. smeg</i> growth (cells removed)
0			
1			
5			
24			

Growth (columns 2, 3, and 4). The yellow or orange V<sub>10</sub> is shown from the left with 10 and 3.3 mM samples, and the colorless V<sub>1</sub> samples from the left 10 and 3.3 mM in each column. When reduction took place the green color is evident [vanadium (IV)]. See **Supplemental Material** for the pH values of each of these samples.


concentrations of vanadate; for example, at 0.005 and 1.0 mM V-atoms at neutral pH the amount is 75–80% (in terms of mole fraction) of VCit complex. In contrast, at 10 mM V-atoms at neutral pH the amount is ~10%. However, if the amounts are calculated, this would correspond to 0.0004, and 1.0 mM VCit complex in the media. These observations are consistent with the very strong formation constant for VCit. The PV complex, however, is much less stable, and even though there is much more phosphate in the assay, the PV complex is only observed in a significant concentration at high V and P<sub>i</sub> concentrations (5 and 24 mM, respective). Although the PV complex is only observed in high concentrations at mM V-treatments the shifting of the V<sub>1</sub> signal attests to the presence of the PV adduct in the assays with *M. tb* and *M. smeg*.

The results shown in **Figure 7** are in general agreement with the experimental data obtained in **Figures 2, 3, 5, 6** with regard to formation of the VCit and PV complexes. **Figure 7** shows that there are four different VCit complexes formed at low vanadium concentration, but at higher concentration and in the presence of phosphate two major VCit complexes formed. There are several assumptions on which these estimations are based, including the differences in ionic strengths, the low concentration of the vanadium used, and the fact that some reduction takes place during the growth experiment to be changing the concentrations somewhat of the VCit and PV complexes. Regardless, our general considerations demonstrate that VCit and PV complexes should form as predicted by the speciation

analysis (Pettersson et al., 1983; Ehde et al., 1989; Selling et al., 1994) and reflect the equilibrium mixture observed by NMR analysis.

## DISCUSSION

Vanadium derivatives have been considered as therapeutic agents since 1899 (Lyonnet et al., 1899; Crans et al., 2018). Recent studies demonstrate that vanadium compounds and salts are effective antidiabetic agents (Sakurai, 2002; Thompson and Orvig, 2006; Kiss et al., 2008; Smith et al., 2008; Thompson et al., 2009; Willsky et al., 2011, 2013; Crans, 2015; Trevino et al., 2015; Boulmier et al., 2017; Levina and Lay, 2017b) but their therapeutic properties have been expanded for these materials to be anticancer agents (Lu and Zhu, 2011; Leon et al., 2017; Medina et al., 2017; Crans et al., 2018), immunotherapy agents (Lu and Zhu, 2011; Selman et al., 2018) as well as treatment for other diseases (Zwolak, 2014; Pessoa et al., 2015). Many of the compounds are coordination complexes, which form vanadate during processing after administration (Thompson et al., 2009; Willsky et al., 2011). Considering the ease with which vanadium undergoes redox chemistry (Crans et al., 2010), it is known that some of the vanadium is converted from V(V) to lower oxidation states upon administration (Thompson et al., 2009; Willsky et al., 2011). Furthermore, studies were carried out on the salts prior to investigation of coordination complexes (Sakurai et al., 1990; Goldfine et al., 2000; Willsky et al., 2001,



**FIGURE 7** | The evaluation of the speciation at different vanadium concentration using HySS Program (vs. 2009) (Alderighi et al., 1999). The speciation diagram shown was calculated at several vanadium (V-atom) concentrations in the presence of 0.48 mM citrate and 24 mM phosphate found in the growth media and **(A)** 5.0  $\mu\text{M}$  vanadate, **(B)** 3.3 mM vanadate, and **(C)** 10 mM vanadate. Note, the concentrations of different species regardless of nuclearity are here shown in terms of V-atoms.

2013; Smith et al., 2008). The biological activity of homoxovanadates have been of considerable interest considering the varied biological effects depending on the specific vanadium species (Sakurai et al., 1990; Willsky et al., 2001; Postal et al., 2016). Information on cellular uptake by vanadium species is

of interest, and the studies presented here compare the effect of monomeric vanadate and decavanadate. Such direct comparison is important to understand the interconversion between these species, and despite the many reports in the literature (reviewed in Aureliano and Crans, 2009), no similar study showing data for

direct comparison between the two different species have been reported.

Cellular studies have been reported in different types of organisms that consider the uptake of vanadate. Willsky reported studies with yeast (Willsky et al., 1984b, 1985; Crans et al., 2002), and concluded that  $V_1$  was the species entering the cells; however, upon entry processing took place; forming decavanadate in the lysosomes, and forming V(IV) through redox processes. The treatment of the yeast cells with vanadate solutions containing mainly vanadate monomer and oligomers is evidenced by  $^{51}V$  NMR spectroscopy and EPR spectroscopy. These studies were very elegant and truly important, because they documented the formation of decavanadate in cells from monomeric vanadate for the first time (Willsky et al., 1984b). Recently, Zakrzewska and Zivic reported a series of studies in fungi exploring the uptake of both vanadate and V(IV) using  $^{51}V$  and EPR spectroscopy, as well as polarographic studies (Zizic et al., 2013, 2016; Hadzibrahimovic et al., 2017). The initial  $^{51}V$  NMR studies on *Phycomyce blakesleeanus* mycelium were supplemented with polarography, which allowed for the assignment of the signal at  $-535$  ppm accumulating inside the fungi cells to  $V_1$  even though the pH of the system was defined by HEPES buffer at 7.2 where  $V_1$  generally has a peak position at a higher frequency (Zizic et al., 2016). The authors attribute the signal shifting to intracellular complexation of  $V_1$ . Alternatively, these observations are consistent with a hydrophobic environment having been reported in some cases to alter composition and speciation and impact the shifting of observed signals. Combined, these studies showed convincingly that the cell wall is not responsible for reduction of vanadate (Hadzibrahimovic et al., 2017) and these studies also confirm that tetrameric vanadate is not able to enter the cells (Zizic et al., 2016) as reported in yeast as well (Crans et al., 2002) (Willsky et al., 1984a,b).

Cellular studies in mammalian systems have also been carried out, and many of these studies are linked to the stability and speciation of vanadium compounds in blood (Delgado et al., 2005; Zhang et al., 2006; Li et al., 2009; Jakusch et al., 2011, 2014; Willsky et al., 2013; Sanna et al., 2014, 2017; Jakusch and Kiss, 2017). Although bacterial cells do not contain blood, they do infect mammals and thus the vanadium chemistry in blood is relevant to bacterial infections as well. These studies include detailed investigations into various complexes, and projections based on studies in model systems what would be observed in mammalian systems and *in vivo*. Although such studies remain model studies, they are important because they begin to provide a view of how it will look like in the *in vivo* cells. Studies were also reported by Garner and coworkers using  $^1H$  NMR spin echo and  $^{51}V$  NMR spectroscopy to characterize the uptake of vanadate and subsequent reduction to form V(IV) in erythrocytes (Garner et al., 1997). The reduction of the vanadate was attributed to glutathione as demonstrated in a time-dependent response using both  $^{51}V$  NMR and  $^1H$  spin echo NMR (Garner et al., 1997). The addition of the blocking agent 4,4'-diisothio-cyanostilbene-2,2b-disulfonic acid (DIDS) specifically blocks the anion transporter preventing the vanadate from entering the erythrocytes. Adding the blocker prevented

vanadate entering, and the depletion of glutathione was arrested. Thus, depletion of intracellular glutathione could be correlated with entering vanadate and the authors proposed that the reduced vanadium [presumably V(IV)] formed complexed with cellular components, because no or little V(IV) formed (Garner et al., 1997).

These studies stand in direct contrast to studies in which decavanadate is associated with proteins and have been characterized by X-ray crystallography (Winkler et al., 2017). However, consideration of both solution and solid state is necessary to characterize both the effects of decavanadate and simple oxovanadates. The studies shown here demonstrate three important findings. First, the fact that decavanadate is a potent inhibitor of growth of two mycobacterial species and more so than monomeric vanadate, which is a known potent phosphatase inhibitor (Crans, 2015; McLauchlan et al., 2015). Indeed, phosphatase inhibition has recently been reported by  $V_{10}$  of the Leishmania acid phosphatase as well as earlier reports (Aureliano and Crans, 2009; Dorsey et al., 2018). Second, the fact that monomeric vanadate is not as potent inhibitor for *M. tb* and *M. smeg*. Third, these studies also demonstrate that under conditions in which the decavanadate would remain stable, in the presence of *Mycobacterium* spp. the decavanadate undergoes hydrolysis. These combined observations suggest that some type of mechanism that involves the bacterium (or something excreted by the bacterium) interacting with the decavanadate and causes the conversion to oxovanadates. This is particularly interesting because similar response is not observed in the media without bacteria.

## CONCLUSION

The studies presented demonstrate that decavanadate ( $V_{10}$ ) inhibits the growth of two mycobacterial species, *M. tb* and *M. smeg*, whereas the oxovanadates prepared from  $NaVO_3$  inhibit growth with 10-100-fold less potency. The inhibition was observed in media containing citrate and phosphate, resulting in the formation of a VCit complex and PV complexes. However, neither of these complexes appeared to interfere with the observed inhibition for  $V_{10}$  and the inhibition by  $V_1$  was still 10-100-fold less. The greater inhibitory selectivity of  $V_{10}$  (or a component of  $V_{10}$ ) result is important because it demonstrates that simple vanadium salts can have different effects on cells, and that a vanadium compound other than the potent phosphatase inhibitor, monomeric vanadate ( $V_1$ ), can inhibit growth. Importantly, these studies suggest that other oxometalates may have the ability to inhibit growth of mycobacteria and other pathogens.

In addition, these studies demonstrated that mycobacteria or some component excreted by the mycobacteria which catalyzes the hydrolysis of  $V_{10}$  and the same effect was not observed in media in the absence of growing cells. This process is particularly interesting because it implies that the  $V_{10}$  interacts with the some cell component, possibly through some mechanism involving  $V_{10}$  delivering a V-atom to such a cell component. This type of mechanism is novel and further investigation into such process

will be of general interest and important for the mode of action of POMs in biological systems.

## AUTHOR CONTRIBUTIONS

NS carried out the  $^{51}\text{V}$  NMR studies, prepared the figures and wrote part of the manuscript. ZA carried out the growth studies with vanadate and decavanadate and recorded some of the NMR spectra. ZA also collected the experiments with the colors of the various media. HM did the speciation calculations. SK and DCri advised and oversaw the biological studies. DCra combined the chemical and biological components of the work, advised and oversaw the NMR studies and wrote the manuscript with the assistance of NS. All authors contributed to the editing of the manuscript.

## FUNDING

NS was funded by Rajabhat Rajanagarindra University to take a summer leave to carry out research. ZA was funded by a fellowship from Libyan Ministry of Education. Supplies were

obtained in part from funds from the Arthur P. Cope Foundation administered by the American Chemical Society (to DCC) and the NSF grant (CHE-1709564) (to DCri and DCra).

## ACKNOWLEDGMENTS

We thank the Core Facilities at Colorado State University for their assistance in carrying out the work described in this manuscript. NS thanks Rajabhat Rajanagarindra University for the support to travel and carry out research.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fchem.2018.00519/full#supplementary-material>

More figures are included in the supplementary materials showing additional spectra supporting the results presented in the manuscript and include NMR experiments, growth experiments and additional information on the speciation calculations.

## REFERENCES

Alderighi, L., Gans, P., Ienco, A., Peters, D., Sabatini, A., and Vacca, A. (1999). Hyperquad simulation and speciation (HySS): a utility program for the investigation of equilibria involving soluble and partially soluble species. *Coor. Chem. Rev.* 184, 311–318. doi: 10.1016/S0010-8545(98)00260-4

Andersson, I., Gorzsas, A., Kerezsi, C., Toth, I., and Pettersson, L. (2005). Speciation in the aqueous  $\text{H}^+/\text{H}_2\text{VO}_4^-\text{H}_2\text{O}_2/\text{phosphate}$  system. *Dalton Trans.* 22, 3658–3666. doi: 10.1039/b508273k

Auerbach-Nevo, T., Zarivach, R., Peretz, M., and Yonath, A. (2005). Reproducible growth of well diffracting ribosomal crystals. *Acta Crystallogr. Sec. D* 61, 713–719. doi: 10.1107/S0907444905006311

Aureliano, M. (2014). Decavanadate contribution to vanadium biochemistry: *in vitro* and *in vivo* studies. *Inorg. Chim. Acta* 420, 4–7. doi: 10.1016/j.ica.2013.10.010

Aureliano, M. (2016). Decavanadate toxicology and pharmacological activities: V-10 or V-1, both or none? *Oxid. Med. Cell Longev.* 2016:6103457. doi: 10.1155/2016/6103457

Aureliano, M., and Crans, D. C. (2009). Decavanadate ( $\text{V}_{10}\text{O}_{28}^{6-}$ ) and oxovanadates: oxometalates with many biological activities. *J. Inorg. Biochem.* 103, 536–546. doi: 10.1016/j.jinorgbio.2008.11.010

Aureliano, M., Fraqueza, G., and Ohlin, C. A. (2013). Ion pumps as biological targets for decavanadate. *Dalton Trans.* 42, 11770–11777. doi: 10.1039/c3dt50462j

Aureliano, M., and Ohlin, C. A. (2014). Decavanadate *in vitro* and *in vivo* effects: facts and opinions. *J. Inorg. Biochem.* 137, 123–130. doi: 10.1016/j.jinorgbio.2014.05.002

Aureliano, M., Ohlin, C. A., Vieira, M. O., Marques, M. P. M., Casey, W. H., and De Carvalho, L. (2016). Characterization of decavanadate and decanioate solutions by Raman spectroscopy. *Dalton Trans.* 45, 7391–7399. doi: 10.1039/C5DT04176G

Baes, C. F. (1976). *The Hydrolysis of Cations*. New York, NY: John Wiley & Sons.

Baruah, B., Roden, J. M., Sedgwick, M., Correa, N. M., Crans, D. C., and Levinger, N. E. (2006). When is water not water? Exploring water confined in large reverse micelles using a highly charged inorganic molecular probe. *J. Am. Chem. Soc.* 128, 12758–12765. doi: 10.1021/ja0624319

Bashan, A., and Yonath, A. (2008). The linkage between ribosomal crystallography, metal ions, heteropolytungstates and functional flexibility. *J. Mol. Struct.* 890, 289–294. doi: 10.1016/j.molstruc.2008.03.043

Bbl<sup>tm</sup>, D. (2018). “Middlebrook 7H9 Broth • Middlebrook 7H9 Broth with Glycerol • Middlebrook 7H9 Broth with Polysorbate 80 • Middlebrook ADC Enrichment.” 2nd ed.

Beresford, N., Mulhearn, D., Szczepankiewicz, B., Liu, G., Johnson, M. E., Fordham-Skelton, A., et al. (2009). Inhibition of MptpB phosphatase from *Mycobacterium tuberculosis* impairs mycobacterial survival in macrophages. *J. Antimicrob. Chemother.* 63, 928–936. doi: 10.1093/jac/dkp031

Boulmier, A., Feng, X. X., Oms, O., Mialane, P., Riviere, E., Shin, C. J., et al. (2017). Anticancer activity of Polyoxometalate-Bisphosphonate complexes: synthesis, characterization, *in vitro* and *in vivo* results. *Inorg. Chem.* 56, 7558–7565. doi: 10.1021/acs.inorgchem.7b01114

Carsella, J. S., Sánchez-Lombardo, I.; Bonetti, S. J., and Crans, D. C. (2017). Selenium speciation in the Fountain Creek Watershed correlates with water hardness, Ca and Mg levels *Molecules* 22, 708, 1–16. doi: 10.3390/molecules22050708

Chasteen, N. D. (1983). “The biochemistry of vanadium,” in *Copper, Molybendum, and Vanadium in Biological Systems*, eds M. J. Clarke, J. B. Goodenough, J. A. Ibers, C. K. Jørgensen, D. M. P. Mingos, J. B. Neilands, G. A. Palmer, D. Reinen, P. J. Sadler, R. Weiss, and R.J.P. Williams (New York, NY: Springer-Verlag), 105–138. doi: 10.1007/BFb0111304

Chen, C. Y., Chen, M. L., Chen, H. B., Wang, H. X., Cramer, S. P., and Zhou, Z. H. (2014). alpha-Hydroxy coordination of mononuclear vanadyl citrate, malate and S-citramalate with N-heterocycle ligand, implying a new protonation pathway of iron-vanadium cofactor in nitrogenase. *J. Inorg. Biochem.* 141, 114–120. doi: 10.1016/j.jinorgbio.2014.08.003

Chen, C. Y., Zhou, Z. H., Mao, S. Y., and Wan, H. L. (2007). Asymmetric dinuclear hydroxyl and ethoxyl citrato dioxovanadates(V). *J. Coord. Chem.* 60, 1419–1426. doi: 10.1080/00958970601029420

Clare, B. W., Kepert, D. L., and Watts, D. W. (1973a). Acid decompositon of decavanadate-specific salt effects. *J. Chem. Soc., Dalton Trans.* 22, 2481–2487.

Clare, B. W., Kepert, D. L., and Watts, D. W. (1973b). Kinetic study of the acid decompositon of decavanadate. *J. Chem. Soc. Dalton Trans.* 22, 2479–2487. doi: 10.1039/dt9730002479

Comba, P., and Helm, L. (1988). The solution structure and reactivity of decavanadate. *Helv. Chim. Acta* 71, 1406–1420. doi: 10.1002/hclca.19880710605

Correia, I., Adão, P., Roy, S., Wahba, M., C., M., Maurya, M. R., Marques, F., et al. (2014). Hydroxyquinoline derived vanadium(IV and V) and copper(II) complexes as potential anti-tuberculosis and anti-tumor agents. *J. Inorg. Biochem.* 141, 83–93. doi: 10.1016/j.jinorgbio.2014.07.019

Correia, I., Roy, S., Matos, C. P., Borovic, S., Butenko, N., Cavaco, I., et al. (2015). Vanadium(IV) and copper(II) complexes of salicylaldimines and aromatic heterocycles: Cytotoxicity: DNA binding and DNA cleavage properties. *J. Inorg. Biochem.* 147, 134–146. doi: 10.1016/j.jinorgbio.2015.02.021

Costello, R. L., and Hedgecock, L. W. (1959). Effect of metavanadate ion on the growth *in vitro* of mycobacterium tuberculosis. *J. Bacteriol.* 77, 794–799.

Crans, D., Yang, L., Haase, A., and Yang, X. (2018). Health benefits of Vanadium and its potential as an anticancer agent. *Met. Ions Life Sci.* 18, 251–279. doi: 10.1515/9783110470734-015

Crans, D. C. (2000). Chemistry and insulin-like properties of vanadium(IV) and vanadium(V) compounds. *J. Inorg. Biochem.* 80, 123–131. doi: 10.1016/S0162-0134(00)00048-9

Crans, D. C. (2005). Fifteen years of dancing with vanadium. *Pure Appl. Chem.* 77, 1497–1527. doi: 10.1351/pac200577091497

Crans, D. C. (2015). Antidiabetic, chemical, and physical properties of organic Vanadates as Presumed transition-state inhibitors for Phosphatases. *J. Org. Chem.* 80, 11899–11915. doi: 10.1021/acs.joc.5b02229

Crans, D. C., Bedi, H. S., Li, S., Zhang, B., Nomiya, K., Kasuga, N. C., et al. and Willsky, G.R. (2002). “Tetravanadate, decavanadate, Keggins and Dawson oxotungstates inhibit growth of *S. cerevisiae*” in *Polyoxometalate Chemistry for Nano-Composite Design, [Symposium]* (Honolulu, HI), 181–195.

Crans, D. C., Peters, B. J., Wu, W., and McLauchlan, C. C. (2017). Does Anion-Cation Organization in  $\text{Na}^+$ -containing X-ray crystal structures relate to solution interactions in inhomogeneous nanoscale environments: Sodium-decavanadate in solid state materials, minerals and microemulsions. *Coord. Chem. Rev.* 344, 115–130. doi: 10.1016/j.ccr.2017.03.016

Crans, D. C., Rithner, C. D., Baruah, B., Gourley, B. L., and Levinger, N. E. (2006). Molecular probe location in reverse micelles determined by NMR dipolar interactions. *J. Am. Chem. Soc.* 128, 4437–4445. doi: 10.1021/ja0583721

Crans, D. C., Rithner, C. D., and Theisen, L. A. (1990). Application of time-resolved  $^{51}\text{V}$  2D NMR for quantitation of kinetic exchange pathways between vanadate monomer, dimer, tetramer, and pentamer. *J. Am. Chem. Soc.* 112, 2901–2908. doi: 10.1021/ja00164a009

Crans, D. C., Smee, J. J., Gaidamauskas, E., and Yang, L. Q. (2004). The chemistry and biochemistry of vanadium and the biological activities exerted by vanadium compounds. *Chem. Rev.* 104, 849–902. doi: 10.1021/cr020607t

Crans, D. C., Trujillo, A. M., Pharazyn, P. S., and Cohen, M. D. (2011). How environment affects drug activity: localization, compartmentalization and reactions of a vanadium insulin-enhancing compound, dipicolinatooxovanadium(V). *Coord. Chem. Rev.* 255, 2178–2192. doi: 10.1016/j.ccr.2011.01.032

Crans, D. C., Woll, K. A., Prusinskas, K., Johnson, M. D., and Norkus, E. (2013). Metal speciation in health and medicine represented by iron and vanadium. *Inorg. Chem.* 52, 12264–12275. doi: 10.1021/ic4007873

Crans, D. C., Yang, L. Q., Jakusch, T., and Kiss, T. (2000). Aqueous chemistry of ammonium (dipicolinato)oxovanadate(V): the first organic vanadium(V) insulin-mimetic compound. *Inorg. Chem.* 39, 4409–4416. doi: 10.1021/ic9908367

Crans, D. C., Zhang, B., Gaidamauskas, E., Keramidas, A. D., Willsky, G. R., and Roberts, C. R. (2010). Is Vanadate reduced by thiols under biological conditions? Changing the redox potential of  $\text{V}(\text{V})/\text{V}(\text{IV})$  by complexation in aqueous solution. *Inorg. Chem.* 49, 4245–4256. doi: 10.1021/ic100080k

David, S., Barrosa, V., Cruzb, C., and Delgado, R. (2005). *In vitro* effect of free and complexed indium(III) against Mycobacterium tuberculosis. *FEMS Microbiol. Lett.* 251, 119–124. doi: 10.1016/j.femsle.2005.07.044

Delgado, T. C., Tomaz, A. I., Correia, I., Pessoa, J. C., Jones, J. G., Geraldes, C. F., et al. (2005). Uptake and metabolic effects of insulin mimetic oxovanadium compounds in human erythrocytes. *J. Inorg. Biochem.*, 99, 2328–2339. doi: 10.1016/j.jinorgbio.2005.08.014

Dorsey, B. M., McLauchlan, C. C., and Jones, M. A. (2018). Evidence that speciation of oxovanadium complexes does not solely account for inhibition of Leishmania Acid Phosphatases. *Front. Chem.* 8:109. doi: 10.3389/fchem.2018.00109

Druskovich, D. M., and Kepert, D. L. (1975). Base decomposition of decavanadate. *J. Chem. Soc. Dalton Trans.* 22, 947–951. doi: 10.1039/dt9750000947

Dutta, N. K., He, R., Pinn, M. L., He, Y., Burrows, F., Zhang, Z.-Y., et al. (2016). Mycobacterial protein tyrosine phosphatase A and B inhibitors augment the bactericidal activity of the standard anti-tuberculosis regimen. *ACS Infect. Dis.* 2, 231–239. doi: 10.1021/acsinfecdis.5b00133

Ehde, P., Andersson, I., and Pettersson, L. (1989). Multicomponent polyanions. 43. A study of aqueous equilibria in the vanadocitrate system. *Acta Chem. Scand.* 43, 136–143. doi: 10.3891/acta.chem.scand.43-0136

Esbak, H., Enyedy, E. A., Kiss, T., Yoshikawa, Y., Sakurai, H., Garribba, E., et al. (2009). Aminoacid-derivatised picolinato-oxovanadium(IV) complexes: characterisation, speciation and *ex vivo* insulin-mimetic potential. *J. Inorg. Biochem.* 103, 590–600. doi: 10.1016/j.jinorgbio.2008.11.001

Evans, H. T. Jr. (1966). The molecular structure of the isopoly complex ion, decavanadate ( $\text{V}_{10}\text{O}_{28}^6$ ). *Inorg. Chem.* 5, 967–977. doi: 10.1021/ic50040a004

Fraqueza, G., De Carvalho, L., Marques, M. P. M., Maia, L., Ohlin, C. A., Casey, W. H., et al. (2012). Decavanadate, decaniobate, tungstate and molybdate interactions with sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase: quercentin prevents cysteine oxidation by vanadate but does not reverse ATPase inhibition. *Dalton Trans.* 41, 12749–12758. doi: 10.1039/c2dt31688a

Fu, D. Y., Zhang, S. M., Qu, Z. Y., Yu, X. H., Wu, Y. Q., and Wu, L. X. (2018). Hybrid assembly toward enhanced thermal stability of virus-like particles and antibacterial activity of Polyoxometalates. *ACS Appl. Mater. Interf.* 10, 6137–6145. doi: 10.1021/acsami.7b17082

Gajera, S. B., Mehta, J. I., and Patel, M. N. (2015). DNA interaction, cytotoxicity, antibacterial and antituberculosis activity of oxovanadium(IV) complexes derived from  $\gamma$ -uroquinolones and 4-hydroxy-5-((4-hydroxyphenyl)diazenyl)thiazole-2(3H)-thione $^T$ . *RSC Adv.* 5, 21710–21719. doi: 10.1039/C5RA01222H

Garner, M., Reglinski, J., Smith, W. E., McMurray, J., Abdullah, I., and Wilson, R. (1997). A  $^1\text{H}$  spin echo and  $^{51}\text{V}$  NMR study of the interaction of vanadate with intact erythrocytes. *J. Biol. Inorg. Chem.* 2, 235–241. doi: 10.1007/s007750050129

Goldfine, A. B., Patti, M.-E., Zuberi, L., Goldstein, B. J., Leblanc, R., Landaker, E. J., et al. (2000). Metabolic effects of vanadyl sulfate in humans with non-insulin-dependent diabetes mellitus: *in vivo* and *in vitro* studies. *Metabolism* 49, 400–410. doi: 10.1016/S0026-0495(00)90418-9

Gresser, M. J., Tracey, A. S., and Parkinson, K. M. (1986). Vanadium(V) oxyanions: the interaction of vanadate with pyrophosphate, phosphate, and arsenate. *J. Am. Chem. Soc.* 108, 6229–6234. doi: 10.1021/ja00280a020

Gumerova, N., Krivosudsky, L., Fraqueza, G., Breibek, J., Al-Sayed, E., Tanuhadi, E., et al. (2018). The P-type ATPase inhibiting potential of polyoxotungstates. *Metalomics* 10, 287–295. doi: 10.1039/C7MT00279C

Hadzibrahimovic, M., Suznjevic, D., Pastor, F., Antic, T. C., Zizic, M., Zakrzewska, J., et al. (2017). The interactions of vanadate monomer with the mycelium of fungus *Phycomyces blakesleeanus*: reduction or uptake? *Anton. V. Leeuw. Inter. J. Gen. Mol. Microb.* 110, 365–373. doi: 10.1007/s10482-016-0808-0

Hider, R. C., and Kong, X. (2010). Chemistry and biology of siderophores. *Nat. Prod. Rep.* 27, 637–657. doi: 10.1039/b906679a

Hill, C. L., Hartnup, M., Faraj, M., Weeks, M., Prosser-Mccartha, C. M., Brown, R. B. Jr., et al. (1990). Polyoxometalates as inorganic anti-HIV-1 compounds. Structure-activity relationships. *Adv. Chemotherapy AIDS*, 33–41.

Jakusch, T., Enyedy, E. A., Kozma, K., Paár, Z., Bényei, A., and Kiss, T. (2014). Vanadate complexes of 3-hydroxy-1, 2-dimethyl-pyridinone: Speciation, structure and redox properties. *Inorg. Chim. Acta* 420, 92–102. doi: 10.1016/j.ica.2013.12.034

Jakusch, T., and Kiss, T. (2017). *In vitro* study of the antidiabetic behavior of vanadium compounds. *Coord. Chem. Rev.* 351, 118–126. doi: 10.1016/j.ccr.2017.04.007

Jakusch, T., Pessoa, J. C., and Kiss, T. (2011). The speciation of vanadium in human serum. *Coord. Chem. Rev.* 255, 2218–2226. doi: 10.1016/j.ccr.2011.02.022

Kaliva, M., Kyriakakis, E., and Safifoglou, A. (2002). Reactivity investigation of dinuclear vanadium(IV,V)-citrate complexes in aqueous solutions. A closer look into aqueous vanadium-citrate interconversions. *Inorg. Chem.* 41, 7015–7023. doi: 10.1021/ic020323r

Kaliva, M., Raptopoulou, C. P., Terzis, A., and Safifoglou, A. (2003). Systematic studies on pH-dependent transformations of dinuclear vanadium(V)-citrate complexes in aqueous solutions - A perspective relevance to aqueous vanadium(V)-citrate speciation. *J. Inorg. Biochem.* 93, 161–173. doi: 10.1016/S0162-0134(02)00563-9

Kioseoglou, E., Gabriel, C., Petanidis, S., Pscharis, V., Raptopoulou, C. P., Terzis, A., et al. (2013). Binary Decavanadate-Betaine composite materials of potential anticarcinogenic activity. *Z. Anorg. Allg. Chem.* 639, 1407–1416. doi: 10.1002/zaac.201300144

Kioseoglou, E., Petanidis, S., Gabriel, C., and Salifoglou, A. (2015). The chemistry and biology of vanadium compounds in cancer therapeutics. *Coord. Chem. Rev.* 301, 87–105. doi: 10.1016/j.ccr.2015.03.010

Kiss, T., Jakusch, T., Hollender, D., Dornyei, A., Enyedy, E. A., Pessoa, J. C., et al. (2008). Biospeciation of antidiabetic VO(IV) complexes. *Coord. Chem. Rev.* 252, 1153–1162. doi: 10.1016/j.ccr.2007.09.011

Kustin, K. (2015). Aqueous vanadium ion dynamics relevant to bioinorganic chemistry: a review. *J. Inorg. Biochem.* 147, 32–38. doi: 10.1016/j.jinorgbio.2014.12.009

Le, M., Rathje, O., Levina, A., and Lay, P. A. (2017). High cytotoxicity of vanadium (IV) complexes with 1, 10-phenanthroline and related ligands is due to decomposition in cell culture medium. *J. Biol. Inorg. Chem.* 22, 663–672. doi: 10.1007/s00775-017-1453-4

Leon, I. E., Cadavid-Vargas, J. F., Di Virgilio, A. L., and Etcheverry, S. B. (2017). Vanadium, Ruthenium and Copper compounds: a new class of Nonplatinum Metallodrugs with Anticancer activity. *Curr. Med. Chem.* 24, 112–148. doi: 10.2174/092986732366616082416254

Leon, I. E., Porro, V., Astrada, S., Egusquiza, M. G., Cabello, C. I., Bollati-Fogolin, M., et al. (2014). Polyoxometalates as antitumor agents: bioactivity of a new polyoxometalate with copper on a human osteosarcoma model. *Chem. Biol. Interact.* 222, 87–96. doi: 10.1016/j.cbi.2014.10.012

Levina, A., Crans, D. C., and Lay, P. (2017a). Speciation of metal drugs, supplements and toxins in media and bodily fluids controls *in vitro* activities. *Coord. Chem. Rev.* 352, 473–498. doi: 10.1016/j.ccr.2017.01.002

Levina, A., and Lay, P. A. (2017b). Stabilities and Biological Activities of Vanadium Drugs: What is the Nature of the Active Species?. *Chem. Asian J.* 12, 1692–1699. doi: 10.1002/asia.201700463

Li, M., Ding, W., Smee, J. J., Baruah, B., Willsky, G. R., and Crans, D. C. (2009). Anti-diabetic effects of vanadium (III, IV, V) - chlorodipicolinate complexes in streptozotocin-induced diabetic rats, *BioMetals* 103, 585–905. doi: 10.1007/s10534-009-9241-4

Lodyga-Chruscinska, E., Sanna, D., Garribba, E., and Micera, G. (2008). Potentiometric, spectroscopic, electrochemical and DFT characterization of oxovanadium(IV) complexes formed by citrate and tartrates in aqueous solution at high ligand to metal molar ratios: the effects of the trigonal bipyramidal distortion in bis-chelated species and biological implications. *Dalton Trans.* 28, 4903–4916. doi: 10.1039/b803520b

Lu, L. P., and Zhu, M. L. (2011). Metal-based inhibitors of protein Tyrosine Phosphatases. *Anti Cancer Agents Med. Chem.* 11, 164–171. doi: 10.2174/187152011794941271

Lyonnet, B., Martz, S., and Martin, E. (1899). L'emploi thérapeutique des dérivés du vanadium. *La Presse Méd.* 1, 191–192.

Maiti, A., and Ghosh, S. (1989). Synthesis and reactivity of the Oxovanadium(IV) Complexes of Two N-O donors and potentiation of the antituberculosis activity of one of them on Chelation to metal ions: part I. *V. J. Inorg. Biochem.* 36, 131–139. doi: 10.1016/0162-0134(89)80020-0

Martins, P. G., Mori, M., Chiaradia-Delatorre, L. D., Menegatti, A. C., Mascarello, A., Botta, B., et al. (2015). Exploring oxido-vanadium (IV) complexes as YopH inhibitors: mechanism of action and modeling studies. *ACS Med. Chem. Lett.* 6, 1035–1040. doi: 10.1021/acsmmedchemlett.5b00267

Mclaughlan, C. C., Peters, B. J., Willsky, G. R., and Crans, D. C. (2015). Vanadium-phosphatase complexes: Phosphatase inhibitors favor the trigonal bipyramidal transition state geometries. *Coord. Chem. Rev.* 301–302, 163–199. doi: 10.1016/j.ccr.2014.12.012

Medina, J. J. M., Naso, L. G., Pérez, A. L., Rizzi, A., Ferrer, E. G., and Williams, P. A. (2017). Antioxidant and anticancer effects and bioavailability studies of the flavonoid baicalin and its oxido-vanadium (IV) complex. *J. Inorg. Biochem.* 166, 150–161. doi: 10.1016/j.jinorgbio.2016.11.005

Moskovitz, B. L., and Group, A. T. H.-C. S. (1988). Clinical trial and tolerance of HPA-23 in patients with acquired immunodeficiency syndrome. *Antimicrob. Agents Chemother.* 32, 1300–1303. doi: 10.1128/AAC.32.9.1300

Noeske, J., Wasserman, M. R., Terry, D. S., Altman, R. B., Blanchard, S. C., and Cate, J. H. D. (2015). High-resolution structure of the *Escherichia coli* ribosome. *Nat. Struct. Mol. Biol.* 22, 336–341. doi: 10.1038/nsmb.2994

Pessoa, J. C., Etcheverry, S., and Gambino, D. (2015). Vanadium compounds in medicine. *Coord. Chem. Rev.* 301, 24–48. doi: 10.1016/j.ccr.2014.12.002

Pessoa, J. C., and Tomaz, I. (2010). Transport of therapeutic Vanadium and Ruthenium complexes by blood plasma components. *Curr. Med. Chem.* 17, 3701–3738. doi: 10.2174/092986710793213742

Pettersson, L., Andersson, I., and Hedman, B. (1985). Multicomponent polyanions. 37. A potentiometric and 51V-NMR study of equilibria in the  $H^+-HVO_4^{2-}$  system in 3.0 M-Na(ClO<sub>4</sub>) medium covering the range 1E-1g[H+]-10. *Chem. Scr.* 25, 309–317.

Pettersson, L., Hedman, B., Andersson, I., and Ingrı, N. (1983). Multicomponent polyanions. 34. P potentiometric and 51V NMR study of equilibria in the  $H^+-HVO_4^{2-}$  system in the 0.6 M Na(Cl) medium covering the range 1E-1g[H+]-10. *Chem. Scrip.* 22, 254–264.

Pluskey, S., Mahroof-Tahir, M., Crans, D. C., and Lawrence, D. S. (1996). Vanadium oxoanions and cAMP-dependent protein kinase: an anti-substrate inhibitor. *Biochem. J.* 321, 333–339. doi: 10.1042/bj3210333

Pope, M. T., and Müller, A. (1991). Polyoxometalate chemistry: an old field with new dimensions in several disciplines. *Angew. Chem. Int. Ed. Engl.* 30, 34–48. doi: 10.1002/anie.199100341

Postal, K., Maluf, D. F., Valdameri, G., Rüdiger, A. L., Hughes, D. L., de Sá, E. L., et al. (2016). Chemoprotective activity of mixed valence polyoxovanadates against diethylsulphate in *E. coli* cultures: insights from solution speciation studies. *RSC Adv.* 6, 114955–114866. doi: 10.1039/c6ra15826a

Rehder, D. (1991). The bioinorganic chemistry of vanadium. *Angew. Chem. Int. Ed. Engl.* 30, 148–167. doi: 10.1002/anie.199101481

Rehder, D., Pessoa, J. C., Geraldes, C., Castro, M., Kabanos, T., Kiss, T., et al. (2002). *In vitro* study of the insulin-mimetic behaviour of vanadium(IV, V) coordination compounds. *J. Biol. Inorg. Chem.* 7, 384–396. doi: 10.1007/s00775-001-0311-5

Rhule, J. T., Hill, C. L., Judd, D. A., and Schinazi, R. F. (1998). Polyoxometalates in medicine. *Chem. Rev.* 98, 327–358. doi: 10.1021/cr960396q

Sakurai, H. (2002). A new concept: the use of vanadium complexes in the treatment of diabetes mellitus. *Chem. Rec.* 2, 237–248. doi: 10.1002/tcr.10029

Sakurai, H., Tsuchiya, K., Nukatsuka, M., Kawada, J., Ishikawa, S., Yoshida, H., et al. (1990). Insulin-mimetic action of vanadyl complexes. *J. Clin. Biochem. Nutr.* 8, 193–200. doi: 10.3164/jcbn.8.193

Samart, N., Saeger, J., Haller, K., Manuel Aureliano, M., and Crans, D. (2014). Interaction of decavanadate with interfaces and biological model membrane systems: characterization of soft oxometalate systems. *J. Mol. Eng. Mat.* 2, 1–21. doi: 10.1142/S2251237314400073

Sambandamurthy, V. K., Derrick, S. C., Hsu, T., Chen, B., Larsen, M. H., Jalapathy, K. V., et al. (2006). *Mycobacterium tuberculosis* ΔRD1 ΔpanCD: a safe and limited replicating mutant strain that protects immunocompetent and immunocompromised mice against experimental tuberculosis. *Vaccine* 24, 6309–6320. doi: 10.1016/j.vaccine.2006.05.097

Sanchez-Lombardo, I., Baruah, B., Alvarez, S., Werst, K. R., Segaline, N. A., Levinger, N. E., et al. (2016). Size and shape trump charge in interactions of oxovanadates with self-assembled interfaces: application of continuous shape measure analysis to the decavanadate anion. *New J. Chem.* 40, 962–975. doi: 10.1039/C5NJ01788B

Sanna, D., Serra, M., Micera, G., and Garribba, E. (2014). Interaction of antidiabetic vanadium compounds with hemoglobin and red blood cells and their distribution between plasma and erythrocytes. *Inorg. Chem.* 53, 1449–1464. doi: 10.1021/ic402366x

Sanna, D., Ugone, V., Micera, G., Buglyó, P., Bíró, L., and Garribba, E. (2017). Speciation in human blood of Metvan, a vanadium based potential anti-tumor drug. *Dalton Trans.* 46, 8950–8967. doi: 10.1039/C7DT00943G

Selling, A., Andersson, I., Pettersson, L., and Schramm, C. M., S.L., D., and J.H., G. (1994). Multicomponent Polyanions. 47. The aqueous Vanadophosphate system. *J. Inorg. Chem.* 33, 3141–3150. doi: 10.1021/ic00092a021

Selman, M., Russo, C., Bergeron, A., Son, H., Krishnan, R., El-Sayes, N., et al. (2018). Multi-modal potentiation of oncolytic virotherapy by Vanadium compounds. *Mol. Ther.* 26, 56–69. doi: 10.1016/j.ymthe.2017.10.014

Shah, H. S., Joshi, S. A., Haider, A., Kortz, U., Ur-Rehman, N., and Iqbal, J. (2015). Synthesis of chitosan-coated polyoxometalate nanoparticles against cancer and its metastasis. *RSC Adv.* 5, 93234–93242. doi: 10.1039/C5RA18489D

Smith, D. M., Pickering, R. M., and Lewith, G. T. (2008). A systematic review of vanadium oral supplements for glycaemic control in type 2 diabetes mellitus. *Qjm—an Inter. J. Med.* 101, 351–358. doi: 10.1093/qjmed/hcn003

Sun, T. D., Cui, W., Yan, M., Qin, G., Guo, W., Gu, H. X., et al. (2016). Target delivery of a novel Antitumor Organoplatinum(IV)-substituted Polyoxometalate Complex for Safer and more effective colorectal cancer therapy *in vivo*. *Adv. Mater.* 28, 7397–7404. doi: 10.1002/adma.201601778

Thompson, K. H., Licher, J., Lebel, C., Scaife, M. C., McNeill, J. H., and Orvig, C. (2009). Vanadium treatment of type 2 diabetes: a view to the future. *J. Inorg. Biochem.* 103, 554–558. doi: 10.1016/j.jinorgbio.2008.12.003

Thompson, K. H., and Orvig, C. (2006). Vanadium in diabetes: 100 years from Phase 0 to Phase, I. *J. Inorg. Biochem.* 100, 1925–1935. doi: 10.1016/j.jinorgbio.2006.08.016

Trevino, S., Sánchez-Lara, E., Sarmiento-Ortega, V. E., Sánchez-Lombardo, I., Flores-Hernández, J. A., Pérez-Benítez, A., et al. (2015). Hypoglycemic, lipid-lowering and metabolic regulation activities of metforminium decavanadate (H<sub>2</sub>Metf) [V10O<sub>28</sub>]·8H<sub>2</sub>O using hypercaloric-induced carbohydrate and lipid deregulation in Wistar rats as biological model. *J. Inor. Biochem.* 147, 85–92. doi: 10.1016/j.jinorgbio.2015.04.002

Turian, G. (1951). Action plasmogène du fer chez les Mycobactéries. Le bacille de la fèole, indicateur biologique du fer. 34,917–920. *Helv. Chim. Acta* 34, 917–920. doi: 10.1002/hlca.19510340325

Turner, T. L., Nguyen, V. H., McLauchlan, C. C., Dymon, Z., Dorsey, B. M., Hooker, J. D., et al. (2012). Inhibitory effects of decavanadate on several enzymes and Leishmania tarentolae *in vitro*. *J. Inorg. Biochem.* 108, 96–104. doi: 10.1016/j.jinorgbio.2011.09.009

Upadhyay, A., Fontes, F., Gonzalez-Juarrero, M., McNeil, M. R., Crans, D. C., Jackson, M., et al. (2015). MenI, a novel menaquinone reductase in *Mycobacterium tuberculosis* is required for survival in macrophages. *ACS Central Sci.* 2015, 1, 292–302. doi: 10.1021/acscentsci.5b00212

Vilas Boas, L. V., and Costa Pessoa, J. (1987). "Vanadium," in *Comprehensive Coordination Chemistry. The Synthesis, Reactions, Properties & Applications of Coordination Compounds*, eds G. Wilkinson, Sir, R. D. Gillard, and J. A. McCleverty (New York, NY: Pergamon Press), 453–583.

Wang, L., Zhou, B. B., and Liu, J. R. (2013). Anticancer Polyoxometalates. *Prog. Chem.* 25, 1131–1141.

Weinstein, S., Jahn, W., Glotz, C., Schlinzen, F., Levin, I., Janell, D., et al. and Yonath, A. (1999). Metal compounds as tools for the construction and the interpretation of medium-resolution maps of ribosomal particles. *J. Struct. Biol.* 127, 141–151. doi: 10.1006/jsb1.1999.4135

Willsky, G. R., Chi, L.-H., Godzala, M. III, Kostyniak, P. J., Smee, J. J., Trujillo, A. M., et al. (2011). Anti-diabetic effects of a series of vanadium dipicolinate complexes in rats with streptozotocin-induced diabetes. *Coord. Chem. Rev.* 255, 2258–2269. doi: 10.1016/j.ccr.2011.06.015

Willsky, G. R., Goldfine, A. B., Kostyniak, P. J., McNeill, J. H., Yang, L. Q., Khan, H. R., et al. (2001). Effect of vanadium(IV) compounds in the treatment of diabetes: *in vivo* and *in vitro* studies with vanadyl sulfate and bis(maltolato)oxovanadium(IV). *J. Inorg. Biochem.* 85, 33–42. doi: 10.1016/S0162-0134(00)00226-9

Willsky, G. R., Halvorsen, K., Godzala, M. E., Iii, C. hi, L.-H., Most, M. J., Kaszynski, P., et al. (2013). Coordination chemistry may explain pharmacokinetics and clinical response of vanadyl sulfate in type 2 diabetic patients. *Metallomics* 5, 1491–1502. doi: 10.1039/c3mt00162h

Willsky, G. R., Leung, J. O., Offermann, P. V. Jr., Plotnick, E. K., and Dosch, S. F. (1985). Isolation and characterization of vanadate-resistant mutants of *Saccharomyces cerevisiae*. *J. Bacteriol.* 164, 611–617.

Willsky, G. R., Preischel, D. A., and McCabe, B. C. (1984a). Vanadium metabolism in *S-Cerevisiae*. *Biophys. J.* 45, A76–A76.

Willsky, G. R., White, D. A., and McCabe, B. C. (1984b). Metabolism of added Ortho-Vanadate to Vanadyl and High- molecular-weight Vanadates by *Saccharomyces-Cerevisiae*. *J. Biol. Chem.* 259, 3273–3281.

Winkler, P. A., Huang, Y., Sun, W., Du, J., and and, Lü, W. (2017). Electron cryo-microscopy structure of a human TRPM4 channel. *Nature* 552, 200–204. doi: 10.1038/nature24674

Zhai, F. Y., Wang, X. H., Li, D. L., Zhang, H., Li, R., and Song, L. S. (2009). Synthesis and biological evaluation of decavanadate Na(4)Co(H<sub>2</sub>O)(6)V(10)O(28)center dot 18H<sub>2</sub>O. *Biomed. Pharmacother.* 63, 51–55. doi: 10.1016/j.biopha.2008.01.006

Zhang, Y., Yang, X. D., Wang, K., and Crans, D. C. (2006). The permeability and cytotoxicity of insulin-mimetic vanadium (III,IV,V)-dipicolinate complexes, *J. Inorg. Biochem.* 100, 80–87. doi: 10.1016/j.jinorgbio.2005.10.006

Zhou, Z. H., Zhang, H., Jiang, Y. Q., Lin, D. H., Wan, H. L., and Tsai, K. R. (1999). Complexation between vanadium(V) and citrate: spectroscopic and structural characterization of a dinuclear vanadium(V) complex. *Transit. Metal Chem.* 24, 605–609. doi: 10.1023/A:1006947218366

Zizic, M., Miladinovic, Z., Stanic, M., Hadzibrahimovic, M., Zivic, M., and Zakrzewska, J. (2016). V-51 NMR investigation of cell-associated vanadate species in *Phycomyces blakesleeanus* mycelium. *Res. Microbiol.* 167, 521–528. doi: 10.1016/j.resmic.2016.04.012

Zizic, M., Zivic, M., Spasojevic, I., Bogdanovic Pristov, J., Stanic, M., Cvetic-Antic, T., et al. (2013). The interactions of vanadium with *Phycomyces blakesleeanus* mycelium: enzymatic reduction, transport and metabolic effects. *Res. Microbiol.* 164, 61–69. doi: 10.1016/j.resmic.2012.08.007

Zwolak, I. (2014). Vanadium carcinogenic, immunotoxic and neurotoxic effects: a review of *in vitro* studies. *Toxicol. Mech. Method.* 24, 1–12. doi: 10.3109/15376516.2013.843110

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Samart, Arhouma, Kumar, Murakami, Crick and Crans. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.