

# Palladium Catalyzed Chemoselective Activation of $sp^3$ vs $sp^2$ C–H Bonds: Oxidative Coupling to Form Quaternary Centers

Gang Hong,<sup>†,‡</sup> Pradip D. Nahide,<sup>†</sup> Uday Kumar Neelam,<sup>†</sup> Peter Amadeo,<sup>†</sup> Arjun Vijeta,<sup>†</sup> John M. Curto,<sup>†</sup> Charles E. Hendrick,<sup>†</sup> Kelsey F. VanGelder,<sup>†</sup> and Marisa C. Kozlowski<sup>\*,†</sup>

<sup>†</sup>Department of Chemistry, Roy and Diana Vagelos Laboratories, University of Pennsylvania, Philadelphia, Pennsylvania 19104, United States

<sup>†</sup>Key Laboratory for Advanced Materials, Institute of Fine Chemicals and School of Chemistry & Molecular Engineering, East China University of Science and Technology, 130 Meilong Road, Shanghai, 200237, P. R. China

## Supporting Information Placeholder

**ABSTRACT:** The oxidative activation of alkyl C–H bonds vs arene C–H bonds with  $\text{Pd}(\text{OAc})_2$  has been found to be generalizable to a number of nucleophilic substrates allowing the formation of a range of hindered quaternary centers. The substrates share a common mechanistic path wherein  $\text{Pd}(\text{II})$  initiates an oxidative dimerization. The resultant dimer modifies the palladium catalyst to favor activation of alkyl C–H bonds in contrast to the trends typically observed via a concerted metalation deprotonation mechanism. Notably, insertion occurs at the terminus of the alkyl arene for hindered substrates. Two different oxidant systems were discovered that turn over the process. Parameters have been identified that predict, which substrates are productive in this reaction.

**KEY WORDS:** cross dehydrogenative coupling; oxidative coupling; palladium; chemoselective activation; quaternary


---

## Introduction

The use of aromatic hydrocarbons as precursors to the construction of more complex structures has many intrinsic advantages. Aromatic hydrocarbons are inexpensive building blocks readily available from petroleum. Their direct functionalization to desired targets avoids the introduction of functional groups such as halides that require the use of hazardous materials or generate undesired byproducts. The aromatic hydrocarbons are also more stable than the corresponding functionalized equivalents such as benzylic halides. For example, naphthyl halide analogs are not stable and are not readily amenable to  $S_N2$  displacement.

For these reasons, much research has focused on the effective use of aromatic hydrocarbons as substrates and metal catalyzed C–H activation chemistry has played a key role.<sup>1</sup> The use of palladium, in particular, has proven highly effective especially in the activation of  $sp^2$  C–H bonds of arenes (e.g. Scheme 1, eq 1),<sup>2</sup> even allowing C–H activation of two reaction partners where the only byproduct is dihydrogen.<sup>3</sup> On the other hand, the activation of the  $sp^3$  C–H bonds of aromatic hydrocarbons by palladium occurs much less readily to the point where toluene derivatives can undergo selective  $sp^2$  C–H functionalization in the presence of one or more benzylic centers.<sup>4</sup> Strategies to achieve benzylic activation have relied on directing groups (Scheme 1, eq 2),<sup>5</sup> generation of benzylic radicals (Scheme 1, eq 3),<sup>6</sup> or the lower acidity of benzylic sites (Scheme 1, eq 4).<sup>7</sup>

Recently, we have reported an alternative strategy to activation of aromatic hydrocarbons wherein dehydrogenative coupling forms a hindered bond (Scheme 1, eq 5).<sup>8</sup> One unique feature of this process is the catalytic activation of benzylic C–H bonds under oxidizing conditions with palladium that typically result in  $sp^2$  C–H activation. Another unusual feature is that coupling occurs at the terminus of the alkyl arene, presumably via a migration of the palladium center. In this article, we disclose our discovery of a range of acyclic and cyclic substrates that can participate in this process to generate hindered bonds selectively (eq 6). From the studies undertaken, guidelines are now available to predict which carbon nucleophiles




will be effective.

## Results and Discussion


Our analysis of the reaction from eq 5 revealed the intermediacy of a dimer from the azlactone, which readily forms under a range of oxidative conditions.<sup>9</sup> Notably, others have found reactions that also proceed through dimers similar to those that we reported.<sup>10</sup> Reasoning that other substrates which form similar dimeric intermediates might be subject to a similar reaction

**Scheme 1. Pd-Catalyzed Activation of Arenes and Toluenes**


Typical Pd CDC via arene insertion




Directed Pd CDC benzylic insertion




Radical Pd CDC benzylic insertion




Deprotonative toluene arylation




Previous work



This work




profile (eq 7), the literature was surveyed for compounds that readily undergo oxidative dimerization. A surprisingly large number of compounds with similar behavior was identified including malonates,<sup>11</sup> malononitrile,<sup>12</sup> cyanoacetate,<sup>13</sup> oxindoles,<sup>14</sup> isooxindoles,<sup>15</sup> benzofurans,<sup>16</sup> fluorenes,<sup>17</sup> and hydroxycoumarins.<sup>18</sup>



**Acyclic Substrates: Malononitriles.** A primary concern was whether the reaction was restricted to cyclic nucleophiles. Thus, the initial priority was to explore acyclic substrates capable of oxidative dimerization. Malononitriles are also well-known to undergo oxidative dimerization.<sup>12</sup> To probe the fundamental reactivity, trials were first conducted with stoichiometric Pd(OAc)<sub>2</sub> using dinitrile **1a** along with toluene (Scheme 2).


**Scheme 2. Variable Yields in Dinitrile Toluene Coupling**



Good reactivity was observed, but the outcome was strongly affected by the batch of palladium used. Older batches of palladium were highly effective in the stoichiometric conversion of dinitrile **1a** to **3aa** with toluene (Scheme 2), while new batches were not. Different sources of Pd(OAc)<sub>2</sub> are known to give much different results due to different composition arising from the preparation,

such as the red-brown [Pd<sub>3</sub>(OAc)<sub>5</sub>NO<sub>2</sub>]. [Pd<sub>3</sub>(OAc)<sub>6</sub>] is a dark purple solid whereas [Pd(OAc)<sub>2</sub>]<sub>n</sub> polymer is a light grey-purple powder.<sup>19</sup>


**Scheme 3. Catalytic Palladium Oxidative Coupling of Malononitriles with Methylarenes (eq 8)**



Proceeding with older batches of Pd(OAc)<sub>2</sub>, the *para*-methoxyphenyl and *para*-methylphenyl malononitriles were found to be effective partners in the coupling of different toluenes (Scheme 3). Further, the same stoichiometric oxidant discovered for the azlactones, dimethylbenzoquinone (DMBQ), was found to be effective in turning over the process with catalytic amounts of Pd(OAc)<sub>2</sub>. In most cases, similar yields were obtained for the catalytic vs stoichiometric palladium conditions. Amongst the xylene isomers, *para*-xylene (27-36%) was less effective than *meta*-xylene (59-65%) or *ortho*-xylene (32-50%). In all cases, reactions were conducted until all starting material was consumed and the mass balance is decomposition with no specific byproducts being isolable. Notably, higher yields were obtained under catalytic conditions for *para*-methoxytoluene. We speculate that stoichiometric Pd(OAc)<sub>2</sub> can cause direct oxidation of the *para*-methoxytoluene, which was mitigated with lower amounts of Pd(OAc)<sub>2</sub> in the presence of the mild oxidant DMBQ.


For ethylbenzene, a divergence in behavior was seen relative to our prior report with azlactone (eq 5)<sup>8</sup> where exclusive functionalization of the terminal CH<sub>3</sub> position occurred. With the malononitrile, a mixture of the terminal (methyl) vs internal (benzylic) products was observed (eq 9). The terminal product is proposed to arise from initial benzylic insertion followed by  $\beta$ -hydride elimination and migratory insertion.<sup>20</sup> We hypothesize that the driving force for rearrangement to the terminal position is largely controlled by formation of the less hindered alkyl palladium species. However, these results indicate that the nucleophile also plays a role, where sterically small nucleophiles can react at either the  $\alpha$ - or  $\beta$ -positions. Under stoichiometric conditions, there is a preference for the latter **3ag'**, whereas the former **3ag** predominates under catalytic conditions. Under catalytic conditions, the higher effective

concentration of the nucleophile relative to the palladium species may cause greater capture of the benzylic intermediate prior to rearrangement.



**Acyclic Substrates:  $\alpha$ -Cyanoacetates.**  $\alpha$ -Cyanoacetates are also well known to undergo such dimerizations, although somewhat more slowly.<sup>11</sup> To probe the fundamental reactivity, trials were first conducted with stoichiometric  $\text{Pd}(\text{OAc})_2$  using cyanoacetate **4a** along with 20 equivalents of alkylarene in dioxane (eq 10, Scheme 4). At 95 °C, the reaction with toluene proceeded well (81% yield). Again, *para*-xylene (38%) was less effective than *meta*-xylene (69%) or *ortho*-xylene (61%). Substrates with electron-donating groups (*para*-methoxytoluene) proceeded with moderate conversion (42%), but those with electron-withdrawing groups, such as *para*-trifluorotoluene or *para*-cyanotoluene gave no product. 2-Methylnaphthalene proceeded well (67%), whereas the more hindered 1-methylnaphthalene did not work (<10%). In contrast to the malononitriles, ethylbenzene gave the product arising exclusively from functionalization of the terminal position of the alkyl (**5ag**).


**Scheme 4. Stoichiometric Palladium Oxidative Coupling of Cyanoacetates with Alkylarenes (eq 10)**



Initial trials using the same DMBQ oxidant as described above for the malononitriles led to poor yields. On the whole, the  $\alpha$ -cyanoacetates were less reactive than malononitriles. As such, high throughput screening was undertaken to identify an oxidant that would permit turnover of the process. With varying amounts of

$\text{Pd}(\text{OAc})_2$ , several oxidants were examined (Scheme 5) and  $\text{K}_2\text{S}_2\text{O}_8$  was found to be superior. Having observed that carboxylic additives improved the yields with the azlactone substrate,<sup>8</sup> presumably by ligating the palladium to prevent precipitation during the reduction, both acetic acid and pivalic acid additives were examined. Further, several solvents (DMF, NMP, 1,4-dioxane, 2-butanol, diglyme, *i*-PrOAc) were screened to minimize the amount of alkylarene employed (see SI). Overall acetic acid in dioxane performed best. Further examination of temperature (95, 110, 120 °C) revealed that higher temperatures (120 °C) were well tolerated causing little decomposition and higher yields (see SI).



**Scheme 5. Optimization of Oxidant for Cyanoacetates with Alkylarenes.<sup>a,b</sup>**



<sup>a</sup>Substrate **4a**,  $\text{Pd}(\text{OAc})_2$ , 0.1 M toluene, 95 °C, 15 h. <sup>b</sup>Sphere size correlates with conversion vs an internal standard (see SI for quantitative values).

Applying the optimized conditions (eq 11) to a range of toluene substrates (top of Scheme 6) provided results as good as for the stoichiometric transformation with the exception of *para*-methoxytoluene, which likely undergoes an oxidation side reaction with the  $\text{K}_2\text{S}_2\text{O}_8$ . Yet again, *para*-xylene (48%) was less effective than *meta*-xylene (68%) or *ortho*-xylene (62%). Further, the catalytic conditions performed well for several different cyanoacetates (bottom of Scheme 6) including those bearing electron-withdrawing groups (**5ba**) or electron-donating groups (**5da**).

**Scheme 6. Catalytic Palladium Oxidative Coupling of Cyanoacetates with Alkylarenes (eq 11)**



entry	X	Y	additive (equiv)	yield (%) <sup>b</sup>
1	0.3	2		98 (97) <sup>c</sup>
2	0.3	1		76
3	0.2	2		64
4	0.2	2	PivOH (1)	72
5	0.3	2	AcOH (1)	80
6	0.2	2	PivOH (1)	96 <sup>d</sup>
7	0.3	2	AcOH (1)	77 <sup>d</sup>
8	0.2	2	$\text{K}_2\text{CO}_3$ (1)	51
9	0.2	2	NaOAc (1)	44


<sup>a</sup>Reaction conditions: **6a** (0.15 mmol), **2a** (0.1 M),  $\text{Pd}(\text{OAc})_2$  (X mol %),  $\text{K}_2\text{S}_2\text{O}_8$  (Y equiv) at 120 °C for 24 h under argon. <sup>b</sup>Isolated yield. <sup>c</sup>Run twice <sup>d</sup>Activated charcoal (10x weight of Pd) was added.

Additional 3-substituted oxindoles were synthesized following literature protocols<sup>23</sup> and were examined in the coupling with toluene (eq 13, Scheme 8). Different 3-aryl groups with both

**Scheme 8. Palladium Oxidative Coupling of Oxindoles with Toluene (eq 13)**


**Cyclic Substrates: Oxindoles.** Moving away from acyclic substrates, oxindoles are another class of compounds that readily dimerize.<sup>14</sup> In addition, oxindoles bearing two substituents at the 3-position are found in natural products and pharmaceutical targets (Scheme 7).<sup>21</sup> Notably, 3-aryl-3-benzyl motif that maps onto this method has been utilized by Hoffmann-LaRoche.<sup>21c</sup> Functionalization of oxindoles has been the focus of much work, including oxidative methods.<sup>22</sup>

**Scheme 7. Bioactive Oxindole Structures**



Very good initial results were obtained in the coupling of 3-phenyloxindole (**6a**) with toluene (eq 13, Scheme 8) with stoichiometric  $\text{Pd}(\text{OAc})_2$  (see SI). A survey of oxidants revealed  $\text{K}_2\text{S}_2\text{O}_8$  as highly effective while minimizing byproduct formation (see SI). In this case, using dioxane as a solvent was not as effective (see SI), so reactions were conducted in 0.1 M alkylarene. Further optimization (Table 1) revealed that the palladium catalyst loading could be lower when a pivalic acid additive and activated charcoal were employed (entry 6).

**Table 1. Optimization of 3-Phenylloxindole Toluene Coupling (eq 12)**




electron-donating and electron-withdrawing substituents were well tolerated leading to high yields under both stoichiometric and catalytic conditions (**7aa-7ma**). In comparison, generation of the oxindole cation initiates a Friedel-Crafts reaction with arenes; such a pathway cannot incorporate toluene at the benzylic carbon. Notably, this approach to **7ba**, **7fa** and similar structures is orthogonal to the oxidative coupling method reported by Li and workers<sup>24</sup> wherein selective coupling of the oxindole occurs via a Friedel-Crafts reaction resulting in functionalization of the arene C–H of toluene rather than the benzylic C–H. The 2-naphthyl substrate

formed product (**7ma**) in high yield whereas the hindered 1-naphthyl substrate reacted poorly.

Heterocyclic groups could also be employed at the 3-position including a case with a nitrogen substituent (**7na-7pa**). The oxindole could also be employed directly without protection of the nitrogen (**7qa**) or with other protecting groups (**7ra**, **7sa**); however, the Boc group was somewhat unstable at the high reaction temperatures.


Similar trends were observed again with different arylalkanes with improved outcomes for ethylbenzene (eq 14, Scheme 9). Varying toluene, poor reactivity was again observed for substrates with electron-withdrawing groups such as *para*-methylbenzonitrile and *para*-chlorotoluene. The reaction was also unsuccessful for 2-methylfuran. To probe the  $\beta$ -hydride elimination and rearrangement, isobutylbenzene was examined. As expected, the addition of steric hindrance abrogated reactivity.

**Scheme 9. Palladium Oxidative Coupling of Parent Oxindole with Alkylarenes (eq 14)**



**Cyclic Substrates: Benzofuranones.** Benzofuranones are isoelectronic with oxindoles and also exhibit oxidative dimerization behavior.<sup>16</sup> They are found in a number of natural products and pharmaceutical targets (Scheme 10).<sup>25</sup>

**Scheme 10. Bioactive Benzofuranone Structures**




Using the same conditions developed for the oxindoles, toluene coupled very well with the parent 3-phenylbenzofuranone to generate **9aa** under stoichiometric or catalytic conditions in 94% and 91% yield, respectively. Substrates varying the core ring system

substituents (top Scheme 11) or the 3-aryl group (middle Scheme 11) were generated<sup>26</sup> and were found to couple with toluene very well. Electron-donating (9fa, 9ja-9la) and electron-withdrawing groups (9ga, 9ha) could be employed. Notably, chloro groups could be employed providing an opportunity for further derivatization (9ha). Lower catalyst loadings could also be employed, but required longer reaction or resulted in lower yields (for 9ba with 5 mol % Pd(OAc)<sub>2</sub>, 59% and 75% product were obtained at reaction times of 24 h and 48 h, respectively).

A similar scope was seen with different alkylarenes as had been observed in the cases above (bottom, Scheme 11).

**Scheme 11. Palladium Oxidative Coupling of Benzofuranones with Alkylarenes (eq 15)<sup>a</sup>**




<sup>a</sup>For 9ba, using conditions B with 5 mol % Pd(OAc)<sub>2</sub>, 59% and 75% product were obtained at reaction times of 24 h and 48 h, respectively.

**Substrate Properties Correlated with Reactivity.** In our prior report, the dimer of the azlactone (eq 5) was found to form under reaction conditions. When isolated and resubjected to the reaction conditions, this dimer also converted to product. The corresponding dimers were observed in the conversion of substrates 1, 4, 6, and 8

although very little of the dimer built up during the conversion of oxindoles 6 or benzofuranones 8. The dimer (10) of  $\alpha$ -cyanoacetate 4a was readily formed with oxidants and was independently synthesized (Scheme 12) with Cu(TMEDA)Cl(OH). Resubjection of dimer 10 to the reaction conditions with Pd(OAc)<sub>2</sub> and toluene again afforded the product 5aa (Scheme 12).

**Scheme 12. Dimer Intermediate from Cyanoacetate Substrate**




Heating the dimer with toluene alone did not form the product. Similarly, subjection of substrate under the catalytic conditions without Pd(OAc)<sub>2</sub> did not form the product. For example, when oxindole 6a was treated with 2 equiv K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>, 1 equiv PivOH, charcoal, in 0.1 M toluene for 24 h at 120 °C, the outcome was 73% unreacted 6a, 25% dimer, and 0% product. Thus, it appears that Pd(OAc)<sub>2</sub> plays a role in the C–H activation event of the toluene as well as in dimerization of the nucleophilic substrate.

Calculations of the bond dissociation energy (BDE) values for a range of potential nucleophilic substrates was undertaken (Table 2). If the ability to undergo dimerization via nonionic processes is the most important factor in determining reactivity, then the substrate C–H bond dissociation energy should correlate with reactivity in the overall process. A survey of all the substrates in Table 2, revealed that only entries 3–11, possessing C–H bond dissociation energies in the range of 61.0–71.6 kcal/mol, were effective in the transformation. Substrates with similarly weak C–H bonds (entries 1–2, and to a lesser extent, entries 12, 14), were found to be unreactive. All of these compounds have weaker bond dissociation energies than toluene (90 kcal/mol),<sup>27</sup> which accounts for why dimer alone (see above) is unable to activate toluene.

To further probe the importance of the initial oxidation to the dimer, the dimer of 6u was independently synthesized<sup>28</sup> and resubjected to the reaction conditions in a similar manner as outlined in Scheme 8. Similar to monomer 6u, the dimer of 6u did not convert to the product in the presence of Pd(OAc)<sub>2</sub> (Scheme 13) indicating that dimerization is necessary, but not sufficient in order for the process to occur.

**Scheme 13. 3-Benzylloxindole Dimer with Pd(OAc)<sub>2</sub>**




The dimer BDE values were also computed and correlated more strongly with reactivity (Table 2). Substrates with dimers BDE values either below 2 kcal/mol or above 15 kcal/mol did not form product. That compounds with stronger dimer BDE values did not generate product, even when the dimer was preformed (see Scheme

13) indicates that the oxidizing power of the dimer is not the most important factor. Furthermore, the relative reactivity of the substrates in entries 3-11 followed the order of the dimer BDE values as judged by TLC monitoring.

Competition experiments between substrates were performed to probe the role of BDE values (Scheme 14). Reaction from the monomer and dimer gave different outcomes, consistent with two separate steps contributing to the overall rate: dimerization by palladium and reaction of the dimer with a palladium alkyl. For all pairs, the substrates with the stronger BDE values were less reactive.

**Scheme 14. Competition Experiments**



**Table 2. Bond Dissociation Energies of A-H and A-A**

Entry	A-H	monomer BDE <sup>a</sup> (kcal/mol)	dimer BDE <sup>a,b</sup> (kcal/mol)
1		71.9	0.8
2		71.0	1.4
3		60.8	2.4
4		65.1	5.3
5		67.9	7.4

6		68.1	8.7
7		69.0	10.0
8		69.5	10.5
9		69.6	10.8
10		69.8	11.4
11		70.7	13.0
12		71.7	22.6
13		72.1	22.7
14		79.1	24.1
15		74.7	26.3
16		81.9	27.9
17		78.2	38.6

<sup>a</sup>Computed at uB3LYP/6-31G\*. <sup>b</sup>Meso diastereomer.

All told, it appears that the dimer must be able to fragment readily, hence the poor results with compounds with higher dimer BDE (Table 2, entries 12-17). The poor results with dimers that cleave very readily (entries 1-2) are consistent with two scenarios. The dimer may play a partial role in assisting the palladium activation of toluene, in which case a modest oxidizing power is needed which the dimers of **11** and **12** lack. Alternately, the captodative radical from homolysis of the dimer must engage the palladium alkyl intermediate and cannot do so if it is too stable, as manifested in very low dimer BDE values.

### Conclusions

The discovery of a range of nucleophilic substrates that can engage in the oxidative activation of alkyl C-H bonds vs arene C-H bonds with  $\text{Pd}(\text{OAc})_2$  has been described. This process uses inexpensive alkylarene precursors derived from petroleum and allows the formation of a range of hindered quaternary centers. All of the substrates react by an initial oxidative dimerization initiated by the  $\text{Pd}(\text{OAc})_2$  and/or the terminal oxidant. A number of terminal oxidants are compatible with the process with DMBQ or  $\text{K}_2\text{S}_2\text{O}_8$  being the most effective to date. The resultant dimer modifies the

palladium catalyst to favor activation of alkyl C–H bonds in contrast to the trends typically observed via a concerted metalation deprotonation mechanism. Notably, insertion occurs at the terminus of the alkyl arene for hindered substrates via  $\beta$ -hydride elimination/migratory insertion, a pathway that is not typically seen in oxidative palladium chemistry. The bond dissociation energies of the dimeric intermediates have been identified that predict which substrates are productive in this reaction. These guidelines allow rationale selection of substrates to employ in this method and further mechanistic studies are underway to understand the discrete steps of this unusual and complex mechanism.

## ASSOCIATED CONTENT

**Supporting Information.** Experimental procedures for all experiments and characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## AUTHOR INFORMATION

(1) For reviews on metal-catalyzed C–H activation, see: (a) C–H Activation. In *Topics in Current Chemistry* Vol. 292; Yu, J.-Q.; Shi, Z., Eds. Springer: Berlin, 2010. (b) Roudesly, F.; Oble, J.; Poli, G. Metal-catalyzed CH activation/functionalization: The fundamentals. *J. Mol. Cat. A: Chemical* **2017**, *426*, 275–296. (c) Chen, X.; Engle, K. M.; Wang, D. H.; Yu, J.-Q. Palladium(II)-Catalyzed C–H Activation/C–C Cross-Coupling Reactions: Versatility and Practicality. *Angew. Chem. Int. Ed.* **2009**, *48*, 5094–5115. (d) Arockiam, P. B.; Bruneau, C.; Dixneuf, P. H.; Ruthenium(II)-Catalyzed C–H Bond Activation and Functionalization. *Chem. Rev.* **2012**, *112*, 5879–5918. (e) Pototschnig, G.; Maulide, N.; Schnürch, M. Direct Functionalization of C–H Bonds by Iron, Nickel, and Cobalt Catalysis. *Chem. Eur. J.* **2017**, *23*, 1–28. (f) Usman, M.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. Recent Developments in Cobalt Catalyzed Carbon–Carbon and Carbon–Heteroatom Bond Formation via C–H Bond Functionalization. *Synthesis* **2017**, *49*, 1419–1443.

(2) Cook, A. K.; Sanford, M. S. Mechanism of the Palladium-Catalyzed Arene C–H Acetoxylation: A Comparison of Catalysts and Ligand Effects. *J. Am. Chem. Soc.* **2015**, *137*, 3109–3118.

(3) For reviews on CDC reaction, see: (a) Lyons, T. M.; Sanford, M. S. Palladium-Catalyzed Ligand-Directed C–H Functionalization Reactions. *Chem. Rev.* **2010**, *110*, 1147–1169. (b) Wencel-Delord, J.; Dröge, T.; Liu, F.; Glorius, F. Towards Mild Metal-catalyzed C–H Bond Activation. *Chem. Soc. Rev.* **2011**, *40*, 4740–4761. (c) Liu, C.; Zhang, H.; Shi, W.; Lei, A. W. Bond Formations between Two Nucleophiles: Transition Metal Catalyzed Oxidative Cross-Coupling Reactions. *Chem. Rev.* **2011**, *111*, 1780–1824. (d) Kozlowski, M. C. Oxidative Coupling in Complexity Building Transforms. *Acc. Chem. Res.* **2017**, *50*, 638–643.

(4) (a) Li, B.-J.; Tian, S.-L.; Fang, Z.; Shi, Z.-J. Multiple C–H Activations To Construct Biologically Active Molecules in a Process Completely Free of Organohalogen and Organometallic Components. *Angew. Chem. Int. Ed.* **2008**, *47*, 1115–1118. (b) Potavathri, S.; Pereira, K. C.; Gorelsky, S. I.; Pike, A.; LeBris, A. P.; DeBoef, B. Regioselective Oxidative Arylation of Indoles Bearing N-Alkyl Protecting Groups: Dual C–H Functionalization via a Concerted Metalation–Deprotonation Mechanism. *J. Am. Chem. Soc.* **2010**, *132*, 14676–14681. (c) Balcells, D.; Clot, E.; Eisenstein, C–H Bond Activation in Transition Metal Species from a Computational Perspective. *Chem. Rev.* **2010**, *110*, 749–823. (d) Wang, D.-Y.; Guo, S. H.; Pan, G.-F.; Zhu, X.-Q.; Gao, Y.-R.; Wang, Y.-Q. Direct Dehydrogenative Arylation of Benzaldehydes with Arenes Using Transient Directing Groups. *Org. Lett.* **2018**, *20*, 1794–1797.

(5) (a) Guin, S.; Rout, S. K.; Banerjee, A.; Nandi, S.; Patel, B. K. Four Tandem C–H Activations: A Sequential C–C and C–O Bond Making via a Pd-Catalyzed Cross Dehydrogenative Coupling (CDC) Approach.

## Corresponding Authors

marisa@sas.upenn.edu

## ACKNOWLEDGMENT

We are grateful to the NSF (CHE1464778, CHE1764298) and the NIH (GM-08765) for financial support of this research. Partial instrumentation support was provided by the NIH and NSF (1S10RR023444, 1S10RR022442, CHE-0840438, CHE-0848460, 1S10OD011980) and computational support by was provided by XSEDE (TG-CHE120052). G.H., P.N., U.N.K., and A.V. thank the Chinese Scholarship Council, the University of Guanajuato and CONACyT, Dr. Reddy's Laboratories (Dr. Rakeshwar Bandichhor), and the S. N. Bose Scholarship Program of India, respectively, for financial support. Drs. Rakesh Kohli and Charles W. Ross III are acknowledged for obtaining accurate mass data.

## REFERENCES

*Org. Lett.* **2012**, *14*, 5294–5297. (b) Kianmehr, E.; Faghih, Nasser.; Khan, K. M. Palladium-Catalyzed Regioselective Benzylation–Annulation of Pyridine N–Oxides with Toluene Derivatives via Multiple C–H Bond Activations: Benzylation versus Arylation. *Org. Lett.* **2015**, *17*, 414–417.

(6) (a) Xie, P.; Xie, Y. J.; Qian, B.; Zhou, H.; Xia, C.; Huang, H. M. Palladium-Catalyzed Oxidative Carbonylation of Benzylic C–H Bonds via Nondirected C(sp<sup>3</sup>)–H Activation. *J. Am. Chem. Soc.* **2012**, *134*, 9902–9905. (b) Xie, P.; Xia, C.; Huang, H. M. Palladium-Catalyzed Oxidative Aminocarbonylation: A New Entry to Amides via C–H Activation. *Org. Lett.* **2013**, *15*, 3370–3373. (c) Liu, H.; Laurenczy, G.; Yan, N.; Dyson, P. J. Amide Bond Formation via C(sp<sup>3</sup>)–H Bond Functionalization and CO Insertion. *Chem. Commun.* **2014**, *50*, 341–343. (d) Tanaka, T.; Hashiguchi, K.; Tanaka, T.; Yazaki, R.; Ohshima, T. Chemoselective Catalytic Dehydrogenative Cross-Coupling of 2-Acylimidazoles: Mechanistic Investigations and Synthetic Scope. *ACS Catal.* **2018**, *8*, 8430–8440.

(7) Sha, S.-C.; Teyrulnikov, S.; Li, M.; Hu, B.; Fu, Y.; Kozlowski, M. C.; Walsh, P. J. Cation– $\pi$  Interactions in the Benzylic Arylation of Toluenes with Bi-metallic Catalysts. *J. Am. Chem. Soc.* **2018**, *140*, 12415–12423.

(8) Curto, J. M.; Kozlowski, M. C. Chemoselective Activation of sp<sup>3</sup> vs sp<sup>2</sup> C–H Bonds with Pd(II). *J. Am. Chem. Soc.* **2015**, *137*, 18–21.

(9) (a) Dixit, V. M.; Bhat, V.; Trozzolo, A. M.; George, M. V. Sensitized photooxygenations of  $\Delta^2$ -Oxazolin-5-ones and Related Studies. *J. Org. Chem.* **1979**, *44*, 4169–4173. (b) Kato, H.; Tani, K.; Kurumisawa, H.; Tamura, Y. Photooxidation of Some Mesoionic and Related Systems. *Chem. Lett.* **1980**, *9*, 717–720. (c) Arlandini, E.; Clerici, F.; Erba, E.; Rossi, L. M. Reaction of 5-Amino-4,5-dihydro-4-methylene-1,2,3-triazoles with 2,4-Diaryl-5(4*H*)-oxazolone-4-yl Radicals. *Chem. Ber.* **1990**, *123*, 217–220. (d) Rodriguez, H.; Marquez, A.; Chuaqui, C. A.; Gomez, B. Oxidation of Mesoionic Oxazolones by Oxygen. *Tetrahedron* **1991**, *47*, 5681–5688.

(10) (a) Andersen, K. K.; Gloster, D. F.; Bray, D. D.; Shoja, M.; Kjær, A. Synthesis of Symmetrical 2,2',4,4'-tetrasubstituted[4,4'-bioxazole]-5,5'(4*H,4'**H*)-diones and Their Reactions with Some Nucleophiles. *J. Heterocycl. Chem.* **1998**, *35*, 317–324. (b) Tanaka, T.; Tanaka, T.; Tsuji, T.; Yazaki, R.; Ohshima, T. Strategy for Catalytic Chemoselective Cross-Enolate Coupling Reaction via a Transient Homocoupling Dimer. *Org. Lett.* **2018**, *20*, 3541–3544.

(11) (a) De Jongh, H. A. P.; De Jongh, C. R. H. I.; Sinnige, H. J. M.; De Klein, W. J.; Huysmans, W. G. B.; Mijs, W. J.; Van Den Hoek, W. J.; Smidt, J. Oxidative carbon–carbon coupling. II. Effect of ring substituents on the oxidative carbon–carbon coupling of arylmalonic

esters, arylmalodinitriles, and arylcyanoacetic esters *J. Org. Chem.* **1972**, *37*, 1960–1966. (b) De Jongh, H. A. P.; De Jongh, C. R. H. I.; Huysmans, W. G. B.; Sinnige, H. J. M.; De Klein, W. J. Mijs, W. J.; Jaspers, H. Radical Initiation of Vinyl Polymerization By  $\alpha$ ,  $\alpha$ ,  $\alpha'$ ,  $\alpha'$ -Tetrasubstituted Dibenzyls. *Makromolekulare Chemie*. **1972**, *157*, 279–298.

(12) (a) Hartzler, H. D. Polycyano Radicals. *J. Org. Chem.* **1966**, *31*, 2654–2658. (b) Huang, X. L.; Dannenberg, J. J. Theoretical Studies of Radical Recombination Reactions. 4. An AM1/CI Study of Reactions of Benzyllic and Allylic Radicals. An Intrinsic Barrier to Bond Formation. *J. Org. Chem.* **1991**, *56*, 6367–6371.

(13) (a) De Jongh, H. A. P.; De Jongh, C. R. H. I.; Mijs, W. J. *J. Org. Chem.* Oxidative carbon–carbon coupling. I. Oxidative coupling of alpha-substituted benzylcyanides **1971**, *36*, 3160–3168. (b) Kozlowski, M. C.; Divirgilio, E. S.; Malolanarasimhan, K.; Mulrooney, C. A. Oxidation of Chiral  $\alpha$ -phenylacetate Derivatives: Formation of Dimers with Contiguous Quaternary Stereocenters versus Tertiary Alcohols. *Tetrahedron: Asymmetry*. **2005**, *16*, 3599–3605.

(14) (a) Klasek, A.; Lycka, A.; Rouchal, M.; Rudolf, O.; Ruzicka, A. Reduction of 3-Aminoquinoline-2,4(1*H*,3*H*)-diones and Deamination of the Reaction Products. *Helvetica Chimica Acta*. **2014**, *97*, 595–612. (b) Wu, H.-R.; Huang, H.-Y.; Ren, C.-L.; Liu, L.; Wang, D.; Li, C.-J. Fe<sup>III</sup>-Catalyzed Cross-Dehydrogenative Arylation (CDA) between Oxindoles and Arenes under an Air Atmosphere. *Chem.-Eur. J.* **2015**, *21*, 16744–16748. (c) Ghosh, S.; Chaudhuri, S.; Bisai, A. Oxidative Dimerization of 2-Oxindoles Promoted by KOtBu–I<sub>2</sub>: Total Synthesis of ( $\pm$ )-Folicanthine. *Org. Lett.* **2015**, *17*, 1373–1376. (d) Bleith, T.; Deng, Q.-H.; Wadeppohl, H.; Gade, L. H. Radical Changes in Lewis Acid Catalysis: Matching Metal and Substrate. *Angew. Chem., Int. Ed.* **2016**, *55*, 7852–7856. (e) Wu, H.-R.; Cheng, L.; Kong, D.-L.; Huang, H.-Y.; Gu, C.-L.; Liu, L.; Wang, D.; Li, C.-J. FeCl<sub>3</sub>-Mediated Radical Tandem Reactions of 3-Benzyl-2-oxindoles with Styrene Derivatives for the Stereoselective Synthesis of Spirocyclohexene Oxindoles. *Org. Lett.* **2016**, *18*, 1382–1385. (f) Uraguchi, D.; Torii, M.; Ooi, T. Acridinium Betaine as a Single-Electron-Transfer Catalyst: Design and Application to Dimerization of Oxindoles. *ACS Catal.* **2017**, *7*, 2765–2769.

(15) Forrester, A. R.; Ingram A. S.; Thomson, R. H. Persulphate Oxidations. Part VII. Oxidation of o-benzyl- and o-benzoyl-benzamides. *J. Chem. Soc., Perkin Trans. 1*. **1972**, *0*, 2853–2857.

(16) (a) Scaiano, J. C.; Martin, A.; Yap, G. P. A.; Ingold, K. U. A Carbon-Centered Radical Unreactive Toward Oxygen: Unusual Radical Stabilization by a Lactone Ring. *Org. Lett.* **2000**, *2*, 899–901. (b) Frenette, M.; MacLean, P. D.; Barclay, L. R. C.; Scaiano, J. C. Radically Different Antioxidants: Thermally Generated Carbon-Centered Radicals as Chain-Breaking Antioxidants. *J. Am. Chem. Soc.* **2006**, *128*, 16432–16433. (c) Dhotare, B. B.; Kumar, M.; Nayak, S. K. Catalytic Oxidation of 3-Arylbenzofuran-2(3H)-ones with PCC–H<sub>5</sub>IO<sub>6</sub>: Syntheses of 3-Aryl-3-hydroxy/3-amido-3-arylbenzofuran-2(3H)-ones. *J. Org. Chem.* **2018**, *83*, 10089–10096.

(17) (a) Kitagawa, T.; Miyabo, A.; Fujii, H.; Okazaki, T.; Mori, T.; Matsudou, M.; Sugie, T.; Takeuchi, K. Self-Initiated Autoxidation of a Sterically Crowded Cycloheptatriene Derivative via Norcaradienylloxyl Radicals. *J. Org. Chem.* **1997**, *62*, 888–892. (b) Frenette, M.; Aliaga, C.; Sanchis, E. F.; Scaiano, J. C. Bond Dissociation Energies for Radical Dimers Derived from Highly Stabilized Carbon-Centered Radicals. *Org. Lett.* **2004**, *6*, 2579–2582. (c) Al-Afyouni, M. H.; Huang, T. A.; Hung-Low, F.; Bradley, C. A. Synthesis of Bifluorenes via Cobalt Halide Radical Coupling. *Tetrahedron Lett.* **2011**, *52*, 3261–3265.

(18) Li, J.-S.; Fu, D.-M.; Xue, Y.; Li, Z.-W.; Li, D.-L.; Da, Y.-D.; Yang, F.; Zhang, L.; Lu, C.-H.; Li, G. One-step Synthesis of Furocoumarins via Oxidative Annulation of 4-Hydroxycoumarins with DDQ. *Tetrahedron* **2015**, *71*, 2748–2752.

(19) Carole, W. A.; Colacot, T. J. Understanding Palladium Acetate from a User Perspective. *Chem. Eur. J.* **2016**, *22*, 7686–7695.

(20) For examples of Pd migration via  $\beta$ -hydride elimination/migration insertion, see: (a) Sommer, H.; Juliá-Hernández, F. Martin, R.; Marek, I. Walking Metals for Remote Functionalization. *ACS Central Science* **2018**, *4*, 153–165. (b) Le Bras, J.; Muzart, J. Palladium-Catalyzed Domino Dehydrogenation/Heck-Type Reactions of Carbonyl Compounds. *Adv. Synth. Catal.* **2018**, *360*, 2411–2428. (c) Franzoni, I.; Mazet, C. Recent trends in Pd-catalyzed remote functionalization of carbonyl compounds. *Org. Biomol. Chem.* **2014**, *12*, 233–241.

(21) (a) Takasugi, M.; Monde, K.; Katsui, N.; Shirata, A. Spirobrassinin, a Novel Sulfur-containing Phytoalexin from the Daikon Raphanus sativus L. var. hortensis (Cruciferae). *Chem. Lett.* **1987**, *16*, 1631–1632. (b) Pedras, M. S. C.; Okanga, F. I.; Zaharia, I. L. Khan, A. Q. Phytoalexins from Crucifers: Synthesis, Biosynthesis, And Biotransformation. *Phytochemistry* **2000**, *53*, 161–176. (c) Katayev, D.; Kundig, E. P. Catalytic Enantioselective Synthesis of a 3-Aryl-3-benzylloxindole ( $=$ 3-Aryl-3-benzyl-1,3-dihydro-2*H*-indol-2-one) Exhibiting Antitumor Activity. *Helv. Chim. Acta* **2012**, *95*, 2287–2295.

(22) Selected references: (a) Huang, H.-Y.; Wu, H.-R.; Wei, F.; Wang, D.; Liu, L. Iodine-Catalyzed Direct Olefination of 2-Oxindoles and Alkenes via Cross-Dehydrogenative Coupling (CDC) in Air. *Org. Lett.* **2015**, *17*, 3702–3705. (b) Hu, R.-B.; Wang, C.-H.; Ren, W.; Zhong, L.; Yang, S.-D. Direct Allylic C–H Bond Activation To Synthesize [Pd( $\eta^3$ -cin)(IPr)Cl] Complex: Application in the Allylation of Oxindoles. *ACS Catal.* **2017**, *7*, 7400–7404. (c) Gao, S.; Liu, H.; Yang, C.; Fu, Z.; Yao, H.; Lin, A. Accessing 1,3-Dienes via Palladium-Catalyzed Allylic Alkylation of Pronucleophiles with Skipped Enynes. *Org. Lett.* **2017**, *19*, 4710–4713. (d) Klare, H. F. T.; Goldber, A. F. G.; Duquette, D. C.; Stoltz, B. M. Oxidative Fragmentations and Skeletal Rearrangements of Oxindole Derivatives. *Org. Lett.* **2017**, *19*, 988–991.

(23) (a) Susanti, D.; Ng, L. L. R.; Chan, P. W. H. Silica Gel-Mediated Hydroamination/Semipinacol Rearrangement of 2-Alkylaminophenylprop-1-yn-3-ols: Synthesis of 2-Oxindoles from Alkynes and 1-(2-Aminophenyl) Ketones. *Adv. Synth. Catal.* **2014**, *356*, 353–358. (b) Li, Y.; Wang, K.; Ping, Y.; Wang, Y.; Kong, W. Nickel-Catalyzed Domino Heck Cyclization/Suzuki Coupling for the Synthesis of 3,3-Disubstituted Oxindoles. *Org. Lett.* **2018**, *20*, 921–924. (c) Panyam, P. K. R.; Ugale, B.; Gandhi, T. Palladium(II)/N-Heterocyclic Carbene Catalyzed One-Pot Sequential  $\alpha$ -Arylation/Alkylation: Access to 3,3-Disubstituted Oxindoles. *J. Org. Chem.* **2018**, *83*, 7622–7632.

(24) Wu, H.-R.; Huang, H.-Y.; Ren, C.-L.; Liu, L.; Wang, D.; Li, C.-J. Fe<sup>III</sup>-Catalyzed Cross-Dehydrogenative Arylation (CDA) between Oxindoles and Arenes under an Air Atmosphere. *Chem.-Eur. J.* **2015**, *21*, 16744–16748.

(25) (a) Nicolaou, K. C.; Snyder, S. A.; Huang, X.; Simonsen, K. B.; Koumbis, A. E.; Bigot, A. Studies toward Diazonamide A: Initial Synthetic Forays Directed toward the Originally Proposed Structure. *J. Am. Chem. Soc.* **2004**, *126*, 10162–10173. (b) Piacente, S.; Montoro, P.; Oleszek, W.; Pizza, C. Yucca schidigera Bark: Phenolic Constituents and Antioxidant Activity. *J. Nat. Prod.* **2004**, *67*, 882–885. (c) Gu, Q.; Wang, R.-R.; Zhang, X.-M.; Wang, Y.-H.; Zheng, Y.-T.; Zhou, J.; Chen, J.-J. A new benzofuranone and anti-HIV constituents from the stems of *Rhus chinensis* *Planta. Med.* **2007**, *73*, 1–4. (d) Nicolaou, K. C.; Wu, T. R.; Kang, Q.; Chen, D. T. K.; Total Synthesis of Hopeahainol A and Hopeanol. *Angew. Chem., Int. Ed.* **2009**, *48*, 3440–3443.

(26) (a) Yang, M.; Jiang, X.; Shi, W.-J.; Zhu, Q.-L.; Shi, Z.-J. Direct Lactonization of 2-Arylacetic Acids through Pd(II)-Catalyzed C–H Activation/C–O Formation. *Org. Lett.* **2013**, *15*, 690–693. (b) Cheng, X.-F.; Li, Y.; Su, Y.-M.; Yin, F.; Wang, J.-Y.; Sheng, J.; Vora, H. U.; Wang, X.-S.; Yu, J.-Q. Pd(II)-Catalyzed Enantioselective C–H Activation/C–O Bond Formation: Synthesis of Chiral Benzofuranones. *J. Am. Chem. Soc.* **2013**, *135*, 1236–1239. (c) Cho, B. S.; Chung, Y. K. Palladium-catalyzed Bisarylation of 3-Alkylbenzofurans to 3-Arylalkyl-2-Arylbenzofurans On Water: Tandem C(sp<sup>3</sup>)–H And C(sp<sup>2</sup>)–H Activation Reactions Of 3-Alkylbenzofurans. *Chem. Commun.* **2015**, *51*, 14543–14546.

(27) Blanksby, S. J.; Ellison, G. B. Bond Dissociation Energies of Organic Molecules *Acc. Chem. Res.* **2003**, *36*, 255-263.

(28) (a) Lee, H. J.; Lee, S.; Lim, J. W.; Kim, J. N. An Expedient Synthesis of Oxindole Dimers by Direct Oxidative Dimerization of Oxindoles. *Bull. Korean Chem. Soc.* **2013**, *34*, 2446-2450.