

Evaluation of DESS as a storage medium for microbial community analysis

Kevin M. Lee, Madison Adams, and Jonathan L Klassen¹

University of Connecticut, Department of Molecular and Cell Biology, Storrs CT USA

¹To whom correspondence should be addressed: jonathan.klassen@uconn.edu

9 Abstract

10 Microbial ecology research requires sampling strategies that accurately represent the
11 microbial community under study. These communities must typically be transported from
12 the collection location to the laboratory and then stored until they can be processed.
13 However, there is a lack of consensus on how best to preserve microbial communities
14 during transport and storage. Here, we evaluated DESS (Dimethyl sulfoxide,
15 Ethylenediamine tetraacetic acid, Saturated Salt) solution as a broadly applicable
16 preservative for microbial ecology experiments. We stored fungus gardens grown by the
17 ant *Trachymyrmex septentrionalis* in DESS, 15% glycerol, and phosphate buffered saline
18 (PBS) to test the ability of these preservatives to maintain the structure of fungus garden
19 microbial communities. Variation in microbial community structure due to differences in
20 preservative type was minimal when compared to variation between ant colonies.
21 Additionally, DESS preserved the structure of a defined mock community more faithfully
22 than either 15% glycerol or PBS. DESS is inexpensive, easy to transport, and effective in
23 preserving microbial community structure. We therefore conclude that DESS is a valuable
24 preservative for use in microbial ecology research.

25 Introduction

26 Microbial ecologists frequently use culture-independent methods to study the structure of
27 microbial communities (Hugerth and Andersson, 2017). Such experiments face many
28 design challenges that must be considered so that data produced matches the structure
29 of the original microbial community (Vandeputte et al., 2017; Pollock et al., 2018). Sample
30 collection, handling, and storage are the first steps in any culture-independent study, and
31 decisions made during this phase of a study can strongly affect downstream analyses
32 (Hugerth & Andersson, 2017; Vandeputte et al., 2017). Ideally, DNA should be extracted
33 from samples immediately following collection (Rochelle et al., 1994; Cardona et al.,
34 2012). However, sampling typically occurs outside of the laboratory and samples can
35 therefore experience prolonged and sometimes poorly controlled transportation and
36 storage conditions that permit nucleic acid degradation and/or microbial growth (Amir et
37 al., 2017). Either of these processes will alter the structure of the sampled microbial
38 community such that obtained the experimental data does not match the original
39 community structure in those samples.

40 Cold storage (at -80°C, -20°C, or 4°C, in order of preference) is the accepted gold
41 standard for protecting samples from potentially adverse conditions that compromise
42 microbial community structure (Rissanen et al., 2010; Williamson et al., 2011; Choo,
43 Leong & Rogers, 2015; Hale et al., 2016; Kia et al., 2016; Song et al., 2016, but see also
44 Bahl, Bergström & Licht, 2012). One disadvantage of cold storage is that cold chains are
45 fragile, especially when collection sites are remote (Vandeputte et al., 2017). Electrically
46 powered cooling requires substantial and robust physical infrastructure. Alternatively,
47 refrigerants such as ice, dry ice, and liquid nitrogen can be difficult to obtain, must be
48 refreshed routinely, and (in the case of dry ice and liquid nitrogen) are regulated or
49 prohibited on many modes of transportation. These issues limit the deployment and

robustness of cold storage for preserving microbial community samples, despite its effectiveness.

Many studies have used preservative media as an alternative solution to overcome the logistical difficulties of cold transport and storage. These include proprietary media such as DNAgard, RNAlater, OMNIgene.GUT, and LifeGuard, and less-expensive non-proprietary media such as Tris-EDTA, ethanol, and phenol-chloroform (Rissanen et al., 2010; Gaither et al., 2011; Gray, Pratte & Kellogg, 2013; Tatangelo et al., 2014; Choo, Leong & Rogers, 2015). As with cold storage, these buffers attempt to preserve the structure of the microbial community being sequenced. Many studies have compared the effectiveness of preservative media for a variety of sample types (Rissanen et al., 2010; Vlčková et al., 2012; Gray, Pratte & Kellogg, 2013; Tatangelo et al., 2014; Choo, Leong & Rogers, 2015; Anderson et al., 2016; Blekhman et al., 2016; Amir et al., 2017; Vogtmann et al., 2017), and typically conclude that most preservatives alter microbial community structure to a greater or lesser degree. Some preservatives such as ethanol and phenol-chloroform are also hazardous and therefore face travel restrictions, limiting their use for field collections. There is therefore an ongoing need to identify storage media that are effective, inexpensive, and well-suited to field collections.

Dimethyl sulfoxide (DMSO)-ethylenediamine tetraacetic acid (EDTA)-saturated salt solution (DESS, also known as SSD) is a non-proprietary, non-hazardous storage medium that shows strong potential for use as a preservative in microbial ecology studies. In this medium, DMSO permeates cells and facilitates the rapid entry of EDTA that suppresses nuclease activity by chelating divalent cations. Sodium chloride salt further suppresses enzymatic activity and contributes sodium ions that stabilize the negatively charged phosphate backbone of DNA (Seutin, White & Boag, 1991). DESS was first used to preserve avian blood samples by Seutin et al. (1991) and has since been used to preserve the morphology and DNA of nematodes (Yoder et al., 2006), corals, and small marine invertebrates (Gaither et al., 2011). DESS did not strongly alter microbial community structure when used to preserve mock microbial communities created from environmental isolates (Gray, Pratte & Kellogg, 2013) or soil and water samples, with and without cold storage (Tatangelo et al., 2014). Other DMSO-based preservatives have similar properties (Kerckhof et al., 2014). DESS therefore shows promise as a broadly applicable preservative that can overcome the fragility or lack of cold chains during sample transport between collection sites and the laboratory.

Here, we evaluated DESS as a preservative using the microbial communities found in fungus gardens grown by the ant *Trachymyrmex septentrionalis* as a model system studied by our research group. We typically collect these samples in hot and humid locations that are far from the lab, meaning that our cold chain is susceptible to failure. We further validated our field-based observations using a mock microbial community with a defined structure. Our results suggest that DESS is an excellent preservative of microbial community structure that is useful for field collections where cold transport and storage are challenging.

Materials and Methods

Sample Collection

T. septentrionalis colonies were collected in New Jersey, Florida, Georgia, and North Carolina during 2014 and 2015. Permits for collecting samples were obtained from the corresponding state department: State of New Jersey Department of Environmental Protection Division of Parks and Forestry State Park Service Unnumbered Letter of Authorization; North Carolina Division of Parks and Recreation Scientific Research and Collecting Permit 2015_0030; Florida Department of Agriculture and Consumer Services unnumbered Letter of Authorization; Georgia Department of Natural Resources State Parks & Historic Sites Scientific Research and Collection Permit 032015; Department of Natural Resources Wildlife Resources Division unnumbered Letter of Authorization. Colonies were identified by their distinctive “half-moon” mound shape and the presence of *T. septentrionalis* worker ants. An initial ~25cm deep trench was dug beside the colony entrance and then expanded until the fungus garden chamber was gently breached. After expanding this opening, the fungus garden was removed using a flame-sterilized spoon. Fungus gardens were naturally homogenized during collection by crumbling due to their fragility. Approximately 200mg of each fungus garden was subsampled into DESS (20% DMSO (v/v), 250mM EDTA, saturated with sodium chloride), 15% (v/v) glycerol, or PBS (137mM NaCl, 2.7mM KCl, 10mM Na₂HPO₄, 2mM KH₂PO₄, pH 7.4), frozen immediately on dry ice, and then transferred to -80°C storage upon return to the laboratory. Glycerol was chosen for comparison based on its potential use as a cryoprotectant that would allow cells to be cultured upon return to the lab. PBS served as a baseline that was not expected to facilitate sample preservation beyond maintaining osmotic balance.

Sample Processing

***T. septentrionalis* fungus gardens:** Samples were warmed to -20°C overnight and then thawed at 4°C just prior to processing (2-5 hours). Replicate subsamples were pooled to obtain ~1g wet mass of fungus garden (equivalent to 250-500mg dry mass). Preliminary experiments using DNA extracted directly from *T. septentrionalis* fungus garden inconsistently amplified 16S rRNA genes using PCR (data not shown). This was likely due to the high amount of fungal biomass in these samples compared to bacterial biomass and the presence of inhibitory chemical compounds (Ishak et al., 2011). We therefore adapted the protocol of (Apajalahti et al., 1998) to enrich for bacterial cells prior to DNA extraction, using medium speed vortexing to homogenize samples instead of rotational shaking and a final centrifugation at 15,000g for 15 minutes instead of 30,000g to pellet bacterial cells. DNA was extracted from the resulting cell pellets using a CTAB/bead beating protocol modified from Cafaro & Currie, 2005 by using three two-minute cycles of bead beating (Biospec Minibeadbeater) separated by 2.5 min cooling on ice for cell lysis, and 24:1 chloroform:isoamyl alcohol for DNA extraction. DNA extracts were resuspended in nuclease-free water and quantified spectrophotometrically using a BioSpec EON plate reader with a Take3Trio plate. Negative DNA extraction controls that contained water instead of sample were included alongside each extraction batch. DNA

extracts with $A_{260/280}$ or $A_{260/230}$ ratios ≤ 1.5 were purified using Agencourt XP magnetic beads (Beckman Coulter) following the manufacturer's protocol.

Mock Community: A ZymoBIOMICS Microbial Community Standard (Zymo Research Corporation) comprised of five Firmicutes (Gram positive) and three Proteobacteria (Gram negative) was used to validate our field-based experiments. Although not as diverse, the major constituents of the mock community belong to the predominant phyla that we observed in the *T. septentrionalis* fungus gardens (see Results) and are facultative anaerobes, like the common taxa that we observe in our fungus garden samples. One vial of this mock community was divided into 18 equal aliquots and each aliquot was centrifuged at 5000g for 5 minutes to pellet the cells. The supernatant containing the proprietary Zymo preservative was removed and the cells were resuspended in 1mL of DESS, PBS, or 15% glycerol by vortexing. DNA was extracted from two samples resuspended in each preservative as described above either immediately (before freezing; t_0) or after one (t_1) or two (t_2) months of storage at -80°C . These DNA extracts were quality checked as described above but none required magnetic bead cleaning. Negative controls were not used for this experiment because of the known composition of the mock communities.

PCR Screening: All quality-checked DNA extracts were screened by PCR amplification of the V4 region of the 16S rRNA gene before sequencing on an Illumina MiSeq. Each 25 μl reaction used 1X GoTaq reaction buffer (Promega), 0.3 μM each of primers 515F and 806R (Caporaso et al., 2011; Invitrogen), 1.25 units of GoTaq DNA polymerase (Promega), 300ng/ μl bovine serum albumin (New England BioLabs), 50ng of template DNA, and nuclease-free water (Thomas Scientific). PCR reactions were run on a T-100 Thermal Cycler (BioRad) for 3 min at 95°C , 30 cycles of: 30 sec at 95°C , 30 sec at 50°C , and 60 sec at 72°C , followed by a single 5 min cycle at 72°C . Bands were visualized using agarose gel electrophoresis. PCR reactions lacking the expected 350bp product were re-cleaned with magnetic beads as described above and screened by PCR a second time.

Community Amplicon Sequencing

Quality-checked, PCR-screened DNA extracts, including negative controls that lacked visible bands on an agarose gel, were submitted to the University of Connecticut Microbial Analysis, Resources, and Services (MARS) facility for sequencing. Submitted DNA samples were quantified fluorometrically using a PicoGreen (Invitrogen) assay in 384-well plates read on a Synergy plate reader (BioTek). After quantification, 30ng of sample DNA was added to 1x Phusion High Fidelity master mix (New England Biolabs) containing 1 μM indexed sequencing primers with Illumina adapters (Kozich et al., 2013; Invitrogen), 4nM non-Illumina primers (515F/806R; Caporaso et al., 2011; Invitrogen) to a final volume of 50 μl . These reactions were split into three equal aliquots and PCR amplified using settings: 94°C for 3 minutes initial denaturation, 30 cycles of 94°C for 45 seconds denaturation, 50°C for 1 minute annealing, 72°C for 1.5 minutes extension, 72°C for 10 minutes final extension. These PCR reactions were re-pooled and quantified using a QIAxcel instrument (Qiagen). Sample libraries with a PCR product concentration >

0.5ng/μl and peak(s) at 400bp (+/- 15%) were pooled by adding equal masses of PCR product from each sample, except for the negative controls that all had low PCR product concentrations and for which the largest possible mass of DNA was added instead. These pooled libraries were cleaned using Mag-Bind RXNPure Plus beads (OMEGA bio-tek), resuspended in 25μl molecular biology grade water, quantified using a Qubit assay (Invitrogen), and adjusted to 4nM (1.5ng/μl when using MARS adaptors). Amplicon libraries were diluted to 6pM in Illumina HT1 buffer with 30% PhiX phage DNA added and sequenced using a V2 (2x250) cartridge on an Illumina MiSeq instrument.

Bioinformatic Analysis

16S rRNA gene sequences were processed in R v3.4.3 (R Core Team, 2017) using the DADA2 v1.7.0 pipeline (Callahan et al., 2016) following the guidelines at https://benjjneb.github.io/dada2/tutorial_1_7.html (accessed 7/2/2018). Processed sequences and accompanying metadata were imported into phyloseq v1.22.3 (McMurdie & Holmes, 2013) and screened for potential contaminants using decontam v0.20.0 (Davis et al., 2017) using the decontam “prevalence protocol” with the P* threshold set to 0.5. Amplicon Sequence Variants (ASVs) that were not classified as bacteria were removed, as were bacterial sequences that were not classified to at least the phylum level. Decontam identified 36 fungus garden ASVs as contaminants (ranging from 0-10.2% of the sequences in each sample, mean 1.4%), which were removed from this dataset. After excluding four samples with low read counts, the final fungus garden dataset contained 26 samples from 10 *T. septentrionalis* colonies (minimum: 11,379 reads; mean: 46,125 reads). Instead of using decontam, seven ASVs (ranging from 0.4 – 3.4% of the sequences in each sample, mean 1.6%) were manually detected and removed from the mock community dataset because they did not match known members of that community. All 16 mock community samples were included in the final mock community dataset (minimum: 24,891 reads; mean: 36,445 reads). The raw sequence data have been deposited in the National Center for Biotechnology Information database under BioProject ID PRJNA479679.

The resulting phyloseq-compatible R data object was subsampled to the lowest read count among all libraries and ASV counts were converted to relative abundances. Weighted and Unweighted UniFrac distances were calculated, ordinated, and visualized using the distance, ordinate, and plot_ordination functions in phyloseq, respectively. Phyloseq was also used to calculate Shannon’s diversity, Chao1 diversity, and the observed diversity metrics using the estimate_richness function. Permutational Analysis of Variance (PERMANOVA) tests were done using Adonis in the vegan R package v2.4-5 (Oksanen et al., 2017). The aov, kruskal.test, and TukeyHSD commands implemented in the base R package were used to perform Analysis of Variance (ANOVA), Kruskal-Wallis and Tukey’s Honest Significant Difference testing, respectively. The code and metadata files to reproduce these analyses are available in Supplemental File 1: DESS_manuscript_code_and_metadata.zip.

Results

***T. septentrionalis* fungus gardens**

T. septentrionalis ants tend symbiotic fungus gardens as their food source, which also host distinct bacterial microbiomes. Our taxonomic profiles of *T. septentrionalis* fungus garden microbial communities suggest these communities are highly variable, regardless of preservative (Figure 1A). Of the 10 sampled fungus gardens, four were dominated by Firmicutes, two were dominated by Proteobacteria, and one was dominated by Planctomycetes. The remaining three colonies had high amounts of both Firmicutes and Proteobacteria (Figure 1A). Acidobacteria, Actinobacteria, Verrucomicrobia, and (except for colony JKH095) Planctomycetes had low abundances in most fungus gardens.

The type of preservative in which samples were stored only minimally correlated with variation in microbial community structure. Principal coordinate analyses (PCoA) of both Weighted and Unweighted UniFrac distances show that samples from the same colony group together. In contrast, samples stored in different media are distributed throughout each PCoA plot (Figure 1B – Weighted UniFrac, Supplemental Figure 1 – Unweighted UniFrac). Colony of origin accounted for the largest amount of the between-sample variation in β -diversity (PERMANOVA Weighted UniFrac, $F = 52.496$, $R^2 = 0.799$, $p = 0.001$, Unweighted UniFrac $F = 2.2136$, $R^2 = 0.555$, $p = 0.001$) with storage medium accounting for a very small percentage of this variation (PERMANOVA Weighted UniFrac, $F = 11.78$, $R^2 = 0.039$, $p = 0.006$, Unweighted UniFrac $F = 0.84939$, $R^2 = 0.069$, $p = 0.764$) that was linked to colony of origin (colony-by-storage medium interaction: PERMANOVA Weighted UniFrac, $F = 8.37$, $R^2 = 0.156$, $p = 0.010$, Unweighted UniFrac $F = 1.0959$, $R^2 = 0.311$, $p = 0.317$). Weighted UniFrac distances between samples from the same colony that were stored in different preservatives did not significantly differ from each other (Kruskal-Wallis, $Df = 2$, $\chi^2 = 4.779$, $p = 0.090$), implying that one preservative type did not alter microbial community structure more than the others. Like β -diversity, the α -diversity of samples collected from the same colony did not vary between preservative types (Figure 1C; Kruskal-Wallis, Shannon: $Df = 2$, $\chi^2 = 0.934$, $p = 0.627$; Chao1 and Observed: $Df = 2$, $\chi^2 = 1.127$, $p = 0.569$). Collectively, these data indicate that preservative type had at most a minor effect on the observed structure of *T. septentrionalis* fungus garden microbial communities.

Mock Community Samples

Our experiments with *T. septentrionalis* fungus gardens suggested that preservative type may have a small effect on community structure. However, the lack of a reference standard prevented us from determining which samples best represented the true structure of *T. septentrionalis* fungus garden microbial communities and which represented artificial community structures that were biased by the effects of storage. We therefore used a mock community of known composition to isolate the effect of preservative on microbial community structure. Storage at -80°C for 1 or 2 months changed the relative abundances of all taxa in all preservatives when compared to the corresponding unfrozen (t_0) samples (Figure 2A). The PCoA of the Weighted UniFrac distances showed three distinct clusters, containing either: (1) all t_0 samples; (2) t_1 and t_2

DESS samples; or (3) t_1 and t_2 glycerol and PBS samples (Figure 2B). Variation in mock community structure correlated weakly with preservative type (PERMANOVA, $F_{(2,17)} = 163.8$, $R^2 = 0.133$, $p = 0.001$). Instead, variation in mock community structure correlated strongly to storage time (PERMANOVA, $F_{(2,17)} = 977.5$, $R^2 = 0.794$, $p = 0.010$). However, the mean Weighted UniFrac distances between t_1 and t_2 samples frozen in the same preservative were very small (DESS = 0.054, glycerol = 0.055, PBS = 0.036) and not significantly different from each other (ANOVA, $F_{(2,18)} = 1.959$, $p = 0.175$). These small distances imply that “time” actually represents changes induced by freezing and/or any amount of frozen storage.

Mock community samples frozen in DESS changed less during storage at -80°C compared to the samples that were stored in either PBS or glycerol (Figure 2C). *Bacillus*, *Enterococcus*, *Listeria*, and *Staphylococcus* were all overrepresented in mock communities stored in all preservatives relative to the t_0 baseline samples. Similarly, *Escherichia*, *Lactobacillus*, *Pseudomonas*, and *Salmonella* were all underrepresented in mock communities stored in all preservatives relative to the t_0 baseline samples. The relative abundance of six out of eight taxa differed from the t_0 baseline samples less when frozen in DESS compared to glycerol and PBS (Figure 2C). The exceptions were *Lactobacillus*, whose relative abundance changed equally in all preservatives, and *Bacillus*, whose relative abundance changed more in DESS compared to the glycerol and PBS. Changes in taxon relative abundance compared to the t_0 baseline did not differ between taxa stored in glycerol or PBS (Tukey’s Honest Significance with Bonferroni correction, $p > 0.050$), except for *Staphylococcus* (Tukey’s Honest Significance with Bonferroni correction, $p = 0.016$). Collectively, these data show that the structure of mock community samples stored in DESS were more similar to that of unfrozen t_0 baseline samples compared to samples stored in glycerol or PBS.

Discussion

T. septentrionalis fungus garden microbial communities were more diverse than what has been reported for fungus gardens from related ant species. Fungus-growing ants from the genera *Atta* and *Acromyrmex* raise fungus gardens that contain low microbial diversity and are dominated by Gammaproteobacteria, as are fungus gardens grown by the only other *Trachymyrmex* species studied thus far, *T. zetekii* (Suen et al., 2010; Aylward et al., 2012, 2014). However, our results agree with the limited data available for other *T. septentrionalis* fungus gardens (Ishak et al., 2011) and some other ant genera (Kellner et al., 2015). These differences could be due to species-specific differences in habitat and foraging behavior (De Fine Licht & Boomsma, 2010), which might lead to different microbes entering fungus gardens via different substrates. Understanding how these dynamics contribute to the diversity of the *T. septentrionalis* fungus garden microbial community is an active and ongoing area of research in our lab.

Preservative type had only a small impact on *T. septentrionalis* fungus garden microbial community structure. The ~4% of variation in fungus garden microbial community structure attributed to preservative by PERMANOVA was dwarfed by the >75% of

variation attributed to the colony of origin (values are for Weighted Unifrac tests; Figure 1B). Preservative type also did not correlate with changes in the alpha-diversity of *T. septentrionalis* fungus garden microbial communities (Figure 1C). This minimal effect of preservative parallels other studies where between-host differences in microbial community structure greatly exceeded differences caused by variation in storage strategy (Bai et al., 2012; Hill et al., 2016; Song et al., 2016).

Because our *T. septentrionalis* fungus gardens were all frozen in the field, we lack a reference community against which to compare the slightly different microbial community structures of the fungus gardens stored in DESS compared to those stored in glycerol or PBS. We therefore used a mock community to more precisely assess changes in microbial community structure caused by each preservative. In these experiments, all mock communities changed relative to t_0 samples, likely indicating either a consistently negative effect of storage for 1-2 months at -80°C (Bahl, Bergström & Licht, 2012; Kia et al., 2016) and/or the consistently negative effect of the additional freeze/thaw cycle undergone by the t_1 and t_2 samples but not the t_0 samples (Cardona et al., 2012; Gorzelak et al., 2015). Four of the five Firmicutes strains in the mock community were overrepresented after storage at -80°C . Such increases in Firmicutes following cold storage has been reported previously (Bahl, Bergström & Licht, 2012; Anderson et al., 2016; Hill et al., 2016; Song et al., 2016). However, these changes were less severe for samples stored in DESS compared to those stored in glycerol or PBS, confirming previous results (Gray, Pratte & Kellogg, 2013; Tatangelo et al., 2014). These results suggest that changes caused by storage are less severe for samples stored in DESS compared to other preservatives.

Conclusions

In summary, our results suggest that DESS is an excellent storage medium for microbial ecology samples. Storage in DESS did not obscure the ecological differences between the microbial communities of fungus gardens collected from different ant colonies. DESS also preserved the community structure of a mock community more faithfully than either glycerol or PBS. Future research should focus on how microbial community structure changes when samples stored in DESS are repeatedly frozen and thawed, as might occur in a broken cold chain. DESS is inexpensive and non-hazardous, making it easy to transport. Together, these attributes suggest that DESS is a versatile and economical preservative that is suitable for the transport and storage of microbial communities.

330 **Acknowledgements**

331 We thank the members of the Klassen lab for assistance with sampling, and the University
332 of Connecticut MARS facility for assistance with library preparation and DNA sequencing.
333 We would also like to thank the personnel from the many state forests and parks from
334 which we collected ants for their valuable insights and enthusiasm for the scientific
335 research conducted in their forests and parks, and the anonymous reviewers for their
336 constructive feedback on this manuscript.

337

References

- Amir A., McDonald D., Navas-Molina JA., Debelius J., Morton JT., Hyde E., Robbins-Pianka A., Knight R. 2017. Correcting for microbial blooms in fecal samples during room-temperature shipping. *mSystems* 2:e00199-16. DOI: 10.1128/mSystems.00199-16.
- Anderson EL., Li W., Klitgord N., Highlander SK., Dayrit M., Seguritan V., Yooseph S., Biggs W., Venter JC., Nelson KE., Jones MB. 2016. A robust ambient temperature collection and stabilization strategy: Enabling worldwide functional studies of the human microbiome. *Scientific Reports* 6:31731. DOI: 10.1038/srep31731.
- Apajalahti JH., Särkilahti LK., Mäki BR., Heikkinen JP., Nurminen PH., Holben WE. 1998. Effective recovery of bacterial DNA and percent-guanine-plus-cytosine-based analysis of community structure in the gastrointestinal tract of broiler chickens. *Applied and Environmental Microbiology* 64:4084–4088. DOI: 10.1128/AEM.71.7.3911-3916.2005.
- Aylward FO., Burnum KE., Scott JJ., Suen G., Tringe SG., Adams SM., Barry KW., Nicora CD., Piehowski PD., Purvine SO., Starrett GJ., Goodwin LA., Smith RD., Lipton MS., Currie CR. 2012. Metagenomic and metaproteomic insights into bacterial communities in leaf-cutter ant fungus gardens. *ISME Journal* 6:1688–1701. DOI: 10.1038/ismej.2012.10.
- Aylward FO., Suen G., Biedermann PHW., Adams AS., Scott JJ., Malfatti SA., Del Rio TG., Tringe SG., Poulsen M., Raffa KF., Klepzig KD., Currie CR. 2014. Convergent bacterial microbiotas in the fungal agricultural systems of insects. *mBio* 5:e02077-14. DOI: 10.1128/mBio.02077-14.
- Bahl MI., Bergström A., Licht TR. 2012. Freezing fecal samples prior to DNA extraction affects the Firmicutes to Bacteroidetes ratio determined by downstream quantitative PCR analysis. *FEMS Microbiology Letters* 329:193–197. DOI: 10.1111/j.1574-6968.2012.02523.x.
- Bai G., Gajer P., Nandy M., Ma B., Yang H., Sakamoto J., Blanchard MH., Ravel J., Brotman RM. 2012. Comparison of storage conditions for human vaginal microbiome studies. *PLoS ONE* 7:e36934. DOI: 10.1371/journal.pone.0036934.
- Blekhman R., Tang K., Archie EA., Barreiro LB., Johnson ZP., Wilson ME., Kohn J., Yuan ML., Gesquiere L., Grieneisen LE., Tung J. 2016. Common methods for fecal sample storage in field studies yield consistent signatures of individual identity in microbiome sequencing data. *Scientific Reports* 6:31519. DOI: 10.1038/srep31519.
- Cafaro MJ., Currie CR. 2005. Phylogenetic analysis of mutualistic filamentous bacteria associated with fungus-growing ants. *Canadian Journal of Microbiology* 51:441–446. DOI: 10.1139/w05-023.
- Callahan BJ., McMurdie PJ., Rosen MJ., Han AW., Johnson AJA., Holmes SP. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* 13:581–583. DOI: 10.1038/nmeth.3869.

378 Caporaso JG., Lauber CL., Walters WA., Berg-Lyons D., Lozupone CA., Turnbaugh P.J.,
379 Fierer N., Knight R. 2011. Global patterns of 16S rRNA diversity at a depth of millions
380 of sequences per sample. *Proceedings of the National Academy of Sciences of the*
381 *United States of America* 108:4516–4522. DOI: 10.1073/pnas.1000080107.

382 Cardona S., Eck A., Cassellas M., Gallart M., Alastrue C., Dore J., Azpiroz F., Roca J.,
383 Guarner F., Manichanh C. 2012. Storage conditions of intestinal microbiota matter in
384 metagenomic analysis. *BMC Microbiology* 12:1471–2180. DOI: 10.1186/1471-2180-
385 12-158.

386 Choo JM., Leong LEX., Rogers GB. 2015. Sample storage conditions significantly
387 influence faecal microbiome profiles. *Scientific Reports* 5:16350. DOI:
388 10.1038/srep16350.

389 Davis NM., Proctor D., Holmes SP., Relman DA., Benjamin J. 2017. Simple statistical
390 identification and removal of contaminant sequences in marker-gene and
391 metagenomics data. *bioRxiv*:221499. DOI: <http://dx.doi.org/10.1101/221499>.

392 De Fine Licht HH., Boomsma JJ. 2010. Forage collection, substrate preparation, and diet
393 composition in fungus-growing ants. *Ecological Entomology* 35:259–269. DOI:
394 10.1111/j.1365-2311.2010.01193.x.

395 Gaither MR., Szabó Z., Crepeau MW., Bird CE., Toonen RJ. 2011. Preservation of corals
396 in salt-saturated DMSO buffer is superior to ethanol for PCR experiments. *Coral*
397 *Reefs* 30:329–333. DOI: 10.1007/s00338-010-0687-1.

398 Gorzelak MA., Gill SK., Tasnim N., Ahmadi-Vand Z., Jay M., Gibson DL. 2015. Methods
399 for improving human gut microbiome data by reducing variability through sample
400 processing and storage of stool. *PLoS ONE* 10:e0134802. DOI:
401 10.1371/journal.pone.0134802.

402 Gray MA., Pratte ZA., Kellogg CA. 2013. Comparison of DNA preservation methods for
403 environmental bacterial community samples. *FEMS Microbiology Ecology* 83:468–
404 477. DOI: 10.1111/1574-6941.12008.

405 Hale VL., Tan CL., Niu K., Yang Y., Cui D., Zhao H., Knight R., Amato KR. 2016. Effects
406 of field conditions on fecal microbiota. *Journal of Microbiological Methods* 130:180–
407 188. DOI: 10.1016/j.mimet.2016.09.017.

408 Hill CJ., Brown JRM., Lynch DB., Jeffery IB., Ryan CA., Ross RP., Stanton C., O'Toole
409 PW. 2016. Effect of room temperature transport vials on DNA quality and
410 phylogenetic composition of faecal microbiota of elderly adults and infants.
411 *Microbiome* 4:19. DOI: 10.1186/s40168-016-0164-3.

412 Hugerth LW., Andersson AF. 2017. Analysing microbial community composition through
413 amplicon sequencing: From sampling to hypothesis testing. *Frontiers in Microbiology*
414 8:1561. DOI: 10.3389/fmicb.2017.01561.

415 Ishak HD., Miller JL., Sen R., Dowd SE., Meyer E., Mueller UG. 2011. Microbiomes of ant
416 castes implicate new microbial roles in the fungus-growing ant *Trachymyrmex*
417 *septrionalis*. *Scientific Reports* 1:00204. DOI: 10.1038/srep00204.

418 Kellner K., Ishak HD., Linksvayer TA., Mueller UG. 2015. Bacterial community
419 composition and diversity in an ancestral ant fungus symbiosis. *FEMS Microbiology*
420 *Ecology* 91:fiv073. DOI: 10.1093/femsec/fiv073.

421 Kerckhof FM., Courtens ENP., Geirnaert A., Hoefman S., Ho A., Vilchez-Vargas R.,
422 Pieper DH., Jauregui R., Vlaeminck SE., Van De Wiele T., Vandamme P., Heylen K.,
423 Boon N. 2014. Optimized cryopreservation of mixed microbial communities for
424 conserved functionality and diversity. *PLoS ONE* 9:e99517. DOI:
425 10.1371/journal.pone.0099517.

426 Kia E., MacKenzie BW., Middleton D., Lau A., Waite DW., Lewis G., Chan YK., Silvestre
427 M., Cooper GJS., Poppitt SD., Taylor MW. 2016. Integrity of the human faecal
428 microbiota following long-term sample storage. *PLoS ONE* 11:e0163666. DOI:
429 10.1371/journal.pone.0163666.

430 Kozich JJ., Westcott SL., Baxter NT., Highlander SK., Schloss PD. 2013. Development
431 of a dual-index sequencing strategy and curation pipeline for analyzing amplicon
432 sequence data on the miseq illumina sequencing platform. *Applied and*
433 *Environmental Microbiology* 79:5112–5120. DOI: 10.1128/AEM.01043-13.

434 McMurdie PJ., Holmes S. 2013. Phyloseq: An R Package for Reproducible Interactive
435 Analysis and Graphics of Microbiome Census Data. *PLoS ONE* 8:e61217. DOI:
436 10.1371/journal.pone.0061217.

437 Oksanen J., Blanchet FG., Friendly M., Kindt R., Legendre P., McGlinn D., Minchin PR.,
438 O'Hara RB., Simpson GL., Solymos P., Stevens MHH., Szoecs E., Wagner H. 2017.
439 vegan: Community Ecology Package. *R package version 2.4-5*.

440 Pollock J., Glendinning L., Wisedchanwet T., Watson M. 2018. The madness of
441 microbiome: Attempting to find consensus “best practice” for 16S microbiome
442 studies. *Applied and Environmental Microbiology* 84:026627-17. DOI:
443 10.1128/AEM.02627-17.

444 R Core Team. 2017. R Development Core Team. *R: A Language and Environment for*
445 *Statistical Computing*. DOI: <http://www.R-project.org>.

446 Rissanen AJ., Kurhela E., Aho T., Oittinen T., Tirola M. 2010. Storage of environmental
447 samples for guaranteeing nucleic acid yields for molecular microbiological studies.
448 *Applied Microbiology and Biotechnology* 88:977–984. DOI: 10.1007/s00253-010-
449 2838-2.

450 Rochelle PA., Cragg BA., Fry JC., John Parkes R., Weightman AJ. 1994. Effect of sample
451 handling on estimation of bacterial diversity in marine sediments by 16S rRNA gene
452 sequence analysis. *FEMS Microbiology Ecology* 15:215–225. DOI: 10.1111/j.1574-
453 6941.1994.tb00245.x.

454 Seutin G., White BN., Boag PT. 1991. Preservation of avian blood and tissue samples for
455 DNA analyses. *Canadian Journal of Zoology*. DOI: 10.1139/z91-013.

456 Song SJ., Amir A., Metcalf JL., Amato KR., Xu ZZ., Humphrey G., Knight R. 2016.
457 Preservation methods differ in fecal microbiome stability, affecting suitability for field

studies. *mSystems* 1:e00021-16. DOI: 10.1128/mSystems.00021-16.

Suen G., Scott JJ., Aylward FO., Adams SM., Tringe SG., Pinto-Tomás AA., Foster CE., Pauly M., Weimer PJ., Barry KW., Goodwin LA., Bouffard P., Li L., Osterberger J., Harkins TT., Slater SC., Donohue TJ., Currie CR. 2010. An insect herbivore microbiome with high plant biomass-degrading capacity. *PLoS Genetics* 6:e1001129. DOI: 10.1371/journal.pgen.1001129.

Tatangelo V., Franzetti A., Gandolfi I., Bestetti G., Ambrosini R. 2014. Effect of preservation method on the assessment of bacterial community structure in soil and water samples. *FEMS Microbiology Letters* 356:32–38. DOI: 10.1111/1574-6968.12475.

Vandeputte D., Tito RY., Vanleeuwen R., Falony G., Raes J. 2017. Practical considerations for large-scale gut microbiome studies. *FEMS Microbiology Reviews* 41:S154–S167. DOI: 10.1093/femsre/fux027.

Vlčková K., Mrázek J., Kopečný J., Petrželková KJ. 2012. Evaluation of different storage methods to characterize the fecal bacterial communities of captive western lowland gorillas (*Gorilla gorilla gorilla*). *Journal of Microbiological Methods* 91:45–51. DOI: 10.1016/j.mimet.2012.07.015.

Vogtmann E., Chen J., Amir A., Shi J., Abnet CC., Nelson H., Knight R., Chia N., Sinha R. 2017. Comparison of collection methods for fecal samples in microbiome studies. *American Journal of Epidemiology* 185:115–123. DOI: 10.1093/aje/kww177.

Williamson KE., Kan J., Polson SW., Williamson SJ. 2011. Optimizing the indirect extraction of prokaryotic DNA from soils. *Soil Biology and Biochemistry* 43:736–748. DOI: 10.1016/j.soilbio.2010.04.017.

Yoder M., Tandingan De Ley I., King IW., Mundo-Ocampo M., Mann J., Blaxter M., Poiras L., De Ley P. 2006. DESS: A versatile solution for preserving morphology and extractable DNA of nematodes. *Nematology* 8:367–376. DOI: 10.1163/156854106778493448.

Figure 1

Preservative type does not alter the community structure of *T. septentrionalis* fungus gardens. (A) Relative abundances of phyla in *T. septentrionalis* fungus gardens, grouped by colony. Phyla with relative abundances <15% in all samples are grouped into a single “Other” category. (B) PCoA of Weighted UniFrac distances between *T. septentrionalis* fungus garden bacterial communities. Colors indicate preservative types, and shapes indicate samples from different colonies. (C) Alpha-diversity (Shannon’s, Observed, and Chao1) in *T. septentrionalis* fungus gardens. Center bars in the box plot indicate the median, top and bottom of boxes indicate the 25th and 75th quartile, respectively, and whiskers indicate ± 1.5 times the standard error. Single points represent outliers.




Figure 2
 DESS preserves microbial mock community structure better than PBS or glycerol. (A) Relative abundances of genera in mock community samples. (B) PCoA of Weighted UniFrac distances between mock communities stored in DESS, PBS, or glycerol, indicated by different colors, for 0, 1, or 2 months, indicated by different shapes. (C) Heatmap showing log2 fold changes in relative abundance of genera in the mock community compared to t0 samples. *Indicate where these changes were significantly different between preservatives (Bonferroni corrected p-values: *p < 0.01, ***p < 0.001).

