



Mathematics of Operations Research

Publication details, including instructions for authors and subscription information:
<http://pubsonline.informs.org>

Infinite-Horizon Average Optimality of the N-Network in the Halfin-Whitt Regime

Ari Arapostathis, Guodong Pang

To cite this article:

Ari Arapostathis, Guodong Pang (2018) Infinite-Horizon Average Optimality of the N-Network in the Halfin-Whitt Regime. *Mathematics of Operations Research* 43(3):838-866. <https://doi.org/10.1287/moor.2017.0886>

Full terms and conditions of use: <https://pubsonline.informs.org/page/terms-and-conditions>

This article may be used only for the purposes of research, teaching, and/or private study. Commercial use or systematic downloading (by robots or other automatic processes) is prohibited without explicit Publisher approval, unless otherwise noted. For more information, contact permissions@informs.org.

The Publisher does not warrant or guarantee the article's accuracy, completeness, merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications, or inclusion of an advertisement in this article, neither constitutes nor implies a guarantee, endorsement, or support of claims made of that product, publication, or service.

Copyright © 2017, INFORMS

Please scroll down for article—it is on subsequent pages

INFORMS is the largest professional society in the world for professionals in the fields of operations research, management science, and analytics.

For more information on INFORMS, its publications, membership, or meetings visit <http://www.informs.org>

Infinite-Horizon Average Optimality of the N-Network in the Halfin–Whitt Regime

Ari Arapostathis,^a Guodong Pang^b

^aDepartment of Electrical and Computer Engineering, The University of Texas at Austin, Austin, Texas 78712; ^bThe Harold and Inge Marcus Department of Industrial and Manufacturing Engineering, College of Engineering, Pennsylvania State University, University Park, Pennsylvania 16802

Contact: ari@ece.utexas.edu  <http://orcid.org/0000-0003-2207-357X> (AA); gup3@psu.edu,  <http://orcid.org/0000-0001-6468-0429> (GP)

Received: February 10, 2016

Revised: September 30, 2016; March 23, 2017

Accepted: June 17, 2017

Published Online in Articles in Advance:
February 12, 2018

MSC2010 Subject Classification: 60K25,
68M20, 90B22, 90B36

OR/MS Subject Classification: Primary:
stochastic models, stochastic networks,
queueing; secondary: optimization

<https://doi.org/10.1287/moor.2017.0886>

Copyright: © 2018 INFORMS

Abstract. We study the infinite-horizon optimal control problem for N-network queueing systems, which consists of two customer classes and two server pools, under average (ergodic) criteria in the Halfin–Whitt regime. We consider three control objectives: (1) minimizing the queueing (and idleness) cost, (2) minimizing the queueing cost while imposing a constraint on idleness at each server pool, and (3) minimizing the queueing cost while requiring fairness on idleness. The running costs can be any nonnegative convex functions having at most polynomial growth. For all three problems, we establish asymptotic optimality; namely, the convergence of the value functions of the diffusion-scaled state process to the corresponding values of the controlled diffusion limit. We also present a simple state-dependent priority scheduling policy under which the diffusion-scaled state process is geometrically ergodic in the Halfin–Whitt regime, and some results on convergence of mean empirical measures, which facilitate the proofs.

Funding: This research was supported in part by the Army Research Office [Grant W911NF-17-1-0019]. The work of the first author was also supported in part by the Office of Naval Research [Grant N00014-14-1-0196]. The work of the second author is also supported in part by the Marcus Endowment Grant at the Harold and Inge Marcus Department of Industrial and Manufacturing Engineering at Penn State.

Keywords: parallel-server network • N-network • reneging/abandonment • Halfin–Whitt (QED) regime • diffusion scaling • long-time average control • ergodic control • ergodic control with constraints • geometric ergodicity • stable Markov optimal control • asymptotic optimality


1. Introduction

Parallel server networks in the Halfin–Whitt regime have been very actively studied in recent years. Many important insights have been gained in their performance, design, and control. One important question that has mostly remained open is optimal control under the long-run average expected cost (ergodic) criterion. Since it is prohibitive to exactly solve the discrete state Markov decision problem, the plausible approach is to solve the control problem for the limiting diffusion in the Halfin–Whitt regime and use this as an approximation. However, the results in the existing literature for ergodic control of diffusions (see a good review in Arapostathis et al. [2]) cannot be directly applied to the class of diffusion models arising from the parallel server networks in the Halfin–Whitt regime. Recently, Arapostathis et al. [3] and Arapostathis and Pang [1] have developed the basic tools needed to tackle this class of ergodic control problems.

Given an optimal solution to the control problem for the diffusion limit, the important task that remains is to show it gives rise to a scheduling policy for the network and establish that any sequence of such scheduling policies is asymptotically optimal in the Halfin–Whitt regime. Under the discounted cost criterion, this task has been accomplished in Atar et al. [8] for the multiclass *V-model* (or *V-network*), which consists of multiple customer classes that are catered by servers in a single pool, and in Atar [7] for multiclass multipool networks with certain tree topologies. Under the ergodic criterion, the problem becomes much more difficult because it is intertwined with questions concerning the ergodicity of the diffusion-scaled state process under the scheduling policies. This relates to various open questions on the stochastic stability of parallel server networks in the Halfin–Whitt regime.

Stability of the multiclass *V-model* in the Halfin–Whitt regime is well treated in Gamarnik and Stolyar [14]. Stolyar [23] has recently proved the tightness of the stationary distributions of the diffusion-scaled state process for the so-called *N-network* (or *N-model*), depicted in Figure 1, with no abandonment under a static priority policy. For the *V-network*, Arapostathis et al. [3] have shown that a sequence of scheduling policies constructed from the optimal solution to the diffusion control problem under the ergodic criterion is asymptotically optimal.

Figure 1. The N-network.



In this construction, the state space is divided into a compact subset with radius in the order of the square root of the number of servers around the steady state, and its complement. An approximation to the optimal control for the diffusion is used inside this set, and a static priority policy is employed in its complement. It follows from the results of Arapostathis et al. [3] that under this sequence of scheduling policies, the state process is geometrically ergodic. The proof of asymptotic optimality takes advantage of the fact that, under the static priority scheduling policy, the state process of the V-model in the Halfin-Whitt regime is geometrically ergodic. In fact, such a static priority policy for the V-model also corresponds to a constant Markov control, under which the limiting diffusion is geometrically ergodic.

However, for multiclass multipool networks, although the optimal control problem for the limiting diffusion has been thoroughly solved in Arapostathis and Pang [1], the lack of sufficient understanding of the stochastic stability properties of the diffusion-scaled state process has been the critical obstacle to establishing asymptotic optimality. It is worth noting that this difficulty is related to the so-called joint work conservation (JWC) condition, which plays a key role in the study of multiclass multipool networks as shown in Atar [6, 7]. Although the JWC condition holds for the limiting diffusions over the entire state space, it generally holds only in a bounded subset of the state space for the diffusion-scaled process, whose radius is in the order of the number of servers around the steady state. Thus an optimal control derived from the limiting diffusion does not translate well to a scheduling policy, which is compatible with the controlled dynamics of the network on the entire state space. At the same time, although as shown in Arapostathis and Pang [1], there exists a constant Markov control under which the limiting diffusion of multiclass multipool networks is geometrically ergodic, it is unclear if this is also the case for the diffusion-scaled state processes under the corresponding static priority scheduling policy. Therefore the limiting diffusion does not offer much help in the synthesis of a suitable scheduling policy on the part of the state space where the JWC condition does not hold, and as a result constructing stable policies for multiclass multipool networks is quite a challenge.

In this paper, we address these challenging problems for the N-network. We study three ergodic control problems: (P1) minimizing the queueing (and idleness) cost, (P2) minimizing the queueing cost while imposing a constraint on the idleness of each server pool (e.g., the long-run average idleness cannot exceed a specified threshold), and (P3) minimizing the queueing cost while requiring fairness on idleness (e.g., the average idleness of the two server pools satisfies a fixed ratio condition). The running cost can be any nontrivial nonnegative convex functions having at most polynomial growth. Under its usual parameterization, the control specifies the number of customers from each class that are scheduled to each server pool, and we refer to it as a “scheduling” policy. However, the control can be also parameterized in a way so as to specify which class of customers should be scheduled to server pool 2 if it has any available servers (“scheduling” control), and which of the server pools should class 1 customers be routed to, if both pools have available servers (“routing” control). The optimal control problems for the limiting diffusion corresponding to (P1)–(P3) are well posed and in the case of (P1)–(P2), the solutions can be fully characterized via Hamilton-Jacobi-Bellman (HJB) equations, following the methods in Arapostathis and Pang [1], Arapostathis et al. [3]. The dynamic programming characterization for (P3) is more difficult. This is one of those rare examples in ergodic control where the running cost is not

bounded below or above, and there is no blanket stability property. In this paper, we establish the existence of a solution to the HJB equation, and the usual characterization of optimality for this problem.

We first present a Markov scheduling policy, for the N-network under which the diffusion-scaled state processes are geometrically ergodic in the Halfin–Whitt regime (see Section 3.2). Unlike the V-model, this scheduling policy is a state-dependent priority (SDP) policy, i.e., priorities change as the system state varies—yet it is simple to describe. This result is significant since it indicates that the ergodic control problems for the diffusion-scaled processes in the Halfin–Whitt regime have finite values. Moreover, it can be used as a scheduling policy outside a bounded subset of the state space where the JWC property might fail to hold. On the other hand, it follows from the theory in Arapostathis and Pang [1] that the controlled diffusion limit is geometrically ergodic under some constant Markov control (see Theorem 4.2 in Arapostathis and Pang [1]). In this paper, we show that a much stronger result applies for the N-network (Lemma 4.1): as long as the scheduling control is a constant Markov control with pool 2 prioritizing class 2 over 1, the controlled diffusion limit is geometrically ergodic, uniformly over all routing controls (e.g., class 1 customers prioritizing server pool 1 over 2, or an SDP policy, or even a nonstationary one).

The main results of the paper center around the proof of convergence of the value functions, which is accomplished by establishing matching lower and upper bounds (see Theorems 5.1–5.2). To prove the lower bound, the key is to show that as long as the long-run average first-order moment of the diffusion-scaled state process is finite, the associated mean empirical measures are tight and converge to an ergodic occupation measure corresponding to a stationary stable Markov control for the limiting diffusion (Lemma 7.1). In fact, we can show that for the N-network, under any admissible (work conserving) scheduling policy, the long-run average m th ($m \geq 1$) moment of the diffusion-scaled state process is bounded by the long-run average m th moment of the diffusion-scaled queue under that policy (Lemma 8.1). The lower bounds can then be deduced from these observations. It is worth noting that to establish asymptotic optimality for the fairness problem (P3), we must relax the equality in the constraint and show instead that the constraint is asymptotically feasible.

To establish the upper bound, a Markov scheduling policy is synthesized, which is the concatenation of a Markov policy induced by the solution of the ergodic control problem for the diffusion limit, and which is applied on a bounded subset of the state space where the JWC condition holds, and the SDP policy, which is applied on the complement of this set.

The proof involves the following key components. First, we apply the spatial truncation approximation technique developed in Arapostathis et al. [3] and Arapostathis and Pang [1] for the ergodic control problem for the diffusion limit. This provides us with an ϵ -optimal continuous precise control. Second, we show that under the concatenation of the Markov scheduling policy induced by this ϵ -optimal control and the SDP policy, the diffusion-scaled state processes are geometrically ergodic (Lemma 9.1). Then, we prove that the mean empirical measures of the diffusion-scaled process and control, converge to the ergodic occupation measure of the diffusion limit associated with the ϵ -optimal precise control originally selected (Lemma 7.2). Uniform integrability implied by the geometric ergodicity takes care of the rest.

1.1. Literature Review

In a certain way, the N-network has been viewed as the benchmark of multiclass multipool networks, mainly because it is simple to describe, yet it has complicated enough dynamics. There are several important studies on stochastic control of parallel server networks, focusing on N-networks. Xu et al. [30] studied the Markovian single-server N-network and showed that a threshold scheduling policy is optimal under the expected discounted and long-run average linear holding cost, using a Markov decision process approach. In the conventional (single-server) heavy-traffic regime, the N-network with two single servers, was first studied in Harrison [19], under the assumption of Poisson arrivals and deterministic services, and a “discrete-review” policy is shown to be asymptotically optimal under an infinite-horizon discounted linear queueing cost. The N-model with renewal arrival processes and general service time distributions was then studied in Bell and Williams [10], as a Brownian control problem under an infinite-horizon discounted linear queueing cost, and a threshold policy is shown to be asymptotically optimal. Ghamami and Ward [15] studied the N-network with renewal arrival processes, general service time distributions, and exponential patience times, and showed that a two-threshold scheduling policy is asymptotically optimal via a Brownian control problem under an infinite-horizon discounted linear queueing cost. Brownian control models for multiclass networks were pioneered in Harrison [18, 20] and have been extended to many interesting networks; see Williams [29] for an extensive review of that literature.

In the many-server Halfin–Whitt regime, Atar [6, 7] pioneered the study of multiclass multipool networks with abandonment (of a certain tree topology) via the corresponding control problems for the diffusion limit

under an infinite-horizon discounted cost. Gurvich and Whitt [16, 17] have studied queue-and-idleness-ratio controls for multiclass multipool networks (including the N-network) in the Halfin–Whitt regime by establishing a state-space-collapse property, under certain assumptions on the network structure and the system parameters. The N-network with many-server pools and abandonment has been recently studied in Tezcan and Dai [26], where a static priority policy is shown to be asymptotically optimal in the Halfin–Whitt regime under a finite-time horizon cost criterion. In Ward and Armony [27], some blind fair routing policies are proposed for some multiclass multipool networks (including the N-network), where the control problems are formulated to minimize the average queueing cost under a fairness constraint on the idleness.

On the other hand, most of the existing results on the stochastic control of multiclass multipool networks in the Halfin–Whitt regime have only considered either discounted cost criteria (Atar [6, 7], Atar et al. [9]) or finite-time horizon cost criteria (Dai and Tezcan [12, 13]). There is only limited work of multiclass networks under ergodic cost criteria. Arapostathis et al. [3] have recently studied the multiclass V-model under ergodic cost in the Halfin–Whitt regime. The inverted V-model is studied in Armony [4], and it is shown that the Fastest server first policy is asymptotically optimal for minimizing the steady-state expected queue length and waiting time. For the same model, Armony and Ward [5] showed that a threshold policy is asymptotically optimal for minimizing the steady-state expected queue length and waiting time subject to a “fairness” constraint on the workload division. Biswas [11] has recently studied a multiclass multipool network with “help” under an ergodic cost criterion, where each server pool has a dedicated stream of a customer class, and can help with other customer classes only when it has idle servers. The N-network does not belong to the class of models considered in Biswas [11]. For general multiclass multipool networks, Arapostathis and Pang [1] have thoroughly studied ergodic control problems for the limiting diffusion. However, as mentioned earlier, asymptotic optimality has remained open. This work makes a significant contribution in that direction, by studying the N-network. The fairness problem we study fills, in some sense (our formulation is more general), the asymptotic optimality gap in Ward and Armony [27], where the associated approximate diffusion control problems are studied via simulations.

We also feel that this work contributes to the understanding of the stability of multiclass multipool networks in the Halfin–Whitt regime. In this topic, in addition to the stability studies of the V- and N-networks in Gamarnik and Stolyar [14] and Stolyar [23], it is worthwhile mentioning the following relevant work. Stolyar and Yudovina [25] studied the stability of multiclass multipool networks under a load balancing scheduling and routing policy, “longest-queue freest-server” (LQFS-LB). They showed that the fluid limit may be unstable in the vicinity of the equilibrium point for certain network structures and system parameters, and that the sequence of stationary distributions of the diffusion-scaled processes may not be tight in the underloaded regime and the Halfin–Whitt regime. They also provided positive answers to the stability and exchange of limit results in the diffusion scale for one special class of networks. Stolyar and Yudovina [24] proved the tightness of the sequence of stationary distributions of multiclass multipool networks under a *leaf activity priority* policy (assigning static priorities to the activities in the order of sequential “elimination” of the tree leaves) in the scale $n^{1/2+\varepsilon}$ (n is the scaling parameter) for all $\varepsilon > 0$, which was extended to the diffusion scale $n^{1/2}$ in Stolyar [23]. The stability/recurrence properties for general multiclass multipool networks under other scheduling policies remain open.

As alluded above, the main challenge to establish asymptotic optimality for general multiclass multipool networks is to understand the stochastic stability/recurrence properties of the diffusion-scaled state processes in the Halfin–Whitt regime. Despite the recent development in Stolyar and Yudovina [24, 25], Stolyar [23], these are far from being adequate for proving the asymptotic optimality for general multiclass multipool networks. The stochastic stability/recurrence properties may depend critically upon the network topology and/or parameter assumptions. We believe that the methodology developed here for the N-network will provide some important insights on what stochastic stability properties are required and the roles they may play in proving asymptotic optimality.

1.2. Organization of the Paper

The notation used in this paper is summarized in Section 1.3. A detailed description of the N-network model is given in Section 2. We define the control objectives in Section 3.1 and present an SDP policy that is geometrically stable in Section 3.2. We state the corresponding ergodic control problems for the limiting diffusion, as well as the results on the characterization of optimality in Section 4. The asymptotic optimality results are stated in Section 5. We describe the system dynamics and an equivalent control parameterization in Section 6. In Section 7, we establish convergence results for the mean empirical measures for the diffusion-scaled state processes. We then prove the lower and upper bounds in Sections 8 and 9, respectively. The proof of geometric stability of the SDP policy is given in Appendices A, and B is concerned with the proof of Theorem 4.3.

1.3. Notation

The following notation is used in this paper. The symbol $\mathbb{R}$ denotes the field of real numbers, and $\mathbb{R}_+$, $\mathbb{N}$, and $\mathbb{Z}$ denote the sets of nonnegative real numbers, natural numbers, and integers, respectively. Given two real numbers a and b , the minimum (maximum) is denoted by $a \wedge b$ ($a \vee b$), respectively. Define $a^+ := a \vee 0$ and $a^- := -(a \wedge 0)$. The integer part of a real number a is denoted by $\lfloor a \rfloor$. We also let $e := (1, 1)^\top$.

For a set $A \subset \mathbb{R}^d$, we use $\bar{A}$, A^c , and $\mathbb{1}_A$ to denote the closure, the complement, and the indicator function of A , respectively. A ball of radius $r > 0$ in $\mathbb{R}^d$ around a point x is denoted by $B_r(x)$, or simply as B_r if $x = 0$. The Euclidean norm on $\mathbb{R}^d$ is denoted by $|\cdot|$, $x \cdot y$ denotes the inner product of $x, y \in \mathbb{R}^d$, and $\|x\| := \sum_{i=1}^d |x_i|$.

For a nonnegative function $g \in \mathcal{C}(\mathbb{R}^d)$, we let $\mathcal{O}(g)$ denote the space of functions $f \in \mathcal{C}(\mathbb{R}^d)$ satisfying $\sup_{x \in \mathbb{R}^d} |f(x)|/(1 \vee g(x)) < \infty$. We also let $\mathfrak{D}(g)$ denote the subspace of $\mathcal{O}(g)$ consisting of those functions f satisfying $\limsup_{|x| \rightarrow \infty} |f(x)|/(1 \vee g(x)) = 0$. Abusing the notation, $\mathcal{O}(x)$ and $\mathfrak{D}(x)$ occasionally denote generic members of these sets.

We let $\mathcal{C}_c^\infty(\mathbb{R}^d)$ denote the set of smooth real-valued functions on $\mathbb{R}^d$ with compact support. Given any Polish space $\mathcal{X}$, we denote by $\mathcal{P}(\mathcal{X})$ the set of probability measures on $\mathcal{X}$ and we endow $\mathcal{P}(\mathcal{X})$ with the Prokhorov metric. For $\nu \in \mathcal{P}(\mathcal{X})$ and a Borel measurable map $f: \mathcal{X} \rightarrow \mathbb{R}$, we often use the abbreviated notation $\nu(f) := \int_{\mathcal{X}} f d\nu$. The quadratic variation of a square integrable martingale is denoted by $\langle \cdot, \cdot \rangle$. For any path $X(\cdot)$ of a càdlàg process, we use the notation $\Delta X(t)$ to denote the jump at time t .

2. Model Description

All stochastic variables introduced below are defined on a complete probability space $(\Omega, \mathfrak{F}, \mathbb{P})$. The expectation w.r.t. $\mathbb{P}$ is denoted by $\mathbb{E}$.

2.1. The N-Network Model

Consider an N-network with two classes of jobs (or customers) and two server pools, as depicted in Figure 1. Jobs of each class arrive according to a Poisson process with rates λ_i^n , $i = 1, 2$. There are two server pools, each of which have multiple statistically identical servers, and servers in pool 1 can only serve class 1 jobs, while servers in pool 2 can serve both classes of jobs. Let N_j^n be the number of servers in pool j , $j = 1, 2$. The service times of all jobs are exponentially distributed, where jobs of class 1 are served at rates μ_{11}^n and μ_{12}^n by servers in pools 1 and 2, respectively, while jobs of class 2 are served at a rate μ_{22}^n by servers in pool 2. Throughout the paper, we set $\mu_{21}^n \equiv 0$ and $\mu_{22}^n \equiv 0$. Jobs may abandon while waiting in queue, with an exponential patience time with rate γ_i^n for $i = 1, 2$. We study a sequence of such networks indexed by an integer n , which is the order of the number of servers and let $n \rightarrow \infty$.

Throughout the paper, we assume that the parameters satisfy the following conditions.

Assumption 2.1 (Halfin–Whitt Regime). *As $n \rightarrow \infty$, the following hold:*

$$\begin{aligned} \frac{\lambda_i^n}{n} &\rightarrow \lambda_i > 0, & \frac{\lambda_i^n - n\lambda_i}{\sqrt{n}} &\rightarrow \hat{\lambda}_i, & \gamma_i^n &\rightarrow \gamma_i \geq 0, & i = 1, 2, \\ \frac{N_j^n}{n} &\rightarrow v_j > 0, & \sqrt{n}(n^{-1}N_j^n - v_j) &\rightarrow 0, & j = 1, 2, \\ \mu_{ij}^n &\rightarrow \mu_{ij} > 0, & \sqrt{n}(\mu_{ij}^n - \mu_{ij}) &\rightarrow \hat{\mu}_{ij}, & i, j = 1, 2. \end{aligned}$$

We also have

$$\lambda_1 > \mu_{11}v_1, \quad \frac{\lambda_1 - \mu_{11}v_1}{\mu_{12}v_2} + \frac{\lambda_2}{\mu_{22}v_2} = 1. \quad (1)$$

Note that (1) implies that class 1 jobs are overloaded for server pool 1, class 2 jobs are underloaded for server pool 2, and the overload of class 1 jobs can be served by server pool 2, so that both server pools are critically loaded. This assumption is referred to as the *complete resource pooling* condition (Williams [28], Atar [7]).

Let ξ^* be a constant matrix

$$\xi^* := \begin{bmatrix} 1 & \frac{\lambda_1 - \mu_{11}v_1}{\mu_{12}v_2} \\ 0 & \frac{\lambda_2}{\mu_{22}v_2} \end{bmatrix}. \quad (2)$$

The quantity ξ_{ij}^* can be interpreted as the steady-state fraction of service allocation of pool j to class i jobs in the fluid scale. Define $x^* = (x_i^*)_{i=1,2}$ and $z^* = (z_{ij}^*)_{i,j=1,2}$ by

$$x_1^* := \xi_{11}^*v_1 + \xi_{12}^*v_2, \quad x_2^* := \xi_{22}^*v_2, \quad (3)$$

$$z^* = (z_{ij}^*) := (\xi_{ij}^* v_j) = \begin{bmatrix} v_1 & \frac{\lambda_1 - \mu_{11} v_1}{\mu_{12}} \\ 0 & \frac{\lambda_2}{\mu_{22}} \end{bmatrix}. \quad (4)$$

Then, x_i^* can be interpreted as the steady-state total number of class i jobs, and z_{ij}^* can be interpreted as the steady-state number of class i jobs receiving service in pool j , in the fluid scale. It is easy to check that $e \cdot x^* = e \cdot v$, where $v := (v_1, v_2)^\top$.

For each $i = 1, 2$, let $X_i^n = \{X_i^n(t) : t \geq 0\}$ and $Q_i^n = \{Q_i^n(t) : t \geq 0\}$ be the total number of class i jobs in the system and in the queue, respectively. For each $j = 1, 2$, let $Y_j^n = \{Y_j^n(t) : t \geq 0\}$ be the number of idle servers in server pool j . For $i, j = 1, 2$, let $Z_{ij}^n = \{Z_{ij}^n(t) : t \geq 0\}$ be the number of class i jobs being served in server pool j , and note that $Z_{21}^n \equiv 0$. The following fundamental balance equations hold:

$$\begin{aligned} X_1^n(t) &= Q_1^n(t) + Z_{11}^n(t) + Z_{12}^n(t), & N_1^n &= Y_1^n(t) + Z_{11}^n(t), \\ X_2^n(t) &= Q_2^n(t) + Z_{22}^n(t), & N_2^n &= Y_2^n(t) + Z_{12}^n(t) + Z_{22}^n(t), \\ X_i^n(t) &\geq 0, & Q_i^n(t) &\geq 0, & Y_j^n(t) &\geq 0, & Z_{ij}^n(t) &\geq 0, & i, j &= 1, 2, \end{aligned} \quad (5)$$

for each $t \geq 0$. We let $Z^n = (Z_{ij}^n)_{i,j=1,2}$, $X^n = (X_i^n)_{i=1,2}$, and analogously define Q^n and Y^n .

2.2. Scheduling Control

For the control problem, we only consider work conserving policies, which are nonanticipative and preemptive. Work conservation requires that the processes Q^n and Y^n satisfy

$$Q_1^n(t) \wedge Y_j^n(t) = 0, \quad \forall j = 1, 2 \quad \text{and} \quad Q_2^n(t) \wedge Y_2^n(t) = 0, \quad \forall t \geq 0.$$

In other words, no server will idle if there is any job in a queue that the server can serve. Service preemption is allowed; that is, jobs in service at pool 2 can be interrupted and resumed at a later time to serve jobs from the other class.

Let

$$\begin{aligned} q_1(x, z) &:= x_1 - z_{11} - z_{12}, & y_1^n(x, z) &:= N_1^n - z_{11}, \\ q_2(x, z) &:= x_2 - z_{22}, & y_2^n(x, z) &:= N_2^n - z_{12} - z_{22}. \end{aligned}$$

We define the action set $\mathcal{X}^n(x)$ as

$$\begin{aligned} \mathcal{X}^n(x) &:= \{z \in \mathbb{Z}_+^{2 \times 2} : z_{21} = 0, q_1(x, z) \wedge q_2(x, z) \wedge y_1^n(x, z) \wedge y_2^n(x, z) \geq 0, \\ &\quad q_1(x, z) \wedge (y_1^n(x, z) + y_2^n(x, z)) = 0, q_2(x, z) \wedge y_2^n(x, z) = 0\}. \end{aligned}$$

Define the σ -fields

$$\begin{aligned} \mathcal{F}_t^n &:= \sigma\{X^n(0), \tilde{A}_i^n(s), \tilde{S}_{ij}^n(s), \tilde{R}_i^n(s) : i, j = 1, 2, 0 \leq s \leq t\} \vee \mathcal{N}, \\ \mathcal{G}_t^n &:= \sigma\{\delta \tilde{A}_i^n(t, r), \delta \tilde{S}_{ij}^n(t, r), \delta \tilde{R}_i^n(t, r) : i, j = 1, 2, r \geq 0\}, \end{aligned}$$

where $\mathcal{N}$ is the collection of all $\mathbb{P}$ -null sets, and

$$\begin{aligned} \tilde{A}_i^n(t) &:= A_i^n(\lambda_i^n t), & \delta \tilde{A}_i^n(t, r) &:= \tilde{A}_i^n(t+r) - \tilde{A}_i^n(t), \\ \tilde{S}_{ij}^n(t) &:= S_{ij}^n\left(\mu_{ij}^n \int_0^t Z_{ij}^n(s) ds\right), & \delta \tilde{S}_{ij}^n(t, r) &:= S_{ij}^n\left(\mu_{ij}^n \int_0^t Z_{ij}^n(s) ds + \mu_{ij}^n r\right) - \tilde{S}_{ij}^n(t), \\ \tilde{R}_i^n(t) &:= R_i^n\left(\gamma_i^n \int_0^t Q_i^n(s) ds\right), & \delta \tilde{R}_i^n(t, r) &:= R_i^n\left(\gamma_i^n \int_0^t Q_i^n(s) ds + \gamma_i^n r\right) - \tilde{R}_i^n(t). \end{aligned}$$

The processes A_i^n , S_{ij}^n and R_i^n are all rate-1 Poisson processes, representing the arrival, service, and abandonment quantities, respectively. We assume that they are mutually independent, and also independent of the initial condition $X_i^n(0)$. Note that quantities with subscript $i = 2, j = 1$ are all equal to zero. The filtration $\mathbf{F}^n := \{\mathcal{F}_t^n : t \geq 0\}$

represents the information available up to time t , and the filtration $\mathbf{G}^n := \{\mathcal{G}_t^n: t \geq 0\}$ contains the information about future increments of the processes. We say that a scheduling policy Z^n is *admissible* if

- (i) $Z^n(t) \in \mathcal{Z}^n(X^n(t))$ for all $t \geq 0$;
- (ii) $Z^n(t)$ is adapted to $\mathcal{F}_t^n$;
- (iii) $\mathcal{F}_t^n$ is independent of $\mathcal{G}_t^n$ at each time $t \geq 0$;
- (iv) for each $i, j \in \{1, 2\}$, and for each $t \geq 0$, the process $\delta \tilde{S}_{ij}^n(t, \cdot)$ agrees in law with $S_{ij}^n(\mu_{ij}^n \cdot)$, and the process $\delta \tilde{R}_i^n(t, \cdot)$ agrees in law with $R_i^n(\gamma_i^n \cdot)$.

We denote the set of all admissible scheduling policies $(Z^n, \mathbf{F}^n, \mathbf{G}^n)$ by $\mathfrak{Z}^n$. Abusing the notation, we sometimes denote this as $Z^n \in \mathfrak{Z}^n$.

Following Atar [7], we also consider a stronger condition, JWC, for preemptive scheduling policies. Namely, for each $x \in \mathbb{Z}_+^2$, there exists a rearrangement $z \in \mathcal{Z}^n(x)$ of jobs in service such that there is either no job in queue or no idling server in the system, satisfying

$$e \cdot q(x, z) \wedge e \cdot y^n(x, z) = 0. \quad (6)$$

We let $\mathcal{X}^n$ denote the set of all possible values of $\mathbb{Z}_+^2$ for which the JWC condition (6) holds, i.e.,

$$\mathcal{X}^n := \{x \in \mathbb{Z}_+^2: (6) \text{ holds for some } z \in \mathcal{Z}^n(x)\}.$$

Note that the set $\mathcal{X}^n$ may not include all possible scenarios of the system state $X^n(t)$ for finite n at each time $t \geq 0$.

We quote a result from Atar [7], which is used later.

Lemma 2.1 (Lemma 3 in Atar [7]). *There exists a constant $c_0 > 0$ such that, the collection of sets $\tilde{\mathcal{X}}^n$ defined by*

$$\tilde{\mathcal{X}}^n := \{x \in \mathbb{Z}_+^2: \|x - nx^*\| \leq c_0 n\},$$

satisfies $\tilde{\mathcal{X}}^n \subset \mathcal{X}^n$ for all $n \in \mathbb{N}$. Moreover, for any $q, y \in \mathbb{Z}_+^2$ satisfying $e \cdot q \wedge e \cdot y = 0$, and $e \cdot (x - q) = e \cdot (N^n - y) \geq 0$, we have

$$\begin{bmatrix} N_1^n - y_1 & x_1 - q_1 - (N_1^n - y_1) \\ 0 & x_2 - q_2 \end{bmatrix} \in \mathcal{Z}^n(x). \quad (7)$$

We need the following definition.

Definition 2.1. We fix some open ball $\tilde{B}$ centered at the origin such that $n(\tilde{B} + x^*) \subset \tilde{\mathcal{X}}^n$ for all $n \in \mathbb{N}$. The *jointly work conserving action set* $\tilde{\mathcal{Z}}^n(x)$ at x is defined as the subset of $\mathcal{Z}^n(x)$, which satisfies

$$\tilde{\mathcal{Z}}^n(x) := \begin{cases} \{z \in \mathcal{Z}^n(x): e \cdot q(x, z) \wedge e \cdot y^n(x, z) = 0\} & \text{if } x \in n(\tilde{B} + x^*), \\ \mathcal{Z}^n(x) & \text{otherwise.} \end{cases}$$

We also define the associated admissible policies by

$$\begin{aligned} \tilde{\mathfrak{Z}}^n &:= \{Z^n \in \mathfrak{Z}^n: Z^n(t) \in \tilde{\mathcal{Z}}^n(X^n(t)), \forall t \geq 0\}, \\ \mathfrak{Z} &:= \{Z^n \in \tilde{\mathfrak{Z}}^n: n \in \mathbb{N}\}. \end{aligned}$$

We refer to the policies in $\mathfrak{Z}$ as *eventually jointly work conserving* (EJWC).

Remark 2.1. The ball $\tilde{B}$ is fixed in Definition 2.1 only for convenience. We could instead adopt a more general definition of $\mathfrak{Z}$, without affecting the results of the paper. Let $\{D_n, n \in \mathbb{N}\}$ be a collection of domains, which covers $\mathbb{R}^2$ and satisfies $D_n \subset D_{n+1}$, and $\sqrt{n}D_n + nx^* \subset \tilde{\mathcal{X}}^n$ for all $n \in \mathbb{N}$. Then, we redefine $\tilde{\mathcal{Z}}^n$ using Definition 2.1 and replacing $n(\tilde{B} + x^*)$ with $\sqrt{n}D_n + nx^*$ and define $\mathfrak{Z}$ analogously. If $\{Z^n\} \subset \mathfrak{Z}$, then, in the diffusion scale, JWC holds on an expanding sequence of domains that cover $\mathbb{R}^2$. This is the reason behind the terminology EJWC. The EJWC condition plays a crucial role in the derivation of the controlled diffusion limit. Therefore, convergence of mean empirical measures of the diffusion-scaled state process and control, and thus also the lower and upper bounds for asymptotic optimality are established for sequences $\{Z^n, n \in \mathbb{N}\} \subset \mathfrak{Z}$.

3. Ergodic Control Problems

We define the diffusion-scaled processes $\hat{Z}^n = (\hat{Z}_{ij}^n)_{i,j \in \{1,2\}}$, $\hat{X}^n = (\hat{X}_1^n, \hat{X}_2^n)^\top$, and analogously for $\hat{Q}^n$ and $\hat{Y}^n$, by

$$\begin{aligned}\hat{X}_i^n(t) &:= \frac{1}{\sqrt{n}}(X_i^n(t) - nx_i^*), & \hat{Z}_{ij}^n(t) &:= \frac{1}{\sqrt{n}}(Z_{ij}^n(t) - nz_{ij}^*), \\ \hat{Q}_i^n(t) &:= \frac{1}{\sqrt{n}}Q_i^n(t), & \hat{Y}_j^n(t) &:= \frac{1}{\sqrt{n}}Y_j^n(t),\end{aligned}\quad (8)$$

where x^* and z^* are defined in (3)–(4).

3.1. Control Objectives

We consider three control objectives, which address the queueing (delay) and/or idleness costs in the system: (i) *unconstrained problem*, minimizing the queueing (and idleness) cost, (ii) *constrained problem*, minimizing the queueing cost while imposing a constraint on idleness, and (iii) *fairness problem*, minimizing the queueing cost while imposing a constraint on the idleness ratio between the two server pools. The running cost is a function of the diffusion-scaled processes, which are related to the unscaled ones by (8). For simplicity, in all three cost minimization problems, we assume that the initial condition $X^n(0)$ is deterministic and $\hat{X}^n(0) \rightarrow x \in \mathbb{R}^2$ as $n \rightarrow \infty$. Let $\hat{r}: \mathbb{R}_+^2 \times \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ be defined by

$$\hat{r}(q, y) := \sum_{i=1}^2 \xi_i q_i^m + \sum_{j=1}^2 \zeta_j y_j^m, \quad q \in \mathbb{R}_+^2, \quad y \in \mathbb{R}_+^2 \text{ for some } m \geq 1, \quad (9)$$

where $\xi = (\xi_1, \xi_2)^\top$ is a strictly positive vector and $\zeta = (\zeta_1, \zeta_2)^\top$ is a nonnegative vector. In the case $\zeta \equiv 0$, only the queueing cost is minimized. In (P1) below, idleness may be added as a penalty in the objective. We denote by $\mathbb{E}^{Z^n}$ the expectation operator under an admissible policy Z^n .

(P1) (*unconstrained problem*) The running cost penalizes the queueing (and idleness). Let $\hat{r}(q, y)$ be the running cost function as defined in (9). Given an initial state $X^n(0)$, and an admissible scheduling policy $Z^n \in \mathfrak{Z}^n$, we define the diffusion-scaled cost criterion by

$$J(\hat{X}^n(0), Z^n) := \limsup_{T \rightarrow \infty} \frac{1}{T} \mathbb{E}^{Z^n} \left[\int_0^T \hat{r}(\hat{Q}^n(s), \hat{Y}^n(s)) ds \right]. \quad (10)$$

The associated cost minimization problem becomes

$$\hat{V}^n(\hat{X}^n(0)) := \inf_{Z^n \in \mathfrak{Z}^n} J(\hat{X}^n(0), Z^n).$$

(P2) (*constrained problem*). The objective here is to minimize the queueing cost while imposing idleness constraints on the two server pools. Let $\hat{r}_o(q)$ be the running cost function corresponding to $\hat{r}$ in (9) with $\zeta \equiv 0$. The diffusion-scaled cost criterion $J_o(\hat{X}^n(0), Z^n)$ is defined analogously to (10) with running cost $\hat{r}_o(\hat{Q}^n(s))$; that is,

$$J_o(\hat{X}^n(0), Z^n) := \limsup_{T \rightarrow \infty} \frac{1}{T} \mathbb{E}^{Z^n} \left[\int_0^T \hat{r}_o(\hat{Q}^n(s)) ds \right].$$

Also, define

$$J_{c,j}(\hat{X}^n(0), Z^n) := \limsup_{T \rightarrow \infty} \frac{1}{T} \mathbb{E}^{Z^n} \left[\int_0^T (\hat{Y}_j^n(s))^{\tilde{m}} ds \right], \quad j = 1, 2$$

with $\tilde{m} \geq 1$. The associated cost minimization problem becomes

$$\begin{aligned}\hat{V}_c^n(\hat{X}^n(0)) &:= \inf_{Z^n \in \mathfrak{Z}^n} J_o(\hat{X}^n(0), Z^n), \\ \text{subject to } J_{c,j}(\hat{X}^n(0), Z^n) &\leq \delta_j, \quad j = 1, 2,\end{aligned}\quad (11)$$

where $\delta = (\delta_1, \delta_2)^\top$ is a positive vector.

(P3) (*fairness*) Here, we minimize the queueing cost while keeping the average idleness of the two server pools balanced. Let θ be a positive constant and let $1 \leq \tilde{m} < m$. The associated cost minimization problem becomes

$$\begin{aligned}\hat{V}_f^n(\hat{X}^n(0)) &:= \inf_{Z^n \in \mathfrak{Z}^n} J_o(\hat{X}^n(0), Z^n), \\ \text{subject to } J_{c,1}(\hat{X}^n(0), Z^n) &= \theta J_{c,2}(\hat{X}^n(0), Z^n).\end{aligned}$$

We refer to $\hat{V}^n(\hat{X}^n(0))$, $\hat{V}_c^n(\hat{X}^n(0))$, and $\hat{V}_f^n(\hat{X}^n(0))$ as the diffusion-scaled optimal values for the n th system given the initial state $X^n(0)$ for (P1)–(P3), respectively.

Remark 3.1. We choose running costs of the form (10) mainly to simplify the exposition. However, all the results of this paper still hold for more general classes of functions. Let $h_o: \mathbb{R}^2 \rightarrow \mathbb{R}_+$ be a convex function satisfying $h_o(x) \geq c_1|x|^m + c_2$ for some $m \geq 1$ and constants $c_1 > 0$ and $c_2 \in \mathbb{R}$, and $h: \mathbb{R}^2 \rightarrow \mathbb{R}_+$, $h_i: \mathbb{R} \rightarrow \mathbb{R}_+$, $i = 1, 2$, be convex functions that have at most polynomial growth. Then, we can choose $\hat{r}(q, y) = h_o(q) + h(y)$ for the unconstrained problem, and $h_i(y_i)$ as the functions in the constraints in (11) (with $\hat{r}_o = h_o$). For the problem (P3), we require in addition that $h_1 = h_2 \neq 0$ and they are in $\mathcal{O}(|x|^m)$. The analogous running costs can, of course, be used in the corresponding control problems for the limiting diffusion, which are presented later in Section 4.2.

3.2. A Geometrically Stable Scheduling Policy

We introduce a Markov scheduling policy for the N-network that results in geometric ergodicity for the diffusion-scaled state process, and also implies that the diffusion-scaled cost in the ergodic control problem (P1) is bounded, uniformly in $n \in \mathbb{N}$. Let $N_{12}^n := \lfloor \xi_{12}^* N_2^n \rfloor$ and $N_{22}^n := \lceil \xi_{22}^* N_2^n \rceil$. Note that $N_{12}^n + N_{22}^n = N_2^n$.

Definition 3.1. For each n , we define the scheduling policy $\tilde{z}^n = \tilde{z}^n(x)$, $x \in \mathbb{Z}_+^2$ by

$$\begin{aligned}\tilde{z}_{11}^n(x) &= x_1 \wedge N_1^n, \\ \tilde{z}_{12}^n(x) &= \begin{cases} (x_1 - N_1^n)^+ \wedge N_{12}^n & \text{if } x_2 \geq N_{22}^n \\ (x_1 - N_1^n)^+ \wedge (N_2^n - x_2) & \text{otherwise,} \end{cases} \\ \tilde{z}_{22}^n(x) &= \begin{cases} x_2 \wedge N_{22}^n & \text{if } x_1 \geq N_1^n + N_{12}^n \\ x_2 \wedge (N_2^n - (x_1 - N_1^n)^+) & \text{otherwise.} \end{cases}\end{aligned}$$

Note that the scheduling policy $\tilde{z}^n$ is state dependent, and can be interpreted as follows. Class 1 jobs prioritize server pool 1 over 2. Server pool 2 prioritizes the two classes of jobs depending on the system state. Whenever $x_1 \geq N_1^n + N_{12}^n$, server pool 2 allocates no more than N_{22}^n servers to class 2 jobs, while whenever $x_2 \geq N_{22}^n$, it allocates no more than N_{12}^n servers to class 1 jobs. It is easy to check that this policy $\tilde{z}^n$ is work conserving. The resulting queue length and idleness $\tilde{q}^n$ and $\tilde{y}^n$ can be obtained by the balance equations for $x \in \mathbb{Z}_+^2$,

$$\begin{aligned}\tilde{q}_1^n(x) &= x_1 - \tilde{z}_{11}^n(x) - \tilde{z}_{12}^n(x), & \tilde{q}_2^n(x) &= x_2 - \tilde{z}_{22}^n(x), \\ \tilde{y}_1^n(x) &= N_1^n - \tilde{z}_{11}^n(x), & \tilde{y}_2^n(x) &= N_2^n - \tilde{z}_{12}^n(x) - \tilde{z}_{22}^n(x).\end{aligned}$$

Definition 3.2. For each $x \in \mathbb{R}_+^2$, define

$$\tilde{x}^n(x) := (x_1 - nx_1^*, x_2 - nx_2^*), \quad \hat{x}^n(x) := \frac{\tilde{x}^n(x)}{\sqrt{n}}, \quad (12)$$

where x^* is given in (3). Also, define

$$\mathcal{S}^n := \{\hat{x}^n(x): x \in \mathbb{Z}_+^2\} \quad \check{\mathcal{S}}^n := \{\hat{x}^n(x): x \in \tilde{\mathcal{X}}^n\}.$$

For $k \geq 2$ and $\beta > 0$, we let

$$\mathcal{V}_{k,\beta}(x) := |x_1|^k + \beta|x_2|^k, \quad x \in \mathbb{R}^2. \quad (13)$$

The generator of the state process X^n under a scheduling policy z^n takes the form

$$\begin{aligned}\mathcal{L}_n^{z^n} f(x) &:= \sum_{i=1}^2 \lambda_i^n (f(x + e_i) - f(x)) + (\mu_{11}^n z_{11}^n + \mu_{12}^n z_{12}^n) (f(x - e_1) - f(x)) \\ &\quad + \mu_{22}^n z_{22}^n (f(x - e_2) - f(x)) + \sum_{i=1}^2 \gamma_i^n q_i^n (f(x - e_i) - f(x)), \quad x \in \mathbb{Z}_+^2\end{aligned} \quad (14)$$

for $f \in \mathcal{C}_b(\mathbb{R}^2)$. We can write the generator $\hat{\mathcal{L}}_n^{z^n}$ of the diffusion-scaled state process $\hat{X}^n$ using (14) and the function $\hat{x}^n$ in Definition 3.2 as

$$\hat{\mathcal{L}}_n^{z^n} f(\hat{x}) = \mathcal{L}_n^{z^n} f(\hat{x}^n(x)). \quad (15)$$

We have the following proposition.

Proposition 3.1. Let $\hat{X}^n$ denote the diffusion-scaled state process under the scheduling policy $\tilde{z}^n$ in Definition 3.1, and $\hat{\mathcal{L}}_n^{\tilde{z}^n}$ be its generator. For any $k \geq 2$, there exists $\beta_0 > 0$ such that

$$\hat{\mathcal{L}}_n^{\tilde{z}^n} \mathcal{V}_{k,\beta}(\hat{x}) \leq C_1 - C_2 \mathcal{V}_{k,\beta}(\hat{x}), \quad \forall \hat{x} \in \mathcal{S}^n, \quad \forall n \geq n_0, \quad (16)$$

for some positive constants C_1, C_2 , and $n_0 \in \mathbb{N}$, which depend on $\beta \geq \beta_0$ and k . Namely, $\hat{X}^n$ under the scheduling policy $\tilde{z}^n$ is geometrically ergodic. As a consequence, for any $k > 0$, there exists $n_0 \in \mathbb{N}$ such that

$$\sup_{n \geq n_0} \limsup_{T \rightarrow \infty} \frac{1}{T} \mathbb{E}^{\tilde{z}^n} \left[\int_0^T |\hat{X}^n(s)|^k ds \right] < \infty, \quad (17)$$

and the same holds if we replace $\hat{X}^n$ with $\hat{Q}^n$ or $\hat{Y}^n$ in (17). In other words, the diffusion-scaled cost criterion $J(\hat{X}^n(0), Z^n)$ is finite for $n \geq n_0$.

Proof. See Appendix A. $\square$

Remark 3.2. We remark that given (17) for $\hat{X}^n$, the same property may not hold for $\hat{Q}^n$ or $\hat{Y}^n$. It always holds if a scheduling policy satisfies the JWC condition (by the balance Equation (39)). Otherwise, that property needs to be verified under the given scheduling policy. It can easily be checked that if the property holds for any two processes of $\hat{X}^n, \hat{Q}^n$, and $\hat{Y}^n$, then it also holds for the third.

4. Ergodic Control of the Limiting Diffusion

4.1. The Controlled Diffusion Limit

If the action space is $\tilde{\mathcal{X}}^n$, or equivalently, $Z^n \in \tilde{\mathcal{Z}}^n$, the convergence in distribution of the diffusion-scaled processes $\hat{X}^n$ to the limiting diffusion X in (18) is shown in proposition 3 in Atar [7]. For the class of multiclass multipool networks, the drift of the limiting diffusion is given implicitly via a linear map in proposition 3 of Atar [7]. For the N-network, the drift can be explicitly expressed as we show below in (21). In Arapostathis and Pang [1], a leaf elimination algorithm has been developed to provide an explicit expression for the drift of the limiting diffusion of general multiclass multipool networks. In the case of the N-network, the limit process X is a two-dimensional diffusion satisfying the Itô equation

$$dX_t = b(X_t, U_t) dt + \Sigma dW_t, \quad (18)$$

with initial condition $X_0 = x$ and the control $U_t \in \mathbb{U}$, where

$$\mathbb{U} := \{u = (u^c, u^s) \in \mathbb{R}_+^2 \times \mathbb{R}_+^2 : e \cdot u^c = e \cdot u^s = 1\}. \quad (19)$$

In (18), the process W is a two-dimensional standard Wiener process independent of the initial condition $X_0 = x$.

Following the leaf elimination algorithm for the N-network, the drift of the diffusion can be computed as follows. Let

$$\hat{G}[u](x) := \begin{pmatrix} -(e \cdot x)^- u_1^s & x_1 - (e \cdot x)^+ u_1^c + (e \cdot x)^- u_1^s \\ 0 & x_2 - (e \cdot x)^+ u_2^c \end{pmatrix}, \quad u \in \mathbb{U}. \quad (20)$$

Then, the drift $b: \mathbb{R}^2 \times \mathbb{U} \rightarrow \mathbb{R}^2$ takes the form

$$b(x, u) = \begin{pmatrix} -\mu_{11} \hat{G}_{11}[u](x) - \mu_{12} \hat{G}_{12}[u](x) - \gamma_1 (e \cdot x)^+ u_1^c + l_1 \\ \mu_{22} \hat{G}_{22}[u](x) - \gamma_2 (e \cdot x)^+ u_2^c + l_2 \end{pmatrix},$$

which can also be written as (see lemma 4.3 and section 4.2 in Arapostathis and Pang [1])

$$b(x, u) = -B_1(x - (e \cdot x)^+ u^c) + (e \cdot x)^- B_2 u^s - (e \cdot x)^+ \Gamma u^c + l, \quad (21)$$

with

$$B_1 := \text{diag}\{\mu_{12}, \mu_{22}\}, \quad B_2 := \text{diag}\{\mu_{11} - \mu_{12}, 0\}, \quad \Gamma := \text{diag}\{\gamma_1, \gamma_2\}.$$

Here, $l := (l_1, l_2)^T$ defined by

$$l_1 := \hat{\lambda}_1 - \hat{\mu}_{11} z_{11}^* - \hat{\mu}_{12} z_{12}^*, \quad \text{and} \quad l_2 := \hat{\lambda}_2 - \hat{\mu}_{22} z_{22}^*. \quad (22)$$

The covariance matrix is given by $\Sigma := \text{diag}(\sqrt{2\lambda_1}, \sqrt{2\lambda_2})$. The control process U lives in the compact set $\mathbb{U}$ in (19), and $U_t(\omega)$ is jointly measurable in $(t, \omega) \in [0, \infty) \times \Omega$. Moreover, it is *nonanticipative* for $s < t$, $W_t - W_s$ is independent of

$$\mathfrak{F}_s := \text{the completion of } \sigma\{X_0, U_r, W_r, r \leq s\} \text{ relative to } (\mathfrak{F}, \mathbb{P}).$$

Let $\mathbb{U}$ be the set of all such controls, referred to as *admissible* controls. We refer the reader to Section 6.2 on the control parameterization. A mere comparison of (20) with (44) makes it clear how the control process U relates to the control process U^n for the n th system in Definition 6.1.

We remark that (18) can be regarded as a piecewise-linear controlled diffusion. Note that the matrix B_1 is an M -matrix. However, there is an additional term $(e \cdot x)^- B_2 u^s$ in the drift, which differs from the class of piecewise-linear controlled diffusions discussed in section 3.3 of Arapostathis et al. [3]. We refer to (18) as the *limiting diffusion*, or the *diffusion limit*.

The associated limit processes Q , Y , and Z satisfy the following balance equations:

$$\begin{aligned} X_1(t) &= Q_1(t) + Z_{11}(t) + Z_{12}(t), & Y_1(t) + Z_{11}(t) &= 0, \\ X_2(t) &= Q_2(t) + Z_{22}(t), & Y_2(t) + Z_{12}(t) + Z_{22}(t) &= 0 \end{aligned}$$

with $Q_i(t) \geq 0$, $Y_j(t) \geq 0$, $i, j = 1, 2$. Note that these “balance” conditions imply that JWC always holds at the diffusion limit, i.e.,

$$e \cdot Q(t) = (e \cdot X(t))^+, \quad e \cdot Y(t) = (e \cdot X(t))^- , \quad \forall t \geq 0.$$

4.2. Control Problems for the Diffusion Limit

We state the three problems, which correspond to (P1)–(P3) in Section 3.1 for the controlled diffusion in (18). Let $r : \mathbb{R}^2 \times \mathbb{U} \rightarrow \mathbb{R}$ be defined by

$$r(x, u) = r(x, (u^c, u^s)) := \hat{r}((e \cdot x)^+ u^c, (e \cdot x)^- u^s),$$

with the same $\hat{r}$ in (9); that is,

$$r(x, u) = [(e \cdot x)^+]^m \sum_{i=1}^2 \xi_i (u_i^c)^m + [(e \cdot x)^-]^m \sum_{j=1}^2 \zeta_j (u_j^s)^m, \quad m \geq 1 \quad (23)$$

for the given $\xi = (\xi_1, \xi_2)^T$ and $\zeta = (\zeta_1, \zeta_2)^T$ in (9). Let the ergodic cost associated with the controlled diffusion X and the running cost r be defined as

$$J_{x,u}[r] := \limsup_{T \rightarrow \infty} \frac{1}{T} \mathbb{E}_x^U \left[\int_0^T r(X_t, U_t) dt \right], \quad U \in \mathbb{U}.$$

(P1') (*unconstrained problem*). The running cost function $r(x, u)$ is as in (23). The ergodic control problem is then defined as

$$\varrho^*(x) = \inf_{U \in \mathbb{U}} J_{x,u}[r]. \quad (24)$$

(P2') (*constrained problem*). The running cost function $r_o(x, u)$ is as in (23) with $\zeta \equiv 0$. Also, define

$$r_j(x, u) := [(e \cdot x)^- u_j^s]^{\tilde{m}}, \quad j = 1, 2, \quad (25)$$

with $\tilde{m} \geq 1$, and let $\delta = (\delta_1, \delta_2)$ be a positive vector. The ergodic control problem under idleness constraints is defined as

$$\begin{aligned} \varrho_c^*(x) &= \inf_{U \in \mathbb{U}} J_{x,u}[r_o], \\ \text{subject to } & J_{x,u}[r_j] \leq \delta_j, \quad j = 1, 2. \end{aligned} \quad (26)$$

(P3') (*fairness*) The running costs r_o , r_1 , and r_2 are as in (P2'). Let θ be a positive constant, and $1 \leq \tilde{m} < m$. The ergodic control problem under idleness fairness is defined as

$$\begin{aligned} \varrho_f^*(x) &= \inf_{U \in \mathbb{U}} J_{x,u}[r_o], \\ \text{subject to } & J_{x,u}[r_1] = \theta J_{x,u}[r_2]. \end{aligned} \quad (27)$$

The last problem enforces fairness of idleness allocation among the two server pools. Also, note that penalizing only the queueing cost in (P1) raises a well-posedness question, which was resolved in corollaries 4.1–4.2 of Arapostathis and Pang [1].

The quantities $\varrho^*(x)$, $\varrho_c^*(x)$, and $\varrho_t^*(x)$ are called the optimal values of the ergodic control problems (P1')–(P3'), respectively, for the controlled diffusion process X with initial state x . Note that as is shown in section 3 of Arapostathis et al. [3] and sections 3 and 5.4 of Arapostathis and Pang [1], the optimal values $\varrho^*(x)$, $\varrho_c^*(x)$, and $\varrho_t^*(x)$ do not depend on $x \in \mathbb{R}^2$, and thus we remove their dependence on x in the statements below.

Recall that a control is called *Markov* if $U_t = v(t, X_t)$ for a measurable map $v: \mathbb{R}_+ \times \mathbb{R}^2 \rightarrow \mathbb{U}$, and it is called *stationary Markov* if v does not depend on t , i.e., $v: \mathbb{R}^2 \rightarrow \mathbb{U}$. Let $\mathbb{U}_{\text{SM}}$ denote the set of stationary Markov controls. Recall also that a control $v \in \mathbb{U}_{\text{SM}}$ is called *stable* if the controlled process is positive recurrent. We denote the set of such controls by $\mathbb{U}_{\text{SSM}}$, and let μ_v denote the unique invariant probability measure on $\mathbb{R}^2$ for the diffusion under the control $v \in \mathbb{U}_{\text{SSM}}$. We also let $\mathcal{M} := \{\mu_v: v \in \mathbb{U}_{\text{SSM}}\}$, and $\mathcal{G}$ denote the set of ergodic occupation measures corresponding to controls in $\mathbb{U}_{\text{SSM}}$; that is,

$$\mathcal{G} := \left\{ \pi \in \mathcal{P}(\mathbb{R}^2 \times \mathbb{U}): \int_{\mathbb{R}^2 \times \mathbb{U}} \mathcal{L}^u f(x) \pi(dx, du) = 0 \quad \forall f \in \mathcal{C}_c^\infty(\mathbb{R}^2) \right\},$$

where $\mathcal{L}^u f(x)$ is the controlled extended generator of the diffusion X ,

$$\mathcal{L}^u f(x) := \frac{1}{2} \sum_{i,j=1}^2 a_{ij} \partial_{ij} f(x) + \sum_{i=1}^2 b_i(x, u) \partial_i f(x), \quad u \in \mathbb{U},$$

with $a := \Sigma \Sigma^\top$ and $\partial_i := \partial / \partial x_i$ and $\partial_{ij} := \partial^2 / \partial x_i \partial x_j$. The restriction of the ergodic control problem with running cost r to stable stationary Markov controls is equivalent to minimizing

$$\pi(r) = \int_{\mathbb{R}^2 \times \mathbb{U}} r(x, u) \pi(dx, du)$$

over all $\pi \in \mathcal{G}$. If the infimum is attained in $\mathcal{G}$, then we say that the ergodic control problem is *well posed*, and we refer to any $\bar{\pi} \in \mathcal{G}$ that attains this infimum as an *optimal ergodic occupation measure*.

We define the class of admissible controls $\mathbb{U} := \{U = (U^c, U^s): U^c = v^c(x) = (1, 0) \quad \forall x \in \mathbb{R}^2\}$, and we also let

$$\bar{\beta}_k := \frac{(\gamma_1 \vee \mu_{11} \vee \mu_{12})^{k+1}}{\mu_{22}(\gamma_1 \wedge \mu_{11} \wedge \mu_{12})^k}. \quad (28)$$

We have the following lemma.

Lemma 4.1. *Let $\mathcal{V}_{k,\beta}$ be as in (13). There exist positive constants C_1 and C_2 depending only on k and $\beta \geq \bar{\beta}_k$ such that*

$$\mathcal{L}^U \mathcal{V}_{k,\beta}(x) \leq C_1 - C_2 \mathcal{V}_{k,\beta}(x), \quad \forall U \in \mathbb{U}, \quad \forall x \in \mathbb{R}^2.$$

Proof. By (21), we have

$$\begin{aligned} b_1(x, U) &= \begin{cases} -\gamma_1 x_1 + (\mu_{12} - \gamma_1) x_2 + l_1 & \text{if } (e \cdot x)^+ \geq 0 \\ -(\mu_{11} U_1^s + \mu_{12} U_2^s) x_1 - (\mu_{11} - \mu_{12}) U_1^s x_2 + l_1 & \text{otherwise,} \end{cases} \\ b_2(x, U) &= -\mu_{22} x_2 + l_2, \quad \forall x \in \mathbb{R}^2. \end{aligned}$$

Therefore

$$\begin{aligned} \mathcal{L}^U \mathcal{V}_{k,\beta}(x) &\leq -k(\gamma_1 \wedge \mu_{11} \wedge \mu_{12}) |x_1|^k + k(\gamma_1 \vee \mu_{11} \vee \mu_{12}) |x_2| |x_1|^{k-1} - \beta k \mu_{22} |x_2|^k \\ &\quad + k l_1 |x_1|^{k-1} + \beta k l_2 |x_2|^{k-1} + k(k-1)(\lambda_1 |x_1|^{k-2} + \lambda_2 \beta |x_2|^{k-2}). \end{aligned} \quad (29)$$

Let

$$\alpha := \frac{\gamma_1 \wedge \mu_{11} \wedge \mu_{12}}{\gamma_1 \vee \mu_{11} \vee \mu_{12}}.$$

Using Young's inequality, we write

$$|x_2| |x_1|^{k-1} \leq (k-1) \frac{\alpha^{k/(k-1)}}{k} |x_1|^k + \frac{\alpha^{-k}}{k} |x_2|^k \leq (k-1) \frac{\alpha}{k} |x_1|^k + \frac{\alpha^{-k}}{k} |x_2|^k.$$

Thus, by (29), we have

$$\begin{aligned} \mathcal{L}^U \mathcal{V}_{k,\beta}(x) &\leq -(\gamma_1 \wedge \mu_{11} \wedge \mu_{12})|x_1|^k - (k\beta\mu_{22} - \bar{\beta}_k)|x_2|^k + k l_1 |x_1|^{k-1} + \beta k l_2 |x_2|^{k-1} \\ &\quad + k(k-1)(\lambda_1 |x_1|^{k-2} + \lambda_2 \beta |x_2|^{k-2}), \end{aligned}$$

from which the result easily follows. $\square$

As shown in corollary 4.2 of Arapostathis and Pang [1], for any $k \geq 1$, there exists a constant $C = C(k) > 0$ such that any solution X_t of (18) with $X_0 = x_0 \in \mathbb{R}^2$ satisfies

$$\mathbb{E}_x^U \left[\int_0^T |X_t|^k dt \right] \leq C|x_0|^k + CT + C \mathbb{E}_x^U \left[\int_0^T ((e \cdot X_t)^+)^k dt \right], \quad \forall U \in \mathcal{U}, \quad \forall T > 0. \quad (30)$$

This property plays a crucial role in solving (P1')–(P3').

4.3. Optimal Solutions to Problems (P1')–(P3')

The characterization of the optimal solutions to the ergodic control problems (P1')–(P3') has been thoroughly studied in Arapostathis et al. [3] and Arapostathis and Pang [1]. We review some results that are used in the sections, which follow to construct asymptotically optimal scheduling policies and prove asymptotic optimality. We first introduce some notation. Let

$$H_r(x, p) := \min_{u \in \mathcal{U}} [b(x, u) \cdot p + r(x, u)] \quad \text{for } x, p \in \mathbb{R}^2. \quad (31)$$

For $\delta = (\delta_1, \delta_2) \in \mathbb{R}_+^2$, let

$$\mathcal{H}(\delta) := \{\pi \in \mathcal{G}: \pi(r_j) \leq \delta_j, j = 1, 2\} \quad \mathcal{H}^o(\delta) := \{\pi \in \mathcal{G}: \pi(r_j) < \delta_j, j = 1, 2\}.$$

For $\delta \in \mathbb{R}_+^2$, and $\lambda = (\lambda_1, \lambda_2)^T \in \mathbb{R}_+^2$, define the running cost $g_{\delta, \lambda}$ by

$$g_{\delta, \lambda}(x, u) := r_o(x, u) + \sum_{j=1}^2 \lambda_j (r_j(x, u) - \delta_j).$$

We say that the vector $\delta \in (0, \infty)^2$ is *feasible* (or that the constraints in (26) are feasible) if there exists $\pi' \in \mathcal{H}^o(\delta)$ such that $\pi'(r_o) < \infty$. The following is contained in theorem 5.2 of Arapostathis and Pang [1].

Theorem 4.1. *For the ergodic control problem in (24), there exists a unique solution $V \in \mathcal{C}^2(\mathbb{R}^2)$, satisfying $V(0) = 0$ to the associated HJB equation:*

$$\min_{u \in \mathcal{U}} [\mathcal{L}^u V(x) + r(x, u)] = \varrho^*.$$

Moreover, a stationary Markov control $v \in \mathcal{U}_{\text{SSM}}$ is optimal if and only if it satisfies

$$H_r(x, \nabla V(x)) = b(x, v(x)) \cdot \nabla V(x) + r(x, v(x)) \quad \text{a.e. in } \mathbb{R}^2.$$

The following is contained in lemmas 3.3–3.5, and theorems 3.1–3.2 of Arapostathis and Pang [1].

Theorem 4.2. *Suppose that δ is feasible for the ergodic control problem under constraints in (26), i.e., there exists $\pi' \in \mathcal{H}^o(\delta)$ such that $\pi'(r_o) < \infty$. Then, the following hold:*

(a) *There exists $\lambda^* \in \mathbb{R}_+^2$ such that*

$$\inf_{\pi \in \mathcal{H}(\delta)} \pi(r_o) = \inf_{\pi \in \mathcal{G}} \pi(g_{\delta, \lambda^*}) = \varrho_c^*.$$

(b) *If $\pi^* \in \mathcal{H}(\delta)$ attains the infimum of $\pi \mapsto \pi(r_o)$ in $\mathcal{H}(\delta)$, then $\pi^*(r_o) = \pi^*(g_{\delta, \lambda^*})$, and*

$$\pi^*(g_{\delta, \lambda}) \leq \pi^*(g_{\delta, \lambda^*}) \leq \pi(g_{\delta, \lambda^*}) \quad \forall (\pi, \lambda) \in \mathcal{G} \times \mathbb{R}_+^2.$$

(c) *There exists $V_c \in \mathcal{C}^2(\mathbb{R}^2)$ satisfying*

$$\min_{u \in \mathcal{U}} [\mathcal{L}^u V_c(x) + g_{\delta, \lambda^*}(x, u)] = \pi^*(g_{\delta, \lambda^*}) = \varrho_c^*, \quad x \in \mathbb{R}^2.$$

(d) A stationary Markov control $v_c \in \Pi_{\text{SSM}}$ is optimal if and only if it satisfies

$$H_{g_{\delta, \lambda^*}}(x, \nabla V_c(x)) = b(x, v_c(x)) \cdot \nabla V_c(x) + g_{\delta, \lambda^*}(x, v_c(x)) \quad \text{a.e. in } \mathbb{R}^2,$$

where $H_{g_{\delta, \lambda^*}}$ is defined in (31) with r replaced by g_{δ, λ^*} .

(e) The map $\delta \mapsto \inf_{\pi \in \mathcal{H}(\delta)} \pi(r_o)$ is continuous at any feasible point $\hat{\delta}$.

For uniqueness of the solutions V_c , see theorem 3.2 in Arapostathis and Pang [1].

We now turn to the constrained ergodic control problem in (27). Lemma 4.1 implies that assumption 5.1 in Arapostathis and Pang [1] holds, and consequently, the solution of (P3') follows by theorem 5.8 in the same paper. However, the Lagrangian in (P3') is not bounded below in $\mathbb{R}^2$, and since no details were provided in Arapostathis and Pang [1] on the existence of solutions to the HJB equation, we provide a proof in Appendix B.

Theorem 4.3. For any $\theta > 0$, the constraint in (27) is feasible. All the conclusions of Theorem 4.2 hold, provided that we replace $\mathcal{H}(\delta)$ and $g_{\delta, \lambda}$ with

$$\mathcal{H}_\theta(\theta) := \{\pi \in \mathcal{G} : \pi(r_1) = \theta \pi(r_2)\}, \quad (32)$$

and

$$h_{\theta, \lambda}(x, u) := r_o(x, u) + \lambda(r_1(x, u) - \theta r_2(x, u)), \quad \lambda \in \mathbb{R},$$

respectively.

5. Asymptotic Optimality

In this section, we present the main results on asymptotic optimality. We show that the values of the three ergodic control problems in the diffusion scale converge to the values of the corresponding ergodic control problems for the limiting diffusion, respectively. The proofs of the lower and upper bounds are given in Sections 8 and 9, respectively.

Recall the definitions of J , J_o , $\hat{V}^n$, $\hat{V}_c^n$, and $\hat{V}_i^n$ in (P1)–(P3), and the definitions of ϱ^* , ϱ_c^* , and ϱ_i^* in (P1')–(P3').

Theorem 5.1 (Lower Bounds). Let $\hat{X}^n(0) \Rightarrow x \in \mathbb{R}^2$ as $n \rightarrow \infty$. The following hold:

(i) For any sequence $\{Z^n, n \in \mathbb{N}\} \subset \mathfrak{Z}$, the diffusion-scaled cost in (10) satisfies

$$\liminf_{n \rightarrow \infty} J(\hat{X}^n(0), \hat{Z}^n) \geq \varrho^*.$$

(ii) Suppose that under a sequence $\{Z^n, n \in \mathbb{N}\} \subset \mathfrak{Z}$, the constraint in (11) is satisfied for all sufficiently large $n \in \mathbb{N}$. Then,

$$\liminf_{n \rightarrow \infty} J_o(\hat{X}^n(0), \hat{Z}^n) \geq \varrho_c^*,$$

and as a result, we have that $\liminf_{n \rightarrow \infty} \hat{V}_c^n(\hat{X}^n(0)) \geq \varrho_c^*$.

(iii) There exists a positive constant $\hat{C}$, such that if a sequence $\{Z^n, n \in \mathbb{N}\} \subset \mathfrak{Z}$ satisfies

$$\left| \frac{J_{c,1}(\hat{X}^n(0), Z^n)}{J_{c,2}(\hat{X}^n(0), Z^n)} - \theta \right| \leq \epsilon \quad (33)$$

for some $\epsilon \in (0, \theta)$, and all sufficiently large $n \in \mathbb{N}$, then

$$\liminf_{n \rightarrow \infty} J_o(\hat{X}^n(0), Z^n) \geq \varrho_i^* - \hat{C}\epsilon. \quad (34)$$

The proof of the theorem that follows relies on the fact that r and also r_i for $i = 0, 1, 2$, are convex functions of u .

Theorem 5.2 (Upper Bounds). Let $\hat{X}^n(0) \Rightarrow x \in \mathbb{R}^2$ as $n \rightarrow \infty$. The following hold:

(i) $\limsup_{n \rightarrow \infty} \hat{V}^n(\hat{X}^n(0)) \leq \varrho^*$.

(ii) For any $\epsilon > 0$, there exists a sequence $\{Z^n, n \in \mathbb{N}\} \subset \mathfrak{Z}$ such that the constraint in (11) is feasible for all sufficiently large n , and

$$\limsup_{n \rightarrow \infty} J_o(\hat{X}^n(0), \hat{Z}^n) \leq \varrho_c^* + \epsilon.$$

Consequently, we have that $\limsup_{n \rightarrow \infty} \hat{V}_c^n(\hat{X}^n(0)) \leq \varrho_c^*$.

(iii) For any $\epsilon > 0$, there exists a sequence $\{Z^n, n \in \mathbb{N}\} \subset \mathfrak{Z}$ such that (33) holds for all sufficiently large $n \in \mathbb{N}$, and

$$\limsup_{n \rightarrow \infty} J_o(\hat{X}^n(0), Z^n) \leq \varrho_i^* + \epsilon.$$

6. System Dynamics and an Equivalent Control Parameterization

6.1. Description of the System Dynamics

The processes X^n can be represented via rate 1 Poisson processes: for each $i = 1, 2$, and $t \geq 0$, we have

$$\begin{aligned} X_1^n(t) &= X_1^n(0) + A_1^n(\lambda_1^n t) - \sum_{j=1,2} S_{1j}^n \left(\mu_{1j}^n \int_0^t Z_{1j}^n(s) ds \right) - R_1^n \left(\gamma_1^n \int_0^t Q_1^n(s) ds \right), \\ X_2^n(t) &= X_2^n(0) + A_2^n(\lambda_2^n t) - S_{22}^n \left(\mu_{22}^n \int_0^t Z_{22}^n(s) ds \right) - R_2^n \left(\gamma_2^n \int_0^t Q_2^n(s) ds \right). \end{aligned} \quad (35)$$

Recall that the processes A_i^n , S_{ij}^n , and R_i^n are all rate-1 Poisson processes and mutually independent, and independent of the initial quantities $X_i^n(0)$.

By (8) and (35), we can write $\hat{X}_1^n(t)$ and $\hat{X}_2^n(t)$ as

$$\begin{aligned} \hat{X}_1^n(t) &= \hat{X}_1^n(0) + l_1^n t - \mu_{11}^n \int_0^t \hat{Z}_{11}^n(s) ds - \mu_{12}^n \int_0^t \hat{Z}_{12}^n(s) ds - \gamma_1^n \int_0^t \hat{Q}_1^n(s) ds \\ &\quad + \hat{M}_{A,1}^n(t) - \hat{M}_{S,11}^n(t) - \hat{M}_{S,12}^n(t) - \hat{M}_{R,1}^n(t), \end{aligned} \quad (36)$$

$$\hat{X}_2^n(t) = \hat{X}_2^n(0) + l_2^n t - \mu_{22}^n \int_0^t \hat{Z}_{22}^n(s) ds - \gamma_2^n \int_0^t \hat{Q}_2^n(s) ds + \hat{M}_{A,2}^n(t) - \hat{M}_{S,22}^n(t) - \hat{M}_{R,2}^n(t), \quad (37)$$

where for $i = 1, 2$, and $j = 1, 2$,

$$\begin{aligned} \hat{M}_{A,i}^n(t) &:= \frac{1}{\sqrt{n}} (A_i^n(\lambda_i^n t) - \lambda_i^n t), \\ \hat{M}_{S,ij}^n(t) &:= \frac{1}{\sqrt{n}} \left(S_{ij}^n \left(\mu_{ij}^n \int_0^t Z_{ij}^n(s) ds \right) - \mu_{ij}^n \int_0^t Z_{ij}^n(s) ds \right), \\ \hat{M}_{R,i}^n(t) &:= \frac{1}{\sqrt{n}} \left(R_i^n \left(\gamma_i^n \int_0^t Q_i^n(s) ds \right) - \gamma_i^n \int_0^t Q_i^n(s) ds \right), \end{aligned}$$

and $l^n = (l_1^n, l_2^n)^\top$ is defined by

$$l_1^n := \frac{1}{\sqrt{n}} (\lambda_1^n - \mu_{11}^n z_{11}^* n - \mu_{12}^n z_{12}^* n), \quad l_2^n := \frac{1}{\sqrt{n}} (\lambda_2^n - \mu_{22}^n z_{22}^* n),$$

with z_{ij}^* as in (4). It is easy to see that under the assumptions on the parameters in Assumption 2.1, $l^n \rightarrow l$ as $n \rightarrow \infty$, where l is defined in (22). The processes $\hat{M}_{A,i}^n := \{\hat{M}_{A,i}^n(t): t \geq 0\}$, $\hat{M}_{S,ij}^n := \{\hat{M}_{S,ij}^n(t): t \geq 0\}$, and $\hat{M}_{R,i}^n := \{\hat{M}_{R,i}^n(t): t \geq 0\}$ are square integrable martingales w.r.t. the filtration $\mathbf{F}^n$ with quadratic variations

$$\langle \hat{M}_{A,i}^n \rangle(t) := \frac{\lambda_i^n}{n} t, \quad \langle \hat{M}_{S,ij}^n \rangle(t) := \frac{\mu_{ij}^n}{n} \int_0^t Z_{ij}^n(s) ds, \quad \langle \hat{M}_{R,i}^n \rangle(t) := \frac{\gamma_i^n}{n} \int_0^t Q_i^n(s) ds.$$

By (2)–(4), (5), and (8), we obtain the balance equations

$$\begin{aligned} \hat{X}_1^n(t) &= \hat{Q}_1^n(t) + \hat{Z}_{11}^n(t) + \hat{Z}_{12}^n(t), & \hat{Y}_1^n(t) + \hat{Z}_{11}^n(t) &= 0, \\ \hat{X}_2^n(t) &= \hat{Q}_2^n(t) + \hat{Z}_{22}^n(t), & \hat{Y}_2^n(t) + \hat{Z}_{12}^n(t) + \hat{Z}_{22}^n(t) &= 0 \end{aligned} \quad (38)$$

for all $t \geq 0$. The work conservation and JWC conditions translate to the following:

$$\hat{Q}_1^n(t) \wedge \hat{Y}_j^n(t) = 0 \quad \forall j = 1, 2 \quad \text{and} \quad \hat{Q}_2^n(t) \wedge \hat{Y}_2^n(t) = 0, \quad \forall t \geq 0,$$

and $e \cdot \hat{Q}^n(t) \wedge e \cdot \hat{Y}^n(t) = 0$, $t \geq 0$, respectively.

6.2. Control Parameterization

By (38), we obtain

$$e \cdot \hat{X}^n(t) = e \cdot \hat{Q}^n(t) - e \cdot \hat{Y}^n(t), \quad (39)$$

and therefore the JWC condition is equivalent to

$$e \cdot \hat{Q}^n(t) = (e \cdot \hat{X}^n(t))^+ \quad e \cdot \hat{Y}^n(t) = (e \cdot \hat{X}^n(t))^- \quad (40)$$

Definition 6.1. We define the processes $U^{c,n} := (U_1^{c,n}, U_2^{c,n})^\top$, and $U^{s,n} := (U_1^{s,n}, U_2^{s,n})^\top$, $t \geq 0$ by

$$U^{c,n}(t) := \begin{cases} \frac{\hat{Q}^n(t)}{e \cdot \hat{Q}^n(t)} & \text{if } e \cdot \hat{Q}^n(t) > 0, \\ e_1 = (1, 0) & \text{otherwise,} \end{cases} \quad (41)$$

and

$$U^{s,n}(t) := \begin{cases} \frac{\hat{Y}^n(t)}{e \cdot \hat{Y}^n(t)} & \text{if } e \cdot \hat{Y}^n(t) > 0, \\ e_2 = (0, 1) & \text{otherwise,} \end{cases} \quad (42)$$

and let $U^n := (U^{c,n}, U^{s,n})$.

The process $U_i^{c,n}(t)$ represents the proportion of the total queue length in the network at queue i at time t , while $U_j^{s,n}(t)$ represents the proportion of the total idle servers in the network at station j at time t . The control $U^{c,n}(t) = e_1 = (1, 0)$ means that server pool 2 gives strict static priority to class 2 jobs, while the control $U^{s,n}(t) = e_2 = (0, 1)$ means that class 1 jobs strictly prefer service in pool 1.

Given $Z^n \in \mathfrak{Z}^n$, the process U^n is uniquely determined via (38) and (41)–(42) and lives in the set $\mathbb{U}$ in (19). It follows by (38) and (40) that, under JWC, we have that for each $t \geq 0$,

$$\hat{Q}^n(t) = (e \cdot \hat{X}^n(t))^+ U^{c,n}(t), \quad \hat{Y}^n(t) = (e \cdot \hat{X}^n(t))^- U^{s,n}(t). \quad (43)$$

Also, by (43), under the JWC condition, we have

$$\hat{Z}^n = \begin{bmatrix} -(e \cdot \hat{X}^n)^- U_1^{s,n} & \hat{X}_1^n - (e \cdot \hat{X}^n)^+ U_1^{c,n} + (e \cdot \hat{X}^n)^- U_1^{s,n} \\ 0 & \hat{X}_2^n - (e \cdot \hat{X}^n)^+ U_2^{c,n} \end{bmatrix}. \quad (44)$$

7. Convergence of Mean Empirical Measures

For the process X^n under a scheduling policy Z^n , and with U^n as in Definition 6.1, we define the mean empirical measures

$$\Phi_T^{Z^n}(A \times B) := \frac{1}{T} \mathbb{E}^{Z^n} \left[\int_0^T \mathbb{I}_{A \times B}(\hat{X}^n(t), U^n(t)) dt \right] \quad (45)$$

for Borel sets $A \subset \mathbb{R}^2$ and $B \subset \mathbb{U}$. Recall Definition 2.1. The lemma that follows provides a sufficient condition under which the mean empirical measures $\Phi_T^{Z^n}$ are tight and converge to an ergodic occupation measure corresponding to some stationary stable Markov control for the limiting diffusion control problem. The condition simply requires a finite long-run average first-order moment of the diffusion-scaled state process under an EJWC scheduling policy. This lemma is used in Section 8 to prove the lower bounds in Theorem 5.1.

Lemma 7.1. Suppose that under some sequence $\{Z^n, n \in \mathbb{N}\} \subset \mathfrak{Z}$, we have

$$\sup_n \limsup_{T \rightarrow \infty} \frac{1}{T} \mathbb{E}^{Z^n} \left[\int_0^T |\hat{X}^n(s)| ds \right] < \infty. \quad (46)$$

Then, any limit point $\pi \in \mathcal{P}(\mathbb{R}^2 \times \mathbb{U})$ of $\Phi_T^{Z^n}$ defined in (45) as $(n, T) \rightarrow \infty$ satisfies $\pi \in \mathcal{G}$.

Proof. Let $f \in \mathcal{C}_c^\infty(\mathbb{R}^2)$, and define

$$\mathcal{D}f(\hat{X}^n, s) := \Delta f(\hat{X}^n(s)) - \sum_{i=1}^2 \partial_i f(\hat{X}^n(s-)) \Delta \hat{X}_i^n(s) - \frac{1}{2} \sum_{i,i'=1}^2 \partial_{ii'} f(\hat{X}^n(s-)) \Delta \hat{X}_i^n(s) \Delta \hat{X}_{i'}^n(s). \quad (47)$$

By applying Itô's formula (see, e.g., theorem 26.7 in Kallenberg [21]) and using the definition of $\Phi_T^{Z^n}$ in (45), and $\hat{X}^n$ in (36)–(37), we obtain

$$\frac{\mathbb{E}[f(\hat{X}^n(T))] - \mathbb{E}[f(\hat{X}^n(0))]}{T} = \int_{\mathbb{R}^2 \times \mathbb{U}} \mathcal{A} f(\hat{x}, u) \Phi_T^{Z^n}(d\hat{x}, du) + \frac{1}{T} \mathbb{E} \left[\sum_{s \leq T} \mathcal{D}f(\hat{X}^n, s) \right] \quad (48)$$

with $\mathbb{E} = \mathbb{E}^{Z^n}$. Define

$$\begin{aligned}\mathcal{A}_{1,1}^n(\hat{x}, u) &:= -\mu_{12}^n(\hat{x}_1 - (e \cdot \hat{x})^+ u_1^c) + (\mu_{11}^n - \mu_{12}^n)(e \cdot \hat{x})^- u_1^s - \gamma_1^n(e \cdot \hat{x})^+ u_1^c + l_1^n, \\ \mathcal{A}_{2,1}^n(\hat{x}, u) &:= -\mu_{22}^n(\hat{x}_2 - (e \cdot \hat{x})^+ u_2^c) - \gamma_2^n(e \cdot \hat{x})^+ u_2^c + l_2^n, \\ \mathcal{A}_{1,2}^n(\hat{x}, u) &:= \frac{1}{2} \left(\frac{\lambda_1^n}{n} + \mu_{11}^n z_{11}^* + \mu_{12}^n z_{12}^* + \frac{1}{\sqrt{n}} \mu_{12}^n(\hat{x}_1 - (e \cdot \hat{x})^+ u_1^c) \right. \\ &\quad \left. + \frac{1}{\sqrt{n}} (\mu_{11}^n - \mu_{12}^n)(e \cdot \hat{x})^- u_1^s + \frac{1}{\sqrt{n}} \gamma_1^n(e \cdot \hat{x})^+ u_1^c \right), \\ \mathcal{A}_{2,2}^n(\hat{x}, u) &:= \frac{1}{2} \left(\frac{\lambda_2^n}{n} + \mu_{22}^n z_{22}^* + \frac{1}{\sqrt{n}} \mu_{22}^n(\hat{x}_2 - (e \cdot \hat{x})^+ u_2^c) + \frac{1}{\sqrt{n}} \gamma_2^n(e \cdot \hat{x})^+ u_2^c \right).\end{aligned}\tag{49}$$

Since $Z^n \in \mathfrak{Z}^n$, the operator $\mathcal{A}^n: \mathcal{C}_c^\infty(\sqrt{n}\tilde{B}) \rightarrow \mathcal{C}_c^\infty(\sqrt{n}\tilde{B} \times \mathbb{U})$ takes the form

$$\mathcal{A}^n f(\hat{x}, u) := \sum_{i=1}^2 (\mathcal{A}_{i,1}^n(\hat{x}, u) \partial_i f(\hat{x}) + \mathcal{A}_{i,2}^n(\hat{x}, u) \partial_{ii} f(\hat{x})).$$

Let

$$\|f\|_{\mathcal{C}^3} := \sup_{x \in \mathbb{R}^2} \left(|f(x)| + \sum_{i=1,2} |\partial_i f(x)| + \sum_{i,j=1}^2 |\partial_{ij} f(x)| + \sum_{i,j,k=1}^2 |\partial_{ijk} f(x)| \right).$$

By Taylor's formula, using also the fact that the jump size is $1/\sqrt{n}$, we obtain

$$|\mathcal{D}f(\hat{X}^n, s)| \leq \kappa \|f\|_{\mathcal{C}^3} \sum_{i,j,k=1}^2 |\Delta \hat{X}_i^n(s)| |\Delta \hat{X}_j^n(s)| |\Delta \hat{X}_k^n(s)| \leq \frac{\kappa' \|f\|_{\mathcal{C}^3}}{\sqrt{n}} \sum_{i,i'=1}^2 |\Delta \hat{X}_i^n(s) \Delta \hat{X}_{i'}^n(s)|,$$

for some constants κ and κ' that do not depend on $n \in \mathbb{N}$. Let

$$\begin{aligned}\tilde{\mathcal{X}}_1^n(t) &:= \frac{\lambda_1^n}{n} + \frac{1}{n} \mu_{11}^n Z_{11}^n(t) + \frac{1}{n} \mu_{12}^n Z_{12}^n(t) + \frac{1}{n} \gamma_1^n Q_1^n(t), \\ \tilde{\mathcal{X}}_2^n(t) &:= \frac{\lambda_2^n}{n} + \frac{1}{n} \mu_{22}^n Z_{22}^n(t) + \frac{1}{n} \gamma_2^n Q_2^n(t)\end{aligned}\tag{50}$$

for $t \geq 0$. Since independent Poisson processes have no simultaneous jumps w.p.1., we have

$$\frac{1}{T} \mathbb{E} \left[\int_0^T \sum_{i,i'=1}^2 |\Delta \hat{X}_i^n(s) \Delta \hat{X}_{i'}^n(s)| ds \right] \leq \frac{1}{T} \mathbb{E} \left[\int_0^T (\tilde{\mathcal{X}}_1^n(s) + \tilde{\mathcal{X}}_2^n(s)) ds \right],$$

and that the right-hand side is uniformly bounded over $n \in \mathbb{N}$, and $T > 0$ by (46). Thus we have

$$\frac{1}{T} \exp \left[\sum_{s \leq T} |\mathcal{D}f(\hat{X}^n, s)| \right] \leq \frac{\kappa' \|f\|_{\mathcal{C}^3}}{T \sqrt{n}} \mathbb{E} \left[\int_0^T \sum_{i,i'=1}^2 |\Delta \hat{X}_i^n(s) \Delta \hat{X}_{i'}^n(s)| ds \right] \rightarrow 0$$

as $(n, T) \rightarrow \infty$. Therefore, taking limits in (48), we obtain

$$\limsup_{(n,T) \rightarrow \infty} \int_{\mathbb{R}^2 \times \mathbb{U}} \mathcal{A}^n f(\hat{x}, u) \Phi_T^{Z^n}(\mathrm{d}\hat{x}, \mathrm{d}u) = 0.$$

Note that for $i = 1, 2$, $\mathcal{A}_{i,1}^n$ tends to the drift of the limiting diffusion b_i , while $\mathcal{A}_{i,2}^n$ tends to λ_i as $n \rightarrow \infty$, uniformly over compact sets in $\mathbb{R}^2 \times \mathbb{U}$.

Let (n_k, T_k) be any sequence along which $\Phi_T^{Z^n}$ converges to some $\pi \in \mathcal{P}(\mathbb{R}^2 \times \mathbb{U})$. Let

$$\mathcal{L}^u f(x) = \sum_{i=1}^2 [\lambda_i \partial_{ii} f(x) + b_i(x, u) \partial_i f(x)].$$

We have

$$\begin{aligned}\int_{\mathbb{R}^2 \times \mathbb{U}} \mathcal{L}^u f(x) \pi(\mathrm{d}x, \mathrm{d}u) - \int_{\mathbb{R}^2 \times \mathbb{U}} \mathcal{A}^n f(\hat{x}, u) \Phi_T^{Z^n}(\mathrm{d}\hat{x}, \mathrm{d}u) &= \int_{\mathbb{R}^2 \times \mathbb{U}} \mathcal{L}^u f(x) (\pi(\mathrm{d}x, \mathrm{d}u) - \Phi_T^{Z^n}(\mathrm{d}x, \mathrm{d}u)) \\ &\quad + \int_{\mathbb{R}^2 \times \mathbb{U}} (\mathcal{L}^u f(\hat{x}) - \mathcal{A}^n f(\hat{x}, u)) \Phi_T^{Z^n}(\mathrm{d}\hat{x}, \mathrm{d}u).\end{aligned}\tag{51}$$

The first term on the right-hand side of (51) converges to 0 as $n \rightarrow \infty$ by the convergence of $\Phi_T^{Z^n}$ to π , while the second term also converges to 0 by the uniform convergence of $\mathcal{L}^u f$ to $\mathcal{A}^n f$ on compact subsets of $\mathbb{R}^2 \times \mathbb{U}$ and the tightness of $\Phi_T^{Z^n}$. Thus we obtain

$$\int_{\mathbb{R}^2 \times \mathbb{U}} \mathcal{L}^u f(x) \pi(dx, du) = 0.$$

This completes the proof. $\square$

Before stating the second lemma, we first introduce a canonical construction of scheduling policies from the optimal control $v \in \mathcal{U}_{\text{SSM}}$ for the diffusion control problems. Recall the notation in Definition 3.2.

Definition 7.1. Let $\omega: \{x \in \mathbb{R}_+^2: e \cdot x \in \mathbb{Z}\} \rightarrow \mathbb{Z}_+^2$ be a measurable map defined by

$$\omega(x) := (\lfloor x_1 \rfloor, e \cdot x - \lfloor x_1 \rfloor), \quad x \in \mathbb{R}^2.$$

For any precise control $v \in \mathcal{U}_{\text{SSM}}$, define the maps $q^n[v]$ and $y^n[v]$ by

$$q^n[v](\hat{x}) := \omega((e \cdot (\sqrt{n}\hat{x} + nx^*))^+ v^c(\hat{x})) \quad y^n[v](\hat{x}) := \omega((e \cdot (\sqrt{n}\hat{x} + nx^*))^- v^s(\hat{x}))$$

for $\hat{x} \in \mathcal{S}^n$. We also define the map (Markov scheduling policy) $z^n[v]$ on $\check{\mathcal{S}}^n$ by

$$z^n[v](\hat{x}) := \begin{bmatrix} N_1^n - y_1^n[v](\hat{x}) & x_1 - q_1^n[v](\hat{x}) - (N_1^n - y_1^n[v](\hat{x})) \\ 0 & x_2 - q_2^n[v](\hat{x}) \end{bmatrix}, \quad \hat{x} \in \check{\mathcal{S}}^n.$$

Compare this to (7).

Corollary 7.1. For any precise control $v \in \mathcal{U}_{\text{SSM}}$, we have

$$e \cdot q^n[v](\hat{x}^n(x)) \wedge e \cdot y^n[v](\hat{x}^n(x)) = 0, \quad \text{and} \quad z^n[v](\hat{x}^n(x)) \in \mathcal{Z}^n(x)$$

for all $x \in \check{\mathcal{X}}^n$, i.e., the JWC condition is satisfied for $x \in \check{\mathcal{X}}^n$.

Proof. This follows from Lemma 2.1 and the definition of the maps $q^n[v]$, $y^n[v]$, and $z^n[v]$. $\square$

The lemma that follows asserts that, if a sequence of EJWC scheduling policies is constructed using any precise stationary stable Markov control in a way that the long-run average moment condition in Lemma 7.1 is satisfied, then any limit of the mean empirical measures of the diffusion-scaled processes agrees with the ergodic occupation measure of the limiting diffusion corresponding to that control. This lemma is used in the proof of upper bounds in Theorem 5.2. Recall Definition 2.1.

Lemma 7.2. Let $v \in \mathcal{U}_{\text{SSM}}$ be a continuous precise control, and $\{Z^n: n \in \mathbb{N}\}$ be any sequence of admissible scheduling policies such that each Z^n agrees with the Markov scheduling policy $z^n[v]$ given in Definition 7.1 on $\sqrt{n}\check{B}$, i.e., $Z^n(t) = z^n[v](\hat{X}^n(t))$ whenever $\hat{X}^n(t) \in \sqrt{n}\check{B}$. For $\hat{x} \in \sqrt{n}\check{B} \cap \mathcal{S}^n$, we define

$$u^{c,n}[v](\hat{x}) := \begin{cases} \frac{q^n[v](\hat{x})}{e \cdot q^n[v](\hat{x})} & \text{if } e \cdot q^n[v](\hat{x}) > 0, \\ v^c(\hat{x}) & \text{otherwise,} \end{cases}$$

and

$$u^{s,n}[v](\hat{x}) := \begin{cases} \frac{y^n[v](\hat{x})}{e \cdot y^n[v](\hat{x})} & \text{if } e \cdot y^n[v](\hat{x}) > 0, \\ v^s(\hat{x}) & \text{otherwise.} \end{cases}$$

For the process X^n under the scheduling policy Z^n , define the mean empirical measures

$$\check{\Phi}_T^{Z^n}(A \times B) := \frac{1}{T} \mathbb{E}^{Z^n} \left[\int_0^T \mathbb{I}_{A \times B}(\hat{X}^n(t), u^n[v](\hat{X}^n(t))) dt \right] \quad (52)$$

for Borel sets $A \subset \sqrt{n}\check{B}$, and $B \subset \mathbb{U}$. Suppose that (46) holds under this sequence $\{Z^n\}$. Then, the ergodic occupation measure π_v of the controlled diffusion in (18) corresponding to v is the unique limit point in $\mathcal{P}(\mathbb{R}^d \times \mathbb{U})$ of $\check{\Phi}_T^{Z^n}$ as $(n, T) \rightarrow \infty$.

Proof. It follows by Corollary 7.1 that $\{Z^n\} \in \mathfrak{Z}$. Also, by the continuity of v , we have

$$\sup_{\hat{x} \in \mathcal{S}^n \cap K} |u^n[v](\hat{x}) - v(\hat{x})| \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (53)$$

for any compact set $K \subset \mathbb{R}^2$. Also, for any $f \in \mathcal{C}_c^\infty(\mathbb{R}^2 \times \mathbb{U})$, it holds that

$$\int_{\mathbb{R}^2 \times \mathbb{U}} f(\hat{x}, u) \tilde{\Phi}_T^{Z^n}(\mathrm{d}\hat{x}, \mathrm{d}u) = \frac{1}{T} \mathbb{E}^{Z^n} \left[\int_0^T f(\hat{X}^n(t), u^n[v](\hat{X}^n(t))) \mathrm{d}t \right],$$

for all sufficiently large n such that the support of f is contained in $\sqrt{n}\tilde{B}$. Therefore, if π^n is any limit point of $\tilde{\Phi}_T^{Z^n}$ as $T \rightarrow \infty$, and we disintegrate π^n as

$$\pi^n(\mathrm{d}\hat{x}, \mathrm{d}u) = \nu^n(\mathrm{d}\hat{x}) \xi^n(\mathrm{d}u \mid \hat{x}), \quad (54)$$

Then, we have

$$\int_{\mathbb{R}^2 \times \mathbb{U}} \mathcal{A}^n f(\hat{x}, u) \pi^n(\mathrm{d}\hat{x}, \mathrm{d}u) = \int_{\mathbb{R}^2} \mathcal{A}^n f(\hat{x}, u^n[v](\hat{x})) \nu^n(\mathrm{d}\hat{x}).$$

By Lemma 7.1, the sequence $\{\nu^n\}$ is tight. Let $\{n\} \in \mathbb{N}$ be any increasing sequence such that $\nu^n \rightarrow \nu \in \mathcal{P}(\mathbb{R}^2)$. To simplify the notation, let $\tilde{\mathcal{A}}^n f(\hat{x}) := \mathcal{A}^n f(\hat{x}, u^n[v](\hat{x}))$. We have

$$\int_{\mathbb{R}^2} \tilde{\mathcal{A}}^n f \mathrm{d}\nu^n - \int_{\mathbb{R}^2} \mathcal{L}^v f \mathrm{d}\nu = \int_{\mathbb{R}^2} (\tilde{\mathcal{A}}^n f - \mathcal{L}^v f) \mathrm{d}\nu^n + \int_{\mathbb{R}^2} \mathcal{L}^v f (\mathrm{d}\nu^n - \mathrm{d}\nu). \quad (55)$$

It follows by (53) that $\tilde{\mathcal{A}}^n f - \mathcal{L}^v f \rightarrow 0$, uniformly as $n \rightarrow \infty$, which implies that the first term on the right-hand side of (55) converges to 0. The second term does the same by the convergence of ν^n to ν . By Lemma 7.1, we have $\int_{\mathbb{R}^2} \tilde{\mathcal{A}}^n f \mathrm{d}\nu^n \rightarrow 0$ as $n \rightarrow \infty$. Therefore we obtain

$$\int_{\mathbb{R}^2} \mathcal{L}^v f(x) \nu(\mathrm{d}x) = 0,$$

and this means that ν is an invariant probability measure for the diffusion associated with the control v . Next, note that, the Markov control ξ^n in (54) agrees with $u^n[v](\hat{x})$ when $\hat{x} \in \sqrt{n}\tilde{B} \cap \mathcal{S}^n$ by definition. In other words, $\xi^n(\mathrm{d}u \mid \hat{x}) = \delta_{u^n[v](\hat{x})}(u)$, where δ denotes the Dirac measure. It then follows by (53) that ξ^n converges to v as $n \rightarrow \infty$ in the topology of Markov controls (Arapostathis et al. [2, section 2.4]). The ergodic occupation measure $\pi_v \in \mathcal{P}(\mathbb{R}^2 \times \mathbb{U})$ is given by $\pi_v(\mathrm{d}x, \mathrm{d}u) := \nu(\mathrm{d}x, \mathrm{d}u) \delta_{v(x)}(u)$. With $g \in \mathcal{C}_c(\mathbb{R}^2 \times \mathbb{U})$, i.e., a continuous function with compact support, we write

$$\begin{aligned} \left| \int_{\mathbb{R}^2 \times \mathbb{U}} g(x, u) (\pi_v(\mathrm{d}x, \mathrm{d}u) - \pi^n(\mathrm{d}x, \mathrm{d}u)) \right| &\leq \left| \int_{\mathbb{U}} \left(\int_{\mathbb{R}^2} g(x, u) (\nu(\mathrm{d}x) - \nu^n(\mathrm{d}x)) \right) \xi^n(\mathrm{d}u \mid x) \right| \\ &\quad + \left| \int_{\mathbb{U}} \left(\int_{\mathbb{R}^2} g(x, u) \nu(\mathrm{d}x) \right) (\xi^n(\mathrm{d}u \mid x) - \delta_{v(x)}(u)) \right|. \end{aligned} \quad (56)$$

The first term on the right-hand side of (56) converges to 0 as $n \rightarrow \infty$ by the convergence of $\nu^n \rightarrow \nu$ in $\mathcal{P}(\mathbb{R}^2)$. Since ν has a continuous density, the second term also converges to 0 as $n \rightarrow \infty$ by Arapostathis et al. [2, lemma 2.4.1]. Therefore (56) shows that $\pi^n \rightarrow \pi_v$ in $\mathcal{P}(\mathbb{R}^2 \times \mathbb{U})$, and this completes the proof. $\square$

8. Proof of the Lower Bounds

In this section, we prove the lower bounds in Theorem 5.1. The following lemma, which applies to the diffusion-scaled process, is analogous to lemma 3.1 (c) for the diffusion limit in Arapostathis and Pang [1].

Lemma 8.1. *There exist constants C_1 and C_2 independent of n such that*

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \mathbb{E}^{Z^n} \left[\int_0^T |\hat{X}^n(s)|^m \mathrm{d}s \right] \leq C_1 + C_2 J_0(\hat{X}^n(0), \hat{Z}^n), \quad \forall n \in \mathbb{N} \quad (57)$$

for any sequence $\{Z^n \in \mathfrak{Z}^n, n \in \mathbb{N}\}$, where $m \geq 1$ is as in (9).

Proof. Let $\mathcal{V}(x) := \mathcal{V}_1(x_1) + \beta \mathcal{V}_2(x_2)$, $x \in \mathbb{R}^2$, where β is a positive constant to be determined later, and $\mathcal{V}_i(x) = |x_i|^{m+1}/\sqrt{1+|x_i|^2}$ for $m \geq 1$. By applying Itô's formula on $\mathcal{V}$ with $\mathbb{E} = \mathbb{E}^{Z^n}$, we obtain from (36) that, for $t \geq 0$,

$$\mathbb{E}[\mathcal{V}(\hat{X}^n(t))] = \mathbb{E}[\mathcal{V}(\hat{X}^n(0))] + \mathbb{E}\left[\int_0^t \mathcal{A}^n \mathcal{V}(\hat{X}^n(s), \hat{Z}^n(s)) ds\right] + \mathbb{E}\left[\sum_{s \leq t} \mathcal{D}\mathcal{V}(\hat{X}^n, s)\right], \quad (58)$$

where $\mathcal{D}\mathcal{V}(\hat{X}^n, s)$ is defined as in (47),

$$\mathcal{A}^n \mathcal{V}(\hat{x}, \hat{z}) := \sum_{i=1}^2 (\mathcal{A}_{i,1}^n(\hat{x}, \hat{z}) \partial_i \mathcal{V}(\hat{x}) + \mathcal{A}_{i,2}^n(\hat{x}, \hat{z}) \partial_{ii} \mathcal{V}(\hat{x})),$$

and

$$\begin{aligned} \mathcal{A}_{1,1}^n(\hat{x}, \hat{z}) &:= l_1^n - \mu_{11}^n \hat{z}_{11} - \mu_{12}^n \hat{z}_{12} - \gamma_1^n (\hat{x}_1 - \hat{z}_{11} - \hat{z}_{12}), \\ \mathcal{A}_{2,1}^n(\hat{x}, \hat{z}) &:= l_2^n - \mu_{22}^n \hat{z}_{22} - \gamma_2^n (\hat{x}_2 - \hat{z}_{22}), \\ \mathcal{A}_{1,2}^n(\hat{x}, \hat{z}) &:= \frac{1}{2} \left[\frac{\lambda_1^n}{n} + (\mu_{11}^n z_{11}^* + \mu_{12}^n z_{12}^*) + \frac{1}{\sqrt{n}} (\mu_{11}^n \hat{z}_{11} + \mu_{12}^n \hat{z}_{12}) + \frac{\gamma_1^n}{\sqrt{n}} (\hat{x}_1 - \hat{z}_{11} - \hat{z}_{12}) \right], \\ \mathcal{A}_{2,2}^n(\hat{x}, \hat{z}) &:= \frac{1}{2} \left[\frac{\lambda_2^n}{n} + \mu_{22}^n z_{22}^* + \frac{1}{\sqrt{n}} \mu_{22}^n \hat{z}_{22} + \frac{\gamma_2^n}{\sqrt{n}} (\hat{x}_2 - \hat{z}_{22}) \right], \end{aligned}$$

for $\hat{x} \in \mathcal{S}^n$, and $\hat{z}_{ij} := (1/\sqrt{n})(z_{ij} - n z_{ij}^*)$ for $z_{ij} \in \mathbb{Z}_+$, and z^* defined in (4). We also use the nonnegative variables $\hat{q}_i$ and $\hat{y}_i$, $i = 1, 2$, which are defined as functions of $\hat{x}$ and $\hat{z}$ via the balance Equation (38), keeping in mind that the work conservation condition holds for these.

Define

$$\begin{aligned} \bar{\mathcal{A}}_{1,1}(x, z) &:= l_1 - \mu_{11} z_{11} - \mu_{12} z_{12} - \gamma_1 (x_1 - z_{11} - z_{12}), \\ \bar{\mathcal{A}}_{2,1}(x, z) &:= l_2 - \mu_{22} z_{22} - \gamma_2 (x_2 - z_{22}) \end{aligned}$$

for $x \in \mathbb{R}^2$ and $z \in \mathbb{R}^{2 \times 2}$.

By the convergence of the parameters in Assumption 2.1, we have that, for $i = 1, 2$,

$$|\bar{\mathcal{A}}_{i,1}(\hat{x}, \hat{z}) - \mathcal{A}_{i,1}^n(\hat{x}, \hat{z})| \leq \kappa_1(n)(\|\hat{x}\| + \|\hat{z}\|) \quad (59)$$

for some constant $\kappa_1(n) \searrow 0$ as $n \rightarrow \infty$.

Let $\hat{\xi} := (e \cdot \hat{q}) \wedge (e \cdot \hat{y})$. We claim that if $\hat{\xi}^n > 0$, then $\hat{q}_1 = 0$, $\hat{y}_2 = 0$ by the work conservation condition. Indeed, since $\hat{q}_i \wedge \hat{y}_2 = 0$ for $i = 1, 2$, then $\hat{\xi}^n > 0$ implies that $\hat{y}_2 = 0$, which, in turn, implies that $\hat{y}_1 > 0$. This, of course, implies that $\hat{q}_1 = 0$.

If $\hat{\xi} = \hat{y}_1$, then by the balance equations, we have $\hat{z}_{11} = -\hat{\xi}$, $\hat{z}_{12} = \hat{x}_1 + \hat{\xi}$, and $\hat{z}_{22} = \hat{x}_2 - \hat{q}_2$. On the other hand, if $\hat{\xi} = \hat{q}_2$, then we obtain $\hat{z}_{11} = \hat{x}_1 + \hat{x}_2 - \hat{\xi}$, $\hat{z}_{12} = \hat{\xi} - \hat{x}_2$, and $\hat{z}_{22} = \hat{x}_2 - \hat{\xi}$. Hence, when $\hat{\xi} > 0$, we have

$$\begin{aligned} \bar{\mathcal{A}}_{1,1}(\hat{x}, \hat{z}) &= -\mu_{12} \hat{x}_1 + (\mu_{11} - \mu_{12}) \hat{\xi} + l_1 & \text{if } \hat{y}_1 < \hat{q}_2, \\ \bar{\mathcal{A}}_{2,1}(\hat{x}, \hat{z}) &= -\mu_{22}(\hat{x}_2 - \hat{q}_2) - \gamma_2 \hat{q}_2 + l_2 \\ \bar{\mathcal{A}}_{1,1}(\hat{x}, \hat{z}) &= -\mu_{11} \hat{x}_1 + (\mu_{11} - \mu_{12})(\hat{\xi} - \hat{x}_2) + l_1 & \text{if } \hat{y}_1 \geq \hat{q}_2, \\ \bar{\mathcal{A}}_{2,1}(\hat{x}, \hat{z}) &= -\mu_{22}(\hat{x}_2 - \hat{\xi}) - \gamma_2 \hat{\xi} + l_2 \end{aligned} \quad (60)$$

and when $\hat{\xi} = 0$, we can use the parameterization $\hat{q} = (e \cdot \hat{x})^+ u^c$ and $\hat{y} = (e \cdot \hat{x})^- u^s$ and (49) to obtain

$$\begin{aligned} \bar{\mathcal{A}}_{1,1}(\hat{x}, \hat{z}) &= -\mu_{12} \hat{x}_1 + (\mu_{12} - \gamma_1) \hat{q}_1 + l_1 & \text{if } (e \cdot \hat{x})^+ > 0, \\ \bar{\mathcal{A}}_{2,1}(\hat{x}, \hat{z}) &= -\mu_{22}(\hat{x}_2 - \hat{q}_2) - \gamma_2 \hat{q}_2 + l_2 \\ \bar{\mathcal{A}}_{1,1}(\hat{x}, \hat{z}) &= -(\mu_{12}(1 - u_1^s) + \mu_{11} u_1^s) \hat{x}_1 - (\mu_{11} - \mu_{12}) \hat{x}_2 u_1^s + l_1 & \text{if } (e \cdot \hat{x})^- \geq 0. \\ \bar{\mathcal{A}}_{2,1}(\hat{x}, \hat{z}) &= -\mu_{22} \hat{x}_2 + l_2 \end{aligned} \quad (61)$$

It follows by the above analysis that

$$|z_{ij}| \in \mathcal{O}(|x| + |q|), \quad i, j \in \{1, 2\}. \quad (62)$$

Hence we have

$$\mathcal{A}_{i,2}^n(\hat{x}, \hat{z}) \in \mathcal{O}(1 + n^{-1/2}|\hat{x}|). \quad (63)$$

Following the steps in the proof of Lemma 4.1, and also using the fact that $\hat{\xi} \leq e \cdot \hat{q}$ and Young's inequality, it follows by (60)–(61) that we can choose $\beta > 0$ and positive constants c_1 and c_2 such that

$$\sum_{i=1}^2 \bar{\mathcal{A}}_{i,1}^n(\hat{x}, \hat{z}) \partial_i \mathcal{V}(\hat{x}) \leq -c_1 \mathcal{V}(\hat{x}) + c_2(1 + |\hat{q}|^m). \quad (64)$$

Thus, by (59), (63), and (64), we obtain

$$\mathcal{A}^n \mathcal{V}(\hat{x}, \hat{z}) \leq -c'_1 \mathcal{V}(\hat{x}) + c'_2(1 + |\hat{q}|^m) \quad (65)$$

for some positive constants c'_1 and c'_2 .

For the jumps in (58), we first note that by the definition of $\mathcal{V}_i$, since there exists a positive constant c_3 such that

$$\sup_{|x'_i - x_i| \leq 1} |\mathcal{V}_i''(x'_i)| \leq c_3(1 + |x_i|^{m-2}), \quad \forall x_i \in \mathbb{R}.$$

Since also the jump size is of order $1/\sqrt{n}$, then by Taylor's expansion, we obtain

$$\Delta \mathcal{V}_i(\hat{X}^n(s)) - \mathcal{V}_i'(\hat{X}^n(s-)) \cdot \Delta \hat{X}_i^n(s) \leq \frac{1}{2} \sup_{|x'_i - \hat{X}_i^n(s-)| \leq 1} |\mathcal{V}_i''(x'_i)| (\Delta \hat{X}_i^n(s))^2,$$

for $i = 1, 2$. Recall the definitions of $\bar{\mathcal{X}}_1^n$ and $\bar{\mathcal{X}}_2^n$ in (50). Thus, for $i = 1, 2$, using also (62), we obtain

$$\begin{aligned} \mathbb{E} \left[\sum_{s \leq t} \mathcal{D} \mathcal{V}_i(\hat{X}^n, s) \right] &\leq \mathbb{E} \left[\sum_{s \leq t} c_3(1 + |\hat{X}_i^n(s-)|^{m-1}) (\Delta \hat{X}_i^n(s))^2 \right] \leq c_3 \mathbb{E} \left[\int_0^t (1 + |\hat{X}_i^n(s)|^{m-1}) \bar{\mathcal{X}}_i^n(s) ds \right] \\ &\leq c_4 \mathbb{E} \left[\int_0^t (1 + |\hat{X}_i^n(s)|^{m-1}) (1 + n^{-1/2}(|\hat{X}^n(s)| + |\hat{Q}^n(s)|)) ds \right] \end{aligned} \quad (66)$$

for some positive constant c_4 . Therefore, by (58), (65), and (66), we can choose positive constants c_5 and c_6 such that

$$\mathbb{E}[\mathcal{V}(\hat{X}^n(t))] \leq \mathbb{E}[\mathcal{V}(\hat{X}^n(0))] + c_6 t - c_5 \mathbb{E} \left[\int_0^t |\hat{X}^n(s)|^m ds \right] + c_6 \mathbb{E} \left[\int_0^t |\hat{Q}^n(s)|^m ds \right].$$

Dividing by t and taking limits as $t \rightarrow \infty$, establishes (57). $\square$

We are now ready to prove Theorem 5.1.

Proof of Theorem 5.1. Let $Z^n \in \mathfrak{Z}^n$, $n \in \mathbb{N}$, be an arbitrary sequence of scheduling policies in $\mathfrak{Z}$, and let $\Phi^n := \Phi^{Z^n}$ as defined in (45). Without loss of generality, we assume that along some increasing sequence $\{n_k\} \subset \mathbb{N}$, we have $\sup_k J(\hat{X}^{n_k}(0), Z^{n_k}) < \infty$; Otherwise, there is nothing to prove. By Lemmas 7.1 and 8.1, the sequence of mean empirical measures $\{\Phi_T^{n_k} : T > 0, k \geq 1\}$ is tight and any subsequential limit as $(n_k, T) \rightarrow \infty$ is in $\mathcal{G}$. Select any subsequence $\{T_k, n'_k\} \subset \mathbb{R}_+ \times \{n_k\}$ with $T_k \rightarrow \infty$ as $k \rightarrow \infty$ and such that

$$J(\hat{X}^{n'_k}(0), Z^{n'_k}) \leq \frac{1}{k} + \liminf_{l \rightarrow \infty} J(\hat{X}^{n'_l}(0), Z^{n'_l}), \quad \text{and} \quad \int_{\mathbb{R}^2 \times \mathbb{U}} r(x, u) \Phi_{T_k}^{n'_k}(dx, du) \leq J(\hat{X}^{n'_k}(0), Z^{n'_k}) + \frac{1}{k}$$

for all $k \in \mathbb{N}$, and extract any further subsequence, also denoted as $\{T_k, n'_k\}$, along which $\Phi_{T_k}^{n'_k} \rightarrow \hat{\pi} \in \mathcal{G}$. Since r is nonnegative, taking limits as $k \rightarrow \infty$, we obtain

$$\liminf_{k \rightarrow \infty} J(\hat{X}^{n_k}(0), Z^{n_k}) \geq \hat{\pi}(r) \geq \varrho^*.$$

This proves part (i).

We next show the lower bound (ii) for the constrained problem. Repeating the same argument as in part (i), suppose that $\sup_k J_0(\hat{X}^{n_k}(0), Z^{n_k}) < \infty$ along some increasing sequence $\{n_k\} \subset \mathbb{N}$. As in the proof of part (i), let $\hat{\pi} \in \mathcal{P}(\mathbb{R}^2 \times \mathbb{U})$ be a limit of Φ_T^n as $(n, T) \rightarrow \infty$. Recall the definition of r_j in (25). Since r_j is bounded below, taking

limits, we obtain $\hat{\pi}(r_j) \leq \delta_j$, $j = 1, 2$. Therefore $\hat{\pi} \in \mathcal{H}(\delta)$, and by optimality, we must have $\hat{\pi}(r_o) \geq \varrho_c^*$. Similarly, we obtain,

$$\liminf_{k \rightarrow \infty} J_o(\hat{X}^{n_k}(0), Z^{n_k}) \geq \hat{\pi}(r_o) \geq \varrho_c^*.$$

This proves part (ii).

The result in part (iii) for the fairness problem follows along the same lines as part (ii). With $\hat{\pi}$ as in part (ii), we have

$$\liminf_{k \rightarrow \infty} J_o(\hat{X}^{n_k}(0), Z^{n_k}) \geq \hat{\pi}(r_o). \quad (67)$$

The uniform integrability of

$$\frac{1}{T} \mathbb{E}^{Z^n} \left[\int_0^T (\hat{Y}_j^n(s))^{\tilde{m}} ds \right], \quad j = 1, 2,$$

which follows by (30) and the assumption that $\tilde{m} < m$, together with (33), imply that

$$(\theta - \epsilon)\hat{\pi}(r_2) \leq \hat{\pi}(r_1) \leq (\theta + \epsilon)\hat{\pi}(r_2).$$

Therefore $\hat{\pi}(r_1) = \tilde{\theta}(\epsilon)\hat{\pi}(r_2)$ for some $\tilde{\theta}(\epsilon)$ satisfying $|\tilde{\theta}(\epsilon) - \theta| \leq \epsilon$. Let

$$\tilde{\varrho} := \inf_{\pi \in \mathcal{H}_1(\tilde{\theta}(\epsilon))} \pi(r_o),$$

and λ^* denote the Lagrange multiplier for the problem in Theorem 4.3. It is clear that $\hat{\pi}(r_o) \geq \tilde{\varrho}$. Writing $\hat{\pi}(r_1) = \tilde{\theta}(\epsilon)\hat{\pi}(r_2)$ as $\hat{\pi}(r_1) - \theta\hat{\pi}(r_2) = (\tilde{\theta}(\epsilon) - \theta)\hat{\pi}(r_2)$, we obtain by Luenberger [22, theorem 1, p. 222] that

$$\varrho_i^* - \tilde{\varrho} \leq |\lambda^*(\tilde{\theta}(\epsilon) - \theta)\hat{\pi}(r_2)| \leq \epsilon |\lambda^*\hat{\pi}(r_2)|. \quad (68)$$

Without loss of generality, we may assume that $\hat{\pi}(r_o) \leq \varrho_i^*$; otherwise, (34) trivially follows by (67). By (30) and Jensen's inequality, we have

$$\hat{\pi}(r_2) \leq \hat{\kappa}(1 + \pi(r_o)^{\tilde{m}/m}) \leq \hat{\kappa}(1 + (\varrho_i^*)^{\tilde{m}/m}) \quad (69)$$

for some constant $\hat{\kappa}$. Therefore combining (68)–(69), we obtain

$$\hat{\pi}(r_o) \geq \tilde{\varrho} \geq \varrho_i^* - \epsilon |\lambda^*| \hat{\kappa}(1 + (\varrho_i^*)^{\tilde{m}/m}),$$

and (34) follows by this estimate and (67). This completes the proof. $\square$

9. Proof of the Upper Bounds

In this section, we prove the upper bounds in Theorem 5.2. We need the following lemma.

Lemma 9.1. Let $\mathcal{V}_{k,\beta}$ be as in (13). Suppose $v \in \mathcal{U}_{\text{SSM}}$ is such that for some positive constants C_1 , C_2 , β , and $k \geq 2$, it holds that

$$\mathcal{L}^v \mathcal{V}_{k,\beta}(x) \leq C_1 - C_2 \mathcal{V}_{k,\beta}(x), \quad \forall x \in \mathbb{R}^2.$$

Let $\hat{X}^n$ denote the diffusion-scaled state process under the scheduling policy $z^n[v]$ in Definition 7.1, and $\hat{\mathcal{L}}_n$ be its generator. Then, there exists $n_0 \in \mathbb{N}$ such that

$$\hat{\mathcal{L}}_n \mathcal{V}_{k,\beta}(\hat{x}) \leq C'_1 - C'_2 \mathcal{V}_{k,\beta}(\hat{x}), \quad \forall \hat{x} \in \tilde{\mathcal{S}}^n$$

for some positive constants C'_1 and C'_2 , and for all $n \geq n_0$.

Proof. See Appendix A. $\square$

We continue with the proof of Theorem 5.2.

Proof of Theorem 5.2. We first prove part (i) for the unconstrained problem. Recall the definition in (13). Let $k = m + 1$. By theorems 5.5 in Arapostathis and Pang [1] and Lemma 4.1, there exists a continuous precise control $v_\epsilon \in \mathcal{U}_{\text{SSM}}$, which is ϵ -optimal for (P1') and satisfies

$$\mathcal{L}^{v_\epsilon} \mathcal{V}_{k,\beta}(x) \leq c_1 - c_2 \mathcal{V}_{k,\beta}(x) \quad \forall x \in \mathbb{R}^2 \quad (70)$$

for any $\beta \geq \bar{\beta}$ defined in (28), and for some positive constants c_1 , c_2 , which depend on β . Recall Definition 2.1. The scheduling policy that we apply to the n th system is as follows: Inside the ball $n\bar{B}$, we apply the Markov

policy in Definition 7.1 $z^n[v_\epsilon]$, while outside this ball, we apply the Markov policy $\tilde{z}^n$ in Definition 3.1. Let Z^n denote this concatenated policy. By Proposition 3.1 and Lemma 9.1, there exist positive constants C_1 , C_2 , β , and $n_0 \in \mathbb{N}$ such that

$$\tilde{\mathcal{L}}_n^{Z^n} \mathcal{V}_{k,\beta}(\hat{x}) \leq C_1 - C_2 \mathcal{V}_{k,\beta}(\hat{x}), \quad \forall \hat{x} \in \mathcal{S}^n, \quad \forall n \geq n_0. \quad (71)$$

Let $\tilde{\Phi}_T^n \equiv \tilde{\Phi}_T^{Z^n}$ as defined in (52). We define

$$\begin{aligned} \hat{q}(\hat{x}) &:= (\hat{x}_1 - Z_{11}^n(\hat{x}) - Z_{12}^n(\hat{x}), \hat{x}_2 - Z_{22}^n(\hat{x})) \\ \hat{y}(\hat{x}) &:= (-Z_{11}^n(\hat{x}) - Z_{12}^n(\hat{x}), -Z_{22}^n(\hat{x})). \end{aligned}$$

By (71), we have $\sup_{n \geq n_0} J(\hat{X}^n(0), Z^n) < \infty$, and by Birkhoff's ergodic theorem for each $n \geq n_0$, there exists $T_n \in \mathbb{R}_+$ such that

$$\left| \int_{\mathbb{R}^2 \times \mathbb{U}} \hat{r}((e \cdot \hat{q}(\hat{x}))^+ u^c, (e \cdot \hat{y}(\hat{x}))^+ u^s) \tilde{\Phi}_T^n(d\hat{x}, du) - J(\hat{X}^n(0), Z^n) \right| \leq \frac{1}{n} \quad (72)$$

for all $T \geq T_n$ and $n \geq n_0$. By (71), the sequence $\{T_n\}$ can be selected so as to also satisfy

$$\sup_{n \geq n_0} \sup_{T \geq T_n} \int_{\mathbb{R}^2 \times \mathbb{U}} \mathcal{V}_{k,\beta}(\hat{x}) \tilde{\Phi}_T^n(d\hat{x}, du) < \infty. \quad (73)$$

Without loss of generality, we assume that $T_n \rightarrow \infty$. Hence, by uniform integrability, which is implied by (73), together with (72) for any $\eta > 0$, there exists a ball B_η such that

$$\left| \int_{B_\eta \times \mathbb{U}} \hat{r}((e \cdot \hat{q}(\hat{x}))^+ u^c, (e \cdot \hat{y}(\hat{x}))^+ u^s) \tilde{\Phi}_T^n(d\hat{x}, du) - J(\hat{X}^n(0), Z^n) \right| \leq \frac{1}{n} + \eta \quad (74)$$

for all $T \geq T_n$ and $n \geq n_0$.

By JWC on $\{\hat{x} \in \sqrt{n}B\}$, we have $(e \cdot \hat{q}(\hat{x}))^+ = (e \cdot \hat{x})^+$ and $(e \cdot \hat{y}(\hat{x}))^+ = (e \cdot \hat{x})^-$ for all $\hat{x} \in B_\eta$, and for all large enough n by Corollary 7.1. On the other hand, $\tilde{\Phi}_T^n$ converges as $(n, T) \rightarrow \infty$, to π_{v_ϵ} in $\mathcal{P}(\mathbb{R}^2 \times \mathbb{U})$ by Lemma 7.2. Therefore

$$\int_{B_\eta \times \mathbb{U}} \hat{r}((e \cdot \hat{q}(\hat{x}))^+ u^c, (e \cdot \hat{y}(\hat{x}))^+ u^s) \tilde{\Phi}_T^n(d\hat{x}, du) \xrightarrow{n \rightarrow \infty} \int_{B_\eta \times \mathbb{U}} r(x, u) \pi_{v_\epsilon}(dx, du). \quad (75)$$

By (74)–(75), we obtain

$$\limsup_{n \rightarrow \infty} J(\hat{X}^n(0), Z^n) \leq \varrho^* + \epsilon + \eta.$$

Since η and ϵ are arbitrary, this completes the proof of part (i).

We next show the upper bound for the constrained problem. Let $\epsilon > 0$ be given. By Theorem 5.7 in Arapostathis and Pang [1] and Lemma 4.1, there exists a continuous precise control $v_\epsilon \in \mathbb{U}_{\text{SSM}}$ and constants $\delta_j^\epsilon < \delta_j$, $j = 1, 2$, satisfying $\pi_{v_\epsilon}(r_0) \leq \varrho_c^* + \epsilon$ and $\pi_{v_\epsilon}(r_j) \leq \delta_j^\epsilon$, $j = 1, 2$, and (70) holds. Let Z^n be the Markov policy constructed in part (i) by concatenating $z^n[v_\epsilon]$ and $\tilde{z}^n$. Following the proof of part (i) and choosing η small enough, i.e., $\eta < \epsilon \wedge \frac{1}{2} \min(\delta_j - \delta_j^\epsilon, j = 1, 2)$, we obtain

$$\begin{aligned} \limsup_{n \rightarrow \infty} J_0(\hat{X}^n(0), Z^n) &\leq \varrho_c^* + 2\epsilon, \\ \limsup_{n \rightarrow \infty} J_{c,j}(\hat{X}^n(0), Z^n) &\leq \frac{1}{2}(\delta_j + \delta_j^\epsilon), \quad j = 1, 2. \end{aligned}$$

This completes the proof of part (ii).

The proof of the upper bound for the fairness problem is analogous to part (ii). By theorem 5.7 and remark 5.1 in Arapostathis and Pang [1], for any $\epsilon > 0$, there exists a continuous precise control $v_\epsilon \in \mathbb{U}_{\text{SSM}}$ for (P3') satisfying

$$\pi_{v_\epsilon}(r_0) \leq \varrho_1^* + \epsilon, \quad \text{and} \quad \pi_{v_\epsilon}(r_1) = \theta \pi_{v_\epsilon}(r_2). \quad (76)$$

Since $\{\pi_{v_\epsilon}, \epsilon \in (0, 1)\}$ is tight, and $(e \cdot x)^-$ is strictly positive on an open subset of B_1 , it follows by the Harnack inequality for the density of the invariant probability measure of the diffusion that

$$\inf_{\epsilon \in (0, 1)} \pi_{v_\epsilon}(r_2) > 0. \quad (77)$$

Arguing as in part (ii), we obtain

$$\begin{aligned} \limsup_{n \rightarrow \infty} J_o(\hat{X}^n(0), Z^n) &\leq \varrho_i^* + \epsilon, \\ \lim_{n \rightarrow \infty} J_{c,j}(\hat{X}^n(0), Z^n) &= \pi_{v_e}(r_j), \quad j = 1, 2. \end{aligned} \quad (78)$$

The result then follows by (76)–(78), thus completing the proof. $\square$

10. Conclusion

We have proved asymptotic optimality for the N-network in the Halfin-Whitt regime. The analysis results in a good understanding of the stability of the diffusion-scaled state processes under certain scheduling policies and the convergence properties of the associated mean empirical measures. The SDP scheduling policy constructed not only gives us a better understanding of the N-network, but also plays a key role in proving the upper bound. In addition, we have identified some important properties of the diffusion-scaled state processes that concern existence of moments, and the convergence of the mean empirical measures. The methodology we followed should help to establish asymptotic optimality for more general multiclass multipool networks in the Halfin-Whitt regime. If this is done, it will nicely complement the results on ergodic control of the limiting controlled diffusion in Arapostathis and Pang [1].

Acknowledgments

The authors thank the referees for the helpful comments that have improved the paper.

Appendix A. Proofs of Proposition 3.1 and Lemma 9.1

In these proofs, we use the fact that the quantities

$$\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n N_{12}^n, \quad n x_1^* - N_1^n - N_{22}^n, \quad \lambda_2^n - \mu_{22}^n N_{22}^n, \quad n x_2^* - N_{22}^n, \quad \text{and} \quad \lambda_2^n - \mu_{22}^n n x_2^*$$

are in $\mathcal{O}(\sqrt{n})$. This is straightforward to verify using Assumption 2.1.

Proof of Proposition 3.1. Simplifying the notation in Definition 3.1, we let $z^n = \tilde{z}^n$, and analogously, for $\tilde{y}^n$ and $\tilde{q}^n$. Fix $k > 2$. Under the scheduling policy in Definition 3.1, the resulting process X^n is Markov with generator

$$\begin{aligned} \mathcal{L}_n^{\tilde{z}^n} f(x) &:= \sum_{i=1}^2 \lambda_i^n (f(x + e_i) - f(x)) + (\mu_{11}^n z_{11}^n + \mu_{12}^n z_{12}^n) (f(x - e_1) - f(x)) \\ &\quad + \mu_{22}^n z_{22}^n (f(x - e_2) - f(x)) + \sum_{i=1}^2 \gamma_i^n q_i^n (f(x - e_i) - f(x)), \quad x \in \mathbb{Z}_+^2, \end{aligned} \quad (A.1)$$

Recall the definition of $\hat{x}$ in (12). Define

$$f_n(x) := |x_1 - n x_1^*|^k + \beta |x_2 - n x_2^*|^k = n^{k/2} (|\hat{x}_1|^k + \beta |\hat{x}_2|^k),$$

for some positive constant β , to be determined later. If we show that

$$\mathcal{L}_n^{\tilde{z}^n} f_n(x) \leq C_1 n^{k/2} - C_2 f_n(x), \quad x \in \mathbb{Z}_+^2, \quad (A.2)$$

for some positive constants C_1 and C_2 , and for all $n \geq n_0$, then by using (15), we obtain (16).

Given (A.2), we easily obtain that

$$\mathbb{E}[f_n(X^n(T))] - f_n(X^n(0)) = \mathbb{E} \left[\int_0^T \mathcal{L}_n f_n(X^n(s)) ds \right] \leq C_1 n^{k/2} T - C_2 \mathbb{E} \left[\int_0^T f_n(X^n(s)) ds \right],$$

which implies that

$$\frac{1}{T} \mathbb{E} \left[\int_0^T \mathcal{V}_{k,\beta}(\hat{X}^n(s)) ds \right] \leq C_1 + \frac{1}{T} \mathcal{V}_{k,\beta}(\hat{X}^n(0)) - \frac{1}{T} \mathbb{E}[\mathcal{V}_{k,\beta}(\hat{X}^n(T))].$$

By letting $T \rightarrow \infty$, this implies that (17) holds.

We now focus on proving (A.2). Note that

$$(a \pm 1)^k - a^k = k a^{k-1} + \mathcal{O}(a^{k-2}) \quad a \in \mathbb{R}.$$

Recall $\tilde{x}$ in (12). Then, by (A.1), we have

$$\mathcal{L}_n^{\tilde{z}^n} f_n(x) = \lambda_1^n (k \tilde{x}_1 |\tilde{x}_1|^{k-2} + \mathcal{O}(|\tilde{x}_1|^{k-2})) + \beta \lambda_2^n (k \tilde{x}_2 |\tilde{x}_2|^{k-2} + \mathcal{O}(|\tilde{x}_2|^{k-2}))$$

$$+ (\mu_{11}^n z_{11}^n + \mu_{12}^n z_{12}^n)(-k\tilde{x}_1|\tilde{x}_1|^{k-2} + \mathcal{O}(|\tilde{x}_1|^{k-2})) + \beta\mu_{22}^n z_{22}^n(-k\tilde{x}_2|\tilde{x}_2|^{k-2} + \mathcal{O}(|\tilde{x}_2|^{k-2})) \\ + \gamma_1^n q_1^n(-k\tilde{x}_1|\tilde{x}_1|^{k-2} + \mathcal{O}(|\tilde{x}_1|^{k-2})) + \beta\gamma_2^n q_2^n(-k\tilde{x}_2|\tilde{x}_2|^{k-2} + \mathcal{O}(|\tilde{x}_2|^{k-2})).$$

Let

$$F_n^{(1)}(x) := (\lambda_1^n + \gamma_1^n q_1^n)\mathcal{O}(|\tilde{x}_1|^{k-2}) + \beta(\lambda_2^n + \gamma_2^n q_2^n)\mathcal{O}(|\tilde{x}_2|^{k-2}) + (\mu_{11}^n z_{11}^n + \mu_{12}^n z_{12}^n)\mathcal{O}(|\tilde{x}_1|^{k-2}) + \beta\mu_{22}^n z_{22}^n\mathcal{O}(|\tilde{x}_2|^{k-2}), \quad (\text{A.3})$$

and

$$F_n^{(2)}(x) := k\tilde{x}_1|\tilde{x}_1|^{k-2}(\lambda_1^n - \gamma_1^n q_1^n) + \beta k\tilde{x}_2|\tilde{x}_2|^{k-2}(\lambda_2^n - \gamma_2^n q_2^n) - k\tilde{x}_1|\tilde{x}_1|^{k-2}(\mu_{11}^n z_{11}^n + \mu_{12}^n z_{12}^n) - \beta k\tilde{x}_2|\tilde{x}_2|^{k-2}\mu_{22}^n z_{22}^n. \quad (\text{A.4})$$

Then,

$$\mathcal{L}_n f_n(x) = F_n^{(1)}(x) + F_n^{(2)}(x).$$

We first study $F_n^{(1)}(x)$. It is easy to observe that for each $i = 1, 2$ and $j = 1, 2$,

$$z_{ij}^n \leq x_i \quad \text{and} \quad q_i^n \leq x_i. \quad (\text{A.5})$$

Thus we obtain

$$\begin{aligned} F_n^{(1)}(x) &\leq (\lambda_1^n + \gamma_1^n x_1)\mathcal{O}(|\tilde{x}_1|^{k-2}) + \beta(\lambda_2^n + \gamma_2^n x_2)\mathcal{O}(|\tilde{x}_2|^{k-2}) \\ &\quad + (\mu_{11}^n + \mu_{12}^n)x_1\mathcal{O}(|\tilde{x}_1|^{k-2}) + \beta\mu_{22}^n x_2\mathcal{O}(|\tilde{x}_2|^{k-2}) \\ &= (\lambda_1^n + \gamma_1^n(nx_1^* + \tilde{x}_1))\mathcal{O}(|\tilde{x}_1|^{k-2}) + \beta(\lambda_2^n + \gamma_2^n(nx_2^* + \tilde{x}_2))\mathcal{O}(|\tilde{x}_2|^{k-2}) \\ &\quad + (\mu_{11}^n + \mu_{12}^n)(nx_1^* + \tilde{x}_1)\mathcal{O}(|\tilde{x}_1|^{k-2}) + \beta\mu_{22}^n(nx_2^* + \tilde{x}_2)\mathcal{O}(|\tilde{x}_2|^{k-2}) \\ &\leq \sum_{i=1}^2 (\mathcal{O}(n)\mathcal{O}(|\tilde{x}_i|^{k-2}) + \mathcal{O}(|\tilde{x}_i|^{k-1})), \end{aligned} \quad (\text{A.6})$$

where the last inequality follows from Assumption 2.1.

We next focus on $F_n^{(2)}(x)$. We consider the following four cases:

Case 1. $x_1 \geq N_1^n + N_{12}^n$, and $x_2 \geq N_{22}^n$. Then,

$$z_{11}^n = N_1^n, \quad z_{12}^n = N_{12}^n, \quad z_{22}^n = N_{22}^n, \quad q_1^n = x_1 - N_1^n - N_{12}^n, \quad q_2^n = x_2 - N_{22}^n.$$

We obtain

$$\begin{aligned} F_n^{(2)}(x) &= k\tilde{x}_1|\tilde{x}_1|^{k-2}[\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n N_{12}^n - \gamma_1^n(nx_1^* - N_1^n - N_{12}^n)] \\ &\quad + \beta k\tilde{x}_2|\tilde{x}_2|^{k-2}[\lambda_2^n - \mu_{22}^n N_{22}^n - \gamma_2^n(nx_2^* - N_{22}^n)] - k\gamma_1^n|\tilde{x}_1|^k - \beta k\gamma_2^n|\tilde{x}_2|^k \\ &= \mathcal{O}(\sqrt{n})(|\tilde{x}_1|^{k-1} + \beta|\tilde{x}_2|^{k-1}) - k\gamma_1^n|\tilde{x}_1|^k - \beta k\gamma_2^n|\tilde{x}_2|^k. \end{aligned} \quad (\text{A.7})$$

Case 2. $x_1 < N_1^n + N_{12}^n$ and $x_2 < N_{22}^n$. Consider two subcases:

Case 2.1. $x_1 > N_1^n$. Then,

$$z_{11}^n = N_1^n, \quad z_{12}^n = x_1 - N_1^n, \quad z_{22}^n = x_2, \quad q_1^n = q_2^n = 0.$$

We have

$$\begin{aligned} F_n^{(2)}(x) &= k\tilde{x}_1|\tilde{x}_1|^{k-2}[\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n(nx_1^* - N_1^n)] + \beta k\tilde{x}_2|\tilde{x}_2|^{k-2}[\lambda_2^n - \mu_{22}^n nx_2^*] - k\mu_{12}^n|\tilde{x}_1|^k - \beta k\mu_{22}^n|\tilde{x}_2|^k \\ &= \mathcal{O}(\sqrt{n})(|\tilde{x}_1|^{k-1} + \beta|\tilde{x}_2|^{k-1}) - k\mu_{12}^n|\tilde{x}_1|^k - \beta k\mu_{22}^n|\tilde{x}_2|^k. \end{aligned} \quad (\text{A.8})$$

Case 2.2. $x_1 < N_1^n$. Then,

$$z_{11}^n = x_1, \quad z_{12}^n = 0, \quad z_{22}^n = x_2, \quad q_1^n = q_2^n = 0.$$

We obtain

$$\begin{aligned} F_n^{(2)}(x) &= k\tilde{x}_1|\tilde{x}_1|^{k-2}[\lambda_1^n - \mu_{11}^n x_1] + \beta k\tilde{x}_2|\tilde{x}_2|^{k-2}[\lambda_2^n - \mu_{22}^n(\tilde{x}_2 + nx_2^*)] \\ &\leq k\tilde{x}_1|\tilde{x}_1|^{k-2}[\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n N_{12}^n] + \beta k\tilde{x}_2|\tilde{x}_2|^{k-2}[\lambda_2^n - \mu_{22}^n nx_2^*] + k\mu_{12}^n N_{12}^n \tilde{x}_1|\tilde{x}_1|^{k-2} - \beta k\mu_{22}^n|\tilde{x}_2|^k. \end{aligned} \quad (\text{A.9})$$

Since $x_1 \leq N_1^n$, we have

$$\mu_{12}^n N_{12}^n \tilde{x}_1 \leq -\frac{\mu_{12}^n N_{12}^n}{N_1^n} |\tilde{x}_1|^2 = -\frac{\mu_{12}^n v_2}{v_1} |\tilde{x}_1|^2 + \mathcal{O}(\sqrt{n})|\tilde{x}_1|.$$

Thus (A.9) takes the form

$$F_n^{(2)}(x) \leq \mathcal{O}(\sqrt{n})(|\tilde{x}_1|^{k-1} + \beta|\tilde{x}_2|^{k-1}) - k\xi_{12}^* \frac{\mu_{12}^n v_2}{v_1} |\tilde{x}_1|^k - \beta k\mu_{22}^n |\tilde{x}_2|^k. \quad (\text{A.10})$$

Case 3. $x_1 \geq N_1^n + N_{12}^n$, and $x_2 < N_{22}^n$. We distinguish the following two subcases.

Case 3.1. $x_1 + x_2 \geq N_1^n + N_2^n$.

Then,

$$z_{11}^n = N_1^n, \quad z_{12}^n = N_2^n - x_2, \quad z_{22}^n = x_2, \quad q_1^n = x_1 + x_2 - N_1^n - N_2^n, \quad q_2^n = 0.$$

We have

$$\begin{aligned} F_n^{(2)}(x) &= k\tilde{x}_1 |\tilde{x}_1|^{k-2} [\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n (N_2^n - nx_2^* - \tilde{x}_2) - \gamma_1^n (x_1 + x_2 - N_1^n - N_2^n)] \\ &\quad + \beta k \tilde{x}_2 |\tilde{x}_2|^{k-2} [\lambda_2^n - \mu_{22}^n (nx_2^* + \tilde{x}_2)] \\ &= k\tilde{x}_1 |\tilde{x}_1|^{k-2} [\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n N_{12}^n + \mu_{12}^n (nx_2^* + N_{22}^n)] \\ &\quad - k\tilde{x}_1 |\tilde{x}_1|^{k-2} \gamma_1^n (nx_1^* + nx_2^* - N_1^n - N_2^n) + \beta k \tilde{x}_2 |\tilde{x}_2|^{k-2} [\lambda_2^n - \mu_{22}^n nx_2^*] \\ &\quad + k(\mu_{12}^n - \gamma_1^n) \tilde{x}_1 \tilde{x}_2 |\tilde{x}_1|^{k-2} - k\gamma_1^n |\tilde{x}_1|^k - \beta k \mu_{22}^n |\tilde{x}_2|^k \\ &= \mathcal{O}(\sqrt{n})(|\tilde{x}_1|^{k-1} + \beta |\tilde{x}_2|^{k-1}) + k(\mu_{12}^n - \gamma_1^n) \tilde{x}_1 \tilde{x}_2 |\tilde{x}_1|^{k-2} - k\gamma_1^n |\tilde{x}_1|^k - \beta k \mu_{22}^n |\tilde{x}_2|^k. \end{aligned} \quad (\text{A.11})$$

Case 3.2. $x_1 + x_2 < N_1^n + N_2^n$. Then,

$$z_{11}^n = N_1^n, \quad z_{12}^n = \tilde{x}_1 + nx_1^* - N_1^n, \quad z_{22}^n = x_2, \quad q_1^n = q_2^n = 0.$$

We have

$$\begin{aligned} F_n^{(2)}(x) &= k\tilde{x}_1 |\tilde{x}_1|^{k-2} [\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n (\tilde{x}_1 + nx_1^* - N_1^n)] + \beta k \tilde{x}_2 |\tilde{x}_2|^{k-2} [\lambda_2^n - \mu_{22}^n (nx_2^* + \tilde{x}_2)] \\ &= k\tilde{x}_1 |\tilde{x}_1|^{k-2} [\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n N_{12}^n - \mu_{12}^n (nx_1^* - N_1^n - N_{12}^n)] \\ &\quad + \beta k \tilde{x}_2 |\tilde{x}_2|^{k-2} [\lambda_2^n - \mu_{22}^n nx_2^*] - k\mu_{12}^n |\tilde{x}_1|^k - \beta k \mu_{22}^n |\tilde{x}_2|^k \\ &= \mathcal{O}(\sqrt{n})(|\tilde{x}_1|^{k-1} + \beta |\tilde{x}_2|^{k-1}) - k\mu_{12}^n |\tilde{x}_1|^k - \beta k \mu_{22}^n |\tilde{x}_2|^k. \end{aligned} \quad (\text{A.12})$$

Case 4. $x_1 < N_1^n + N_{12}^n$, and $x_2 \geq N_{22}^n$. Here, we distinguish four subcases.

Case 4.1. $x_1 \leq N_1^n$ and $x_2 \leq N_2^n$. Using the argument used in Case 2.2, we obtain the same estimate as (A.10).

Case 4.2. $x_1 \leq N_1^n$ and $x_2 > N_2^n$. Then,

$$z_{11}^n = x_1, \quad z_{12}^n = 0, \quad z_{22}^n = N_2^n, \quad q_1^n = 0, \quad q_2^n = x_2 - N_2^n.$$

We use the inequality

$$\mu_{22}^n N_{12}^n + \gamma_2^n (\tilde{x}_2 + nx_2^* - N_2^n) \geq (\mu_{22}^n \wedge \gamma_2^n) \tilde{x}_2 + \mathcal{O}(\sqrt{n}), \quad x_2 > N_2^n$$

to write

$$\begin{aligned} \lambda_2^n - \mu_{22}^n N_2^n - \gamma_2^n (x_2 - N_2^n) &= \lambda_2^n - \mu_{22}^n N_2^n - \gamma_2^n (\tilde{x}_2 + nx_2^* - N_2^n) \\ &\leq \lambda_2^n - \mu_{22}^n N_{22}^n + (\mu_{22}^n \wedge \gamma_2^n) \tilde{x}_2 + \mathcal{O}(\sqrt{n}). \end{aligned}$$

Therefore, as in Case 2.2, we obtain

$$\begin{aligned} F_n^{(2)}(x) &\leq k\mathcal{O}(\sqrt{n})|\tilde{x}_1|^{k-1} + \beta k \tilde{x}_2 |\tilde{x}_2|^{k-2} [\lambda_2^n - \mu_{22}^n N_{22}^n + \mathcal{O}(\sqrt{n})] \\ &\quad - k\xi_{12}^* \frac{\mu_{12}^n \nu_2}{\nu_1} |\tilde{x}_1|^k - \beta k (\mu_{22}^n \wedge \gamma_2^n) |\tilde{x}_2|^k \\ &\leq \mathcal{O}(\sqrt{n})(|\tilde{x}_1|^{k-1} + \beta |\tilde{x}_2|^{k-1}) - k\xi_{12}^* \frac{\mu_{12}^n \nu_2}{\nu_1} |\tilde{x}_1|^k - \beta k (\mu_{22}^n \wedge \gamma_2^n) |\tilde{x}_2|^k. \end{aligned} \quad (\text{A.13})$$

Case 4.3. $x_1 > N_1^n$, and $x_1 + x_2 < N_1^n + N_2^n$. Then,

$$z_{11}^n = N_1^n, \quad z_{12}^n = x_1 - N_1^n, \quad z_{22}^n = x_2, \quad q_1^n = 0, \quad q_2^n = 0.$$

We obtain

$$\begin{aligned} F_n^{(2)}(x) &= k\tilde{x}_1 |\tilde{x}_1|^{k-2} [\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n (\tilde{x}_1 + nx_1^* - N_1^n)] + \beta k \tilde{x}_2 |\tilde{x}_2|^{k-2} [\lambda_2^n - \mu_{22}^n (\tilde{x}_2 + nx_2^*)] \\ &= k\tilde{x}_1 |\tilde{x}_1|^{k-2} [\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n N_{12}^n - \mu_{12}^n (nx_1^* - N_1^n - N_{12}^n)] \\ &\quad + \beta k \tilde{x}_2 |\tilde{x}_2|^{k-2} [\lambda_2^n - \mu_{22}^n N_{22}^n - \mu_{22}^n (nx_2^* - N_{22}^n)] - k\mu_{12}^n |\tilde{x}_1|^k - \beta k \mu_{22}^n |\tilde{x}_2|^k \\ &= \mathcal{O}(\sqrt{n})(|\tilde{x}_1|^{k-1} + \beta |\tilde{x}_2|^{k-1}) - k\mu_{12}^n |\tilde{x}_1|^k - \beta k \mu_{22}^n |\tilde{x}_2|^k. \end{aligned} \quad (\text{A.14})$$

Case 4.4. $x_1 > N_1^n$, and $x_1 + x_2 \geq N_1^n + N_2^n$. Then,

$$Z_{11}^n = N_1^n, \quad Z_{12}^n = x_1 - N_1^n, \quad Z_{22}^n = N_2^n + N_1^n - x_1, \quad Q_1^n = 0, \quad Q_2^n = x_1 + x_2 - N_1^n - N_2^n.$$

Therefore, we obtain

$$F_n^{(2)}(x) = q\beta_1 \tilde{x}_1 |\tilde{x}_1|^{q-2} [\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n (\tilde{x}_1 + nx_1^* - N_1^n)]$$

$$\begin{aligned}
& + q\beta_2 \tilde{x}_2 |\tilde{x}_2|^{q-2} [\lambda_2^n - \mu_{22}^n (N_2^n + N_1^n - x_1) - \gamma_2^n (x_2 - (N_2^n + N_1^n - x_1))] \\
& \leq q\beta_1 \tilde{x}_1 |\tilde{x}_1|^{q-2} [\lambda_1^n - \mu_{11}^n N_1^n - \mu_{12}^n N_{12}^n - \mu_{12}^n (\tilde{x}_1 + nx_1^* - N_1^n - N_{12}^n)] \\
& \quad + q\beta_2 \tilde{x}_2 |\tilde{x}_2|^{q-2} [\lambda_2^n - (\mu_{22}^n \wedge \gamma_2^n)(\tilde{x}_2 + nx_2^*)] \\
& \leq \mathcal{O}(\sqrt{n})(\beta_1 |\tilde{x}_1|^{q-1} + \beta_2 |\tilde{x}_2|^{q-1}) - q\beta_1 \mu_{12}^n |\tilde{x}_1|^q - q\beta_2 (\mu_{22}^n \wedge \gamma_2^n) |\tilde{x}_2|^q,
\end{aligned} \tag{A.15}$$

where the first inequality follows by observing that

$$\mu_{22}^n (N_2^n + N_1^n - x_1) + \gamma_2^n (x_2 - (N_2^n + N_1^n - x_1)) \geq (\mu_{22}^n \wedge \gamma_2^n) x_2,$$

since $x_2 \geq N_2^n + N_1^n - x_1$, and $N_2^n + N_1^n - x_1 > N_{22}^n + N_1^n - x_1 > 0$.

By Young's inequality, we have

$$|\tilde{x}_1|^{k-1} |\tilde{x}_2| \leq \epsilon |\tilde{x}_1|^k + \frac{1}{\epsilon^{k-1}} |\tilde{x}_2|^k, \quad |\tilde{x}_2|^{k-1} |\tilde{x}_1| \leq \epsilon |\tilde{x}_1|^k + \frac{1}{\epsilon^{1/k-1}} |\tilde{x}_2|^k$$

for any $\epsilon > 0$. Using this in (A.11) in combination with (A.7)–(A.8), (A.10), and (A.12)–(A.15), we can choose the constant β properly so that

$$\mathcal{L}_n^{z^n} f_n(x) \leq \sum_{i=1}^2 (\mathcal{O}(n) \mathcal{O}(|\tilde{x}_i|^{k-2}) + \mathcal{O}(\sqrt{n}) \mathcal{O}(|\tilde{x}_i|^{k-1})) - \tilde{C}_2 \sum_{i=1}^2 |\tilde{x}_i|^k \tag{A.16}$$

for some positive constant $\tilde{C}_2$. Now, applying Young's inequality again to the first two terms on the right-hand side of (A.16), we obtain

$$\begin{aligned}
\mathcal{O}(\sqrt{n}) \mathcal{O}(|\tilde{x}_i|^{k-1}) & \leq \epsilon (\mathcal{O}(|\tilde{x}_i|^{k-1}))^{k/(k-1)} + \epsilon^{1-k} (\mathcal{O}(\sqrt{n}))^k, \\
\mathcal{O}(n) \mathcal{O}(|\tilde{x}_i|^{k-2}) & \leq \epsilon (\mathcal{O}(|\tilde{x}_i|^{k-2}))^{k/(k-2)} + \epsilon^{1-k/2} (\mathcal{O}(n))^{k/2}
\end{aligned}$$

for any $\epsilon > 0$. This shows that we can choose β , C_1 , and C_2 appropriately to obtain the claim in (A.2).

Recall $\hat{x}^n$ in (12) and let $\hat{q}_i^n := q_i^n / \sqrt{n}$ for $i = 1, 2$. Concerning the claim in (17) with $\hat{X}^n$ replaced by $\hat{Q}^n$, we observe that in Case 1, $\hat{q}_1^n = \hat{x}_1^n + \mathcal{O}(1)$ and $\hat{q}_2^n = \hat{x}_2^n + \mathcal{O}(1)$, in Case 3.1, $\hat{q}_1^n = \hat{x}_1^n + \hat{x}_2^n + \mathcal{O}(1)$, and $\hat{q}_2^n = 0$, in Case 4.2, $\hat{q}_1^n = 0$, and $\hat{q}_2^n \leq \hat{x}_2^n + \mathcal{O}(1)$, in Case 4.4, $\hat{q}_1^n = 0$, and $\hat{q}_2^n = \hat{x}_1^n + \hat{x}_2^n + \mathcal{O}(1)$, and in all the other cases, $\hat{q}_1^n = \hat{q}_2^n = 0$. The same claim for $\hat{Y}^n$ then follows from the balance Equation (39). The proof of the proposition is complete. $\square$

Proof of Lemma 9.1. We need to show (A.2) holds for $\mathcal{L}_n f_n(x)$ under the scheduling policy $z^n[v]$ in Definition 7.1. We can write $\mathcal{L}_n f_n(x) = F_n^{(1)}(x) + F_n^{(2)}(x)$ with $F_n^{(1)}(x)$ and $F_n^{(2)}(x)$ given by (A.3) and (A.4), respectively. We obtain (A.6) for $F_n^{(1)}(x)$ since (A.5) also holds under the policy $z^n[v]$. For $F_n^{(2)}(x)$, by (A.4) and Definition 7.1, since the control v satisfies (9.1) and $x \in \mathcal{X}^n$ (JWC being satisfied), we easily obtain

$$F_n^{(2)}(x) \leq \mathcal{O}(\sqrt{n})(|\tilde{x}_1|^{k-1} + \beta |\tilde{x}_2|^{k-1}) - \tilde{C}_3 \sum_{i=1}^2 |\tilde{x}_i|^k,$$

for some positive constant $\tilde{C}_3$. Thus, following the argument in the proof of Proposition 3.1, we obtain the claim in (A.2) and hence the result follows by scaling. $\square$

Appendix B. Proof of Theorem 4.3

Recall $\mathcal{H}_i(\theta)$ defined in (32). As in Theorem 4.2, there exists $\lambda^* \in \mathbb{R}$ such that

$$\inf_{\pi \in \mathcal{H}_i(\theta)} \pi(r_o) = \inf_{\pi \in \mathcal{G}} \pi(h_{\theta, \lambda^*}) = \rho_i^*,$$

and the property in (30) implies that the infimum is attained in some $\pi^* \in \mathcal{G}$. Therefore the conclusions analogous to parts (a) and (b) of Theorem 4.2 hold. Part (e) is also standard. It remains to derive the HJB equation and the characterization of optimality corresponding to Theorem 4.2(c)–(d). This is broken in a series of lemmas.

We need to introduce some notation. We denote by τ_δ , $\delta > 0$, the *first exit time* of a process from B_δ^c , i.e.,

$$\tau_\delta := \inf\{t > 0: X_t \notin B_\delta^c\}.$$

We denote by $\mathcal{U}_{\text{SM}}^*$ the class of Markov controls v satisfying $\pi_v(r_o) < \infty$, and by $\mathcal{G}^*$, the corresponding class of ergodic occupation measures.

By the method of proof of (30), there exists inf-compact $\mathcal{V} \in \mathcal{C}^2(\mathbb{R}^2)$ and positive constants κ_1 and κ_2 satisfying

$$\mathcal{L}^\mu \mathcal{V}(x) \leq \kappa_1 - \kappa_2 |x|^m + r_o(x, u), \quad \forall (x, u) \in \mathbb{R}^2 \times \mathbb{U}. \tag{B.1}$$

Moreover, since $(e \cdot x)^- \in \mathcal{D}(|x|^m)$, there exists a constant κ_0 such that

$$(1 + \theta) \lambda^* r_j(x, u) \leq \kappa_0 + \frac{\kappa_2}{2} |x|^m, \quad \forall (x, u) \in \mathbb{R}^2 \times \mathbb{U}, \quad j = 1, 2. \tag{B.2}$$

For $\epsilon > 0$, we define

$$h^\epsilon(x, u) := h_{\theta, \lambda^*}(x, u) + \epsilon \kappa_2 |x|^m.$$

Lemma B.1. *The following hold:*

$$\pi(h_{\theta, \lambda^*}) \leq \kappa_0 + \frac{\kappa_1}{2} + \frac{3}{2}\pi(r_0), \quad \forall \pi \in \mathcal{G}^*, \quad \pi(r_0) \leq \kappa_0 + \frac{\kappa_2}{2} + \pi(h_{\theta, \lambda^*}), \quad \forall \pi \in \mathcal{G}, \quad \pi(h^\epsilon) \leq \epsilon \left(\kappa_0 + \frac{\kappa_1}{2} + \frac{\kappa_2}{2} \right) + (1 + \epsilon)\pi(h_{\theta, \lambda^*}).$$

Proof. This is an easy calculation using (B.1)–(B.2). $\square$

Lemma B.2. *There exists a unique function $V^\epsilon \in \mathcal{C}^2(\mathbb{R}^2)$ with $V^\epsilon(0) = 0$, which is bounded below in $\mathbb{R}^2$, and solves the HJB*

$$\min_{u \in \mathcal{U}} [\mathcal{L}^u V^\epsilon(x) + h^\epsilon(x, u)] = \varrho_\epsilon, \quad x \in \mathbb{R}^2, \quad (\text{B.3})$$

where $\varrho_\epsilon := \inf_{\pi \in \mathcal{G}} \pi(h^\epsilon)$, and the usual characterization of optimality holds. Moreover,

(a) for every $R > 0$, there exists a constant $k_R > 0$ such that

$$\sup_{\epsilon \in (0, 1)} \operatorname{osc}_{B_R} V^\epsilon \leq k_R;$$

(b) if v_ϵ is a measurable a.e. selector from the minimizer of the Hamiltonian in (B.3), then for any $\delta > 0$, we have

$$V^\epsilon(x) \geq \mathbb{E}_x^{v_\epsilon} \left[\int_0^{\tau_\delta} (h^\epsilon(X_s, v_\epsilon(X_s)) - \varrho_\epsilon) ds \right] + \inf_{B_\delta} V^\epsilon;$$

(c) for any stationary control $v \in \mathcal{U}_{\text{SM}}^*$ and for any $\delta > 0$, it holds that

$$V^\epsilon(x) \leq \mathbb{E}_x^v \left[\int_0^{\tau_\delta} (h^\epsilon(X_s, v(X_s)) - \varrho_\epsilon) ds + V^\epsilon(X_{\tau_\delta}) \right].$$

Proof. The proof follows along the lines of theorem 3.3 in Arapostathis et al. [3], using the fact that h^ϵ is inf-compact, for each $\epsilon > 0$, and $\inf_{\pi \in \mathcal{G}} \pi(h^\epsilon) < \infty$ by Lemmas 4.1 and B.1. There is one important difference though: the running cost h^ϵ is not bounded below uniformly in $\epsilon > 0$, and the estimate in part (a) needs special attention. By (B.1)–(B.2), using Itô's formula, we obtain

$$\mathbb{E}_x^U \left[\int_0^\infty e^{-\alpha s} (1 + \theta) \lambda^* r_i(X_s, U_s) ds \right] \leq \mathcal{V}(x) + \mathbb{E}_x^U \left[\int_0^\infty e^{-\alpha s} \left(\kappa_0 + \frac{\kappa_1}{2} + \frac{1}{2} r_0(X_s, U_s) \right) ds \right]$$

for all $U \in \mathcal{U}$, $\alpha > 0$. It follows that, given any ball B_R , the discounted value function

$$\tilde{V}_\alpha^\epsilon(x) := \inf_{U \in \mathcal{U}} \mathbb{E}_x^U \left[\int_0^\infty e^{-\alpha s} (2\kappa_0 + \kappa_1 + h^\epsilon(X_s, U_s)) ds \right]$$

is strictly positive on B_R for all sufficiently small $\alpha > 0$. Therefore, by adding the constant $2\kappa_0 + \kappa_1$ to the running cost, we obtain estimates on the oscillation of $\tilde{V}^\epsilon$ that are uniform over $\epsilon > 0$ by lemmas 3.5 and 3.6 of Arapostathis et al. [3]. $\square$

The next lemma completes the proof of Theorem 4.3.

Lemma B.3. *Let V^ϵ and ϱ_ϵ , for $\epsilon > 0$, be as in Lemma B.2. The following hold:*

(i) *The function V^ϵ converges to some $V_\dagger \in \mathcal{C}^2(\mathbb{R}^2)$, uniformly on compact sets, and $\varrho_\epsilon \rightarrow \varrho_\dagger^*$, as $\epsilon \searrow 0$, and $V_\dagger$ satisfies*

$$\min_{u \in \mathcal{U}} [\mathcal{L}^u V_\dagger(x) + h_{\theta, \lambda^*}(x, u)] = \varrho_\dagger^* = \pi^*(h_{\theta, \lambda^*}). \quad (\text{B.4})$$

Also, any limit point v^* (in the topology of Markov controls) as $\epsilon \searrow 0$ of measurable selectors $\{v_\epsilon\}$ from the minimizer of (B.3) satisfies

$$\mathcal{L}^{v^*} V_\dagger(x) + h_{\theta, \lambda^*}(x, v^*(x)) = \varrho_\dagger^* \quad \text{a.e. in } \mathbb{R}^2.$$

(ii) *A stationary Markov control $v \in \mathcal{U}_{\text{SM}}$ is optimal if and only if it satisfies*

$$H_{h_{\theta, \lambda^*}}(x, \nabla V_\dagger(x)) = b(x, v(x)) \cdot \nabla V_\dagger(x) + h_{\theta, \lambda^*}(x, v(x)) \quad \text{a.e. in } \mathbb{R}^2, \quad (\text{B.5})$$

where $H_{h_{\theta, \lambda^*}}$ is defined in (31) with r replaced by h_{θ, λ^*} .

(iii) *The function V_* has the stochastic representation*

$$V_\dagger(x) = \lim_{\delta \searrow 0} \inf_{v \in \mathcal{U}_{\text{SM}}^*} \mathbb{E}_x^v \left[\int_0^{\tau_\delta} (h_{\theta, \lambda^*}(X_s, v(X_s)) - \varrho_*) ds \right] = \lim_{\delta \searrow 0} \mathbb{E}_x^{\bar{v}} \left[\int_0^{\tau_\delta} (h_{\theta, \lambda^*}(X_s, \bar{v}(X_s)) - \varrho_*) ds \right]$$

for any $\bar{v} \in \mathcal{U}_{\text{SM}}$ that satisfies (B.5).

Proof. We follow the method in the proof of theorem 3.4 in Arapostathis et al. [3]. Since ϱ_ϵ is nonincreasing and bounded below, it converges to some value, which is clearly $\pi^*(h_{\theta, \lambda^*})$ by Lemma B.1. Parts (i) and (iii) then follow as in the proof of lemma 3.9 in Arapostathis et al. [3], and we can follow the method in the proof of lemma 3.10 in the same paper to establish that $V_\dagger^- \in \mathcal{D}(\mathcal{V})$.

Now, let $\hat{v} \in \mathcal{U}_{\text{SM}}$ be any control satisfying (B.5). We modify the estimate in (B.2) and write it as $(1 + \theta)\lambda^* r_i(x, u) \leq \kappa_0 + (\kappa_2/4)|x|^m$ for some constant κ'_0 . An easy calculation using (B.1) then shows that

$$\mathcal{L}^{\hat{v}}(\mathcal{V} + 2V_\dagger) \leq \kappa_0 + \kappa_1 + 2\kappa'_0 - \frac{\kappa_2}{2}|x|^m - h_{\theta, \lambda^*}(x, \hat{v}(x)).$$

Therefore, since $\mathcal{V} + 2V_\dagger$ is inf-compact, we must have $\hat{v} \in \mathcal{U}_{\text{SM}}^*$. Using this and the fact that $V_\dagger^- \in \mathcal{D}(\mathcal{V})$, we deduce that $(1/T)\mathbb{E}_x^{\hat{v}}[V_\dagger^-(X_T)] \rightarrow 0$ as $T \rightarrow \infty$. Hence, by Itô's formula and (B.4), we obtain $\pi_{\hat{v}}(h_{\theta, \lambda^*}) \leq \varrho_\dagger^*$. Thus we must have equality $\pi_{\hat{v}}(h_{\theta, \lambda^*}) = \pi^*(h_{\theta, \lambda^*})$, i.e., $\hat{v}$ is optimal. This completes the proof. $\square$

References

- [1] Arapostathis A, Pang G (2016) Ergodic diffusion control of multiclass multi-pool networks in the Halfin-Whitt regime. *Ann. Appl. Probab.* 26(5):3110–3153.
- [2] Arapostathis A, Borkar VS, Ghosh MK (2012) *Ergodic Control of Diffusion Processes*. Encyclopedia of Mathematics and Its Applications, Vol. 143 (Cambridge University Press, Cambridge, UK).
- [3] Arapostathis A, Biswas A, Pang G (2015) Ergodic control of multi-class $M/M/N + M$ queues in the Halfin-Whitt regime. *Ann. Appl. Probab.* 25(6):3511–3570.
- [4] Armony M (2005) Dynamic routing in large-scale service systems with heterogeneous servers. *Queueing Syst.* 51(3–4):287–329.
- [5] Armony M, Ward AR (2010) Fair dynamic routing in large-scale heterogeneous-server systems. *Oper. Res.* 58(3):624–637.
- [6] Atar R (2005) A diffusion model of scheduling control in queueing systems with many servers. *Ann. Appl. Probab.* 15(1B):820–852.
- [7] Atar R (2005) Scheduling control for queueing systems with many servers: Asymptotic optimality in heavy traffic. *Ann. Appl. Probab.* 15(4):2606–2650.
- [8] Atar R, Mandelbaum A, Reiman MI (2004) Scheduling a multi class queue with many exponential servers: Asymptotic optimality in heavy traffic. *Ann. Appl. Probab.* 14(3):1084–1134.
- [9] Atar R, Mandelbaum A, Shaikh G (2009) Simplified control problems for multiclass many-server queueing systems. *Math. Oper. Res.* 34(4):795–812.
- [10] Bell SL, Williams RJ (2001) Dynamic scheduling of a system with two parallel servers in heavy traffic with resource pooling: Asymptotic optimality of a threshold policy. *Ann. Appl. Probab.* 11(3):608–649.
- [11] Biswas A (2017) An ergodic control problem for many-server multiclass queueing systems with cross-trained servers. *Stochastic Systems* 7(2):264–288.
- [12] Dai JG, Tezcan T (2008) Optimal control of parallel server systems with many servers in heavy traffic. *Queueing Syst.* 59(2):95–134.
- [13] Dai JG, Tezcan T (2011) State space collapse in many-server diffusion limits of parallel server systems. *Math. Oper. Res.* 36(2):271–320.
- [14] Gamarnik D, Stolyar AL (2012) Multiclass multiserver queueing system in the Halfin-Whitt heavy traffic regime: Asymptotics of the stationary distribution. *Queueing Syst.* 71(1–2):25–51.
- [15] Ghamami S, Ward AR (2013) Dynamic scheduling of a two-server parallel server system with complete resource pooling and reneging in heavy traffic: Asymptotic optimality of a two-threshold policy. *Math. Oper. Res.* 38(4):761–824.
- [16] Gurvich I, Whitt W (2009) Queue-and-idleness-ratio controls in many-server service systems. *Math. Oper. Res.* 34(2):363–396.
- [17] Gurvich I, Whitt W (2010) Service-level differentiation in many-server service system via queue-ratio routing. *Oper. Res.* 58(2):316–328.
- [18] Harrison JM (1988) Brownian models of queueing networks with heterogeneous customer populations. Fleming W, Lions PL, eds. *Stochastic Differential Systems, Stochastic Control Theory and Applications*, The IMA Volumes In Mathematics and Its Applications, Vol. 10 (Springer, New York), 147–186.
- [19] Harrison JM (1998) Heavy traffic analysis of a system with parallel servers: Asymptotic optimality of discrete-review policies. *Ann. Appl. Probab.* 8(3):822–848.
- [20] Harrison JM (2000) Brownian models of open processing networks: Canonical representation of workload. *Ann. Appl. Probab.* 10(1):75–103.
- [21] Kallenberg O (2002) *Foundations of Modern Probability*, Probability and Its Applications (New York), 2nd ed. (Springer, New York).
- [22] Luenberger DG (1967) *Optimization by Vector Space Methods* (John Wiley & Sons, New York).
- [23] Stolyar AL (2015) Diffusion-scale tightness of invariant distributions of a large-scale flexible service system. *Adv. Appl. Probab.* 47(1):251–269.
- [24] Stolyar AL, Yudovina E (2012) Tightness of invariant distributions of a large-scale flexible service system under a priority discipline. *Stoch. Syst.* 2(2):381–408.
- [25] Stolyar AL, Yudovina E (2012) Systems with large flexible server pools: Instability of “natural” load balancing. *Ann. Appl. Probab.* 23(5):2099–2183.
- [26] Tezcan T, Dai JG (2010) Dynamic control of N-systems with many servers: Asymptotic optimality of a static priority policy in heavy traffic. *Oper. Res.* 58(1):94–110.
- [27] Ward AR, Armony M (2013) Blind fair routing in large-scale service systems with heterogeneous customers and servers. *Oper. Res.* 61(1):228–243.
- [28] Williams RJ (2000) On dynamic scheduling of a parallel server system with complete resource pooling. McDonald DR, Turner SRE, eds. *Analysis of Communication Networks: Call Centres, Traffic and Performance*. (AMS, Providence, RI), 49–71.
- [29] Williams RJ (2016) Stochastic processing networks. *Annual Rev. Statist. Appl.* 3:323–345.
- [30] Xu SH, Righter R, Shanthikumar JG (1992) Optimal dynamic assignment of customers to heterogeneous servers in parallel. *Oper. Res.* 40(6):1126–1138.