

Opinion piece



Cite this article: Plowright RK, Becker DJ, McCallum H, Manlove KR. 2019 Sampling to elucidate the dynamics of infections in reservoir hosts. *Phil. Trans. R. Soc. B* **374**: 20180336.
<http://dx.doi.org/10.1098/rstb.2018.0336>

Accepted: 12 June 2019

One contribution of 20 to a theme issue
'Dynamic and integrative approaches to
understanding pathogen spillover'.

Subject Areas:

ecology, health and disease and epidemiology

Keywords:

sampling reservoir hosts, emerging infectious
diseases, wildlife disease, zoonoses, spillover

Author for correspondence:

Raina K. Plowright

e-mail: raina.plowright@montana.edu

Electronic supplementary material is available
online at [https://dx.doi.org/10.6084/m9.
figshare.c.4555727](https://dx.doi.org/10.6084/m9.figshare.c.4555727).

Sampling to elucidate the dynamics of infections in reservoir hosts

Raina K. Plowright¹, Daniel J. Becker^{1,2,3}, Hamish McCallum⁴
and Kezia R. Manlove⁵

¹Department of Microbiology and Immunology, Montana State University, Bozeman, MT 59717, USA

²Center for the Ecology of Infectious Diseases, University of Georgia, Athens, GA 30602, USA

³Department of Biology, Indiana University, Bloomington, IN 47405, USA

⁴Environmental Futures Research Institute, Griffith University, Brisbane, Queensland 4111, Australia

⁵Department of Wildland Resources and Ecology Center, Utah State University, Logan, UT 84321, USA

RKP, 0000-0002-3338-6590; DJB, 0000-0003-4315-8628; KRM, 0000-0002-7200-5236

The risk of zoonotic spillover from reservoir hosts, such as wildlife or domestic livestock, to people is shaped by the spatial and temporal distribution of infection in reservoir populations. Quantifying these distributions is a key challenge in epidemiology and disease ecology that requires researchers to make trade-offs between the extent and intensity of spatial versus temporal sampling. We discuss sampling methods that strengthen the reliability and validity of inferences about the dynamics of zoonotic pathogens in wildlife hosts.

This article is part of the theme issue 'Dynamic and integrative approaches to understanding pathogen spillover'.

1. Introduction

At any point in space and time, the risk of pathogen spillover from reservoir hosts to humans, or to other animals, is a function of the intensity of infection within reservoir host populations [1]. Spillover risk is then shaped by a series of processes including the release of infectious particles from reservoir hosts, response of pathogen survival to the environment, behaviours that affect exposure of recipient hosts, and biologically driven susceptibility of recipient hosts [1]. Here, we focus on the dynamics of infection in wildlife reservoirs that determine how pathogen intensity is distributed in space and time.

Quantification of these dynamics is necessary to predict and manage zoonotic transmission [2]. Emerging zoonoses often appear suddenly in novel recipient hosts, and sampling of reservoir hosts (hosts in which the pathogen is maintained [3]) is initiated before the dynamics of the zoonosis are fully characterized. Therefore, sampling often is opportunistic or haphazard and is guided by sparse information on, for example, the observed patterns of spillover and the natural history of the reservoir host. Moreover, information on the pathogen in reservoir hosts, which often tolerate infection with no apparent clinical symptoms or pathology [4], is difficult to obtain. Catching, restraining and sampling hosts, or testing excreta such as urine or faeces, is usually required [5,6], and can be logistically intensive, expensive and hazardous. Therefore, we address how sampling can be designed to minimize these challenges while maximizing information gain.

Design of any sampling strategy requires clear specification of objectives (table 1). The fundamental objective of sampling to inform management of spillover is to identify times and places at which the risk of spillover is elevated. If the reservoir hosts of an emerging pathogen are not well characterized, a wide variety of potential hosts must be sampled to determine whether and the extent to which they can be infected by the pathogen [7]. In other situations, the reservoir hosts are known and are subject to pathogen incursions from other locations (e.g. [8]). Here, we review sampling strategies for situations in which one or more potential reservoir species have been identified, and the pathogen may be endemic. We

Table 1. Illustrative goals and objectives motivating studies of pathogens in reservoir hosts.

goal	objectives	data needed	sampling approaches
identify distribution of reservoir hosts and how those hosts and populations are connected through space and time	estimate the spatial distribution and density of reservoir hosts estimate the spatial and temporal extents of connections between reservoir host units	spatially explicit presence and absence of marked or unmarked animals movements of known and instrumented individuals population-level genetics on individuals and groups	aerial or ground surveys that account for detection probability and use a robust design
identify times and places with high prevalence in reservoir hosts	determine coarse-resolution patterns of prevalence in reservoir hosts, variation in prevalence among reservoir host populations, and the spatial extent of infection estimate the spatial and temporal autocorrelation of infectious animals or populations	spatially and temporally replicated prevalence and seroprevalence within- and among-population prevalence (or seroprevalence if the refractory period is short relative to the host lifespan) in space and time and over life-history stages	probabilistic, spatially and temporally stratified sampling of reservoir host populations adaptive sampling of probabilistically selected, higher-prevalence sites
identify causes of high prevalence in reservoir hosts	estimate the infectious period, exogenous and endogenous covariates associated with infection, and shedding loads	pathogen status, load, immunity and demography of infected and uninfected hosts	probabilistic sampling of populations with high, moderate and low prevalence
identify patterns of transmission in the reservoir host	identify covariates associated with increased susceptibility or transmission estimate rates of change of prevalence to inform the temporal resolution of sampling	time-series of cases or seroconversions in space from same locations longitudinal sampling of individual infection status to identify change in infection state of individuals over time age-stratified prevalence or seroprevalence	longitudinal sampling of individuals and populations
estimate risk of spillover to recipient hosts across space and time to predict future events	investigate the pathogen's potential to persist in the environment describe when, where, and how reservoir and recipient host species interact	biotic and abiotic environmental attributes at small and large spatial extents and resolutions contacts among individuals multispecies (sero)surveys to identify high-prevalence hosts in areas with high prevalence comparative studies of exuded load per host across host species to understand variation in pathogen release among host species	probabilistic, spatially and temporally stratified sampling of reservoir host populations
explore interventions to reduce prevalence or magnitude of an epidemic, or eradicate infection from reservoir hosts	estimate rate of epidemic growth and reproductive number (R_0) estimate rates of effective vaccination or culling design effective implementation strategies (e.g. ring vaccination or culling, treatments at the infection front)	prevalence over time and duration in infection class age-stratified prevalence or seroprevalence and demographic data	longitudinal sampling during epidemics at invasion zones

do not cover well-described statistical approaches such as power and sample size analyses. We also simulate a theoretical wildlife disease to illustrate how spatial and temporal variability or synchrony in infection dynamics can inform sampling decisions and inferences about disease dynamics.

2. Processes driving the distribution and synchrony of pathogens in reservoir hosts

(a) Spatial and temporal distribution and intensity

The distribution and intensity of infection in reservoir hosts vary among individuals and among populations in space and time. This variation is driven by many within- and between-host factors, including host demography and movement, transmission rates, infectious periods and herd immunity. Below we describe the factors that influence spatial and temporal variation. In most instances, there is little *a priori* information to decipher the drivers of variation, and these knowledge gaps must be addressed with sampling (table 1).

When a pathogen first invades a population of hosts, transmission dynamics may be synchronized through the invasion process [9]. Such invasion dynamics may characterize West Nile virus and avian influenza H5N1 invasion in wild birds in the USA, or Zika virus invasion in marmosets (*Callithrix species*) in Brazil [10–12]. During this phase, the basic reproduction number, R_0 , the average number of secondary infections generated over an individual's infectious period when infection is rare [13], is a powerful distillation of the efficiency of pathogen spread. Beyond the initial stage of pathogen invasion, however, R_0 does not capture all fundamental elements of pathogen dynamics that affect the distribution of infection in space and time. For example, a pathogen that produces an acute (short-lived) immunizing infection with high transmission and mortality rates might spread quickly through local host populations and then fade out, to be reintroduced after the pool of susceptible hosts is replenished. Such short-lived epidemics of *Yersinia pestis* in rats, for example, may explain the sporadic outbreaks of bubonic plague in humans in both fourteenth to sixteenth century Europe [14] and modern urban Madagascar [15]. By contrast, another pathogen with a comparable R_0 , but a long infectious period and low transmission rate, may persist in the same population for long periods, eventually producing a spatially and temporally stable infection intensity as exemplified by *Mycobacterium bovis* in white-tailed deer (*Odocoileus virginianus*) in Michigan, USA [16].

Once a pathogen has established in the reservoir host populations, within-host factors (e.g. duration of infection within hosts) interact with the population dynamics, density and movement of reservoir hosts to determine the distribution of infection among hosts [17–19]. The intensity of infection within individuals is governed by immune responses and pathogen life history. Because individuals acquire immunity, they typically have higher pathogen loads during their first infection than during subsequent infections. For example, juvenile *Rousettus aegyptiacus* bats that excreted high levels of Marburg virus during their first infections were linked to spillover of the virus to humans in Uganda [20]. Similarly, pathogen burden and shedding rates can vary over the course of infection as a function of changes in the immune response, microbiome and pathogen

distribution within the host tissues. Bank voles (*Clethrionomys glareolus*) infected with Puumala virus shed high titres of virus during the acute phase of infection and low titres during the chronic phase of infection [21]. Pathogen levels also can rise if host individuals are infected with multiple pathogens or are immunocompromised by physiological or environmental stress. Laboratory mice infected with both worms and bacteria, for example, shed more of both for longer than those infected with either pathogen in isolation [22], and bats (*Pteropus alecto*) are hypothesized to excrete zoonotic viruses during winter, when environmental stress drives reactivation of latent viruses [23–25], but not during summer, when food is abundant [26].

(b) Synchrony

The nature of the transmission process usually leads to synchrony in the distribution of infection among individuals at some spatial scale. Tobler's first law of geography states 'near things are more related than distant things' [27]. Accordingly, the correlation of values of a variable through space (spatial autocorrelation) and time (temporal autocorrelation) is usually positive. Understanding spatial synchrony of infection [28] can help inform sampling design.

Synchrony of infections within populations can arise through myriad processes that drive pathogen transmission (electronic supplementary material, table S1). These include processes that drive synchrony of animal populations (dispersal, social organization and Moran effects (correlation between population size or density that is linearly related to the correlation between their environments), figure 1), processes that drive synchrony of population-immunity (e.g. the strong and directional autocorrelation of invasion, the influx of susceptibles through birth, and synchronization of susceptibility through stress from environmental perturbations), and processes that drive synchrony of exposure, such as pathogen survival (e.g. the response of influenza survival to humidity [29] and of Hendra virus survival to temperature [30]), or behaviour (e.g. winter consumption of date-palm sap by humans in Bangladesh [31]). Synchrony of transmission commonly is seasonal, especially in temperate zones [32]. Efficient sampling of the distribution of infection requires estimating the spatial and temporal scales at which infection dynamics in the reservoir host are synchronized and the extent to which variability or trends in those dynamics are predictable [32].

The spatial extent of synchrony depends on the extent of the process driving synchrony. When host movement drives infection dynamics, spatial autocorrelation is driven by the rate of host movement relative to the infectious period of the pathogen [33]. For example, the more thoroughly mixed the contacts among populations, and the longer the infectious period, the lower the spatial variability in infection dynamics (figure 2c,d). However, acute pathogens in hosts with high connectivity may be as thoroughly mixed as chronic pathogens in hosts with low connectivity [33]. If movement rates of hosts are not high enough to ensure transmission to new subpopulations before recovery from infection, infection can become trapped in subgroups [35]. For example, the spread of measles, an acute disease, is limited in regions where walking is a more common mode of transportation than motorized vehicles [36]. By contrast, measles occurred in waves across the UK prior to vaccination [37] because great




Figure 1. Mechanisms governing spatial and temporal patterns and structure of disease dynamics include dispersal of individuals, social organization and synchrony of host populations and Moran effects. For each mechanism, the color on the spatial prevalence panel indicates the time at which the local outbreak began, and the circles represent the extent of the pathogen in space. The prevalence curves in the local temporal dynamics panel show prevalence through time at the epicentre of each local outbreak. In (a), dispersal of infected hosts (bighorn sheep with red circles) produces spatial and temporal autocorrelation consistent with the movement patterns of the primary host. In (b), synchronous demographic or behavioural dynamics within the host species produce synchrony in spatial and temporal dynamics of prevalence within all host populations. In (c), Moran effects across populations create synchrony among populations experiencing similar environmental conditions. Here, we imagine that limited nutritional availability consistently increases host susceptibility to infection during autumn in forest populations. This leads to synchronous outbreaks in all forest populations that experience the nutrient deficit, without simultaneous outbreaks in populations in locations that are not forested. Cases are infected individuals.




Figure 2. A simulation of pathogen dynamics in reservoir host populations with varying autocorrelation of prevalence in space (populations) and time (days). Prevalence falls along spatial and temporal gradients of variability. In (a), variability is high over space but low over time (e.g. chronic infections with highly variable prevalence over locations but stable prevalence over time such as hepatitis B in human populations [34]). In (b), variability is high over space and time (e.g. acute pathogens such as canine distemper virus in carnivores at the extent of the USA). In (c), variability is low over space and time, as with chronic, endemic pathogens in highly connected populations, such as herpes simplex virus in human populations. In (d), variability is low over space but high over time, as in highly contagious infections with seasonal transmission such as influenza. Methods for the simulations are described in electronic supplementary material, Methods.

distances could be travelled by car or train within the two-week infectious period. High movement rates relative to infectious periods create synchronous dynamics across populations, maintenance of herd immunity and dampening of epidemics [38]. However, high synchrony also can lead

to synchronous pathogen extinction across all infected populations. Understanding host movement relative to the pathogen's infectious period is critical to inform sampling.

Unpredictable spatial synchrony in transmission can be generated through synchronous environmental stochasticity

[39]. Stochastic perturbations in climate or extreme events such as hurricanes and fires have synchronized parasitic disease in grouse (*Lagopus lagopus scoticus*) [39,40] and have been hypothesized to cause pulses of viral shedding from bats (*Pteropus* species) [38,41]. Comparing the extent of synchronous infection dynamics with the extent of environmental variation can be useful for causal inference [42].

Truly decoupled dynamics across space or time, where dynamics vary chaotically among populations or depend on factors that are limited in space and time (figure 2b), also can occur. Isolation, for example on islands, may foster uncoupling from other populations, and reduce the likelihood of pathogen persistence, particularly if populations are small and infectious periods are short relative to host lifespans [33]. Complex transmission feedbacks within populations may drive spatial asynchrony. For example, longer infectious periods in older bighorn sheep (*Ovis canadensis*) may drive feedbacks that result in either high or low endemic states of pneumonia-causing *Mycoplasma ovipneumoniae* depending on the age-structure of the population at first infection [43]. These feedbacks could create dynamics that are relatively independent across space but have temporal trends. Thus, sampling multiple populations across space, with many sampling points through time, is essential to make inferences that can be generalized across all infected bighorn sheep populations [44,45].

3. Sampling the distribution of infection in reservoir hosts

(a) Basic sampling principles

Designing a sampling strategy that identifies how pathogens are distributed in space and time is challenging. In particular, trade-offs between the spatial extent of sampling and intensity of sampling are ubiquitous. Sampling intensity should be informed by temporal and spatial variability in prevalence (the proportion of infected individuals in a population) and by spatial and temporal autocorrelation. In general, more sampling is required in more variable and less autocorrelated systems. The observed autocorrelation depends on resolution of observation and units for sampling and analysis (e.g. individuals, populations or regions; days, months or years), which must reflect the objectives of the study (table 1). For example, canine distemper virus in wolves (*Canis lupus*) might appear patchily distributed at coarse resolution (e.g. counties within Montana), but synchronous at fine resolution (e.g. Yellowstone National Park [46]). If the resolution of data collection is coarse, one may infer that infection dynamics are spatially independent (figure 2a,b), whereas finer-resolution sampling within a more limited spatial extent may indicate spatial homogeneity (figure 2c,d). Neither choice is erroneous, as long as the inferences are aligned with the scale of investigation. Ultimately, however, trade-offs between spatial versus temporal sampling, and decisions on the resolution and units of analysis, are decided by the prioritization of objectives and resources available for sampling.

To draw valid statistical inferences from field data, theory generally prescribes sampling randomly in space and time [47]. However, for logistical reasons, probabilistic sampling (random selection of samples in which all units have equal probability of being selected) is rare in animal epidemiology.

For example, social hosts, such as bats that roost in a common location, are not randomly distributed, and it is also challenging to randomly sample bats within these roosts [48]. Stratified random sampling or other generalized random stratified designs may be more feasible while achieving a spatially well-balanced random sample and valid inferences [49–52]. The stratification can be informed by knowledge of seasonal peaks in prevalence and concentrations of infection to focus efforts in high-risk places and times. Sampling also continues at times and in locations where risk of infection is thought to be low. Designs with unequal probability selection based on an auxiliary variable also can be considered in such cases.

Stratified random sampling is less feasible for pathogens for which prevalence responds to ephemeral environmental drivers, or to transient dynamics in the reservoir host. In these circumstances, one may wish to implement adaptive sampling, in which probabilistic sampling is complemented by more-intensive spatial and temporal sampling during an outbreak or spillover. Opportunistic sampling often is deployed following spillover in an effort to isolate pathogens or identify reservoir hosts [53]. However, if opportunistic sampling is accompanied by some probabilistic sampling, it typically provides more insight into the spatial and temporal dynamics of infection [54–58].

Although adaptive sampling designs are valuable for maximizing data collection around mortality events or spillover, or for sampling when there is little *a priori* information, simulations suggest that statistical power to detect temporal trends in infection dynamics is greater when populations are sampled repeatedly and consistently over a long period of time (e.g. monthly over a few years) than when a given population is sampled for a short period of time, albeit repeatedly [59] (box 1; e.g. daily or weekly over a few months). Alternative sampling designs [59,73,74] are robust to infrequent sampling, distant sites and large spatial extents. These include: rotating panels, which sample each site repeatedly but during temporal windows that do not fully overlap; augmented, serially alternating panels, which complement rotating panels with consistent sampling of one location; and partially augmented, serially alternating panels, in which infrequent sampling of a given location periodically is complemented with frequent sampling of the location (box 1 and table 2).

Another framework that recently has emerged from the disease ecology literature is model-guided fieldwork [75], where mathematical models of pathogen dynamics are developed *a priori* to guide field data collection. Modellers and biologists work together to incorporate multiple hypotheses and uncertainty about the structure of dynamics and then iterate between models and measurement. Such approaches can facilitate transdisciplinary research and lead to more robust inferences.

(b) Targeted approaches to increase information about prevalence

Prevalence is usually inferred from spatially and temporally explicit data on individual infection or exposure status. These data are usually information-weak because the outcome of every sample is binary (infected or not infected), and the outcomes may be subject to error. In addition to simply increasing the number of individuals sampled,

Box 1. Sampling to characterize pathogen prevalence in reservoir hosts.

To illustrate how sampling methods affect interpretation of pathogen dynamics, we used R [60] to simulate infection in a reservoir host as a random realization of a binomial point process [61] (figure 3); see the electronic supplementary material for more information and R code. The kernel-smoothed intensity of this process illustrates spatial and temporal concentrations of infection. We simulated sampling of this hypothetical host over space and time with different designs. Points and thick lines in figure 3 indicate the prevalence estimated by each sampling design. Although this simulation is a clear oversimplification of reality (e.g. no population structure, no underlying mechanistic model of infection dynamics), this serves as a heuristic tool to illustrate the variation in inference about prevalence estimates from different spatio-temporal sampling designs.

In opportunistic or haphazard designs, nearby populations are sampled following a spillover event (A). For example, the emergence of severe acute respiratory syndrome in 2002 in Guangdong Province, China was followed rapidly by surveys of mammals in wet markets [62]. Similarly, the emergence of Hendra virus in Brisbane, Australia in 1994 was followed by surveys of wild and domestic animals through Queensland [63]. Although important for isolating virus and identifying reservoir hosts, opportunistic or haphazard sampling may overestimate prevalence, may not capture both spatial and temporal variation in infection dynamics, and may limit the validity of inferences about the population [47]. Repeated (i.e. longitudinal) sampling of single or a few populations (B, C) is common in disease ecology [64–67]. However, longitudinal sampling across a broad number of spatially replicated populations is rare during epidemiological surveillance of wildlife [68]. Opportunistic sampling of multiple populations may be complemented with repeated sampling of one population (e.g. [69]). Longitudinal sampling often is conducted at regular intervals (e.g. every four months), and such designs can capture or consistently fail to detect temporal peaks in viral shedding from reservoirs.

Because pathogen transmission processes are temporally dynamic, even coarse-resolution spatial sampling must be temporally explicit. Estimates of prevalence from one population at one point in time may be misleading if disease dynamics are sufficiently rapid that prevalence changes substantially over the study period. For example, a cross-sectional sample of a travelling wave epidemic could bias estimates of spatial variance if different populations are at troughs and peaks of prevalence. Nevertheless, estimates of prevalence that are based on pooling of samples in space or time frequently are reported in the literature [70].

Random sampling can reduce bias that results from sampling at regular intervals (D). Although sampling with random designs may reduce bias in estimates of spatial and temporal infection dynamics, it may not capture temporal trends within a given location. Furthermore, random samples may be clustered in time and space. Additionally, random selection of sampling locations may have less statistical power than intentional selection of sampling locations [71]. Moreover, truly random sampling may not make sense for certain taxa, such as central-place foragers (e.g. many bats), which are most easily sampled at locations that are not randomly distributed (e.g. roosts) [48]. Stratified random sampling in space and time [72] may be more effective. For example, in E, two samples are drawn from each region. Another alternative to random sampling is adaptive sampling (F), in which random sampling is augmented by more-intensive sampling in the spillover region. This design reduces bias associated with longitudinal surveys while capitalizing on opportunistic sampling following spillover (e.g. virus isolation).

Panels F through I illustrate designs from the sampling literature [59,73,74]. Rotating panels (G) sample each site a finite number of times; as sampling of each site ceases, sampling of another site begins. Although rotating panels can help infer fine-resolution temporal dynamics efficiently over space, they also can restrict broader longitudinal analyses and may change the state of the epidemiological system if sampling of a given site occurs too frequently [73]. Serially alternating sampling is similar to rotating sampling but increases the interval between samples of each site. Both the rotating and the serially alternating designs can be augmented with longitudinal sampling of single or multiple sites (e.g. H). The partially augmented, serially alternating design replaces longitudinal sampling of one site with sampling of multiple sites at consecutive intervals (I). Prior simulations suggested that the power of augmented, serially alternating and partially augmented, serially alternating sampling to detect temporal trends is greater than that of rotating sampling [59]. However, given that these designs include replicated sampling over time per multiple sites, their implementation can require ample sampling effort and therefore resources in terms of personnel, time and funding.

the information content of wildlife field samples can be augmented in several ways.

Information on ages of sampled animals is useful because age–seroprevalence or age–prevalence curves can be used to estimate transmission rates [17]. Seroconversion of juveniles provides clear evidence of ongoing pathogen transmission within a population. If juveniles seroconvert each year, the pathogen is likely to be persistent and endemic in that population rather than infrequent and oscillatory. Serosurveys of juveniles are particularly useful in systems where long-lived circulating antibodies are the only measurable indication of exposure (e.g. African bat henipaviruses, in which RNA

rarely is found, and virus has not yet been isolated [76,77]), as long as maternal immunity is not mistaken for juvenile exposure [78]. Sampling of isolated populations (e.g. [76]) similarly can help distinguish between pathogen persistence at the population level versus spatially and temporally patchy transmission.

Seroprevalence can be useful for monitoring spatial and temporal trends in prevalence if few individuals are infected at any point in time (e.g. with infectious periods short or transmission rates low), or if detection is difficult (e.g. lethal sampling is required to test whether lyssaviruses are present in most mammals [79]). However, if antibodies

Table 2. Sampling designs used in studies of pathogen dynamics in reservoir hosts.^a

sampling design	description	advantages	disadvantages
opportunistic	nearby populations are sampled following a spillover event	isolating pathogen, identify reservoir hosts, pragmatic	overestimate prevalence, cannot capture spatial or temporal variation, skew inference of prevalence
single longitudinal	repeated sampling of single population over time	infer some temporal dynamics	cannot capture spatial variation, regular intervals could consistently miss shedding pulses
replicated longitudinal	repeated sampling of multiple populations over time	infer some spatial and temporal dynamics	logistically challenging, regular intervals could consistently miss shedding pulses
random sample	random distribution of sampling events over space and time	reduce bias from sampling at regular intervals	may not capture spatial or temporal variation when truly random, may not be feasible for many species
random stratified	random sampling from predetermined regions in space and time	more likely to obtain a representative spatial and temporal sample	may require greater effort than a simple random sample
adaptive sampling	random sampling augmented by intensive spatio-temporal sampling near outbreaks	reduce bias while capturing benefits of opportunistic sampling (e.g. isolating pathogen)	uncertainty in final sample size
rotating panel	each site is sampled x number of times and then the next site is sampled x times	infer fine-resolution temporal dynamics efficiently over space	few longitudinal samples from any one population, can modify system if sampling of given sites is too frequent
augmented serial panel	increases the between-site interval from a rotating panel design, adds a longitudinal study for one site	higher power for trend detection, longitudinal analysis possible, less likely to modify system	may require greater effort in terms of time and funding
partially augmented serial	replaces the longitudinal sampling of the augmented serial panel with repeated sampling of multiple sites	higher power for trend detection, longitudinal analysis possible, minimized bias	may require greater effort in terms of time and funding

^aSee box 1 for references.

persist for long periods relative to the lifespan of individual hosts, seroprevalence can remain relatively stable over time, even if pathogen prevalence oscillates or the pathogen is extinct locally [80].

Longitudinal sampling of even a small number of known animals over the course of their infections can place preliminary constraints on disease process parameters, which in turn may prove useful for identifying the duration of infection and immunity with stratified or adaptive sampling of populations. This strategy has been used to study wildlife diseases in diverse hosts, from bighorn sheep [43] to bats [81,82]), and is essential for elucidating the within-host dynamics of poorly understood bat viruses [23].

One may identify spatial extents and resolutions for investigation of spillover risk by focusing early sampling at invasion fronts, as suggested for non-native invasive species [9,83,84]. Informal adaptive sampling often is employed following spillover events, but it would be valuable to use formal adaptive sampling [54–58]. Focused sampling at invasion fronts facilitates explicit estimation of transmission, recovery and disease-induced mortality rates before herd

immunity shapes dynamics. Moreover, higher public health burdens are often observed at the invasion front because epidemic curves in the reservoir hosts peak at those fronts, exerting high pathogen pressure. Moreover, human populations at invasion fronts rarely are well prepared to reduce spillover [11]. Sampling at the invasion front can be informed by an iterative process of data assessment, dynamic modelling, spatial and temporal forecasting and model validation [60]. Adaptive sampling, which complements random or random stratified sampling across space and time with focused sampling (e.g. in the region and months following a spillover event), also could be informative, but rarely has been implemented [56,85–87].

(c) Characterization of spatial and temporal dynamics

If there is no recent outbreak epicentre, various statistical approaches can be used to estimate the spatial and temporal structure of a pathogen to inform the sampling design. At most scales of observation, infection dynamics have some level of spatial and temporal dependence that decreases




Figure 3. Allocation of sampling effort over time (months) and space (e.g. along a latitudinal gradient) and consequences for inference of pathogen prevalence. Grey shading in panel (a) denotes the underlying spatial and temporal pattern in infection prevalence (the kernel-smoothed intensity of a random realization of a binomial point process). Open circles represent sampling locations. Colours indicate sampling location. Panel (b) illustrates the observed and true temporal trends in infection prevalence across sampling sites. Thin lines indicate the known infection prevalence over the annual cycle, whereas filled circles are estimated prevalence values given each design and ignoring error in estimation of the prevalence. Thick lines indicate the observed time-series of infection prevalence and only connect points from a single location; locations that are only sampled once within a design have no corresponding thick line. Heuristic sampling designs are as follows: opportunistic (A), single longitudinal (B), replicated longitudinal (C), randomized (D), random stratified (E), adaptive (F), rotating (G), augmented, serially alternating (H) and partially augmented, serially alternating (I). Sampling effort is held relatively constant across A to F and G to I (for which designs can require higher sampling effort). Methods for the simulations are given in electronic supplementary material, Methods.

with distance or time. Understanding the extent of spatial and temporal autocorrelation can inform the allocation of sampling effort [59,73,74]; this can be accomplished formally through exploratory tools such as correlograms (e.g. plotting the between-site correlation as a function of between-site distance) or semivariograms, for which semivariance is modelled as a function of spatial and temporal lags [88]. Pairs or clusters of samples within regions can be compared with samples among regions to estimate spatial autocorrelation. For example, Tobin [84] demonstrated that within a given spatial extent, sampling clusters of points estimated spatial autocorrelation more accurately than sampling evenly distributed but distant points.

The larger the extent of spatial dependence, the further the sampling units (e.g. sampled populations) can be from each other to make general inferences about risk across a landscape. For example, systems with little spatial variability—e.g. highly connected systems and those in which the infectious period is long relative to the lifespan of the host (e.g. figure 2c,d)—require sampling of a small number of spatially disparate populations. In this case, more frequent sampling of one or two locations may be more informative than infrequent sampling of many locations. By contrast, spatially asynchronous infections may require infrequent sampling of many locations across the ranges of the pathogen and host.

In theory, the lower the temporal variance, the less frequently one needs to sample. Surveillance of the temporal dynamics of infection often is systematic, with sampling at regular intervals within single or multiple populations, and this is often dictated by logistical and funding constraints. For example, the Soay sheep (*Ovis aries*) population of St Kilda, Scotland, UK was sampled annually to reveal

long-term fluctuations in parasite prevalence [64], and vampire bats (*Desmodus rotundus*) were sampled annually and biannually to reveal endemic viral and bacterial pathogens [65,66]. However, systematic sampling of a population in which prevalence is either extrinsically driven and seasonal or endogenous and epidemic may not detect temporal peaks in prevalence, particularly if the interval between samples is similar to or longer than the periodicity in the pathogen cycles or the peaks in epidemics. Random or rotating sampling designs can reduce the likelihood of this type of bias (box 1 and table 2). When temporal variance is high, as in cases with seasonal oscillations or multiple-year peaks, the sampling interval should reflect the periodicity of the disease. In such cases, generalized random stratified approaches may be used to avoid the pitfalls of systematic sampling and the clumping of simple random sampling [49].

The most challenging pathogens to sample, and therefore those that require the most-intense sampling, are those with highly localized infection dynamics [38]. If only one population is sampled, one erroneously might infer that all populations have similar dynamics. Characterization of infection structure in these systems is best captured with random, rotating or augmented sampling (box 1 and figure 3).

4. Conclusion

Uncertainty in predictions of spillover risk is reduced by knowledge of the spatial and temporal distribution of infection among populations of reservoir hosts. Financial and logistical constraints often force one to make inferences on the basis of small sample sizes and wide confidence intervals, and to

make trade-offs between spatial and temporal replication. Adopting wise design choices that are appropriate to the background dynamics of a particular system can extend the utility of even sparse data, and is essential to efficiently understand prevalence dynamics in reservoir host populations.

Data accessibility. This article has no additional data.

Authors' contributions. All authors wrote the first draft and contributed to writing and editing the manuscript. D.J.B., K.R.M. and R.K.P. made the figures.

Competing interests. We have no competing interests.

Acknowledgements. We thank Megan Higgs, Paul Cross and Kathi Irvine for comments on the manuscript, and Jamie Lloyd-Smith and Peter Hudson for conversations that inspired the manuscript.

Funding. This work was funded by the Defense Advanced Research Projects Agency (Young Faculty Award D16AP00113 and the DARPA PREEMPT Cooperative Agreement no. D18AC00031), US National Science Foundation (DEB-1716698), US National Institute of General Medical Sciences of the US National Institutes of Health (P20GM103474 and P30GM110732), and the USDA National Institute of Food and Agriculture (Hatch project 1015891). D.J.B. was also supported by an appointment to the Intelligence Community Postdoctoral Research Fellowship Program at Indiana University, administered by Oak Ridge Institute for Science and Education through an interagency agreement between the US Department of Energy and the Office of the Director of National Intelligence. The content of the information does not necessarily reflect the position or the policy of the US government, and no official endorsement should be inferred.

References

- Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, Lloyd-Smith JO. 2017 Pathways to zoonotic spillover. *Nat. Rev. Microbiol.* **15**, 502–510. (doi:10.1038/nrmicro.2017.45)
- Kuiken T, Leighton F, Fouchier R, LeDuc J, Peiris J, Schudel A, Stöhr K, Osterhaus AD. 2005 Pathogen surveillance in animals. *Science* **309**, 1680–1681. (doi:10.1126/science.1113310)
- Viana M, Mancy R, Biek R, Cleaveland S, Cross PC, Lloyd-Smith JO, Haydon DT. 2014 Assembling evidence for identifying reservoirs of infection. *Trends Ecol. Evol.* **29**, 270–279. (doi:10.1016/j.tree.2014.03.002)
- Bean AG, Baker ML, Stewart CR, Cowled C, Deffrasnes C, Wang L-F, Lowenthal JW. 2013 Studying immunity to zoonotic diseases in the natural host—keeping it real. *Nat. Rev. Immunol.* **13**, 851. (doi:10.1038/nri3551)
- Stallknecht DE. 2007 Impediments to wildlife disease surveillance, research, and diagnostics. In *Wildlife and emerging zoonotic diseases: the biology, circumstances and consequences of cross-species transmission*, pp. 445–461. Berlin: Springer.
- Giles JR, Peel AJ, Wells K, Plowright RK, McCallum H, Restif O. 2018 Optimizing non-invasive sampling of an infectious bat virus. *bioRxiv* (doi:10.1101/401968)
- Plowright RK, Becker DJ, Crowley DE, Washburne AD, Huang T, Nameer PO, Gurley ES, Han BA. 2019 Prioritizing surveillance of Nipah virus in India. *PLoS Negl. Trop. Dis.* **13**, e0007393. (doi:10.1371/journal.pntd.0007393)
- East I, Wicks R, Martin P, Sergeant E, Randall L, Garner MJ. 2013 Use of a multi-criteria analysis framework to inform the design of risk based general surveillance systems for animal disease in Australia. *Prev. Vet. Med.* **112**, 230–247. (doi:10.1016/j.prevetmed.2013.09.012)
- Williams PJ, Hooten MB, Womble JN, Esslinger GG, Bower MR. 2018 Monitoring dynamic spatio-temporal ecological processes optimally. *Ecology* **99**, 524–535. (doi:10.1002/ecy.2120)
- Rohani P, Breban R, Stallknecht DE, Drake JM. 2009 Environmental transmission of low pathogenicity avian influenza viruses and its implications for pathogen invasion. *Proc. Natl Acad. Sci. USA* **106**, 10 365–10 369. (doi:10.1073/pnas.0809026106)
- Kilpatrick AM, Randolph SE. 2012 Drivers, dynamics, and control of emerging vector-borne zoonotic diseases. *Lancet* **380**, 1946–1955. (doi:10.1016/S0140-6736(12)61151-9)
- Terzian ACB *et al.* 2018 Evidence of natural Zika virus infection in neotropical non-human primates in Brazil. *Sci. Rep.* **8**, 16034. (doi:10.1038/s41598-018-34423-6)
- Diekmann O, Heesterbeek J, Metz JA. 1990 On the definition and the computation of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous populations. *J. Math. Biol.* **28**, 365–382. (doi:10.1007/BF00178324)
- Keeling MJ, Gilligan CA. 2000 Metapopulation dynamics of bubonic plague. *Nature* **407**, 903–906. (doi:10.1038/35038073)
- Vogler AJ, Chan F, Nottingham R, Andersen G, Drees K, Beckstrom-Sternberg SM, Wagner DM, Chanteau S, Lenski R. 2013 A decade of plague in Mahajanga, Madagascar: insights into the global maritime spread of pandemic plague. *mBio* **4**, e00623-12. (doi:10.1128/mBio.00623-12)
- O'Brien DJ, Schmitt SM, Fitzgerald SD, Berry DE, Hickling GJ. 2006 Managing the wildlife reservoir of *Mycobacterium bovis*: the Michigan, USA, experience. *Vet. Microbiol.* **112**, 313–323. (doi:10.1016/j.vetmic.2005.11.014)
- Hudson PJ, Rizzoli P, Grenfell BT, Heesterbeek JAP, Dobson AP. 2002 *The ecology of wildlife diseases*. Oxford, UK: Oxford University Press.
- Ostfeld RS, Keesing F, Eviner VT. 2010 *Infectious disease ecology: effects of ecosystems on disease and of disease on ecosystems*. Princeton, NJ: Princeton University Press.
- Keeling MJ, Rohani P. 2008 *Modeling infectious diseases in humans and animals*. Princeton, NJ: Princeton University Press.
- Amman BR *et al.* 2012 Seasonal pulses of Marburg virus circulation in juvenile *Rousettus aegyptiacus* bats coincide with periods of increased risk of human infection. *PLoS Pathog.* **8**, e1002877. (doi:10.1371/journal.ppat.1002877)
- Bernshtein A, Apekina N, Mikhailova T, Myasnikov YA, Khlyap L, Korotkov YS, Gavrilovskaya IN. 1999 Dynamics of Puumala hantavirus infection in naturally infected bank voles (*Clethrionomys glareolus*). *Arch. Virol.* **144**, 2415–2428. (doi:10.1007/s007050050654)
- Lass S, Hudson PJ, Thakar J, Saric J, Harvill E, Albert R, Perkins SE. 2013 Generating super-shedders: co-infection increases bacterial load and egg production of a gastrointestinal helminth. *J. R. Soc. Interface* **10**, 20120588. (doi:10.1098/rsif.2012.0588)
- Plowright RK, Peel AJ, Streicker DG, Gilbert AT, McCallum H, Wood J, Baker ML, Restif O. 2016 Transmission or within-host dynamics driving pulses of zoonotic viruses in reservoir–host populations. *PLoS Negl. Trop. Dis.* **10**, e0004796. (doi:10.1371/journal.pntd.0004796)
- Plowright RK *et al.* 2015 Ecological dynamics of emerging bat virus spillover. *Proc. R. Soc. B* **282**, 20142124. (doi:10.1098/rspb.2014.2124)
- Kessler MK *et al.* 2018 Changing resource landscapes and spillover of henipaviruses. *Ann. NY Acad. Sci.* **1429**, 79–99. (doi:10.1111/nyas.13910)
- Field H *et al.* 2015 Spatiotemporal aspects of Hendra virus infection in pteropid bats (flying-foxes) in eastern Australia. *PLoS ONE* **10**, e0144055. (doi:10.1371/journal.pone.0144055)
- Tobler WR. 1970 A computer movie simulating urban growth in the Detroit region. *Econ. Geogr.* **46**, 234–240. (doi:10.2307/143141)
- Walter JA, Sheppard LW, Anderson TL, Kastens JH, Bjørnstad ON, Liebhold AM, Reuman DC. 2017 The geography of spatial synchrony. *Ecol. Lett.* **20**, 801–814. (doi:10.1111/ele.12782)
- Lowen AC, Mubareka S, Steel J, Palese P. 2007 Influenza virus transmission is dependent on relative humidity and temperature. *PLoS Pathog.* **3**, e151. (doi:10.1371/journal.ppat.0030151)
- Martin G, Plowright R, Chen C, Kault D, Selleck P, Skerratt LF. 2015 Hendra virus survival does not explain spillover patterns and implicates relatively

- direct transmission routes from flying foxes to horses. *J. Gen. Virol.* **96**, 1229–1237. (doi:10.1099/vir.0.000073)
31. Luby SP *et al.* 2009 Recurrent zoonotic transmission of Nipah virus into humans, Bangladesh, 2001–2007. *Emerg. Infect. Dis.* **15**, 1229. (doi:10.3201/eid1508.081237)
 32. Bjornstad ON. 2018 *Seasonality. Epidemics: models and data using R*, pp. 81–94. Cham, Switzerland: Springer International Publishing.
 33. Cross PC, Lloyd-Smith JO, Johnson PLF, Getz WM. 2005 Duelling timescales of host movement and disease recovery determine invasion of disease in structured populations. *Ecol. Lett.* **8**, 587–595. (doi:10.1111/j.1461-0248.2005.00760.x)
 34. Medley GF, Lindop NA, Edmunds WJ, Nokes DJ. 2001 Hepatitis-B virus endemicity: heterogeneity, catastrophic dynamics and control. *Nat. Med.* **7**, 619–624. (doi:10.1038/87953)
 35. Sah P, Leu ST, Cross PC, Hudson PJ, Bansal S. 2017 Unraveling the disease consequences and mechanisms of modular structure in animal social networks. *Proc. Natl Acad. Sci. USA* **14**, 4165–4170. (doi:10.1073/pnas.1613616114)
 36. Bharti N, Djibo A, Ferrari M, Grais R, Tatem A, McCabe CA, Bjornstad ON, Grenfell BT. 2010 Measles hotspots and epidemiological connectivity. *Epidemiol. Infect.* **138**, 1308–1316. (doi:10.1017/S0950268809991385)
 37. Grenfell BT, Bjornstad ON, Kappey J. 2001 Travelling waves and spatial hierarchies in measles epidemics. *Nature* **414**, 716–723. (doi:10.1038/414716a)
 38. Plowright RK, Foley P, Field HE, Dobson AP, Foley JE, Eby P, Daszak P. 2011 Urban habituation, ecological connectivity and epidemic dampening: the emergence of Hendra virus from flying foxes (*Pteropus* spp.). *Proc. R. Soc. B* **278**, 3703–3712. (doi:10.1098/rspb.2011.0522)
 39. Cattadori IM, Haydon DT, Hudson PJ. 2005 Parasites and climate synchronize red grouse populations. *Nature* **433**, 737–741. (doi:10.1038/nature03276)
 40. Hudson PJ, Cattadori IM, Boag B, Dobson AP. 2006 Climate disruption and parasite–host dynamics: patterns and processes associated with warming and the frequency of extreme climatic events. *J. Helminthol.* **80**, 175–782. (doi:10.1079/JOH2006357)
 41. Peel A, Eby P, Kessler M, Lunn T, Breed A, Plowright R. 2017 Hendra virus spillover risk in horses: heightened vigilance and precautions being urged this winter. *Aust. Vet. J.* **95**, N20–N21.
 42. Plowright RK, Sokolow SH, Gorman ME, Daszak P, Foley JE. 2008 Causal inference in disease ecology: investigating ecological drivers of disease emergence. *Front. Ecol. Environ.* **6**, 420–429. (doi:10.1890/070086)
 43. Plowright RK, Manlove K, Besser TE, Paez D, Andrews KR, Matthews PE, Waits LP, Hudson PJ, Cassirer EF. 2017 Age-structured variation in infectious period explains epidemiological features of pneumonia in bighorn sheep. *Ecol. Lett.* **20**, 1325–1336. (doi:10.1111/ele.12829)
 44. Cassirer EF *et al.* 2017 Pneumonia in bighorn sheep: risk and resilience. *J. Wildl. Manag.* **82**, 32–45. (doi:10.1002/jwmg.21309)
 45. Cassirer EF, Manlove KR, Plowright RK, Besser TE. 2016 Evidence for strain-specific immunity to pneumonia in bighorn sheep. *J. Wildl. Manag.* **81**, 133–143. (doi:10.1002/jwmg.21172)
 46. Almberg ES, Cross PC, Smith DW. 2010 Persistence of canine distemper virus in the Greater Yellowstone Ecosystem's carnivore community. *Ecol. Appl.* **20**, 2058–2074. (doi:10.1890/09-1225.1)
 47. Smith AN, Anderson MJ, Pawley MD. 2017 Could ecologists be more random? Straightforward alternatives to haphazard spatial sampling. *Ecography* **40**, 1251–1255. (doi:10.1111/ecog.02821)
 48. O'Shea TJ, Ellison LE, Stanley TR. 2004 Survival estimation in bats: historical overview, critical appraisal, and suggestions for new approaches. In *Sampling rare or elusive species: concepts, designs, and techniques for estimating population parameters* (ed. W Thompson), pp. 297–336. Washington, DC: Island Press.
 49. Stevens Jr DL, Olsen AR. 2004 Spatially balanced sampling of natural resources. *J. Am. Stat. Assoc.* **99**, 262–278. (doi:10.1198/016214504000000250)
 50. Yoccoz NG, Nichols JD, Boulinier T. 2001 Monitoring of biological diversity in space and time. *Trends Ecol. Evol.* **16**, 446–453. (doi:10.1016/S0169-5347(01)02205-4)
 51. Nusser SM, Clark WR, Otis DL, Huang L. 2008 Sampling considerations for disease surveillance in wildlife populations. *J. Wildl. Manag.* **72**, 52–60. (doi:10.2193/2007-317)
 52. Ver Hoef JM. 2008 Spatial methods for plot-based sampling of wildlife populations. *Environ. Ecol. Stat.* **15**, 3–13. (doi:10.1007/s10651-007-0035-y)
 53. Morner T, Obendorf D, Artois M, Woodford M. 2002 Surveillance and monitoring of wildlife diseases. *Rev. Sci. Tech.* **21**, 67–76. (doi:10.20506/rst.21.1.1321)
 54. Thompson SK. 1991 Adaptive cluster sampling: designs with primary and secondary units. *Biometrics* **47**, 1103–1115. (doi:10.2307/2532662)
 55. Thompson SK. 1991 Stratified adaptive cluster sampling. *Biometrika* **78**, 389–397. (doi:10.1093/biomet/78.2.389)
 56. Thompson SK. 1990 Adaptive cluster sampling. *J. Am. Stat. Assoc.* **85**, 1050–1059. (doi:10.1080/01621459.1990.10474975)
 57. Smith DR, Conroy MJ, Brakhage DH. 1995 Efficiency of adaptive cluster sampling for estimating density of wintering waterfowl. *Biometrics* **51**, 777–788. (doi:10.2307/2532964)
 58. Conroy MJ, Runge JP, Barker RJ, Schofield MR, Fonnesebeck CJ. 2008 Efficient estimation of abundance for patchily distributed populations via two-phase, adaptive sampling. *Ecology* **89**, 3362–3370. (doi:10.1890/07-2145.1)
 59. Urquhart NS, Kincaid TM. 1999 Designs for detecting trend from repeated surveys of ecological resources. *J. Agric. Biol. Environ. Stat.* **4**, 404–414. (doi:10.2307/1400498)
 60. R Development Core Team. 2006 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. See <http://www.R-project.org/>.
 61. Baddeley A, Rubak E, Turner R. 2015 *Spatial point patterns: methodology and applications with R*. Boca Raton, FL: CRC Press.
 62. Guan Y *et al.* 2003 Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. *Science* **302**, 276–278. (doi:10.1126/science.1087139)
 63. Young PL, Halpin K, Selleck PW, Field H, Gravel JL, Kelly MA, Mackenzie JS. 1996 Serologic evidence for the presence in *Pteropus* bats of a paramyxovirus related to equine morbillivirus. *Emerg. Infect. Dis.* **2**, 239–240. (doi:10.3201/eid0203.960315)
 64. Graham A *et al.* 2016 Exposure to viral and bacterial pathogens among Soay sheep (*Ovis aries*) of the St Kilda archipelago. *Epidemiol. Infect.* **144**, 1879–1888. (doi:10.1017/S0950268816000017)
 65. Streicker DG *et al.* 2012 Ecological and anthropogenic drivers of rabies exposure in vampire bats: implications for transmission and control. *Proc. R. Soc. B* **279**, 3384–3392. (doi:10.1098/rspb.2012.0538)
 66. Volokhov D, Becker D, Bergner L, Camus M, Orton R, Chizhikov V, Altizer SM, Streicker DG. 2017 Novel hemotropic mycoplasmas are widespread and genetically diverse in vampire bats. *Epidemiol. Infect.* **145**, 3154–3167. (doi:10.1017/S095026881700231X)
 67. Gorsich EE, Ezenwa VO, Cross PC, Bengis RG, Jolles AE. 2015 Context-dependent survival, fecundity and predicted population-level consequences of brucellosis in African buffalo. *J. Anim. Ecol.* **84**, 999–1009. (doi:10.1111/1365-2656.12356)
 68. Becker DJ, Crowley DE, Washburne AD, Plowright RK. 2019 Temporal and spatial limitations in global surveillance for bat filoviruses and henipaviruses. *bioRxiv*, 674655. (doi:10.1101/674655)
 69. Li W *et al.* 2005 Bats are natural reservoirs of SARS-like coronaviruses. *Science* **310**, 676–679. (doi:10.1126/science.1118391)
 70. O'Brien C, Van Riper III C, Myers DE. 2009 Making reliable decisions in the study of wildlife diseases: using hypothesis tests, statistical power, and observed effects. *J. Wildl. Dis.* **45**, 700–712. (doi:10.7589/0090-3558-45.3.700)
 71. Dzul MC, Dixon PM, Quist MC, Dinsmore SJ, Bower MR, Wilson KP, Gaines DB. 2013 Using variance components to estimate power in a hierarchically nested sampling design. *Environ. Monit. Assess.* **185**, 405–414. (doi:10.1007/s10661-012-2562-8)
 72. Artois M *et al.* 2009 Wildlife disease surveillance and monitoring. In *Management of disease in wild mammals* (eds RJ Delahay, GC Smith, MR Hutchings), pp. 187–213. New York, NY: Springer.
 73. Fuller WA. 1999 Environmental surveys over time. *J. Agric. Biol. Environ. Stat.* **4**, 331–345. (doi:10.2307/1400493)
 74. McDonald TL. 2003 Review of environmental monitoring methods: survey designs. *Environ. Monit. Assess.* **85**, 277–292. (doi:10.1023/A:1023954311636)

75. Restif O *et al.* 2012 Model-guided study design: a practical framework for the design and analysis of multi-disciplinary ecological studies. *Ecol. Lett.* **15**, 1083–1094. (doi:10.1111/j.1461-0248.2012.01836.x)
76. Peel AJ *et al.* 2012 Henipavirus neutralising antibodies in an isolated island population of African fruit bats. *PLoS ONE* **7**, e30346. (doi:10.1371/journal.pone.0030346)
77. Peel AJ, Baker KS, Hayman DT, Broder CC, Cunningham AA, Fooks AR, Garnier R, Wood JL, Restif O. 2018 Support for viral persistence in bats from age-specific serology and models of maternal immunity. *Sci. Rep.* **8**, 3859. (doi:10.1038/s41598-018-22236-6)
78. Plowright RK, Field HE, Smith C, Divljan A, Palmer C, Tabor GM, Daszak P, Foley JE. 2008 Reproduction and nutritional stress are risk factors for Hendra virus infection in little red flying foxes (*Pteropus scapulatus*). *Proc. R. Soc. B* **275**, 861–869. (doi:10.1098/rspb.2007.1260)
79. Gilbert AT *et al.* 2013 Deciphering serology to understand the ecology of infectious diseases in wildlife. *EcoHealth* **10**, 298–313. (doi:10.1007/s10393-013-0856-0)
80. Cross PC, Caillaud D, Heisey DM. 2013 Underestimating the effects of spatial heterogeneity due to individual movement and spatial scale: infectious disease as an example. *Landsc. Ecol.* **28**, 247–257. (doi:10.1007/s10980-012-9830-4)
81. Glennon EE *et al.* 2019 What is stirring in the reservoir? Modelling mechanisms of henipavirus circulation in fruit bat hosts. *Phil. Trans. R. Soc. B* **374**, 20190021. (doi:10.1098/rstb.2019.0021)
82. Brook CE *et al.* 2019 Disentangling serology to elucidate henipa- and filovirus transmission in Madagascar fruit bats. *J. Anim. Ecol.* **88**, 1001–1016. (doi:10.1111/1365-2656.12985)
83. Bjørnstad ON, Liebhold AM, Johnson DM. 2008 Transient synchronization following invasion: revisiting Moran's model and a case study. *Popul. Ecol.* **50**, 379–389. (doi:10.1007/s10144-008-0105-5)
84. Tobin PC. 2004 Estimation of the spatial autocorrelation function: consequences of sampling dynamic populations in space and time. *Ecography* **27**, 767–775. (doi:10.1111/j.0906-7590.2004.03977.x)
85. Flanagan ML, Leighton TJ, Dudley JP. 2011 *Anticipating viral species jumps: bioinformatics and data needs*. State College, PA: Pennsylvania State University.
86. Kabaghe AN, Chipeta MG, McCann RS, Phiri KS, Van Vugt M, Takken W, Diggle P, Terlouw AD. 2017 Adaptive geostatistical sampling enables efficient identification of malaria hotspots in repeated cross-sectional surveys in rural Malawi. *PLoS ONE* **12**, e0172266. (doi:10.1371/journal.pone.0172266)
87. Chipeta MG, Terlouw DJ, Phiri KS, Diggle PJ. 2016 Adaptive geostatistical design and analysis for prevalence surveys. *Spatial Stat.* **15**, 70–84. (doi:10.1016/j.spasta.2015.12.004)
88. Isaaks EH, Srivastava MR. 1989 *An introduction to applied geostatistics*. New York, NY: Oxford University Press.