




Translation of Diverse Aramid- and 1,3-Dicarbonyl-peptides by Wild Type Ribosomes *in Vitro*

Omer Ad,[†] Kyle S. Hoffman,[‡] Andrew G. Cairns,[†] Aaron L. Featherston,[†] Scott J. Miller,^{*,†} Dieter Söll,^{*,‡} and Alanna Schepartz^{*,†,‡}

[†]Department of Chemistry and [‡]Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, Connecticut 06520, United States

Supporting Information

ABSTRACT: Here, we report that wild type *Escherichia coli* ribosomes accept and elongate precharged initiator tRNAs acylated with multiple benzoic acids, including aramid precursors, as well as malonyl (1,3-dicarbonyl) substrates to generate a diverse set of aramid-peptide and polyketide-peptide hybrid molecules. This work expands the scope of ribozyme- and ribosome-catalyzed chemical transformations, provides a starting point for *in vivo* translation engineering efforts, and offers an alternative strategy for the biosynthesis of polyketide-peptide natural products.



INTRODUCTION

As far as we know, ribosomes have evolved for billions of years to perform a single reaction—formation of an amide bond between two α -amino acid substrates brought into proximity by tRNAs within the ribosome active site, the peptidyl transferase center (PTC). In cells and extracts, the chemistry possible within a wild type ribosome PTC has expanded to include reactions of more than 200 different nonproteinogenic α -amino and hydroxy acids;^{1–4} ribosomes containing remodeled PTCs support amide bond formation to and from a small number of β -amino acids^{5–7} and dipeptides^{8,9} with limited efficiency. The combination of cell-free *in vitro* translation systems and ribozyme-catalyzed tRNA acylation reactions offers the opportunity for even greater reaction diversity, including the introduction of multiple *N*-alkyl,¹⁰ *D*- α ,^{11,12} α -hydroxy,¹³ and β -amino acids,^{14,15} the precursors of β -peptide foldamers.^{16–19}

A second family of foldamer-like molecules are aramids, oligomers of substituted aminobenzoic acids.²⁰ Aramids possess remarkably varied properties. Kevlar, a polymer of 1,4-phenylenediamine and terephthaloyl chloride, is a strong and heat-resistant fiber,²¹ whereas cystobactamids are DNA gyrase inhibitors active against Gram-negative bacteria.²² Many other aramid foldamers with diverse and significant properties have been reported.^{23–25} Recently, wild type *E. coli* ribosomes were shown to accept and elongate initiator tRNAs precharged with aromatic foldamer-dipeptide appendages.²⁶ Notably, in this case the foldamer monomers did not themselves react within the PTC, being displaced from the reaction center by a Gly–Phe dipeptide spacer.^{26,27} Here, we report that wild type *E. coli* ribosomes accept and elongate precharged initiator tRNAs acylated directly with multiple benzoic acids, including

aramid precursors, as well as malonyl (1,3-dicarbonyl) substrates. The result is a diverse set of aramid-peptide and polyketide-peptide hybrid molecules. This work provides new knowledge about the generality of flexizyme-promoted tRNA acylation reactions, expands the scope of ribosome-catalyzed chemical transformations, provides a starting point for *in vivo* translation engineering efforts, and offers an alternative strategy for biosynthesis of polyketide-peptide natural products.

RESULTS AND DISCUSSION

As the first step toward the ribosomal synthesis of aramid-like peptides, we made use of an established microhelix (MH) gel-shift assay²⁸ and high-resolution mass spectrometry (Figure 1A) to evaluate whether the cyanomethyl esters of unsubstituted aminobenzoic acids were substrates for the flexizyme ribozyme eFx.²⁹ Incubation of cyanomethyl esters 1–3 (5 mM) with 25 μ M microhelix MH and 25 μ M eFx (Table S1) in bicine buffer at pH 9 for 48 h showed little or no evidence of MH acylation when the reaction products were evaluated on an acid-urea PAGE gel (Figure 1B). A low level of MH acylation by the *o*- and *om*-analogues 1 and 2 (and a trace with 3) could be observed using a highly sensitive RNase A/LC-HRMS assay³⁰ that detects the acylated adenine nucleoside (Figure 1C). We also investigated the extent of tRNA acylation using the alternative ribozyme dFx³¹ and the 1,3-dinitrobenzyl esters of *p*- and *o*-aminobenzoic acid (4 and 5, respectively, as shown in Scheme S1). These substrates also failed to yield the expected MH products when incubated with dFx under standard conditions¹⁴ and analyzed using acid-urea gels or

Received: May 9, 2019

Published: June 26, 2019

FLAG polypeptide was detected in the presence of DNA template and *m*-AN-tRNA.

Aminobenzoate esters hydrolyze exceptionally slowly,³⁴ suggesting that the electron-rich aromatic ring contributes to the low reactivity of 1–3. In addition, the structure of the ethyl ester of *L*-phenylalanine bound to Fx (as an Fx-tRNA fusion)³⁵ shows π -stacking between Fx base guanine 24 and the *L*-phenylalanine aromatic ring; this stacking would be less favorable with an electron-rich arene.^{36,37} To investigate whether reactivity in eFx-promoted reactions was correlated with arene electron density, we prepared a diverse set of substituted benzoic acid cyanomethyl esters (Figure 3A) and




Figure 3. Probing structure–activity relationships for cyanomethyl esters of substituted benzoic acids in eFx-promoted acylation reactions. (A) Substituted benzoic acid cyanomethyl esters studied herein. (B) Acid-urea gel-shift analysis of MH acylation by cyanomethyl esters 6 and 8–15 in the presence of ribozyme eFx. Yield was estimated by UV densitometry. (C) LC-HRMS analysis of MH acylation reactions containing cyanomethyl esters 6 and 8–15 after RNase A digestion. Exact masses are reported in Table S2.

evaluated the extent to which eFx reactivity correlated with the sign and magnitude of the relevant sigma factor, which measures the inductive effect of the aromatic substituent.³⁸ As expected, benzoic acid cyanomethyl esters possessing strong electron-withdrawing substituents, such as penta-fluoro 8, *p*-nitro 9 ($\sigma = +0.78$), or *p*-Cl 10 ($\sigma = +0.23$), were excellent eFx substrates in model MH reactions, with 78–99% yields (Figure 3B,C). However, other factors are clearly important: a benzoic

acid cyanomethyl ester possessing a weak electron-withdrawing substituent, such as *p*-azido 11 ($\sigma = +0.08$), was also an excellent substrate (yield of acylated MH = 74%), as were analogues possessing both strong and weak electron-donating substituents, such as *p*-methoxy 12 ($\sigma = -0.27$; yield of acylated MH = 62%) and *p*-methyl 13 ($\sigma = -0.17$; yield of acylated MH = 54%). Notably, the poorest yields were observed in eFx-promoted reactions of substrates 6 (yield of acylated MH = 25%) and 15 (yield of acylated MH = 23%), all of which contain one or more acidic protons/hydrogen bond donors, just like amino benzoic acids 1, 2, and 3. These results imply that the presence of hydrogen bond donors in certain positions contributes to the poor reactivity of amino benzoic acids 1–3. Consistent with this notion, *p*-hydroxybenzoic acid 16 [pKa = 8.3 (*p*-hydroxybenzoic acid methyl ester)] was a poor substrate, whereas alcohol 17 [pKa = 15 (benzyl alcohol)] and aldehyde 18 reacted well (Scheme S1 and Figure S4). It is possible that certain hydrogen bond donors alter the position of the aromatic ring in the eFx active site or coordinate and inactivate functional groups involved in catalysis. Determining the exact nature of these interactions is beyond the scope of this discussion but will be essential to effectively engineer new ribozymes that accept diverse foldamer building blocks *in vitro* and *in vivo*.

With a new set of aramid substrates in hand, we used the PURExpress Δ (aa, tRNA) *in vitro* translation kit to evaluate if initiator tRNAs acylated with diverse benzoic acids could be accommodated in the ribosomal P-site and initiate translation of an AR-VF-FLAG polypeptide carrying an aramid monomer (AR) at the N-terminus (Figure 4). Every benzoic acid cyanomethyl ester that acylated the microhelix MH with a yield >50% in an eFx-promoted reaction (Figure 3) was used to acylate fMetT, and translation reactions were performed and analyzed as described above. With one exception, every single AR-fMetT initiated translation of an AR-VF-FLAG peptide whose mass corresponded to incorporation of the prescribed substituted benzoic acid. The singular exception was *p*-azidobenzoic acid 11; in this case the mass of the isolated polypeptide was consistent with *in situ* reduction of the azide to an amine. These results demonstrate that diverse aramid-like monomers can be accommodated directly within the ribosomal P-site and act as acceptors for a natural α -amino acid in the A-site. They show further that use of *p*-azidobenzoic acid 11 effectively circumvents the poor reactivity of *p*-aminobenzoic acid 3 to generate a polypeptide with a *p*-aminobenzoic acid monomer at the N-terminus. The observation that wild type *E. coli* ribosomes can initiate translation using tRNAs acylated with diverse aramid-like monomers significantly expands the scope of *in vitro* translation reactions beyond that of Kawakami³⁹ and lays the initial groundwork for the biosynthesis of genetically encoded, sequence-defined polyaramid oligomers.

We next sought to evaluate the relative efficiency of PURExpress reactions initiated with differentially acylated fMetT derivatives. To begin, we monitored the yield of fMet-VF-FLAG (approximated as the extracted ion abundance) as a function of time in PURExpress Δ reactions supplemented with either 50 μ M precharged fMetT-fMet (charged using the dFx substrate fMet-DBE) or 50 μ M *L*-methionine. The bulk of both reactions was complete within 100 min, but the yield of fMet-VF-FLAG in reactions supplemented with precharged fMetT-fMet was 1.5% of that obtained in reactions supplemented with *L*-methionine (Figure S5A). Next, we




Figure 4. Initiator tRNAs acylated with diverse benzoic acids are accommodated in the ribosomal P-site and are elongated into AR-VF-FLAG polypeptides. LC-HRMS analysis of reaction products whose masses correspond to AR-VFDYKDDDDK (AR-VF-FLAG) polypeptides containing diverse substituted benzoic acid monomers.

compared the extracted ion abundance (after 30–90 min) of the AR-VF-FLAG peptide initiated with fMetT precharged with benzoic acid ester 8. The yield of this AR-VF-FLAG polypeptide (C6F5-VF-FLAG) was 25–30% of the yield of fMet-VF-FLAG (generated in reactions supplemented with precharged fMetT-fMet) (Figure S5B) and within the range observed when translation was initiated with fMetT precharged with natural amino acids.⁴⁰ The relative yields of AR-VF-FLAG peptides initiated with other precharged fMetT derivatives were also comparable (Figure S5D), suggesting similar initiation efficiencies. We note that when ValT was precharged with β -Phe, the yield of fMet- β -Phe-F-FLAG was 5-fold higher than the yield of fMet-VF-FLAG generated with precharged fMetT (Figure S5C). As initiation complex assembly is the rate limiting step during translation,⁴¹ the higher yield of fMet- β -Phe-F-FLAG relative to fMet-VF-FLAG is likely due to the difficulty assembling the translation initiation complex using non-natural fMetT derivatives. Benzoic acid monomers that were poor eFx substrates (yields <50%) in model MH reactions, such as 6 and 15 (Figure 3), failed to detectably initiate peptide synthesis from WT ribosomes *in vitro*. This observation suggests that the ribosome is largely agnostic of

aramid structure, and that the concentration of non-natural fMetT derivative, rather than monomer structure, determines the reaction yield in PURExpress reactions.⁴²

Like aramid natural products,²² polyketide-peptide hybrid molecules are biosynthesized by mega-assemblies of complex protein enzymes;^{43–45} the combination of peptide and polyketide-based functionality can translate into highly unique biological functions.^{46–48} To evaluate whether wild type *E. coli* ribosomes are capable of biosynthesizing a polyketide-peptide hybrid, we prepared malonate derivatives 19–23 (Figure 5A).




Figure 5. Wild type *E. coli* ribosomes support the biosynthesis of polyketide-peptide hybrid molecules. (A) Malonic esters 19–23 evaluated as substrates for eFx or dFx. (B) Acid-urea gel-shift analysis of MH acylation by esters 19–23 in the presence of eFx (19, 22) or dFx (20, 21, 23). Yield was estimated by UV densitometry. (C) LC-HRMS analysis of MH acylation reactions containing esters 19–23 after RNase A digestion. (D) LC-HRMS analysis of reaction products whose masses corresponds to Mal-VFDYKDDDDK (Mal-VF-FLAG) polypeptides containing methyl and nitrobenzyl malonates 23 and 22. ND = not determined due to lack of separation from unacylated microhelix. Exact masses are reported in Table S2.

Model microhelix MHacylation reactions were analyzed using acid-urea gels (Figure 5B) and RNase A/LC-HRMS (Figure 5C) as described above. Although the malonic acid half esters 19, 20, and 21 were poor substrates for the requisite Fx analogue, cyanomethyl ester 22 was a moderate substrate, acylating the acylated MH in 40% yield. Although no gel-shift was observed in the eFx-promoted MH acylation reaction of cyanomethyl ester 23 (perhaps because of low molecular weight and/or polarity),⁴⁹ strong evidence for reaction was observed using RNase A/LC-HRMS (Figure 5C). Indeed, the addition of fMetT derivatives acylated with 22 and 23 (50–100 μM) to PURExpress Δ (aa, tRNA) *in vitro* translation

reactions led to the isolation of polypeptides carrying malonates **22** and **23** (22-VF-FLAG and 23-VF-FLAG, respectively), whose masses were confirmed by LC-HRMS (Figure S5D). The yield of 23-VF-FLAG, estimated as described above, was approximately 20% of the yield of fMet-VF-FLAG produced in reactions supplemented with precharged fMetT (Figure S5A). We conclude that extant *E. coli* ribosomes have the capacity to biosynthesize simple polyketide-peptide hybrid molecules.

In summary, here we report that wild type *E. coli* ribosomes accept precharged initiator tRNAs acylated with multiple substituted benzoic acids, including the monomeric unit of Kevlar, as well as malonyl (1,3-dicarbonyl) substrates. The ribosome then elongates these substrates to generate a diverse set of aramid-peptide and polyketide-peptide hybrid molecules. This work expands the scope of reactions catalyzed by both flexizyme and wild type ribosomes, provides a starting point for *in vivo* translation engineering efforts, and offers an alternative strategy for biosynthesis of polyketide-peptide natural products.

Safety Statement. No unexpected or unusually high safety hazards were encountered during the execution of these experiments.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acscentsci.9b00460.

Synthesis and characterization of flexizyme substrates; and procedures for formation, characterization, purification, and analysis of tRNA acylation and IVT reactions (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: Scott.Miller@yale.edu.

*E-mail: Dieter.Söll@yale.edu.

*E-mail: Alanna.Schepartz@yale.edu.

ORCID

Aaron L. Featherston: 0000-0003-0918-067X

Scott J. Miller: 0000-0001-7817-1318

Dieter Söll: 0000-0002-3077-8986

Alanna Schepartz: 0000-0003-2127-3932

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the Center for Genetically Encoded Materials, an NSF Center for Chemical Innovation (NSF CHE-1740549). O.A. was supported in part by Agilent Technologies as an Agilent Fellow.

■ REFERENCES

- (1) Guo, J.; Wang, J.; Anderson, J. C.; Schultz, P. G. Addition of an α -hydroxy acid to the genetic code of bacteria. *Angew. Chem., Int. Ed.* **2008**, *47*, 722–725.
- (2) Chin, J. W. Expanding and reprogramming the genetic code. *Nature* **2017**, *550*, 53–60.
- (3) Vargas-Rodriguez, O.; Sevostyanova, A.; Söll, D.; Crnković, A. Upgrading aminoacyl-tRNA synthetases for genetic code expansion. *Curr. Opin. Chem. Biol.* **2018**, *46*, 115–122.
- (4) Young, D. D.; Schultz, P. G. Playing with the molecules of life. *ACS Chem. Biol.* **2018**, *13*, 854–870.
- (5) Maini, R.; Nguyen, D. T.; Chen, S.; Dedkova, L. M.; Chowdhury, S. R.; Alcalá-Torano, R.; Hecht, S. M. Incorporation of β -amino acids into dihydrofolate reductase by ribosomes having modifications in the peptidyltransferase center. *Bioorg. Med. Chem.* **2013**, *21*, 1088–1096.
- (6) Maini, R.; Chowdhury, S. R.; Dedkova, L. M.; Roy, B.; Daskalova, S. M.; Paul, R.; Chen, S.; Hecht, S. M. Protein synthesis with ribosomes selected for the incorporation of β -amino acids. *Biochemistry* **2015**, *54*, 3694–3706.
- (7) Melo Czekster, C.; Robertson, W. E.; Walker, A. S.; Söll, D.; Schepartz, A. *In vivo* biosynthesis of a β -amino acid-containing protein. *J. Am. Chem. Soc.* **2016**, *138*, 5194–5197.
- (8) Maini, R.; Dedkova, L. M.; Paul, R.; Madathil, M. M.; Chowdhury, S. R.; Chen, S.; Hecht, S. M. Ribosome-mediated incorporation of dipeptides and dipeptide analogues into proteins *in vitro*. *J. Am. Chem. Soc.* **2015**, *137*, 11206–11209.
- (9) Chen, S.; Ji, X.; Gao, M.; Dedkova, L. M.; Hecht, S. M. *In cellulo* synthesis of proteins containing a fluorescent oxazole amino acid. *J. Am. Chem. Soc.* **2019**, *141*, 5597–5601.
- (10) Subtelny, A. O.; Hartman, M. C. T.; Szostak, J. W. Ribosomal synthesis of *N*-methyl peptides. *J. Am. Chem. Soc.* **2008**, *130*, 6131–6136.
- (11) Dedkova, L. M.; Fahmi, N. E.; Golovine, S. Y.; Hecht, S. M. Enhanced D-amino acid incorporation into protein by modified ribosomes. *J. Am. Chem. Soc.* **2003**, *125*, 6616–6617.
- (12) Goto, Y.; Murakami, H.; Suga, H. Initiating translation with D-amino acids. *RNA* **2008**, *14*, 1390–1398.
- (13) Ohta, A.; Murakami, H.; Higashimura, E.; Suga, H. Synthesis of polyester by means of genetic code reprogramming. *Chem. Biol.* **2007**, *14*, 1315–1322.
- (14) Fujino, T.; Goto, Y.; Suga, H.; Murakami, H. Ribosomal synthesis of peptides with multiple β -amino acids. *J. Am. Chem. Soc.* **2016**, *138*, 1962–1969.
- (15) Katoh, T.; Suga, H. Ribosomal incorporation of consecutive β -amino acids. *J. Am. Chem. Soc.* **2018**, *140*, 12159–12167.
- (16) Wang, P. S.; Schepartz, A. β -peptide bundles: Design. Build. Analyze. Biosynthesize. *Chem. Commun.* **2016**, *52*, 7420–7432.
- (17) Checchio, J. W.; Gellman, S. H. Targeting recognition surfaces on natural proteins with peptidic foldamers. *Curr. Opin. Struct. Biol.* **2016**, *39*, 96–105.
- (18) Montalvo, G.; Waegle, M. M.; Shandler, S.; Gai, F.; DeGrado, W. F. Infrared signature and folding dynamics of a helical β -peptide. *J. Am. Chem. Soc.* **2010**, *132*, 5616–8.
- (19) Seebach, D.; Gardiner, J. Beta-peptidic peptidomimetics. *Acc. Chem. Res.* **2008**, *41*, 1366–1375.
- (20) García, J. M.; García, F. C.; Serna, F.; de la Peña, J. L. High-performance aromatic polyamides. *Prog. Polym. Sci.* **2010**, *35*, 623–686.
- (21) Tanner, D.; Fitzgerald, J. A.; Phillips, B. R. The kevlar story - an advanced materials case study. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 649–654.
- (22) Baumann, S.; Herrmann, J.; Raju, R.; Steinmetz, H.; Mohr, K. I.; Hättel, S.; Harmrolfs, K.; Stadler, M.; Mäller, R. Cystobactamids: myxobacterial topoisomerase inhibitors exhibiting potent antibacterial activity. *Angew. Chem., Int. Ed. Engl.* **2014**, *53*, 14605–14609.
- (23) Saraogi, I.; Incarvito, C. D.; Hamilton, A. D. Controlling curvature in a family of oligoamide α -helix mimetics. *Angew. Chem., Int. Ed.* **2008**, *47*, 9691–9694.
- (24) Meisel, J. W.; Hu, C. T.; Hamilton, A. D. Mimicry of a β -hairpin turn by a nonpeptidic laterally flexible foldamer. *Org. Lett.* **2018**, *20*, 3879–3882.
- (25) Saha, S.; Kauffmann, B.; Ferrand, Y.; Huc, I. Selective encapsulation of disaccharide xylobiose by an aromatic foldamer helical capsule. *Angew. Chem., Int. Ed.* **2018**, *57*, 13542–13546.
- (26) Rogers, J. M.; Kwon, S.; Dawson, S. J.; Mandal, P. K.; Suga, H.; Huc, I. Ribosomal synthesis and folding of peptide-helical aromatic foldamer hybrids. *Nat. Chem.* **2018**, *10*, 405–412.

- (27) Schepartz, A. Foldamers wave to the ribosome. *Nat. Chem.* **2018**, *10*, 377–379.
- (28) Goto, Y.; Suga, H. Flexizymes as a tRNA acylation tool facilitating genetic code reprogramming. *Methods Mol. Biol.* **2012**, *848*, 465–478.
- (29) Murakami, H.; Saito, H.; Suga, H. A versatile tRNA aminoacylation catalyst based on RNA. *Chem. Biol.* **2003**, *10*, 655–662.
- (30) McMurry, J. L.; Chang, M. C. Y. Fluorothreonyl-tRNA deacylase prevents mistranslation in the organofluorine producer *Streptomyces cattleya*. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114*, 11920–11925.
- (31) Murakami, H.; Ohta, A.; Ashigai, H.; Suga, H. A highly flexible tRNA acylation method for non-natural polypeptide synthesis. *Nat. Methods* **2006**, *3*, 357–359.
- (32) Nawrot, B.; Sprinzl, M. Aminoacyl-tRNA analogues; synthesis, purification and properties of 3-anthraniloyl oligoribonucleotides. *Nucleosides Nucleotides* **1998**, *17*, 815–829.
- (33) Mortimer, S. A.; Weeks, K. M. A fast-acting reagent for accurate analysis of RNA secondary and tertiary structure by SHAPE chemistry. *J. Am. Chem. Soc.* **2007**, *129*, 4144–4145.
- (34) Drossman, H.; Johnson, H.; Mill, T. Structure activity relationships for environmental processes 1: Hydrolysis of esters and carbamates. *Chemosphere* **1988**, *17*, 1509–1530.
- (35) Xiao, H.; Murakami, H.; Suga, H.; Ferré-D'Amaré, A. R. Structural basis of specific tRNA aminoacylation by a small *in vitro* selected ribozyme. *Nature* **2008**, *454*, 358–361.
- (36) Hunter, C. A.; Lawson, K. R.; Perkins, J.; Urch, C. J. Aromatic interactions. *J. Chem. Soc. Perkins Trans. 2* **2001**, *2*, 651–669.
- (37) Meyer, E. A.; Castellano, R. K.; Diederich, F. Interactions with aromatic rings in chemical and biological recognition. *Angew. Chem., Int. Ed.* **2003**, *42*, 1210–1250.
- (38) Hansch, C.; Leo, A.; Taft, R. W. A survey of Hammett substituent constants and resonance and field parameters. *Chem. Rev.* **1991**, *91*, 165–195.
- (39) Kawakami, T.; Ogawa, K.; Hatta, T.; Goshima, N.; Natsume, T. Directed evolution of a cyclized peptoid-peptide chimera against a cell-free expressed protein and proteomic profiling of the interacting proteins to create a protein-protein interaction inhibitor. *ACS Chem. Biol.* **2016**, *11*, 1569–1577.
- (40) Goto, Y.; Ohta, A.; Sako, Y.; Yamagishi, Y.; Murakami, H.; Suga, H. Reprogramming the translation initiation for the synthesis of physiologically stable cyclic peptides. *ACS Chem. Biol.* **2008**, *3*, 120–129.
- (41) Gualerzi, C. O.; Pon, C. L. Initiation of mRNA translation in bacteria: structural and dynamic aspects. *Cell. Mol. Life Sci.* **2015**, *72*, 4341–4367.
- (42) Goto, Y.; Katoh, T.; Suga, H. Flexizymes for genetic code reprogramming. *Nat. Protoc.* **2011**, *6*, 779–790.
- (43) Staunton, J.; Weissman, K. J. Polyketide biosynthesis: a millennium review. *Nat. Prod. Rep.* **2001**, *18*, 380–416.
- (44) Dutta, S.; Whicher, J. R.; Hansen, D. A.; Hale, W. A.; Chemler, J. A.; Congdon, G. R.; Narayan, A. R. H.; Håkansson, K.; Sherman, D. H.; Smith, J. L.; Skiniotis, G. Structure of a modular polyketide synthase. *Nature* **2014**, *510*, 512–517.
- (45) Robbins, T.; Liu, Y.-C.; Cane, D. E.; Khosla, C. Structure and mechanism of assembly line polyketide synthases. *Curr. Opin. Struct. Biol.* **2016**, *41*, 10–18.
- (46) Du, L.; Sanchez, C.; Shen, B. Hybrid peptide-polyketide natural products: biosynthesis and prospects toward engineering novel molecules. *Metab. Eng.* **2001**, *3*, 78–95.
- (47) Silakowski, B.; Nordsiek, G.; Kunze, B.; Blöcker, H.; Mäller, R. Novel features in a combined polyketide synthase/non-ribosomal peptide synthetase: the myxalamid biosynthetic gene cluster of the myxobacterium *Stigmatella aurantiaca* Sga151. *Chem. Biol.* **2001**, *8*, 59–69.
- (48) Walsh, C. T. Polyketide and nonribosomal peptide antibiotics: modularity and versatility. *Science* **2004**, *303*, 1805–1810.
- (49) Fujino, T.; Kondo, T.; Suga, H.; Murakami, H. Exploring of minimal RNA substrate of flexizymes. *ChemBioChem.* **2019**, in press. DOI: 10.1002/cbic.201900150