

NON-COMMUTATIVE RESOLUTIONS OF TORIC VARIETIES

ELEONORE FABER, GREG MULLER, AND KAREN E. SMITH

ABSTRACT. Let R be the coordinate ring of an affine toric variety. We prove, using direct elementary methods, that the endomorphism ring $\text{End}_R(\mathbb{A})$, where \mathbb{A} is the (finite) direct sum of all (isomorphism classes of) conic R -modules, has finite global dimension equal to the dimension of R . This gives a precise version, and an elementary proof, of a theorem of Špenko and Van den Bergh implying that $\text{End}_R(\mathbb{A})$ has finite global dimension. Furthermore, we show that $\text{End}_R(\mathbb{A})$ is a non-commutative crepant resolution if and only if the toric variety is simplicial. For toric varieties over a perfect field k of prime characteristic, we show that the ring of differential operators $D_k(R)$ has finite global dimension.

1. INTRODUCTION

Consider a local or graded ring R which is commutative and Noetherian. A well-known theorem of Auslander-Buchsbaum and Serre states that R is regular if and only if R has finite global dimension—that is, if and only if every R -module has finite projective dimension.

While the standard definition of regularity does not extend to non-commutative rings, the definition of global dimension does, so this suggests that finite global dimension might play the role of regularity for non-commutative rings. This is an old idea going back at least to Dixmier [Dix63], though since then our understanding of connection between regularity and finite global dimension has been refined by the works of Auslander, Artin, Shelter, Van den Bergh and others. One particularly relevant connection here is the following: *If R is a regular ring of finite type over a perfect field k , then the ring of differential operators $D_k(R)$ has finite global dimension.*¹ This theorem is due to Roos in characteristic zero [Roo72, Cha74], and Paul Smith in characteristic p [Smi87].

Date: April 15, 2019.

2010 Mathematics Subject Classification. 13C14, 13A35, 16E10, 16S32 .

Key words and phrases. toric variety, non-commutative resolution, finite global dimension, non-commutative crepant resolution, rings of differential operators, strong F-regularity, conic module.

The first author was funded by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 789580. The third author was partially supported by the US National Science Foundation, grant DMS-1501625.

¹We provide a counterexample to the converse in Theorem 6.13.

In this paper, we establish the finite global dimension of several non-commutative algebras associated to a *normal toric algebra* R over a field k , a type of algebra which is commutative but rarely regular. These algebras are:

- (1) $\text{End}_R(\mathbb{A})$, where \mathbb{A} is a *complete sum of conic modules*. ‘Conic modules’ are certain combinatorial defined R -modules first studied systematically by Bruns and Gubaladze [BG03], and a direct sum of conic modules is ‘complete’ if every conic module is isomorphic to a summand.
- (2) $\text{End}_R(R^{1/q})$, assuming q is sufficiently large. Here, $R^{1/q}$ is the k -algebra spanned by q -th roots of monomials in R . When k is perfect and q is a power of the characteristic of k , $R^{1/q}$ is the ring of q -th roots of all elements in R .
- (3) The ring of k -linear differential operators $D_k(R)$ of R , assuming k is perfect with positive characteristic.

Our primary result is an explicit projective resolution for each simple module for the algebra $\text{End}_R(\mathbb{A})$ in (1), from which it follows that the global dimension is equal to the dimension of R ; see Theorem 6.1. Similar results follow for the algebras $\text{End}_R(R^{1/q})$ in case (2), which we show in Proposition 4.15 are Morita equivalent to $\text{End}_R(\mathbb{A})$. The result for differential operators follows by taking a direct limit; see Theorem 6.13. Our proofs are elementary, direct, self-contained, and constructive. It should be noted that in a very nice paper, Špenko and Van den Bergh independently found less constructive proofs for the finiteness of global dimension in settings (1) and (2), albeit without explicit bounds on the global dimension [ŠVdB17b, 1.3.6]; they also beat us to publication. See also [RŠVdB17].

Because these algebras have finite global dimension, they should be regarded as ‘regular’ in an appropriate non-commutative sense. Concretely, we may say the first two are **non-commutative resolutions** of the toric algebra R , as defined by Bondal and Orlov [BO02]; see also [Leu12, Wem16, DITV15].

Finally, we consider when these non-commutative resolutions are **crepant**, a condition introduced in [VdB04] which ensures the module category of the resolution is sufficiently close to the module category of the original algebra. We conclude that $\text{End}_R(\mathbb{A})$ is a non-commutative crepant resolution if and only if the toric algebra R is *simplicial* (Theorem 7.2).

This work naturally leads to several questions for future research.

- What conditions on a ring R ensure that $D_k(R)$ has finite global dimension?
- For an arbitrary algebra R of characteristic p , what conditions on R imply the global dimension of $\text{End}_R(R^{1/p^e})$ is finite for sufficiently large e ?
- When is there some (non-complete) sum \mathbb{B} of conic modules over a toric ring such that the endomorphism ring $\text{End}_R(\mathbb{B})$ is a non-commutative crepant resolution?

- Can our work on the finite global dimension of $D_k(R)$ in prime characteristic be used to prove finite global dimension of $D_k(R)$ in characteristic zero (when R is a toric algebra)? Recent work of Jeffries [Jef17] shows that the obstruction of reduction to prime characteristic for differential operators introduced in [SVdB97] may vanish for all toric algebras. See also [MVdB98] for a more direct approach to showing that rings of differential operators on toric rings of characteristic zero have finite global dimension.

1.1. The proof: Conic modules of toric algebras. The preceding results are proven by carefully considering the *conic modules* of a normal toric algebra R . These are fractional monomial ideals of R whose monomial support is given by shifting the defining cone of R . Conic modules are rank one and Cohen-Macaulay, and each toric R has finitely many conic modules up to isomorphism. Conic modules may be parameterized by polyhedral **chambers of constancy** inside a hyperplane arrangement. These properties of conic modules have been systematically studied by Bruns and Gubeladze [BG03, Bru05].

For each chamber of constancy Δ , we use the combinatorics of the faces in Δ to construct a chain complex K_Δ^\bullet of sums of conic modules. The essential property of these complexes is the Acyclicity Lemma (Lemma 5.4), which states that the complex $\text{Hom}_R(A_{\Delta'}, K_\Delta^\bullet)$ is either acyclic or a resolution of the ground field k , depending on whether or not $A_\Delta \simeq A_{\Delta'}$.

The benefit of this lemma is as follows. A direct sum of conic modules \mathbb{A} is *complete* if every conic module of R is isomorphic to a summand of \mathbb{A} . Our motivating example is the ring $R^{1/n}$ spanned by formal n th roots of monomials in R , which is a complete sum of conic modules provided n is large enough (Proposition 4.15).

Given a complete sum of conic modules \mathbb{A} and a chamber of constancy Δ , the Acyclicity Lemma implies that the complex $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet)$ is a finite projective resolution of a simple $\text{End}_R(\mathbb{A})$ -module. Using standard arguments on global dimension (and a nice choice of grading), we can show that every finitely generated $\text{End}_R(\mathbb{A})$ -module has a finite projective resolution; and thus, the finiteness of the global dimension.

1.2. Relation with the Frobenius map. The classification of commutative rings of characteristic p according to the behavior of Frobenius has a long tradition in commutative algebra. This began with Kunz's theorem that regularity can be characterized by the flatness of Frobenius [Kun69], which led to the development of tight closure and F-regularity [HH90]. Eventually, relationships with the singularities in the minimal model program were discovered and developed [Smi97], [Smi00], [HW02], [Tak04], as well as connections with rings of differential operators [Smi95]

[SVdB97]. Since then, deeper connections with the minimal model program, including the log canonical threshold, multiplier ideals, and log Fano varieties (see for example, [Sch09] [HY03], [ST12], [Bli13], [HX15]), differential operators (see, for example [JNB17]), and F-signature (see [HL02], [Tuc12], [CRST17], for example) have blossomed.

Our work suggests that under some hypothesis, Frobenius may provide a way to construct a *functorial* non-commutative resolution of singularities. The property that a \mathbb{C} -algebra admits a non-commutative crepant resolution is closely related to rational singularities (see [VdB04, SVdB08]). On the other hand, the property of strong-F-regularity is a prime characteristic analog of rational singularities (at least in the Gorenstein case). We speculate that for a strongly F-regular ring R with finite F-representative type (a class of rings including toric rings in prime characteristic), perhaps $\text{End}_R(R^{1/p^e})$ always has finite global dimension. This speculation encompasses the classes of rings arising in representation theory where this idea has been explored from a somewhat different point of view in [ŠVdB17b, RŠVdB17].

Our work corrects a result of Yasuda [Yas10], and is closely related to recent work of Higashitani and Nakajima [HN17]. A different approach to the existence of non-commutative crepant resolutions for three dimensional Gorenstein toric rings in low dimension can be found in [Bro12]; see also [Boc12]. As mentioned earlier, our results overlap considerably with work of Van den Bergh and Špenko, although their machinery is quite different and less constructive; see [ŠVdB17a], [ŠVdB17b]. Our technique, by contrast, is a fairly direct and self-contained, so which should be widely accessible, including to commutative algebraists and algebraic geometers.

1.3. Outline of the paper. After recalling basic definitions and notation associated with toric algebras (also known as normal semi-group rings) in Section 2, we recall the notion *conic modules* from [BG03] and develop it for our purposes in Section 3, culminating in Brun’s cell-decomposition of \mathbb{R}^d induced by any full dimensional pointed rational polyhedral cone C in \mathbb{R}^d [Bru05]. Also in Sections 3 and 4, we introduce the idea of a *open conic module* and its relationship with the cell decomposition, and an instrumental idea in the following sections, and prove several other needed results, for example on the behavior of conic modules under restriction. The technical heart of the paper is in Section 5, where we use this cell decomposition to construct complexes of conic modules over R and prove the crucial Acyclicity Lemma, Lemma 5.4. The benefits of this work is reaped in Section 6, where we show how to induce explicit projective resolutions of simple modules over $\text{End}_R(\mathbb{A})$. Also in Section 6, we finish the proof of Theorem 6.1 that $\text{End}_R(\mathbb{A})$ has finite global dimension, and deduce Corollary 6.13 stating that in prime characteristic, rings of differential operators on toric varieties have finite global dimension.

Finally in Section 7, we prove Theorem 7.2 characterizing the non-commutative resolution $\text{End}_R(\mathbb{A})$ as crepant if and only if R is simplicial.

2. TORIC ALGEBRAS

Fix an arbitrary ground field k . We begin with a recap of toric algebras over k , otherwise known as finitely generated normal semi-group algebras. See [Ful93] for details.

Let M be a finitely generated free abelian group (also called a *lattice*), and let $M_{\mathbb{R}} := \mathbb{R} \otimes M$ be the associated real finite dimensional vector space. A **rational polyhedral cone** C in $M_{\mathbb{R}}$ is a closed set consisting of all non-negative real linear combinations of a finite set of elements in M .

Given a rational polyhedral cone $C \subset M_{\mathbb{R}}$, the associated **toric algebra** $R = k[C \cap M]$ is the k -subalgebra of the Laurent ring $k[M] = k[x^m \mid m \in M]$ defined by

$$R := \text{Span}\{x^m \mid m \in M \cap C\}.$$

Alternatively, the ring R can be defined simply as the semi-group algebra of the semi-group $C \cap M$. The Laurent ring itself is the special case where $C = M_{\mathbb{R}}$.

The algebra R is a domain finitely generated over k , naturally graded by the group M ; its dimension is equal to the rank of the lattice M [Ful93, §1.2]. It is also well-known to be normal and Cohen-Macaulay [Ful93, p29-30].

The algebra $R = k[C \cap M]$ is the coordinate ring of the associated **affine toric variety**

$$X_{\sigma} = \text{Spec } R,$$

where $\sigma \subset N_{\mathbb{R}}$ is the dual cone of C in the dual vector space $N_{\mathbb{R}} = M_{\mathbb{R}}^*$. Typically in the literature, one begins with a finite rank lattice N , and a rational polyhedral cone $\sigma \subset N \otimes \mathbb{R} = N_{\mathbb{R}}$. Our cone C is the dual cone to σ , namely

$$(2.1) \quad C = \{v \in M_{\mathbb{R}} \mid \langle v, w \rangle \geq 0 \text{ for all } w \in \sigma\},$$

where $\langle v, w \rangle$ denotes the natural pairing between elements of $M_{\mathbb{R}}$ and its dual space $N_{\mathbb{R}}$ and M is the dual lattice $\text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$.

Without loss of generality, we can and will assume throughout that our cone C is **full dimensional** in $M_{\mathbb{R}}$. Indeed, we can replace $M_{\mathbb{R}}$ with the span of C if necessary, and the lattice M by its intersection with this span. The assumption that C is full-dimensional is equivalent to the assumption that the dual cone σ is **pointed** (or **strongly convex**), meaning that it contains no subspace of $N_{\mathbb{R}}$. In this case, it is easy that R and $k[M]$ have the same fraction field, so that the Krull dimension of R is equal to the rank of M .

2.1. Primitive Generators. Fix a rational polyhedral cone σ in $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$. Throughout this paper, the notation

$$\Sigma_1 =: \{n_1, n_2, \dots, n_t\} \subset N$$

denotes a set of **minimal primitive** generators for σ . Minimality means that none of the n_i is redundant: removing any one will span a strictly smaller cone; equivalently in the case where σ is pointed, this means that each n_i spans an **extremal ray** of σ in $N_{\mathbb{R}}$. **Primitivity** means that $n_i \in N$ but $\varepsilon n_i \notin N$ for $0 < \varepsilon < 1$. If σ is pointed, the set Σ_1 of minimal primitive generators is uniquely determined.

Equation (2.1) can be rewritten

$$(2.2) \quad \begin{aligned} C &= \{v \in M_{\mathbb{R}} \mid \langle v, n_i \rangle \geq 0 \text{ for all } i = 1, \dots, t\} \\ &= \bigcap_{i=1}^t \{v \in M_{\mathbb{R}} \mid \langle v, n_i \rangle \geq 0\}, \end{aligned}$$

which represents C as a finite intersection of closed half-spaces in $M_{\mathbb{R}}$. In particular, the **facets** (or codimension 1 subcones) of C are indexed by the elements n_i of Σ_1 : each facet of C is precisely the intersection of C with a supporting hyperplane defined by the equation $\langle x, n_i \rangle = 0$ in $M_{\mathbb{R}}$. Abusing terminology slightly, we call the n_i the **primitive inward-pointing normals** to the facets of C .

2.2. Simplicial Toric Rings. Fix a toric ring $R = k[C \cap M]$, where, without loss of generality, we assume that C spans $M_{\mathbb{R}}$. Recall that R is said to be **simplicial** if the primitive generators Σ_1 are *linearly independent*. Simplicial toric varieties are an especially nice class: for example, they have cyclic quotient singularities [Ful93, 2.2].

Toric rings of dimension two are always simplicial, since a pointed full dimensional cone σ in a two dimensional vector space has exactly two extremal rays. However, in higher dimension, the typical toric ring is not simplicial; see Example 4.14.

3. CONIC MODULES

The monomials of a toric algebra R are defined by intersecting a rational polyhedral cone C with a lattice M . If we translate the cone and intersect with M , we instead define the monomials of an R -module, called a *conic module*.

For the formal definition, fix a toric algebra R corresponding to a full-dimensional rational polyhedral cone $C \subset M_{\mathbb{R}}$.

Definition 3.1. Fix a vector $v \in M_{\mathbb{R}}$. The **conic module**² defined by v is the M -graded R -submodule of the Laurent ring $\mathbb{k}[M]$ defined as follows:

$$A_v := \text{Span}\{x^m \mid m \in M \cap (C + v)\}.$$

Proposition 3.2. Let A_v be a conic module for the toric algebra $R = \mathbb{k}[C \cap M]$. Then

- (1) A_v is torsion free and rank one over R .
- (2) A_v is spanned by monomials x^m where $m \in M$ is in the intersection of half spaces

$$\bigcap_{n_i \in \Sigma_1} \{x \in M_{\mathbb{R}} \mid \langle x, n_i \rangle \geq \lceil \langle v, n_i \rangle \rceil\},$$

where the notation $\lceil \lambda \rceil$ denote the smallest integer $\geq \lambda$.

- (3) For any two conic modules A_v and A_w , the module $\text{Hom}_R(A_v, A_w)$ is naturally isomorphic to the M -graded R -submodule of $\mathbb{k}[M]$

$$\text{Span}\{x^m \mid m + (C + v) \cap M \subset (C + w)\}.$$

In particular, the only degree zero R -module homomorphisms are inclusions $A_v \subset A_w$ (composed with scalar multiplication by an element of \mathbb{k}).

Remark 3.3. In addition, conic modules are always Cohen-Macaulay, hence reflexive. This is proved in Corollary 4.16. ■

Remark 3.4. It is tempting to suspect that $\text{Hom}_R(A_v, A_w)$ might be the conic module A_{w-v} , or perhaps another conic module. This is false in general; in Example 4.14, $\text{Hom}_R(A_1, A_2)$ is not even Cohen-Macaulay. However, we show in Proposition 7.2 that $\text{Hom}_R(A_v, A_w)$ is always conic in the special case that R is simplicial. ■

Proof. For (1), note that $A_v \subset \mathbb{k}[M]$ by definition, so it is clearly a torsion-free R -module. Furthermore, tensoring over R with the Laurent ring $\mathbb{k}[M]$, we have $A_v \otimes_R \mathbb{k}[M] \cong \mathbb{k}[M]$, so A_v is rank one.

For (2), observe that the conic module A_v is the span of $x^m \in \mathbb{k}[M]$ where m is in the set

$$\begin{aligned} M \cap (C + v) &= \{m \in M \mid m - v \in C\} \\ &= \{m \in M \mid \langle m - v, n_i \rangle \geq 0 \text{ for all } n_i \in \Sigma_1\} \\ &= \{m \in M \mid \langle m, n_i \rangle \geq \langle v, n_i \rangle \text{ for all } n_i \in \Sigma_1\} \\ &= \{m \in M \mid \langle m, n_i \rangle \geq \lceil \langle v, n_i \rangle \rceil \text{ for all } n_i \in \Sigma_1\} \\ &= M \cap \left(\bigcap_{n_i \in \Sigma_1} \{x \in M_{\mathbb{R}} \mid \langle x, n_i \rangle \geq \lceil \langle v, n_i \rangle \rceil\} \right). \end{aligned}$$

²The name “conic module” seems to have been coined in [BG03], although these were studied earlier, for example, in [Sta82, Don02].

The penultimate equality here follows because $\langle m, n_i \rangle$ is an integer.

For (3), note that since A_v and A_w are finitely generated and M -graded, so is $\text{Hom}_R(A_v, A_w)$. Tensoring with the Laurent ring $\mathbf{k}[M]$, we see that $\text{Hom}_R(A_v, A_w)$ is an R -submodule of $\text{Hom}_{\mathbf{k}[M]}(\mathbf{k}[M], \mathbf{k}[M]) \cong \mathbf{k}[M]$ so that every homogeneous R -module homomorphism $A_v \rightarrow A_w$ is multiplication by some monomial in $\mathbf{k}[M]$. The statement follows. In particular, the only degree-preserving maps are $\mu_\lambda : A_v \rightarrow A_w$ sending each x^m to λx^m , where $\lambda \in \mathbf{k}$. \square

3.1. Open Conic Modules. For any $v \in M_{\mathbb{R}}$, we define the associated **open conic module** to be the M -graded R -submodule of $\mathbf{k}[M]$:

$$\mathring{A}_v := \text{Span} \{x^m \mid m \in M \cap (\mathring{C} + v)\},$$

where $\mathring{C} + v$ denotes the *interior* of the cone $C + v$.

Open conic modules turn out to be conic modules of “nearby” vectors in $M_{\mathbb{R}}$:

Proposition 3.5. *Fix a toric algebra R corresponding to a full-dimensional cone C in $M_{\mathbb{R}}$. For every open conic R module \mathring{A}_v , there exists $w \in M_{\mathbb{R}}$ such that*

$$\mathring{A}_v = A_w.$$

Explicitly, we can take $w = v + \varepsilon p$ where p is any vector in the interior of C and ε is a sufficiently small positive real number.

Proof of Proposition 3.5. Fix any $p \in M_{\mathbb{R}}$ in the interior of C . This means that $\langle p, n_i \rangle > 0$ for all $n_i \in \Sigma_1$. So for any $m \in M$, we have

$$\langle m - v, n_i \rangle > 0 \quad \text{if and only if} \quad \langle m - (v + \varepsilon p), n_i \rangle \geq 0$$

for sufficiently small positive ε . This exactly says that $\mathring{A}_v = A_{v+\varepsilon p}$. \square

4. CHAMBERS OF CONSTANCY AND CELL DECOMPOSITION

Through this section, C denotes a full-dimensional rational polyhedral cone C in $M_{\mathbb{R}} = M \otimes \mathbb{R}$. As usual, $R = \mathbf{k}[C \cap M]$ denotes the corresponding toric ring and Σ_1 the set of primitive inward-pointing normals to its facets.

Different vectors $v \in M_{\mathbb{R}}$ can determine the same conic module over R . This leads to the following partition of $M_{\mathbb{R}}$ into equivalence classes of vectors determining the same conic module:

Definition 4.1. The **chamber of constancy** $\Delta \subset M_{\mathbb{R}}$ containing $v \in M_{\mathbb{R}}$ is the set of all $w \in M_{\mathbb{R}}$ such that $A_v = A_w$. Equivalently, w and v belong to the same chamber of constancy if and only if $(C + v) \cap M = (C + w) \cap M$.

Notation 4.2. Because two vectors determine the same conic module if and only if they belong to the same chamber Δ , it is natural to denote a conic module A_Δ .

Proposition 4.3. *The chamber of constancy containing v is the set*

$$(4.1) \quad \begin{aligned} \Delta &= \{x \in M_{\mathbb{R}} \mid \lceil \langle v, n_i \rangle \rceil - 1 < \langle x, n_i \rangle \leq \lceil \langle v, n_i \rangle \rceil \text{ for all } n_i \in \Sigma_1\} \\ &= \bigcap_{n_i \in \Sigma_1} \{x \in M_{\mathbb{R}} \mid \lceil \langle v, n_i \rangle \rceil - 1 < \langle x, n_i \rangle \leq \lceil \langle v, n_i \rangle \rceil\} \end{aligned}$$

where n_1, \dots, n_t runs through the set Σ_1 of primitive inward-pointing normal vectors to the facets of C .

Before proving this, we note the following immediate corollary.

Corollary 4.4. *Fix a full-dimensional rational polyhedral cone $C \subset M_{\mathbb{R}}$. The corresponding chambers of constancy decompose $M_{\mathbb{R}}$ into disjoint locally closed polyhedral regions; these chambers are bounded if C is pointed.*

Proof of Corollary. Proposition 4.3 expresses a chamber Δ as an intersection of finitely many half-open strips formed by the intersection of a closed half space and a parallel open half space. Thus it is locally closed and polyhedral. If C is pointed, then its dual σ is full-dimensional, so Σ_1 spans $N_{\mathbb{R}}$, making Δ bounded. \square

Proof of Proposition 4.3. Fix $v \in M_{\mathbb{R}}$. The Lemma is equivalent to the claim that $A_w = A_v$ if and only if the two lists of integers

$$\lceil \langle v, n_1 \rangle \rceil, \lceil \langle v, n_2 \rangle \rceil, \dots, \lceil \langle v, n_t \rangle \rceil$$

and

$$\lceil \langle w, n_1 \rangle \rceil, \lceil \langle w, n_2 \rangle \rceil, \dots, \lceil \langle w, n_t \rangle \rceil$$

are the same. Clearly, if $\lceil \langle v, n_i \rangle \rceil = \lceil \langle w, n_i \rangle \rceil$ for all $n_i \in \Sigma_1$, then Proposition 3.2 (2) ensures that $A_v = A_w$. It remains to prove the converse.

Let us assume, without loss of generality, that $\lceil \langle v, n_1 \rangle \rceil < \lceil \langle w, n_1 \rangle \rceil$. We need to show that $A_v \neq A_w$. For this, we will find a monomial $x^m \in A_v$ but not in A_w .

Pick any $m \in M \cap \mathcal{H}$, where \mathcal{H} is the hyperplane in $M_{\mathbb{R}}$ whose equation is $\langle x, n_1 \rangle = \lceil \langle v, n_1 \rangle \rceil$. Note that m is *not* in the closed half space

$$\{x \in M_{\mathbb{R}} \mid \langle x, n_1 \rangle \geq \lceil \langle w, n_1 \rangle \rceil\},$$

so that $m \notin C + w$ by Proposition 3.2 (2). In particular, $x^m \notin A_w$ for any $m \in M \cap \mathcal{H}$. On the other hand, again by Proposition 3.2, the monomials in A_v are those $m \in M$ such that

$$m \in \bigcap_{i=1}^t \{x \in M_{\mathbb{R}} \mid \langle x, n_i \rangle \geq \lceil \langle v, n_i \rangle \rceil\},$$

a convex subset of $M_{\mathbb{R}}$ which has non-empty intersection with rational hyperplane \mathcal{H} and hence with the lattice $M \cap \mathcal{H}$. Therefore, we can find $m \in (C + v) \cap M \cap \mathcal{H}$. This m satisfies $x^m \in A_v \setminus A_w$. \square

Remark 4.5. The proof of Proposition 4.3 implies that a conic module A_v is uniquely described by the list of integers

$$(4.2) \quad (\lceil \langle v, n_1 \rangle \rceil, \lceil \langle v, n_2 \rangle \rceil, \dots, \lceil \langle v, n_t \rangle \rceil) \in \mathbb{Z}^{\Sigma_1}$$

where the $n_i \in \Sigma_1$ range through the primitive linear forms defining the facets of C . This can be understood in terms of the group of torus-invariant Weil divisors on the toric variety $X_\sigma = \text{Spec } R$, a free group generated by the irreducible codimension one toric subvarieties determined by each of the facets of C , hence is also isomorphic to \mathbb{Z}^{Σ_1} . The conic module A_v is the module of global section of the reflexive sheaf $\mathcal{O}_{X_\sigma}(-D)$, where D is the divisor

$$\lceil \langle v, n_1 \rangle \rceil D_1 + \lceil \langle v, n_2 \rangle \rceil D_2 + \dots + \lceil \langle v, n_t \rangle \rceil D_t,$$

where D_i is the divisor of the facet indexed by $n_i \in \Sigma_1$. The reader is cautioned, however, that the conic modules are a *strict* subset of the set of all reflexive M -graded submodules of the fraction field of $\mathbb{k}[M]$ in general, so not every list of integers corresponds to a conic module.³ ■

4.1. CW Decomposition of $M_{\mathbb{R}}$. The cone C in $M_{\mathbb{R}}$ determines a CW decomposition of $M_{\mathbb{R}}$ whose maximal dimensional cells are the interiors of the chambers of constancy. More precisely:

Proposition 4.6. [Bru05] *Fix a full dimensional rational polyhedral cone C in $M_{\mathbb{R}}$. Each of its chambers of constancy in $M_{\mathbb{R}}$ is a disjoint union of open polyhedral cells (of different dimensions). Together, ranging over all chambers, these cells define a CW decomposition of $M_{\mathbb{R}}$.*

Remark 4.7. We caution the reader that a chamber of constancy Δ need not have cells of every dimension $\leq \dim M_{\mathbb{R}}$. See Section 7. ■

Proof of Proposition 4.6. Fix a chamber of constancy Δ containing a vector $v \in M_{\mathbb{R}}$. We will explicitly describe Δ as a finite union of open polyhedral cells.

There is exactly one open cell of codimension zero, namely the interior of Δ . This is the set of vectors in $v \in M_{\mathbb{R}}$ satisfying

$$(4.3) \quad \lceil \langle v, n_i \rangle \rceil - 1 < \langle x, n_i \rangle < \lceil \langle v, n_i \rangle \rceil$$

for all $n_i \in \Sigma_1$, the set of primitive inward-pointing normals to the facets of C . We can think of these inequalities as the complete set of inequalities defining Δ as in Proposition 4.3 but in which *all inequalities are strict*.

The open cells of codimension 1 are the interiors of the *facets* of Δ . These cells are defined as the points of $M_{\mathbb{R}}$ satisfying the same set of strict linear inequalities

³In general not even all rank 1 maximal Cohen-Macaulay modules are conic, see [Bae04].

as above in (4.3), but with exactly one of the linear inequalities replaced by an *equality* of the form $\langle x, n_i \rangle = \lceil \langle v, n_i \rangle \rceil$ for some $n_i \in \Sigma_1$.

The open cells of codimension 2 are similarly defined by the same set of strict linear inequalities (4.3) but with two equalities. These cells are open polytopes in the codimension two linear spaces in $M_{\mathbb{R}}$ obtained by intersecting two supporting hyperplanes of Δ . In a similar way, each cell of codimension i of Δ is an open polytope in a codimension i linear space of $M_{\mathbb{R}}$ obtained by intersecting i supporting hyperplanes of Δ .

Summarizing formally, we can describe each open cell of Δ as follows. Pick any $v \in \Delta$ and set $d_j = \lceil \langle v, n_j \rangle \rceil$ for each $n_j \in \Sigma_1$. Then

$$(4.4) \quad \tau_{\Omega} := \left\{ x \in M_{\mathbb{R}} \mid \begin{array}{ll} d_j - 1 < \langle x, n_j \rangle < d_j & \text{for } n_j \in \Omega, \\ \langle x, n_j \rangle = d_j & \text{for } n_j \notin \Omega \end{array} \right\}.$$

where Ω is some subset of the set Σ_1 of primitive inward-pointing normals of C .

The space $M_{\mathbb{R}}$ is clearly the disjoint union of these polyhedral cells, as we range over all cells of all chambers. It is straightforward to check that this cell decomposition satisfies the definition of a CW complex. (See, for example, [Mun84, §38].) \square

Remark 4.8. The cell decomposition of Δ is not a CW complex because some boundary cells are missing. However, the closure $\overline{\Delta}$ of Δ is a CW subcomplex of $M_{\mathbb{R}}$ in an obvious way. \blacksquare

The CW decomposition of $M_{\mathbb{R}}$ can be clarified using open conic modules:

Corollary 4.9. *With notation as above, two vectors $v, w \in M_{\mathbb{R}}$ lie in the same open cell of the CW decomposition of $M_{\mathbb{R}}$ if and only if*

$$A_v = A_w \quad \text{and} \quad \mathring{A}_v = \mathring{A}_w.$$

Moreover, for v and w lying in the same chamber of constancy Δ we have

$$\mathring{A}_v \subset \mathring{A}_w$$

if and only if the open cell containing v is contained in the boundary of the open cell containing w .

Proof. Fix $v, w \in \Delta$. It suffices to show that v and w are in the same open cell of Δ if and only if $\mathring{A}_v = \mathring{A}_w$.

Fix p in the interior of Δ . According to Proposition 3.5, $\mathring{A}_v = \mathring{A}_w$ if and only if

$$(4.5) \quad \lceil \langle v + \varepsilon p, n_j \rangle \rceil = \lceil \langle w + \varepsilon p, n_j \rangle \rceil$$

for all $n_j \in \Sigma_1$ and sufficiently small ε . But (4.5) holds if and only if

$$\langle v, n_j \rangle \in \mathbb{Z} \iff \langle w, n_j \rangle \in \mathbb{Z}$$

in which case both values are $d_j = \lceil \langle v, n_j \rangle \rceil = \lceil \langle w, n_j \rangle \rceil$. But this means exactly that both v and w lie in the open cell

$$(4.6) \quad \left\{ x \in M_{\mathbb{R}} \mid \begin{array}{ll} d_j - 1 < \langle x, n_j \rangle < d_j & \text{for } j \in \Omega \\ \langle x, n_j \rangle = d_j & \text{for } j \notin \Omega \end{array} \right\}.$$

where Ω is the set of indices for which $\langle v, n_j \rangle$ and $\langle w, n_j \rangle$ are *not* integers.

For the second statement, we observe that τ is in the boundary of τ' precisely when τ is the intersection of τ' with some supporting hyperplanes for Δ defined by linear equations $\langle x, n_k \rangle = \lceil \langle v, n_k \rangle \rceil$. So clearly if $m \in M$ satisfies all the defining (in)equalities to be in the interior of $C + v$ (where $v \in \tau$), then it also satisfies the less restrictive conditions to be in the interior of $C + w$ (where $w \in \tau'$). This says exactly that $\mathring{A}_v \subset \mathring{A}_w$. \square

Notation 4.10. In light of Corollary 4.9, it is reasonable to denote an open conic module \mathring{A}_v by \mathring{A}_τ where τ is the open cell containing v . However, it can happen that $\mathring{A}_\tau = \mathring{A}_{\tau'}$ for distinct cells τ and τ' , so we will use this notation only when considering open conic modules \mathring{A}_v for v in a fixed chamber of constancy Δ .

4.2. Isomorphism Classes of Conic Modules. Isomorphic conic modules can be understood in terms of the chamber decomposition of $M_{\mathbb{R}}$:

Proposition 4.11. *Two conic R -modules A_Δ and $A_{\Delta'}$ are isomorphic if and only if there is an $m \in M$ such that $\Delta = \Delta' + m$.*

Proof. Because conic modules are M -graded rank one torsion-free modules over the normal ring R , any M -graded isomorphism between them is given by multiplication by some monomial x^m of the Laurent ring $k[x^M]$. So $A_v \cong A_w$ if and only if $A_w = A_{v+m}$ for some $m \in M$. \square

Proposition 4.11 says that M acts naturally by translation on the set of chambers of constancy. The orbits of this action index the distinct isomorphism classes of conic modules just as the chambers themselves index distinct conic modules. Put differently, there is an induced CW decomposition of the real torus $M_{\mathbb{R}}/M$ whose maximal dimensional cells are in one-one-correspondence with the isomorphism classes of conic modules for R . The compactness of $M_{\mathbb{R}}/M$ has the following immediate consequence:

Corollary 4.12. [BG03, Prop. 3.6] *For any toric algebra R , there are finitely many isomorphism types of conic R -modules.*⁴

Example 4.13. The cone C in Figure 1a induces the CW decomposition of the plane in Figure 1b. The chambers of constancy in $M_{\mathbb{R}} = \mathbb{R}^2$ are the connected unions of same-colored cells.




FIGURE 1. A cone and the corresponding chambers.

The ring $R = \mathbb{k}[C \cap \mathbb{Z}^2]$ has presentation

$$R = \mathbb{k}[x, y, z]/(xz - y^2).$$

There are two isomorphism classes of conic R -modules, represented by $A_0 = R$ (the blue chambers) and $A = Ry + Rz$ (the red chambers). ■

Example 4.14. Consider the cone C in \mathbb{R}^3 over the unit square depicted in Figure 2. The corresponding toric algebra $R = \mathbb{k}[C \cap \mathbb{Z}^3]$ has presentation $\mathbb{k}[x, y, z, w]/(xz - yw)$. There are three isomorphism types of conic R -modules, represented by $R = A_0$, $(x, y)R = A_1$ and $(x, w)R = A_2$. These are represented (up to translation by \mathbb{Z}^3) by the three chambers depicted in Figure 3. Figure 4 depicts several subsets of the cellular neighborhood of a point in \mathbb{R}^3 . ■

4.3. Frobenius and Cohen-Macaulayness. Fix a positive integer q . Let $q^{-1}\mathbb{Z}$ be the cyclic subgroup of \mathbb{Q} generated by $\frac{1}{q}$ and let $q^{-1}M$ be the abelian group $q^{-1}\mathbb{Z} \otimes M$. Note that $q^{-1}M$ is a lattice containing M as an index q^d subgroup, where d is the rank of M , and that $q^{-1}M_{\mathbb{R}} = M_{\mathbb{R}}$.

Consider the toric algebra

$$R^{1/q} := \mathbb{k}[C \cap q^{-1}M] = \text{Span}\{x^{m/q} \mid \frac{m}{q} \in C \cap q^{-1}M\}.$$

⁴For a different interpretation cf. [Bri93, VdB93] where a criterion for Cohen-Macaulayness of modules of covariants is given, and [SVdB97, Prop. 3.2.3] that shows that $R^{1/q}$ is a direct sum of modules of covariants belonging to so-called strongly critical characters.



FIGURE 2. The cone C over a square.

FIGURE 3. The three types of Chambers of Constancy.




FIGURE 4. Unions of chambers

This notation is meant to be suggestive: if $q = p^e$ where the ground field k is a perfect of characteristic p , then $R^{1/q}$ is the ring of p^e -th roots of all elements in $R = k[C \cap M]$. However, the toric algebra $R^{1/q}$ makes sense in general for all natural numbers q , even in characteristic zero.

The toric algebra $R^{1/q}$ is naturally graded by the lattice $q^{-1}M$. It contains the sub-algebra $R = k[C \cap M]$ as the subring of elements whose degrees are in the sublattice M .

Proposition 4.15. [Bru05] *The R -module $R^{1/q}$ decomposes as a direct sum of (finitely many) modules isomorphic to conic R -modules. For sufficiently large q , every conic module appears (up to isomorphism) as a summand of $R^{1/q}$.*

Proof. Considering $R^{1/q}$ as a module over the subring $R = \mathbf{k}[C \cap M]$, there is a natural grading by the quotient group $q^{-1}M/M$. The graded pieces give an R -module decomposition

$$R^{1/q} = \bigoplus_{[\frac{v}{q}] \in q^{-1}M/M} [R^{1/q}]_{[\frac{v}{q}]} = \bigoplus_{[\frac{v}{q}] \in q^{-1}M/M} \text{Span}\{x^{\frac{v}{q}+m} \mid \frac{v}{q} + m \in (\frac{v}{q} + M) \cap C\}$$

indexed by the classes $[\frac{v}{q}] = \frac{v}{q} + M$ of the quotient group $q^{-1}M/M$. Note that $R^{1/q}$ is finite over its degree 0-graded piece, $R = \mathbf{k}[C \cap M]$.

The $q^{-1}M/M$ -graded pieces of this decomposition are all (isomorphic to) conic modules. Indeed, fixing any representative $\frac{v}{q}$ for a coset in $q^{-1}M/M$, the natural map

$$A_{-\frac{v}{q}} \rightarrow [R^{1/q}]_{[\frac{v}{q}]}$$

given by multiplication by $x^{v/q}$ is easily seen to be an isomorphism.

To see that every conic module appears as a summand of $R^{1/q}$, we need only observe that we can refine the lattice by a sufficiently large q so that each chamber of constancy Δ contains a lattice point in $q^{-1}M$. \square

Corollary 4.16. *The conic modules of a toric algebra are all Cohen-Macaulay.*

Proof. Toric algebras are Cohen-Macaulay. Thus $R^{1/q}$ is a Cohen-Macaulay ring. Being finite over the subring R , $R^{1/q}$ is Cohen-Macaulay as an R -module. Thus each of the R -module direct summands $[R^{1/q}]_{[\frac{v}{q}]}$ is a Cohen-Macaulay R -module. For sufficiently large q , these include all the isomorphism types of conic modules. \square

4.4. The Poset of Chambers of Constancy. The set of conic R -modules has a natural partial ordering given by inclusion. Since the conic modules are naturally indexed by the chambers of constancy, we have a corresponding partial ordering of the set of chambers: $\Delta \preceq \Delta'$ if and only if $A_\Delta \subseteq A_{\Delta'}$.

Proposition 3.2(3) can be rephrased using this language as follows:

Proposition 4.17. *Let Δ and Δ' be chambers of constancy for the toric ring R . Then*

$$\text{Hom}_R(A_\Delta, A_{\Delta'}) = \text{Span}\{x^m \mid m + \Delta \preceq \Delta'\}.$$

Remark 4.18. The partial ordering on chambers can be described in terms of the list of integers

$$(\lceil \langle v, n_1 \rangle \rceil, \lceil \langle v, n_2 \rangle \rceil, \dots, \lceil \langle v, n_t \rangle \rceil)$$

determining Δ as in Remark 4.5. That is, $\Delta \preceq \Delta'$ if and only if their corresponding integer lists (d_1, d_2, \dots, d_t) and $(d'_1, d'_2, \dots, d'_t)$ satisfy $-d_i \leq -d'_i$ for every i . That is, $\Delta \preceq \Delta'$ if and only if $\lceil \langle v, n_i \rangle \rceil \geq \lceil \langle v', n_i \rangle \rceil$ for all $n_i \in \Sigma_1$ and any (equivalently, every) $v \in \Delta$ and $v' \in \Delta'$. \blacksquare

We assign to each chamber of constancy a numerical value which helps us understand its place in the poset:

Definition 4.19. The **degree** of the chamber of constancy Δ is

$$\deg(\Delta) = - \sum_{n_i \in \Sigma_1} \lceil \langle v, n_i \rangle \rceil,$$

where v is any vector in Δ and the n_i run through the finite set $\Sigma_1 \subset \text{Hom}_{\mathbb{Z}}(M, \mathbb{Z})$ of primitive inward-pointing normals to the facets of C .

Remark 4.20. The degree of a chamber of constancy Δ is equal to the degree of the corresponding torus-invariant divisor D on the affine toric variety $X_\sigma = \text{Spec } R$ with the property that A_Δ is the module of global sections of the associated reflexive sheaf $\mathcal{O}_{X_\sigma}(D)$. See Remark 4.5. ■

The technical heart of our main argument proceeds by induction on the degree. The following is a crucial step:

Lemma 4.21. *Given chambers of constancy $\Delta \succeq \Delta'$, the following are equivalent.*

- (1) $\deg(\Delta) - \deg(\Delta') = 1$.
- (2) *The chain $\Delta \succ \Delta'$ is saturated in the poset of chambers of constancy; that is, if Δ'' is a chamber of constancy with $\Delta \succeq \Delta'' \succeq \Delta'$, then either $\Delta'' = \Delta$ or $\Delta'' = \Delta'$.*
- (3) Δ and Δ' are adjacent; that is, they are on opposite sides of a unique hyperplane.

Proof. The proof is straightforward when one considers the linear inequalities defining Δ and Δ' given by Proposition 4.3. In order to have their degrees differ by exactly one, the only possibility is that exactly one of these linear inequalities differ by exactly 1. So we have exactly one $n_i \in \Sigma_1$ such that Δ' has an (intersecting) supporting hyperplane defined by $\langle x, n_i \rangle = d$ (where $d = \lceil \langle v', n_i \rangle \rceil$ for $v' \in \Delta'$) and Δ has (intersecting) supporting hyperplane $\langle x, n_i \rangle = d + 1$; This means that Δ and Δ' are on opposite sides of their shared supporting hyperplane whose equation is $\langle x, n_i \rangle = d$, and no other module $A_{\Delta''}$ can be squeezed between A_Δ and $A_{\Delta'}$. □

4.5. Restriction to a Hyperplane. Let H be a supporting hyperplane of our rational polyhedral cone C in $M_{\mathbb{R}}$. Then $C \cap H$ is a polyhedral cone in the vector space H , rational with respect to the lattice $M \cap H$. The cone $C \cap H$ is full-dimensional and/or pointed (in H) if C is (in $M_{\mathbb{R}}$).

As usual, we index the facets of C by the set $\Sigma_1 = \{n_1, \dots, n_t\}$ of primitive inward-pointing normals. We can choose an appropriate hyperplane H so that the cone $C \cap H$ is one of these facets, say with corresponding normal n_t . The facets of

the cone $C \cap H$ are the intersections of the facets of C with H , and hence indexed by the (restrictions to H of the) linear functionals $\{n_1, \dots, n_{t-1}\}$. These are the primitive inward-pointing normals to the facets of $C \cap H$.

Consider the corresponding toric algebra of the cone $C \cap H$ in H :

$$S = \mathbb{k}[(C \cap H) \cap (M \cap H)] := \text{Span}\{x^m \mid m \in (C \cap H) \cap (M \cap H)\}.$$

There is a natural surjection of \mathbb{k} -algebras

$$R \twoheadrightarrow S \quad x^m \mapsto \begin{cases} x^m & \text{if } m \in H \\ 0 & \text{if } m \notin H, \end{cases}$$

whose kernel is the height one monomial ideal of R

$$(4.7) \quad P_{n_t} := \text{Span}\{x^m \mid \langle m, n_t \rangle > 0\}.$$

Proposition 4.22. *Let H be a supporting hyperplane of our full-dimensional rational polyhedral cone C in $M_{\mathbb{R}}$ and let*

$$R \twoheadrightarrow S$$

be the induced surjection of toric algebras corresponding to the rational polyhedral cones C and $C \cap H$, respectively. Then

- (1) *The toric S -modules are in one-to-one correspondence with toric R -modules of vectors v lying in the subspace H of $M_{\mathbb{R}}$.*
- (2) *The map $\Delta \mapsto \Delta \cap H$ induces a bijection between the chambers of constancy in $M_{\mathbb{R}}$ (indexing conic R -modules) that meet H and the chambers of constancy in H (indexing conic S -modules).*
- (3) *The bijection in (2) respects the partial order and the cell decomposition of chambers. In particular, the open cells of H are the same as the open cells of $M_{\mathbb{R}}$ contained in H .*

Proof. All statements follow immediately from the definitions and by applying Propositions 3.2, 3.5, and 4.3, describing chambers and cells in terms of the elements of Σ_1 . The point is that the restrictions to H of the linear functionals $\{n_1, \dots, n_{t-1}\}$ in Σ_1 for C is the set of primitive inward-pointing normals to $C \cap H$. \square

5. CHAIN COMPLEXES OF CONIC MODULES

Throughout this section, R denotes a toric algebra corresponding to a pointed full-dimensional rational polyhedral cone C in $M_{\mathbb{R}} = M \otimes \mathbb{R}$.

Fix a conic module A_{Δ} over R , with the chamber of constancy $\Delta \subset M_{\mathbb{R}}$. Our goal is to write down a complex K_{Δ}^{\bullet} of R -modules, abutting to A_{Δ} , and consisting only of direct sums of conic modules. In §6, we will construct from this complex

a projective resolution of a simple module over the endomorphism ring $\text{End}_R(\mathbb{A})$ where \mathbb{A} is a sum of conic modules.

5.1. The Chain Complex of a Chamber. Fix a chamber of constancy Δ . For each open cell of Δ , arbitrarily fix some orientation. Note that if τ_i is a cell of Δ of codimension $i \neq 0$, then τ_i is contained in the boundary of some cell τ_{i-1} of codimension $i-1$ of Δ . Our choice of orientation on τ_{i-1} induces an orientation on its boundary components, including τ_i , which may or may not agree with our choice of orientation for τ_i .

We define the **chain complex of the chamber Δ** as the complex K_Δ^\bullet of M -graded R -modules

$$(5.1) \quad \cdots \longrightarrow \bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau)=2}} \mathring{A}_\tau \longrightarrow \bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau)=1}} \mathring{A}_\tau \longrightarrow \bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau)=0}} \mathring{A}_\tau = A_\Delta$$

where each map in the complex is a signed sum of signed inclusion maps $\mathring{A}_{\tau_{i+1}} \hookrightarrow \mathring{A}_{\tau_i}$ (whenever $\mathring{A}_{\tau_{i+1}} \subset \mathring{A}_{\tau_i}$) and zero maps (whenever $\mathring{A}_{\tau_{i+1}} \not\subset \mathring{A}_{\tau_i}$). The sign is determined by our chosen orientations: the sign is 1 if our chosen orientation on τ_{i+1} agrees with the orientation induced by our chosen orientation on τ_i by virtue of its being a boundary component of τ_i , and the sign is -1 otherwise.

Lemma 5.1. K_Δ^\bullet is an M -graded complex of R -modules.

Proof. We need only verify that K_Δ^\bullet is a complex, that is, that $d^2 = 0$. For this, we need to follow just one monomial $x^m \in A_{\tau_{i+1}}$ in one summand of an arbitrary term

$$\bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau)=i+1}} \mathring{A}_\tau$$

of K_Δ^\bullet . Under d , this monomial is sent to the list of signed monomials $\pm x^m$ and zeros in $\bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau)=i}} \mathring{A}_\tau$, according to whether or not $\tau_{i+1} \subset \partial\tau_i$ (and if so, whether or not the orientations agree). Applying d again, the reader can check that the signed monomials come in pairs so cancel out to produce $d^2(x^m) = 0$. Indeed, the calculation is exactly the same as that to verify that the cellular cochain complex of the associated CW complex of $\overline{\Delta}$ is a complex; to see this note that if τ and τ' are open cells of $\overline{\Delta}$ such that $\tau \subset \partial\tau'$, and if τ is a cell of Δ , then so is τ' . Thus the differential of the complex K_Δ^\bullet applied to some monomial x^m in some summand A_τ is the same as it would be in the cellular cohomology complex of the closed cell complex $\overline{\Delta}$. See [Mun84]. \square

Remark 5.2. The chain complex K_Δ^\bullet has length bounded by the vector space dimension of $M_{\mathbb{R}}$, which (because C is full dimensional) is the same as the Krull dimension of R . In general, the length of K_Δ^\bullet is equal to the codimension of the

smallest dimensional cell appearing in Δ . Example 5.3 shows that the length can be strictly smaller than the dimension of R because there can be chambers Δ with no zero-dimensional cells. ■

Example 5.3. Consider again the three-dimensional toric algebra R obtained from the cone over the square in \mathbb{R}^3 as in Example 4.14. This non-simplicial toric ring admits three isomorphism types of conic modules, whose (representative) chambers are pictured below in Figure 5, with boundary cells in each chamber colored darker. Note that the chamber of type (A) contains a cell of dimension zero (point) but the chambers of types (B) and (C) do not. Therefore the complexes K_{Δ}^{\bullet} corresponding




FIGURE 5. The three types of chambers with shaded boundary cells.

to chambers of type (A) will have length three while the length of the complexes corresponding to any other chamber will have length strictly less. The complex K_{Δ}^{\bullet} corresponding to the first type of chamber is

$$A_0 \longrightarrow A_0^{\oplus 4} \longrightarrow A_1^{\oplus 2} \oplus A_2^{\oplus 2} \longrightarrow A_0,$$

with M -graded structure and mappings given by signed inclusions between chambers, as given in Figure 6.

The complex of R -modules corresponding to the latter two types of chambers is

$$\begin{aligned} A_2 &\longrightarrow A_0^{\oplus 2} \longrightarrow A_1 \\ A_1 &\longrightarrow A_0^{\oplus 2} \longrightarrow A_2, \end{aligned}$$

with M -graded structure described by signed inclusions between chambers, as given in Figure 7. ■

5.2. The Acyclicity Lemma. Our eventual goal will be to apply the functor $\text{Hom}_R(\mathbb{A}, -)$ to complexes of the form K_{Δ}^{\bullet} in order to construct minimal projective resolutions of simple modules over the endomorphism ring $\text{End}_R(\mathbb{A})$, where \mathbb{A} is a finite sum of conic modules. The following **Acyclicity Lemma** is crucial.

Lemma 5.4 (The Acyclicity Lemma). *Fix a toric algebra R coming from a pointed full dimensional rational polyhedral cone $C \subset M_{\mathbb{R}}$. Let A_{Δ} and $A_{\Delta'}$ be two conic modules for R , with associated chambers of constancy Δ and Δ' .*

- (1) *The complex $\text{Hom}_R(A_{\Delta'}, K_{\Delta}^{\bullet})$ is exact if $A_{\Delta} \not\cong A_{\Delta'}$;*




FIGURE 6. The complex of an octahedral chamber




FIGURE 7. The complexes of tetrahedral chambers

- (2) If $A_\Delta \simeq A_{\Delta'}$, then $\text{Hom}_R(A_{\Delta'}, K_\Delta^\bullet)$ has one-dimensional homology, which appears (only) in homological degree zero and M -degree m , where m is the unique element of M such that $\Delta = m + \Delta'$.

Proof of the Acyclicity Lemma. Because the complex is M -graded, it suffices to prove acyclicity in each degree $m \in M$. It is enough to prove the Acyclicity Lemma only in degree $m = 0 \in M$, because of the identification of \mathbf{k} -vector spaces

$$[\text{Hom}_R(A_{\Delta'}, A_{\tilde{\Delta}})]_m = [\text{Hom}_R(x^m A_{\Delta'}, A_{\tilde{\Delta}})]_0 = [\text{Hom}_R(A_{\Delta'+m}, A_{\tilde{\Delta}})]_0$$

for all $m \in M$ and all chambers $\Delta', \tilde{\Delta}$.

We prove the Acyclicity Lemma in M -degree 0 in two cases:

- (i) $\Delta = \Delta'$
- (ii) $\Delta \neq \Delta'$

Case (i): Since $\Delta = \Delta'$ and all the modules appearing in the terms of K_Δ^\bullet are of the form \mathring{A}_τ , where τ is a cell of Δ , we have

$$A_\Delta \supset \mathring{A}_\tau;$$

the inclusion is *proper* unless τ has codimension zero, in which case $A_\Delta = \mathring{A}_\tau$. So there are no degree preserving maps $A_{\Delta'} = A_\Delta \rightarrow \mathring{A}_\tau$ except in homological degree zero. That is, the entire chain complex $[\text{Hom}_R(A_{\Delta'}, K_\Delta^\bullet)]_0$ reduces to the zero complex, except in homological degree zero, where we have $[\text{Hom}_R(A_\Delta, A_\Delta)]_0 = k$. This completes the proof of Case (i).

Case (ii): Suppose $\Delta \neq \Delta'$. To show that the complex $[\text{Hom}_R(A_{\Delta'}, K_\Delta^\bullet)]_0$ is exact, we first examine this complex of k -vector spaces carefully. From Proposition 3.2(3), we have that

$$(5.2) \quad [\text{Hom}_R(A_{\Delta'}, A_\Delta)]_0 = \begin{cases} k & \text{if } A_{\Delta'} \subset A_\Delta \\ 0 & \text{otherwise.} \end{cases}$$

This implies

Lemma 5.5. *The zero-th graded piece $[\text{Hom}_R(A_{\Delta'}, K_\Delta^\bullet)]_0$ of the M -graded complex $\text{Hom}_R(A_{\Delta'}, K_\Delta^\bullet)$ is*

$$\begin{array}{ccccc} \cdots & \longrightarrow & \bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau) = 2 \\ A_{\Delta'} \subset \mathring{A}_\tau}} k & \longrightarrow & \bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau) = 1 \\ A_{\Delta'} \subset \mathring{A}_\tau}} k & \longrightarrow & \bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau) = 0 \\ A_{\Delta'} \subset \mathring{A}_\tau}} k \end{array}$$

consisting only of those summands indexed by cells τ of Δ for which $A_{\Delta'} \subset \mathring{A}_\tau$.

Proceeding with the proof of Case (ii), observe that without loss of generality, we may assume $A_\Delta \supset A_{\Delta'}$, because otherwise, there are no degree-preserving maps $A_{\Delta'} \rightarrow \mathring{A}_\tau$ for $\tau \subset \Delta$, so the entire complex is zero. So fix Δ and Δ' with $\Delta' \prec \Delta$. We induce on the quantity

$$\deg \Delta - \deg \Delta',$$

where $\deg \Delta$ denotes the degree of the chamber Δ as defined in Definition 4.19. Our assumption that $\Delta' \prec \Delta$ but $\Delta \neq \Delta'$ ensures that this difference is at least 1.

5.2.1. Base Case of Induction. Assume $\deg \Delta - \deg \Delta' = 1$. By Lemma 4.21, it follows that $\Delta' \prec \Delta$ is a saturated chain in the poset of chambers, and Δ and Δ' are on opposite sides of a supporting hyperplane \mathcal{H} of Δ . This means that $A_{\Delta'} = \mathring{A}_\tau$ where τ is the interior cell of the facet of $\Delta \cap \mathcal{H}$. So $A_{\Delta'}$ is a submodule of A_Δ and of \mathring{A}_τ but not of any other $\mathring{A}_{\tau'}$ appearing in the chain complex K_Δ^\bullet . So $[\text{Hom}_R(A_{\Delta'}, K_\Delta^\bullet)]_0$ degenerates to the complex with non-zero terms only in homological degrees 0 and 1:

$$[\text{Hom}_R(A_{\Delta'}, A_\Delta)]_0 \longrightarrow [\text{Hom}_R(A_{\Delta'}, A_{\Delta'})]_0$$

both of which are k . This complex is the isomorphism $k \longrightarrow k$, hence exact.

5.2.2. Inductive Step. We now fix $d > 0$ and assume that we have proved the Acyclicity Lemma in degree $0 \in M$ on *all* toric varieties and all chambers $\Delta' \prec \Delta$ satisfying $\deg \Delta - \deg \Delta' < d$.

Given $\Delta \succ \Delta'$ with $\deg \Delta - \deg \Delta' = d \geq 2$, use Lemma 4.21 to find a chamber Δ'' such that

$$\Delta \succ \Delta'' \succ \Delta'$$

with the chain $\Delta'' \succ \Delta'$ *saturated*. Let \mathcal{H} be the supporting hyperplane separating the chambers Δ'' and Δ' .

Consider the inclusion of conic modules $A_{\Delta'} \subset A_{\Delta''}$. The cokernel is annihilated by the ideal P_{n_t} generated by monomials x^m such that $\langle m, n_t \rangle > 0$, where n_t is the inward-pointing normal of \mathcal{H} . Thus the cokernel is naturally a module over the ring R/P_{n_t} , the toric algebra S of the cone $C \cap H$ in the subspace H , the supporting hyperplane of C with inward pointing normal n_t (parallel to \mathcal{H}). We claim that this cokernel complex can be understood as the complex of a chamber of constancy over the toric algebra S :

Claim: *There is a short exact sequence of M -graded complexes*

$$0 \longrightarrow \mathrm{Hom}_R(A_{\Delta''}, K_{\Delta}^{\bullet}) \hookrightarrow \mathrm{Hom}_R(A_{\Delta'}, K_{\Delta}^{\bullet}) \longrightarrow \mathrm{Hom}_S(A_{\delta''}, K_{\delta}^{\bullet})[1] \longrightarrow 0$$

where S is the toric algebra of the cone $C \cap H$ and $\delta'', \delta \subset H$ are the chambers of constancy for S -modules as described by Proposition 4.22:

$$\delta := (\Delta - m) \cap H \quad \delta'' := (\Delta'' - m) \cap H.$$

Here $m \in \mathcal{H} \cap M$ is any lattice point used to translate \mathcal{H} to the subspace H , and the notation $C^{\bullet}[1]$ denotes a homological shift of the complex C^{\bullet} .

Before establishing the claim, we note its sufficiency to complete the proof of the Acyclicity Lemma. Indeed, since both $\deg \Delta - \deg \Delta''$ and $\deg \delta - \deg \delta''$ are precisely one less than $\deg \Delta - \deg \Delta'$, the inductive assumption implies that both complexes $\mathrm{Hom}_R(A_{\Delta''}, K_{\Delta}^{\bullet})$ and $\mathrm{Hom}_S(A_{\delta''}, K_{\delta}^{\bullet})$ are exact in degree zero. Whence so is the complex in the middle.

To prove the claim, first note that by restricting any map $A_{\Delta''} \rightarrow K_{\Delta}^{\bullet}$ to $A_{\Delta'}$, we get an inclusion of M -graded complexes of R -modules

$$\mathrm{Hom}_R(A_{\Delta''}, K_{\Delta}^{\bullet}) \hookrightarrow \mathrm{Hom}_R(A_{\Delta'}, K_{\Delta}^{\bullet}).$$

The cokernel \mathcal{C}^\bullet is annihilated by the prime ideal P_{n_t} and hence is a complex of modules over the toric algebra S . Its degree zero subcomplex is

$$(5.3) \quad \cdots \longrightarrow \bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau) = 2 \\ A_{\Delta'} \subset \mathring{A}_\tau \\ \tau \subset \mathcal{H}}} \mathbf{k} \longrightarrow \bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau) = 1 \\ A_{\Delta'} \subset \mathring{A}_\tau \\ \tau \subset \mathcal{H}}} \mathbf{k} \longrightarrow \bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau) = 0 \\ A_{\Delta'} \subset \mathring{A}_\tau \\ \tau \subset \mathcal{H}}} \mathbf{k}$$

because of the following lemma:

Lemma 5.6. *Let $\Delta, \Delta', \Delta''$ be chambers of constancy such that $\Delta \succ \Delta'' \succ \Delta'$, with Δ' and Δ'' adjacent on either side of hyperplane \mathcal{H} . Let τ be an open cell of Δ . The natural map*

$$\left[\text{Hom}_R(A_{\Delta''}, \mathring{A}_\tau) \right]_0 \longrightarrow \left[\text{Hom}_R(A_{\Delta'}, \mathring{A}_\tau) \right]_0$$

induced by restricting maps from $A_{\Delta''}$ to the submodule $A_{\Delta'}$ is either an isomorphism (when $\tau \not\subset \mathcal{H}$) or the zero map (when $\tau \subset \mathcal{H}$) because the source is zero.

Proof. The statement amounts to saying that an inclusion $A_{\Delta'} \subset \mathring{A}_\tau$ fails to factor as

$$A_{\Delta'} \subset A_{\Delta''} \subset \mathring{A}_\tau$$

if and only if $\tau \subset \mathcal{H} \cap \Delta$.

To see this, note that because Δ' and Δ'' lie on opposite sides of the single separating hyperplane \mathcal{H} , a monomial x^m is in $A_{\Delta''}$ but not $A_{\Delta'}$ if and only if $m \in \mathcal{H} \cap M$. So any inclusion $A_{\Delta'} \subset \mathring{A}_\tau$ will extend to $A_{\Delta''}$ unless some $m \in \mathcal{H} \cap M$ is not in $\mathring{C} + v$ where $v \in \tau$. This happens if and only if \mathcal{H} is a supporting hyperplane of the translated cone $C + v$, that is, if and only if $v \in \mathcal{H}$. \square

Finally, we need to show that the cokernel complex $[\mathcal{C}^\bullet]_0$ is, up to homological shift, the same the complex of conic modules over the toric algebra $S = \mathbf{k}[(C \cap H) \cap (M \cap H)]$ described in the claim. Pick any $m \in \mathcal{H} \cap M$, and consider the translation

$$M_{\mathbb{R}} \rightarrow M_{\mathbb{R}} \quad \text{sending} \quad x \mapsto x - m.$$

This map translates \mathcal{H} to H , and respects the partial ordering on chambers as well as the CW decomposition of $M_{\mathbb{R}}$. In particular, the chambers $\Delta \succ \Delta'' \succ \Delta'$ are translated to chambers $\delta \succ \delta'' \succ \delta'$, and every open cell of Δ is translated to a corresponding open cell of δ . Intersecting with the subspace H , we have

$$\delta \cap H \succ \delta'' \cap H \succ \delta' \cap H = \emptyset$$

and now the chambers $\delta \cap H \succ \delta'' \cap H$ are chambers of constancy for the conic modules over S by Proposition 4.22; likewise the translations of the open cells $\tau \in \mathcal{H}$ become the open cells of the conic-induced CW decomposition of H . Thus the cokernel complex $[\mathcal{C}^\bullet]_0$ from (5.3) above can be identified with the zero-th

graded piece of the complex of S -modules

$$\mathrm{Hom}_S(A_{\delta''}, K_{\delta}^{\bullet}).$$

The proof is complete. \square

6. GLOBAL DIMENSION OF ENDOMORPHISM ALGEBRAS

Throughout this section, R denotes a toric algebra defined by a pointed full dimensional cone $C \subset M_{\mathbb{R}}$. For any ring Λ , the **global dimension** $\mathrm{gl.dim}(\Lambda)$ is the maximum projective dimension (possibly infinite) of any left or right Λ -module.

A fundamental theorem of Auslander-Buchsbaum and Serre states that a *commutative* Noetherian local ring Λ has finite global dimension if and only if it is regular. When Λ is not regular, it may still admit a finitely generated faithful module M such that the ring $\mathrm{End}_{\Lambda}(M)$ has finite global dimension. Such an endomorphism ring is called a **non-commutative resolution of singularities** of the commutative ring Λ .

The main result of this section is a constructive proof that toric rings admit natural non-commutative resolutions:

Theorem 6.1. *Let $\mathbb{A} = A_{\Delta_1} \oplus \cdots \oplus A_{\Delta_s}$, where the A_{Δ_i} run through each isomorphism class of conic R -modules. Then the global dimension of $\mathrm{End}_R(\mathbb{A})$ is equal to the Krull dimension of R .*

As an immediate corollary, we deduce that the same is true for any Morita equivalent ring. In particular, invoking Proposition 4.15,

Corollary 6.2. *For any natural number q , consider the R -module*

$$R^{1/q} = \mathbf{k}\left[\frac{1}{q}M \cap C\right].$$

Then provided that q is sufficiently large⁵, the global dimension of $\mathrm{End}_R(R^{1/q})$ is equal to the dimension of R .

From Corollary 6.2, we deduce also that in prime characteristic p , the ring of differential operators $D_{\mathbf{k}}(R)$ on a toric ring has finite global dimension; see Theorem 6.13. We do not know if the same is true in characteristic zero.

The rest of this section is devoted to the proofs of these results, as well as Theorem 6.13 on the finite global dimension of $D_{\mathbf{k}}(R)$ for toric R over a perfect field of characteristic p .

⁵In practice, it is easy to see how large is sufficiently large; see the proof of Proposition 4.15.

6.1. Modules over Endomorphism Rings. We review some of the general module theory for endomorphism rings in our special setting at hand. See also [SVdB97, 4.1] or [HGK04, 11.1].

Fix an R -module \mathbb{A} , of the form

$$(6.1) \quad \mathbb{A} := \bigoplus_{i=1}^m A_{\Delta_i}^{d_i},$$

where the d_i are positive integers and every conic module is isomorphic to a unique A_{Δ_i} appearing in the sum. We call such a module \mathbb{A} a **complete sum of conic modules**. Our motivating example is $R^{1/q}$, which is isomorphic to a complete sum of conic modules when q is large enough (Proposition 4.15). When we choose an isomorphism from $R^{1/q}$ to a complete sum of conic modules, we make a choice of the representatives A_{Δ_i} appearing in the sum. The resulting M -gradings on $R^{1/q}$ and its endomorphism ring depend on this choice, and so they should be regarded as artificial (but very useful!).

Consider the endomorphism algebra $\text{End}_R(\mathbb{A})$. Because \mathbb{A} is finitely generated, the ring $\text{End}_R(\mathbb{A})$ is Noetherian. Thus the maximal length of a minimal projective resolution in the category of left and right $\text{End}_R(\mathbb{A})$ -modules is the same, see [Aus55]. Therefore, we focus attention on the *right* modules over $\text{End}_R(\mathbb{A})$.

6.2. Right modules over $\text{End}_R(\mathbb{A})$. For any R -module B , $\text{Hom}_R(\mathbb{A}, B)$ is naturally a right $\text{End}_R(\mathbb{A})$ -module with action given by composition on the first factor. This induces a functor

$$\text{Hom}_R(\mathbb{A}, -) : \text{Mod}(R) \longrightarrow \text{Mod}(\text{End}_R(\mathbb{A}))$$

where $\text{Mod}(\text{End}_R(\mathbb{A}))$ denotes the category of right $\text{End}_R(\mathbb{A})$ -modules. This functor restricts to an equivalence between the full subcategories

$$(6.2) \quad \text{add}(\mathbb{A}) \xrightarrow{\sim} \text{proj}(\text{End}_R(\mathbb{A}))$$

of finite direct sums of (graded) summands of \mathbb{A} and finitely generated (graded) projective right $\text{End}_R(\mathbb{A})$ modules. To see this, observe that the functor $G_* := \text{Hom}_R(\mathbb{A}, -)$ has left adjoint $G^* := - \otimes_{\text{End}_R(\mathbb{A})} \mathbb{A}$, and that the adjunction maps define isomorphisms

$$G^* G_*(\mathbb{A}) \xrightarrow{\sim} \mathbb{A} \quad \text{and} \quad \text{End}_R(\mathbb{A}) \xrightarrow{\sim} G_* G^*(\text{End}_R(\mathbb{A})) .$$

Both functors respect the M -grading and commute with direct sums, so they induce an equivalence $\text{add}(\mathbb{A}) \simeq \text{add}(\text{End}_R(\mathbb{A}))$, the latter of which equals $\text{proj}(\text{End}_R(\mathbb{A}))$. By restricting to the M -graded category, we have a functorial bijection between conic R -modules and indecomposable projective M -graded modules over $\text{End}_R(\mathbb{A})$. Summarizing and continuing, we have

Proposition 6.3. *For any Δ ,*

- (1) *The $\text{End}_R(\mathbb{A})$ -module*

$$P_\Delta := \text{Hom}_R(\mathbb{A}, A_\Delta)$$

is a graded indecomposable projective module, and every graded indecomposable projective $\text{End}_R(\mathbb{A})$ -module is isomorphic to a module of this form.

- (2) *The projective module P_Δ has a unique maximal graded submodule. Every graded simple $\text{End}_R(\mathbb{A})$ -module is isomorphic to a quotient of some P_Δ by this maximal submodule.*

We let S_Δ denote the unique graded simple quotient of P_Δ .

Although Proposition 6.3 can be found in the literature (eg. see [SVdB97, Lemma 4.1.1] or [HGK04, Prop. 11.1.1]), we keep the paper self-contained and our proof easy to follow by explicitly describing the unique maximal (right) submodule of P_Δ —called the *radical* of P_Δ —in Section 6.3 below.

Remark 6.4. The word *graded* in Proposition 6.3 above refers to the natural M -grading on all the objects; all maps are homogeneous in this grading. However, because C is pointed, the toric ring R can also be given a (non-canonical) connected \mathbb{Z} -grading by fixing a group homomorphism $M \rightarrow \mathbb{Z}$ taking strictly positive values on $M \cap (C \setminus \{0\})$. This grading induces compatible gradings on conic modules, on \mathbb{A} , and on $\text{End}_R(\mathbb{A})$. Both statements in Proposition 6.3 are valid for this connected \mathbb{Z} -grading as well. The same is true of Theorem 6.5 below. \blacksquare

The following theorem (proven in the next section) states that the functor G_* sends the complex K_Δ^\bullet to a projective resolution of the simple quotient module S_Δ .

Theorem 6.5. *The complex $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet)$ is a graded $\text{End}_R(\mathbb{A})$ -projective resolution of the simple module S_Δ , and this projective resolution has minimal length.*

6.3. Radicals. We review some general theory in preparation to prove Theorem 6.5. By definition, a homogeneous R -module homomorphism $f : A_\Delta \rightarrow A_{\Delta'}$ is **radical** if it is *not* an isomorphism. The radical homomorphisms generate a graded R -submodule $\text{Rad}(A_\Delta, A_{\Delta'}) \subseteq \text{Hom}_R(A_\Delta, A_{\Delta'})$. By Proposition 4.17,

$$\text{Rad}(A_\Delta, A_{\Delta'}) = \text{Span}\{x^m \mid m \in M \text{ such that } \Delta + m \prec \Delta'\},$$

where the symbol \prec indicates *strictly less* in the poset on chambers. In particular, $\text{Hom}_R(A_\Delta, A_{\Delta'})/\text{Rad}(A_\Delta, A_{\Delta'})$ is one dimensional over k if $A_\Delta \simeq A_{\Delta'}$ and zero otherwise.

More generally, we may represent a graded homomorphism from a direct sum \mathbb{A} of n conic modules A_i to a direct sum \mathbb{B} of m conic modules B_j by an $m \times n$ matrix $[\phi_{ij}]$ where $\phi_{ij} \in \text{Hom}_R(A_j, B_i)$; such a homomorphism is **radical** if each of its

entries ϕ_{ij} is.⁶ In particular, the elements of $\text{Hom}_{\text{End}_R(\mathbb{A})}(\mathbb{A}, A_\Delta)$ are represented by row vectors with entries in $\text{Hom}_R(A_i, A_\Delta)$; the homogenous maps in its radical have entries in the radical of each $\text{Hom}_R(A_i, A_\Delta)$. This is clearly a maximal graded right submodule: if some homogeneous row matrix (ϕ_j) is not in $\text{Rad}(\mathbb{A}, A_\Delta)$, then some entry $\phi_j \in \text{Hom}_R(A_j, A_\Delta)$ is a graded isomorphism. Since elements of $\text{End}_R(\mathbb{A})$ act on the right by column operations on the row vector (ϕ_j) , the right module generated by (ϕ_j) contains the row vector $(\psi_j) \in \text{Hom}_R(\mathbb{A}, A_\Delta)$ which has zeros in every spot except the j -th, which is the identity map. Since this idempotent is a $\text{End}_R(\mathbb{A})$ generator for the right module $\text{Hom}_{\text{End}_R(\mathbb{A})}(\mathbb{A}, A_\Delta)$, we conclude that $\text{Rad}(\mathbb{A}, A_\Delta)$ is maximal.

Put differently, $\text{End}_R(\mathbb{A})$ is *graded semiperfect* in the language of [Däs92].

Proof of Theorem 6.5. Fix a chamber Δ , and corresponding simple $\text{End}_R(\mathbb{A})$ -module S_Δ . Recall that $K_\Delta^0 = A_\Delta$ by construction. It follows that the $\text{Hom}_R(\mathbb{A}, K_\Delta^0) = P_\Delta$, and so there is a natural surjection $\text{Hom}_R(\mathbb{A}, K_\Delta^0) \rightarrow S_\Delta$.

The natural surjection $\text{Hom}_R(\mathbb{A}, K_\Delta^0) = P_\Delta \rightarrow S_\Delta$ makes the complex $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet)$ into a projective resolution of S_Δ . Indeed, since each K_Δ^i is a direct sum of conic modules, each $\text{Hom}_R(\mathbb{A}, K_\Delta^i)$ is a projective module over $\text{End}_R(\mathbb{A})$. It remains to show that this complex is *exact*, and that there is no shorter projective resolution.

The acyclicity Lemma 5.4 tells us that this complex $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet)$ is exact except possibly at the penultimate spot

$$\bigoplus_{\substack{\text{cells } \tau \in \Delta \\ \text{codim}(\tau) = 1}} \text{Hom}_R(\mathbb{A}, \mathring{A}_\tau) \rightarrow \text{Hom}_R(\mathbb{A}, A_\Delta) = P_\Delta \rightarrow S_\Delta .$$

Since the kernel of the natural projection map $P_\Delta \rightarrow S_\Delta$ is the radical of P_Δ , we need to check that this is the image of the incoming map. But this is immediate from Lemma 5.4(2).

It remains to show that S_Δ has no shorter projective resolution. This follows from the following proposition, whose proof will appear below:

Proposition 6.6. *For any pair of chambers of constancy Δ and Δ' ,*

$$\dim(\text{Ext}_{\text{End}_R(\mathbb{A})}^i(S_\Delta, S_{\Delta'})) = \# \text{ of codim. } i \text{ cells } \tau \text{ in } \Delta \text{ with } \mathring{A}_\tau \simeq A_{\Delta'}$$

To see that $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet)$ is a minimal projective resolution of S_Δ , let τ be a cell of maximal codimension in Δ , and let Δ' be the chamber of constancy such that $A_{\Delta'} = \mathring{A}_\tau$. The length of $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet)$ is $\text{codim}(\tau)$, and from the proof of Proposition 6.6 it follows that

$$\text{Ext}_{\text{End}_R(\mathbb{A})}^{\text{codim}(\tau)}(S_\Delta, S_{\Delta'}) \neq 0 .$$

⁶Here, we consider the elements of \mathbb{A} and \mathbb{B} as column vectors with entries in A_i and B_j respectively, and the matrix acts on the left.

Therefore, S_Δ cannot have a projective resolution shorter than $\text{length codim}(\tau)$. \square

Proof of Proposition 6.6. Using the resolution $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet) \rightarrow S_\Delta$ to compute the Ext groups, this Ext group is the i th cohomology of the complex

$$\text{Hom}_{\text{End}_R(\mathbb{A})}(\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet), S_{\Delta'})$$

By Lemma 6.7 below, this is equal to

$$\bigoplus_{\tau \in \Delta} \text{Hom}_R(\mathring{A}_\tau, A_{\Delta'}) / \text{Rad}(\mathring{A}_\tau, A_{\Delta'})$$

as a graded vector space. On each component, the boundary map is given by a sum of compositions with proper homogeneous injections, and so consequently every component of the image is 0. Hence, the boundary map is zero on the above graded vector space. The subspace of cohomological degree i corresponds to τ of codimension i . As the τ -indexed summand is 1-dimensional if $\mathring{A}_\tau \simeq A_{\Delta'}$ and is 0 otherwise, the dimension of Ext counts the desired τ . This will conclude the proof of Proposition 6.6 as soon as we prove Lemma 6.7. \square

Lemma 6.7. *For any Δ and Δ' , there is a graded k -vector space isomorphism*

$$\text{Hom}_{\text{End}_R(\mathbb{A})}(P_\Delta, S_{\Delta'}) \simeq \text{Hom}_R(A_\Delta, A_{\Delta'}) / \text{Rad}(A_\Delta, A_{\Delta'})$$

These spaces are dimension one or zero, depending on whether $A_\Delta \simeq A_{\Delta'}$ or not.

Proof. By the projectivity of P_Δ , there is a short exact sequence

$$0 \rightarrow \text{Hom}(P_\Delta, \text{Rad}(P_{\Delta'})) \rightarrow \text{Hom}(P_\Delta, P_{\Delta'}) \rightarrow \text{Hom}(P_\Delta, S_{\Delta'}) \rightarrow 0$$

The equivalence of categories (6.2) induced by $\text{Hom}_R(\mathbb{A}, -)$ guarantees that every homogenous map $g : P_\Delta \rightarrow P_{\Delta'}$ is given by applying $\text{Hom}_R(\mathbb{A}, -)$ to a homogenous map $g' : A_\Delta \rightarrow A_{\Delta'}$. The composition $\mathbb{A} \xrightarrow{f} A_\Delta \xrightarrow{g'} A_{\Delta'}$ is in $\text{Rad}(P_{\Delta'})$ for all f if and only if $g' \in \text{Rad}(A_\Delta, A_{\Delta'})$. Hence, the isomorphism $\text{Hom}(A_\Delta, A_{\Delta'}) \simeq \text{Hom}(P_\Delta, P_{\Delta'})$ induces a bijection between $\text{Rad}(A_\Delta, A_{\Delta'})$ and the image of $\text{Hom}(P_\Delta, \text{Rad}(P_{\Delta'}))$. By the above short exact sequence, $\text{Hom}(P_\Delta, S_{\Delta'}) \simeq \text{Hom}(A_\Delta, A_{\Delta'}) / \text{Rad}(A_\Delta, A_{\Delta'})$. \square

Remark 6.8. The resolution $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet)$ is ‘minimal’ in a stronger sense. Specifically, it lifts to a homogeneous split injection $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet) \hookrightarrow P^\bullet$ along any graded projective resolution $P^\bullet \rightarrow S_\Delta$. This is the graded analog of a ‘minimal projective resolution’, and can be proven by observing the morphisms in $\text{Hom}_R(\mathbb{A}, K_\Delta^\bullet)$ have coordinates in $\text{Rad}(\text{End}_R(\mathbb{A}))$ in an appropriate sense. \blacksquare

Corollary 6.9. *The projective dimension of S_Δ is the maximum codimension of cells in Δ . In particular, $\text{pdim}(S_\Delta) \leq \text{rank}(M)$, with equality when A_Δ is free.*

Example 6.10. Consider again the three-dimensional toric algebra R coming from the cone over the square in \mathbb{R}^3 , cf. Example 4.14. We have already calculated the three types of conic complexes for the conic modules A_0, A_1, A_2 . Now consider $\mathbb{A} = A_0 \oplus A_1 \oplus A_2$ and $\text{End}_R(\mathbb{A})$. This ring has three graded simples $S_{A_i} = \text{Hom}_R(\mathbb{A}, A_i)/\text{Rad}(\mathbb{A}, A_i)$, $i = 0, 1, 2$. We obtain projective resolutions for the simple modules S_{A_i} by applying the functor $\text{Hom}_R(\mathbb{A}, -)$ to the conic complexes $K_{\Delta_i}^\bullet$. Thus we see that the endomorphism ring $\text{End}_R(\mathbb{A})$ has two graded simple modules of projective dimension 2 defined by the tetrahedral chambers, and one graded simple of projective dimension 3 defined by the octahedral chamber. The same is true for the Morita equivalent ring $\text{Hom}_R(R^{1/q}, R^{1/q})$, for $q \gg 0$. ■

We can finally prove the opening theorem of the section.

Proof of Theorem 6.1. The bound $\text{gl.dim}(\text{End}_R(\mathbb{A})) \geq \text{rank}(M)$ follows from Corollary 6.9.

Since $\text{End}_R(\mathbb{A})$ is a finitely generated module over a commutative Noetherian ring R , a theorem of Auslander [Aus55] implies the left and right global dimensions coincide, and a subsequent theorem of Bass [Bas68, III, Prop. 6.7] implies the following.

$$\text{gl.dim}(\text{End}_R(\mathbb{A})) = \sup\{\text{pdim}_{\text{End}_R(\mathbb{A})}(S) \mid S \text{ is a simple right } \text{End}_R(\mathbb{A})\text{-module}\}$$

The remainder of the proof will reduce to *graded* simple right $\text{End}_R(\mathbb{A})$ -modules.

Let S be a simple right $\text{End}_R(\mathbb{A})$ -module (with no grading). The annihilator $\text{Ann}(S)$ in $\text{End}_R(\mathbb{A})$ is a two-sided maximal ideal, and so $\mathfrak{m} := \text{Ann}(S) \cap R$ is a maximal ideal in R . This implies R/\mathfrak{m} is a field, and S is a finite dimensional vector space over R/\mathfrak{m} . Since R/\mathfrak{m} must be a finite extension of the ground field k , S must also be finite dimensional over k .

Next, fix a connected \mathbb{Z} -grading on R , which induces a \mathbb{Z} -grading on $\text{End}_R(\mathbb{A})$ (see Remark 6.4). This \mathbb{Z} -grading may be weakened to a descending \mathbb{Z} -filtration, by setting

$$\mathcal{F}_i(\text{End}_R(\mathbb{A})) := \bigoplus_{j \geq i} \text{End}_R(\mathbb{A})_{\deg=j}$$

Note that the associated graded algebra $gr(\text{End}_R(\mathbb{A}))$ is canonically identified with $\text{End}_R(\mathbb{A})$.

Choose a surjection $\text{End}_R(\mathbb{A}) \rightarrow S$, which induces a compatible filtration on S . Since the filtrations on $\text{End}_R(\mathbb{A})$ and S are *left-limited*, a theorem of [NVO79, Cor. 7.6] implies that

$$\text{pdim}_{\text{End}_R(\mathbb{A})}(S) \leq \text{pdim}_{\text{End}_R(\mathbb{A})}(gr(S)) .$$

The associated graded module $gr(S)$ is still finite dimensional over k , and so it has a finite composition sequence as a \mathbb{Z} -graded $\text{End}_R(\mathbb{A})$ -module, whose subquotients are \mathbb{Z} -graded simple $\text{End}_R(\mathbb{A})$ -modules. As the projective dimension of $gr(S)$ is bounded by the maximum projective dimension of the subquotients (see e.g. the proof of Bass' theorem [Bas68, III, Prop. 6.7]), we deduce that

$$\text{pdim}(S) \leq \text{pdim}(gr(S)) \leq \sup_{\Delta} \{\text{pdim}(S_{\Delta})\} = \text{rank}(M)$$

where the last equality is Corollary 6.9. \square

Remark 6.11. Our results carry over to the complete case: if \widehat{R} is the completion of a pointed toric algebra R at its maximal graded ideal, then $\text{End}_{\widehat{R}}(\widehat{R} \otimes \mathbb{A})$ has global dimension equal to the dimension of \widehat{R} . This follows from a general properties about the behavior of global dimension under completion; cf. e.g. [SVdB17b, 3.4]. \blacksquare

6.4. Global dimension of differential operators in positive characteristic.

Given a commutative algebra R over a field k , the algebra of **differential operators on R over k** is defined as (see e.g. the discussion in [SVdB97, 2.1])

$$D_k(R) := \bigcup_{n \in \mathbb{N}} D_k^n(R)$$

where $D_k^0(R) := R$ and

$$D_k^n(R) := \{\theta \in \text{End}_k(R) \mid \text{for all } r \in R, [r, \theta] \in D_k^{n-1}(R)\}.$$

If R is the coordinate ring of a smooth complex variety, then $D_{\mathbb{C}}(R)$ is the classical ring of regular differential operators generated by multiplication operators and derivatives along vector fields.

When k has positive characteristic, the ring $D_k(R)$ exhibits several behaviors without a natural geometric analog. Most relevant for us is the following.

Lemma 6.12. [Yek92] *Let k be a perfect field of characteristic p , and let R be finitely generated as an R^p -module. Then, as subalgebras of $\text{End}_k(R)$,*

$$\bigcup_{e \in \mathbb{N}} \text{End}_{R^{p^e}}(R) = D_k(R).$$

That is, every R^{p^e} -linear map from R to itself is a differential operator, and every differential operator is R^{p^e} -linear for sufficiently large e . Note that $D_k^e(R)$ almost never equals $\text{End}_{R^{p^e}}(R)$, and $D_k^e(R)$ is almost never an algebra.

Theorem 6.13. *Let R be a pointed toric ring of dimension d over a perfect field k of prime characteristic p . Then the ring $D_k(R)$ of differential operators on R has finite global dimension.*

Proof. Let R^{1/p^e} denote the ring of p^e -th roots of elements in R . The Frobenius map $F^e : R \rightarrow R^{p^e}$ induces an isomorphism

$$\mathrm{End}_R(R^{1/p^e}) \xrightarrow{\sim} \mathrm{End}_{R^{p^e}}(R) .$$

Consequently, Corollary 6.2 implies that, for e sufficiently large,

$$\mathrm{gl.dim}(\mathrm{End}_{R^{p^e}}(R)) = \mathrm{gl.dim}(\mathrm{End}_R(R^{1/p^e})) = d .$$

By a theorem of Berštein [Ber58],

$$\mathrm{r.gl.dim}(D_{\mathbf{k}}(R)) = \mathrm{rgldim}\left(\bigcup_{e \in \mathbb{N}} \mathrm{End}_{R^{p^e}}(R)\right) \leq 1 + \lim_{e \in \mathbb{N}} (\mathrm{rgldim}(\mathrm{End}_{R^{p^e}}(R))) = d+1$$

A symmetric application of Berštein's theorem implies that $\mathrm{l.gl.dim}(D_{\mathbf{k}}(R)) \leq d+1$ as well, implying that $\mathrm{gl.dim}(D_{\mathbf{k}}(R)) \leq d+1$. \square

Remark 6.14. The proof implies that $\mathrm{gl.dim}(D_{\mathbf{k}}(R)) \leq d+1$. We suspect that in fact $\mathrm{gl.dim}(D_{\mathbf{k}}(R)) = d$, but as yet have not proved it. If R is smooth over \mathbf{k} , then Paul Smith's theorem [Smi87] implies that $\mathrm{gl.dim}(D_{\mathbf{k}}(R)) = d$. The general case would follow if we could show that for $q_0 < q$ sufficiently large, then $\mathrm{End}_R(R^{1/q})$ is flat (as a left or right) module over $\mathrm{End}_R(R^{1/q_0})$. \blacksquare

7. NON-COMMUTATIVE CREPANT RESOLUTIONS

Non-commutative *crepant* resolutions (NCCRs) were introduced by Michel Van den Bergh in his foundational paper [VdB04]. A natural question arises now that we have shown $\mathrm{End}_R(\mathbb{A})$ is a non-commutative resolution: when are these resolutions crepant?

First we recall the definition.

Definition 7.1. Let S be a normal Cohen-Macaulay domain of dimension d , and let B be a finitely generated reflexive S -module. The algebra $\mathrm{End}_S(B)$ is a **non-commutative crepant resolution (NCCR)** of S if every simple $\mathrm{End}_S(B)$ -module has projective dimension d .

Van den Bergh first defined NCCRs only for Gorenstein normal domains S ; in this case, he showed that $\mathrm{End}_S(B)$ is an NCCR if and only if $\mathrm{End}_S(B)$ has finite global dimension and is Cohen-Macaulay as an S -module; see [VdB04, Lemma 4.2]. If we assume only that S is a normal Cohen-Macaulay domain, Iyama and Wemyss showed that Van den Bergh's alternate characterization of an NCCR continues to be valid, as long as S admits a canonical module [IW14, 2.17]. For a discussion of the justifications and applications of the larger Cohen-Macaulay generality, consult [IW14, DFI15]. See also [Leu12] for a survey on NCCRs.

The main result of this section is the following characterization of when $\text{End}_R(\mathbb{A})$ is crepant:

Theorem 7.2. *For a toric algebra R and a complete sum of conic modules \mathbb{A} (such as $R^{1/q}$ for large enough q), $\text{End}_R(\mathbb{A})$ is an NCCR of R if and only if R is a simplicial toric algebra.*

This theorem will be proven in two parts, as Propositions 7.5 and 7.9.

7.1. Simplicial toric algebras. Much of the story of this paper simplifies significantly in the simplicial case.

Proposition 7.3. *If R is simplicial, then every chamber of constancy $\Delta \subset M_{\mathbb{R}}$ contains a unique 0-cell.*

Proof. Since $|\Sigma_1| = \dim(M_{\mathbb{R}})$, a 0-cell in the CW decomposition of $M_{\mathbb{R}}$ is a point v such that $\langle v, n_i \rangle \in \mathbb{Z}$ for all i . For this 0-cell to be in Δ , we must have $\langle v, n_i \rangle = d_i$ for all i , where $d_i := \lceil \langle w, n_i \rangle \rceil$ for any $w \in \Delta$. Since Σ_1 is a basis, there is a unique solution v to this system. \square

Lemma 7.4. *Let R be a simplicial toric algebra. For all Δ, Δ' ,*

$$\text{Hom}_R(A_{\Delta}, A_{\Delta'}) = A_{w-v}$$

where $v \in \Delta$ and $w \in \Delta'$ are the unique 0-cells in the respective chambers of constancy.

Proof. If $u \in C + (w - v)$, then $u + (C + v) \subset C + w$. Therefore, for any $x^u \in A_{w-v}$, $x^u A_v \subset A_w$ and so $x^u \in \text{Hom}_R(A_{\Delta}, A_{\Delta'})$.

Consider $u \in M$ such that $x^u \in \text{Hom}_R(A_{\Delta}, A_{\Delta'})$. For each $n_i \in \Sigma_1$, there exist $m \in M \cap (C + v)$ and $m' \in M \cap (C + w)$ such that $\langle m, n_i \rangle = \langle v, n_i \rangle$ and $\langle m', n_i \rangle = \langle w, n_i \rangle$. Since $x^u \cdot x^m = x^{m+u} \in A_w$,

$$\langle m + u, n_i \rangle \geq \langle w, n_i \rangle \Rightarrow \langle v, n_i \rangle + \langle u, n_i \rangle \geq \langle w, n_i \rangle \Rightarrow \langle u, n_i \rangle \geq \langle w - v, n_i \rangle.$$

Therefore, $x^u \in A_{w-v}$. \square

Proposition 7.5. *Let R be a simplicial toric algebra of Krull dimension d , and let \mathbb{A} be a complete sum of conic modules. Then $\text{End}_R(\mathbb{A})$ is an NCCR of R .*

Proof. Assume R is simplicial. Then by Lemma 7.4, $\text{End}_R(\mathbb{A})$ is a sum of conic modules, hence Cohen-Macaulay. On the other hand, each chamber of constancy contains a zero-cell, so all graded simple right $\text{End}_R(\mathbb{A})$ -modules have projective dimension d by Corollary 6.9. We show that $\text{End}_R(\mathbb{A})$ is homologically homogeneous, that is, any simple S of $\text{End}_R(\mathbb{A})$ has projective dimension d . Therefore,

assume that S has projective dimension $k < d$. This means that we can find an $\text{End}_R(\mathbb{A})$ -projective resolution

$$0 \rightarrow P_k \rightarrow \cdots \rightarrow P_0 \rightarrow S \rightarrow 0 ,$$

where each P_i is a finitely generated projective $\text{End}_R(\mathbb{A})$ -module. Since by Lemma 7.4 $\text{End}_R(\mathbb{A})$ has depth d as an R -module, any R -direct summand must also have depth equal to d . Thus, since any P_i is a finite direct sum of direct summands of $\text{End}_R(\mathbb{A})$, it follows that $\text{depth}_R(P_i) = d$ for all $i = 0, \dots, k$. With the same argument as in the proof of Theorem 6.1 we see that S has finite length as an R -module, which implies that $\text{depth}_R(S) = 0$. But the behavior of depth in exact sequences is well-understood: this implies $\text{depth}_R(P_k) = k < d$ (see e.g., [Eis95, Cor.18.6]), a contradiction. \square

Remark 7.6. When R is simplicial, R is isomorphic to the ring of invariant polynomials $\mathbb{k}[x_1, \dots, x_d]^G$ for the action of a finite abelian group G on \mathbb{k}^d . As an R -module, the ring $\mathbb{A} = \mathbb{k}[x_1, \dots, x_d]$ is a complete sum of conic modules, and the corresponding endomorphism ring $\text{End}_R(\mathbb{A})$ is isomorphic to the skew group ring $\mathbb{k}[x_1, \dots, x_d] * G$. In [CQV12, Section 4], so-called *cellular resolutions* are constructed for these skew group rings. However, it is not clear how to extend their construction to the non-simplicial case. \blacksquare

Example 7.7. Let R be a singular toric algebra of dimension 2. Without loss of generality (see [Ful93, 2.2]), the cone C can be taken to be generated by e_1 and $ke_1 + me_2$, with $0 < k < m$ and $(k, m) = 1$, and where e_i are the two standard basis vectors of $M = \mathbb{Z}^2$. Thus C is simplicial and the inner normals are $n_1 = e_2$ and $n_2 = me_1 - ke_2$.

Let us first determine all isomorphism classes of conic modules: since the half-open square $(-1, 0]^2$ is a fundamental domain for the action of M , we find a chamber of constancy for each isomorphism class in it, by Proposition 4.11. There are m different chambers, which can be indexed by their unique 0-cells $v_i = -\frac{i}{m}e_1$ for $0 \leq i < m$ (Proposition 7.3). In particular, the chamber of constancy $\Delta_i := \Delta_{v_i}$ consists of all $x = (x_1, x_2)$ such that

$$-1 < x_2 \leq 0 \quad \text{and} \quad -i - 1 < mx_1 - kx_2 \leq -i .$$

Denote the corresponding conic module by A_i . A general chamber of constancy can be indexed by its 0-cell $v_{ij} = (-\frac{i}{m}, j)$ for $i, j \in \mathbb{Z}$. The corresponding conic module is then $A_{ij} := A_{v_{ij}}$. Now we are ready to describe the conic complex $\mathcal{K}_{\Delta_i}^\bullet$: it is of the form

$$(7.1) \quad A_{i-k-1,1} \longrightarrow A_{i-1} \oplus A_{i-k,1} \longrightarrow A_i ,$$

with the first index $\bmod m$. It is easy to see that $A_{i-k,1} \simeq A_{i-k}$ and $A_{i-k-1,1} \simeq A_{i-k-1}$.




FIGURE 8. The fundamental domain $(-1, 0]^2$ for $m = 3$, $k = 2$ with chambers Δ_0 , Δ_1 , Δ_2 .

The module

$$\mathbb{A} = \bigoplus_{i=0}^{m-1} A_i$$

is a complete sum of conic modules as in (6.1). The graded simples of $\text{End}_R(\mathbb{A})$ all have projective dimension 2, with projective resolution $\text{Hom}_R(\mathbb{A}, \mathcal{K}_{\Delta_i}^\bullet)$ coming from (7.1). This implies that $\text{End}_R(\mathbb{A})$ is an NCCR of R ; in fact the only NCCR up to Morita equivalence, see [IW13, Prop. 2.6]. \blacksquare

7.2. The non-simplicial case. The preceding argument always fails in the non-simplicial case for the following geometric reason.

Lemma 7.8. *If R is not simplicial, then there exists a chamber of constancy Δ containing no 0-cells.*

Proof. We proceed by induction on $|\Sigma_1|$. For the base case, assume $|\Sigma_1| - 1 = d := \text{rank}(M)$ and enumerate the elements as n_1, n_2, \dots, n_{d+1} . Choose an identification $\det : \Lambda^d N \xrightarrow{\sim} \mathbb{Z}$, and set

$$\det(\widehat{n}_i) := \det(n_1 \wedge n_2 \wedge \dots \wedge \widehat{n}_i \wedge \dots \wedge n_{d+1})$$

The identity $n_1 \wedge n_2 \wedge \dots \wedge n_i \wedge \dots \wedge n_{d+1} = 0$ in the exterior algebra of N implies

$$(7.2) \quad \sum_{i=1}^{d+1} (-1)^i \det(\widehat{n}_i) n_i = 0.$$

Pick an i such that $|\det(\widehat{n}_i)|$ is maximized; in particular, $\det(\widehat{n}_i) \neq 0$. First, we claim there is a chamber of constancy Δ given by

$$\Delta = \{x \in M_{\mathbb{R}} \mid \delta_{ij} - 1 < \langle x, n_j \rangle \leq \delta_{ij} \text{ for all } j = 1, \dots, d+1\}$$

where δ_{ij} is the Kronecker δ .⁷ To see this, consider a point w in the interior of the facet of C with normal n_i . This implies that $\langle w, n_j \rangle \geq 0$, with equality if and only if $i = j$. For sufficiently small $\varepsilon > 0$, we have $-1 < \langle -\varepsilon w, n_j \rangle \leq 0$, with equality if and only if $i = j$. Adding a small perturbation p , we find a point $-\varepsilon w + p$ with

$$\lceil \langle -\varepsilon w + p, n_j \rangle \rceil = \delta_{ij} \text{ for all } j = 1, \dots, d+1$$

and so $-\varepsilon w + p$ is in the desired chamber of constancy Δ .

Now, assume that Δ does have a 0-cell, at a point $v \in M_{\mathbb{R}}$. This means v is in d -many linearly independent hyperplanes, or equivalently, we can find a $j \in \{1, 2, \dots, d+1\}$ such that $\det(\widehat{n}_j) \neq 0$ and

$$\langle v, n_k \rangle = \lceil \langle v, n_k \rangle \rceil \text{ for all } k \neq j.$$

By the defining equations of Δ , $\lceil \langle v, n_k \rangle \rceil = 1$ if $k = i$ and 0 otherwise. Therefore, applying $\langle v, - \rangle$ to (7.2) yields

$$0 = \sum_{k=1}^{d+1} (-1)^k \det(\widehat{n}_k) \langle v, n_k \rangle = (-1)^i \det(\widehat{n}_i) + (-1)^j \det(\widehat{n}_j) \langle v, n_j \rangle.$$

Since $\det(\widehat{n}_i) \neq 0$, this equation implies that both $\det(\widehat{n}_j)$ and $\langle v, n_j \rangle$ are non-zero. Solving for $\langle v, n_j \rangle$, we find

$$\langle v, n_j \rangle = (-1)^{j-i+1} \frac{\det(\widehat{n}_i)}{\det(\widehat{n}_j)},$$

so rounding up, we have $\langle v, n_j \rangle \neq 0$. This contradicts the assumption that v was in the chamber Δ . Therefore, Δ has no 0-cells.

Next, assume the proposition is true whenever $|\Sigma_1| = i$ for some $i > d$. Consider a C with $|\Sigma_1| = i+1$, and enumerate the elements n_1, n_2, \dots, n_{i+1} . Since C is pointed, the set of normals is a spanning set, and so there is an i -tuple of them which is also a spanning set. Without loss of generality, assume n_1, n_2, \dots, n_i is a spanning set, and let C' be the cone they define.

Consider the chambers of constancy associated to the cone C' . By the inductive hypothesis, there is a chamber Δ' which does not contain a 0-cell. The chamber Δ' is a union of chambers of constancy of C , each corresponding to a different value of $\lceil \langle -, n_{i+1} \rangle \rceil$. Let Δ be the chamber of C contained in Δ' on which $\lceil \langle -, n_{i+1} \rangle \rceil$ is maximal.

A 0-cell of Δ would have to be the intersection of a 1-cell in Δ' with a hyperplane of the form $\langle v, n_{i+1} \rangle = a$ for some $a \in \mathbb{Z}$. However, if the 1-cell is transverse to the hyperplane, then there are points in Δ' on which $\langle v, n_{i+1} \rangle > a$, contradicting maximality. If the 1-cell is contained in the hyperplane, then their intersection is not a point. Therefore, Δ does not contain a 0-cell, completing the induction. \square

⁷Equivalently, the negative standard basis vector $-e_i$ is a conic divisor; see Remark 4.5.

Proposition 7.9. *If R is a non-simplicial toric algebra and \mathbb{A} is a complete sum of conic modules, then $\text{End}_R(\mathbb{A})$ is not an NCCR of R .*

Proof. By the preceding lemma, there is a chamber of constancy Δ with no 0-cell. By Corollary 6.9, the simple $\text{End}_R(\mathbb{A})$ -module S_Δ has projective dimension strictly less than $\dim(R)$. Hence, not every simple $\text{End}_R(\mathbb{A})$ -module has the same projective dimension, and so it is not an NCCR. \square

Example 7.10. Look again at the cone over the square in \mathbb{R}^3 from Examples 5.3 and 6.10. Here $\text{End}_R(\mathbb{A})$, for $\mathbb{A} = A_0 \oplus A_1 \oplus A_2$, is of global dimension 3 but has two simples of projective dimension 2, coming from the two tetrahedral chambers of constancy. This reproves the fact that $\text{End}_R(\mathbb{A})$ is not an NCCR.⁸

However, this can be fixed by omitting one of the two tetrahedral conic modules. For example, if we omit A_2 , we can build analogous conic complexes

$$\begin{aligned} A_0 &\longrightarrow A_1 \oplus A_0^{\oplus 4} \longrightarrow A_0^{\oplus 4} \oplus A_1^{\oplus 2} \longrightarrow A_0, \\ A_1 &\longrightarrow A_0^{\oplus 2} \longrightarrow A_0^{\oplus 2} \longrightarrow A_1, \end{aligned}$$

which the functor $\text{Hom}_R(A_0 \oplus A_1, -)$ takes to minimal length projective resolutions of the only two graded simple $\text{End}_R(A_0 \oplus A_1)$ -modules. By an analogous argument to the simplicial case, this implies $\text{End}_R(A_0 \oplus A_1)$ is an NCCR of R . \blacksquare

8. ACKNOWLEDGEMENTS

The authors are grateful to Colin Ingalls for his interest and stimulating discussions while he was visiting Ann Arbor. The first author wishes to thank Michael Wemyss for helpful discussions, as well, and the third author Kevin Tucker and Michel Van den Bergh. We also want to thank the referee for valuable comments and suggestions.

REFERENCES

- [Aus55] M. Auslander. On the dimension of modules and algebras. III. Global dimension. *Nagoya Math. J.*, 9:67–77, 1955.
- [Bae04] Cornel Baetica. Cohen-Macaulay classes which are not conic. *Comm. Algebra*, 32(3):1183–1188, 2004.
- [Bas68] Hyman Bass. *Algebraic K-theory*. W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- [Ber58] I. Beršteĭn. On the dimension of modules and algebras. IX. Direct limits. *Nagoya Math. J.*, 13:83–84, 1958.
- [BG03] Winfried Bruns and Joseph Gubeladze. Divisorial linear algebra of normal semigroup rings. *Algebr. Represent. Theory*, 6(2):139–168, 2003.
- [Bli13] Manuel Blickle. Test ideals via algebras of p^{-e} -linear maps. *J. Algebraic Geom.*, 22(1):49–83, 2013.

⁸See Chapter 5 of [Qua05] or Theorem P.2 of [Leu12].

- [BO02] A. Bondal and D. Orlov. Derived categories of coherent sheaves. In *Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002)*, pages 47–56. Higher Ed. Press, Beijing, 2002.
- [Boc12] R. Bocklandt. Generating toric noncommutative crepant resolutions. *J. Algebra*, 364:119–147, 2012.
- [Bri93] Michel Brion. Sur les modules de covariants. *Ann. Sci. École Norm. Sup. (4)*, 26(1):1–21, 1993.
- [Bro12] Nathan Broomhead. Dimer models and Calabi-Yau algebras. *Mem. Amer. Math. Soc.*, 215(1011):viii+86, 2012.
- [Bru05] Winfried Bruns. Conic divisor classes over a normal monoid algebra. In *Commutative algebra and algebraic geometry*, volume 390 of *Contemp. Math.*, pages 63–71. Amer. Math. Soc., Providence, RI, 2005.
- [Cha74] Stephen U. Chase. On the homological dimension of algebras of differential operators. *Comm. Algebra*, 1:351–363, 1974.
- [CQV12] Alastair Craw and Alexander Quintero Vélez. Cellular resolutions of noncommutative toric algebras from superpotentials. *Adv. Math.*, 229(3):1516–1554, 2012.
- [CRST17] J. Carvajal-Rojas, K. Schwede, and K. Tucker. Bertini Theorems for F -signature. *preprint*, [arXiv:1710.01277](https://arxiv.org/abs/1710.01277), 2017.
- [Däs92] Sorin Dăscălescu. Graded semiperfect rings. *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)*, 36(84)(3-4):247–254, 1992.
- [DFI15] H. Dao, E. Faber, and C. Ingalls. Noncommutative (crepant) desingularizations and the global spectrum of commutative rings. *Algebr. Represent. Theory*, 18(3):633–664, 2015.
- [DITV15] H. Dao, O. Iyama, R. Takahashi, and C. Vial. Non-commutative resolutions and Grothendieck groups. *J. Noncommut. Geom.*, 9(1):21–34, 2015.
- [Dix63] J. Dixmier. Représentations irréductibles des algèbres de Lie nilpotentes. *An. Acad. Brasil. Ci.*, 35:491–519, 1963.
- [Don02] X. Dong. Canonical modules of semigroup rings and a conjecture of Reiner. *Discrete Comput. Geom.*, 27(1):85–97, 2002. Geometric combinatorics (San Francisco, CA/Davis, CA, 2000).
- [Eis95] David Eisenbud. *Commutative algebra. With a view toward algebraic geometry*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [Ful93] William Fulton. *Introduction to Toric Varieties*, volume 131 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, 1993.
- [HGK04] M. Hazewinkel, N. Gubareni, and V. V. Kirichenko. *Algebras, rings and modules. Vol. 1*, volume 575 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 2004.
- [HH90] Melvin Hochster and Craig Huneke. Tight closure, invariant theory, and the Briançon-Skoda theorem. *J. Amer. Math. Soc.*, 3(1):31–116, 1990.
- [HL02] Craig Huneke and Graham J. Leuschke. Two theorems about maximal Cohen-Macaulay modules. *Math. Ann.*, 324(2):391–404, 2002.
- [HN17] A. Higashitani and Y. Nakajima. Conic divisorial ideals of Hibi rings and their applications to non-commutative crepant resolutions. *preprint*, [arXiv:1702.07058](https://arxiv.org/abs/1702.07058), 2017.
- [HW02] Nobuo Hara and Kei-Ichi Watanabe. F -regular and F -pure rings vs. log terminal and log canonical singularities. *J. Algebraic Geom.*, 11(2):363–392, 2002.

- [HX15] Christopher D. Hacon and Chenyang Xu. On the three dimensional minimal model program in positive characteristic. *J. Amer. Math. Soc.*, 28(3):711–744, 2015.
- [HY03] Nobuo Hara and Ken-Ichi Yoshida. A generalization of tight closure and multiplier ideals. *Trans. Amer. Math. Soc.*, 355(8):3143–3174, 2003.
- [IW13] Osamu Iyama and Michael Wemyss. On the noncommutative Bondal-Orlov conjecture. *J. Reine Angew. Math.*, 683:119–128, 2013.
- [IW14] O. Iyama and M. Wemyss. Maximal modifications and Auslander-Reiten duality for non-isolated singularities. *Invent. Math.*, 197(3):521–586, 2014.
- [Jef17] Jack Jeffries. Derived functors of differential operators. *preprint*, [arXiv:1711.03960](#), 2017.
- [JNB17] Jack Jeffries and Luis Núñez Betancourt. Quantifying singularities with differential operators. 2017. in preparation.
- [Kun69] Ernst Kunz. Characterizations of regular local rings for characteristic p . *Amer. J. Math.*, 91:772–784, 1969.
- [Leu12] Graham J. Leuschke. Non-commutative crepant resolutions: scenes from categorical geometry. In *Progress in commutative algebra 1*, pages 293–361. de Gruyter, Berlin, 2012.
- [Mun84] James R. Munkres. *Elements of algebraic topology*. Addison-Wesley Publishing Company, Menlo Park, CA, 1984.
- [MVB98] Ian M. Musson and Michel Van den Bergh. Invariants under tori of rings of differential operators and related topics. *Mem. Amer. Math. Soc.*, 136(650):viii+85, 1998.
- [NVO79] Constantin Năstăsescu and F. Van Oystaeyen. *Graded and filtered rings and modules*, volume 758 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979.
- [Qua05] C. Quarles. Krull–Schmidt rings and noncommutative resolutions of singularities. Master’s thesis, University of Washington, 2005.
- [Roo72] Jan-Erik Roos. Détermination de la dimension homologique globale des algèbres de Weyl. *C. R. Acad. Sci. Paris Sér. A-B*, 274:A23–A26, 1972.
- [RŠVdB17] T. Raedschelders, Š. Špenko, and M. Van den Bergh. The Frobenius morphism in invariant theory. *preprint*, [arXiv:1705.01832](#), 2017.
- [Sch09] Karl Schwede. F -injective singularities are Du Bois. *Amer. J. Math.*, 131(2):445–473, 2009.
- [Smi87] S. P. Smith. The global homological dimension of the ring of differential operators on a nonsingular variety over a field of positive characteristic. *J. Algebra*, 107(1):98–105, 1987.
- [Smi95] Karen E. Smith. The D -module structure of F -split rings. *Math. Res. Lett.*, 2(4):377–386, 1995.
- [Smi97] Karen E. Smith. Vanishing, singularities and effective bounds via prime characteristic local algebra. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 289–325. Amer. Math. Soc., Providence, RI, 1997.
- [Smi00] Karen E. Smith. Globally F -regular varieties: applications to vanishing theorems for quotients of Fano varieties. *Michigan Math. J.*, 48:553–572, 2000. Dedicated to William Fulton on the occasion of his 60th birthday.
- [ST12] Karl Schwede and Kevin Tucker. A survey of test ideals. In *Progress in commutative algebra 2*, pages 39–99. Walter de Gruyter, Berlin, 2012.
- [Sta82] Richard P. Stanley. Linear Diophantine equations and local cohomology. *Invent. Math.*, 68(2):175–193, 1982.

- [SVdB97] Karen E. Smith and Michel Van den Bergh. Simplicity of rings of differential operators in prime characteristic. *Proc. London Math. Soc. (3)*, 75(1):32–62, 1997.
- [SVdB08] J. T. Stafford and M. Van den Bergh. Noncommutative resolutions and rational singularities. *Michigan Math. J.*, 57:659–674, 2008. Special volume in honor of Melvin Hochster.
- [ŠVdB17a] Š. Špenko and M. Van den Bergh. Non-commutative resolutions of some toric varieties I. *preprint*, [arXiv:1701.05255](#), 2017.
- [ŠVdB17b] Špela Špenko and Michel Van den Bergh. Non-commutative resolutions of quotient singularities for reductive groups. *Invent. Math.*, 210(1):3–67, 2017.
- [Tak04] Shunsuke Takagi. An interpretation of multiplier ideals via tight closure. *J. Algebraic Geom.*, 13(2):393–415, 2004.
- [Tuc12] Kevin Tucker. F -signature exists. *Invent. Math.*, 190(3):743–765, 2012.
- [VdB93] Michel Van den Bergh. Cohen-Macaulayness of semi-invariants for tori. *Trans. Amer. Math. Soc.*, 336(2):557–580, 1993.
- [VdB04] Michel Van den Bergh. Non-commutative crepant resolutions. In *The legacy of Niels Henrik Abel*, pages 749–770. Springer, Berlin, 2004.
- [Wem16] Michael Wemyss. Noncommutative resolutions. In *Noncommutative algebraic geometry*, volume 64 of *Math. Sci. Res. Inst. Publ.*, pages 239–306. Cambridge Univ. Press, New York, 2016.
- [Yas10] T. Yasuda. Noncommutative resolution of toric singularities: An application of Frobenius morphism of noncommutative blowup. *preprint*, [arXiv:1002.0181v1](#), 2010.
- [Yek92] Amnon Yekutieli. An explicit construction of the Grothendieck residue complex. *Astérisque*, (208):127, 1992. With an appendix by Pramathanath Sastry.

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS, LS2 9JT, UK

E-mail address: e.m.faber@leeds.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OK 73019, USA

E-mail address: gmuller@math.ou.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109, USA

E-mail address: kesmith@umich.edu