

ON THE CHENG-YAU GRADIENT ESTIMATE FOR CARNOT GROUPS AND SUB-RIEMANNIAN MANIFOLDS

FABRICE BAUDOUIN^{*}, MARIA GORDINA^{†‡}, AND PHANUEL MARIANO[†]

ABSTRACT. In this note we show how results in [4, 6, 11] yield the Cheng-Yau estimate on two classes of sub-Riemannian manifolds: Carnot groups and sub-Riemannian manifolds satisfying a generalized curvature-dimension inequality.

CONTENTS

1. Introduction	1
2. Carnot groups	2
2.1. Preliminaries	2
2.2. The Cheng-Yau estimate	3
3. Sub-Riemannian manifolds	4
3.1. Preliminaries	4
3.2. Examples	6
3.3. The Cheng-Yau estimate	6
References	8

Primary 58J35; Secondary 53C17, 35H10

1. INTRODUCTION

Let M be a d -dimensional Riemannian complete non-compact manifold with the Ricci curvature bounded below by $-(d-1)K$. Let u be a positive harmonic function in a Riemannian ball $B(x_0, 2r)$, then we say that u satisfies the Cheng-Yau estimate if

$$(1.1) \quad \sup_{B(x_0, r)} |\nabla \log u| \leq C_d \left(\frac{1}{r} + \sqrt{K} \right),$$

where C_d is a global constant depending only on the dimension d . In particular, when $K = 0$ this estimate shows that positive harmonic functions are constant. This estimate was formulated in a more general form in [10, 28], and stated as in (1.1) in [24, Theorem 3.1]. Sharp versions of the Cheng-Yau inequality were given

Key words and phrases. Cheng-Yau estimate, Carnot groups, sub-Riemannian manifolds, curvature-dimension inequality.

^{*} This research was supported in part by NSF Grant DMS-1660031.

[‡] Research was supported in part by the Simons Fellowship.

[†] This research was supported in part by NSF Grants DMS-1405169, DMS-1712427.

in [21, 22]. The stability of (1.1) under certain perturbations of the metric was considered in [29]

The standard curvature arguments are not easily available in the case when M is replaced by a sub-Riemannian manifold. Nevertheless, there has been significant progress in geometric analysis on sub-Riemannian manifolds in [1, 4, 5, 7]. Even in the absence of a Riemannian structure, [7] developed new techniques to prove a number of results which in the Riemannian setting go back to the work of Yau and Li-Yau. The main tool in [7] relied on a generalized curvature-dimension inequality on a class of sub-Riemannian manifolds with transverse symmetries.

Carnot groups also form a large and interesting class of sub-Riemannian manifolds. These are Lie groups whose Lie algebra admits a stratified structure. This stratified structure allows for Hörmander's condition [18, Theorem 1.1] to be satisfied. Hörmander's theorem guarantees that the sub-Laplacian associated with the structure of a Carnot group is hypoelliptic. In particular, this gives us the existence of a smooth heat kernel for this Laplacian. Most Carnot groups do not satisfy a generalized curvature-dimension inequality, so one needs to employ different techniques than in [7].

The Cheng-Yau estimate was proved in [2, Corollary 4.6] for the simplest non-commutative Carnot group, the Heisenberg group, using probabilistic (coupling) techniques. The purpose of this note is to show that this estimate can be proven on two classes of sub-Riemannian manifolds, namely, sub-Riemannian manifolds satisfying a generalized curvature-dimension inequality and Carnot groups, by relying on results from [4, 6, 11]. In particular, this recovers the known fact that global non-negative harmonic functions in these two settings have to be constant (see [8, Theorem 5.8.1] and [5, Theorem 5.1] in more generality).

2. CARNOT GROUPS

2.1. Preliminaries. We recall that a Carnot group of step N is a simply connected Lie group $\mathbb{G}$ whose Lie algebra can be written as

$$\mathfrak{g} = \mathcal{V}_1 \oplus \cdots \oplus \mathcal{V}_N,$$

where

$$[\mathcal{V}_i, \mathcal{V}_j] = \mathcal{V}_{i+j}$$

and $\mathcal{V}_k = 0$ for $k > N$. In particular, Carnot groups are nilpotent.

Let $V_1, \dots, V_d$ be a linear basis for the vector space $\mathcal{V}_1$. The V_i s can be viewed as left-invariant vector fields on $\mathbb{G}$. The left-invariant sub-Laplacian on $\mathbb{G}$ is the operator

$$(2.2) \quad L = \sum_{i=1}^d V_i^2.$$

Let μ be the bi-invariant Haar measure on $\mathbb{G}$. Since Carnot groups are complete the operator L in (2.2) is essentially self-adjoint on $L^2(\mathbb{G}, \mu)$, with domain being the space of smooth and compactly supported functions $f : \mathbb{G} \rightarrow \mathbb{R}$ denoted by $C_c^\infty(\mathbb{G})$. We abuse notation and denote by L the Friedrichs extension of this operator to a unique non-positive self-adjoint operator on $L^2(\mathbb{G}, \mu)$. Then the heat semigroup $(P_t)_{t \geq 0}$ on $\mathbb{G}$ can be defined through the spectral theorem. As L is hypoelliptic,

P_t admits a positive smooth fundamental solution to the heat equation called the heat kernel $p_t(g, g')$.

Denote by $\nabla = (V_1, \dots, V_d)$ the gradient determined by this basis, and denote by $\|\cdot\|$ the usual Euclidean norm. The *carré du champ* operator of L is defined by

$$\Gamma(f, f) := \frac{1}{2} (Lf^2 - 2fLf) = \|\nabla f\|^2 = \sum_{i=1}^d (V_i f)^2,$$

and it is often thought of as the square of the *length of the gradient* ∇ . We let d be the Carnot-Carathéodory distance on $\mathbb{G}$ making $(\mathbb{G}, d)$ a metric space. We refer the reader to [8] for more details and results on Carnot groups.

2.2. The Cheng-Yau estimate. We say a function $u : \mathbb{G} \rightarrow \mathbb{R}$ is *harmonic* in a domain $D \subset \mathbb{G}$ if $Lu = 0$ on $D \subset \mathbb{G}$.

Theorem 2.1. *If u is any positive harmonic function for L in a ball $B(x, 2r) \subset \mathbb{G}$, then there exists a constant $C > 0$ not dependent on u, r and x such that*

$$(2.3) \quad \sup_{B(x, r)} \|\nabla \log u\| \leq \frac{C}{r}.$$

Moreover, if u is a positive harmonic function on $\mathbb{G}$, then u must be equal to a constant.

Proof. In the proof, C will denote a generic positive constant that does not depend on u, r and x whose value might change from line to line. First recall the reverse Poincaré inequality for the heat semigroup obtained for Carnot groups in [4, Proposition 2.5]

$$(2.4) \quad \|\nabla P_t f(g)\|^2 \leq \frac{C}{t} \left(P_t f^2(g) - (P_t f)^2(g) \right)$$

for functions $f \in C_c^\infty(\mathbb{G})$. Note that for functions $f \in L^\infty(\mathbb{G})$ we have

$$(2.5) \quad P_t f^2(g) \leq \|f\|_{L^\infty(G)}^2,$$

therefore combining (2.4) and (2.5) we obtain

$$(2.6) \quad \|\nabla P_t f(g)\|^2 \leq \frac{C}{t} \|f\|_{L^\infty(G)}^2.$$

Taking a square root in (2.6) implies that

$$(2.7) \quad \|\nabla P_t\|_{\infty \rightarrow \infty} \leq \frac{C}{\sqrt{t}}.$$

Applying [11, Theorem 1.2], in particular that (iii) implies (i), shows that (2.7) implies that there exists a $C > 0$ such that, for every ball $B(x, r)$ and every function u that is harmonic in $B(x, 2r)$ we have

$$(2.8) \quad \|\nabla u\|_{L^\infty(B(x, r))} \leq \frac{C}{r\mu(B(x, 2r))} \int_{B(x, 2r)} |u| d\mu.$$

Then we can apply [11, Lemma 2.3] to show that (2.8) implies the Cheng-Yau estimate (2.3).

We rely on results in [11] that require several assumptions that are satisfied for Carnot groups as follows. Their first assumption is that the underlying space is

a non-compact doubling Dirichlet metric measure space. The space $\mathbb{G}$ is doubling since by [16, Proposition 11.15] (or more classically by [14, 20]) there exists a $C > 0$ independent of $x \in \mathbb{G}, r > 0$ such that $|B(x, r)| = Cr^Q$ where $Q = \sum_{j=1}^N j \dim(\mathcal{V}_j)$ is the homogeneous dimension of the $\mathbb{G}$. Here $|E|$ denotes the Lebesgue measure of the set E and recall that the Haar measure μ is Lebesgue measure up to a constant. We also have that $\mathbb{G}$ is Dirichlet space as described in [25, Section 3, pp.233-234]. This is actually true in general for Hörmander's type operators with bounded measurable coefficients on Lie groups having polynomial volume growth in the sense of [23]. Another ingredient in [11] is upper Gaussian bounds on the heat kernel which follow from [27, Theorem VIII.2.9]. The space also supports a local scale-invariant L^2 -Poincaré inequality by [16, Proposition 11.17].

Finally, if u is a positive harmonic function on all of $\mathbb{G}$, taking $r \rightarrow \infty$ in (2.3) gives us that u must be constant. $\square$

3. SUB-RIEMANNIAN MANIFOLDS

3.1. Preliminaries. In this section we study the setting similar to [5]. We state relevant details here for completeness. Let $(\mathbb{M}, \mu)$ be a measure space, where $\mathbb{M}$ is an n -dimensional C^∞ connected manifold endowed with a smooth measure μ . Recall (e.g. [26, p.85]) that the measure μ on a smooth manifold $\mathbb{M}$ is called a *smooth measure* if μ is a Radon measure which has a smooth Radon-Nikodym derivative with respect to the Lebesgue measure when viewed in coordinates, that is, for any smooth coordinate chart $\varphi : U \rightarrow V$, $U \subset \mathbb{M}$, $V \subset \mathbb{R}^n$, the pushforward measure $\varphi_*(\mu)$ has a smooth Radon-Nikodym derivative with respect to the Lebesgue measure on V . Let L be a second order diffusion operator on $\mathbb{M}$ which is locally subelliptic (in the sense of [13, 19]). We refer to [3, Section 1] for a detailed account on properties of locally subelliptic operators and associated distances that we are going to use in the sequel. In addition, we assume that

$$\begin{aligned} L1 &= 0, \\ \int_{\mathbb{M}} f Lg d\mu &= \int_{\mathbb{M}} g Lf d\mu, \\ \int_{\mathbb{M}} f Lf &\leq 0 \end{aligned}$$

for every $f, g \in C_c^\infty(\mathbb{M})$, where as before $C_c^\infty(\mathbb{M})$ denotes the space of smooth compactly supported functions on $\mathbb{M}$.

The space $\mathbb{M}$ is endowed with a *carré du champ* operator defined by

$$\Gamma(f, g) := \frac{1}{2} (L(fg) - fLg - gLf), \quad f, g \in C^\infty(\mathbb{M}).$$

We denote $\Gamma(f) = \Gamma(f, f)$. It is not too hard to see that $\Gamma(f) \geq 0$ for all $f \in C^\infty(\mathbb{M})$. We will also assume the existence of a symmetric, first-order differential bilinear form $\Gamma^Z : C^\infty(\mathbb{M}) \times C^\infty(\mathbb{M}) \rightarrow C^\infty(\mathbb{M})$ that satisfies

$$\begin{aligned} \Gamma^Z(fg, h) &= f\Gamma^Z(g, h) + g\Gamma^Z(f, h), \\ \Gamma^Z(f) &= \Gamma^Z(f, f) \geq 0, \end{aligned}$$

for all $f, g, h \in C^\infty(\mathbb{M})$. Given the first order bi-linear forms Γ and Γ^Z on $\mathbb{M}$, we can introduce the following second-order differential forms

$$\Gamma_2(f, g) = \frac{1}{2} (L\Gamma(f, g) - \Gamma(f, Lg) - \Gamma(g, Lf))$$

and

$$\Gamma_2^Z(f, g) = \frac{1}{2} (L\Gamma^Z(f, g) - \Gamma^Z(f, Lg) - \Gamma^Z(g, Lf)).$$

Similar to Γ , we will use the notation $\Gamma_2(f) := \Gamma_2(f, f)$, $\Gamma_2^Z(f) := \Gamma_2^Z(f, f)$.

We suppose the following assumptions to hold throughout this section.

- (I) There exists an increasing sequence $h_k \in C_c^\infty(\mathbb{M})$ such that $h_k \uparrow 1$ on $\mathbb{M}$, and

$$\|\Gamma(h_k)\|_\infty + \|\Gamma^Z(h_k)\|_\infty \rightarrow 0, \text{ as } k \rightarrow \infty.$$

- (II) For any $f \in C^\infty(\mathbb{M})$ one has

$$\Gamma(f, \Gamma^Z(f)) = \Gamma^Z(f, \Gamma(f)).$$

- (III) The *generalized curvature-dimension inequality* $CD(\rho_1, \rho_2, \kappa, d)$ is satisfied with $\rho_1 \geq 0$. That is, there exist constants $\rho_1 \geq 0, \rho_2 > 0, \kappa \geq 0$, and $d \geq 2$ such that the following inequality holds

$$\Gamma(f) + \nu \Gamma_2^Z(f) \geq \frac{1}{d} (Lf)^2 + \left(\rho_1 - \frac{\kappa}{\nu}\right) \Gamma(f) + \rho_2 \Gamma^Z(f),$$

for all $f \in C^\infty(\mathbb{M})$ and every $\nu > 0$.

- (IV) The heat semigroup generated by L , which will be denoted P_t , is stochastically complete, that is, for $t \geq 0$, $P_t 1 = 1$ and for every $f \in C_c^\infty(\mathbb{M})$ and $T \geq 0$, one has

$$\sup_{t \in [0, T]} \|\Gamma(P_t f)\|_\infty + \|\Gamma^Z(P_t f)\|_\infty < +\infty.$$

- (V) Given any two points $x, y \in \mathbb{M}$, there exists a subunit curve (in the sense of [12]), joining them.

- (VI) The metric space $(\mathbb{M}, d)$ is complete with respect to the intrinsic distance defined by

$$(3.9) \quad d(x, y) := \sup \{|f(x) - f(y)| : f \in C^\infty(\mathbb{M}), \|\Gamma(f)\|_\infty \leq 1\},$$

for all $x, y \in \mathbb{M}$ and where we define $\|g\|_\infty = \text{ess sup}_{\mathbb{M}} |g|$.

Note that by [9, Lemma 5.29] and [5, Equation (2.4)] we know that Assumption (V) implies that d is indeed a metric on $\mathbb{M}$.

As a consequence of Assumption (VI), the operator L is essentially self-adjoint on $C_c^\infty(\mathbb{M})$ (e.g. [3, Proposition 1.20, Proposition 1.21]). Thus, the pre-Dirichlet form $\mathcal{E}(f, g)$ defined on $C_c^\infty(\mathbb{M})$ by

$$\mathcal{E}(f, g) = \int_{\mathbb{M}} \Gamma(f, g) d\mu,$$

has a unique closure as a Dirichlet form, and the generator of this Dirichlet form is the Friedrichs extension of L . We define the Sobolev space $W^{1,2}(\mathbb{M})$ to be the domain $\mathcal{D}$ of $\mathcal{E}$ with the norm on $W^{1,2}(\mathbb{M})$ given by

$$\|f\|_{W^{1,2}(\mathbb{M})} = \sqrt{\|f\|_2^2 + \mathcal{E}(f, f)}.$$

It is a consequence of Assumption (III) that the metric measure space $(\mathbb{M}, d, \mu)$ satisfies the volume doubling property and supports a scale invariant L^2 -Poincaré

inequality on metric balls (see [5]). In particular, by [17, Chapter 8] locally Lipschitz continuous functions form a dense subclass in $W^{1,2}(\mathbb{M})$.

For an open set $U \subset \mathbb{M}$, one can define the local Sobolev space $W_{\text{loc}}^{1,2}(U)$ to be the space of all functions f such that for any compact set $K \subset U$ there exists $F \in \mathcal{D}$ satisfying $f = F$ a.e. on K . For each $p \geq 2$, we define $W^{1,p}(U)$ to be space of functions $f \in W_{\text{loc}}^{1,2}(U)$ satisfying $f, \sqrt{\Gamma(f)} \in L^p(U)$.

3.2. Examples. We remark that so far the approach has been purely analytical as we have not mentioned any geometric structure of these sub-Riemannian manifolds. In fact $\mathbb{M}$ and L are rather general, even though we have sub-Riemannian manifolds in mind for $\mathbb{M}$.

3.2.1. Sum of squares operators. We start by recalling a natural setting where assumption (V) is satisfied. Let us consider L that are *sums of squares* operators in the form of

$$(3.10) \quad L = \sum_{i=1}^m X_i^2 + X_0,$$

where the X_i are C^∞ vector fields. We refer the reader to [15] for more details on operators of the form given in (3.10) in the context of sub-Riemannian manifolds. Consider the following assumption.

Assumption 3.1. (*Hörmander's condition*) We will say that L satisfies Hörmander's (bracket generating) condition if the vector fields $\{X_1, \dots, X_m\}$ with their Lie brackets span the tangent space $T_x \mathbb{M}$ at every point $x \in \mathbb{M}$.

Hörmander's condition guarantees analytic and topological properties such as hypoellipticity of L and topological properties of $\mathbb{M}$. The Chow-Rashevski theorem says that Hörmander's condition is sufficient to ensure that any two points in $\mathbb{M}$ can be connected by a finite length sub-unit curve. Thus, operators L of the form (3.10) that satisfy Hörmander's condition automatically satisfy assumption (V).

3.2.2. Other examples. We note that the assumptions (I)-(VI) are satisfied for a large class of sub-Riemannian manifolds. Such a class includes all Sasakian manifolds whose horizontal Webster-Tanaka-Ricci curvature is bounded below, a wide subclasses of principal bundles over Riemannian manifolds whose Ricci curvature is bounded below, and Carnot groups of step 2. We remark that in general we do not know if $CD(\rho_1, \rho_2, \kappa, d)$ is satisfied for Carnot groups of an arbitrary step. This shows the need to treat Carnot groups separately in Theorem 2.1. We refer the reader to [7] for a comprehensive treatment on sub-Riemannian manifolds satisfying assumptions (I)-(VI).

3.3. The Cheng-Yau estimate. Recall that we say a function $u : \mathbb{M} \rightarrow \mathbb{R}$ is *harmonic* in a domain $D \subset \mathbb{M}$ if $Lu = 0$ on $D \subset \mathbb{M}$. We can now state the main result of this section.

Theorem 3.2. *Suppose assumptions (I)-(VI) hold for $\mathbb{M}$ and L . If u is any positive harmonic function for L in a ball $B(x, 2r) \subset \mathbb{M}$, then there exists a constant $C > 0$ not dependent on u, r and x such that*

$$(3.11) \quad \sup_{B(x,r)} \sqrt{\Gamma(\log u)} \leq \frac{C}{r}.$$

Moreover, if u is any positive harmonic function on $\mathbb{M}$, then u must be equal to a constant.

Proof. In the proof, C, C_2 will denote generic positive constants that do not depend on u, r and x_0 , whose values might change from line to line. First we check that the assumptions of the results in [11, Theorem 1.2, Lemma 2.3] are satisfied. The results in [11] require the assumption that the underlying space is a non-compact doubling Dirichlet metric measure space. Doubling is shown in (1.11) of [5, Theorem 1.5]. Further, we need $\mathbb{M}$ to satisfy upper Gaussian bounds on the heat kernel, which is given in [5, Theorem 4.1]. Finally we need $\mathbb{M}$ to support a local L^2 -Poincaré inequality of the form

$$(3.12) \quad \frac{1}{\mu(B(x, r))} \int_{B(x, r)} |f - f_B| d\mu \leq C_2 r \left(\frac{1}{\mu(B(x, r))} \int_{B(x, r)} \Gamma(f) d\mu \right)^{\frac{1}{2}},$$

for some $C_2 > 0$, for every ball $B(x, r)$ and each $f \in W^{1,2}(B(x, r))$. Here we used the notation $f_B = \mu(B)^{-1} \int_B f d\mu$. To see this, by (1.12) in [5, Theorem 1.5] we have that there exists constant $C_2 > 0$, depending only on $\rho_1, \rho_2, \kappa, d$, for which one has for every $x \in \mathbb{M}$ and every $r > 0$,

$$(3.13) \quad \int_{B(x, r)} |f - f_B|^2 \leq C_2 r^2 \int_{B(x, r)} \Gamma(f) d\mu,$$

for every $f \in C^1(\overline{B}(x, r))$. Using Cauchy-Schwarz inequality, followed by (3.13) we have that

$$\begin{aligned} \frac{1}{\mu(B(x, r))} \int_{B(x, r)} |f - f_B| d\mu &\leq \frac{1}{\mu(B(x, r))} \left(\int_{B(x, r)} |f - f_B|^2 d\mu \right)^{\frac{1}{2}} \sqrt{\mu(B(x, r))} \\ &\leq \frac{1}{\sqrt{\mu(B(x, r))}} \left(C_2 r^2 \int_{B(x, r)} \Gamma(f) d\mu \right)^{\frac{1}{2}} \\ &= \sqrt{C_2} r \left(\frac{1}{\mu(B(x, r))} \int_{B(x, r)} \Gamma(f) d\mu \right)^{\frac{1}{2}}. \end{aligned}$$

This shows (3.12) holds for all $f \in C^1(\overline{B}(x, r))$. By a density argument we can show that (3.12) holds for all $f \in W^{1,2}(B(x, r))$, as needed.

To finish off the proof, we note that Corollary 3.5 in [6] shows that

$$(3.14) \quad \left\| \sqrt{\Gamma(P_t f)} \right\|_{\infty \rightarrow \infty} \leq \frac{C}{\sqrt{t}},$$

where $C = n \sqrt{\frac{(2\kappa + \rho_2)}{2\rho_2}}$. Using the estimate (3.14), the rest of the proof becomes similar to the proof of Theorem 2.1. $\square$

Acknowledgement. *The authors would like to thank the anonymous referee for the careful review of the paper. P. Mariano is also grateful for helpful and insightful conversations with O. Munteanu.*

REFERENCES

1. Dominique Bakry, Fabrice Baudoin, Michel Bonnefont, and Djalil Chafaï, *On gradient bounds for the heat kernel on the Heisenberg group*, J. Funct. Anal. **255** (2008), no. 8, 1905–1938. MR 2462581 (2010m:35534)
2. Sayan Banerjee, Maria Gordina, and Phanuel Mariano, *Coupling in the Heisenberg group and its applications to gradient estimates*, Ann. Probab. **46** (2008), no. 6, 3275–3312. MR 3857856
3. Fabrice Baudoin, *Geometric inequalities on Riemannian and sub-Riemannian manifolds by heat semigroups techniques*, <https://arxiv.org/abs/1801.05702>, 2018.
4. Fabrice Baudoin and Michel Bonnefont, *Reverse Poincaré inequalities, isoperimetry, and Riesz transforms in Carnot groups*, Nonlinear Anal. **131** (2016), 48–59. MR 3427969
5. Fabrice Baudoin, Michel Bonnefont, and Nicola Garofalo, *A sub-Riemannian curvature-dimension inequality, volume doubling property and the Poincaré inequality*, Math. Ann. **358** (2014), no. 3–4, 833–860. MR 3175142
6. Fabrice Baudoin and Nicola Garofalo, *A note on the boundedness of Riesz transform for some subelliptic operators*, Int. Math. Res. Not. IMRN (2013), no. 2, 398–421. MR 3010694
7. ———, *Curvature-dimension inequalities and Ricci lower bounds for sub-Riemannian manifolds with transverse symmetries*, J. Eur. Math. Soc. (JEMS) **19** (2017), no. 1, 151–219. MR 3584561
8. A. Bonfiglioli, E. Lanconelli, and F. Uguzzoni, *Stratified Lie groups and potential theory for their sub-Laplacians*, Springer Monographs in Mathematics, Springer, Berlin, 2007. MR 2363343
9. E. A. Carlen, S. Kusuoka, and D. W. Stroock, *Upper bounds for symmetric Markov transition functions*, Ann. Inst. H. Poincaré Probab. Statist. **23** (1987), no. 2, suppl., 245–287. MR 898496
10. S. Y. Cheng and S. T. Yau, *Differential equations on Riemannian manifolds and their geometric applications*, Comm. Pure Appl. Math. **28** (1975), no. 3, 333–354. MR 0385749
11. Thierry Coulhon, Renjin Jiang, Pekka Koskela, and Adam Sikora, *Gradient estimates for heat kernels and harmonic functions*, <https://arxiv.org/abs/1703.02152>, 2017.
12. C. Fefferman and D. H. Phong, *Subelliptic eigenvalue problems*, Conference on harmonic analysis in honor of Antoni Zygmund, Vol. I, II (Chicago, Ill., 1981), Wadsworth Math. Ser., Wadsworth, Belmont, CA, 1983, pp. 590–606. MR 730094
13. Charles L. Fefferman and Antonio Sánchez-Calle, *Fundamental solutions for second order subelliptic operators*, Ann. of Math. (2) **124** (1986), no. 2, 247–272. MR 855295
14. G. B. Folland and Elias M. Stein, *Hardy spaces on homogeneous groups*, Mathematical Notes, vol. 28, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1982. MR 657581
15. Maria Gordina and Thomas Laetsch, *Sub-Laplacians on Sub-Riemannian Manifolds*, Potential Anal. **44** (2016), no. 4, 811–837. MR 3490551
16. Piotr Hajlasz and Pekka Koskela, *Sobolev met Poincaré*, Mem. Amer. Math. Soc. **145** (2000), no. 688, x+101. MR 1683160 (2000j:46063)
17. Juha Heinonen, Pekka Koskela, Nageswari Shanmugalingam, and Jeremy T. Tyson, *Sobolev spaces on metric measure spaces*, New Mathematical Monographs, vol. 27, Cambridge University Press, Cambridge, 2015, An approach based on upper gradients. MR 3363168
18. Lars Hörmander, *Hypoelliptic second order differential equations*, Acta Math. **119** (1967), 147–171. MR 0222474 (36 #5526)
19. David Jerison and Antonio Sánchez-Calle, *Subelliptic, second order differential operators*, Complex analysis, III (College Park, Md., 1985–86), Lecture Notes in Math., vol. 1277, Springer, Berlin, 1987, pp. 46–77. MR 922334
20. Ron Karidi, *Geometry of balls in nilpotent Lie groups*, Duke Math. J. **74** (1994), no. 2, 301–317. MR 1272979
21. Peter Li and Jiaping Wang, *Complete manifolds with positive spectrum. II*, J. Differential Geom. **62** (2002), 143–162. MR 1987380
22. Ovidiu Munteanu, *On the gradient estimate of Cheng and Yau*, Proc. Amer. Math. Soc. **140** (2012), no. 4, 1437–1443. MR 2869128
23. L. Saloff-Coste and D. W. Stroock, *Opérateurs uniformément sous-elliptiques sur les groupes de Lie*, J. Funct. Anal. **98** (1991), no. 1, 97–121. MR 1111195

24. R. Schoen and S.-T. Yau, *Lectures on differential geometry*, Conference Proceedings and Lecture Notes in Geometry and Topology, I, International Press, Cambridge, MA, 1994, Lecture notes prepared by Wei Yue Ding, Kung Ching Chang [Gong Qing Zhang], Jia Qing Zhong and Yi Chao Xu, Translated from the Chinese by Ding and S. Y. Cheng, Preface translated from the Chinese by Kaising Tso. MR MR1333601 (97d:53001)
25. Karl-Theodor Sturm, *On the geometry defined by Dirichlet forms*, Seminar on Stochastic Analysis, Random Fields and Applications (Ascona, 1993), Progr. Probab., vol. 36, Birkhäuser, Basel, 1995, pp. 231–242. MR 1360279
26. Terence Tao, *Hilbert’s fifth problem and related topics*, Graduate Studies in Mathematics, vol. 153, American Mathematical Society, Providence, RI, 2014. MR 3237440
27. N. Th. Varopoulos, L. Saloff-Coste, and T. Coulhon, *Analysis and geometry on groups*, Cambridge University Press, Cambridge, 1992. MR 95f:43008
28. Shing Tung Yau, *Harmonic functions on complete Riemannian manifolds*, Comm. Pure Appl. Math. **28** (1975), 201–228. MR 0431040
29. Qi S. Zhang, *Stability of the Cheng-Yau gradient estimate*, Pacific J. Math. **225** (2006), 379–398. MR 2233741

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269, USA
Email address: `fabrice.baudoin@uconn.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269, USA
Email address: `maria.gordina@uconn.edu`

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, USA
Email address: `pmariano@purdue.edu`