

Enantio- and Diastereoselective Synthesis of Homoallylic α -Trifluoromethyl Amines by Catalytic Hydroalkylation of Dienes

Chibueze I. Onyeagusi, Xinxin Shao, and Steven J. Malcolmsen*



Cite This: *Org. Lett.* 2020, 22, 1681–1685



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: We describe a strategy for the enantio- and diastereoselective synthesis of homoallylic α -trifluoromethyl amines by the catalytic hydroalkylation of terminal dienes. Trifluoromethyl-substituted isatin-derived azadienolate nucleophiles undergo γ -selective alkylation with a Pd–DTBM–SEGPHOS catalyst, which additionally promotes regioselective addition to the diene and delivers products in up to 86% yield, 10:1 dr, and 97.5:2.5 er.




Chiral amines bearing an α -trifluoromethyl group hold a significant place among classes of medicinally important N-containing molecules. The CF_3 group modulates a number of pharmacological parameters, alters the amine basicity and polarity significantly, and stands in as a non-hydrolyzable amide surrogate.¹ Despite the beneficial properties this motif might impart upon drug-like molecules, methods for the catalytic enantioselective synthesis of α -trifluoromethyl amines are fairly uncommon, with most approaches to these enantioenriched compounds relying on chiral auxiliary chemistry.²

A valuable subset of these compounds are α -trifluoromethyl homoallylic amines. Recently, a number of groups have investigated allylic substitution approaches to these molecules utilizing azaallyl anion building blocks.³ With an isatin⁴ activating group for the nucleophile (Scheme 1), an Ir-catalyzed procedure gives rise to branch-selective coupling of the allylic carbonate exclusively at the azadienolate's α -position; however, this product spontaneously undergoes aza-Cope rearrangement to deliver a net γ -allylation. As a result, the unsaturated amines bear only one stereogenic center.^{4,5} The same class of nucleophiles have been utilized in a γ -selective allylation with Morita–Baylis–Hillman-type allylic carbonates by employing a chiral nucleophilic catalyst (Scheme 1).⁶ These coupling processes can yield homoallylic amines with vicinal *syn* stereogenic centers;^{6a} however, the product scope is limited to aryl-substituted allylic centers and carbonyl-containing alkenes.^{7,8}


Given our longstanding interests in both umpolung synthesis of amines⁹ and diene hydrofunctionalization reactions,¹⁰ we envisioned that the merger of these strategies would allow for the enantio-, diastereo-, and regioselective preparation of α - CF_3 homoallylic amines that bear vicinal stereogenic centers and internal alkenes,^{10a} potentially enabling the *anti* diastereomer to be accessed (Scheme 1). Several challenges had to be met and overcome for the successful realization of this idea.

Scheme 1. Prior Catalytic Enantioselective α - CF_3 Homoallylic Amine Synthesis and Proposed Strategy


■ Ir-Catalyzed α -Selective Allylation/Aza-Cope Rearrangement



■ *syn*- & γ -Selective Allylation with MBH Carbonates




■ Proposal: *anti*- & γ -Selective Azadienolate–Diene Coupling



Received: January 25, 2020

Published: February 4, 2020



Table 1. Optimization for γ -Selective Addition of an Isatin-Derived Azadienolate to Phenylbutadiene^a

entry	L	NaBARF ₄ (Y/N)	3a:3a' ^b	conv to 3a/3a' ^b (%)	dr of 3a ^b	er of 3a ^c
1	BINAP	N	1:2.5	48	1:1	nd
2	MeO-BIPHEP	N	1:2.5	66	1.5:1	nd
3	DTBM-MeO-BIPHEP	N	>20:1	49	7:1	nd
4	SEGPPOS	N	1:1.1	30	3:1	nd
5	DM-SEGPPOS	N	1:1.5	38	3:1	nd
6	DTBM-SEGPPOS	N	>20:1	>98	9:1	91.5:8.5
7	DTBM-SEGPPOS	Y	>20:1	>98	9:1	93:7
8 ^d	DTBM-SEGPPOS	N	>20:1	>98	8:1	94:6
9 ^{d,e}	DTBM-SEGPPOS	Y	>20:1	97 (80) ^f	8:1	95:5

^aReactions run under N₂ with 0.15 mmol of diene **2a** (0.75 M). ^bDetermined by 376 MHz ¹⁹F NMR spectroscopy of the unpurified mixture.


^cDetermined by HPLC analysis of purified **3a**. ^d1,4-Dioxane as solvent. ^eReaction for 12 h. ^fIsolated yield of purified **3a**. nd = not determined.

Primary among these was regioselectivity. In contrast to previous metal-catalyzed allylic substitution methods, our goal was to develop a kinetically γ -selective allylation of isatin azadienolates, obviating the aza-Cope rearrangement and thus giving rise to a fundamentally different product connectivity than would otherwise be available. Could a catalyst be found that is γ -selective, and would this process still be efficient? Would the same catalyst also provide regioselectivity with respect to the diene? Finally, would such a catalyst allow for control of the relative and absolute stereochemistry of the homoallylic amine products? Only recently has a diene hydroalkylation that sets two stereogenic centers in one bond-forming event been reported.¹¹ Herein, we illustrate that Pd–DTBM-SEGPPOS promotes the γ -selective coupling of azadienolates with the terminal olefin of dienes, generating the *anti* diastereomer of homoallylic α -CF₃ amines with excellent levels of stereocontrol.¹²

We initiated our investigations by exploring the coupling of *N*-trifluoroethyl imine **1** and phenylbutadiene **2a** (Table 1).¹³ Whereas most ligands favor azadienolate α -alkylation product **3a'** under Pd catalysis, bis(phosphines) comprised of 3,5-di-*tert*-butyl-4-methoxy (DTBM) aryl groups at phosphorus exclusively deliver the desired γ -alkylation product **3a** (compare entries 3 and 6 with entries 1–2 and 4–5).¹⁴ Notably, the diastereomeric ratio for **3a** is also considerably higher with the more γ -selective catalysts, affording the *anti* diastereomer as the major isomer (entries 3 and 6). Use of DTBM-SEGPPOS provides the greatest conversion to **3a**, which is isolated in 9:1 dr and 91.5:8.5 er (entry 6).¹⁵ We discovered that the enantioselectivity could be improved by the addition of 10 mol % NaBARF₄ (93:7 er, entry 7).¹⁶ By switching the solvent from diethyl ether to 1,4-dioxane, the γ -alkylation product **3a** is isolated in 80% yield, 8:1 dr, and 95:5 er (entry 9).

Taking the conditions in Table 1, entry 9, as optimal, we next explored the scope of aryl-substituted terminal dienes for

coupling with **1** (Table 2). Both electron-rich (e.g., entries 1 and 6–7) and electron-poor (entries 2–3) dienes generate the homoallylic amine products with $\geq 87\%$ conversion, $\geq 7:1$ dr, and moderate to good enantioselectivity. Heterocycles are tolerated, with furyl-substituted **3i** (entry 8) and pyridyl-containing **3j** (entry 9) isolated in good yields and stereo-

Table 2. Aryl Diene Scope in Azadienolate Coupling^a

entry	product (3); R ¹	conv to 3a ^b (%) yield of 3 ^c (%)	dr of 3 ^b	er of 3 ^d
1	3b ; 4-(MeO)(C ₆ H ₄)	87; 68	10:1	94:6
2	3c ; 4-Cl(C ₆ H ₄)	98; 75	7:1	90.5:9.5
3	3d ; 4-(F ₃ C)(C ₆ H ₄)	94; 74	8:1	90:10
4	3e ; 3-Me(C ₆ H ₄)	89; 72	6:1	93:7
5 ^e	3f ; 2-Me(C ₆ H ₄)	63; 59	2:1	92.5:7.5
6	3g ; 2-naphthyl	94; 62	8:1	94:6
7	3h ; 3,4-dioxolato(C ₆ H ₃)	94; 86	10:1	91.5:8.5
8	3i ; 2-furyl	69; 59	8:1	93:7
9	3j ; 3-pyridyl	83; 82	7:1	94:6

^aReactions run under N₂ with 0.15 mmol of diene **2** (0.75 M).

^bDetermined by 376 MHz ¹⁹F NMR or 400 MHz ¹H NMR spectroscopy of the unpurified mixture. ^cIsolated yield of purified **3**.

^dDetermined by HPLC analysis of purified **3**. ^eReaction run without NaBARF₄; 9:1 **3f**:**3f'**.

selectivities. The majority of dienes lead exclusively to the γ -alkylation product; however, an *ortho* substituent on the arene results in a more sluggish reaction (44% conversion in 12 h) that furnishes approximately 10% α -coupling of the azadienolate (entry 5). Improving reaction efficiency required the omission of $\text{NaBAR}^{\text{F}}_4$, and while diastereoselectivity was lower, enantioselectivity remained high (92.5:7.5 er). In all, aryl-substituted dienes readily participate in couplings with **1** at room temperature, affording homoallylic α -trifluoromethyl amines **3b–j** in 59–86% yield.

Alkyl-substituted terminal dienes are also effective coupling partners; however, the Pd–DTBM–SEGPHOS-catalyzed processes with imine **1** require elevated temperature (50 °C) to proceed effectively (Table 3). Consequently, we also omitted

Table 3. Azadienolate–Alkyl Diene Coupling Scope^a


entry	product (3); R ¹	conv to 3 (%); ^b yield of 3 (%) ^c	dr of 3 ^b	er of 3 ^d
1 ^e	3k ; Ph(CH ₂) ₂	82; 69	4:1	88:12
2	3l ; <i>n</i> -hexyl	65; 60	4:1	81.5:18.5
3	3m ; EtO ₂ C(CH ₂) ₂	84; 83	3:1	91:9
4	3n ; Cy	61; 43	3:1	90:10
5 ^f	3o ;	83; 83	3:1	86.5:13.5

^aReactions run under N₂ with 0.15 mmol of diene **2** (0.75 M).
^bDetermined by 376 MHz ¹⁹F NMR or 400 MHz ¹H NMR spectroscopy of the unpurified mixture. ^cIsolated yield of purified **3**.
^dDetermined by HPLC analysis of purified **3**. ^eA 10:1 mixture of γ -alkylation products **3k** and **3k'** was formed; see ref 17. ^f11:1 **3o**:**3o'**.

the $\text{NaBAR}^{\text{F}}_4$ additive to enable the reaction to proceed at a higher rate. Both linear (entries 1–3) and α -branched (entries 4–5) dienes participate in the reactions, affording homoallylic amines **3k–o** in 43–83% yield in up to 4:1 dr and 91:9 er. Products derived from isomerization of the diene along the alkyl chain (“chain walking”) prior to enolate addition could not be detected, including with phenethyl **2k** or heptadienoate **2m**. Alkyl dienes largely or solely lead to γ -alkylation of the azadienolate, although it is notable that piperidine **2o** affords ca. 9% α product **3o'**. We also observed roughly 8% of an additional γ -alkylation product **3k''** with phenethyl diene **2k**. Homoallylic amine **3k''** bears a different connectivity with respect to the diene-derived fragment, and a series of experiments suggest that **3k''** is formed from the aza-Cope rearrangement of α -alkylation product **3k'**.¹⁷

Intriguingly, in the course of our alkyl diene studies, we discovered that hexadienoate **2p** (Scheme 2) undergoes diene isomerization into conjugation with the ester prior to its coupling with **1**, furnishing the ethyl-substituted stereogenic center of homoallylic amine **3p**. The process is reasonably

Scheme 2. Isomerization/Alkylation of 3,5-Hexadienoate and Comparison to Its Internal Diene Isomer



efficient, with the α -CF₃ amine obtained in 48% yield, 12:1 dr, and 97.5:2.5 er. Comparatively, its internal diene analogue **4** also delivers acrylate **3p** with similar levels of stereoselectivity but lower conversion.

We have explored a number of additional reaction partners to expand the scope of the hydrofunctionalization (Figure 1).





Figure 1. Limitations in Reaction Partners.

Imine **5** was tested in a coupling with diene **2a** in order to access α -difluoromethyl amines, but unfortunately, the nucleophile undergoes complete decomposition without alkylation. Substituted imines, such as **6**, would form products bearing tetrasubstituted stereogenic centers^{5,6b,8} but were unreactive. Other dienes were also investigated. Alcohol- and silyl ether-containing alkyl dienes **2q** and **2r** lead to a complex mixture of products, which we surmise to be a combination of the desired γ -alkylation **3**, the regiomer α -alkylation **3'**, and the aza-Cope rearrangement products **3''**, all as a mixture of diastereomers, rather than products attributable to chain walking.

The diene hydroalkylation with imine **1** can be performed on a 1.0 mmol scale to furnish the α -trifluoromethyl isatin-protected homoallylic amine **3a** in 82% yield (Scheme 3). Additionally, hydrolysis of the isatin moiety under mildly acidic conditions delivers the free amine **7** in 77% yield.

Catalytic enantioselective diene hydrofunctionalization provides an enabling route toward highly valuable chemical building blocks that are not readily prepared by other methods. Here, in combination with imine umpolung,³ we have shown that important homoallylic α -trifluoromethyl amines bearing contiguous stereogenic centers and an internal olefin can be accessed for the first time. Utilizing an isatin auxiliary, we have discovered that, in contrast to other transition-metal-catalyzed processes, palladium ligated with DTBM–SEGPHOS allows for regioselective γ -alkylation of the derived azadienolate, generat-

Scheme 3. Scale Up of Azadienolate Hydroalkylation and Product Imine Hydrolysis



ing the *anti* diastereomer of the homoallylic $\alpha\text{-CF}_3$ amines with high levels of stereocontrol. This catalytic process should open up new chemical space for drug discovery.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c00342>.

Experimental procedures, analytical data for new compounds, X-ray crystallographic data, and NMR spectra (PDF)

Accession Codes

CCDC 1978720 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

Steven J. Malcolmson – Department of Chemistry, Duke University, Durham, North Carolina 27708, United States;
 orcid.org/0000-0003-3229-0949;
 Email: steven.malcolmson@duke.edu

Authors

Chibueze I. Onyeagusi – Department of Chemistry, Duke University, Durham, North Carolina 27708, United States
 Xinxin Shao – Department of Chemistry, Duke University, Durham, North Carolina 27708, United States

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.orglett.0c00342>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was generously supported by the NIH (GM124286) and the NSF (CHE-1800012). C.I.O. is grateful to the Duke Chemistry Department for a Burroughs-Wellcome fellowship. All X-ray crystallographic measurements were made in the Molecular Education, Technology, and Research Innovation Center (METRIC) at NC State University; we thank Dr. Roger Sommer (NC State) for assistance with analysis.

■ REFERENCES

- (1) (a) Meanwell, N. A. Fluorine and Fluorinated Motifs in the Design and Application of Bioisosteres for Drug Design. *J. Med. Chem.* **2018**, *61*, 5822. (b) Gillis, E. P.; Eastman, K. J.; Hill, M. D.; Donnelly, D. J.; Meanwell, N. A. Applications of Fluorine in Medicinal Chemistry. *J. Med. Chem.* **2015**, *58*, 8315.
- (2) For reviews, see: (a) Fioravanti, S. Trifluoromethyl Aldimines: An Overview in the Last Ten Years. *Tetrahedron* **2016**, *72*, 4449. (b) Nie, J.; Guo, H.-C.; Cahard, D.; Ma, J.-A. Asymmetric Construction of Stereogenic Carbon Centers Featuring a Trifluoromethyl Group from Prochiral Trifluoromethylated Substrates. *Chem. Rev.* **2011**, *111*, 455. For additional examples not highlighted in these reviews, see: (c) Grellepois, F.; Ben Jamaa, A.; Saraiva Rosa, N. α -Trifluoromethylated Tertiary Homoallylic Amines: Diastereoselective Synthesis and Conversion into β -Aminoesters, γ - and δ -Aminoalcohols, Azetidines and Pyrrolidines. *Org. Biomol. Chem.* **2017**, *15*, 9696. (d) Grellepois, F.; Ben Jamaa, A.; Gassama, A. Diastereoselective Addition of Organomagnesium and Organolithium Reagents to Chiral Trifluoromethyl *N*-tert-Butanesulfinyl Hemiaminals. *Eur. J. Org. Chem.* **2013**, *2013*, 6694. (e) Guo, T.; Song, R.; Yuan, B.-H.; Chen, X.-Y.; Sun, X.-W.; Lin, G.-Q. Highly Efficient Asymmetric Construction of Quaternary Carbon-Containing Homoallylic and Homopropargylic Amines. *Chem. Commun.* **2013**, *49*, 5402.
- (3) For a review on azaallyl anions as imine umpolung reagents, see: (a) Tang, S.; Zhang, X.; Sun, J.; Niu, D.; Chruma, J. J. 2-Azaallyl Anions, 2-Azaallyl Cations, 2-Azaallyl Radicals, and Azomethine Ylides. *Chem. Rev.* **2018**, *118*, 10393. For other reviews on umpolung chemistry, see: (b) Mahatthananchai, J.; Bode, J. W. On the Mechanism of N-Heterocyclic Carbene-Catalyzed Reactions Involving Acyl Azoliums. *Acc. Chem. Res.* **2014**, *47*, 696. (c) Boyce, G. R.; Greszler, S. N.; Johnson, J. S.; Linghu, X.; Malinowski, J. T.; Nicewicz, D. A.; Satterfield, A. D.; Schmitt, D. C.; Steward, K. M. Silyl Glyoxylates. Conception and Realization of Flexible Conjunctive Reagents for Multicomponent Coupling. *J. Org. Chem.* **2012**, *77*, 4503. (d) Izquierdo, J.; Hutson, G. E.; Cohen, D. T.; Scheidt, K. A. A Continuum of Progress: Applications of N-Heterocyclic Carbene Catalysis in Total Synthesis. *Angew. Chem., Int. Ed.* **2012**, *51*, 11686. (e) Smith, A. B., III; Wuest, W. M. Evolution of Multi-Component Anion Relay Chemistry (ARC): Construction of Architecturally Complex Natural and Unnatural Products. *Chem. Commun.* **2008**, *44*, 5883. (f) Seebach, D. Methods of Reactivity Umpolung. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 239.
- (4) Shi, L.-M.; Sun, X.-S.; Shen, C.; Wang, Z.-F.; Tao, H.-Y.; Wang, C.-J. Catalytic Asymmetric Synthesis of α -Trifluoromethyl Homoallylic Amines via Umpolung Allylation/2-Aza-Cope Rearrangement: Stereoselectivity and Mechanistic Insight. *Org. Lett.* **2019**, *21*, 4842.
- (5) Azaallyl anions formed from fluorenyl imines have likewise been shown to undergo enantioselective allylation followed by 2-aza-Cope rearrangement; see: (a) Wang, Y.; Deng, L.-F.; Zhang, X.; Niu, D. Catalytic Asymmetric Synthesis of α -Tetrasubstituted α -Trifluoromethyl Homoallylic Amines by Ir-Catalyzed Umpolung Allylation of Imines. *Org. Lett.* **2019**, *21*, 6951. (b) Shen, C.; Wang, R.-Q.; Wei, L.; Wang, Z.-F.; Tao, H.-Y.; Wang, C.-J. Catalytic Asymmetric Umpolung Allylation/2-Aza-Cope Rearrangement for the Construction of α -Tetrasubstituted α -Trifluoromethyl Homoallylic Amines. *Org. Lett.* **2019**, *21*, 6940.
- (6) (a) Li, X.; Su, J.; Liu, Z.; Zhu, Y.; Dong, Z.; Qiu, S.; Wang, J.; Lin, L.; Shen, Z.; Yan, W.; Wang, K.; Wang, R. Synthesis of Chiral α -Trifluoromethylamines with 2,2,2-Trifluoroethylamine as a “Building

Block. *Org. Lett.* **2016**, *18*, 956. For a similar reaction, see: (b) Chen, P.; Yue, Z.; Zhang, J.; Lv, X.; Wang, L.; Zhang, J. Phosphine Catalyzed Asymmetric Umpolung Addition of Trifluoromethyl Ketimines to Morita–Baylis–Hillman Carbonates. *Angew. Chem., Int. Ed.* **2016**, *55*, 13316. For a related method with allenolate electrophiles, see: (c) Gao, Y.-N.; Shi, M. Enantioselective Synthesis of Isatin-Derived α -(Trifluoromethyl)imine Derivatives: Phosphine-Catalyzed γ -Addition of α -(Trifluoromethyl)imines and Allenolates. *Eur. J. Org. Chem.* **2017**, 2017, 1552.

(7) For additional examples of enantioselective homoallylic amine synthesis involving stereoselective sigmatropic rearrangement, see: (a) Wei, L.; Zhu, Q.; Xiao, L.; Tao, H.-Y.; Wang, C.-J. Synergistic Catalysis for Cascade Allylation and 2-aza-Cope Rearrangement of Azomethine Ylides. *Nat. Commun.* **2019**, *10*, 1594. (b) Huo, X.; Zhang, J.; Fu, J.; He, R.; Zhang, W. Ir/Cu Dual Catalysis: Enantio- and Diastereodivergent Access to α,α -Disubstituted α -Amino Acids Bearing Vicinal Stereocenters. *J. Am. Chem. Soc.* **2018**, *140*, 2080. (c) Wei, L.; Zhu, Q.; Xu, S.-M.; Chang, X.; Wang, C.-J. Stereodivergent Synthesis of α,α -Disubstituted α -Amino Acids via Synergistic Cu/Ir Catalysis. *J. Am. Chem. Soc.* **2018**, *140*, 1508.

(8) For additional examples of catalytic enantioselective synthesis of α -trifluoromethyl amines not covered in refs 2 and 4–6, see: (a) Hu, B.; Deng, L. Direct Catalytic Asymmetric Synthesis of Trifluoromethylated γ -Amino Esters/Lactones via Umpolung Strategy. *J. Org. Chem.* **2019**, *84*, 994. (b) Zhu, J.; Huang, L.; Dong, W.; Li, N.; Yu, X.; Deng, W.-P.; Tang, W. Enantioselective Rhodium-Catalyzed Addition of Arylboroxines to N-Unprotected Ketimines: Efficient Synthesis of Cipargamin. *Angew. Chem., Int. Ed.* **2019**, *58*, 16119. (c) Hu, B.; Bezpalko, M. W.; Fei, C.; Dickie, D. A.; Foxman, B. M.; Deng, L. Origin of and a Solution for Uneven Efficiency by Cinchona Alkaloid-Derived, Pseudoenantiomeric Catalysts for Asymmetric Reactions. *J. Am. Chem. Soc.* **2018**, *140*, 13913. (d) Li, Z.; Hu, B.; Wu, Y.; Fei, C.; Deng, L. Control of Chemoselectivity in Asymmetric Tandem Reactions: Direct Synthesis of Chiral Amines Bearing Nonadjacent Stereocenters. *Proc. Natl. Acad. Sci. U. S. A.* **2018**, *115*, 1730. (e) Hu, B.; Deng, L. Catalytic Asymmetric Synthesis of Trifluoromethylated γ -Amino Acids through the Umpolung Addition of Trifluoromethyl Imines to Carboxylic Acid Derivatives. *Angew. Chem., Int. Ed.* **2018**, *57*, 2233. (f) Hu, L.; Wu, Y.; Li, Z.; Deng, L. Catalytic Asymmetric Synthesis of Chiral γ -Amino Ketones via Umpolung Reactions of Imines. *J. Am. Chem. Soc.* **2016**, *138*, 15817. (g) Kawai, H.; Kusuda, A.; Nakamura, S.; Shiro, M.; Shibata, N. Catalytic Enantioselective Trifluoromethylation of Azomethine Imines with Trimethyl-(trifluoromethyl)silane. *Angew. Chem., Int. Ed.* **2009**, *48*, 6324.

(9) (a) Malcolmson, S. J.; Li, K.; Shao, X. 2-Azadienes as Enamine Umpolung Synthons for the Preparation of Chiral Amines. *Synlett* **2019**, 30, 1253. (b) Shao, X.; Malcolmson, S. J. Catalytic Enantio- and Diastereoselective Cyclopropanation of 2-Azadienes for the Synthesis of Aminocyclopropanes Bearing Quaternary Carbon Stereogenic Centers. *Org. Lett.* **2019**, *21*, 7380. (c) Daniel, P. E.; Onyeagusi, C. I.; Ribeiro, A. A.; Li, K.; Malcolmson, S. J. Palladium-Catalyzed Synthesis of α -Trifluoromethyl Benzylic Amines via Fluoroarylation of *gem*-Difluoro-2-azadienes Enabled by Phosphine-Catalyzed Formation of an Azaallyl–Silver Intermediate. *ACS Catal.* **2019**, *9*, 205. (d) Shao, X.; Li, K.; Malcolmson, S. J. Enantioselective Synthesis of *anti*-1,2-Diamines by Cu-Catalyzed Reductive Couplings of Azadienes with Aldimines and Ketimines. *J. Am. Chem. Soc.* **2018**, *140*, 7083. (e) Li, K.; Shao, X.; Tseng, L.; Malcolmson, S. J. 2-Azadienes as Reagents for Preparing Chiral Amines: Synthesis of 1,2-Amino Tertiary Alcohols by Cu-Catalyzed Enantioselective Reductive Couplings with Ketones. *J. Am. Chem. Soc.* **2018**, *140*, 598. (f) Li, K.; Weber, A. E.; Tseng, L.; Malcolmson, S. J. Diastereoselective and Enantiospecific Synthesis of 1,3-Diamines via 2-Azaallyl Anion Benzylic Ring-Opening of Aziridines. *Org. Lett.* **2017**, *19*, 4239. (g) Daniel, P. E.; Weber, A. E.; Malcolmson, S. J. Umpolung Synthesis of 1,3-Amino Alcohols: Stereoselective Addition of 2-Azaallyl Anions to Epoxides. *Org. Lett.* **2017**, *19*, 3490.

(10) (a) Adamson, N. J.; Malcolmson, S. J. Catalytic Enantio- and Regioselective Addition of Nucleophiles in the Intermolecular

Hydrofunctionalization of 1,3-Dienes. *ACS Catal.* **2020**, *10*, 1060. (b) Park, S.; Adamson, N. J.; Malcolmson, S. J. Brønsted Acid and Pd–PHOX Dual-Catalyzed Enantioselective Addition of Activated C–Pronucleophiles to Internal Dienes. *Chem. Sci.* **2019**, *10*, 5176. (c) Park, S.; Malcolmson, S. J. Development and Mechanistic Investigations of Enantioselective Pd-Catalyzed Intermolecular Hydroaminations of Internal Dienes. *ACS Catal.* **2018**, *8*, 8468. (d) Adamson, N. J.; Wilbur, K. C. E.; Malcolmson, S. J. Enantioselective Intermolecular Pd-Catalyzed Hydroalkylation of Acyclic 1,3-Dienes with Activated Pronucleophiles. *J. Am. Chem. Soc.* **2018**, *140*, 2761. (e) Adamson, N. J.; Hull, E.; Malcolmson, S. J. Enantioselective Intermolecular Addition of Aliphatic Amines to Acyclic Dienes with a Pd–PHOX Catalyst. *J. Am. Chem. Soc.* **2017**, *139*, 7180.

(11) This report utilizes a dual metal-catalyzed strategy; see: (a) Zhang, Q.; Yu, H.; Shen, L.; Tang, T.; Dong, D.; Chai, W.; Zi, W. Stereodivergent Coupling of 1,3-Dienes with Aldimine Esters Enabled by Synergistic Pd and Cu Catalysis. *J. Am. Chem. Soc.* **2019**, *141*, 14554. For transformations with allenes/alkynes, see: (b) Bora, P. P.; Sun, G.-J.; Zheng, W.-F.; Kang, Q. Rh/Lewis Acid Catalyzed Regio-, Diastereo- and Enantioselective Addition of 2-Acyl Imidazoles with Allenes. *Chin. J. Chem.* **2018**, *36*, 20. (c) Cruz, F. A.; Dong, V. M. Stereodivergent Coupling of Aldehydes and Alkynes via Synergistic Catalysis Using Rh and Jacobsen's Amine. *J. Am. Chem. Soc.* **2017**, *139*, 1029. (d) Zhou, H.; Wei, Z.; Zhang, J.; Yang, H.; Xia, C.; Jiang, G. From Palladium to Brønsted Acid Catalysis: Highly Enantioselective Regiodivergent Addition of Alkoxyallenes to Pyrazolones. *Angew. Chem., Int. Ed.* **2017**, *56*, 1077. (e) Trost, B. M.; Xie, J.; Sieber, J. D. The Palladium Catalyzed Asymmetric Addition of Oxindoles and Allenes: An Atom-Economical Versatile Method for the Construction of Chiral Indole Alkaloids. *J. Am. Chem. Soc.* **2011**, *133*, 20611. (f) Trost, B. M.; Jäkel, C.; Plietker, B. Palladium-Catalyzed Asymmetric Addition of Pronucleophiles to Allenes. *J. Am. Chem. Soc.* **2003**, *125*, 4438.

(12) A preprint of this manuscript was published on January 20, 2020; see: Onyeagusi, C. I.; Shao, X.; Malcolmson, S. J. Enantio- and Diastereoselective Synthesis of Homoallylic α -Trifluoromethyl Amines by Catalytic Hydroalkylation of Dienes. 2020. (*ChemRxiv*). <https://doi.org/10.26434/chemrxiv.11653143.v1>.

(13) For additional screening data, see the Supporting Information.

(14) The transformations in Table 1 utilize a preformed Pd–bis(phosphine) complex derived from $[\text{Pd}(\eta^3\text{-cinnamyl})\text{Cl}]_2$, bis(phosphine), and $\text{NaBAR}^{\text{F}_4}$. When conducting reactions directly with *in situ*-generated catalyst, we observe a significant induction period in the reaction that is avoided by employing an isolated complex. See the Supporting Information for further details.

(15) Pd–DTBM-SEGPHOS has been shown to be the optimal catalyst in a handful of enantioselective diene hydrophosphinylation reactions; see: Nie, S.-Z.; Davison, R. T.; Dong, V. M. Enantioselective Coupling of Dienes and Phosphine Oxides. *J. Am. Chem. Soc.* **2018**, *140*, 16450.

(16) Although $\text{NaBAR}^{\text{F}_4}$ improves enantioselectivity, it greatly reduces the reaction rate. Under the conditions shown in Table 1, entry 6 without $\text{NaBAR}^{\text{F}_4}$, the reaction is complete within 6 h, but longer reaction times are needed the more $\text{NaBAR}^{\text{F}_4}$ is added. The data in Table 1 are all shown for 16 h reaction time for comparative purposes. See the Supporting Information for additional details.

(17) See the Supporting Information for a detailed discussion.

