

Constraining the fraction of core-collapse supernovae harbouring choked jets with high-energy neutrinos

Dafne Guetta,¹★ Roi Rahin,² Imre Bartos³★ and Massimo Della Valle^{4,5}

¹*ORT Braude, 216002 Karmiel, Israel*

²*Technion, 3200003 Haifa, Israel*

³*Department of Physics, University of Florida, PO Box 118440, Gainesville, FL 32611-8440, USA*

⁴*Capodimonte Observatory, INAF-Naples, Salita Moiariello 16, I-80131 Naples, Italy*

⁵*European Southern Observatory, Karl-Schwarzschild-Strae 2, D-85748 Garching bei München, Germany*

Accepted 2019 November 17. Received 2019 November 3; in original form 2019 June 18

ABSTRACT

The joint observation of core-collapse supernovae with gamma-ray bursts shows that jets can be launched in the aftermath of stellar core collapse, likely by a newly formed black hole that accretes matter from the star. Such gamma-ray bursts have only been observed accompanying Type Ibc supernovae, indicating a stellar progenitor that lost its hydrogen envelope before collapse. According to recent hypothesis, it is possible that jets are launched in core-collapse events even when the progenitors still retain their hydrogen envelopes; however, such jets are not able to burrow through the star and will be stalled into the interior of the progenitor star before escaping. These jets are called choked jets. High-energy neutrinos produced by such choked jets could escape the stellar envelope and could be observed. Here, we examine how multimessenger searches for high-energy neutrinos and core-collapse supernovae can detect or limit the fraction of stellar collapses that produce jets. We find that a high fraction of jet production is already limited by previous observational campaigns. We explore possibilities with future observations using Large Synoptic Survey Telescope, IceCube, and Km3NET.

Key words: gravitational waves – stars: neutron.

1 INTRODUCTION

The mechanism driving core-collapse supernova (CCSN) explosions is still unclear. A common explanation is neutrino-driven outflows (Colgate & Petschek 1982); however, a recent idea is that jets could play an important role in SN explosions (Piran et al. 2019). CCSNe associated with long duration gamma-ray bursts (GRBs), sometimes called ‘hypernovae’, provide a clear evidence for strongly aspherical explosions (Maeda et al. 2008), suggesting a jet-like activity. Observations strongly suggest that a tiny fraction of CC (GRB-SNe) is powered by jets (see Berger et al. 2003; Soderberg et al. 2006; Guetta & Della Valle 2007). Recently, we have observed in GRB171205A/SN 2017iuk the associated ‘cocoon’ (Izzo et al. 2019). This fact opens the interesting possibility that also broad-lines SNe without GRBs, which are energetically similar to GRB-SNe ($\sim 10^{52}$ erg) may be powered by ‘cocoon’.

Piran et al. (2019) explored that a similar mechanism might be active also in ‘standard’ CCSNe types not associated with GRBs. Such a scenario is suggested by observations indicating energy deposition by the so-called ‘choked’ jets in a cocoon within the stellar envelope.

Oded & Soker (2011) discusses a scenario characterized by non-relativistic jets in CCSNe. The cocoon eventually breaks out from the star releasing energetic material at very high, yet subrelativistic, velocities. It is possible to identify this fast-moving material by looking at the very early-time supernova spectra, as this component has a unique signature. Recently, a clear evidence has been provided indicating such a deposition by Izzo et al. (2019), who detected spectroscopic signatures of iron group elements moving at $\sim 120\,000$ km s⁻¹ originating from the innermost parts of the exploding progenitor star of SN 2017iuk. However, whether or not CCSNe that do not harbour GRBs might be powered by jet/cocoon activity is still a matter of active discussion within the community. The most natural interpretation of the energetic fast-moving component observed in the early spectra of the supernovae by Piran et al. (2019) is that this is the cocoon’s matter that breaks out from the progenitor. In a few cases such as SN 2008D/GRB 080109 (Mazzali et al. 2008), SN 2006aj/GRB 060218 (Campana et al. 2006) and GRB 171205A/2017iuk (Izzo et al. 2019), we might have even seen the direct thermal emission of this hot cocoon material. In this case, Piran’s interpretation implies the existence of powerful jets within these supernovae. These jets have difficulty to penetrate the progenitor; therefore, these objects are simply observed as energetic supernovae or hypernovae in the optical band and do not have accompanying gamma-ray emission. The

* E-mail: dafneguetta@braude.ac.il (DG); imrebartos@ufl.edu (IB)

jets that do not succeed to drill the envelope are labelled ‘choked’ jets (CJ).

It may be that a significant fraction of CCSNe, and possibly all those that have not lost their hydrogen envelope, harbour choked jets. The shocks involved in these hidden jets may be the source of high-energy neutrinos observed by IceCube (see Senno, Murase & Meszaros 2016; He et al. 2018).

In choked jets, neutrinos and gamma-rays are produced by the decay of pions produced in the interaction of accelerated protons and thermal photons. While these sources are transparent to neutrinos, they are opaque to gamma-ray photons as these particles are produced inside the stellar envelope. This can justify the lack of association between observed GRBs and IceCube neutrinos. Therefore, these sources are dark in GeV–TeV gamma-rays, and do not contribute to the Fermi diffuse gamma-ray background implying that the Waxman–Bahcall bound and the Fermi constraint on the diffuse gamma-ray emission do not apply to these sources (Murase, Guetta & Ahlers 2016).

These characteristics have been used by several authors (e.g. Murase & Ioka 2013; Murase et al. 2016; Senno et al. 2016; Denton & Tamborra 2017; Senno, Murase & Meszaros 2017; Esmaili & Murase 2018) to show that choked jets’ so-called failed GRBs may be the possible sources for the observed diffuse neutrino flux.

In this paper, we check if we can use the neutrino emission to test the hypothesis that most supernovae produce jets (e.g. Piran et al. 2019). If only a small fraction of CCSNe is powered by jets (e.g. only GRB-SNe), the number of neutrinos revealed by neutrino detectors will be smaller than the number expected if the totality of CCSNe (including, type II, Ibc and GRB-SNe) is powered by jets.

We investigate the detectability of high-energy neutrinos from choked jets within CCSNe, and the limits set by non-detection on the fraction of CCSNe that harbour jets. For this analysis, we assume that the conditions set by Senno et al. (2016) are satisfied and use the model of He et al. (2018) to calculate the expected neutrino flux from a single source. In their estimate they include the relevant microphysical processes such as multipion production in pp and p-gamma interactions, as well as the energy losses of mesons and muons.

2 NEUTRINO EMISSION ESTIMATE FROM CHOKED JETS

We consider the possibility that a fraction of rapidly rotating massive stars at the end of their lives undergo core collapse, form a compact star or a black hole, and produce a jet that it is stalled before it breaks through the star (MacFadyen, Woosley & Heger 2001; Meszaros & Waxman 2001). We take the stellar parameters given in Meszaros & Waxman (2001).

Following Senno et al. (2016), we assume that the photons are free to move inside the jet as the plasma is optically thin inside the jet but cannot escape as the envelope or circumstellar medium outside is largely optically thick to Thomson scattering. Since the photons are trapped inside the jet, the protons can interact with these thermal photons very efficiently. We can assume that the fraction of protons converted in pions, the so-called f_π , is almost 1 in this process. The radiation constraints apply to shocks in the envelope material as well as those in the choked jet. With their model, Senno et al. (2016) have estimated the diffuse neutrino flux from choked jets. In this paper, we are interested in estimating the flux from a single source.

We consider the possibility that electrons and protons are accelerated in the internal shock model and estimate the neutrino flux and spectrum from a single source following the model of He et al. (2018). The model of He et al. (2018) is valid if the conditions described in Senno et al. (2016) are met: the jet is choked, protons are accelerated to high energies efficiently, and thermal photons are produced in the jet head and propagate into the internal shock region. In this model, the accelerated protons interact with the photons from the choked jet head and produce pions. The pions decay into high-energy neutrinos. He et al. (2018) estimate the neutrino spectra numerically, taking into account microphysical processes. There is a low-energy cut-off in the neutrino spectra due to the threshold of the photo–meson interaction. The other cut-off is at high energy due to the photo–meson cooling of protons, or the synchrotron cooling of pions and muons.

In this model, the parameters that most affect the neutrino flux are the Lorentz factor, the duration of the jet, and the luminosity of the jet. In our analysis, we use the same values for these parameters as taken in He et al. (2018). These are $\Gamma = 100$, 10 , $t = 100$, 3.3×10^4 s, and $L = 10^{51}$, 3×10^{48} erg s⁻¹. He et al. (2018) show how different set of parameters imply different neutrino fluxes and spectra; see fig. 3 of He et al. (2018). Their results are consistent with the diffuse flux found by Senno et al. (2016) and therefore with the upper limits given from IceCube on point source analysis and on diffuse flux analysis.

3 NEUTRINO DETECTION

High-energy neutrinos interact with nucleons present in the detector producing secondary particles, which travel faster than the speed of light in the sea or ice, and therefore induce the emission of Cherenkov light.

Currently operating high-energy neutrino detectors include: IceCube, a cubic-kilometre observatory located at the geographic South Pole (Aartsen et al. 2014a); the ANTARES detector deep in the Mediterranean sea (Ageron et al. 2011); and KM3Net (Adrian-Martinez et al. 2016), also in the Mediterranean, currently under construction.

We determine the total number of expected neutrinos by a source by integrating the neutrino fluence convoluted with the effective energy of the detector over the energy range 1–100 TeV

$$N(r) = \int_{1\text{TeV}}^{100\text{TeV}} \frac{dN_\nu(E(1+z))}{dE_\nu} A(E_\nu, \delta) dE_\nu \quad (1)$$

where r is the luminosity distance, $\frac{dN_\nu}{dE_\nu}$ is the neutrino spectral fluence, $A(E_\nu, \delta)$ is the effective area of the neutrino detector, as a function of the neutrino energy E_ν and of the source declination. In our analysis, we consider the IceCube effective area for time-dependent searches using muon neutrinos from point sources (Aartsen et al. 2014b). The area for point sources is found using the results of Aartsen et al. (2015) and the link <https://icecube.wisc.edu/science/data/PS-3years>.

The number of neutrinos expected from a source at distance of ~ 1 Gpc is consistent with the upper limits found by IceCube as shown in He et al. (2018).

4 FRACTION OF SUPERNOVA OBSERVATIONS WITH DETECTED NEUTRINO COUNTERPART

We now estimate the fraction f_ν of CCSNe detected electromagnetically that will also be detected via high-energy neutrinos. We convolve $N(r)$ with the frequency of occurrence of CCSNe as a

function of redshift. Due to the shortness of the lifetime of the progenitors of CCSNe, typically a few million years, we can use star formation history as proxy of the history of the CCSN rate (e.g. Hopkins & Beacom 2011; Madau & Dickinson 2014). We find that the expected fraction of supernovae within distance r_{\max} that will have a detected astrophysical high-energy neutrino counterpart is

$$f_v = \frac{f_{\text{jet}} f_b \int_0^{r_{\max}} (1 - \text{Pois}(0, N(r))) \rho(r) 4\pi r^2 dr}{\int_0^{r_{\max}} \rho(r) 4\pi r^2 dr} \quad (2)$$

where f_b is the jet beaming factor, defined as the fraction of the sky in which the jet emits high-energy neutrinos, f_{jet} is the fraction of CCSNe that produces jets, $\text{Pois}(k, \lambda)$ is the Poisson probability of measuring k for average value λ , and r_{\max} is the maximum luminosity distance out to which the CCSN can be detected optically. This fraction depends on the neutrino detection threshold through $N(r)$ defined in equation (1). In this definition, we do not include the possibility that a background neutrino can be coincident by chance with the supernova, which we separately account for below.

Here, we consider r_{\max} for two representative facilities for ongoing and future observations: the Zwicky Transient Facility (ZTF; Rauch 2019) and the Large Synoptic Survey Telescope (LSST; <https://www.lsst.org/scientists/scibook>). For ZTF, we assume a magnitude limit of $R \sim 19$ for a typical 30 s exposure (Bellm & Kulkarni 2017; Rauch 2019). For LSST, we assume a magnitude limit of $R \sim 24$ for a typical exposure of 2×15 s (Abell et al. 2009). After assuming an absolute magnitude at maximum for CCSNe of $M = -17.5$ (Patat et al. 1994), we find that they can be detected out to $r \sim 200$ Mpc with ZTF, and out to $r \sim 1.5$ Gpc with LSST. The cosmological effects are taken into account in the analysis. For the LSST case, the redshift effects become important.

With these observational limits, using equation (2) and the model parameters (Model 1) $\Gamma = 100$, $t = 100$ s, and $L = 10^{51}$ erg s $^{-1}$, we find that a fraction $f_v^{\text{ZTF}} \approx 0.5 f_b$ of CCSNe detected by ZTF might produce neutrinos detectable with IceCube or Km3NET. For LSST, this fraction becomes $f_v^{\text{LSST}} \approx 0.01 f_b$. In order to check the effects of the model parameters on our results, we consider another set of model parameters (Model 2) like $\Gamma = 100$, $t = 3.3 \times 10^4$ s, and $L = 3.3 \times 10^{48}$ erg s $^{-1}$, we find that a fraction $f_v^{\text{ZTF}} \approx 0.03 f_b$ of CCSNe detected by ZTF might produce neutrinos detectable with IceCube or Km3NET. For LSST, this fraction becomes $f_v^{\text{LSST}} \approx 5 \times 10^{-4} f_b$. For another set of parameters (Model 3) like $\Gamma = 10$, $t = 3.3 \times 10^4$ s, and $L = 3.3 \times 10^{48}$ erg s $^{-1}$, we find that a fraction $f_v^{\text{ZTF}} \approx 0.01 f_b$ of CCSNe detected by ZTF might produce neutrinos detectable with IceCube or Km3NET. For LSST, this fraction becomes $f_v^{\text{LSST}} \approx 10^{-4} f_b$.

The same estimate can be performed for SNe-Ibc, which are a fraction of CC, about 30 percent (Botticella et al. 2017) but, on average, slightly brighter (up to $M \approx -18$) and so detectable with LSST up to 2.3 Gpc and with ZTF to 280 Mpc.

We find that for the model parameters of Model 1, a fraction $f_v^{\text{ZTF}} \approx 0.2 f_b$ of SNe-Ibc detected by ZTF might have IceCube or Km3NET neutrino detections. For SNe-Ibc detected by LSST, this fraction is $f_v^{\text{LSST}} \approx 4 \times 10^{-3} f_b$. For the other set of parameters, Model 2, we find a fraction $f_v^{\text{ZTF}} \approx 0.02 f_b$ of SNe-Ibc detected by ZTF might have IceCube or Km3NET neutrino detections. For SNe-Ibc detected by LSST, this fraction is $f_v^{\text{LSST}} \approx 4 \times 2 \times 10^{-4} f_b$. In the case of Model 3, we find a fraction $f_v^{\text{ZTF}} \approx 0.007 f_b$ of SNe-Ibc detected by ZTF might have IceCube or Km3NET neutrino detections. For SNe-Ibc detected by LSST, this fraction is $f_v^{\text{LSST}} \approx 4 \times 6.2 \times 10^{-5} f_b$

Finally, we consider hypernovae (without GRBs) that are about 7 per cent of SNe-Ibc (Guetta & Della Valle 2007). These sources are of particular interest because possibly they are the most suitable candidates to harbour choked jets. HNe are as bright as -19 , and so detectable up to 3 Gpc (to 400 Mpc with ZTF).

We find that for Model 1 a fraction $f_v^{\text{ZTF}} \approx 0.02 f_b$ of hypernovae detected by ZTF may produce neutrinos detectable by IceCube or Km3NET. For hypernovae detected by LSST, this fraction is $f_v^{\text{LSST}} \approx 7 \times 10^{-4} f_b$. For Model 2, we find that a fraction $f_v^{\text{ZTF}} \approx 0.008 f_b$ of hypernovae detected by ZTF may produce neutrinos detectable by IceCube or Km3NET. For hypernovae detected by LSST, this fraction is $f_v^{\text{LSST}} \approx 9 \times 10^{-5} f_b$. For Model 3, we find that a fraction $f_v^{\text{ZTF}} \approx 0.003 f_b$ of hypernovae detected by ZTF may produce neutrinos detectable by IceCube or Km3NET. For hypernovae detected by LSST, this fraction is $f_v^{\text{LSST}} \approx 3 \times 10^{-5} f_b$.

All these factors are smaller by a factor of about 10 for ANTARES.

The fractions of CCSNe producing neutrinos detectable with IceCube or Km3NET decrease from ZTF to LSST because the latter telescope will be able to discover an increasing number of supernovae out to very large distances and therefore hardly recognizable by neutrino detectors.

5 NEUTRINO BACKGROUND RATE

The main component for the background is the flux of atmospheric neutrinos, which is caused by the interaction of cosmic rays, high-energy protons and nuclei, with the Earth's atmosphere. Decay of charged pions and kaons produced in cosmic ray interactions generates the flux of atmospheric neutrinos and muons. Their energy spectrum is about one power steeper than the spectrum of the parent cosmic rays at Earth, due to the energy-dependent competition between meson decay and interaction in the atmosphere.

Supernovae have evolutionary time-scale of the order of dozens of days. However, we expect that a jet will be driven and neutrinos will be emitted from SNe only for a time frame comparable to the duration of long GRBs, i.e. about a minute. The time of core collapse is much more uncertain than this time frame, with characteristic uncertainties of hours–days. This means that the relevant time frame for background accumulation is hours–days. Here, we consider 1 d as a characteristic uncertainty. Given that IceCube detects about 100 000 neutrino candidates per year (Aartsen et al. 2019), and assuming 1 deg 2 directional uncertainty, the expected number of background neutrinos coincident with a given CCSN is $N_{\text{atm}} \sim 10^{-2}$.

6 HOW MANY SUPERNOVA FOLLOW-UPS WILL LEAD TO A MULTIMESSENGER DETECTION?

We first consider a search for neutrinos coincident with a population of N_{sn} detected CCSNe. We want to know how many supernovae will lead to a 3σ detection of neutrinos, assuming that a fraction f_{jet} of the supernovae produces jets and therefore high-energy neutrinos with luminosities estimated above. The expected number of background neutrino candidates detected in coincidence with the N_{sn} supernovae is $10^{-2} N_{\text{sn}}$. The expected number of signal neutrinos from these supernovae is $N_{\text{sn}} f_v$. To discover a connection at 3σ significance level, we require the expected number of signal neutrinos to be equal to 3σ uncertainty of the number of background neutrinos, i.e.

Table 1. Expected number of electromagnetically detected that need to be followed up with IceCube or KM3NeT to identify them as sources of high-energy neutrinos at 3σ significance level for different source types (column #1), considering sources electromagnetically identified by ZTF (column #2) or LSST (column #3). For models SN-Ibc and hypernova, we adopted source parameters from our Model 1. We assumed that a fraction $f_{\text{jet}} = 1$ of the sources produces jets (and therefore neutrinos) with beaming factor $f_b \approx 0.1$. For other parameters, all results in this table scale as $f_{\text{jet}}^{-2} f_b^{-2}$.

	ZTF	LSST
CCSN	40	10^5
SN-Ibc	200	6×10^5
Hypernova	2×10^4	2×10^7

the following equation needs to be satisfied.

$$N_{\text{sn}} f_v = 3 \sqrt{10^{-2} N_{\text{sn}}} \quad (3)$$

To obtain quantitative results for the different models considered here, we adopt a beaming factor $f_b \approx 0.1$ (Liang et al. 2007).

The obtained numbers of supernovae of different types required for detection at 3σ significance level are listed in Table 1. We see that, for our CCSN model with rate density $7 \times 10^{-5} \text{Mpc}^{-1} \text{yr}^{-1}$, the required number of CCSN detections can be achieved in less than a year by either ZTF or LSST if the fraction of CCSNe producing jets is ~ 1 .

7 HOW MANY NEUTRINO FOLLOW-UPS WILL LEAD TO A MULTIMESSENGER DETECTION?

Let f_{SN} denote the currently unknown fraction of astrophysical neutrinos detected by IceCube/KM3Net that were produced by SNe. Given that the density of CCSNe follows the cosmic star formation rate, we can calculate the fraction of neutrinos from SNe that come from within a distance threshold. For ZTF's 200 Mpc, this fraction is 2 per cent, while for LSST's 1500 Mpc it is 15 per cent (Bartos et al. 2017). Therefore, the fraction of detected astrophysical neutrinos that originates from SNe detectable by ZTF is $0.02 f_{\text{SN}}$, while for LSST it is $0.15 f_{\text{SN}}$. Assuming optimistically that a significant fraction of astrophysical neutrinos comes from SN jets, i.e. $f_{\text{SN}} \sim 1$, we find that about 50 (6) neutrinos need to be followed up to find a coincident CCSN with ZTF (LSST). This number is proportionally higher if not all neutrinos originate from CCSNe.

After assuming a temporal coincidence time window of 1 d and neutrino localization uncertainty of 1° , the probability of chance coincidence between a neutrino and a CCSN is 5×10^{-4} for ZTF and 0.2 for LSST. For ZTF, following up about 100 neutrinos is expected to lead to two coincident CCSNe, which would be a statistically significant deviation (3σ) from the background. For LSST, following up 80 neutrinos would lead to an expected statistically significant (3σ) excess of signal neutrinos over the background.

8 DETECTION WITH NEUTRINO MULTIPLETS

Detecting a single or even a double neutrino in coincidence with a SN is not by itself sufficient for discovery. A higher number of neutrinos coincident with a single SN would be needed to establish an astrophysical origin. Approximating the number fraction of SNe that results in the detection of three or more astrophysical neutrinos as f_v^3 , we find that the inverse of this fraction is significantly larger than the number of SNe needed for detection with a single neutrino (see Table 1). Therefore, we conclude that the follow-up of neutrino multiplets is not competitive with the follow-up of multiple supernovae with a single detected neutrino.

9 CONCLUSION

We investigated the prospects of probing jet production by a large fraction of CCSNe below the stellar envelope, as proposed by Piran et al. (2019). We calculated the expected high-energy neutrino flux from choked jets below the stellar envelope and calculated their detectability with IceCube, Km3NET, and ANTARES. We computed the number of follow-ups that need to be carried out in order to find coincident supernovae–neutrinos and obtain the fraction of supernovae that produces jets. We find that our results depend on the adopted parameters, such as Lorentz factor, burst duration, and luminosity. We have explored the set of parameters of Model 1, which we consider to be the most realistic case. Our conclusions are the following:

(i) If all CCSNe produce jets with a beaming factor of 0.1, then we need about 40 CCSNe detected by a ZTF-like telescope, or 10^5 CCSNe detected by LSST, to be able to establish neutrino emission and therefore jets from CCSNe. This can be achieved within 1 yr of observations with either ZTF or LSST. If not all CCSNe produce jets, then these numbers are proportionally higher.

(ii) Considering the electromagnetic follow-up of astrophysical neutrinos, about 100 astrophysical neutrinos need to be followed up by a ZTF-like telescope or LSST in order to establish a statistically significant excess of temporally and directionally coincident CCSNe, according to our neutrino emission model. The required number of follow-ups is greater due to the fraction of high-energy neutrinos that are not astrophysical, and if not all CCSNe drive jets.

(iii) Searching for neutrino doublet coincident with CCSNe can lead to discovery even with a single such association. We find that the search for such doublets will lead to discovery on a similar time frame as the accumulation of singlets.

(iv) The results show that the search for distant hypernovae will add a lot of noise and few neutrinos; therefore, if we include hypernovae it will take many years to make a detection. We recommend a search strategy that does not look beyond CCSN distances in order to optimize signal-to-noise ratio.

The results found in this paper are consistent with the ones already cited in this paper, Senno et al. (2016, 2017) and He et al. (2018), that estimate the diffuse neutrino flux obtained from choked jets and use the observed IceCube neutrino flux to constrain the neutrino production mechanism model. Our results are also consistent with the ones found in a recent work of Esmaili & Murase (2018), who searched for temporal and spatial coincidences between high-energy neutrino events in 6 yr of IceCube data and SNe Ib/c that occurred in the same time period. They did not find any significant correlation. However, they used their analysis to put a constraint on the total neutrino energy and therefore on the isotropic-equivalent energy of

cosmic rays. Their results are consistent with the constraints found by Senno et al. (2017), who used 1 yr sample of muon neutrinos in IceCube. Another interesting work on a similar topic is the one of Denton & Tamborra (2017), who perform a detailed analysis of the internal shock model for the production of high-energy neutrinos and use the IceCube results to constrain the model parameters. Our results on the effect of the model parameters are consistent with their conclusions.

REFERENCES

- Aartsen M. G. et al., 2014a, *Phys. Rev. Lett.*, 113, 10
 Aartsen M. G. et al., 2014b, *ApJ*, 796, 10
 Aartsen M. G. et al., 2015, *ApJ*, 807, 46
 Aartsen M. G. et al., 2019, *Phys. Rev. Lett.*, 122, 10
 Abell P. A. et al., 2009, LSST Science Book, LSST Corporation, Tucson, p. 245
 Adrian-Martinez S., Ageron M., Aharonian F., Aiello S., Albert, 2016, *Eur. Phys. J. C*, 76, 12
 Ageron M., Aguilar J. A., Al Samarai I., Albert A., Ameli F., 2011, *Nucl. Instrum. Methods Phys. Res.*, 656, 11
 Bartos I., Ahrens M., Finley C., Márka S., 2017, *Phys. Rev.*, 96, 023003
 Bellm E., Kulkarni S., 2017, *Nat. Astron.*, 1, 0071
 Berger E., Kulkarni S. R., Frail D. A., Soderberg A. M., 2003, *ApJ*, 599, 408
 Botticella M. T. et al., 2017, *A&A*, 598, A50
 Campana S. et al., 2006, *Nature*, 442, 1008
 Colgate S. A., Petschek G., 1982, *Nature*, 296, 802
 Denton P. B., Tamborra I., 2017, *Apj*, 855, 37
 Esmaili A., Murase K., 2018, *J. Cosmol. Astropart. Phys.*, 1812, 008
 Guetta D., Della Valle M., 2007, *ApJ*, 657, L73
 He H.-N., Kusenko A., Nagataki S., Fan Y.-Z., Wei D.-M., 2018, *ApJ*, 856, 119
 Hopkins A., Beacom J. F., 2011, *ApJ*, 651, 142
 Izzo L. et al., 2019, *Nature*, 565, 324
 Liang E., Zhang B., Virgili F., Dai Z. G., 2007, *ApJ*, 662, 1111
 MacFadyen A. I., Woosley S. E., Heger A., 2001, *ApJ*, 550, 410
 Madau P., Dickinson M., 2014, *ARA&A*, 52, 415
 Maeda K. et al., 2008, *Science*, 319, 1220
 Mazzali P. A. et al., 2008, *Science*, 321, 1185
 Mészáros P., Waxman E., 2001, *Phys. Rev. Lett.*, 87, 171102
 Murase K., Ioka K., 2013, *Phys. Rev. Lett.*, 111, 121102
 Murase K., Guetta D., Ahlers M., 2016, *Phys. Rev. Lett.*, 116, 071101
 Oded P., Soker N., 2011, *MNRAS*, 416, 1697
 Patat F., Barbon R., Cappellaro E., Turatto M., 1994, *A&A*, 282, 731
 Piran T., Nakar E., Mazzali P., Pian E., 2019, *ApJ*, 871, L25
 Rauch L., 2019, in EPJ Web Conf., 207, 20
 Senno N., Murase K., Mészáros P., 2016, *Phys. Rev. D*, 93, 83003
 Senno N., Murase K., Meszaros P., 2017, *J. Cosmol. Astropart. Phys.*, 1801, 025
 Soderberg A. M., Nakar E., Kulkarni S. R., Berger E., 2006, *Apj*, 638, 930

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.