

Local Properties of Riesz Minimal Energy Configurations and Equilibrium Measures

D. P. Hardin¹, A. Reznikov^{2,*}, E. B. Saff¹, and A. Volberg³

¹Center for Constructive Approximation, Department of Mathematics, Vanderbilt University, 37240, USA, ²Department of Mathematics, Florida State University, 32306, USA, and ³Department of Mathematics, Michigan State University, 48824, USA

**Correspondence to be sent to: e-mail: reznikov@math.fsu.edu*

We investigate separation properties of N -point configurations that minimize discrete Riesz s -energy on a compact set $A \subset \mathbb{R}^p$. When A is a smooth $(p-1)$ -dimensional manifold without boundary and $s \in [p-2, p-1]$, we prove that the order of separation (as $N \rightarrow \infty$) is the best possible. The same conclusions hold for the points that are a fixed positive distance from the boundary of A whenever A is any p -dimensional set. These estimates extend a result of Dahlberg for certain smooth $(p-1)$ -dimensional surfaces when $s = p-2$ (the harmonic case). Furthermore, we obtain the same separation results for "greedy" s -energy points. We deduce our results from an upper regularity property of the s -equilibrium measure (i.e., the measure that solves the continuous minimal Riesz s -energy problem), and we show that this property holds under a local smoothness assumption on the set A .

1 Introduction

In this article we study, respectively, the properties of separation and regularity for minimal discrete and for continuous Riesz energy. For a measure μ supported on a compact set A in Euclidean space and $s > 0$, its *Riesz s -potential* and *Riesz s -energy* are

Received July 28, 2017; Revised September 26, 2017; Accepted October 3, 2017

Communicated by Prof. Misha Sodin

defined by

$$U_s^\mu(x) := \int_A \frac{d\mu(y)}{|x - y|^s}, \quad I_s[\mu] := \int_A U_s^\mu(x) d\mu(x), \quad (1.1)$$

and its *Riesz log-potential* and *Riesz log-energy* by

$$U_{\log}^\mu(x) := \int_A \log \frac{1}{|x - y|} d\mu(y), \quad I_{\log}[\mu] := \int_A U_{\log}^\mu(x) d\mu(x).$$

The constant $W_s(A) := \inf I_s[\mu]$, where the infimum is taken over all probability measures μ supported on A , is called the *s-Wiener constant* of the set A , and the *s-capacity* of A is given by

$$\text{cap}_s(A) := \frac{1}{W_s(A)}, \quad s > 0, \quad \text{cap}_{\log}(A) := \exp(-W_{\log}(A)).$$

If $W_s(A) < \infty$, it is known that there exists a unique probability measure μ_s that attains $W_s(A)$ and we call μ_s the *s-equilibrium measure* for A (see [16]).

The problem of minimizing $I_s[\mu]$ has a discrete analogue. Namely, for an integer $N \geq 2$ we set

$$\mathcal{E}_s(A, N) := \min_{\omega_N \subset A} E_s(\omega_N),$$

where the infimum is taken over all N -point configurations $\omega_N = \{x_1, \dots, x_N\} \subset A$ and

$$E_s(\omega_N) := \sum_{i \neq j} \frac{1}{|x_i - x_j|^s}.$$

By $\omega_N^* = \omega_{N,s}^* = \{x_1^*, \dots, x_N^*\}$ we denote any *optimal N-point s-energy configuration*; i.e., a configuration that attains $\mathcal{E}_s(A, N)$. It is known that if $W_s(A) < \infty$, then

$$\frac{1}{N} \sum_{j=1}^N \delta_{x_j^*} \xrightarrow{*} \mu_s,$$

where δ_x denotes the unit point mass at x , and the convergence is in the weak* topology. Thus, for sets of positive *s*-capacity, by solving the discrete minimization problem, we "discretize" the measure μ_s that solves the continuous problem.

We shall study properties of ω_N^* , especially its *separation distance* given by

$$\delta(\omega_N^*) := \min_{i \neq j} |x_i^* - x_j^*|. \quad (1.2)$$

In the theory of approximation and interpolation, the separation distance is often associated with some measure of stability of the approximation. In [6] Dahlberg proved that for a $C^{1+\epsilon}$ -smooth d -dimensional manifold $A \subset \mathbb{R}^{d+1}$ without boundary and $s = d - 1$ (the harmonic case), there exists a constant $c > 0$ such that

$$\delta(\omega_N^*) \geq cN^{-1/d}, \quad \forall N \geq 2. \quad (1.3)$$

For such a set A , the order $N^{-1/d}$ for separation of N -point configurations is best possible. (More generally, this is true for any set A that is lower d -regular with respect to some finite measure μ , see Definition 2.1.) For the special case $A = \mathbb{S}^d := \{x \in \mathbb{R}^{d+1} : |x| = 1\}$, Kuijlaars, Saff, and Sun [15] extended Dahlberg's result by proving (1.3) for $s \in [d-1, d]$ and in [4], Brauchart, Dragnev, and Saff extended the range of s to $s \in (d-2, d)$ with explicit values for the constant c . Our first goal is to extend the results from [6] and [15] to all C^∞ -smooth d -dimensional manifolds for $s \in [d-1, d]$ and to interior points of d -dimensional bodies for $s \in (d-2, d)$. More generally, we show that (1.3) holds whenever the s -equilibrium measure of the manifold is upper regular (see Theorem 2.3).

Since the problem of determining the minimum $\mathcal{E}_s(A, N)$ requires solving an extremal problem in N variables, it is natural to consider a somewhat simpler discretization method, namely, the computation of *greedy s -energy points* defined below which involves minimization in only a single variable. For the logarithmic kernel on $A \times A$ where $A \subset \mathbb{C}$, such points were introduced by Edrei [10] and extensively explored by Leja [17] and his students. For general kernels they were investigated by López and Saff [18].

Definition 1.1. A sequence $\omega_\infty^* = \{a_j^*\}_{j=1}^\infty \subset A$ is called a *sequence of greedy s -energy points* if $a_1^* \in A$ and for every $N > 1$ we have

$$\sum_{j=1}^{N-1} \frac{1}{|a_N^* - a_j^*|^s} = \inf_{y \in A} \sum_{j=1}^{N-1} \frac{1}{|y - a_j^*|^s}. \quad \square$$

Notice that if $\omega_{N-1} := \{a_1^*, \dots, a_{N-1}^*\}$ is already determined, then a_N^* is chosen to minimize $E_s(\omega_{N-1} \cup \{y\})$ over all $y \in A$. It is known [18] that if $W_s(A) < \infty$ and $\omega_\infty^* = \{a_j^*\}_{j=1}^\infty$ is a sequence of greedy s -energy points, then

$$\frac{1}{N} \sum_{j=1}^N \delta_{a_j^*} \xrightarrow{*} \mu_s.$$

Some computational aspects of using the greedy s -energy points for numerical integration can be found in [11]. Our second goal, which is achieved in Theorem 2.4 and

Corollary 3.2, is to prove that for a smooth d -dimensional manifold A and $s \in (d-1, d)$ or $s > d$, there exists a constant $c > 0$ such that, for every $i < j$, we have

$$|a_i^* - a_j^*| \geq cj^{-1/d}.$$

In particular, this implies that $\delta(\{a_1^*, \dots, a_N^*\}) \geq CN^{-1/d}$. Moreover, when $s > d$ we also prove that for some constant $C > 0$ the *covering radius* η for such points satisfies

$$\eta(\{a_1^*, \dots, a_N^*\}, A) := \max_{y \in A} \min_{j=1, \dots, N} |y - a_j^*| \leq CN^{-1/d}.$$

For configurations that attain the minimal discrete energy $\mathcal{E}_s(A, N)$, this was done in [13] for $s > d$ and in [6] for $s = d - 1$.

Since the method of proof for the above results utilizes the regularity properties of the measure μ_s (see Definition 2.1), our third goal is to obtain sufficient conditions for this regularity. As we show in Theorem 2.7, compact C^∞ -smooth d -dimensional manifolds $A \subset \mathbb{R}^{d+1}$ without boundary satisfy our conditions (we anticipate, however, that the same result holds for C^2 -smooth manifolds). In the case $s = d - 1$, such a result is proved in [20]. Another result of this type was proved in [24] under an assumption that the potential U_s^μ of the measure μ satisfies an appropriate Hölder condition in the whole space \mathbb{R}^{d+1} . We derive our result, Theorem 2.7, using only smoothness of the manifold A by applying the theory of pseudo-differential operators.

The article is organized as follows. The main results in the integrable case, which include separation properties of minimal energy and greedy energy points, are stated in Section 2 and proved in Sections 5 and 7. In Section 3 we state the separation and covering properties of greedy energy points in the non-integrable case, which are proved in Subsections 5.2 and 5.3. In Section 4 we cite some known results from potential theory that we need to prove our main results, and in Section 6 we give a short introduction to the theory of pseudo-differential operators, which we need for the proof of Theorem 2.7 in Section 7.

2 Main Results in the Integrable Case

In this section, we state and discuss our main results for integrable Riesz kernels. Their proofs are given in Sections 5 and 7. We shall work primarily with a class of ℓ -regular sets, which are defined as follows.

Definition 2.1. A compact set A is called ℓ -regular, $\ell > 0$, if for some measure λ supported on A there exists a positive constant C such that for any $x \in A$ and $r < \text{diam}(A)$ we have

$$C^{-1}r^\ell \leq \lambda(B(x, r)) \leq Cr^\ell,$$

where $B(x, r)$ denotes the open ball $B(x, r) := \{y \in \mathbb{R}^p : |y - x| < r\}$. The set A is called ℓ -regular at $x \in A$ if for some positive number r_1 , the set $A \cap B(x, r_1)$ is ℓ -regular.

Further, we call a measure μ upper d -regular at x if for some constant $c(x)$ and any $r > 0$ we have

$$\mu(B(x, r)) \leq c(x)r^d. \quad (2.1)$$

□

As the next example shows, a set A can be ℓ -regular with $\ell \in \mathbb{N}$, but its s -equilibrium measure μ_s can be d -regular with $d < \ell$.

Example 2.2. For the closed unit ball $\mathbb{B}^\ell := \{x \in \mathbb{R}^\ell : |x| \leq 1\}$, which is ℓ -regular, and $s \in (\ell - 2, \ell)$ the s -equilibrium measure is given by (see, e.g., [16] or [3])

$$d\mu_s = M(1 - |x|^2)^{(s-\ell)/2} dx, \quad M = \frac{\Gamma(1 + s/2)}{\pi^{\ell/2} \Gamma(1 + (s - \ell)/2)}.$$

We notice that μ_s is ℓ -regular at every interior point of \mathbb{B}^ℓ . However, for x on the boundary $\partial\mathbb{B}^\ell = \mathbb{S}^{\ell-1}$, the measure μ_s satisfies

$$C^{-1}r^{(\ell+s)/2} \leq \mu_s(B(x, r)) \leq Cr^{(\ell+s)/2},$$

so that μ_s is not ℓ -regular at $x \in \partial\mathbb{B}^\ell$.

□

We now present our main results which include the possibility of different regularities for the set A and the measure μ_s . Although stated only for $s > 0$, they remain valid for $\ell = 1$ and $s = \log$.

Theorem 2.3. Let $A \subset \mathbb{R}^p$ be a compact set of positive s -capacity, $0 \leq p - 2 < s < d \leq \ell \leq p$, and μ_s be the s -equilibrium measure on A . Assume $A' \subset A$ is such that A is ℓ -regular at every $x \in A' \subset A$ and μ_s is upper d -regular at every $x \in A'$ with $\sup_{x \in A'} c(x) \leq c$ for some $c > 0$. Then there exists a positive constant C such that for any optimal N -point

s -energy configuration $\omega_N^* = \{x_1^*, \dots, x_N^*\}$, any $x_j^* \in A'$ and any $x_k^* \in A$ with $k \neq j$ we have

$$|x_j^* - x_k^*| > CN^{-1/d}. \quad (2.2)$$

□

In particular, (2.2) holds in the following cases (see Corollaries 2.8 and 2.9 and Example 2.2):

- $A \subset \mathbb{R}^{\ell+1}$ is a compact ℓ -regular C^∞ -smooth manifold without boundary, $s \in [\ell-1, \ell)$, and $A' = A$ with $d = \ell$;
- $A \subset \mathbb{R}^\ell$ is compact, $s \in (\ell-2, \ell)$, and $A' = \{x \in A : \text{dist}(x, \partial A) \geq \varepsilon\}$ with $\varepsilon > 0$ and $d = \ell$;
- $A = \mathbb{B}^\ell$, $s \in (\ell-2, \ell)$, and $A' = \{x \in \mathbb{R}^\ell : |x| \leq 1 - \varepsilon\}$ with $\varepsilon \in (0, 1)$ and $d = \ell$;
- $A = \mathbb{B}^\ell$, $s \in (\ell-2, \ell)$, and $A' = \partial \mathbb{B}^\ell$ with $d = (s + \ell)/2$.

Remark. In the case $\ell = 1$ and $s = \log$, our results imply the sharp estimate that when $x_j^* = \pm 1$ and $x_k^* \neq x_j^*$,

$$|x_k^* - x_j^*| \geq cN^{-2}. \quad (2.3)$$

Indeed, in this case the optimal log-energy configurations ω_N^* consist of *Fekete points*; i.e., the roots of $(1 - x^2)P'_{N-1}(x)$, where P_N is the N th degree Legendre polynomial (see e.g., [21]), for which it is known that (2.3) cannot be improved for x_k^* near ± 1 . □

The next theorem concerns greedy energy points defined in Definition 1.1.

Theorem 2.4. Let $A \subset \mathbb{R}^{\ell+1}$ be a compact C^∞ -smooth ℓ -dimensional manifold without boundary, $\ell-1 \leq s < \ell$. If $\omega_\infty^* = \{a_j^*\}_{j=1}^\infty$ is a sequence of greedy s -energy points on A , then there exists a positive constant $c(A, s)$ such that, for any $i < j$,

$$|a_i^* - a_j^*| \geq c(A, s)j^{-1/\ell}.$$

□

Theorems 2.3 and 2.4 are immediate consequences of Theorem 2.5 stated below and the following trivial observation: if $\omega_N^* = \{x_1^*, \dots, x_N^*\}$ is an optimal N -point s -energy configuration, then for any $k = 1, \dots, N$ we have

$$\sum_{j \neq k} \frac{1}{|x_k^* - x_j^*|^s} = \inf_{y \in A} \sum_{j \neq k} \frac{1}{|y - x_j^*|^s}.$$

Theorem 2.5. Let $A \subset \mathbb{R}^p$ be a compact set of positive s -capacity and μ_s be the s -equilibrium measure on A . Let $\omega_N = \{x_1, \dots, x_N\}$ be any N -point configuration in A , and $y^* \in A$ satisfy (The right-hand side of (2.4) is called the s -polarization (see, e.g., [1]) of ω_N .)

$$\sum_{j=1}^N \frac{1}{|y^* - x_j|^s} = \inf_{y \in A} \sum_{j=1}^N \frac{1}{|y - x_j|^s}. \quad (2.4)$$

If $0 \leq p - 2 < s < d \leq \ell \leq p$, A is ℓ -regular at y^* and μ_s is upper d -regular at y^* , then for every $j = 1, \dots, N$

$$|y^* - x_j| \geq (c_1 c(y^*) + 1)^{-1/s} \cdot N^{-1/d}, \quad (2.5)$$

where the constant $c(y^*)$ is from (2.1) and the positive constant c_1 depends only on A and s . \square

Our next goal is to present a sufficient condition for Theorem 2.5 to hold. We begin with the following definition.

Definition 2.6. Let $A \subset \mathbb{R}^p$ be a compact set d -regular at a point $x_0 \in A$. We say that A is (d, C^∞) -smooth at x_0 if there exists a positive number r_0 and a C^∞ -smooth invertible function $\varphi: B(x_0, r_0) \cap A \rightarrow \mathbb{R}^d$ such that $\varphi(B(x_0, r_0) \cap A)$ is open in \mathbb{R}^d and φ^{-1} is also C^∞ -smooth. \square

Our next theorem is a local result showing that if a manifold is C^∞ -smooth at a point, then the s -equilibrium measure is upper d -regular at this point.

Theorem 2.7. Let $A \subset \mathbb{R}^p$ be a compact set of positive s -capacity, where $p \in \{d, d + 1\}$ and $s \in [p - 2, d)$, and μ_s be the s -equilibrium measure on A . If A is (d, C^∞) -smooth at a point $x_0 \in A$, then μ_s is upper d -regular at x_0 ; i.e., inequality (2.1) holds for any $r > 0$. \square

Example 2.2 illustrates the sharpness of this theorem. We note that if y^* is as in (2.4) and the assumptions of Theorem 2.7 hold with x_0 replaced by y^* , then the conclusion of Theorem 2.5 follows.

The next corollary follows from Theorem 2.7 and the fact that, if $p = d$, then A is (p, C^∞) -smooth at $x_0 \in A$ if and only if x_0 is an interior point of A .

Corollary 2.8. Let $A \subset \mathbb{R}^d$ be compact, $s \in [d - 2, d)$ and x_0 be an interior point of A . If μ_s is the s -equilibrium measure on A , then μ_s is upper d -regular at x_0 . \square

Obviously, a C^∞ -smooth manifold without boundary satisfies the conditions of Theorem 2.7; therefore, we have the following consequence.

Corollary 2.9. Let $A \subset \mathbb{R}^{d+1}$ be a compact C^∞ -smooth d -dimensional manifold without boundary, $d-1 \leq s < d$ and μ_s be the s -equilibrium measure on A . Then μ_s is uniformly upper d -regular on A . \square

3 Main Results in the Non-Integrable Case

In this section, we state an analogue of Theorem 2.5 for the case $s > d$ under very weak assumptions on the set A . As a consequence, we deduce separation and covering properties of greedy energy points in this case. These properties are proved in Section 5. Below \mathcal{H}_d denotes the d -dimensional Hausdorff measure normalized by $\mathcal{H}_d([0, 1]^d) = 1$. By $\overline{\mathcal{M}}_d$ we denote the *upper d -dimensional Minkowskii content*; i.e., for a compact set $A \subset \mathbb{R}^p$, set

$$\overline{\mathcal{M}}_d(A) := \limsup_{\varepsilon \rightarrow 0^+} \frac{\mathcal{L}_p(\{x \in \mathbb{R}^p : \text{dist}(x, A) < \varepsilon\})}{\beta_{p-d} \varepsilon^{p-d}}, \quad (3.1)$$

where \mathcal{L}_p is the Lebesgue measure on \mathbb{R}^p and β_{p-d} is the volume of a $(p-d)$ -dimensional unit ball (for $p = d$, we set $\beta_0 := 1$).

Proposition 3.1. If $A \subset \mathbb{R}^p$ is a compact set with $\mathcal{H}_d(A) > 0$ ($d \leq p$) and $s > d$, then there exists a constant $c > 0$ such that for any N -point configuration $\omega_N = \{x_1, \dots, x_N\} \subset A$ and $y^* \in A$ satisfying

$$\sum_{j=1}^N \frac{1}{|y^* - x_j|^s} = \inf_{y \in A} \sum_{j=1}^N \frac{1}{|y - x_j|^s},$$

we have, for every $j = 1, \dots, N$,

$$|y^* - x_j| \geq c \cdot N^{-1/d}. \quad (3.2)$$

\square

Corollary 3.2. With the assumptions of Theorem 3.1, there exists a constant $c > 0$ such that for any sequence $\omega_\infty^* = \{a_j^*\}_{j=1}^\infty$ of greedy energy points and any $i < j$, we have

$$|a_i^* - a_j^*| \geq c j^{-1/d}. \quad (3.3)$$

If, in addition, $A \subset \tilde{A}$ for a d -regular set \tilde{A} and $\overline{\mathcal{M}}_d(A) < \infty$, then for some $c > 0$ and every $N \geq 2$, the covering radius of $\omega_N^* := \{a_1^*, \dots, a_N^*\} \subset \omega_\infty^*$ satisfies

$$\eta(\omega_N^*, A) = \max_{y \in A} \min_{j=1, \dots, N} |y - a_j^*| \leq cN^{-1/d}. \quad (3.4)$$

□

4 Some Facts from Potential Theory

For the convenience of the reader, we state several known results from potential theory that will be used in the proofs of the above formulated theorems. The following theorem can be found, for example, in [16, p. 136] or [3, Theorems 4.2.15 and 4.5.11].

Theorem 4.1. If $A \subset \mathbb{R}^p$ is a compact set of positive s -capacity, then the s -equilibrium measure μ_s is unique. Moreover, the inequality $U_s^{\mu_s}(x) \leq W_s(A)$ holds μ_s -a.e. and the inequality $U_s^{\mu_s}(x) \geq W_s(A)$ holds s -quasi-everywhere; i.e., if $F \subset \{x \in A : U_s^\mu(x) < W_s(A)\}$ is compact, then $W_s(F) = \infty$. Furthermore, if $s \in [p-2, p)$, then $U_s^{\mu_s}(x) \leq W_s(A)$ for every $x \in \mathbb{R}^p$. □

The following theorem is a special case of [19, Theorem 2.5].

Theorem 4.2. Let $s < d$ and μ be a measure supported on $A \subset \mathbb{R}^p$, where A is d -regular. If for some constant M the inequality $U_s^\mu(x) \geq M$ holds s -quasi-everywhere on A , then it holds everywhere on A . □

We conclude this section with two results from the theory of non-integrable Riesz potentials. The first result can be found in [12, Theorem 2.4] and [2, Proposition 2.5], while the second is a consequence of the proof of [13, Theorem 3].

Theorem 4.3. Assume $A \subset \mathbb{R}^p$, $\mathcal{H}_d(A) > 0$ and $s > d$. Then there exists two positive constants $c_1(s)$ and $c_2(s)$ such that for any N -point configuration $\omega_N = \{x_1, \dots, x_N\} \subset A$ we have

$$\inf_{y \in A} \sum_{j=1}^N \frac{1}{|y - x_j|^s} \leq c_1(s)N^{s/d}$$

and

$$E_s(\omega_N) = \sum_{i \neq j} \frac{1}{|x_i - x_j|^s} \geq c_2(s)\overline{\mathcal{M}}_d(A)^{-s/d}N^{1+s/d}. \quad \square$$

Theorem 4.4. Suppose the compact set $A \subset \mathbb{R}^p$ with $\mathcal{H}_d(A) > 0$ is contained in some d -regular compact set \tilde{A} and $s > d$. If $\omega_N = \{x_1, \dots, x_N\} \subset A$ is an N -point configuration with separation distance $\delta(\omega_N) \geq \tau N^{-1/d}$ for some $\tau > 0$, then for some constant $R(s, \tau, p_s)$,

$$\eta(\omega_N, A) := \max_{y \in A} \min_{j=1, \dots, N} |y - x_j| \leq R(s, \tau, p_s) N^{-1/d}, \quad (4.1)$$

where p_s is any positive constant such that

$$\inf_{y \in A} \sum_{j=1}^N \frac{1}{|y - x_j|^s} \geq p_s N^{s/d}. \quad (4.2)$$

□

5 Proofs of Theorem 2.5 and Proposition 3.1

For $x = (x(1), \dots, x(p)) \in A$, set $x_r := (x(1), \dots, x(p), r) \in \mathbb{R}^{p+1}$ and consider A as a subset of \mathbb{R}^{p+1} with $x = x_0$; i.e., $x(p+1) = 0$.

The next lemma is related to results of Carleson [5] for $s \in [d-1, d)$ and Wallin [24].

Lemma 5.1. Assume the measure μ on A is upper d -regular at $x \in A$. If $d-2 < s < d$, then there exists a constant c_1 that depends only on s and d such that

$$U_s^\mu(x_r) \geq U_s^\mu(x) - c_1 \cdot c(x) \cdot r^{d-s},$$

where $c(x)$ is a constant from (2.1). □

Proof. We first notice that for $x, y \in A$ we have $|y - x_r|^2 = |y - x|^2 + r^2$. Therefore,

$$\begin{aligned} U_s^\mu(x) - U_s^\mu(x_r) &= \int_A \frac{(|y - x|^2 + r^2)^{s/2} - |y - x|^s}{(|y - x|^2 + r^2)^{s/2} \cdot |y - x|^s} d\mu(y) \\ &= \int_{|y-x| \leq 2r} \frac{(|y - x|^2 + r^2)^{s/2} - |y - x|^s}{(|y - x|^2 + r^2)^{s/2} \cdot |y - x|^s} d\mu(y) \\ &\quad + \int_{|y-x| > 2r} \frac{(|y - x|^2 + r^2)^{s/2} - |y - x|^s}{(|y - x|^2 + r^2)^{s/2} \cdot |y - x|^s} d\mu(y) =: I_1 + I_2. \end{aligned} \quad (5.1)$$

We have

$$I_1 \leq \int_{|y-x| \leq 2r} \frac{d\mu(y)}{|y - x|^s} = \int_0^\infty \mu\{y: |y - x| \leq 2r, |y - x|^{-s} > t\} dt$$

$$\begin{aligned}
&= \int_0^{(2r)^{-s}} \mu\{y: |y-x| \leq 2r\} dt + \int_{(2r)^{-s}}^{\infty} \mu\{y: |y-x| < t^{-1/s}\} dt \\
&\leq c(x)(2r)^{d-s} + c(x) \frac{s}{d-s} (2r)^{d-s} = 2^{d-s} \cdot \frac{d}{d-s} \cdot c(x) \cdot r^{d-s} = c_2 \cdot c(x) \cdot r^{d-s},
\end{aligned} \tag{5.2}$$

where the constant c_1 depends only on s and d .

To estimate I_2 we need the following inequality. For every positive t there exists a constant c , such that for every $0 < \varepsilon < 1/4$ we have

$$(1 + \varepsilon)^t \leq 1 + c\varepsilon. \tag{5.3}$$

This estimate is trivial since the function $\varepsilon \mapsto ((1 + \varepsilon)^t - 1)/\varepsilon$ is continuous on the closed interval $[0, 1/4]$. Using (5.3) with $\varepsilon = r^2/|x-y|^2$ and $t = s/2$, we get

$$\begin{aligned}
I_2 &= \int_{|y-x|>2r} \frac{(|y-x|^2 + r^2)^{s/2} - |y-x|^s}{(|y-x|^2 + r^2)^{s/2} \cdot |y-x|^s} d\mu(y) \\
&\leq cr^2 \int_{|y-x|>2r} \frac{d\mu(y)}{|y-x|^{s+2}} \leq cr^2 \int_0^{(2r)^{-s-2}} \mu\{y: |y-x| < t^{-1/(s+2)}\} dt \\
&\leq c_3 \cdot c(x) \cdot r^2 \int_0^{(2r)^{-s-2}} t^{-d/(s+2)} dt = c_4 \cdot c(x) \cdot r^{d-s}.
\end{aligned} \tag{5.4}$$

Equality (5.1) combined with estimates (5.2) and (5.4) imply the lemma. ■

5.1 Proof of Theorem 2.5

Set

$$\gamma_N := \sum_{j=1}^N \frac{1}{|Y^* - x_j|^s} = \inf_{y \in A} \sum_{j=1}^N \frac{1}{|y - x_j|^s}.$$

Since by Theorem 4.1 we have $U_s^{\mu_s}(x) \leq W_s(A)$ for every $x \in \mathbb{R}^p$, we deduce that

$$\gamma_N \leq W_s(A)N. \tag{5.5}$$

Setting $\nu(\omega_N) := \frac{1}{N} \sum_{j=1}^N \delta_{x_j}$, we obtain for $y \in A$ that

$$U_s^{\nu(\omega_N)}(Y) \geq \frac{1}{N} \frac{\gamma_N}{W_s(A)} W_s(A) \geq \frac{1}{N} \frac{\gamma_N}{W_s(A)} U_s^{\mu_s}(Y),$$

which by the domination principle for potentials (see [14]) and Lemma 5.1 implies for $r := N^{-1/d}$ that

$$U_s^{\nu(\omega_N)}(Y_r^*) \geq \frac{1}{N} \frac{\gamma_N}{W_s(A)} U_s^{\mu_s}(Y_r^*) \geq \frac{1}{N} \frac{\gamma_N}{W_s(A)} (U_s^{\mu_s}(Y^*) - c_1 \cdot c(Y^*) N^{-1+s/d}). \quad (5.6)$$

Since A is ℓ -regular at y^* , for a small positive number r the set $A \cap \overline{B(y^*, r)}$ is ℓ -regular; applying Theorem 4.2 to this set, we obtain $U_s^{\mu_s}(Y^*) \geq W_s(A)$; thus, it follows from (5.5) and (5.6) that

$$U_s^{\nu(\omega_N)}(Y_r^*) \geq \frac{\gamma_N}{N} - c_1 \cdot c(Y^*) N^{-1+s/d},$$

or

$$\sum_{j=1}^N \frac{1}{|Y_r^* - x_j|^s} \geq \gamma_N - c_1 \cdot c(Y^*) N^{s/d}.$$

Without loss of generality, we prove (2.5) for $j = 1$. Since $|Y_r^* - x_1| \geq r = N^{-1/d}$ and $|Y_r^* - x| \geq |Y^* - x|$ for every $x \in A$, we have

$$\begin{aligned} \gamma_N - c_1 \cdot c(Y^*) N^{s/d} &\leq \sum_{j=1}^N \frac{1}{|Y_r^* - x_j|^s} = \sum_{j=2}^N \frac{1}{|Y_r^* - x_j|^s} + \frac{1}{|Y_r^* - x_1|^s} \\ &\leq \sum_{j=2}^N \frac{1}{|Y^* - x_j|^s} + N^{s/d} = \sum_{j=1}^N \frac{1}{|Y^* - x_j|^s} - \frac{1}{|Y^* - x_1|^s} + N^{s/d} \\ &= \gamma_N - \frac{1}{|Y^* - x_1|^s} + N^{s/d}. \end{aligned} \quad (5.7)$$

Therefore,

$$|Y^* - x_1| \geq (c_1 c(Y^*) + 1)^{-1/s} \cdot N^{-1/d}. \quad \blacksquare$$

5.2 Proof of Proposition 3.1

The proof is immediate. We merely observe that, by Theorem 4.3 we have for every $j = 1, \dots, N$,

$$c_1(s) N^{s/d} \geq \sum_{j=1}^N \frac{1}{|Y^* - x_j|^s} \geq |Y^* - x_j|^{-s};$$

therefore,

$$|Y^* - x_j| \geq c_1(s)^{-1/s} N^{-1/d}. \quad \blacksquare$$

5.3 Proof of Corollary 3.2

We notice that the estimate (3.3) follows from Proposition 3.1 and the fact that for every j we have

$$\sum_{i=1}^{j-1} \frac{1}{|a_j^* - a_i^*|^s} = \inf_{y \in A} \sum_{i=1}^{j-1} \frac{1}{|y - a_i^*|^s}.$$

In view of inequality (4.2) in Theorem 4.4, to deduce (3.4) it is enough to show that the inequality

$$\inf_{y \in A} \sum_{j=1}^N \frac{1}{|y - a_j^*|^s} \geq p_s N^{s/d} \quad (5.8)$$

holds for some positive constant p_s independent of N . For this purpose, observe that Theorem 4.3 implies that for some positive c that does not depend on N we have, for $\omega_N = \{a_1^*, \dots, a_N^*\}$,

$$E_s(\omega_N) \geq c N^{1+s/d}. \quad (5.9)$$

Hence, for every $j = 1, \dots, N$,

$$\sum_{i=1}^{j-1} \frac{1}{|a_j^* - a_i^*|^s} = \inf_{y \in A} \sum_{i=1}^{j-1} \frac{1}{|y - a_i^*|^s} \leq \sum_{i=1}^{j-1} \frac{1}{|a_N^* - a_i^*|^s} \leq \sum_{i=1}^{N-1} \frac{1}{|a_N^* - a_i^*|^s},$$

and so

$$E_s(\omega_N) = 2 \sum_{j=2}^N \sum_{i=1}^{j-1} \frac{1}{|a_j^* - a_i^*|^s} \leq 2N \sum_{i=1}^{N-1} \frac{1}{|a_N^* - a_i^*|^s} = 2N \inf_{y \in A} \sum_{i=1}^{N-1} \frac{1}{|y - a_i^*|^s}.$$

In view of (5.9), we get

$$\inf_{y \in A} \sum_{i=1}^{N-1} \frac{1}{|y - a_i^*|^s} \geq c_2 N^{s/d}.$$

Applying this estimate for N instead of $N - 1$, inequality (5.8) follows with $p_s = c_2$. \blacksquare

6 Some Facts From the Theory of Pseudo-Differential Operators

In order to prove Theorem 2.7 we need some facts from the theory of pseudo-differential operators. We give a brief introduction to the results we need in this section.

Let $\mathcal{S}(\mathbb{R}^d)$ be the class of Schwartz functions on \mathbb{R}^d and $\mathcal{S}'(\mathbb{R}^d)$ be the set of tempered distributions. For an open set Ω , we denote by $\mathcal{E}'(\Omega)$ the class of tempered distributions with compact support in Ω . The Fourier transform is denoted by \mathcal{F} and defined on $\mathcal{S}(\mathbb{R}^d)$ by the formula

$$\mathcal{F}(f)(\xi) := \int_{\mathbb{R}^d} f(x) e^{-2\pi i x \cdot \xi} dx, \quad f \in \mathcal{S}(\mathbb{R}^d).$$

We next introduce a class of functions (or *symbols*) that define standard pseudo-differential operators.

Definition 6.1. For a number $m \in \mathbb{R}$, we say that a function $p(x, \xi): \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ belongs to the class $S^m(\Omega)$ if $p \in C^\infty(\Omega \times \mathbb{R}^d)$ and for every compact set $K \subset \Omega$ and multi-indices α, β there exists a constant $C(K, \alpha, \beta)$ such that

$$|D_\xi^\alpha D_x^\beta p(x, \xi)| \leq C(K, \alpha, \beta) |\xi|^{m-|\alpha|}, \quad x \in \Omega, \quad |\xi| > 1, \quad (6.1)$$

where we use the notation

$$D_\xi^\alpha p(x, \xi) := \frac{\partial^{|\alpha|}}{\partial \xi^\alpha} p(x, \xi), \quad D_x^\beta p(x, \xi) := \frac{\partial^{|\beta|}}{\partial x^\beta} p(x, \xi). \quad \square$$

The Paley–Schwartz–Wiener theorem implies that if $f \in \mathcal{E}'(\mathbb{R}^d)$, then its Fourier transform $\mathcal{F}(f)$ is a function with

$$|\mathcal{F}(f)(\xi)| \leq C(1 + |\xi|)^N, \quad \xi \in \mathbb{R}^d$$

for some positive constants C and N . If p belongs to $S^m(\Omega)$ and $f \in \mathcal{E}'(\Omega)$, then, for a fixed x , we can view $p(x, \xi) \mathcal{F}(f)(\xi)$ as a tempered distribution. We define an operator P on $\mathcal{E}'(\Omega)$ by

$$P(f)(x) := \mathcal{F}^{-1}(p(x, \cdot) \mathcal{F}(f)(\cdot))(x), \quad x \in \Omega. \quad (6.2)$$

We further set

$$\Psi^m(\Omega) := \{P: p \in S^m(\Omega)\}, \quad \Psi^{-\infty}(\Omega) := \bigcap_{m \in \mathbb{R}} \Psi^m(\Omega).$$

We continue with the definition of Sobolev spaces. For every $s \in \mathbb{R}$ and $p \in (1, \infty)$ set

$$W_0^{s,p}(\Omega) := \{f \in \mathcal{E}'(\Omega) : \mathcal{F}^{-1}[(1 + |\xi|^2)^{s/2} \cdot \mathcal{F}(f)(\xi)] \in L^p(\mathbb{R}^d)\}$$

and

$$W_{loc}^{s,p} = \{f \in \mathcal{S}'(\mathbb{R}^d) : \varphi f \in W_0^{s,p}(\mathbb{R}^d) \text{ for any } \varphi \in C_0^\infty(\mathbb{R}^d)\}.$$

As with the usual Sobolev spaces (i.e., with integer s), the following embedding property holds (see, e.g., [7] or [8]).

Theorem 6.2. Assume Ω is an open set in \mathbb{R}^d with smooth boundary. If $sp > d$ and $f \in W_0^{s,p}(\Omega)$, then $f \in L^\infty(\Omega)$. \square

The following theorem about the action of pseudo-differential operators on Sobolev spaces can be found in [23, Theorem 2.1] or [22, Theorem 2.1D].

Theorem 6.3. If $P \in \Psi^m(\Omega)$ and $f \in W_0^{s,p}(\Omega)$, then $P(f) \in W_{loc}^{s-m,p}(\Omega)$. Moreover, if $P \in \Psi^{-\infty}(\Omega)$ and $f \in \mathcal{E}'(\Omega)$, then $P(f) \in C^\infty(\Omega)$. \square

We further discuss regularity properties of solutions of the equation $Pu = f$. We say that the function $p: \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ is *elliptic of order m* if $p \in S^m(\Omega)$ and for every $x \in \Omega$ there are two positive constants $c(x)$ and $r(x)$, such that

$$|p(x, \xi)| \geq c(x)|\xi|^m, \text{ for every } \xi \text{ with } |\xi| > r(x).$$

The following theorem can be found in [23, Corollary 4.3].

Theorem 6.4. Let p be an elliptic function of order m and $P \in \Psi^m(\Omega)$ be the corresponding operator defined as in (6.2). Then there exist $Q \in \Psi^{-m}(\Omega)$ and $R \in \Psi^{-\infty}(\Omega)$ such that

$$QP = I + R,$$

where I is the identity operator. \square

7 Proof of Theorem 2.7

Case $p = d$.

Since A is (d, C^∞) -smooth at x_0 , we conclude that x_0 is an interior point of A ; Theorem 4.2 applied to a small neighbourhood of x_0 implies that $U_s^{\mu_s}(x) = W_s(A)$ in $B(x_0, r_0)$ for sufficiently small r_0 . We now apply the inversion formula (see e.g., [9]): in the sense of distributions, for every $x \in B(x_0, r_0/2)$,

$$d\mu_s(x) = c_{d,s} \cdot \text{p.v.} \int_{\mathbb{R}^d} \frac{U_s^{\mu_s}(x) - U_s^{\mu_s}(y)}{|x - y|^{2d-s}} dy = c_{d,s} \cdot \text{p.v.} \int_{\mathbb{R}^d} \frac{W_s(A) - U_s^{\mu_s}(y)}{|x - y|^{2d-s}} dy.$$

Since $U_s^{\mu_s}(y) = W_s(A)$ for every $y \in B(x_0, r_0)$, we get

$$d\mu_s(x) = c_{d,s} \cdot \int_{\mathbb{R}^d \setminus B(x_0, r_0)} \frac{W_s(A) - U_s^{\mu_s}(y)}{|x - y|^{2d-s}} dy,$$

which is a bounded function when $x \in B(x_0, r_0/2)$. Therefore, the density of $d\mu_s$ with respect to the Lebesgue measure is bounded from above in the ball $B(x_0, r_0/2)$. \blacksquare

Case $p = d + 1$.

The case $s = d - 1$ is done in [20], thus we focus on the case $s > d - 1$. Since $A \subset \mathbb{R}^{d+1}$ is d -regular at x_0 and $s \in (d - 1, d)$, we obtain from Theorem 4.2 that $U_s^\mu(x) = W_s(A)$ for any $x \in A \cap B(x_0, r_1)$ for some $r_1 > 0$. Since A is C^∞ -smooth at x_0 , there exists a C^∞ -smooth map $\varphi: B(x_0, r_0) \rightarrow \tilde{B}$ such that $\tilde{B} \subset \mathbb{R}^d$ is open and $\psi := \varphi^{-1}$ is C^∞ -smooth in \tilde{B} . Without loss of generality, we assume $r_0 < r_1/2$. Set

$$d\mu^1 := \mathbb{1}_{B(x_0, r_0)} d\mu_s, \quad \mu^2 := \mu_s - \mu^1, \quad (7.1)$$

and

$$\nu := \mu^1 \circ \psi.$$

We notice that for $\tilde{x} \in \psi^{-1}(B(x_0, r_0/2))$ we have

$$U_s^{\mu^1}(\psi(\tilde{x})) = W_s(A) - U_s^{\mu^2}(\psi(\tilde{x}))$$

and the right-hand side is a smooth function. Therefore, $U^{\mu^1}(\psi(\tilde{x})) \in C^\infty(\psi^{-1}(B(x_0, r_0/2)))$.

We further write

$$U_s^{\mu^1}(\psi(\tilde{x})) = \int_{B(x_0, r_0)} \frac{d\mu^1(y)}{|y - \psi(\tilde{x})|^s} = \int_{\tilde{B}} \frac{d\nu(\tilde{y})}{|\psi(\tilde{y}) - \psi(\tilde{x})|^s}. \quad (7.2)$$

Our next goal is to write the Taylor formula for $|\psi(\tilde{y}) - \psi(\tilde{x})|^{-s}$ when \tilde{y} is in the neighbourhood of \tilde{x} . Since $\psi \in C^\infty$, there exists a C^∞ matrix $a(\tilde{x})$ and a C^∞ vector-valued function $w_1(\tilde{x}, \tilde{y})$ such that

$$\psi(\tilde{y}) - \psi(\tilde{x}) = a(\tilde{x}) \cdot (\tilde{y} - \tilde{x}) + w_1(\tilde{x}, \tilde{y})$$

and for some constant C and any component

$$|w_1(\tilde{x}, \tilde{y})| \leq C|\tilde{x} - \tilde{y}|^2, \quad \|\nabla_{\tilde{x}} w_1(\tilde{x}, \tilde{y})\|_\infty \leq C|\tilde{x} - \tilde{y}|,$$

where $\nabla_{\tilde{x}} w_1(\tilde{x}, \tilde{y})$ is the matrix of gradients of w_1 in the first variable, and $\|\cdot\|_\infty$ is the ℓ^∞ matrix norm. Therefore,

$$|\psi(\tilde{y}) - \psi(\tilde{x})|^2 = |a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^2 + w_2(\tilde{x}, \tilde{y}),$$

where w_2 is a real-valued C^∞ function with

$$|w_2(\tilde{x}, \tilde{y})| \leq C|\tilde{x} - \tilde{y}|^3, \quad |\nabla_{\tilde{x}} w_2(\tilde{x}, \tilde{y})| \leq C|\tilde{x} - \tilde{y}|^2.$$

If r_0 is small enough and $\tilde{y}, \tilde{x} \in B(x_0, r_0/2)$, then

$$\left| \frac{w_2(\tilde{x}, \tilde{y})}{|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^2} \right| \leq 1/2.$$

Consequently,

$$|\psi(\tilde{y}) - \psi(\tilde{x})|^{-s} = |a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^{-s} \cdot \left(1 + \frac{w_2(\tilde{x}, \tilde{y})}{|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^2} \right)^{-s/2}. \quad (7.3)$$

We notice that

$$w_3(\tilde{x}, \tilde{y}) := \frac{w_2(\tilde{x}, \tilde{y})}{|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^2} \in C^1$$

with $|\nabla_{\tilde{x}} w_3(\tilde{x}, \tilde{y})|$ bounded. Therefore, (7.3) implies

$$|\psi(\tilde{y}) - \psi(\tilde{x})|^{-s} = |a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^{-s} + w_4(\tilde{x}, \tilde{y}),$$

where

$$|w_4(\tilde{x}, \tilde{y})| \leq C_1 |\tilde{y} - \tilde{x}|^{-s+1}, \quad |\nabla_{\tilde{x}} w_4(\tilde{x}, \tilde{y})| \leq C_2 |\tilde{y} - \tilde{x}|^{-s}.$$

We plug this into (7.2) to get

$$U_s^{\mu_1}(\psi(\tilde{x})) = \int_{\tilde{B}} \frac{d\nu(\tilde{y})}{|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^s} + \int_{\tilde{B}} w_4(\tilde{x}, \tilde{y}) d\nu(\tilde{y}).$$

Since

$$\int_{\tilde{B}} |\nabla_{\tilde{x}} w_4(\tilde{x}, \tilde{y})| d\nu(\tilde{y}) \leq C_2 \int_{\tilde{B}} \frac{d\nu(\tilde{y})}{|\tilde{y} - \tilde{x}|^s} \leq C_3 \int_{B(x_0, r_0)} \frac{d\mu(y)}{|y - x|^s} \leq C_3 W_s(A),$$

we see that the function $\tilde{x} \mapsto \int_{\tilde{B}} w_4(\tilde{x}, \tilde{y}) d\nu(\tilde{y})$ belongs to $W^{1,\infty}(\psi^{-1}(B(x_0, r_0/4)))$. Let u be a Schwartz function equal to 1 in $\psi^{-1}(B(x_0, r_0/4))$ and to 0 outside of $\psi^{-1}(B(x_0, r_0/2))$. Then

$$u(\tilde{x}) \int_{\tilde{B}} \frac{d\nu(\tilde{y})}{|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^s} = u(\tilde{x}) U_s^{\mu_1}(\psi(\tilde{x})) - u(\tilde{x}) \int_{\tilde{B}} w_4(\tilde{x}, \tilde{y}) d\nu(\tilde{y}) =: w(\tilde{x}) \in W_0^{1,\infty}(\mathbb{R}^d). \quad (7.4)$$

We next show that the operator

$$P: v \mapsto u(\tilde{x}) \int_{\tilde{B}} \frac{d\nu(\tilde{y})}{|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^s} \quad (7.5)$$

is pseudo-differential. Namely, we use the Plancherel identity to obtain

$$\int_{\tilde{B}} \frac{d\nu(\tilde{y})}{|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^s} = \int_{\mathbb{R}^d} \mathcal{F}(v)(\xi) \overline{\mathcal{F}_{\tilde{y}}(|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^{-s})(\xi)} d\xi. \quad (7.6)$$

By definition of the Fourier Transform, we have

$$\mathcal{F}_{\tilde{y}}(|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^{-s})(\xi) = \int_{\mathbb{R}^d} |a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^{-s} e^{-2\pi i \tilde{y} \cdot \xi} d\tilde{y}.$$

Since the matrix $a(\tilde{x})$ is a $d \times (d+1)$ matrix of rank d , we observe that the set $\{a(\tilde{x}) \cdot \tilde{y} : \tilde{y} \in \mathbb{R}^d\}$ is a d -dimensional linear subspace of \mathbb{R}^{d+1} . Take a rotation R that maps this set to $\{y = (y(1), \dots, y(d+1)) \in \mathbb{R}^{d+1} : y(d+1) = 0\}$ and an operator T that maps the latter space to \mathbb{R}^d by erasing the $(d+1)$ 'st coordinate. We make a change of variables

$$\tilde{z} = T \cdot R \cdot a(\tilde{x}) \cdot (\tilde{y} - \tilde{x}).$$

By definition of T and R , we have

$$|\tilde{z}| = |T \cdot R \cdot a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})| = |a(\tilde{x})(\tilde{y} - \tilde{x})|,$$

and therefore, setting $b(\tilde{x}) := (T \cdot R \cdot a(\tilde{x}))^{-1}$, we get

$$\begin{aligned}
\mathcal{F}_{\tilde{y}}(|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^{-s})(\xi) &= \int_{\mathbb{R}^d} |a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^{-s} e^{-2\pi i \tilde{y} \cdot \xi} d\tilde{y} \\
&= e^{-2\pi i \tilde{x} \cdot \xi} \int_{\mathbb{R}^d} |\tilde{z}|^{-s} e^{-2\pi i (b(\tilde{x}) \tilde{z}) \cdot \xi} |\det(b(\tilde{x}))| d\tilde{z} \\
&= |\det(b(\tilde{x}))| e^{-2\pi i \tilde{x} \cdot \xi} \mathcal{F}(|\tilde{z}|^{-s})((b^t(\tilde{x}))\xi) \\
&= |\det(b(\tilde{x}))| e^{-2\pi i \tilde{x} \cdot \xi} |b^t(\tilde{x})\xi|^{s-d}.
\end{aligned} \tag{7.7}$$

We plug (7.7) into (7.6):

$$\int_{\tilde{B}} \frac{d\nu(\tilde{y})}{|a(\tilde{x}) \cdot (\tilde{y} - \tilde{x})|^s} = \int_{\mathbb{R}^d} \mathcal{F}(\nu)(\xi) |\det(b(\tilde{x}))| \cdot |b^t(\tilde{x})\xi|^{s-d} e^{2\pi i \tilde{x} \cdot \xi} d\xi \tag{7.8}$$

$$= \mathcal{F}^{-1} \left(\mathcal{F}(\nu)(\xi) |\det(b(\tilde{x}))| \cdot |b^t(\tilde{x})\xi|^{s-d} \right) (\tilde{x}). \tag{7.9}$$

Setting

$$p(\tilde{x}, \xi) := u(\tilde{x}) |\det(b(\tilde{x}))| \cdot |b^t(\tilde{x})\xi|^{s-d},$$

we obtain that the operator P defined in (7.5) is an elliptic pseudo-differential with symbol $p \in S^{s-d}(\tilde{B})$. We apply Theorem 6.4 to equation (7.4). Since $P\nu = w$, we get

$$\nu + R\nu = Qw, \quad R\nu \in C^\infty(\tilde{B}). \tag{7.10}$$

Further, since $w \in W_0^{1,\infty}(\tilde{B})$, we get from Theorem 6.3 that $Qw \in W_{loc}^{1+s-d,p}(\tilde{B})$ for any $p > 1$. By Theorem 6.2, we obtain that $Qw \in L^\infty(\psi^{-1}(B(x_0, r_0/4)))$, and from (7.10) we get $\nu \in L^\infty(\psi^{-1}(B(x_0, r_0/4)))$. Since the measure μ_1 defined in (7.1) is an image of ν under a smooth map ψ^{-1} , we deduce that for $r < r_0/4$

$$\mu(B(x_0, r)) = \nu(\psi^{-1}(B(x_0, r))) \leq C_1 \mathcal{H}_d(\psi^{-1}(B(x_0, r))) \leq C_2 r^d. \quad \blacksquare$$

Funding

This work was supported, in part, by National Science Foundation grant DMS-1516400 to D.P.H., A.R., E.B.S and also this work was supported, in part, by National Science Foundation grant DMS-1600065 to A.V.

Acknowledgements

The authors would like to thank the referees for carefully reading this article and providing very helpful remarks and suggestions.

References

- [1] Borodachov, S. V., D. P. Hardin, A. Reznikov, and E. B. Saff. "Optimal discrete measures for Riesz potentials." *Trans. Amer. Math. Soc.* (to appear) (2016): preprint arXiv:1606.04128.
- [2] Borodachov, S. V., D. P. Hardin, and E. B. Saff. "Asymptotics of best-packing on rectifiable sets." *Proc. Amer. Math. Soc.* 135, no. 8 (2007): 2369–80.
- [3] Borodachov, S. V., D. P. Hardin, and E. B. Saff. *Minimal Discrete Energy on Rectifiable Sets*. Springer, 2018 (to appear).
- [4] Brauchart, J. S., P. D. Dragnev, and E. B. Saff. "Riesz external field problems on the hypersphere and optimal point separation." *Potential Anal.* 41, no. 3 (2014): 647–78.
- [5] Carleson, L. "Removable singularities of continuous harmonic functions in R^m ." *Math. Scand.* 12 (1963): 15–8.
- [6] Dahlberg, B. "On the distribution of Fekete points." *Duke Math. J.* 45, no. 3 (1978): 537–42.
- [7] Demengel, F. and G. Demengel. *Functional Spaces for the Theory of Elliptic Partial Differential Equations*. Universitext. Springer, London; EDP Sciences, Les Ulis, 2012. Translated from the 2007 French original by Reinie Erné.
- [8] Di Nezza, E., G. Palatucci, and E. Valdinoci. "Hitchhiker's guide to the fractional Sobolev spaces." *Bull. Sci. Math.* 136, no. 5 (2012): 521–73.
- [9] Dunkels, A. "An inversion formula and a note on the Riesz kernel." *Ann. Inst. Fourier (Grenoble)* 26, no. 4 (1976): 197–205.
- [10] Edrei, A. "Sur les déterminants récurrents et les singularités d'une fonction donnée par son développement de Taylor." *Compositio Math.* 7 (1939): 20–88.
- [11] Elefante, G. and S. De Marchi. "Quasi-Monte Carlo integration on manifolds with mapped low-discrepancy points and greedy minimal Riesz s -energy points." Available at <http://www.math.unipd.it/~demarchi/papers/PaperDeMElef3.pdf>, 2016.
- [12] Erdélyi, T. and E. B. Saff. "Riesz polarization inequalities in higher dimensions." *J. Approx. Theory* 171 (2013): 128–47.
- [13] Hardin, D. P., E. B. Saff, and J. T. Whitehouse. "Quasi-uniformity of minimal weighted energy points on compact metric spaces." *J. Complexity* 28, no. 2 (2012): 177–91.
- [14] Itô, M. "Remarks on Ninomiya's domination principle." *Proc. Japan Acad.* 40 (1964): 743–6.
- [15] Kuijlaars, A. B. J., E. B. Saff, and X. Sun. "On separation of minimal Riesz energy points on spheres in Euclidean spaces." *J. Comput. Appl. Math.* 199, no. 1 (2007): 172–80.
- [16] Landkof, N. S. *Foundations of Modern Potential Theory. Die Grundlehren der mathematischen Wissenschaften, Band*, vol. 180. New York-Heidelberg: Springer, 1972. Translated from the Russian by A. P. Doohovskoy.
- [17] Leja, F. "Sur certaines suites liées aux ensembles plans et leur application à la représentation conforme." *Ann. Polon. Math.* 4 (1957): 8–13.

- [18] López García, A. and E. B. Saff. "Asymptotics of greedy energy points." *Math. Comp.* 79, no. 272 (2010): 2287–316.
- [19] Reznikov, A., E. B. Saff, and O. V. Vlasiuk. "A minimum principle for potentials with application to Chebyshev constants." *Potential Anal.* (to appear) (2016): preprint arXiv:1607.07283.
- [20] Sjögren, P. "Estimates of mass distributions from their potentials and energies." *Ark. Mat.* 10 (1972): 59–77.
- [21] Szegő, G. *Orthogonal Polynomials*, 4th ed. Providence, RI: American Mathematical Society, 1975. Amer. Math. Soc., Colloquium Publications, vol. XXIII.
- [22] Taylor, M. E. *Pseudodifferential Operators and Nonlinear PDE*, vol. 100. Progress in Mathematics. Boston, MA: Birkhäuser Boston, Inc., 1991.
- [23] Trèves, F. *Introduction to Pseudodifferential and Fourier Integral Operators*, vol. 1. New York-London: Plenum Press, 1980. Pseudodifferential operators, The University Series in Mathematics.
- [24] Wallin, H. "Existence and properties of Riesz potentials satisfying Lipschitz conditions." *Math. Scand.* 19 (1966): 151–60.