

## SCHUBERT CALCULUS AND TORSION EXPLOSION

GEORDIE WILLIAMSON,  
WITH A JOINT APPENDIX WITH ALEX KONTOROVICH AND PETER J. McNAMARA

*Dedicated to Meg and Gong.*

### 1. INTRODUCTION

Let  $G$  be a connected algebraic group over an algebraically closed field. A basic question in representation theory asks for the dimensions and characters of the simple rational  $G$ -modules. Structure theory of algebraic groups allows one to assume that  $G$  is reductive. If the ground field is of characteristic zero, then the theory runs parallel to the well-understood theory for compact Lie groups. In positive characteristic  $p$ , Steinberg’s tensor product theorem, the linkage principle and Jantzen’s translation principle reduce this to a question about finitely many modules which occur in the same block as the trivial module (the “principal block”). For these modules Lusztig has proposed a conjecture if  $p > h$ , where  $h$  denotes the Coxeter number of the root system of  $G$  [Lus80].<sup>1</sup> He conjectures an expression for the characters of the simple modules in terms of affine Kazhdan-Lusztig polynomials and the (known) characters of standard modules.

Lusztig’s conjecture has been shown to hold for  $p$  large (without an explicit bound) thanks to the work of Andersen, Jantzen and Soergel [AJS94], Kashiwara and Tanisaki [KT95, KT96], Kazhdan and Lusztig [KL93, KL94a, KL94b] and Lusztig [Lus94]. Alternative proofs for large  $p$  have been given by Arkhipov, Bezrukavnikov and Ginzburg [ABG04], Bezrukavnikov, Mirković and Rumynin [BMR08, BM13] (in the broader context of Lie algebra representations) and Fiebig [Fie11]. Fiebig also gives an explicit enormous<sup>2</sup> bound [Fie12] and establishes the multiplicity one case [Fie10]. For any fixed  $G$  and “reasonable”  $p$  very little is known: the case of rank 2 groups can be deduced from Jantzen’s sum formula, and intensive computational efforts have checked the conjecture for small  $p$  and certain groups, all of rank  $\leq 5$ . There is no conjecture as to what happens if  $p$  is smaller than the Coxeter number.

---

Received by the editors June 22, 2015 and, in revised form, May 5, 2016 and August 4, 2016.

2010 *Mathematics Subject Classification*. Primary 20C20, 20G05; Secondary 14N15, 14M15.

The first author of the Appendix is partially supported by an NSF CAREER grant DMS-1455705 and an Alfred P. Sloan Research Fellowship.

<sup>1</sup>Lusztig first proposed his conjecture under a restriction equivalent to  $p \geq 2h - 3$  (see [Jan08, Section 4] and [Jan03, Section 8.22] for a discussion). Kato [Kat85, Section 5] proved that if Lusztig’s conjecture holds for restricted weights, then it holds for all weights in the Jantzen region (Lusztig’s original formulation). Since Kato’s work  $p > h$  has been widely regarded as a realistic bound [Jan08, Section 4].

<sup>2</sup>For example, at least of the order of  $p \gg n^{n^2}$  for  $\mathrm{SL}_{n+1}$ .

In [Soe00] Soergel introduced a subquotient of the category of rational representations, dubbed the “subquotient around the Steinberg weight,” as a toy model for the study of Lusztig’s conjecture. Whilst the full version of Lusztig’s conjecture is based on the combinatorics of alcoves and the affine Weyl group, the subquotient around the Steinberg weight is controlled by the finite Weyl group and behaves like a modular version of category  $\mathcal{O}$ . Lusztig’s conjecture implies that the multiplicities in the subquotient around the Steinberg weight are given by finite Kazhdan-Lusztig polynomials. Thus Lusztig’s conjecture implies that “the subquotient around the Steinberg weight satisfies the Kazhdan-Lusztig conjecture.”

In [Soe00] Soergel goes on to explain how the subquotient around the Steinberg weight is controlled by Soergel bimodules. This allows him to relate this category to the category of constructible sheaves on the Langlands dual flag variety, with coefficients in the field of definition of  $G$ . Using Soergel’s results and the theory of parity sheaves [JMW14], one can see that a part of Lusztig’s conjecture for  $p > h$  is equivalent to the absence of  $p$  torsion in the stalks and costalks of integral intersection cohomology complexes of Schubert varieties in the flag variety. It has been known since the birth of the theory of intersection cohomology that 2-torsion occurs in type  $B_2$ , and 2- and 3-torsion occurs in type  $G_2$ . For over a decade no other examples of torsion were known. In 2002 Braden discovered 2-torsion in the stalks of integral intersection cohomology complexes on flag varieties of types  $D_4$  and  $A_7$  (see Braden’s appendix to [WB12]). In 2011 Polo discovered 3-torsion in the cohomology of the flag variety of type  $E_6$  and  $n$ -torsion in a flag variety of type  $A_{4n-1}$ . Polo’s (as yet unpublished) results are significant, as they emphasize how little we understand in high rank (see the final lines of [Wil12]).

In general these topological calculations appear extremely difficult. Recently Elias and the author found a presentation for the monoidal category of Soergel bimodules by generators and relations [EW], building on the work of Libedinsky [Lib10], Elias-Khovanov [EK10] and Elias [Eli16]. One of the applications of this theory is that one can decide whether a given intersection cohomology complex has  $p$ -torsion in its stalks or costalks (the bridge between intersection cohomology and Soergel bimodules is provided by the theory of parity sheaves).<sup>3</sup> The basic idea is as follows: given any pair  $(\underline{w}, x)$ , where  $x, w \in W$  and  $\underline{w}$  is a reduced expression for  $w \in W$ , one has an “intersection form,” an integral matrix. The stalks of the intersection cohomology complex corresponding to  $w$  are free of  $p$ -torsion if no elementary divisors of the intersection forms associated to all elements  $x \leq w$  are divisible by  $p$ . In principle, this gives an algorithm to decide whether Lusztig’s conjecture is correct around the Steinberg weight.<sup>4</sup> This algorithm (in a slightly different form) was discovered independently by Libedinsky [Lib15].

The generators and relations approach certainly makes calculations easier. However, this approach still has its difficulties: the diagrammatic calculations remain extremely subtle, and the “light leaves” basis in which the intersection form is calculated depends on additional choices which seem difficult to make canonical. Recent

---

<sup>3</sup>One can also perform this calculation using the theory of moment graphs [FW14]. However, the computations using generators and relations are generally much simpler.

<sup>4</sup>One can extend this to the full version of Lusztig’s conjecture by using a certain subset of the affine Weyl group, thanks to the work of Fiebig [Fie11]. Although it seems likely that the converse holds, at present one knows only one implication: the absence of  $p > h$  torsion implies the truth of Lusztig’s conjecture in characteristic  $p$ .

progress in this direction has been made by Xuhua He and the author [HW], who discovered that certain entries in the intersection form (which in some important examples are all entries) are canonical and may be evaluated in terms of expressions in the nil Hecke ring.

The main result of this paper may be seen as an example of this phenomenon. We observe that one may embed certain structure constants of Schubert calculus for  $SL_n$  as the entries of the  $1 \times 1$  intersection forms associated to pairs  $(\underline{w}, x)$  in (much) higher rank groups. In this way one can produce many new examples of torsion which grow exponentially in the rank. For example, using Schubert calculus for the flag variety of  $SL_4$  we observe that the Fibonacci number  $F_{i+1}$  occurs as torsion in  $SL_{3i+5}$ . We deduce that there is no linear function  $f(n)$  of  $n$  such that Lusztig's conjecture holds for all  $p \geq f(n)$  for  $SL_n$ . In the Appendix (by Kontorovich, McNamara and the author) we apply recent results of Bourgain-Kontorovich [BK14] in number theory to deduce that the torsion in  $SL_n$  grows exponentially in  $n$ .

Finally, there is a related conjecture due to James [Jam90] concerning the simple representations of the symmetric group in characteristic  $p$ . When combined with known results about the decomposition numbers for Hecke algebras at roots of unity, the James conjecture would yield the decomposition numbers for symmetric groups  $S_n$  in characteristic  $p > \sqrt{n}$ . In the final section of the paper we explain why our counter-examples to Lusztig's conjecture for  $SL_N$  with  $p > \binom{N}{2}$  imply that the James conjecture is incorrect for  $S_{p(\binom{N}{2})}$ . (Parts of this section were explained to me by Joe Chuang.)

**1.1. Main result.** Let  $R = \mathbb{Z}[x_1, x_2, \dots, x_n]$  be a polynomial ring in  $n$  variables. We regard  $R$  as a graded ring with  $\deg x_i = 2$  (we double degrees for reasons coming from Soergel bimodules). Let  $W = S_n$  denote the symmetric group on  $n$  letters viewed as a Coxeter group with simple reflections  $S$  consisting of the simple transpositions. Then  $W$  acts by permutation of variables on  $R$ . Let  $s_1, \dots, s_{n-1}$  denote the simple transpositions of  $S_n$ , and let  $\ell$  denote the corresponding length function. Let  $\partial_i$  denote the  $i^{\text{th}}$  divided difference operator,

$$\partial_i(f) = \frac{f - s_i f}{x_i - x_{i+1}} \in R.$$

For any element  $w \in S_n$  we obtain well-defined operators  $\partial_w = \partial_{i_1} \cdots \partial_{i_m}$  where  $w = s_{i_1} \cdots s_{i_m}$  is a reduced expression for  $w$  in the generators  $S$ .

Consider elements of the form

$$\kappa = \partial_{w_m}(x_1^{a_m} x_n^{b_m} \partial_{w_{m-1}}(x_1^{a_{m-1}} x_n^{b_{m-1}} \cdots \partial_{w_1}(x_1^{a_1} x_n^{b_1}) \cdots)),$$

where  $w_i \in S_n$  are arbitrary. We assume that  $\sum \ell(w_i) = a + b$  where  $a = \sum a_i$  and  $b = \sum b_i$  so that  $\kappa \in \mathbb{Z}$  for degree reasons. Given a subset  $I \subset \{1, \dots, n-1\}$  let  $w_I$  denote the longest element in the parabolic subgroup  $\langle s_j \rangle_{j \in I}$ . Our main theorem is the following.

**Theorem 1.1.** *Suppose that  $n \geq 1$  and  $a, b \geq 0$  are as above, and that  $\kappa \neq 0$ . Then there exists a reduced expression  $\underline{w}$  for an element of  $S_{a+n+b}$  such that the intersection form in degree zero of  $\underline{w}$  at  $w_I$ , where  $I = \{1, 2, \dots, a+n+b-1\} \setminus \{a, a+n\}$ , is the  $1 \times 1$  matrix  $((-1)^a \kappa)$ .*

The construction of the expression  $\underline{w}$  is explicit and combinatorial based on  $w_1, \dots, w_m, a_1, \dots, a_m$  and  $b_1, \dots, b_m$ . We will also see that for certain choices of  $a_i, b_i, w_i$  the prime factors of the numbers  $\kappa$  grow exponentially in  $h = n + a + b$ .

**1.2. Schubert calculus.** We explain why “Schubert calculus” occurs in the title. Consider the coinvariant ring  $C$  for the action of  $W = S_n$  on  $R$ . That is,  $C$  is equal to  $R$  modulo the ideal generated by  $W$ -invariant polynomials of positive degree. The Borel isomorphism gives a canonical identification of  $C$  with the integral cohomology of the complex flag variety of  $\mathrm{SL}_n$ .

The divided difference operators  $\partial_w$  act on  $C$ , as do elements of  $R$ . The coinvariant ring  $C$  has a graded  $\mathbb{Z}$ -basis given by the Schubert classes  $\{X_w \mid w \in S_n\}$  (normalized with  $X_{w_0} = x_1^{n-1} x_2^{n-2} \cdots x_{n-1}$  and  $X_w = \partial_{ww_0} X_{w_0}$ ). We have

$$(1.1) \quad \partial_i X_w = \begin{cases} X_{s_i w} & \text{if } s_i w < w, \\ 0 & \text{otherwise.} \end{cases}$$

The action of multiplication by  $f \in R$  of degree 2 is given as follows (the Chevalley formula):

$$(1.2) \quad f \cdot X_w = \sum_{\substack{t \in T \\ \ell(tw) = \ell(w) + 1}} \langle f, \alpha_t^\vee \rangle X_{tw}.$$

(Here  $T$  denotes the set of reflections (transpositions) in  $S_n$ , and if  $t = (i, j) \in T$  with  $i < j$ , then  $\alpha_t^\vee = \varepsilon_i - \varepsilon_j$  where  $\{\varepsilon_i\}$  is the dual basis to  $x_1, \dots, x_n$ .)

Now consider the numbers one may obtain as coefficients in the basis of Schubert classes by multiplication by  $x_1$  and  $x_n$  and by applying Demazure operators, starting with  $X_{\mathrm{id}}$ . Because  $\partial_w X_{w^{-1}} = X_{\mathrm{id}} = 1$ , any coefficient of any Schubert class that we obtain in this way can be realized as the coefficient of  $X_{\mathrm{id}}$ . Now Theorem 1.1 says that any such number occurs as torsion in  $\mathrm{SL}_{n+a+b}$  where  $a$  (resp.  $b$ ) counts the number of times that one has applied the operator of multiplication by  $x_1$  (resp.  $x_n$ ).

**1.3. Note to the reader.** This paper is entirely algebraic in that it relies only on Soergel (bi)modules, their diagrammatics and connections to representation theory (due to Soergel). Except in remarks, we neither explain nor use the relation to constructible sheaves and torsion. An alternative geometric proof of the main theorem (discovered a year after this paper was first circulated) is given in [Wil].

#### 1.4. Structure of the paper.

**Sections 2–4:** Contain background on Soergel (bi)modules and intersection forms.

**Section 5:** We prove the main theorem.

**Section 6:** We use our main theorem for  $n = 4, 5$  to give examples of torsion.

**Section 7:** We explain the connection to the Lusztig conjecture.

**Section 8:** We explain the connection to the James conjecture.

**Appendix A:** We (AK, PM and GW) prove the exponential growth of torsion.

## 2. SOERGEL BIMODULES

In the first three sections we recall what we need from the theory of Soergel (bi)modules and intersection forms. This paper is not self-contained. The main references are [Soe90, Soe92, Soe07, EK10, Eli16, EW].

Fix  $n \geq 1$  and let  $W = S_n$  denote the symmetric group on  $n$  letters. Throughout we view  $W$  as a Coxeter group with simple reflections  $S = \{(i, i+1) \mid 1 \leq i < n\}$ , and we denote by  $\ell$  its length function and  $\leq$  its Bruhat order. Let  $\mathcal{H}$  denote the Hecke algebra of  $(W, S)$  over  $\mathbb{Z}[v^{\pm 1}]$  normalized as in [Soe97]. Let  $\{H_x\}_{x \in W}$  and  $\{\underline{H}_x\}_{x \in W}$  denote its standard and Kazhdan-Lusztig bases.

Fix a field  $\mathbb{k}$  of characteristic  $p > 2$ , and let  $R = \mathbb{k}[x_1, \dots, x_n]$ . Then  $W$  acts by permutation of variables on  $R$  (graded algebra automorphisms). The reader may easily check (see, e.g., [Lib15, Lemma 7.4]) that this action is reflection faithful in the sense of [Soe07, Definition 1.5]. Given  $s \in S$  we denote by  $R^s \subset R$  the invariant subring.

Given a  $\mathbb{Z}$ -graded object (vector space, module, bimodule)  $M = \bigoplus M^i$  we let  $M(j)$  denote the shifted object:  $M(j)^i = M^{i+j}$ .

The category of *Soergel bimodules*  $\mathcal{B}$  is the full additive monoidal graded Karoubian subcategory of graded  $R$ -bimodules generated by  $B_s = R \otimes_{R^s} R(1)$  for all  $s \in S$ . In other words, the indecomposable Soergel bimodules are the shifts of the indecomposable direct summands of the *Bott-Samelson bimodules*

$$B_{\underline{w}} = B_{s_1} \otimes_R B_{s_2} \otimes_R \cdots \otimes_R B_{s_m}(m)$$

for all expressions  $\underline{w} = s_1 s_2 \cdots s_m$  in  $S$ . For any  $w \in S_n$  let  $B_w$  denote the indecomposable self-dual Soergel bimodule which occurs as a summand of  $B_{\underline{w}}$  for any reduced expression  $\underline{w}$  for  $w$  and is not isomorphic to a summand of  $B_{\underline{w}'}$  for any shorter  $\underline{w}'$ . The set  $\{B_w\}_{w \in W}$  coincides with the set of all indecomposable Soergel bimodules, up to shifts in the grading [Soe07].

*Remark 2.1.* In [Soe07] Soergel develops the theory of Soergel bimodules for a reflection faithful representation  $V$  over an infinite field of characteristic  $\neq 2$ . We have remarked above that the reflection faithful hypothesis is always satisfied. The assumption that  $\mathbb{k}$  is infinite is made in order to identify  $R$  with the polynomial functions on  $V$ . However, all the results of [Soe07] hold if one simply defines  $R$  to be the symmetric algebra on  $V^*$ , as we do. Alternatively, the reader may assume that  $\mathbb{k}$  is infinite throughout.

We denote by  $[\mathcal{B}]$  the split Grothendieck group of  $\mathcal{B}$  (i.e.,  $[B] = [B'] + [B'']$  if  $B \cong B' \oplus B''$ ). We make  $[\mathcal{B}]$  into a  $\mathbb{Z}[v^{\pm 1}]$  algebra via  $v[M] := [M(-1)]$ ,  $[B][B'] := [B \otimes_R B']$ . In [Soe07] Soergel proves that there exists a unique isomorphism of  $\mathbb{Z}[v^{\pm 1}]$ -algebras

$$\text{ch} : [\mathcal{B}] \xrightarrow{\sim} \mathcal{H}$$

such that  $\text{ch}(R(-1)) = v$  and  $\text{ch}(B_s) = \underline{H}_s$  for all  $s \in S$ .

*Remark 2.2.* Under our assumptions  $B_w$  may be realized as the equivariant intersection cohomology of the indecomposable parity sheaf [JMW14] of the Schubert variety labelled by  $w$  in the flag variety [Fie08, FW14]. In particular, if  $\mathbb{k}$  is of characteristic zero, then  $B_w$  is the equivariant intersection cohomology of a Schubert variety. In fact, the whole theory of Soergel bimodules can be seen as providing an algebraic description of the Hecke category.

Set  ${}^p\underline{H}_x := \text{ch}(B_x) \in \mathcal{H}$ . Then  $\{{}^p\underline{H}_x\}_{x \in W}$  is a basis which depends only on the characteristic  $p$  of  $\mathbb{k}$ , the  $p$ -canonical basis (see [Wil12, JW]). Let us write

$$(2.1) \quad \underline{H}_x = \sum h_{y,x} H_y, \quad {}^p\underline{H}_x = \sum {}^p h_{y,x} H_y, \quad {}^p\underline{H}_x = \sum {}^p a_{y,x} \underline{H}_y,$$

for polynomials  $h_{y,x} \in \mathbb{Z}[v]$  and  ${}^p h_{y,x}, {}^p a_{y,x} \in \mathbb{Z}[v^{\pm 1}]$ . The polynomials  $h_{y,x}$  are (normalizations of) Kazhdan-Lusztig polynomials and have non-negative coefficients. The polynomials  ${}^p h_{y,x}, {}^p a_{y,x}$  also have non-negative coefficients [JW, Proposition 4.2].

Throughout this paper we will say that  $p$  occurs as torsion in  $\text{SL}_n$  if there exists  $x \in W$  such that  ${}^p\underline{H}_x \neq \underline{H}_x$ .

### 3. SOERGEL MODULES

In this section we assume that  $p > n$ , so that the results of [Soe00] are available. Let  $R_+^W \subset R$  denote the  $W$ -invariants of positive degree,  $\langle R_+^W \rangle$  the ideal they generate and  $C = R/\langle R_+^W \rangle$  the coinvariant algebra, which inherits an (even) grading and a  $W$ -action from  $R$ . Let  $\mathcal{C}$  denote the category of *Soergel modules* consisting of all

$$D_{\underline{w}} := C \otimes_{C^{s_m}} \cdots \otimes_{C^{s_2}} C \otimes_{C^{s_1}} \mathbb{k}(m)$$

for expressions  $\underline{w} = s_1 s_2 \cdots s_m$  in  $S$ , together with their shifts, direct sums and summands inside the category of graded  $C$ -modules. (Note the order of tensor factors.)

For a reduced expression  $\underline{x}$  for  $x$  let  $D_x$  denote the unique summand of  $D_{\underline{x}}$  which does not occur as a summand of  $D_{\underline{x}'}$  for any shorter expression  $\underline{x}'$ . The set  $\{D_x \mid x \in W\}$  is well-defined and gives a set of representatives for the isomorphism classes of indecomposable Soergel modules (up to shift) [Soe00, Theorem 2.8.1].

How to go from Soergel bimodules to Soergel modules? Given a right  $R$ -module  $M$  which is killed by  $R_+^W$  the canonical map  $M \otimes_{R^s} R \rightarrow M \otimes_C C$  is an isomorphism. Hence we have an isomorphism of graded right  $C$ -modules,

$$\mathbb{k} \otimes_R R \otimes_{R^s} R \otimes_{R^t} \cdots \otimes_{R^u} R \cong \mathbb{k} \otimes_C C \otimes_{C^s} C \otimes_{C^t} \cdots \otimes_{C^u} C.$$

It follows that if we compose the functor  $M \mapsto \mathbb{k} \otimes_R M$  with the equivalence between right and left  $C$ -modules ( $C$  is commutative), we obtain a functor

$$c : \mathcal{B} \rightarrow \mathcal{C}$$

with  $c(B_{\underline{w}}) = D_{\underline{w}}$ .

**Lemma 3.1.**  $c(B_x) \cong D_x$ .

*Proof. Step 1:* We claim that the natural map provides an isomorphism,

$$(3.1) \quad \mathbb{k} \otimes_R \text{Hom}_{\mathcal{B}}^{\bullet}(B_{\underline{x}}, B_{\underline{y}}) \xrightarrow{\sim} \text{Hom}_C^{\bullet}(c(B_{\underline{x}}), c(B_{\underline{y}})).$$

(Here and in the rest of the proof,  $\text{Hom}^{\bullet}$  denotes the graded module of morphisms of all degrees.) By repeated application of the biadjunction  $(\otimes_R B_s(1), \otimes_R B_s(1))$  we may assume that  $\underline{x}$  is the empty sequence. The map  $\phi \mapsto \phi(1)$  gives a canonical identification of  $\text{Hom}_{\mathcal{B}}^{\bullet}(R, B_{\underline{y}})$  with the submodule of invariants

$$\Gamma_{\text{id}} B_{\underline{y}} := \{m \in B_{\underline{y}} \mid rm = mr \text{ for all } r \in R\}.$$

Now  $\Gamma_{\text{id}} B_{\underline{y}}$  is the first step in the filtration  $\Gamma_{\leq 0} B_{\underline{y}} \subset \Gamma_{\leq 1} B_{\underline{y}} \subset \cdots$  considered after the proof of [Soe07, Proposition 5.7], and from [Soe07, Proposition 5.9] we deduce

that the subquotients of this filtration are free as left  $R$ -modules. Thus  $\Gamma_{\text{id}} B_{\underline{y}}$  is a summand of  $B_{\underline{y}}$  as a left  $R$ -module. The injectivity of (3.1) follows.

We deduce the surjectivity of (3.1) by showing that both sides have the same (finite) dimension. If  $\underline{y} = s_1 s_2 \cdots s_m$ , let us write  $\underline{H}_{s_1} \underline{H}_{s_2} \cdots \underline{H}_{s_m} = \sum g_x H_x$  for some  $g_x \in \mathbb{Z}[v^{\pm 1}]$ . In the notation of [Soe07] we have, by [Soe07, Proposition 5.7],

$$\sum_{m \in \mathbb{Z}} (B_{\underline{y}} : \nabla_{\text{id}}[m]) v^{-m} = g_{\text{id}}$$

and  $(R : \Delta_x[m]) = \delta_{x,\text{id}} \delta_{m,0}$  (Kronecker's  $\delta$ ). Now we apply [Soe07, Theorem 5.15] to deduce that  $\text{Hom}^\bullet(R, B_{\underline{y}})$  is free of rank  $g_{\text{id}}(1)$  over  $R$ . On the other hand, if  $\langle -, - \rangle : \mathbb{Z}W \times \mathbb{Z}W \rightarrow \mathbb{Z}$  denotes the pairing with  $\langle x, y \rangle = \delta_{x,y}$ , then by [Soe00, Lemma 2.11.2], we have

$$\dim \text{Hom}_C^\bullet(\mathbb{k}, c(B_{\underline{y}})) = \langle \text{id}, \sum g_x(1)x \rangle = g_{\text{id}}(1).$$

The surjectivity follows.

*Step 2:* Because  $c(B_w) = D_w$  we can appeal to the defining properties of  $B_x$  and  $D_x$  to see that it is enough to show: if  $B$  is indecomposable, then so is  $c(B)$ . By the previous step  $\text{End}_C(c(B)) = \mathbb{k} \otimes_R \text{End}_B(B)$ , and so  $\text{End}_C(c(B))$  is a quotient of the local ring  $\text{End}_B(B)$ . Now the result follows as a non-zero quotient of a local ring is local.  $\square$

*Remark 3.2.* The above proof uses representation theory, via [Soe00, Lemma 2.11.2]. Soergel has found an algebraic proof of the above lemma, valid for any finite Coxeter group (unpublished).

Denote by  $f$  the functor of forgetting the grading on a  $C$ -module, and let  $f\mathcal{C}$  denote the essential image of  $\mathcal{C}$  under  $f$ . By [Soe00, Theorem 2.8.1], the indecomposable objects in  $f\mathcal{C}$  are precisely the  $\{fD_x\}$ . We denote by  $[f\mathcal{C}]$ ,  $[\mathcal{C}]$  the split Grothendieck groups of  $f\mathcal{C}$  and  $\mathcal{C}$ , respectively. Because  $\mathcal{C}$  is graded,  $[\mathcal{C}]$  is naturally a  $\mathbb{Z}[v^{\pm 1}]$ -module via  $v[M] := [M(-1)]$  as above. These observations, together with the above lemma, show that we have a commutative diagram

$$\begin{array}{ccccc} [f\mathcal{C}] & \xleftarrow{f} & [\mathcal{C}] & \xleftarrow[\sim]{c} & [\mathcal{B}] \\ \beta \downarrow \sim & & \downarrow \sim & = & \downarrow \text{ch} \\ \mathbb{Z}S_n & \xleftarrow{1 \leftrightarrow v} & \mathcal{H} & & \mathcal{H} \end{array}$$

where  $\beta$  is defined by

$$(3.2) \quad \beta(fD_x) = \text{ch}(B_x)_{|v=1} = {}^p \underline{H}_x|_{v=1}.$$

#### 4. INTERSECTION FORMS

Let  $\mathcal{B}$  denote the category of Soergel bimodules defined above. Given any ideal  $I \subset W$  (i.e.,  $x \leq y \in I \Rightarrow x \in I$ ) we denote by  $\mathcal{B}_I$  the ideal of  $\mathcal{B}$  generated by all morphisms which factor through a Bott-Samelson bimodule  $B_{\underline{y}}$ , where  $\underline{y}$  is a reduced expression for  $y \in I$ .

Given  $x \in W$  we denote by  $\mathcal{B}^{\geq x}$  the quotient category  $\mathcal{B}/\mathcal{B}_{\not\geq x}$  where  $\not\geq x := \{y | y \not\geq x\}$ . We write  $\text{Hom}_{\geq x}$  for (degree zero) morphisms in  $\mathcal{B}^{\geq x}$ . All Bott-Samelson bimodules  $B_{\underline{x}}$  corresponding to reduced expressions  $\underline{x}$  for  $x$  become canonically isomorphic to  $B_x$  in  $\mathcal{B}^{\geq x}$ . We have  $\text{End}_{\geq x}(B_x) = \mathbb{k}$ . Given any expression  $\underline{w}$  in  $S$  the *intersection form*<sup>5</sup> is the canonical pairing

$$I_{x,\underline{w},d}^{\mathbb{k}} : \text{Hom}_{\geq x}(B_x(d), B_{\underline{w}}) \times \text{Hom}_{\geq x}(B_{\underline{w}}, B_x(d)) \rightarrow \text{End}_{\geq x}(B_x(d)) = \mathbb{k}.$$

The following is standard (see, e.g., [JMW14, Lemma 3.1] for a similar situation).

**Lemma 4.1.** *The multiplicity of  $B_x(d)$  as a summand of  $B_{\underline{w}}$  equals the rank of  $I_{x,\underline{w},d}^{\mathbb{k}}$ .*

In the papers [EK10, EW] the category of Soergel bimodules is presented by generators and relations. More precisely, a diagrammatic category is defined by generators and relations, and it is proved that its Karoubi envelope is equivalent to Soergel bimodules, as a graded monoidal category. We will not repeat the rather complicated list of generators and relations here; see [EW, Section 1.4] or [HW, Section 2.7].

In the category  $\mathcal{D}$  the intersection form is explicit and amenable to computation: see [HW, Section 2.10] for examples. From the diagrammatic approach it is clear that  $I_{x,\underline{w},d}^{\mathbb{k}}$  is defined over  $\mathbb{Z}$ , in the sense that there exists an integral form  $I_{x,\underline{w},d}$  on a pair of free  $\mathbb{Z}$ -modules such that  $I_{x,\underline{w},d}^{\mathbb{k}} = I_{x,\underline{w},d} \otimes_{\mathbb{Z}} \mathbb{k}$  for any field  $\mathbb{k}$ .

**Corollary 4.2.** *The following are equivalent:*

- (1) *The indecomposable Soergel bimodules in characteristic  $p$  categorify the Kazhdan-Lusztig basis. That is,  ${}^p H_x = H_x$  for all  $x \in W$ .*
- (2) *For all (reduced) expressions  $\underline{w}$ , all  $x \in W$  and all  $m \in \mathbb{Z}$  the graded ranks of  $I_{x,\underline{w},m} \otimes_{\mathbb{Z}} \mathbb{Q}$  and of  $I_{x,\underline{w},m} \otimes_{\mathbb{Z}} \mathbb{k}$  agree.*
- (3) *For all reduced expressions  $\underline{w}$  and all  $x \in W$  the graded ranks of  $I_{x,\underline{w},0} \otimes_{\mathbb{Z}} \mathbb{Q}$  and of  $I_{x,\underline{w},0} \otimes_{\mathbb{Z}} \mathbb{k}$  agree.*

*Proof.* Soergel's theorem [Soe01, Lemma 5] implies that the indecomposable Soergel bimodules categorify the Kazhdan-Lusztig basis if  $\mathbb{k}$  is of characteristic zero (see [EW14] for an algebraic proof of this fact). Now Lemma 4.1 says that (2) holds if and only if  $B_{\underline{w}}$  decomposes the same way over  $\mathbb{Q}$  as it does over  $\mathbb{k}$ . Hence (1) and (2) are equivalent and (1) implies (3).

It remains to be seen whether (3) implies (1). We show the contrapositive. So assume that (1) is not satisfied, and let  $w$  be of minimal length such that  ${}^p H_w \neq H_w$ . For any  $s \in S$  with  $ws < w$ ,  $B_w$  is a summand of  $B_{ws}B_s$ . By our minimality assumption  $\text{ch}(B_{ws}) = H_{ws}$ , and hence  $\text{ch}(B_{ws}B_s) = H_{ws}H_s = \sum g_x H_x$  for some  $g_x \in \mathbb{Z}_{\geq 0}$ . Hence  $\text{ch}(B_w) = {}^p H_w = \sum {}^p a_{x,w} H_x$  with  ${}^p a_{x,w} \in \mathbb{Z}_{\geq 0}$ . By Lemma 4.1,

---

<sup>5</sup>The terminology “intersection form” comes from geometry: in de Cataldo and Migliorini's Hodge theoretic proof of the decomposition theorem, a key role is played by certain intersection forms associated to the fibers of proper maps [dCM02, dCM05]. In our setting, these intersection forms are associated to the fibers of Bott-Samelson resolutions of Schubert varieties. The relevance of these forms for the study of torsion in intersection cohomology was pointed out in [JMW14].

if  $x < w$  is such that  ${}^p a_{x,w} \neq 0$ , then the ranks of  $I_{x,\underline{w},0} \otimes_{\mathbb{Z}} \mathbb{Q}$  and  $I_{x,\underline{w},0} \otimes_{\mathbb{Z}} \mathbb{k}$  differ, for any reduced expression  $\underline{w}$  for  $w$ . Thus (3)  $\Rightarrow$  (1).  $\square$

*Remark 4.3.* The intersection form and the above proposition are some of the tools used by Fiebig to establish his bound [Fie12].

## 5. PROOF OF THE THEOREM

Let  $W$  denote the symmetric group on  $\{1, 2, \dots, a+n+b\}$  with Coxeter generators  $S = \{s_1, s_2, \dots, s_{a+n+b-1}\}$  the simple transpositions. Given a subset  $I \subset S$  let  $W_I$  denote the corresponding standard parabolic subgroup and  $w_I$  its longest element. Consider the sets

$$A = \{s_1, s_2, \dots, s_{a-1}\}, M = \{s_{a+1}, \dots, s_{a+n-1}\}, B = \{s_{a+n+1}, \dots, s_{a+n+b-1}\}.$$

Then  $W_A$  (resp.  $W_M$ , resp.  $W_B$ ) is the subgroup of permutations of  $\{1, \dots, a\}$  (resp.  $\{a+1, \dots, a+n\}$ , resp.  $\{a+n+1, \dots, a+n+b\}$ ).

We use the notation of Section 1.1 except we shift all indices by  $a$ . That is, we regard  $S_n$  as embedded in  $S_{a+n+b}$  as the standard parabolic subgroup  $W_M$ . We rename  $R = \mathbb{Z}[x_1, \dots, x_{a+n+b}]$  and write  $\alpha_i = x_i - x_{i+1}$  for the simple root corresponding to  $s_i$ . Fix

$$(5.1) \quad \kappa = \partial_{w_m}(x_{a+1}^{a_m} x_{a+n}^{b_m} \partial_{w_{m-1}}(x_{a+1}^{a_{m-1}} x_{a+n}^{b_{m-1}} \cdots \partial_{w_1}(x_{a+1}^{a_1} x_{a+n}^{b_1}) \cdots)),$$

which we assume is a non-zero integer. (Now  $w_1, \dots, w_m \in W_M$ , and the fact that  $\kappa$  is a non-zero integer implies that  $\sum \ell(w_i) = a+b$ .)

We now perform some preliminary simplifications of the right hand side of (5.1). By replacing each  $x_{a+1}^{a_i} x_{a+n}^{b_i}$  with  $x_{a+1}^{a_i} \partial_{\text{id}} x_{a+n}^{b_i}$  we may assume that for all  $i$ , either  $a_i$  or  $b_i$  is zero. Let  $M' = M \setminus \{s_{a+1}, s_{a+n-1}\}$ . If  $w \in W_{M' \cup \{s_{a+1}\}}$ , then  $\partial_w$  commutes with the operator of multiplication with  $x_{a+n}$ . Thus if  $a_i$  is zero, then we may assume that  $w_i$  is minimal in its coset  $w_i W_{M' \cup \{s_{a+1}\}}$ . Similarly, if  $b_i$  is zero, then we may assume that  $w_i$  is minimal in its coset  $w_i W_{M' \cup \{s_{a+n-1}\}}$ . From now on we assume that the right hand side of (5.1) has been simplified in this way. Finally, the minimal coset representatives of  $W_M / W_{M' \cup \{s_{a+n-1}\}}$  are the elements

$$\text{id}, s_{a+1}, s_{a+2}s_{a+1}, \dots, s_{a+n-1}s_{a+n-2} \cdots s_{a+2}s_{a+1}.$$

Similarly, the minimal coset representatives of  $W_M / W_{M' \cup \{s_{a+1}\}}$  are the elements

$$\text{id}, s_{a+n-1}, s_{a+n-2}s_{a+n-1}, \dots, s_{a+1}s_{a+2} \cdots s_{a+n-2}s_{a+n-1}.$$

Thus each  $w_i$  belongs to the first (resp. second) list if  $b_i = 0$  (resp.  $a_i = 0$ ).

Fix a reduced expression  $\underline{w}_M$  for  $w_M$  and reduced expressions  $\underline{w}_i$  for each  $w_i$ . (In fact, following the reductions of the previous paragraph each  $w_i$  has a unique reduced expression.) Let  $\underline{w}$  be the sequence

$$\underline{w} = \underline{w}_m \underline{w}_m \underline{v}_m \cdots \underline{w}_2 \underline{w}_2 \underline{v}_2 \underline{w}_1 \underline{u}_1 \underline{v}_1 \underline{w}_M,$$

where

$$\begin{aligned} \underline{u}_1 &= (s_a \cdots s_{a-a_1+1}) \cdots (s_a s_{a-1})(s_a) \\ \underline{u}_2 &= (s_a \cdots s_{a-a_1-a_2+1}) \cdots (s_a \cdots s_{a-a_1-1})(s_a \cdots s_{a-a_1}) \\ &\quad \vdots \\ \underline{u}_m &= (s_a \cdots s_1) \cdots (s_a \cdots s_{a-a_1-\cdots-a_{m-1}-1})(s_a \cdots s_{a-a_1-\cdots-a_{m-1}}) \end{aligned}$$

(subscripts fall by 1 within each set of parentheses, and  $s_a$  occurs  $a_i$  times in  $\underline{u}_i$ ) and

$$\begin{aligned}\underline{v}_1 &= (s_{a+n} \cdots s_{a+n+b_1-1}) \cdots (s_{a+n} s_{a+n+1}) (s_{a+n}) \\ \underline{v}_2 &= (s_{a+n} \cdots s_{a+n+b_1+b_2-1}) \cdots (s_{a+n} \cdots s_{a+n+b_1+1}) (s_{a+n} \cdots s_{a+n+b_1}) \\ &\quad \vdots \\ \underline{v}_m &= (s_{a+n} \cdots s_{a+n+b-1}) \cdots (s_{a+n} \cdots s_{a+n+b_1+\cdots+b_{m-1}})\end{aligned}$$


(subscripts rise by 1 within each set of parentheses, and  $s_{a+n}$  occurs  $b_i$  times in  $\underline{v}_i$ ).

**Remark 5.1.** The sequence  $\underline{u}_m \cdots \underline{u}_2 \underline{u}_1$  (resp.  $\underline{v}_m \cdots \underline{v}_2 \underline{v}_1$ ) is a reduced expression for the longest element of  $W_{A \cup \{s_a\}}$  (resp.  $W_{\{s_{a+n}\} \cup B}$ ). If we denote by  $\underline{u}'_i$  (resp.  $\underline{v}'_i$ ) the expression obtained from  $\underline{u}_i$  (resp.  $\underline{v}_i$ ) by deleting every occurrence of  $s_a$  (resp.  $s_{a+n}$ ), then  $\underline{u}'_m \cdots \underline{u}'_2 \underline{u}'_1$  (resp.  $\underline{v}'_m \cdots \underline{v}'_2 \underline{v}'_1$ ) is a reduced expression for the longest element of  $W_A$  (resp.  $W_B$ ).

**Example 5.2.** We give a real-life example. We take  $n = 4$  and consider the operator  $F : f \mapsto \partial_{23}(x_4^2(\partial_1(x_1 f)))$  on  $\mathbb{Z}[x_1, x_2, x_3, x_4]$  (we write  $\partial_{23} := \partial_2 \partial_3$ ). In the next section we will see that  $F$  is a “Fibonacci operator”; in particular,

$$\partial_1 F^3(x_1) = \partial_{123}(x_4^2 \partial_1(x_1 \partial_{23}(x_4^2 \partial_1(x_1 \partial_{23}(x_4^2 \partial_1(x_1^2)))))) = 3.$$

In the notation of Section 1.1 we have  $w_1 = w_3 = w_5 = s_1$ ,  $w_2 = w_4 = s_2 s_3$ ,  $w_6 = s_1 s_2 s_3$ ,  $a_1 = 2$ ,  $a_3 = a_5 = 1$ ,  $a_2 = a_4 = a_6 = 0$ ,  $b_1 = b_3 = b_5 = 0$  and  $b_2 = b_4 = b_6 = 2$ . Hence  $a = 4$ ,  $b = 6$  and  $a + n + b = 14$ . We can depict  $\underline{w}$  as follows:



The rest of this section will be occupied with the proof of the following theorem.

**Theorem 5.3.** *The degree zero intersection form of  $\underline{w}$  at  $w_{A \cup M \cup B}$  is the  $1 \times 1$ -matrix  $((-1)^a \kappa)$ .*

In the proof we will need the notion of subexpression and defect together with the main result of [HW]. Fix a word  $\underline{y} = s_{i_1} \cdots s_{i_m}$  in  $S$ . A *subexpression* of  $\underline{y}$  is a sequence  $\underline{e} = e_1 \cdots e_m$  with  $e_i \in \{0, 1\}$  for all  $i$ . We set  $\underline{y}^{\underline{e}} := s_{i_1}^{e_1} \cdots s_{i_m}^{e_m} \in \bar{W}$ .

Any subexpression  $\underline{e}$  determines a sequence  $y_0, y_1, \dots, y_m \in W$  via  $y_0 := \text{id}$ ,  $y_j := s_{i_{m+1-j}}^{e_{m+1-j}} y_{j-1}$  for  $1 \leq i \leq m$  (so  $y_m = \underline{y}^{\underline{e}}$ ). Given a subexpression  $\underline{e}$  we associate a sequence  $d_j \in \{U, D\}$  (for *Up*, *Down*) via

$$d_j := \begin{cases} U & \text{if } s_{i_j} y_{m-j} > y_{m-j}, \\ D & \text{if } s_{i_j} y_{m-j} < y_{m-j}. \end{cases}$$

Usually we view  $\underline{e}$  as the decorated sequence  $(d_1 e_1, \dots, d_m e_m)$ . The *defect* of  $\underline{e}$  is

$$\text{df}(\underline{e}) := |\{i \mid d_i e_i = U0\}| - |\{i \mid d_i e_i = D0\}|.$$

*Remark 5.4.* See [EW, Section 2.4] for examples and motivation. We warn the reader that in this paper we work from right to left to define the defect, rather than from left to right as in [EW, Section 2.4] and [HW, Section 2.3]. This change of conventions is necessary to have the operators  $\partial_i$  act on polynomials on the left. One may easily pass between the two possible choices via the symmetry on Soergel bimodules which interchanges left and right actions. (In the diagrammatic language of [EW] this corresponds to flipping diagrams about the  $y$ -axis.)

Recall that the nil Hecke ring  $NH$  is defined to be the algebra generated by  $R$  and symbols  $\delta_i$  for each  $s_i \in S$ , and subject to the relation  $\delta_i^2 = 0$  for all  $s_i \in S$ , the braid relations and the nil Hecke relation

$$\delta_i f = s_i(f) \delta_i + \partial_i(f) \quad \text{for all } s_i \in S \text{ and } f \in R.$$

As left  $R$ -modules  $NH$  is free with basis  $\{\delta_w\}_{w \in W}$ , where  $\delta_w := \delta_{i_1} \cdots \delta_{i_k}$  for any reduced expression  $w = s_{i_1} \cdots s_{i_k}$ . The grading on  $R$  extends to a grading on  $NH$  with  $\deg \delta_w = -2\ell(w)$  for all  $w \in W$ .

Equipped with this notation we can now give the proof.

**Lemma 5.5.**  $\underline{w}$  is reduced.

*Proof.* Let us fix an element  $x \in W_Q$  where  $Q = \{s_p, s_{p+1}, \dots, s_{q-1}, s_q\}$  (for some  $p, q$  with  $1 < p \leq q < a+n+b-1$ ) and a reduced expression  $\underline{x}$  for  $x$ . Then for any  $j$  the expressions

$$s_j s_{j-1} \cdots s_{p-1} \underline{x} \quad \text{and} \quad s_j s_{j+1} \cdots s_{q+1} \underline{x}$$

are reduced. (For example, one can write a formula for how the displayed elements act on  $1, 2, \dots, a+n+b$  in terms of  $x$ , and verify that their lengths differ from  $\ell(x)$  by  $j-p+2$  (resp.  $q-j+2$ ) by counting inversions. It follows that the lengths of the displayed expressions agree with the lengths of the underlying elements, and thus they are reduced.)

From the definition of  $\underline{w}$  it follows that there exists a sequence of expressions  $\emptyset = \underline{x}_0, \underline{x}_1, \dots, \underline{x}_r = \underline{w}$  such that each  $\underline{x}_i$  is obtained from  $\underline{x}_{i-1}$  by the procedure of the previous paragraph. Thus  $\underline{w}$  is reduced as claimed.  $\square$

Write  $\underline{w} = s_{i_1} \cdots s_{i_l}$ .

**Lemma 5.6.** Any subexpression  $\underline{e}$  of  $\underline{w}$  with  $\underline{w}^{\underline{e}} = w_{A \cup M \cup B}$  has  $e_j = 0$  if  $s_{i_j} \in \{s_a, s_{a+n}\}$  and  $e_j = 1$  if  $s_{i_j} \in A \cup B$ .

*Proof.* Let  $\underline{e}$  denote a subexpression of  $\underline{w}$  with  $\underline{w}^{\underline{e}} = w_{A \cup M \cup B}$ .

Any expression  $\underline{y}$  for  $w_A$  contains a subsequence of the form  $s_{a-1} s_{a-2} \cdots s_1$  (think about what happens to  $1 \in \{1, \dots, a+n+b\}$ ). In  $\underline{w}$ ,  $s_1$  occurs only once. To the left of  $s_1$  there is only one occurrence of  $s_{a-1}, s_{a-2}$ , etc. We conclude that

the restriction of  $\underline{e}$  to  $(s_a s_{a-1} \cdots s_2 s_1)$  in  $\underline{u}_i$  is equal to  $(01 \cdots 11)$ , where  $i$  is the maximal index with  $\underline{u}_i \neq \emptyset$ . Now any expression for  $w_A$  starting in  $s_{a-1} \cdots s_1$  has to contain a subsequence to the right of the form  $s_{a-1} \cdots s_2$  (think about what happens to  $2 \in \{1, \dots, a+n+b\}$ ). Continuing in this way we see that the restriction of  $\underline{e}$  to each  $\underline{u}_j$  has the form

$$(01 \cdots 1) \cdots (01 \cdots 1)(01 \cdots 1)$$

(with the same bracketing as in the definition of each  $\underline{u}_i$ ). Similar arguments apply to each  $\underline{v}_i$ , and the result follows.  $\square$

**Lemma 5.7.** *There is a unique subexpression  $\underline{e}$  of  $\underline{w}$  such that  $\underline{w}^{\underline{e}} = w_{A \cup M \cup B}$  and  $\underline{e}$  has defect zero.*

*Proof.* By the previous lemma we must have  $e_j = 0$  (resp. 1) if  $s_{i_j} \in \{s_a, s_{a+n}\}$  (resp.  $s_{i_j} \in A \cup B$ ). Because each  $e_j$  with  $s_{i_j} \in \{s_a, s_{a+n}\}$  is  $U0$  and because  $W_{A \cup B}$  and  $W_M$  commute, we only have to understand subexpressions  $\underline{e}'$  of

$$\underline{w}' = \underline{w}_m \underline{w}_{m-1} \cdots \underline{w}_1 \underline{w}_M$$

of defect  $-(a+b) = -\sum_{i=1}^m \ell(w_i)$  such that  $(\underline{w}')^{\underline{e}'} = w_M$ . Now  $\ell(\underline{w}') = \ell(w_M) + a+b$ , and hence any subexpression  $\underline{e}'$  of  $\underline{w}'$  with  $(\underline{w}')^{\underline{e}'} = w_M$  has at most  $a+b$  zeroes. Moreover, if  $\underline{e}'$  has defect  $-a-b$ , then  $\underline{e}'$  must have exactly  $a+b$  zeroes, all of which have to be  $D0$ . Now, using that  $\underline{w}_M$  is reduced, the only subexpression of  $\underline{w}'$  fulfilling these requirements is

$$(0 \cdots 0)(0 \cdots 0) \cdots (0 \cdots 0)(1 \cdots 1). \quad \square$$

*Proof of Theorem 5.3.* We conclude from the previous two lemmas and their proofs that the unique defect zero subexpression  $\underline{e}$  of  $\underline{w}$  with  $\underline{w}^{\underline{e}} = w_A w_B w_M$  is

$$\underline{e} = \underline{e}_m \underline{f}_m \underline{g}_m \cdots \underline{e}_2 \underline{f}_2 \underline{g}_2 \underline{e}_1 \underline{f}_1 \underline{g}_1 \underline{e}_0,$$

where  $\underline{e}_0$  (resp.  $\underline{e}_i$ ) is a subexpression of  $\underline{w}_M$  (resp.  $\underline{w}_i$ ) given by

$$\underline{e}_0 = (U1, U1, \dots, U1) \quad (\text{resp. } \underline{e}_i = (D0, D0, \dots, D0))$$

and  $\underline{f}_i$  (resp.  $\underline{g}_i$ ) is a subexpression of  $\underline{u}_i$  (resp.  $\underline{v}_i$ ) given by

$$(U0, U1, \dots, U1)(U0, U1, \dots, U1) \cdots (U0, U1, \dots, U1).$$

(we use the same bracketing as in the definition of  $\underline{u}_i$  and  $\underline{v}_i$ ).

Hence the intersection form of  $\underline{w}$  at  $w_{A \cup M \cup B}$  for degree  $d = 0$  is indeed a  $1 \times 1$  matrix. Applying [HW, Theorem 5.1] its unique entry is given by the coefficient of  $\delta_{w_{A \cup M \cup B}} = \delta_{w_A} \delta_{w_M} \delta_{w_B}$  in<sup>6</sup>

$$E := E_m F_m G_m \cdots E_2 F_2 G_2 E_1 F_1 G_1 E_0,$$

<sup>6</sup>Actually, as noted in Remark 5.4, here we use a “right to left” convention, rather than the “left to right” convention of [HW]. One can check that [HW, Theorem 5.1] holds in either convention. Alternatively one can proceed as follows. Let  $\underline{w}^r = s_{i_\ell} \cdots s_{i_1}$  denote the reversed sequence, and let  $\iota : NH \rightarrow NH$  denote the anti-involution with  $\iota(f) = f$  for  $f \in R$  and  $\iota(\delta_x) = \delta_{x^{-1}}$  for  $x \in W$ . Then [HW, Theorem 5.1] implies that the intersection form of  $\underline{w}^r$  at  $w_{A \cup M \cup B}$  is the  $1 \times 1$  matrix given by the coefficient of  $\delta_{w_A} \delta_{w_M} \delta_{w_B}$  in  $\iota(E)$ . This implies the statement because the intersection forms of  $\underline{w}$  and  $\underline{w}^r$  at  $w_{A \cup M \cup B}$  agree.

where  $E_0 = \delta_{w_M}$ ,  $E_i = \delta_{w_i}$  for  $1 \leq i \leq m$  and the  $F_i, G_i$  are given by

$$F_1 = (\alpha_a \delta_{a-1} \cdots \delta_{a-a_1+1}) \cdots (\alpha_a \delta_{a-1})(\alpha_a)$$

$$F_2 = (\alpha_a \delta_{a-1} \cdots \delta_{a-a_1-a_2+1}) \cdots (\alpha_a \delta_{a-1} \cdots \delta_{a-a_1-1})(\alpha_a \delta_{a-1} \cdots \delta_{a-a_1})$$

⋮

$$F_m = (\alpha_a \delta_{a-1} \cdots \delta_1) \cdots (\alpha_a \delta_{a-1} \cdots \delta_{a-a_1-\cdots-a_{m-1}-1})(\alpha_a \delta_{a-1} \cdots \delta_{a-a_1-\cdots-a_{m-1}})$$

$$G_1 = (\alpha_{a+n} \delta_{a+n+1} \cdots \delta_{a+n+b_1-1}) \cdots (\alpha_{a+n} \delta_{a+n+1})(\alpha_{a+n})$$

⋮

$$G_m = (\alpha_{a+n} \delta_{a+n+1} \cdots \delta_{a+n+b-1}) \cdots (\alpha_{a+n} \delta_{a+n+1} \cdots \delta_{a+n+b_1+\cdots+b_{m-1}}).$$

In  $NH$  we can write  $E = \sum_{y \in W_{A \cup M \cup B}} f_y \delta_y$ . After noting that

$$\deg E = 2(-\ell(w_A) - \ell(w_B) + a + b - \sum \ell(w_i) - \ell(w_M)) = -2\ell(w_{A \cup M \cup B}),$$

we see that, in fact,  $E = \kappa' \delta_{w_{A \cup M \cup B}}$  for some  $\kappa' \in \mathbb{Z}$ . In particular, whenever we apply a nil Hecke relation  $f \delta_i = \delta_i s_i(f) + \partial_i(f)$  with  $s_i \in A \cup B$  to reduce  $E$ , the term involving  $\partial_i(f)$  does not contribute. (It would lead to a term which is zero for degree reasons.) Hence we can write

$$E = \delta_{w_A} \delta_{w_B} (\delta_{w_m} \gamma_m \delta_m) \cdots (\delta_{w_2} \gamma_2 \delta_2) (\delta_{w_1} \gamma_1 \delta_1) \delta_{w_M},$$

where each  $\gamma_i$  (resp.  $\delta_i$ ) is a product of  $a_i$  (resp.  $b_i$ ) roots of the form  $x_k - x_{a+1}$  with  $k < a+1$  (resp.  $x_{a+n} - x_k$  for  $k > a+n$ ). Hence we have

$$\begin{aligned} E &= \delta_{w_A w_B} (\delta_{w_m} (-x_{a+1})^{a_m} x_{a+n}^{b_m}) \cdots (\delta_{w_2} (-x_{a+1})^{a_2} x_{a+n}^{b_2}) (\delta_{w_1} (-x_{a+1})^{a_1} x_{a+n}^{b_1}) \delta_{w_M} \\ &= (-1)^a \kappa \cdot \delta_{w_{A \cup M \cup B}}, \end{aligned}$$

where the first (resp. second) equality follows from Lemma 5.8 (resp. 5.9) below. The theorem follows.  $\square$

**Lemma 5.8.** *Let  $w_1, \dots, w_m \in W_M$  and  $\zeta_1, \dots, \zeta_m \in R$ . Assume that for some  $1 \leq i \leq m$  we can write  $\zeta_i = \zeta_i^M \zeta'_i$  for some  $W_M$ -invariant  $\zeta_i^M$  of positive degree, and that  $\sum \deg \zeta_i = \sum \ell(w_i)$ . Then  $\delta_{w_m} \zeta_m \cdots \delta_{w_1} \zeta_1 \delta_{w_M} = 0$ .*

*Proof.* We have  $\delta_{w_m} \zeta_m \cdots \delta_{w_i} \zeta'_i \cdots \delta_{w_1} \zeta_1 \delta_{w_M} \in \bigoplus_{y \in W_M} R \delta_y$ , and hence

$$\delta_{w_m} \zeta_m \cdots \delta_{w_i} \zeta'_i \cdots \delta_{w_1} \zeta_1 \delta_{w_M} = 0$$

because it is of degree  $< -2\ell(w_M)$ . As  $\zeta_i^M$  is  $W_M$ -invariant,

$$0 = \zeta_i^M (\delta_{w_m} \zeta_m \cdots \delta_{w_i} \zeta'_i \cdots \delta_{w_1} \zeta_1 \delta_{w_M}) = \delta_{w_m} \zeta_m \cdots \delta_{w_1} \zeta_1 \delta_{w_M}. \quad \square$$

**Lemma 5.9.** *With  $w_i, a_i, b_i, \kappa$  as above we have*

$$(\delta_{w_m} x_{a+1}^{a_m} x_{a+n}^{b_m}) (\delta_{w_{m-1}} x_{a+1}^{a_{m-1}} x_{a+n}^{b_{m-1}}) \cdots (\delta_{w_1} x_{a+1}^{a_1} x_{a+n}^{b_1}) \delta_{w_M} = \kappa \cdot \delta_{w_M}.$$

*Proof.* It is well known that  $\delta_i \mapsto \partial_i, f \mapsto (f \cdot)$  makes  $R$  into an  $NH$ -module. In  $NH$  we can write

$$(\delta_{w_m} x_{a+1}^{a_m} x_{a+n}^{b_m}) (\delta_{w_{m-1}} x_{a+1}^{a_{m-1}} x_{a+n}^{b_{m-1}}) \cdots (\delta_{w_1} x_{a+1}^{a_1} x_{a+n}^{b_1}) = K + \sum_{\text{id} \neq w \in W_M} f_w \delta_w,$$

where  $K \in \mathbb{Z}$  for degree reasons. By applying this identity to  $1 \in R$  we deduce that  $K = \kappa$ . The lemma now follows because if  $w \in W_M$ , then  $\delta_w \delta_{w_M} = 0$  unless  $w = \text{id}$ .  $\square$

## 6. (COUNTER)-EXAMPLES

We use the notation of Section 1.2 and write  $\partial_{12} := \partial_1 \partial_2$ ,  $X_1 := X_{s_1}$ ,  $X_{12} := X_{s_1 s_2}$ , etc.

6.1.  $n < 4$ : One checks easily using (1.1) and (1.2) that for  $n = 2, 3$  one can only obtain  $\kappa = \pm 1$ .

6.2.  $n = 4$ : Using (1.1) and (1.2) we see that in  $C$  we have

$$(6.1) \quad X_1 = x_1 \quad \text{and} \quad X_3 = -x_4.$$

Consider the (degree zero) operator  $F : C \rightarrow C$  given by

$$F : h \mapsto \partial_{23}(x_4^2(\partial_1(x_1 h))).$$

Using (1.1) and (1.2) one checks that  $F$  preserves the submodule  $\mathbb{Z}X_1 \oplus \mathbb{Z}X_3$  and in the basis  $X_1, X_3$  is given by

$$F = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}.$$

This matrix determines the Fibonacci recursion. Hence for  $i \geq 1$  we have

$$F^i(x_1) = F_{i+1}X_1 + F_iX_3,$$

where  $F_1 = 1$ ,  $F_2 = 1$ ,  $F_3 = 2$ ,  $F_4 = 3$ , etc., denote the Fibonacci numbers. In particular,

$$\partial_1(F^i(x_1)) = F_{i+1}.$$

We conclude from the main theorem that any prime dividing the Fibonacci number  $F_{i+1}$  occurs as torsion in  $\mathrm{SL}_{3i+5}$ . By Carmichael's theorem [Car14] the first  $n \gg 1$  Fibonacci numbers have at least  $n$  distinct prime factors. By the prime number theorem we conclude that the torsion in  $\mathrm{SL}_n$  grows at least as fast as some constant times  $n \log n$ . Hence no linear bound is sufficient for Lusztig's conjecture.

It is a well-known conjecture that infinitely many Fibonacci numbers are prime. By the above results, this conjecture would immediately imply that the torsion in  $\mathrm{SL}_n$  grows exponentially in  $n$ . Unfortunately, little seems to be known about the rate of growth of prime factors of Fibonacci numbers.

In the Appendix we work with different operators in order to establish exponential growth of torsion. If  $U_l$  (resp.  $U_u$ ) denotes the operator  $h \mapsto \partial_{21}(x_1^2(\partial_1(x_1 h)))$  (resp.  $h \mapsto \partial_{23}(x_4^2(\partial_3(x_4 h))))$ , then  $U_l$  and  $U_u$  preserve the submodule  $\mathbb{Z}X_1 \oplus \mathbb{Z}X_3$  and in the basis  $X_1, X_3$  are given by

$$U_l = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad U_u = \begin{pmatrix} -1 & -1 \\ 0 & -1 \end{pmatrix}.$$

It follows from our main theorem that any prime dividing any matrix coefficient of any word of length  $\ell$  in the generators  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$  occurs as torsion in  $\mathrm{SL}_{3\ell+5}$ . Indeed, given any word  $\omega_1 \omega_2 \cdots \omega_r$  in the operators  $U_l$  and  $U_u$  we may obtain all four coefficients (up to sign) of the corresponding product of the matrices  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$  as  $\partial_i(\omega_1(\cdots(\omega_r(x_j))\cdots))$  for  $i \in \{1, 3\}$  and  $j \in \{1, 4\}$  (use (6.1) and the fact that  $\partial_1(X_1) = 1 = \partial_3(X_3)$ ).

6.3.  $n = 5$ : In the following table we list some examples of  $p$  torsion in  $\mathrm{SL}_N$  found using  $n = 5$ . The entries in the list were found by random computer searches,

$N$	14	17	20	22	25	30	40	50	70	100
$p$	3	7	13	23	53	197	2237	34183	4060219	470858183

(These entries were found as follows. Consider the eight degree zero operators:

$$\partial_{4321}x_1^4, \partial_{321}x_1^3, \partial_{21}x_1^2, \partial_1x_1, \partial_{1234}x_5^4, \partial_{234}x_5^3, \partial_{34}x_5^2, \partial_4x_5.$$

It is not difficult to calculate the matrices of these operators acting on any homogeneous component of  $C$  in the Schubert basis. The above entries were obtained as prime factors of coefficients obtained by repeated application of these operators to  $x_1^3$  and  $x_1^2x_5 \in C^6$ .)

## 7. LUSZTIG CONJECTURE

This section consists of connections and complements to [Soe00], with which we assume the reader is familiar. In keeping with the setting of this paper, we work with  $G = \mathrm{SL}_n$  throughout; however, analogous statements are true (with the same proofs) for any connected reductive group.

As in Section 3 we assume in this and the following section that  $p > n$ . Let  $\mathcal{O}$  denote the “regular subquotient around the Steinberg weight” as defined in [Soe00, Section 2.3]. We denote by  $\Delta(x), P(x)$  the standard and projective objects in  $\mathcal{O}$  and by  $\theta_s : \mathcal{O} \rightarrow \mathcal{O}$  for  $s \in S$  the translation functor [Soe00, Section 2.5]. Let  $[\mathcal{O}]$  denote the Grothendieck group of  $\mathcal{O}$  and

$$\alpha : [\mathcal{O}] \xrightarrow{\sim} \mathbb{Z}[W]$$

the isomorphism with  $\alpha([\Delta(x)]) = x$  for all  $x \in W$  ( $\alpha$  is denoted  $A$  in [Soe00, Section 2.10]). As observed in [Soe00], Lusztig’s conjecture implies that

$$\alpha(P(x)) = \underline{H}_{x|v=1} \quad \text{for all } x \in W.$$

*Remark 7.1.* This observation should be compared with a much earlier theorem of Jantzen [Jan79, Anhang, Corollar] matching multiplicities of simple modules in Weyl modules in sufficiently large characteristic  $p$  and multiplicities of simple modules in Verma modules in characteristic 0. This observation, together with Jantzen’s calculations in rank 2, were the main ingredients that led to the formulation of the Lusztig conjecture.

**Proposition 7.2.** *We have  $\alpha(P(x)) = {}^p\underline{H}_{x|v=1}$ . In particular, if  ${}^p\underline{H}_x \neq \underline{H}_x$  with  $p > n$ , then Lusztig’s conjecture fails for  $\mathrm{SL}_n$  in characteristic  $p$ .*

*Remark 7.3.* Recall that  ${}^p a_{y,x}, {}^p h_{y,x} \in \mathbb{Z}_{\geq 0}[v^{\pm 1}]$  (see Section 2). In particular,

$${}^p \underline{H}_x = \underline{H}_x \Leftrightarrow {}^p \underline{H}_{x|v=1} = \underline{H}_{x|v=1}.$$

*Proof.* Let  $p\mathcal{O}$  denote the full subcategory of projective objects in  $\mathcal{O}$ , and  $[p\mathcal{O}]$  its split Grothendieck group. Because  $\mathcal{O}$  has a finite homological dimension, the map  $[p\mathcal{O}] \rightarrow [\mathcal{O}]$  induced by the inclusion is an isomorphism. Recall the commutative

diagram

$$\begin{array}{ccc} [p\mathcal{O}] & \xrightarrow{\mathbb{V}} & [f\mathcal{C}] \\ \alpha \downarrow & & \beta' \downarrow \\ \mathbb{Z}S_n & = & \mathbb{Z}S_n \end{array}$$

from [Soe00, Section 2.11] (see Section 3 for the definition of  $f\mathcal{C}$ ).

We claim that  $\beta'$  above agrees with the  $\beta$  defined in Section 3. If  $\underline{w} = st \cdots u$  we have, by [Soe00, Theorem 2.6.2],

$$\mathbb{V}(\theta_u \cdots \theta_t \theta_s M_{\text{id}}) \cong fD_{\underline{w}}.$$

Thus, by [Soe00, Sections 2.5 and 2.10],

$$\beta'([fD_{\underline{w}}]) = \alpha(\theta_u \cdots \theta_t \theta_s M_{\text{id}}) = (1+s)(1+t) \cdots (1+u).$$

By the commutativity of the diagram in Section 3,

$$\beta([fD_{\underline{w}}]) = \text{ch}([B_s B_t \cdots B_u])_{v=1} = (\underline{H}_s \underline{H}_t \cdots \underline{H}_u)_{v=1} = (1+s)(1+t) \cdots (1+u).$$

Hence  $\beta = \beta'$  as claimed, as  $[f\mathcal{C}]$  is generated by  $[fD_{\underline{w}}]$  over all expressions  $\underline{w}$ .

Now we are done: by [Soe00, Theorem 2.8.2] we have  $\text{VP}(x) = fD_x$ , and the proposition follows from (3.2).  $\square$

## 8. JAMES CONJECTURE

In this section we explain why the results of the previous section yield counter-examples to the James conjecture [Jam90] on the decomposition numbers of Schur algebras and the symmetric group.

Fix positive integers  $N$  and  $r$ . Let  $\Lambda^+(N, r)$  denote the set of partitions of  $r$  into at most  $N$  parts; that is, sequences  $(\lambda_1, \dots, \lambda_N)$  such that  $\lambda_1 \geq \cdots \geq \lambda_N \geq 0$  with  $r = \sum \lambda_i$ . Then  $\Lambda^+(N, r)$  is a partially ordered set with respect to the dominance order  $\leq$ .

Let  $S(N, r)$  denote the Schur algebra over  $\mathbb{Z}$  (see, e.g., [Gre81]). Its category of representations is equivalent to the category of polynomial representations of the group scheme  $GL_N$  of fixed degree  $r$ . Fix a field  $\mathbb{k}$  of characteristic  $p > N$ , and let  $S_{\mathbb{k}}(N, r)$  denote the Schur algebra over  $\mathbb{k}$ . The category  $\text{Rep } S_{\mathbb{k}}(N, r)$  of finitely generated  $S_{\mathbb{k}}(N, r)$ -modules is a highest weight category with simple modules indexed by  $\Lambda^+(N, r)$ . Given  $\lambda \in \Lambda^+(N, r)$  we denote by  $L(\lambda)$  (resp.  $\Delta(\lambda)$ ,  $\nabla(\lambda)$ ,  $P(\lambda)$ ,  $T(\lambda)$ ) the simple (resp. standard, costandard, indecomposable projective, indecomposable tilting) module indexed by  $\lambda$ .

Let  $S_q(N, r)$  denote the  $q$ -Schur algebra and  $S_{\varepsilon}(N, r)$  its specialization at a fixed primitive  $p^{\text{th}}$ -root of unity  $\varepsilon \in \mathbb{C}$  (see, e.g., [Don98]). Then the category of finitely generated  $S_{\varepsilon}(N, r)$ -modules is highest weight. As above we write  $L_{\varepsilon}(\lambda)$  (resp.  $\Delta_{\varepsilon}(\lambda)$ , etc.) for the simple (resp. standard, etc.) module corresponding to  $\lambda \in \Lambda^+(N, r)$ . Given a module  $M$  for  $S_{\varepsilon}(N, r)$  we may choose a stable  $\mathbb{Z}[\varepsilon]$ -lattice and reduce to obtain a module over  $S_{\mathbb{k}}(N, r)$ . In this way we obtain the decomposition map on Grothendieck groups,

$$d : [\text{Rep } S_{\varepsilon}(N, r)] \rightarrow [\text{Rep } S_{\mathbb{k}}(N, r)].$$

One has  $d([\nabla_\varepsilon(\lambda)]) = [\nabla(\lambda)]$ . The James conjecture [Jam90] predicts that

$$(8.1) \quad d([L_\varepsilon(\lambda)]) = [L(\lambda)]$$

if  $p > \sqrt{r}$ .<sup>7</sup>

Let  $\rho = (N-1, \dots, 1, 0) \in \Lambda^+(N, \binom{N}{2})$  and let  $\text{st} := (p-1)\rho$  denote the “Steinberg weight.” Let  $S_N$  denote the symmetric group of  $N$  letters, acting by permutation on  $\mathbb{Z}^N$ . Lusztig’s quantum character formula gives (see (2.1) for notation)

$$(8.2) \quad [\nabla_\varepsilon(\text{st} + x\rho) : L_\varepsilon(\text{st} + y\rho)] = h_{x,y}(1)$$

as modules for  $S_\varepsilon(N, p\binom{N}{2})$ . (Actually, Lusztig’s quantum character formula gives the multiplicity for the quantum group of  $\mathfrak{sl}_N$  specialized at  $\varepsilon$  in terms of an affine Kazhdan-Lusztig polynomial. The translation of his formula to yield the above multiplicity is standard but technical. Alternatively, one can appeal to [AJS94] and the  $p \gg 0$  version of (8.3) below.)

Similarly, [Soe00, Theorem 1.2.2] gives (again see (2.1) for notation)

$$(8.3) \quad [\nabla(\text{st} + x\rho) : L(\text{st} + y\rho)] = {}^p h_{x,y}(1)$$

as modules for  $S_{\mathbb{k}}(N, p\binom{N}{2})$ . (Actually, Soergel’s result gives this multiplicity for rational modules for  $\text{SL}_N(\mathbb{k})$ . The translation to  $GL_N(\mathbb{k})$  and hence to modules for the Schur algebra is standard.)

We conclude that whenever  ${}^p \underline{H}_x \neq \underline{H}_x$  for some  $x$  the characters of the simple modules for  $S_\varepsilon(N, p\binom{N}{2})$  and  $S_{\mathbb{k}}(N, p\binom{N}{2})$  are different, because the simple and costandard modules both give bases for the Grothendieck group. In particular, there exists  $\lambda$  such that  $d([L_\varepsilon(\lambda)]) \neq [L(\lambda)]$ . Hence any  $p$  appearing on the table in Section 6 with  $p > \binom{N}{2}$  contradicts the James conjecture for  $S(N, p\binom{N}{2})$ .

*Remark 8.1.* A straightforward computation in  $[\text{Rep } S_\varepsilon(N, r)]$  and  $[\text{Rep } S_{\mathbb{k}}(N, r)]$  shows that

$$d([L_\varepsilon(\text{st} + x\rho)]) = \sum a_{x,y}(1) [L(\text{st} + y\rho)],$$

and so the  $a_{x,y}$  evaluated at 1 give part of James’ “adjustment matrix.”

However, amongst weights of the form  $\text{st} + x\rho$  for  $x \in S_N$  only  $\text{st} + w_0\rho$  is  $p$ -restricted ( $w_0$  denotes the longest element of  $S_N$ ). Hence the above non-trivial decomposition numbers are invisible to the symmetric group, as all simple modules corresponding to non- $p$ -restricted weights are killed by the Schur functor.

To get counter-examples in the symmetric group we can use the Ringel self-duality of the Schur algebra and modular category  $\mathcal{O}$ . (Parts of the following argument are due to Joe Chuang.) Given any  $N' \geq N$  we have an obvious embedding  $\Lambda^+(N, r) \hookrightarrow \Lambda^+(N', r)$  obtained by appending 0’s to the partition. There is a quotient functor  $f : \text{Rep } S(N', r) \rightarrow \text{Rep } S(N, r)$  which preserves simple, standard, costandard modules and indecomposable tilting modules corresponding to  $\lambda$  in  $\Lambda^+(N, r) \subset \Lambda^+(N', r)$  (see [Gre81, Section 6.5] and [Don98, Section A4.5]).

---

<sup>7</sup>A stronger version requires that  $p$  be larger than the weight of  $\lambda$ . It reduces to the condition  $p > \sqrt{r}$  for the principal block, which will be the only case considered below.

Now consider a variant of the subquotient around the Steinberg weight discussed in the previous section. Consider the Serre subquotient  $\mathcal{O} := \mathcal{A}/\mathcal{N}$  of  $\text{Rep } S_{\mathbb{k}}(N, p\binom{N}{2})$  where  $\mathcal{A}$  is the Serre subcategory generated by simple modules  $L(\lambda)$  such that  $\lambda \leq st + \rho$  and  $\lambda$  lies in the same block as  $st + \rho$ , and  $\mathcal{N}$  denotes the Serre subcategory generated by those simples  $L(\lambda) \in \mathcal{A}$  which are not of the form  $L(st + x\rho)$  for some  $x \in S_N$ . The definition of  $\mathcal{O}_\varepsilon$  is obtained by replacing  $L(\lambda)$  by  $L_\varepsilon(\lambda)$  in the definition of  $\mathcal{O}$ .

Let us denote the images of  $L(st + x\rho)$ ,  $\Delta(st + x\rho)$ , etc., in  $\mathcal{O}$  as  $L(x)$ ,  $\Delta(x)$ , etc., and similarly for  $\mathcal{O}_\varepsilon$ . Then  $\mathcal{O}$  is the highest weight category with simple, standard, etc., and tilting objects  $L(x)$ ,  $\Delta(x)$ , etc., and similarly for  $\mathcal{O}_\varepsilon$ . It is known that both categories are Ringel self-dual. (The proof of this fact seems not to be explicit in the literature. However, a proof may be obtained by adapting ideas of [BBM04]. One shows that one has a braid group action on  $D^b(\mathcal{O})$  (resp.  $D^b(\mathcal{O}_\varepsilon)$ ) and a lift of the longest element interchanges a projective and tilting generator.) Applying Bernstein-Gelfand-Gelfand (BGG) reciprocity and then Ringel self-duality for  $\mathcal{O}$  and  $\mathcal{O}_\varepsilon$  we obtain

$$h_{x,y}(1) = [\nabla_\varepsilon(x) : L_\varepsilon(y)] = (P_\varepsilon(y) : \Delta_\varepsilon(x)) = (T_\varepsilon(yw_0) : \nabla_\varepsilon(xw_0)),$$

$${}^p h_{x,y}(1) = [\nabla(x) : L(y)] = (P(y) : \Delta(x)) = (T(yw_0) : \nabla(xw_0)).$$

Applying Ringel self-duality of  $S(p\binom{N}{2}, p\binom{N}{2})$  [Don93] and of  $S_\varepsilon(p\binom{N}{2}, p\binom{N}{2})$  [Don98] we have

$$h_{x,y}(1) = (T_\varepsilon(st + yw_0\rho) : \nabla_\varepsilon(st + xw_0\rho)) = (P_\varepsilon((st + yw_0\rho)') : \Delta_\varepsilon((st + xw_0\rho)')),$$

$${}^p h_{x,y}(1) = (T(st + yw_0\rho) : \nabla(st + xw_0\rho)) = (P((st + yw_0\rho)') : \Delta((st + xw_0\rho)')).$$

Finally, again by BGG reciprocity

$$h_{x,y}(1) = [\nabla_\varepsilon((st + xw_0\rho)') : L_\varepsilon((st + yw_0\rho)'),]$$

$${}^p h_{x,y}(1) = [\nabla((st + xw_0\rho)') : L((st + yw_0\rho)')].$$

The partitions  $(st + xw_0\rho)'$  and  $(st + yw_0\rho)'$  are  $p$ -restricted. Hence, after applying the Schur functor, the first (resp. second) number can be interpreted as a decomposition number for the Hecke algebra specialized at  $\varepsilon$  (resp. the symmetric group in characteristic  $p$ ). It follows that the results of the previous section also produce counter-examples for the symmetric group.

*Remark 8.2.* Consulting the table of counter-examples in Section 6 we see that the smallest counter-example produced by the above methods occurs in  $S_{N'}$  with

$$N' = p\binom{N}{2} = 2237\binom{40}{2} = 1\,744\,860.$$

The size of this number is a relic of our method (in particular, the fact that we cannot say anything about  $p$ -restricted weights). It is an important question as to where the first counter-examples occur.

## APPENDIX A. EXPONENTIAL GROWTH OF TORSION

A.1. **Statement of the theorem.** Let

$$(A.1) \quad \Gamma := \left\langle \left( \begin{array}{cc} 1 & 1 \\ 0 & 1 \end{array} \right), \left( \begin{array}{cc} 1 & 0 \\ 1 & 1 \end{array} \right) \right\rangle^+$$

be the sub-semi-group of  $\mathrm{SL}(2, \mathbb{Z})$  generated (freely) by the matrices displayed. For a matrix  $\gamma \in \Gamma$ , let  $\ell(\gamma)$  be its word length in the generators of  $\Gamma$ . In the main body of the paper, the third author proves that any prime  $p$  dividing any entry  $\gamma_{ij}$  of any matrix

$$\gamma = \left( \begin{array}{cc} \gamma_{11} & \gamma_{12} \\ \gamma_{21} & \gamma_{22} \end{array} \right) \in \Gamma$$

occurs as torsion in  $\mathrm{SL}_{3\ell+5}$ , where  $\ell = \ell(\gamma)$  is the word length (see Section 6.2). The purpose of this Appendix is to show the existence of exponentially large (relative to word length) prime divisors of matrix coefficients in  $\Gamma$ , thus giving exponentially large counter-examples to the expected bounds in Lusztig's conjecture.

In fact, the stated purpose can be accomplished by an almost<sup>8</sup> direct application of the affine sieve [BGS06, BGS10, SGS13]; see also, e.g., [Kon14]. It turns out that one can do much more using recent progress on “local-global” problems in “thin orbits” (see, e.g., the discussion in [Kon13]); namely, one can produce not just prime divisors but actual primes in the entries of  $\Gamma$  and moreover give explicit estimates for their exponential growth rates (which are far superior compared to those which would come from an affine sieve analysis). Our main result is the following

**Theorem A.1.** *There are absolute constants  $\tau > 0$  and  $c > 1$  so that, for all  $L$  large, there exists  $\gamma \in \Gamma$  of word length  $\ell(\gamma) \leq L$  and top-left entry  $\gamma_{11} = p$  prime with  $p > \tau c^L$ . In fact, there are many primes arising this way,*

$$(A.2) \quad \# \{p > \tau c^L : \exists \gamma \in \Gamma \text{ with } \ell(\gamma) \leq L \text{ and } \gamma_{11} = p\} \gg \frac{c^L}{L}.$$

*The implied constant above is absolute and effective.*

Throughout this Appendix,  $p$  always denotes a prime. The notation  $f(L) \gg g(L)$  means that  $g = O(f)$ ; i.e.,  $|g(L)| \leq M|f(L)|$  for a fixed  $M > 0$  and all large  $L$ . In this case,  $M$  is the *implied constant* referred to above.

Exact estimates for  $\tau$  and  $c$  can readily be determined; the value coming from our proof is  $c = (\frac{1+\sqrt{5}}{2})^{1/5} \approx 1.101\dots$ , and we can take  $\tau = 5/7$ ; see (A.7). It turns out that Theorem A.1 is a nearly immediate consequence of recent advances toward Zaremba's conjecture on continued fractions with bounded partial quotients.

Given  $A \geq 1$ , let  $\Gamma_A$  be the sub-semi-group,

$$(A.3) \quad \Gamma_A := \left\langle \left( \begin{array}{cc} a & 1 \\ 1 & 0 \end{array} \right) \cdot \left( \begin{array}{cc} b & 1 \\ 1 & 0 \end{array} \right) : 1 \leq a, b \leq A \right\rangle^+.$$

(In fact,  $\Gamma_A$  is freely generated by the displayed elements.)

---

<sup>8</sup>For the application to be immediate,  $\Gamma$  would need to be a Zariski-dense *group* and not just a semi-group; minor modifications are needed to handle this case.

**Theorem A.2** (Bourgain-Kontorovich [BK14]). *There exists  $A_0$  and an absolute constant  $c < \infty$  so that, for  $A \geq A_0$  and all  $N$  large,*

$$\#\{n \leq N : \exists \gamma \in \Gamma_A \text{ with } \gamma_{11} = n\} = N \left(1 + O\left(e^{-c\sqrt{\log N}}\right)\right),$$

where the implied constant and  $c > 0$  are both absolute.

That is, almost all *integers* (not just primes) arise as top-left entries in the semigroup  $\Gamma_A$ . Bourgain-Kontorovich give  $A_0 = 50$  as an allowable value for  $A$ , and this has since been reduced to  $A_0 = 5$  [FK14, Hua15]; furthermore, Hensley [Hen96] has conjectured that  $A_0 = 2$  is allowable and that the error rate  $O(e^{-c\sqrt{\log N}})$  can be replaced by  $O(1/N)$ . What is most important to our application is that the error rate is asymptotically  $o(1/\log N)$ . This, together with the prime number theorem, has the following immediate corollary.

**Corollary A.3.** *Let notation be as above, and set  $A = 5$ . Then for any fixed constant  $\theta < 1$ ,*

$$(A.4) \quad \#\{p \in (\theta N, N] : \exists \gamma \in \Gamma_A \text{ with } \gamma_{11} = p\} = (1 - \theta) \frac{N}{\log N} \left(1 + o(1)\right),$$

as  $N \rightarrow \infty$ .

Equipped with this estimate, it is a simple matter to give the following proof.

**A.2. Proof of Theorem A.1.** Fix constants  $c > 1$  and  $\tau > 0$  to be chosen later, and let  $\mathcal{S}_1$  denote the set of primes on the left hand side of (A.2),

$$\mathcal{S}_1 := \{p > \tau c^L : \exists \gamma \in \Gamma \text{ with } \ell(\gamma) \leq L \text{ and } \gamma_{11} = p\}.$$

We seek a lower bound on the cardinality of  $\mathcal{S}_1$ .

For a parameter  $A$  (which we will soon set to  $A = 5$ ) and a matrix  $\gamma \in \Gamma_A$ , let  $\ell_A(\gamma)$  denote the word length in the generators of  $\Gamma_A$  given in (A.3). We make the pleasant observation that

$$\begin{pmatrix} a & 1 \\ 1 & 0 \end{pmatrix} \cdot \begin{pmatrix} b & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 \\ b & 1 \end{pmatrix},$$

and hence  $\Gamma_A$  is a sub-semi-group of  $\Gamma$ . Moreover, if  $\gamma \in \Gamma_A \subset \Gamma$ , then the word lengths in the two semi-groups are related by

$$\ell(\gamma) \leq 2A \cdot \ell_A(\gamma),$$

since each generator in  $\Gamma_A$  has word length at most  $2A$  in the generators of  $\Gamma$ . We decrease  $\mathcal{S}_1$  to a smaller set  $\mathcal{S}_2 \subset \mathcal{S}_1$  of primes coming from the top-left entries of  $\Gamma_A$  instead of  $\Gamma$ ,

$$\mathcal{S}_2 := \{p > \tau c^L : \exists \gamma \in \Gamma_A \text{ with } \ell_A(\gamma) \leq L/(2A) \text{ and } \gamma_{11} = p\}.$$

Next we define the archimedean sup-norm

$$\|\gamma\|_\infty := \max(\gamma_{ij}),$$

which for  $\gamma \in \Gamma_A$  is easily seen to be the top-left entry

$$(A.5) \quad \|\gamma\|_\infty = \gamma_{11}.$$

Let

$$\varphi := \frac{1 + \sqrt{5}}{2} \quad \text{and} \quad \bar{\varphi} := \frac{1 - \sqrt{5}}{2}$$

denote the eigenvalues of  $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$ .

For any

$$\gamma = \prod_{i=1}^n \left( \begin{pmatrix} a_i & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} b_i & 1 \\ 1 & 0 \end{pmatrix} \right) \in \Gamma_A,$$

we have

$$\|\gamma\|_\infty = (1 \ 0) \gamma \begin{pmatrix} 1 \\ 0 \end{pmatrix} \geq (1 \ 0) \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^{2n} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = F_{2n+1},$$

where  $F_m$  is the  $m$ th Fibonacci number. Because  $F_{2n+1} = (\varphi^{2n+1} - \bar{\varphi}^{2n+1})/\sqrt{5}$ , if we set  $d := \varphi/\sqrt{5}$ , then for all  $\gamma \in \Gamma_A$ ,

$$\|\gamma\|_\infty \geq d \cdot \varphi^{2\ell_A(\gamma)}.$$

That is, the logarithm of the archimedean norm is controlled (up to a constant) by the word length. Define the “archimedean” parameter  $N$  (with respect to  $L$ ) by

$$(A.6) \quad N := d \cdot \varphi^{L/A}.$$

Replacing the word length condition  $\ell_A(\gamma) \leq L/(2A)$  in  $\mathcal{S}_2$  by the stronger restriction that  $\|\gamma\|_\infty \leq N$  decreases  $\mathcal{S}_2$  to a subset  $\mathcal{S}_3$  defined by

$$\mathcal{S}_3 := \{p > \tau c^L : \exists \gamma \in \Gamma_A \text{ with } \|\gamma\|_\infty \leq N \text{ and } \gamma_{11} = p\}.$$

Since  $\gamma_{11} = p = \|\gamma\|_\infty$ , the condition  $\|\gamma\|_\infty \leq N$  can be replaced by  $p \leq N$ ; hence

$$\mathcal{S}_3 = \{\tau c^L < p \leq N : \exists \gamma \in \Gamma_A \text{ with } \gamma_{11} = p\}.$$

Make the choice

$$(A.7) \quad c = \varphi^{1/A},$$

which is  $(\frac{1+\sqrt{5}}{2})^{1/5} \approx 1.101\dots$  when  $A = 5$ . Then for any  $\theta < 1$ , take  $\tau = \theta d$ . With these choices of parameters, we see that

$$\mathcal{S}_3 = \{\theta N < p \leq N : \exists \gamma \in \Gamma_A \text{ with } \gamma_{11} = p\}.$$

Now we are done: combining the above with (A.6) and (A.4) gives

$$\#\mathcal{S}_1 \geq \#\mathcal{S}_3 \gg \frac{N}{\log N} \gg \frac{c^L}{L},$$

as claimed in (A.2). This completes the proof of Theorem A.1.

#### ACKNOWLEDGEMENTS

The ideas of this paper crystallized after long discussions with Xuhua He. The author thanks him warmly for the invitation to Hong Kong and the many interesting discussions that resulted from the visit. The author also thanks Ben Elias for countless hours (often productive, always enjoyable) getting to know Soergel bimodules. His influence is omnipresent in this paper.

The author thanks Joe Chuang for useful correspondence and explaining how to get counter-examples in the symmetric group. Finally, the author thanks Henning Haahr Andersen, Ben Elias, Peter Fiebig, Anthony Henderson, Daniel Juteau, Nicolas Libedinsky, Kaneda Masaharu and especially Patrick Polo and the referees for valuable comments. These results were announced in June 2013 at ICRT VI in Zhangjiajie, China.

## REFERENCES

- [ABG04] Sergey Arkhipov, Roman Bezrukavnikov, and Victor Ginzburg, *Quantum groups, the loop Grassmannian, and the Springer resolution*, J. Amer. Math. Soc. **17** (2004), no. 3, 595–678, DOI 10.1090/S0894-0347-04-00454-0. MR2053952
- [AJS94] H. H. Andersen, J. C. Jantzen, and W. Soergel, *Representations of quantum groups at a  $p$ th root of unity and of semisimple groups in characteristic  $p$ : independence of  $p$*  (English, with English and French summaries), Astérisque **220** (1994), 321. MR1272539
- [BBM04] A. Beilinson, R. Bezrukavnikov, and I. Mirković, *tilting exercises* (English, with English and Russian summaries), Mosc. Math. J. **4** (2004), no. 3, 547–557, 782. MR2119139
- [BGS06] Jean Bourgain, Alex Gamburd, and Peter Sarnak, *Sieving and expanders* (English, with English and French summaries), C. R. Math. Acad. Sci. Paris **343** (2006), no. 3, 155–159, DOI 10.1016/j.crma.2006.05.023. MR2246331
- [BGS10] Jean Bourgain, Alex Gamburd, and Peter Sarnak, *Affine linear sieve, expanders, and sum-product*, Invent. Math. **179** (2010), no. 3, 559–644, DOI 10.1007/s00222-009-0225-3. MR2587341
- [BK14] Jean Bourgain and Alex Kontorovich, *On Zaremba’s conjecture*, Ann. of Math. (2) **180** (2014), no. 1, 137–196, DOI 10.4007/annals.2014.180.1.3. MR3194813
- [BM13] Roman Bezrukavnikov and Ivan Mirković, *Representations of semisimple Lie algebras in prime characteristic and the noncommutative Springer resolution*, Ann. of Math. (2) **178** (2013), no. 3, 835–919, DOI 10.4007/annals.2013.178.3.2. MR3092472
- [BMR08] Roman Bezrukavnikov, Ivan Mirković, and Dmitriy Rumynin, *Localization of modules for a semisimple Lie algebra in prime characteristic*, Ann. of Math. (2) **167** (2008), no. 3, 945–991, DOI 10.4007/annals.2008.167.945. With an appendix by Bezrukavnikov and Simon Riche. MR2415389
- [Car14] R. D. Carmichael, *On the numerical factors of the arithmetic forms  $\alpha^n \pm \beta^n$* , Ann. of Math. (2) **15** (1913/14), no. 1–4, 49–70, DOI 10.2307/1967798. MR1502459
- [dCM02] Mark Andrea A. de Cataldo and Luca Migliorini, *The hard Lefschetz theorem and the topology of semismall maps* (English, with English and French summaries), Ann. Sci. Éc. Norm. Supér. (4) **35** (2002), no. 5, 759–772, DOI 10.1016/S0012-9593(02)01108-4. MR1951443
- [dCM05] Mark Andrea A. de Cataldo and Luca Migliorini, *The Hodge theory of algebraic maps* (English, with English and French summaries), Ann. Sci. Éc. Norm. Supér. (4) **38** (2005), no. 5, 693–750, DOI 10.1016/j.ansens.2005.07.001. MR2195257
- [Don93] Stephen Donkin, *On tilting modules for algebraic groups*, Math. Z. **212** (1993), no. 1, 39–60, DOI 10.1007/BF02571640. MR1200163
- [Don98] S. Donkin, *The  $q$ -Schur algebra*, London Mathematical Society Lecture Note Series, vol. 253, Cambridge University Press, Cambridge, 1998. MR1707336
- [EK10] Ben Elias and Mikhail Khovanov, *Diagrammatics for Soergel categories*, Int. J. Math. Math. Sci. (2010), Art. ID 978635, 58. MR3095655
- [Eli16] Ben Elias, *The two-color Soergel calculus*, Compos. Math. **152** (2016), no. 2, 327–398, DOI 10.1112/S0010437X15007587. MR3462556
- [EW] B. Elias and G. Williamson. Soergel calculus. Represent. Theory, to appear. arXiv:1309.0865.
- [EW14] Ben Elias and Geordie Williamson, *The Hodge theory of Soergel bimodules*, Ann. of Math. (2) **180** (2014), no. 3, 1089–1136, DOI 10.4007/annals.2014.180.3.6. MR3245013
- [Fie08] Peter Fiebig, *The combinatorics of Coxeter categories*, Trans. Amer. Math. Soc. **360** (2008), no. 8, 4211–4233, DOI 10.1090/S0002-9947-08-04376-6. MR2395170
- [Fie10] Peter Fiebig, *The multiplicity one case of Lusztig’s conjecture*, Duke Math. J. **153** (2010), no. 3, 551–571, DOI 10.1215/00127094-2010-031. MR2667425
- [Fie11] Peter Fiebig, *Sheaves on affine Schubert varieties, modular representations, and Lusztig’s conjecture*, J. Amer. Math. Soc. **24** (2011), no. 1, 133–181, DOI 10.1090/S0894-0347-2010-00679-0. MR2726602
- [Fie12] Peter Fiebig, *An upper bound on the exceptional characteristics for Lusztig’s character formula*, J. Reine Angew. Math. **673** (2012), 1–31, DOI 10.1515/CRELLE.2011.170. MR2999126
- [FK14] Dmitrii A. Frolenkov and Igor D. Kan, *A strengthening of a theorem of Bourgain-Kontorovich II*, Mosc. J. Comb. Number Theory **4** (2014), no. 1, 78–117. MR3284129

- [FW14] Peter Fiebig and Geordie Williamson, *Parity sheaves, moment graphs and the  $p$ -smooth locus of Schubert varieties* (English, with English and French summaries), Ann. Inst. Fourier (Grenoble) **64** (2014), no. 2, 489–536. MR3330913
- [Gre81] James A. Green, *Polynomial representations of  $GL_n$* , Algebra, Carbondale 1980 (Proc. Conf., Southern Illinois Univ., Carbondale, Ill., 1980), Lecture Notes in Math., vol. 848, Springer, Berlin, 1981, pp. 124–140. MR613180
- [Hen96] Douglas Hensley, *A polynomial time algorithm for the Hausdorff dimension of continued fraction Cantor sets*, J. Number Theory **58** (1996), no. 1, 9–45, DOI 10.1006/jnth.1996.0058. MR1387719
- [Hua15] ShinnYih Huang, *An improvement to Zaremba’s conjecture*, Geom. Funct. Anal. **25** (2015), no. 3, 860–914, DOI 10.1007/s00039-015-0327-6. MR3361774
- [HW] X. He and G. Williamson, Soergel calculus and Schubert calculus, Bull. Inst. Math. Acad. Sin. (N.S.), to appear. arXiv:1502.04914.
- [Jam90] Gordon James, *The decomposition matrices of  $GL_n(q)$  for  $n \leq 10$* , Proc. Lond. Math. Soc. (3) **60** (1990), no. 2, 225–265, DOI 10.1112/plms/s3-60.2.225. MR1031453
- [Jan79] Jens Carsten Jantzen, *Moduln mit einem höchsten Gewicht* (German), Lecture Notes in Mathematics, vol. 750, Springer, Berlin, 1979. MR552943
- [Jan03] Jens Carsten Jantzen, *Representations of algebraic groups*, 2nd ed., Mathematical Surveys and Monographs, vol. 107, American Mathematical Society, Providence, RI, 2003. MR2015057
- [Jan08] Jens Carsten Jantzen, *Character formulae from Hermann Weyl to the present*, Groups and analysis, London Math. Soc. Lecture Note Ser., vol. 354, Cambridge Univ. Press, Cambridge, 2008, pp. 232–270, DOI 10.1017/CBO9780511721410.012. MR2528469
- [JMW14] Daniel Juteau, Carl Mautner, and Geordie Williamson, *Parity sheaves*, J. Amer. Math. Soc. **27** (2014), no. 4, 1169–1212, DOI 10.1090/S0894-0347-2014-00804-3. MR3230821
- [JW] L. T. Jensen and G. Williamson, The  $p$ -canonical basis for Hecke algebras, To appear in Perspectives in categorification, arXiv:1304.1448.
- [Kat85] Shin-ichi Kato, *On the Kazhdan-Lusztig polynomials for affine Weyl groups*, Adv. Math. **55** (1985), no. 2, 103–130, DOI 10.1016/0001-8708(85)90017-9. MR772611
- [KL93] D. Kazhdan and G. Lusztig, *Tensor structures arising from affine Lie algebras. I, II*, J. Amer. Math. Soc. **6** (1993), no. 4, 905–947, 949–1011, DOI 10.2307/2152745. MR1186962
- [KL94a] D. Kazhdan and G. Lusztig, *Tensor structures arising from affine Lie algebras. III*, J. Amer. Math. Soc. **7** (1994), no. 2, 335–381, DOI 10.2307/2152762. MR1239506
- [KL94b] D. Kazhdan and G. Lusztig, *Tensor structures arising from affine Lie algebras. IV*, J. Amer. Math. Soc. **7** (1994), no. 2, 383–453, DOI 10.2307/2152763. MR1239507
- [Kon13] Alex Kontorovich, *From Apollonius to Zaremba: local-global phenomena in thin orbits*, Bull. Amer. Math. Soc. (N.S.) **50** (2013), no. 2, 187–228, DOI 10.1090/S0273-0979-2013-01402-2. MR3020826
- [Kon14] Alex Kontorovich, *Levels of distribution and the affine sieve* (English, with English and French summaries), Ann. Fac. Sci. Toulouse Math. (6) **23** (2014), no. 5, 933–966, DOI 10.5802/afst.1432. MR3294598
- [KT95] Masaki Kashiwara and Toshiyuki Tanisaki, *Kazhdan-Lusztig conjecture for affine Lie algebras with negative level*, Duke Math. J. **77** (1995), no. 1, 21–62, DOI 10.1215/S0012-7094-95-07702-3. MR1317626
- [KT96] Masaki Kashiwara and Toshiyuki Tanisaki, *Kazhdan-Lusztig conjecture for affine Lie algebras with negative level. II. Nonintegral case*, Duke Math. J. **84** (1996), no. 3, 771–813, DOI 10.1215/S0012-7094-96-08424-0. MR1408544
- [Lib10] Nicolas Libedinsky, *Presentation of right-angled Soergel categories by generators and relations*, J. Pure Appl. Algebra **214** (2010), no. 12, 2265–2278, DOI 10.1016/j.jpaa.2010.02.026. MR2660912
- [Lib15] Nicolas Libedinsky, *Light leaves and Lusztig’s conjecture*, Adv. Math. **280** (2015), 772–807, DOI 10.1016/j.aim.2015.04.022. MR3350234
- [Lus80] George Lusztig, *Some problems in the representation theory of finite Chevalley groups*, The Santa Cruz Conference on Finite Groups (Univ. California, Santa Cruz, Calif., 1979), Proc. Sympos. Pure Math., vol. 37, Amer. Math. Soc., Providence, R.I., 1980, pp. 313–317. MR604598

- [Lus94] George Lusztig, *Monodromic systems on affine flag manifolds*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **445** (1994), no. 1923, 231–246, DOI 10.1098/rspa.1994.0058. MR1276910
- [SGS13] Alireza Salehi Golsefidy and Peter Sarnak, *The affine sieve*, J. Amer. Math. Soc. **26** (2013), no. 4, 1085–1105, DOI 10.1090/S0894-0347-2013-00764-X. MR3073885
- [Soe90] Wolfgang Soergel, *Kategorie  $\mathcal{O}$ , perverse Garben und Moduln über den Koinvarianten zur Weylgruppe* (German, with English summary), J. Amer. Math. Soc. **3** (1990), no. 2, 421–445, DOI 10.2307/1990960. MR1029692
- [Soe92] Wolfgang Soergel, *The combinatorics of Harish-Chandra bimodules*, J. Reine Angew. Math. **429** (1992), 49–74, DOI 10.1515/crll.1992.429.49. MR1173115
- [Soe97] Wolfgang Soergel, *Kazhdan-Lusztig polynomials and a combinatoric[s] for tilting modules*, Represent. Theory **1** (1997), 83–114 (electronic), DOI 10.1090/S1088-4165-97-00021-6. MR1444322
- [Soe00] Wolfgang Soergel, *On the relation between intersection cohomology and representation theory in positive characteristic*, J. Pure Appl. Algebra **152** (2000), no. 1-3, 311–335, DOI 10.1016/S0022-4049(99)00138-3. Commutative algebra, homological algebra and representation theory (Catania/Genoa/Rome, 1998). MR1784005
- [Soe01] Wolfgang Soergel, *Langlands’ philosophy and Koszul duality*, Algebra—representation theory (Constanta, 2000), NATO Sci. Ser. II Math. Phys. Chem., vol. 28, Kluwer Acad. Publ., Dordrecht, 2001, pp. 379–414, DOI 10.1007/978-94-010-0814-3\_17. MR1858045
- [Soe07] Wolfgang Soergel, *Kazhdan-Lusztig-Polynome und unzerlegbare Bimoduln über Polynomringen* (German, with English and German summaries), J. Inst. Math. Jussieu **6** (2007), no. 3, 501–525, DOI 10.1017/S1474748007000023. MR2329762
- [WB12] Geordie Williamson and Tom Braden, *Modular intersection cohomology complexes on flag varieties*, Math. Z. **272** (2012), no. 3-4, 697–727, DOI 10.1007/s00209-011-0955-y. MR2995137
- [Wil] G. Williamson. On torsion in the intersection cohomology of Schubert varieties. J. Algebra, to appear. arXiv:1512.08295.
- [Wil12] G. Williamson. Some applications of parity sheaves. *Oberwolfach reports*, 2012.

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111, BONN, GERMANY  
*E-mail address:* geordie@mpim-bonn.mpg.de

RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY  
*E-mail address:* alex.kontorovich@rutgers.edu

UNIVERSITY OF QUEENSLAND, BRISBANE, QUEENSLAND, AUSTRALIA  
*E-mail address:* p.mcnamara@uq.edu.au