

SnapShot: Microbial Extremophiles


Amy K. Schmid,^{1,2} Thorsten Allers,³ and Jocelyne DiRuggiero⁴

¹Biology Department, Duke University, Durham, NC, USA; ²Center for Genomics and Computational Biology, Duke University, Durham, NC, USA; ³School of Life Sciences, University of Nottingham, Queen's Medical Centre, Nottingham, UK; ⁴Department of Biology, Johns Hopkins University, Baltimore, MD, USA


Cell



Molecular and cellular adaptations



Limits of life and extreme record holders



SnapShot: Microbial Extremophiles

Amy K. Schmid,^{1,2} Thorsten Allers,³ and Jocelyne DiRuggiero⁴

¹Biology Department, Duke University, Durham, NC, USA; ²Center for Genomics and Computational Biology, Duke University, Durham, NC, USA; ³School of Life Sciences, University of Nottingham, Queen's Medical Centre, Nottingham, UK; ⁴Department of Biology, Johns Hopkins University, Baltimore, MD, USA

Cell

Extremophiles are overwhelmingly single-celled microbes that teach us about the limits of life, both on this planet and beyond. Life evolved under conditions that were harsher than those today and continues to thrive in environments that humans consider extreme. Earthly environments hosting extremophiles resemble extraterrestrial locations; therefore, characterizing extremophile adaptations and mechanisms of survival is essential to guide our search for life elsewhere. Knowledge of extremophiles and their enzymes has driven industrial and technological advances.

Poly-extremophiles are adapted to more than one extreme condition and are surprisingly common. Adaptation to one harsh condition often facilitates resistance to another condition as a by-product. However, the condition-based categories used to classify extremophiles do not adequately reflect their natural environment or phylogenetic history and, instead, reflect convergent evolution. Nonetheless, these categories are useful for illustrating the cellular and molecular adaptations that have evolved for thriving at the extremes, which we delineate in this SnapShot. See the online interactive map for additional detail on biogeography, record holders, and literature references: <http://bit.ly/Schmid-extremophiles-map>.

Psychrophiles | Temperature <15°C

Psychrophiles are found in low-temperature environments, such as polar oceans, lakes, and permafrost (DeMaere et al., 2013). Altered properties of psychrophilic cells include compatible solutes and chaperones, increased membrane fluidity due to shorter side chains and decreased lipid saturation, and exopolysaccharides with cryoprotective roles (Siliakus et al., 2017). To maintain flexibility and activity, psychrophilic proteins are enriched for small amino acids such as glycine; they also have fewer prolines, a reduced hydrophobic core, and a protein surface with fewer charged residues (Reed et al., 2013). These adaptations ensure a lower energy barrier between the various conformations of psychrophilic proteins.

Thermophiles | Temperature >60°C (>80°C for hyperthermophiles)

Thermophiles are found in volcanic hot springs and oceanic hydrothermal vents. Their proteins maintain activity at high temperatures via prominent hydrophobic cores, increased electrostatic interactions, and enrichment in salt and disulfide bridges (Reed et al., 2013). The thermophile-specific enzyme reverse gyrase introduces positive supercoils into DNA to maintain the double-stranded helical structure at high temperatures (Ogawa et al., 2015). The membranes of thermophiles feature lipids that often form monolayers, comprising tetra-ester (bacteria) or tetra-ether (archaea) lipids enriched for branched, saturated hydrocarbon chains (Siliakus et al., 2017). These adaptations ensure thermal stability, thereby maintaining nutrient transport and a chemiosmotic gradient.

Acidophiles | Acidic pH <3

Acidophiles are found in volcanic hot springs and acidic mine drainage. Most acidophiles keep their cytoplasm close to neutral pH using powerful proton efflux pumps and proton-deflecting membranes with reduced permeability (branched ether monolayers resembling those of thermophiles [Siliakus et al., 2017]). Some species have an acidified cytoplasm and use chaperones to maintain correct protein folding under acidic conditions; their proteins have an overabundance of acidic residues on the surface, which minimizes low pH destabilization. Many acidophiles are also resistant to high concentrations of metal ions.

Alkaliphiles | Alkaline pH >9

Alkaliphiles are found in mineral-rich soda lakes and hydrothermal alkaline vents (Banciu and Munyan, 2015). They maintain an intracellular neutral pH by cytosolic acidification, an active mechanism that uses Na^+/H^+ antiporters to accumulate intracellular protons (Padan et al., 2005). A passive mechanism found in some alkaliphilic bacteria is an acidic polymer cell wall matrix. This matrix protects the membrane by preventing the entry of hydroxide ions.

Halophiles | $\text{NaCl} >0.5 \text{ M}$ (or very high sugar for osmophiles)

Halophiles are found in salt lakes and marine salterns and use either "salt-out" or "salt-in" adaptations. The former strategy is found in bacteria and eukaryotes, which accumulate high intracellular concentrations of compatible osmolytes (e.g., glyceral, betaine) to maintain osmotic homeostasis. The latter strategy is found mainly in archaea that accumulate high levels of KCl in their cytoplasm using a K^+/Na^+ antiporter (Gunde-Cimerman et al., 2018). This requires protein adaptions, including an increased negative surface charge due to a high acidic amino acid content (Reed et al., 2013). In high light and low oxygen, many halophiles use bacteriorhodopsin (BR), a light-driven proton pump, to generate ATP to drive the antiporter and maintain osmobalance (Gunde-Cimerman et al., 2018). Photoprotective membrane-associated pigments (e.g., carotenoids) protect against frequent UV damage in hypersaline environments. Many hypersaline-adapted archaea are also polyextremophiles adapted to low temperatures and high pH (Banciu and Munyan, 2015; DeMaere et al., 2013).

Xerophiles | Desiccation, water activity (a_w) <0.8

Xerophiles are found in dry environments such as hot and cold deserts, including the Dry Valleys of Antarctica. Xerophiles use similar adaptations to salt-out halophiles, accumulating high intracellular concentrations of compatible osmolytes (glyceral, betaine, trehalose or glyceral) to cope with low a_w (Lebre et al., 2017). Xerophiles are often radioresistant because the mechanisms for dealing with desiccation render them resistant to ionizing radiation and oxidative stress (Hallsworth, 2018). Xerophiles have elevated levels of antioxidants to protect against desiccation and radiation-induced damage to proteins. For example, *Deinococcus radiodurans* uses manganese complexes as antioxidants (Daly, 2009).

ACKNOWLEDGEMENTS

We are grateful to Schmid lab members Rylee Hackley, Angie Vrugdenhil, Saaz Sakrikar, and Sungmin Hwang for their suggestions. Special thanks to Eric Monson in the Duke University Library Center for Data and Visualization Sciences for advice on the graphics. Funding for this article was provided by NASA 18-EXO18-0091 and NSF DEB1556574 to J.D., NSF MCB1651117 and 1615685 to A.K.S., and Biotechnology and Biological Sciences Research Council (BBSRC) grant numbers BB/N016491/1 and BB/R007543/1 to T.A. Photo credits: Yellowstone Octopus spring, Jocelyne DiRuggiero; Eilat salt flats, Thorsten Allers; Deep sea vent, adapted from Marum Center for Marine Environmental Sciences (CC BY 4.0 license); Atacama desert, Henry Bortman; Deep Lake, Ricardo Cavicchioli; Bacteriorhodopsin, PDB: 1FBB, 10.1038/35020614.

REFERENCES

Banciu, H.L., and Munyan, M.S. (2015). Adaptive strategies in the double-extremophilic prokaryotes inhabiting soda lakes. *Curr. Opin. Microbiol.* 25, 73–79.

Daly, M.J. (2009). A new perspective on radiation resistance based on *Deinococcus radiodurans*. *Nat. Rev. Microbiol.* 7, 237–245.

DeMaere, M.Z., Williams, T.J., Allen, M.A., Brown, M.V., Gibson, J.A., Rich, J., Lauro, F.M., Dyall-Smith, M., Davenport, K.W., Woyke, T., et al. (2013). High level of intergenera gene exchange shapes the evolution of haloarchaea in an isolated Antarctic lake. *Proc. Natl. Acad. Sci. USA* 110, 16939–16944.

Gunde-Cimerman, N., Plemenitaš, A., and Oren, A. (2018). Strategies of adaptation of microorganisms of the three domains of life to high salt concentrations. *FEMS Microbiol. Rev.* 42, 353–375.

Hallsworth, J.E. (2018). Stress-free microbes lack vitality. *Fungal Biol.* 122, 379–385.

Lebre, P.H., De Maayer, P., and Cowan, D.A. (2017). Xerotolerant bacteria: surviving through a dry spell. *Nat. Rev. Microbiol.* 15, 285–296.

Ogawa, T., Yogo, K., Furuike, S., Sutoh, K., Kikuchi, A., and Kinoshita, K., Jr. (2015). Direct observation of DNA overwinding by reverse gyrase. *Proc. Natl. Acad. Sci. USA* 112, 7495–7500.

Padan, E., Bibi, E., Ito, M., and Krulwich, T.A. (2005). Alkaline pH homeostasis in bacteria: new insights. *Biochim. Biophys. Acta* 1717, 67–88.

Reed, C.J., Lewis, H., Trejo, E., Winston, V., and Evilia, C. (2013). Protein adaptions in archaeal extremophiles. *Archaea* 2013, 373275.

Siliakus, M.F., van der Oost, J., and Kengen, S.W.M. (2017). Adaptations of archaeal and bacterial membranes to variations in temperature, pH and pressure. *Extremophiles* 21, 651–670.