



Cite this: *Dalton Trans.*, 2019, **48**, 6319

Received 9th March 2019,
Accepted 26th March 2019

DOI: 10.1039/c9dt01032g

rsc.li/dalton

Successive carbene insertion into 9-phenyl-9-borafluorene†

Tyler A. Bartholome, Kristen R. Bluer and Caleb D. Martin *

The reactions of 9-phenyl-9-borafluorene with trimethylsilyldiazomethane in a 1:1 and 1:2 stoichiometry furnished the corresponding BC₅ and BC₆ heterocycles via the formal insertion of one and two carbene units.

Unsaturated BC₄ heterocycles have recently been recognized as promising reagents for the synthesis of six- to eight-membered boracyclic systems via the formal insertion of one, two, or three atoms into the endocyclic B–C bond.^{1–3} Developing efficient methodologies to access heterocycles featuring tri-coordinate boron centers is of paramount interest as they have applications in electronic materials and pharmaceuticals.^{4–13} In this vein, two types of unsaturated BC₄ systems have been focused upon, boroles (**1**)¹⁴ and their biphenyl-fused variants, 9-borafluorenes (**2**).^{15,16} Boroles are more reactive than 9-borafluorenes as they have greater anti-aromatic character and Lewis acidity.^{17–22} The intermolecular insertion chemistry of boroles has been investigated with a number of unsaturated molecules^{23–44} whereas the corresponding chemistry with 9-borafluorenes is less developed.^{19,45–53}

The first intermolecular insertion reaction of a 9-borafluorene was reported in 2016 by Fukushima and coworkers with the insertion of alkynes into the endocyclic B–C bond of 9-chloro-9-borafluorene (**2Cl**) to furnish unsaturated BC₆ heterocycles (**A** and **B**; Fig. 1a).⁵² Our group, and He, discovered that 9-borafluorenes react with organic azides to generate 9,10-*B,N*-phenanthrenes (**C** and **D**, Fig. 1b) via the insertion of either the α - or γ -nitrogen atom.^{19,51} Investigations with 1,2-dipolar substrates resulted in adducts (imine, nitrile, isocyanide) or seven membered rings (aldehyde, ketone, ketene, isocyanate, carbodiimide, phosphalkyne) in which the negatively polarized atom is bound to boron and the positively polarized atom to carbon (Fig. 1c).^{46–48}

The Ashe, Brown, and Matteson groups have demonstrated that the combination of halomethane/base can serve as a carbene source that inserts into B–C bonds.^{54–62} The insertion of CR₂ units into boron–carbon bonds has also been reported using diazo reagents with notable recent work by the Stephan and Melen groups.^{63–69} Braunschweig and coworkers investigated the reactions of boroles (**1**) with diazo reagents observing two outcomes (Fig. 2).²⁴ The reaction with a bulky diazo species featuring two *para*-tolyl groups on the α -carbon generated a 1,2-azaborine (**F**) via insertion of the terminal nitrogen into the B–C bond whereas the reaction with trimethylsilyldiazomethane resulted in insertion of a CH(SiMe₃) unit into the endocyclic boron–carbon bond of **1** to provide a six-membered ring (**G**) with concomitant release of N₂ gas. Given the insertion chemistry of diazo compounds into boroles, we sought to examine the reactivity of trimethylsilyldiazomethane with 9-phenyl-9-borafluorene (**2Ph**).

The 1:1 stoichiometric reaction of **2Ph** with trimethylsilyldiazomethane at room temperature resulted in gas evol-




Fig. 1 Reported intermolecular insertion reactions of alkynes (a), azides (b), and unsaturated 1,2-dipolar molecules (c) with 9-borafluorenes. TMS = trimethylsilyl.

Baylor University, Department of Chemistry and Biochemistry,

One Bear Place #97348, Waco, TX 76798, USA. E-mail: caleb_d_martin@baylor.edu

†Electronic supplementary information (ESI) available. CCDC 1901552 and 1901553. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9dt01032g





Fig. 2 Reactions of pentaphenylborole (**1**) with diazo reagents (tol = *para*-tolyl).

ution accompanied by a color change from yellow to colorless (Scheme 1). No significant change was observed by *in situ* ^{11}B NMR spectroscopy; however, *in situ* ^1H NMR spectroscopy indicated complete consumption of trimethylsilyldiazomethane within 10 minutes by the disappearance of the trimethylsilyl peak at -0.04 ppm accompanied by the emergence of a singlet at -0.30 ppm. A diagnostic singlet in the aliphatic region at 3.88 ppm was assigned to the proton derived from the diazo α -carbon. Removing the volatiles *in vacuo* gave a pale yellow solid in quantitative yield. An X-ray diffraction study on crystals grown revealed the product as the BC_5 heterocycle (**3**, Fig. 3) from the formal insertion of a carbene unit into one of the B–C bonds. The product crystallizes in the $P\bar{1}$ space group, indicating a 50 : 50 racemate of **3** in the unit cell. The geometry about the boron center is trigonal planar [angular sum = $359.9(2)^\circ$]



Scheme 1 Reaction of trimethylsilyldiazomethane with **2Ph**.




Fig. 3 Solid-state structure of **3**. Hydrogen atoms are omitted for clarity (except at the chiral center), and ellipsoids are drawn at the 50% probability level. Selected bond lengths (Å) and angles ($^\circ$): B(1)–C(1) 1.541(3), C(1)–Si(1) 1.948(2), C(1)–C(2) 1.500(3), C(2)–C(7) 1.406(3), C(7)–C(8) 1.484(3), C(8)–C(13) 1.422(3), C(13)–B(1) 1.551(3), B(1)–C(14) 1.570(3), C(13)–B(1)–C(14) 122.26(18), C(13)–B(1)–C(1) 116.48(16), C(14)–B(1)–C(1) 121.17(16).

and the biphenyl backbone has a slight twist [interplanar angle = $9.9(2)^\circ$].

To determine if another equivalent of trimethylsilyldiazomethane would react with **2Ph**, the 1 : 1 reaction of trimethylsilyldiazomethane with **3** was conducted at room temperature. Gas evolution was observed upon addition, and *in situ* ^{11}B NMR spectroscopy indicated conversion after 20 minutes to a new resonance at 75.4 ppm, shifted downfield from **3** (*cf.* 65.8 ppm). After work up, colorless crystals were isolated in a 45% yield. Alternatively, the same product could be obtained by reacting **2Ph** with trimethylsilyldiazomethane directly in a 1 : 2 stoichiometric ratio. Redissolving the solids in C_6D_6 and acquiring an ^1H NMR spectrum revealed two singlets at -0.07 ppm and -0.59 ppm integrating in a 9 : 9 ratio, shifted upfield from trimethylsilyldiazomethane (-0.04 ppm). Singlets at 3.24 ppm and 2.70 ppm, each integrating to one, indicate the presence of two aliphatic protons. A single crystal X-ray diffraction study unambiguously identified the compound as the BC_6 heterocycle **4** from a second carbene insertion into the other B–C_{biphenyl} bond of **2Ph**. The X-ray crystal structure revealed that the product is a single *meso* isomer in which the trimethylsilyl groups are oriented *cis* with respect to each other (Fig. 4).⁷⁰ The BC_6 ring adopts a boat-like conformation which does not include a plane of symmetry about the boron heteroatom (Fig. S-16†), rationalizing the presence of non-equivalent ^1H NMR signals for the two trimethylsilyl groups and the aliphatic protons assigned to those on the chiral carbon centers adjacent to boron.⁷¹ The geometry at boron is trigonal planar [angular sum = $360.0(2)^\circ$], and the twist in the biphenyl group is significantly more pronounced with an interplanar angle of $48.7(2)^\circ$ [*cf.* **3** = $9.9(2)^\circ$].

The limit of the reactivity of **2Ph** was examined by conducting the reaction with excess trimethylsilyldiazomethane at 80 $^\circ\text{C}$, upon which only the BC_6 product **4** was observed by ^1H




Fig. 4 Solid-state structure of **4**. Hydrogen atoms are omitted for clarity (except at chiral centers), and ellipsoids are drawn at the 50% probability level. Selected bond lengths (Å) and angles ($^\circ$): B(1)–C(1) 1.587(3), C(1)–C(2) 1.523(2), C(2)–C(7) 1.409(3), C(7)–C(8) 1.486(3), C(13)–C(14) 1.509(2), C(14)–Si(2) 1.9302(17), C(14)–B(1) 1.567(3), B(1)–C(15) 1.568(3), C(14)–B(1)–C(15) 119.08(15), C(14)–B(1)–C(1) 117.36(15), C(15)–B(1)–C(1) 123.54(15).

and ^{11}B NMR spectroscopy. Examining the corresponding reaction of excess trimethylsilyldiazomethane with the borole variant, **1**, only gave the BC_5 product (**G**) from reaction with one equivalent. Upon heating at 80 °C for 24 hours, an indiscernible mixture was observed by *in situ* ^{11}B and ^1H NMR spectroscopy.

In summary, the reactions of trimethylsilyldiazomethane with 9-phenyl-9-borafluorene result in the sequential insertion of two carbene units, one into each of the endocyclic B–C bonds of the central BC_4 ring. This represents the first example in which insertion into both B–C bonds is observed for a 9-borafluorene. Interestingly, the chemistry of penta-phenylborole differs as only a single insertion occurs cleanly. The results further demonstrate the potential of 9-borafluorenes to serve as reagents for the preparation of polycyclic boron species.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We are grateful to the Welch Foundation (Grant No. AA-1846) and the National Science Foundation for a CAREER Award (Award No. 1753025) for their generous support of this work.

Notes and references

- J. H. Barnard, S. Yruegas, K. Huang and C. D. Martin, *Chem. Commun.*, 2016, **52**, 9985–9991.
- B. Su and R. Kinjo, *Synthesis*, 2017, **49**, 2985–3034.
- G. Bélanger-Chabot, H. Braunschweig and D. K. Roy, *Eur. J. Inorg. Chem.*, 2017, 4353–4368.
- I. A. Adams and P. A. Rupar, *Macromol. Rapid Commun.*, 2015, **36**, 1336–1340.
- J.-F. Brière and M. Côté, *J. Phys. Chem. B*, 2004, **108**, 3123–3129.
- A. Escande and M. J. Ingleson, *Chem. Commun.*, 2015, **51**, 6257–6274.
- M. Lorenz-Rothe, K. S. Schellhammer, T. Jägeler-Hoheisel, R. Meerheim, S. Kraner, M. P. Hein, C. Schünemann, M. L. Tietze, M. Hummert, F. Ortmann, G. Cuniberti, C. Körner and K. Leo, *Adv. Electron. Mater.*, 2016, **2**, 1600152.
- T. Matsumoto, H. Takamine, K. Tanaka and Y. Chujo, *Mater. Chem. Front.*, 2017, **1**, 2368–2375.
- S. Muhammad, M. R. S. A. Janjua and Z. Su, *J. Phys. Chem. C*, 2009, **113**, 12551–12557.
- K. S. Thanthiriatte and S. R. Gwaltney, *J. Phys. Chem. A*, 2006, **110**, 2434–2439.
- S. Yamaguchi, T. Shirasaka, S. Akiyama and K. Tamao, *J. Am. Chem. Soc.*, 2002, **124**, 8816–8817.
- S. Yamaguchi and A. Wakamiya, *Pure Appl. Chem.*, 2006, **78**, 1413–1424.
- S. J. Cassidy, I. Brettell-Adams, L. E. McNamara, M. F. Smith, M. Bautista, H. Cao, M. Vasiliu, D. L. Gerlach, F. Qu, N. I. Hammer, D. A. Dixon and P. A. Rupar, *Organometallics*, 2018, **37**, 3732–3741.
- J. J. Eisch, N. H. Hota and S. Kozima, *J. Am. Chem. Soc.*, 1969, **91**, 4575–4577.
- R. Köster and G. Benedikt, *Angew. Chem., Int. Ed. Engl.*, 1963, **2**, 323–324.
- P. E. Romero, W. E. Piers, S. A. Decker, D. Chau, T. K. Woo and M. Parvez, *Organometallics*, 2003, **22**, 1266–1274.
- P. A. Chase, P. E. Romero, W. E. Piers, M. Parvez and B. O. Patrick, *Can. J. Chem.*, 2005, **83**, 2098–2105.
- C. J. Berger, G. He, C. Merten, R. McDonald, M. J. Ferguson and E. Rivard, *Inorg. Chem.*, 2014, **53**, 1475–1486.
- S. Yruegas, J. J. Martinez and C. D. Martin, *Chem. Commun.*, 2018, **54**, 6808–6811.
- M. F. Smith, S. J. Cassidy, I. A. Adams, M. Vasiliu, D. L. Gerlach, D. A. Dixon and P. A. Rupar, *Organometallics*, 2016, **35**, 3182–3191.
- B. C. Caputo, Z. J. Manning, J. H. Barnard and C. D. Martin, *Polyhedron*, 2016, **114**, 273–277.
- L. E. Laperriere, S. Yruegas and C. D. Martin, *Tetrahedron*, 2019, **75**, 937–943.
- H. Braunschweig, C. Hörl, L. Mailänder, K. Radacki and J. Wahler, *Chem. – Eur. J.*, 2014, **20**, 9858–9861.
- H. Braunschweig, F. Hupp, I. Krummenacher, L. Mailänder and F. Rauch, *Chem. – Eur. J.*, 2015, **21**, 17844–17849.
- H. Braunschweig, M. A. Celik, F. Hupp, I. Krummenacher and L. Mailänder, *Angew. Chem., Int. Ed.*, 2015, **54**, 6347–6351.
- H. Braunschweig, I. Krummenacher, L. Mailänder and F. Rauch, *Chem. Commun.*, 2015, **51**, 14513–14515.
- H. Braunschweig, M. Celik, T. Dellermann, G. Frenking, K. Hammond, F. Hupp, H. Kelch, I. Krummenacher, F. Lindl, L. Mailänder, J. Müssig and A. Ruppert, *Chem. – Eur. J.*, 2017, **23**, 8006–8013.
- Z. Wang, Y. Zhou, J.-X. Zhang, I. Krummenacher, H. Braunschweig and Z. Lin, *Chem. – Eur. J.*, 2018, **24**, 9612–9621.
- J. H. Barnard, P. A. Brown, K. L. Shuford and C. D. Martin, *Angew. Chem., Int. Ed.*, 2015, **54**, 12083–12086.
- S. Yruegas and C. D. Martin, *Chem. – Eur. J.*, 2016, **22**, 18358–18361.
- S. Yruegas, D. C. Patterson and C. D. Martin, *Chem. Commun.*, 2016, **52**, 6658–6661.
- J. J. Eisch, J. E. Galle, B. Shafii and A. L. Rheingold, *Organometallics*, 1990, **9**, 2342–2349.
- S. A. Couchman, T. K. Thompson, D. J. D. Wilson, J. L. Dutton and C. D. Martin, *Chem. Commun.*, 2014, **50**, 11724–11726.
- K. Huang and C. D. Martin, *Inorg. Chem.*, 2015, **54**, 1869–1875.
- K. Huang, S. A. Couchman, D. J. D. Wilson, J. L. Dutton and C. D. Martin, *Inorg. Chem.*, 2015, **54**, 8957–8968.

- 36 J. H. Barnard, S. Yruegas, S. A. Couchman, D. J. D. Wilson, J. L. Dutton and C. D. Martin, *Organometallics*, 2016, **35**, 929–931.
- 37 K. Huang and C. D. Martin, *Inorg. Chem.*, 2016, **55**, 330–337.
- 38 V. A. K. Adiraju and C. D. Martin, *Chem. – Eur. J.*, 2017, **23**, 11437–11444.
- 39 S. Yruegas, C. Wilson, J. L. Dutton and C. D. Martin, *Organometallics*, 2017, **36**, 2581–2587.
- 40 A. Fukazawa, J. L. Dutton, C. Fan, L. G. Mercier, A. Y. Houghton, Q. Wu, W. E. Piers and M. Parvez, *Chem. Sci.*, 2012, **3**, 1814–1818.
- 41 F. Lindl, S. Lin, I. Krummenacher, C. Lenczyk, A. Stoy, M. Müller, Z. Lin and H. Braunschweig, *Angew. Chem., Int. Ed.*, 2019, **58**, 338–342.
- 42 Y. Li, R. K. Siwatch, T. Mondal, Y. Li, R. Ganguly, D. Koley and C.-W. So, *Inorg. Chem.*, 2017, **56**, 4112–4120.
- 43 C. Fan, W. E. Piers, M. Parvez and R. McDonald, *Organometallics*, 2010, **29**, 5132–5139.
- 44 H. Braunschweig, J. Maier, K. Radacki and J. Wahler, *Organometallics*, 2013, **32**, 6353–6359.
- 45 Y. Shoji, M. Hwang, H. Sugiyama, F. Ishiwari, K. Takenouchi, R. Osuga, J. N. Kondo, S. Fujikawa and T. Fukushima, *Mater. Chem. Front.*, 2018, **2**, 807–814.
- 46 K. R. Bluer, L. E. Laperriere, A. Pujol, S. Yruegas, V. A. K. Adiraju and C. D. Martin, *Organometallics*, 2018, **37**, 2917–2927.
- 47 S. Yruegas, J. H. Barnard, K. Al-Furaiji, J. L. Dutton, D. J. D. Wilson and C. D. Martin, *Organometallics*, 2018, **37**, 1515–1518.
- 48 J. Kashida, Y. Shoji and T. Fukushima, *Chem. – Asian J.*, DOI: 10.1002/asia.201900047.
- 49 Y. Shoji, N. Shigeno, K. Takenouchi, M. Sugimoto and T. Fukushima, *Chem. – Eur. J.*, 2018, **24**, 13223–13230.
- 50 C. Özen, Y. Shoji, T. Fukushima and S. Maeda, *J. Org. Chem.*, 2019, **84**, 1941–1950.
- 51 W. Zhang, G. Li, L. Xu, Y. Zhuo, W. Wan, N. Yan and G. He, *Chem. Sci.*, 2018, **9**, 4444–4450.
- 52 Y. Shoji, N. Tanaka, S. Muranaka, N. Shigeno, H. Sugiyama, K. Takenouchi, F. Hajjaj and T. Fukushima, *Nat. Commun.*, 2016, **7**, 12704.
- 53 W. Zhang, D. Yu, Z. Wang, B. Zhang, L. Xu, G. Li, N. Yan, E. Rivard and G. He, *Org. Lett.*, 2019, **21**, 109–113.
- 54 D. S. Matteson and D. Majumdar, *Organometallics*, 1983, **2**, 1529–1535.
- 55 K. M. Sadhu and D. S. Matteson, *Organometallics*, 1985, **4**, 1687–1689.
- 56 D. S. Matteson and D. Majumdar, *J. Am. Chem. Soc.*, 1980, **102**, 7588–7590.
- 57 H. C. Brown, R. G. Naik, R. K. Bakshi, C. Pyun and B. Singaram, *J. Org. Chem.*, 1985, **50**, 5586–5592.
- 58 H. C. Brown, S. M. Singh and M. V. Rangaishenvi, *J. Org. Chem.*, 1986, **51**, 3150–3155.
- 59 H. C. Brown, A. S. Phadke and M. V. Rangaishenvi, *J. Am. Chem. Soc.*, 1988, **110**, 6263–6264.
- 60 A. D. Rohr, M. M. Banaszak Holl, J. W. Kampf and A. J. Ashe III, *Organometallics*, 2011, **30**, 3698–3700.
- 61 A. J. Ashe III, X. Fang, X. Fang and J. W. Kampf, *Organometallics*, 2001, **20**, 5413–5418.
- 62 J. Chen, Z. Bajko, J. W. Kampf and A. J. Ashe III, *Organometallics*, 2007, **26**, 1563–1564.
- 63 R. L. Melen, *Angew. Chem., Int. Ed.*, 2018, **57**, 880–882.
- 64 A. J. Ruddy, D. M. C. Ould, P. D. Newman and R. L. Melen, *Dalton Trans.*, 2018, **47**, 10377–10381.
- 65 C. Schneider, J. H. W. LaFortune, R. L. Melen and D. W. Stephan, *Dalton Trans.*, 2018, **47**, 12742–12749.
- 66 R. C. Neu and D. W. Stephan, *Organometallics*, 2012, **31**, 46–49.
- 67 R. C. Neu, C. Jiang and D. W. Stephan, *Dalton Trans.*, 2013, **42**, 726–736.
- 68 J. Lam, B. A. R. Günther, J. M. Farrell, P. Eisenberger, B. P. Bestvater, P. D. Newman, R. L. Melen, C. M. Crudden and D. W. Stephan, *Dalton Trans.*, 2016, **45**, 15303–15316.
- 69 C. Tang, Q. Liang, A. R. Jupp, T. C. Johnstone, R. C. Neu, D. Song, S. Grimme and D. W. Stephan, *Angew. Chem., Int. Ed.*, 2017, **56**, 16588–16592.
- 70 From the experiments it is not clear if the observed *meso* isomer is the kinetic or thermodynamic product.
- 71 A variable temperature ^1H NMR spectroscopy experiment from $-60\text{ }^\circ\text{C}$ to $90\text{ }^\circ\text{C}$ in toluene- d_8 did not reveal any signs of coalescence of the signals.