

Gecko diversity: a history of global discovery

Peter Uetz^{a,*}, Alex Slavenko^{b,c}, Shai Meiri^{c,d} and Matthew Heinicke^{e,*}

^aCenter for Biological Data Science, Virginia Commonwealth University, 1000 W Cary St, Richmond, VA 23284, USA

^bDepartment of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

^cSchool of Zoology, Tel Aviv University, 6997801, Tel-Aviv, Israel

^dThe Steinhardt Museum of Natural History, Tel Aviv University, 6997801, Tel-Aviv, Israel

^eDepartment of Natural Sciences, University of Michigan-Dearborn, 4901 Evergreen Rd., Dearborn, MI 48128, USA

Abstract 1935 gecko species (and 224 subspecies) were known in December 2019 in seven families and 124 genera. These nearly 2000 species were described by ~950 individuals of whom more than 100 described more than 10 gecko species each. Most gecko species were discovered during the past 40 years. The primary type specimens of all currently recognized geckos (including subspecies) are distributed over 161 collections worldwide, with 20 collections having about two thirds of all primary types. The primary type specimens of about 40 gecko taxa have been lost or unknown. The phylogeny of geckos is well studied, with DNA sequences being available for ~76% of all geckos (compared to ~63% in other reptiles) and morphological characters now being collected in databases. Geographically, geckos occur on five continents and many islands but are most species-rich in Australasia (which also houses the greatest diversity of family-level taxa), Southeast Asia, Africa, Madagascar, and the West Indies. Among countries, Australia has the highest number of geckos (241 species), with India, Madagascar, and Malaysia being the only other countries with more than 100 described species each. As expected, when correcting for land area, countries outside the tropics have fewer geckos.

Keywords Carphodactylidae; Diplodactylidae; Eublepharidae; Gekkonidae; Gekkota; Phyllodactylidae; Pygopodidae; Sphaerodactylidae

Introduction

Geckos (Sauria: Gekkota; 1935 species) are one of three mega-diverse lineages of squamate reptiles (lizards, snakes, and amphisbaenians), along with the 1685 species of skinks and 1965 species of colubrid snakes (Uetz et al. 2019), that are known today as result of the major squamate radiations that began diversifying about 200 million years ago. All gecko families are relatively old compared to either skinks or colubrids. Molecular clock estimates place the origins of gecko families deep in the Mesozoic (Gamble et al. 2008a, b, 2011; Hedges et al. 2015; Zheng and Wiens 2016), and stem gekkotan fossils dating from the late Jurassic and Cretaceous have been recovered from multiple distant localities in Eurasia (Daza et al. 2014, 2016; Gauthier et al. 2012; Simões et al. 2017). Not all gecko lineages have diversified at the same rate. For example, there are 38 species of Eublepharidae, compared to 1632 species in their sister lineage (Gekkonidae + Phyllodactylidae + Sphaerodactylidae). Thus, the high species richness of geckos has been produced largely by diversification of a subset of successful lineages.

Here we focus on the history of discovery and description of gecko species. In addition, we review the diversity of geckos in terms of species numbers, both taxonomically and geographically, but also in terms of discovery. As mostly small and nocturnal species (Meiri 2020, this volume), many geckos are easy to overlook, though this

is obviously not true for human commensals such as some *Hemidactylus* or conspicuous day geckos such as *Lygodactylus* or *Phelsuma*. Nevertheless, many geckos were described early in the history of herpetology. We finally discuss the factors for species discovery and diversity and how it relates to gecko biology.

A history of gecko discovery

Only three geckos were described by Linnaeus (1758) — the Tokay gecko (*Lacerta Gecko* to Linnaeus, now *Gekko gecko*), Mediterranean house gecko (as *Lacerta turcica*, now *Hemidactylus turcicus*), and Moorish gecko (*Lacerta mauritanica*, now *Tarentola mauritanica*). It then took herpetologists 227 years, from 1758 to 1984, to describe the first 1000 gecko species. It has taken only 35 to describe the next 921 (not counting subspecies). Early descriptions of gecko species commonly appeared in regional monographs or travelogues (e.g. Spix 1825) or else more general zoological works (e.g. Daudin 1802). Some also appeared as stand-alone contributions to journals or society proceedings (e.g. Sparrman 1778). Early descriptions peaked in the mid-19th century with 19 species described in each of 1836, 1845, 1870, and 1885 (Fig. 1). These numbers were driven by the monumental works of André M.C. Duméril and Gabriel Bibron (Duméril and Bibron 1836), John E. Gray (Gray 1845), Richard H. Beddome (Beddome

*Corresponding authors. E-mails: uetz@vcu.edu / heinicke@umich.edu




Figure 1. Gecko species described per year 1758–2019. Some prolific gecko describers from the 19th century are highlighted. The number of new species descriptions has surged in the past 15 years, supported by widely accessible molecular techniques and other advances.

1870a, b) and George A. Boulenger (Boulenger 1885) that were published in those years. These annual totals were only exceeded a century later with 24 species described in 1978. Even though 19th century authors have described most geckos in single publications, only Gray and Boulenger are among the 10 most prolific gecko describers (Table 1). In the late 20th century molecular methods such as karyotyping (e.g. Murphy 1974; King 1982) and allozyme electrophoresis (e.g. Branch et al. 1995) began to be employed to aid in new species discovery. Discoveries skyrocketed in the 21st century (Meiri 2016; Uetz and Stylianou 2018) with the advent of new technologies, such as the internet, DNA sequencing, digital photography, and cheaper travel permitting access to remote areas, as well as the ability for individual researchers to study collections at distant museums. Nevertheless, even in modern times, gecko discovery has been driven by relatively few individuals. Thus, the 1935 gekkotans described since 1758 were authored by about 950 individuals (Uetz and Stylianou 2018), of whom about 100 described more than 10 gecko species each. Eight of the ten most prolific describers of new species are currently active herpetologists, with two (Aaron M. Bauer and L. Lee Grismer) describing more than 130 species each (Table 1).

From the 18th through 20th centuries, most gecko species were described by one or two authors. The earliest gecko species description with more than two authors appeared in 1970 (Minton et al. 1970). Team taxonomy has become the norm in the 21st century, as different scientists are often needed to carry out distinct tasks in the process of

Table 1. Top-11 authors who described the most gecko species still recognized as valid (i.e. 40 or more).

Author	Species
Aaron M. Bauer	143
L. Lee Grismer	132
Perry L. Wood	98
George A. Boulenger	77
Evan S. H. Quah	63
Olivier S. G. Pauwels	54
John E. Gray	49
Montri Sumontha	45
Thomas Ziegler	44
Paul Doughty	40
Paul M. Oliver	40

species discovery such as fieldwork, morphological work, molecular work, specimen comparisons, statistical analysis, and literature review. In some cases this may result in species descriptions with many authors. For instance, several gecko species have been described with more than a dozen authors, such as *Cyrtodactylus phuocbinhensis* Nguyen et al. 2013, *Cyrtodactylus taynguyenensis* Nguyen et al. 2013, *Cyrtodactylus puhuensis* Nguyen et al. 2014, and *Cnemaspis bidongensis* Grismer et al. 2014, each with 14 authors. None of these approach the reptile species with the highest number of authors though, which is the leiosaurid *Enyalius capetinga* Breitman et al. 2018, with 27 authors. Many of the most prolific gecko describers (Table 1) have worked together, thus, for example, almost all the descriptions by Perry Wood and Evan Quah were co-authored by Lee Grismer.

Type specimens of geckos

The primary types of the ~2000 species of geckos are kept in 161 collections worldwide, with 20 collections having about two thirds of all types (see also Uetz et al. 2019, Table 2, Fig. 2). This is important for researchers who describe new species and need to compare them to the types of previously described ones. By far the most gecko primary type specimens are held at the Natural History Museum, London (BMNH; types of 285 taxa). Among its collections are most of the types of species described by Gray, Boulenger, and Beddome in their major 19th century works, along with many types designated by Nicolas Arnold, Albert Günther, Hampton Wildman Parker, Malcolm Smith, and others, and its type specimens originate from across the globe. The Muséum National d'Histoire Naturelle, Paris (MNHN) has a similar global scope and many types dating from the 19th century work of Duméril and Bibron with more recent types, e.g. designated by Aaron Bauer and Georges Pasteur, among others. Major collections often have geographic foci that reflect the work of scientists affiliated with these institutions. The Museum of Comparative Zoology (MCZ), for example, includes many types of African species from the work of former curator Arthur Loveridge, and a large collection of West Indian *Sphaerodactylus* types designated by former director Thomas Barbour (plus Albert Schwartz and Richard Thomas). Interestingly, most of these museums reside in places where no native gecko species are found (Roll et al. 2017, Meiri, 2019, this volume). Only two of the top ten collections are held in locations with native geckos: the Western Australian Museum (WAM) and Ditsong National Museum, Pretoria (DNMNH). All of the gecko

Table 2. The top-10 collections that hold the most gekkotan primary types (species and subspecies). For additional type information see Uetz et al. (2019).

Collection	Taxa with types
BMNH (London, UK)	285
MCZ (Cambridge, USA)	130
MNHN-RA (Paris, France)	107
WAM (Perth, Australia)	100
USNM (Washington, DC, USA)	68
CAS (San Francisco, USA)	69
SMF (Frankfurt, Germany)	68
ZMB (Berlin, Germany)	57
ZFMK (Bonn, Germany)	55
DNMNH (Pretoria, South Africa)	52

primary types held at these two institutions originate from their respective continents. Fifty one institutions have only a single primary gecko type specimen and 21 have two.

The VertNet database (Constable et al. 2010) is the largest meta-database of vertebrate collections, and returned 11,888 entries when searched for gekkotans with type status (in Nov 2019). However, only 568 of these are primary types (holo-, syn-, lecto-, or neotypes) corresponding to 430 species in the Reptile Database (possibly up to ~500 species when all mismatched names such as typos and spelling variants are included, ignoring synonyms). That is, ~25% of all Gekkotans have primary types recorded in VertNet but the vast majority of all VertNet-listed types are secondary types, including 6,542 paratypes, which may be missing from the primary type catalog that Uetz et al. (2019) compiled. VertNet is one of the major North American efforts to consolidate digitized vertebrate collections,




Figure 2. All primary types of the world's geckos are in 161 collections with 20 collections having about two thirds of all types. Type count (X axis) is the number of currently valid taxa (species and subspecies) with primary types.

Table 3. Diversity of geckos in terms of families and species numbers.

Family	species	genera
Carpodactylidae	31	7
Diplodactylidae	154	25
Eublepharidae	38	6
Gekkonidae	1295	57
Phyllodactylidae	148	10
Pygopodidae	45	7
Sphaerodactylidae	224	12
All Gekkota	1935	124
% of all reptiles	17%	10%

and much more advanced than similar projects in other parts of the world. Thus potentially only a small fraction of all collections with gecko specimens have submitted their collection data to meta-databases, though many collections have in-house databases.

A relatively small number of primary gecko type specimens are unknown. We found the types of 40 valid gekkotan taxa (~2%, out of 2159, including subspecies) to be either lost or simply “unlocated” (i.e. their whereabouts were never made clear, even in the original description; e.g. for *Tropiocolotes nattereri* Steindachner 1901)—which means that they are likely lost too. Thus, surprisingly, geckos are less often lost than non-gekkotan types, of which more than 5% are lost or unlocated (Uetz et al. 2019). This is despite the often small size of geckos, but likely due to the fact that most geckos were described only recently and thus had less time to get lost.



Figure 3. Phylogenetic relationship of gecko families. Relationships are based on recent comprehensive molecular phylogenetic studies (Han et al. 2004; Gamble et al. 2008a, b, 2012; Zheng and Wiens 2016).

The diversity of geckos

The nearly 2000 species of geckos represent a tremendous variety of adaptations and lifestyles, too many to be thoroughly reviewed here (see Meiri 2020, this issue, for more details). However, the diversity is reflected by their classification into 7 families and 124 genera (Table 3, Figs. 3, 4). These were traditionally recognized by morphological characters such as feet (absent in pygopods), their eyes and eyelids (true eyelids are present only in eublepharids), and their toepads (carphodactylids and eublepharids both lack adhesive toepads, as do many members of the toepad-bearing families; Bauer 2013). Of the seven currently recognized families, Gekkonidae was the first to be described (Gray 1825), followed by Pygopodidae (as Pygopidae in Gray 1841). Boulenger (1883) recognized Eublepharidae based on differences in vertebral structure as compared to all other geckos, and was the first to note morphological




Figure 4. Species numbers among gecko genera. The 10 most speciose genera (listed) currently contain more than 1,000 species, or about 50% of all geckos, and about 10% of all reptiles. *Cyrtodactylus* is the most speciose genus of geckos, and the most species-rich reptilian genus after *Anolis*. For detailed numbers of smaller genera see the latest release of the Reptile Database and its regularly updated spreadsheet.

similarities between pygopodids and geckos (Boulenger 1884). Subsequent anatomical studies in the 20th century confirmed the status of pygopodids as gekkotans (e.g. Underwood 1957; Kluge 1974). The Carphodactylidae (as Carphodactylini), Diplodactylidae (as Diplodactylinae) and Sphaerodactylidae were described on the basis of anatomical traits (Underwood 1954; Kluge 1967), though for the remainder of the 20th century these groups were often treated as tribes or subfamilies of Gekkonidae and their content changed as new evidence emerged (reviewed by Russell and Bauer 2002). The contemporary seven family classification stems from molecular phylogenetic studies that further clarified the content of the major gekkotan clades and identified the family Phyllodactylidae (Gamble et al. 2008a, b; Han et al. 2004). Within each family there may be a substantial variation in morphological characters, e.g. most phyllodactylid genera can be distinguished by variation in the presence and shape of toe pads. The morphology of the digits (including toepads) and shape of the pupil have historically been the most important characters used in distinguishing gecko genera. More recent molecular work has shown that some of these character states have evolved multiple times and generic classification has been modified accordingly. For example, most leaf-toed geckos were originally placed in the genus *Phyllodactylus* until it was determined that they actually represented over a dozen distinct lineages, across four families (Kluge 1983; Bauer et al. 1997; Heinicke et al. 2014). Conversely, molecular data have also been used to subsume some genera that were previously recognized on the basis of digital morphology, such as the placement of *Colopus* and *Palmatogecko* in the synonymy of *Pachydactylus* (Heinicke et al. 2017). Although the generic and familial classification of geckos is now largely stable, there are still a handful of genera, such as *Cnemaspis* and *Sauromactylus*, that molecular data show to be polyphyletic (Gamble et al. 2012, but also see Javanmardi et al. 2019), implying that some taxonomic revision at the genus level is still required.

Gecko traits

There are no comprehensive databases collecting morphological and life history characters across all geckos, but some efforts have been made to collect body sizes (Meiri 2008; Feldman et al. 2015) and other trait data (Meiri 2018) of use for studying gecko evolution in a phylogenetic context. Some studies have identified morphological synapomorphies of clades using data sets containing hundreds of characters across multiple species belonging to multiple gecko lineages (e.g., Daza and Bauer 2012). Evolutionary patterns of many specific traits of geckos have also been studied. Examples include diurnal activity patterns (Gamble et al. 2015b), gliding adaptations (Heinicke et al. 2012), sex determining mechanisms (Gamble et al. 2015a), habitat-associated diversification and ecomorphology (e.g. Grismer et al. 2015; Heinicke et al. 2017; Oliver et al. 2019; Vidan et al. 2019), and perhaps most notably, digital morphology (Bauer 2019; Gamble et al. 2012; Russell and Gamble 2019). These studies often incorporate data sets comprising a significant fraction of

gecko diversity. For example, Gamble et al. (2012) collected morphological characters of hand and feet of 244 species of geckos representing 107 genera and mapped them to a phylogenetic tree. These authors found that the absence of adhesive toe pads to be the ancestral state for the extant Gekkota as a whole, and their data are consistent with independent origins and losses of adhesive toe pads in the Diplodactylidae, Sphaerodactylidae, Phyllodactylidae, and Gekkonidae, with a strong likelihood of multiple origins in the latter three families.

Geckos and their DNA

With DNA sequences being available for ~76% of all geckos (e.g., Meiri 2018), but only 63% of non-gekkotan reptiles, they are relatively well-studied phylogenetically. For most of these species existing sequence data consist only of a few genes or fragments thereof (most often, ND2, RAG1 and PDC), but broader sequence data sets are now becoming more common (e.g. Skipwith et al. 2019; Wood et al. 2019). More extensive or even complete genome sequences are necessary to address some biological questions. At present, genomes of only a few geckos have been completely sequenced though, including *Gekko japonicus* (Liu et al. 2015), *Paroedura picta* (Hara et al. 2018), and *Eublepharis macularius* (Xiong et al. 2016). Insights into the biology of geckos have begun to emerge from these genome sequences and other high-throughput sequencing projects. For instance, Liu et al. (2015) found specific gene families to be related to the formation of adhesive setae, nocturnal vision and tail regeneration, as well as the diversification of olfactory sensation. In particular, they found that the emergence of setae in geckos is correlated with the duplication and diversification of β -keratin genes.

Geckos of the world – a geographic survey

Geckos are not evenly distributed in the world (Fig. 5, Rösler 2017; Meiri 2019, this issue). Most species are found in the tropics, but geckos also occur in many subtropical and warm temperate regions, especially in arid environments, where they penetrate as far north as the Gobi Desert (*Alsophylax*, *Teratoscincus*) and as far south as Patagonia (*Homonota*). There is extensive regional variation in species richness even when comparing regions of similar latitude and climate. Geckos are most species-rich in the West Indies, southern and eastern Africa, Madagascar, the Middle East, South and Southeast Asia, and Australasia. At least these are the regions where most species have been discovered. The most gecko-rich countries, with more than 100 species each, are Australia (241 species), India (127), Madagascar (120), and Malaysia (104) (Table 4), although some smaller countries have very high species richness, e.g. New Caledonia with 44 species in a land area of only 18,576 km². When correcting for land area, countries outside the tropics have fewer geckos (Fig. 6). Even though tropical countries have more geckos, there is only a weak correlation of latitudes and species numbers, probably because of variation with area, and because tropical Latin American countries, and desert, North African




Figure 5. Number of gecko species per country. Geckos are concentrated in the areas surrounding the Indian Ocean. Compare to species richness map in Meiri (2020, this issue).

Table 4. The Top-10 most gecko-rich countries of the world, each with more than 65 species. Compare to Fig. 6.

Country	Species number
Australia	241
India	136
Madagascar	121
Malaysia	105
Indonesia	97
Vietnam	90
South Africa	85
Thailand	85
Namibia	70
Iran	69

countries, have relatively few geckos (Meiri 2020, this volume, and Fig. 5), but also probably due to an under-count of actual species diversity in the tropics (possibly with the exception of South America which has relatively few geckos). For example, of the ~270 gecko species described in the last 5 years, the vast majority occur in the tropics (and in Australia at tropical, sub-tropical and desert climates), suggesting that as new gecko species are described the proportion of recognized species occurring in the tropics will continue to increase. New descriptions will probably also increase the number of range-restricted species. Currently over 19% of gecko species are known only from their type localities (Meiri et al. 2018). This proportion includes many recently described species which often are discovered in limited habitats such as small islands or areas of exposed karst. The limited ranges of many gecko species also means that local communities are often not nearly as species-rich as country totals indicate. For example, 32 species of *Cyrtodactylus* are known from peninsular Malaysia, Singapore, and adjacent archipelagos, but only one to a few species occurs at any single locality (Grismer and Quah 2019).

The great age and relatively limited fossil record of geckos obscures some of the biogeographic history of the group. The oldest fossils that are unambiguous geckos are all from Eurasia (Daza et al. 2016). Nonetheless, biogeographic reconstructions indicate that geckos were probably also present on most Southern Hemisphere continents including Australia, Africa, and South America at the time of the breakup of Gondwana during the Mesozoic (Gamble et al. 2008a; Oliver and Sanders 2009). Subsequently, gecko lineages have colonized or re-colonized additional landmasses including oceanic islands via dispersal, often across wide barriers (e.g. Gamble et al. 2008b; Nielsen et al. 2011; Heinicke et al. 2011, 2014; Novosolov and Meiri 2013; Skipwith et al. 2016; Oliver et al. 2018). The old age of the clade and high dispersal abilities of geckos results in members of the families Eublepharidae, Gekkonidae, Phyllodactylidae, and Sphaerodactylidae, occurring across multiple continents. Also as a result of this history of dispersals, as many as four families of geckos may occur in sympatry. While the otherwise Australian family Diplodactylidae is also found on New Caledonia and New Zealand — probably due to dispersal after those three land masses split from each other — the Carphodactylidae is entirely restricted to Australia. Only two pygopodids (both species of *Lialis*) occur elsewhere — in nearby New Guinea.

In summary, discovery of geckos continues unabatedly, despite increasing threat from habitat destruction and possibly climate change. There is little indication that the rate of species description will decline soon. Based on past trends, new discoveries are especially likely to come from regions of warm climate, heterogeneous landscape, and limited previous attention from systematic herpetologists. Ironically, with the advent of next-generation sequencing, we will soon have the tools to understand the molecular basis of gecko diversity, both in terms of populations and traits, but possibly only once many species have gone extinct.




Figure 6. The larger a country is, the more geckos it has. As expected, countries outside the tropics (purple-blue) have fewer species (latitude of the geographical centroids of each country encoded by color), but the effect of area is stronger. Country sizes and species numbers are given on a log-10 scale. Bottom right: Russia (3 species).

Materials and methods

Species and author data were derived from the December 2019 version of the Reptile Database. Distribution data and species per country were derived from an updated version of (Roll et al. 2017), using ArcGIS. Latitudinal centroids and countries are from the country08 shapefile of ArcGIS (except South Sudan which was still missing from ArcGIS at the time of writing). Numbers of species were cross-checked with the Reptile Database and corrected if necessary by manual inspection.

Acknowledgments

The authors thank Glenn Shea and Herbert Rösler for critically reading the manuscript. MH is supported by the US National Science Foundation (DEB-1556585 and DEB 1657527).

References

Bauer, A.M. (2019). Gecko adhesion in space and time: a phylogenetic perspective on the scansorial success story. *Integrative and Comparative Biology* 59, pp. 117–130.

Bauer, A.M., Good, D.A., and Branch, W.R. (1997). The taxonomy of the southern African leaf-toed geckos (Squamata: Gekkonidae), with a review of Old World 'Phyllodactylus'. *Proceedings of the California Academy of Sciences* 49, pp. 447–497.

Bauer, A.M., and Lamb, T. (2005). Phylogenetic relationships of southern African geckos in the *Pachydactylus* group (Squamata: Gekkonidae). *African Journal of Herpetology* 54, pp. 105–129.

Beddome, R.H. (1870a). Descriptions of new reptiles from the Madras Presidency. *Madras Monthly Journal of Medical Science* 2, pp. 169–176. [not sure why the article which appeared first is cited as 1870b, when the citations appear in the text in the same citation bracket].

Beddome, R.H. (1870b). Descriptions of some new lizards from the Madras Presidency. *Madras Monthly Journal of Medical Science* 1, pp. 30–35.

Boulenger, G.A. (1883). Remarks on the *Nyctisaura*. *Annals and Magazine of Natural History* (Series 5) 12, p. 308.

Boulenger, G.A. (1884). Synopsis of the families of existing Lizards. *Annals and Magazine of Natural History* (Series 5) 14, pp. 117–122.

Boulenger, G.A. (1885). *Catalogue of the lizards in the British Museum (Natural History), Volume I. Geckonidae, Eublepharidae, Uroplatidae, Pygopodidae, Agamidae*. London: Trustees of the British Museum (Natural History).

Branch, W.R., Bauer, A.M., and Good, D.A. (1995). Species limits in the *Phyllodactylus lineatus* complex (Reptilia: Gekkonidae), with the elevation of two taxa to specific status and the description of two new species. *Journal of the Herpetological Association of Africa* 44, pp. 33–54.

Breitman, M.F., Domingos, F.M.C.B., Bagley, J.C., Wiederhecker, H.C., Ferrari, T.B., Cavalcante, V.H.G.L., Pereira, A.C., Abreu, T.L.S., Soares De-Lima, A.K., Moraes, C.J.S., Prette, A.C.H.d., Silva, I.P.M.C., Mello, R.d., Carvalho, G., Lima, T.M.d., Silva, A.A., Matias, C.A., Carvalho, G.C., Pantoja, J.A.L., Monteiro-Gomes, I., Pinheiro-Paschoal, I., Ferreira-Rodrigues, G., Talarico, A.V.C., Barreto-Lima, A.F., and Colli, G.R. (2018). A New Species of *Enyalius* (Squamata, Leiosauridae) Endemic to the Brazilian Cerrado. *Herpetologica* 74, pp. 355–369.

Constable, H., Guralnick, R., Wieczorek, J., Spencer, C., Peterson, A.T., and VertNet Steering Committee (2010). VertNet: a new model for biodiversity data sharing. *PLoS Biology* 8, p. e1000309.

Daudin, F.M. (1802). *Histoire Naturelle, Générale et Particulière des Reptiles*, Vol 4. Paris, F. Dufart.

Daza, J.D., and Bauer, A.M. (2012). A new amber-embedded sphaerodactyl gecko from Hispaniola, with comments on morphological synapomorphies of the Sphaerodactylidae. *Breviora* 529, pp. 1–28.

Daza, J.D., Bauer, A.M., and Snively, E.D. (2014). On the fossil record of the Gekkota. *The Anatomical Record* 297, pp. 433–462.

Daza, J.D., Stanley, E.L., Wagner, P., Bauer, A.M., and Grimaldi, D.A. (2016). Mid-Cretaceous amber fossils illuminate the past diversity of tropical lizards. *Science Advances* 2, p. e1501080.

Duméril, A.M.C., and Bibron, G. (1836). *Erpetologie Générale ou Histoire Naturelle Complete des Reptiles, Volume 3*. Paris: Librairie Encyclopédique de Roret.

Feldman, A., Pyron, R.A., Sabath, N., Mayrose, I., and Meiri, S. (2015). Body-sizes and diversification rates of lizards, snakes, amphisbaenians and the tuatara. *Global Ecology and Biogeography* 25, pp. 187–197.

Gamble, T., Bauer, A.M., Greenbaum, E., and Jackman, T.R. (2008a). Evidence for Gondwanan vicariance in an ancient clade of gecko lizards. *Journal of Biogeography* 35, pp. 88–104.

Gamble, T., Bauer, A.M., Greenbaum, E., and Jackman, T.R. (2008b). Out of the blue: a novel, trans-Atlantic clade of geckos (Gekkota, Squamata). *Zoologica Scripta* 37, 355–366.

Gamble, T., Colli, G., Rodrigues, M.T., Werneck, F.P., and Simons, A.M. (2011) Phylogeny and cryptic diversity in geckos (*Phyllopezus*; *Phyllodactylidae*; Gekkota) from South America's open biomes. *Molecular Phylogenetics and Evolution* 62, pp. 943–953.

Gamble, T., Coryell, J., Ezaz, T., Lynch, J., Scantlebury, D.P., and Zarkower, D. (2015a). Restriction site-associated DNA sequencing (RAD-seq) reveals an extraordinary number of transitions among gecko sex-determining systems. *Molecular Biology and Evolution* 32, pp. 1296–1309.

Gamble, T., Greenbaum, E., Jackman, T.R., and Bauer, A.M. (2015b). Into the light: diurnality has evolved multiple times in geckos. *Biological Journal of the Linnean Society* 115, pp. 896–910.

Gamble, T., Greenbaum, E., Jackman, T.R., Russell, A.P., and Bauer, A.M. (2012). Repeated origin and loss of adhesive toe-pads in geckos. *PLoS One* 7, p. e39429.

Gauthier, J.A., Kearney, M., Maisano, J.A., Rieppel, O., and Behlke, A.D. (2012). Assembling the squamate tree of life: perspectives from the phenotype and the fossil record. *Bulletin of the Peabody Museum of Natural History* 53, pp. 3–309.

Gray, J.E. (1825) A synopsis of the genera of reptiles and Amphiibia, with a description of some new species. *Annals of Philosophy* 10, pp. 193–217.

Gray, J.E. (1841). A catalogue of the species of reptiles and Amphiibia hitherto described as inhabiting Australia, with a description of some new species from Western Australia. In: G. Grey (ed.), *Journals of Two Expeditions of Discovery in Northwestern and Western Australia, during the years 1837, 38, and 39, under the authority of Her Majesty's Government*, London: T. and W. Boone, pp. 422–449.

Gray, J.E. (1845). *Catalogue of the specimens of lizards in the collection of the British Museum*. London: Trustees of the British Museum/Edward Newman.

Grismer, L.L., and Quah, E.S. (2019). An updated and annotated checklist of the lizards of Peninsular Malaysia, Singapore, and their adjacent archipelagos. *Zootaxa* 4545, pp. 230–248.

Grismer, L.L., Wood, P.L., Ahmad, A.B., Sumarli, A.S., Vazquez, J.J., Ismail, L.H., et al. (2014). A new species of insular Rock Gecko (Genus *Cnemaspis* Strauch, 1887) from the Bidong Archipelago, Terengganu, Peninsular Malaysia. *Zootaxa* 3755, pp. 447–456.

Grismer, L.L., Wood, P.L., Ngo, V.T., and Murdoch, M.L. (2015). The systematics and independent evolution of cave ecomorphology in distantly related clades of Bent-toed Geckos (Genus *Cyrtodactylus* Gray, 1827) from the Mekong Delta and islands in the Gulf of Thailand. *Zootaxa* 3980, pp. 106–126.

Han, D., Zhou, K., and Bauer, A.M. (2004). Phylogenetic relationships among gekkotan lizards inferred from C-mos nuclear DNA sequences and a new classification of the Gekkota. *Biological Journal of the Linnean Society* 83, pp. 353–368.

Hara, Y., Takeuchi, M., Kageyama, Y., Tatsumi, K., Hibi, M., Kiyonari, H., et al. (2018). Madagascar ground gecko genome analysis characterizes asymmetric fates of duplicated genes. *BMC Biology* 16, p. 40.

Hedges, S.B., Marin, J., Suleski, M., Paymer, M., and Kumar, S. (2015). Tree of life reveals clock-like speciation and diversification. *Molecular Biology and Evolution* 32, pp. 835–845.

Heinicke, M.P., Greenbaum, E., Jackman, T.R., and Bauer, A.M. (2011). Phylogeny of a trans-Wallacean radiation (Squamata, Gekkonidae, *Gehyra*) supports a single early colonization of Australia. *Zoologica Scripta* 40 (6), pp. 584–602.

Heinicke, M.P., Greenbaum, E., Jackman, T.R., Bauer, A.M., Heinicke, M.P., Daza, J.D., et al. (2012). Evolution of gliding in Southeast Asian geckos and other vertebrates is temporally congruent with dipterocarp forest development. *Biology Letters* 8, 994–997.

Heinicke, M.P., Daza, J.D., Greenbaum, E., Jackman, T.R., and Bauer, A.M. (2014). Phylogeny, taxonomy and biogeography of a circum-Indian Ocean clade of leaf-toed geckos (Reptilia: Gekkota), with a description of two new genera. *Systematics and Biodiversity* 12 (1), pp. 23–42.

Heinicke, M.P., Jackman, T.R., and Bauer, A.M. (2017). The measure of success: geographic isolation promotes diversification in *Pachydactylus* geckos. *BMC Evolutionary Biology* 17, p. 9.

Javanmardi, S., Vogler, S., and Joger, U. (2019). Phylogenetic differentiation and taxonomic consequences in the *Sauromactylus brosseti* species complex (Squamata: Sphaerodactylidae), with description of four new species. *Zootaxa* 4674, pp. 401–425.

King, M. (1982). Karyotypic Evolution in *Gehyra* (Gekkonidae: Reptilia) II. A New Species from the Alligator Rivers Region in Northern Australia. *Australian Journal of Zoology* 30, pp. 93–101.

Kluge, A.G. (1967). Higher taxonomic categories of gekkonid lizards and their evolution. *Bulletin of the American Museum of Natural History* 135, pp. 1–60..

Kluge, A.G. (1974). A taxonomic revision of the lizard family Pygopodidae. *Miscellaneous Publications, Museum of Zoology, University of Michigan* 147, pp. 1–221.

Kluge, A.G. (1983). Cladistic relationships among gekkonid lizards. *Copeia* 1983, pp. 465–475.

Linnaeus, C. (1758). *Systema naturae*, Volume 1. Stockholm: Laurentii Salvii.

Liu, Y., Zhou, Q., Wang, Y., Luo, L., Yang, J., Yang, L., et al. (2015). *Gekko japonicus* genome reveals evolution of adhesive toe pads and tail regeneration. *Nat Communications* 6, p. 10033.

Meiri, S. (2008). Evolution and ecology of lizard body sizes. *Global Ecology and Biogeography* 17, pp. 724–734.

Meiri, S. (2016). Small, rare and trendy: traits and biogeography of lizards described in the 21st century. *Journal of Zoology* 299, pp. 251–261.

Meiri, S. (2018). Traits of lizards of the world: Variation around a successful evolutionary design. *Global Ecology and Biogeography* 27, pp. 1168–1172.

Meiri, S. (2019) What geckos are – an ecological-biogeographic perspective. *Israel Journal of Ecology & Evolution*, this issue.

Minton, S.A., Anderson, S.C., and Anderson, J.A. (1970). Remarks on some geckos from southwest Asia, with descriptions of three new forms and a key to the genus *Tropiocolotes*. *Proceedings of the California Academy of Sciences* 37, pp. 333–362.

Murphy, R.W. (1974). A New Genus and Species of Eublepharine Gecko (Sauria, Gekkonidae) from Baja California, Mexico. *Proceedings of the California Academy of Sciences* 40, pp. 87–92.

Nguyen, S.N., Le, T.N., Tran, T.A., Orlov, N.L., Lathrop, A., Macculloch, R.D., et al. (2013). Phylogeny of the *Cyrtodactylus irregularis* species complex (Squamata: Gekkonidae) from

Vietnam with the description of two new species. *Zootaxa* 3737, pp. 399–414.

Nguyen, S.N., Yang, J.X., Le, T.N., Nguyen, L.T., Orlov, N.L., Hoang, C.V., et al. (2014). DNA barcoding of Vietnamese bent-toed geckos (Squamata: Gekkonidae: *Cyrtodactylus*) and the description of a new species. *Zootaxa* 3784, pp. 48–66.

Nielsen, S.V., Bauer, A.M., Jackman, T.R., Hitchmough, R.A., and Daugherty, C.H. (2011). New Zealand geckos (Diplodactylidae): cryptic diversity in a post-Gondwanan lineage with trans-Tasman affinities. *Molecular Phylogenetics and Evolution* 59, pp. 1–22.

Novosolov, M. and Meiri, S. (2013). The effect of island type on lizard reproductive traits. *J. Biogeog.*, 40, pp. 2385–2395.

Oliver, P.M., Ashman, L.G., Bank, S., Laver, R.J., Pratt, R.C., Tedeschi, L.G., et al. (2019). On and off the rocks: persistence and ecological diversification in a tropical Australian lizard radiation. *BMC Evolutionary Biology* 19, p. 81.

Oliver, P.M., Brown, R.M., Kraus, F., Rittmeyer, E., Travers, S.L., and Siler, C.D. (2018). Lizards of the lost arcs: mid-Cenozoic diversification, persistence and ecological marginalization in the West Pacific. *Proceedings of the Royal Society B: Biological Sciences* 285, p. 20171760.

Oliver, P.M., and Sanders, K.L. (2009). Molecular evidence for Gondwanan origins of multiple lineages within a diverse Australasian gecko radiation. *Journal of Biogeography* 36, pp. 2044–2055.

Roll, U., Feldman, A., Novosolov, M., Allison, A., Bauer, A.M., Bernard, R., et al. (2017). The global distribution of tetrapods reveals a need for targeted reptile conservation. *Nature Ecology and Evolution* 1, pp. 1677–1682.

Rösler, H. (2017). Gecko-Chorologie (Squamata: Gekkota). *Gekkota* Supplement 4, pp. 1–160.

Russell, A.P., and Bauer, A.M. (2002). Underwood's Classification of the Geckos: A 21st Century Appreciation. *Bulletin of the Natural History Museum London (Zoology)* 68, pp. 113–121.

Russell, A.P., and Gamble, T. (2019). Evolution of the Gekkotan Adhesive System: Does Digit Anatomy Point to One or More Origins? *Integrative and Comparative Biology* 59, pp. 131–147.

Simões, T.R., Pesteguía, S., Hsiou, A.S., and Daza, J.D. (2017). Lepidosaurs from Gondwana: An Introduction. *Journal of Herpetology* 51, pp. 297–299.

Skipwith, P.L., Bauer, A.M., Jackman, T.R., and Sadlier, R.A. (2016). Old but not ancient: coalescent species tree of New Caledonian geckos reveals recent post-inundation diversification. *Journal of Biogeography* 43, pp. 1266–1276.

Skipwith, P.L., Bi, K., and Oliver, P.M. (2019). Relicts and radiations: Phylogenomics of an Australasian lizard clade with east Gondwanan origins (Gekkota: Diplodactyloidea). *Molecular Phylogenetics and Evolution* 140, p. 106589.

Sparrman, A. (1778). Befkrifning och berättelse om LACERTA GEITJE, en obekant och giftig ÖDLA ifrån Goda Hoppa Udden. *Det Götheborgska Wetenskaps och Witterhets Samhällets Handlingar Wetenskaps Afdelningen Forsta Stycket*, Götheborg, pp. 75–78.

Spix, J.B.v. (1825). *Animalia nova sive species nova lacertarum quas in itinere per Brasiliam annis MDCCCXVII–MDCCCXX jussu et auspicio Maximiliani Josephi I Bavariae Regis suscepto collegit et descripsit Dr. J. B. de Spix. Lipsiae: T. O. Weigel; F. S. Hübschmanni, Monachii.*

Steindachner, F. (1901). Berichte der Comission für Oceanographische Forschungen. Expedition S. M. Schiff "Pola" in das Rothe Meer nördliche und südliche Hälften. 1895/96 und 1897/98. Zoologische Ergebnisse. XVII. Bericht über die herpetologischen Aufsammlungen. *Denkschriften der Kaiserlichen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Classe*, Wien 69, pp. 325–335.

Uetz, P., Cherikh, S., Shea, G., Ineich, I., Campbell, P.D., Doronin, I.V., et al. (2019). A global catalog of primary reptile type specimens. *Zootaxa* 4695, pp. 438–450.

Uetz, P., Freed, P., and Hošek, J. (2019). Reptile Database. <http://www.reptile-database.org>, accessed 15 Jan 2020.

Uetz, P., and Stylianou, A. (2018). The original descriptions of reptiles and their subspecies. *Zootaxa* 4375, pp. 257–264.

Underwood, G. (1954). On the classification and evolution of geckos. *Proceedings of the Zoological Society of London* 124, pp. 469–492.

Underwood, G. (1957). On lizards of the family Pygopodidae. A contribution to the morphology and phylogeny of the Squamata. *Journal of Morphology* 100, pp. 207–268.

Vidan, E., Bauer, A.M., Herrera, F.-C., Chirio, L., Nogueira, C.C., Doan, T.M., et al. (2019). The global biogeography of lizard functional groups. *Journal of Biogeography* 46, pp. 2147–2158.

Wood, P.L., Guo, X., Travers, S.L., Su, Y.C., Olson, K.V., Bauer, A.M., et al. (2019). Parachute geckos free fall into synonymy: *Gekko* phylogeny, and a new subgeneric classification, inferred from thousands of ultraconserved elements. *bioRxiv* p. 717520.

Xiong, Z., Li, F., Li, Q., Zhou, L., Gamble, T., Zheng, J., et al. (2016). Draft genome of the leopard gecko, *Eublepharis macularius*. *Gigascience* 5, p. 47.

Zheng, Y., and Wiens, J.J. (2016). Combining phylogenomic and supermatrix approaches, and a time-calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes and 4162 species. *Molecular Phylogenetics and Evolution* 94, pp. 537–547.