



Computation of Chebyshev Polynomials for Union of Intervals

Simon Foucart¹ · Jean Bernard Lasserre²

Received: 30 June 2018 / Revised: 9 March 2019 / Accepted: 20 June 2019 / Published online: 12 August 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Chebyshev polynomials of the first and second kind for a set K are monic polynomials with minimal L_∞ - and L_1 -norm on K , respectively. This article presents numerical procedures based on semidefinite programming to compute these polynomials in case K is a finite union of compact intervals. For Chebyshev polynomials of the first kind, the procedure makes use of a characterization of polynomial non-negativity. It can incorporate additional constraints, e.g. that all the roots of the polynomial lie in K . For Chebyshev polynomials of the second kind, the procedure exploits the method of moments.

Keywords Chebyshev polynomials of the first kind · Chebyshev polynomials of the second kind · Non-negative polynomials · Method of moments · Semidefinite programming

Mathematics Subject Classification 31A15 · 41A50 · 90C22

1 Introduction

The N th Chebyshev polynomial for a compact infinite subset K of \mathbb{C} is defined as the monic polynomial of degree N with minimal max-norm on K . Its uniqueness is a straightforward consequence of the uniqueness of best polynomial approximants to a

Communicated by Thomas Ransford.

The research of Simon Foucart is partially funded by the NSF Grants DMS-1622134 and DMS-1664803. The research of Jean Bernard Lasserre is funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant agreement 666981 TAMING).

✉ Simon Foucart
foucart@tamu.edu

¹ Texas A&M University, College Station, USA

² LAAS-CNRS, Toulouse, France

continuous function (here $z \mapsto z^N$) with respect to the max-norm, see e.g. [4, p. 72, Thm. 4.2]. We shall denote it as \mathcal{T}_N^K , i.e.

$$\mathcal{T}_N^K = \operatorname{argmin}_{P(z)=z^N+\dots} \|P\|_K, \quad \text{where } \|P\|_K = \max_{z \in K} |P(z)|. \quad (1)$$

We reserve the notation T_N^K for the Chebyshev polynomial normalized to have max-norm equal to one on K , i.e.

$$T_N^K = \frac{\mathcal{T}_N^K}{\|\mathcal{T}_N^K\|_K}. \quad (2)$$

With this notation, the usual N th Chebyshev polynomial (of the first kind) satisfies

$$T_N = T_N^{[-1,1]} = 2^{N-1} T_N^{[-1,1]}, \quad N \geq 1. \quad (3)$$

Chebyshev polynomials for a compact subset K of \mathbb{C} play an important role in logarithmic potential theory. For instance, it is known that the capacity $\operatorname{cap}(K)$ of K is related to the Chebyshev numbers $t_N^K := \|\mathcal{T}_N^K\|_K$ via

$$\left(t_N^K\right)^{1/N} \xrightarrow[N \rightarrow \infty]{} \operatorname{cap}(K), \quad (4)$$

see [12, p.163, Thm. 3.1] for a weighted version of this statement. Some recent works [1,2] have studied in greater detail the asymptotics of the convergence (4) in case K is a subset of \mathbb{R} . This being said, the capacity is in general hard to determine—it can be found explicitly in a few specific situations, e.g. when K is the inverse image of an interval by certain polynomials (see [8, Thm. 11]) and otherwise some numerical methods for computing the capacity have been proposed in [11], see also [10, Sect. 5.2]. As for the Chebyshev polynomials, one is tempted to anticipate a worse state of affairs. However, this is not the case for the situation considered in present work, i.e. when $K \subseteq [-1, 1]$ is a finite union of L compact intervals,¹ say

$$K = \bigcup_{\ell=1}^L [a_\ell, b_\ell], \quad -1 = a_1 < b_1 < a_2 < b_2 < \dots < a_L < b_L = 1. \quad (5)$$

There are explicit constructions of Chebyshev polynomials (as orthogonal polynomials with a predetermined weight, see [9, Thm. 2.3]), albeit only under the condition that \mathcal{T}_N^K is a *strict* Chebyshev polynomial (meaning that it possesses $N + L$ points of equioscillation on K —a condition which is verifiable a priori, see [9, Thm. 2.5]). Chebyshev polynomials can otherwise be computed using Remez-type algorithms for finite unions of intervals, see [5].

¹ The assumption $K \subseteq [-1, 1]$ is not restrictive, as any compact subset of \mathbb{R} can be moved into the interval $[-1, 1]$ by an affine transformation.

A first contribution of this article is to put forward an alternative numerical procedure that enables the accurate computation of the Chebyshev polynomials whenever K is a finite union of compact intervals. The procedure, based on semidefinite programming as described in Sect. 2, can also incorporate a weight w (i.e. a continuous and positive function on K), restricted here to be a rational function, and give as output the polynomials

$$\mathcal{T}_N^{K,w} = \operatorname{argmin}_{P(x)=x^N+\dots} \left\| \frac{P}{w} \right\|_K. \quad (6)$$

An appealing feature of this approach is that extra constraints can easily be incorporated into the minimization of (6). For instance, we will show how to compute the N th restricted Chebyshev polynomial on K , i.e. the monic polynomial of degree N having all its roots in K with minimal max-norm on K .

A second contribution of this study is to propose another semidefinite-programming-based procedure to compute weighted Chebyshev polynomials of the second kind, so to speak. By this, we mean polynomials²

$$\mathcal{U}_N^{K,w} \in \operatorname{argmin}_{P(x) \in x^N + \dots} \left\| \frac{P}{w} \right\|_{L_1(K)}. \quad (7)$$

The restriction that the weight w is a rational function is not needed here, but this time the computation is only approximate. Nonetheless, it produces lower and upper bounds for the genuine minimum $\|\mathcal{U}_N^{K,w}/w\|_{L_1(K)}$. Both bounds are shown to converge to the genuine minimum as a parameter $d \geq N$ grows to infinity. Along the way, we shall prove that the Chebyshev polynomial of the second kind for K , if unique, has simple roots all lying inside K .

The procedures for computing Chebyshev polynomials of the first and second kind have been implemented in MATLAB. They rely on the external packages CVX (for specifying and solving convex programs [3]) and Chebfun (for numerically computing with functions [13]). They can be downloaded from the first author's webpage as part of the reproducible file accompanying this article.

2 Chebyshev Polynomials of the First Kind

With K as in (5), we consider a rational³ weight function w taking the form

$$w = \frac{\Sigma}{\Omega}, \quad (8)$$

² The uniqueness of $\mathcal{U}_N^{K,w}$ is not necessarily guaranteed: in the unweighted case, one can, e.g. check that the monic linear polynomials with minimal L_1 -norm on $K = [-1, -c] \cup [c, 1]$ are all the $x - d$, d in the interval $[-c, c]$. We will not delve into conditions ensuring uniqueness of $\mathcal{U}_N^{K,w}$ in this paper.

³ We could also work with piecewise rational weight functions, but we choose not to do so in order to avoid overloading already heavy notation.

where the polynomials Σ and Ω are positive on each $[a_\ell, b_\ell]$. We shall represent polynomials P of degree at most N by their Chebyshev expansions written as

$$P = \sum_{n=0}^N p_n T_n. \quad (9)$$

In this way, finding the N th Chebyshev polynomial of the first kind for K with weight w amounts to solving the optimization problem

$$\underset{p_0, p_1, \dots, p_N \in \mathbb{R}}{\text{minimize}} \max_{\ell=1:L} \left\| \frac{\Omega P}{\Sigma} \right\|_{[a_\ell, b_\ell]} \quad \text{s.to} \quad p_N = \frac{1}{2^{N-1}}. \quad (10)$$

After introducing a slack variable $c \in \mathbb{R}$, this is equivalent to the optimization problem

$$\underset{c, p_0, p_1, \dots, p_N \in \mathbb{R}}{\text{minimize}} \quad c \quad \text{s.to} \quad p_N = \frac{1}{2^{N-1}} \quad \text{and} \quad \left\| \frac{\Omega P}{\Sigma} \right\|_{[a_\ell, b_\ell]} \leq c \quad \text{for all } \ell = 1 : L. \quad (11)$$

The latter constraints can be rewritten as $-c \leq \Omega P / \Sigma \leq c$ on $[a_\ell, b_\ell]$, $\ell = 1 : L$, i.e. as the two polynomial non-negativity constraints

$$c \Sigma(x) \pm \Omega(x) P(x) \geq 0 \quad \text{for all } x \in [a_\ell, b_\ell] \quad \text{and all } \ell = 1 : L. \quad (12)$$

The key to the argument is now to exploit an exact semidefinite characterization of these constraints. This is based on the following result, which was established and utilized in [7], see Theorem 3 there.

Proposition 1 *Given $[a, b] \subseteq [-1, 1]$ and a polynomial $C(x) = \sum_{m=0}^M c_m T_m(x)$ of degree at most M , the non-negativity condition*

$$C(x) \geq 0 \quad \text{for all } x \in [a, b] \quad (13)$$

is equivalent to the existence of semidefinite matrices $\mathbf{Q} \in \mathbb{C}^{(M+1) \times (M+1)}$, $\mathbf{R} \in \mathbb{C}^{M \times M}$ such that

$$\begin{aligned} & \sum_{i-j=m} Q_{i,j} + \alpha \sum_{i-j=m-1} R_{i,j} - \beta \sum_{i-j=m} R_{i,j} + \bar{\alpha} \sum_{i-j=m+1} R_{i,j} \\ &= \begin{cases} \frac{1}{2} c_m, & m = 1 : M \\ c_0, & m = 0 \end{cases}, \end{aligned} \quad (14)$$

where $\alpha = \frac{1}{2} \exp \left(\frac{t}{2} \arccos(a) + \frac{t}{2} \arccos(b) \right)$ and $\beta = \cos \left(\frac{1}{2} \arccos(a) - \frac{1}{2} \arccos(b) \right)$.

In the present situation, we apply this result to the polynomials $C = c\Sigma \pm \Omega P$ required to be non-negative on each $[a_\ell, b_\ell]$. With

$$M := \max \{\deg(\Sigma), \deg(\Omega) + N\}, \quad (15)$$

we write the Chebyshev expansions of Σ and of ΩP as

$$\Sigma = \sum_{m=0}^M \sigma_m T_m, \quad \Omega P = \sum_{m=0}^M (\mathbf{Wp})_m T_m, \quad (16)$$

where $\mathbf{W} \in \mathbb{R}^{(M+1) \times (N+1)}$ is the matrix of the linear map transforming the Chebyshev coefficients of P into the Chebyshev coefficients of ΩP . Our considerations can now be summarized as follows.

Theorem 2 *The N th Chebyshev polynomial $T_N^{K,w}$ for the set K given in (5) and with weight w given in (8) has Chebyshev coefficients p_0, p_1, \dots, p_N that solve the semidefinite program*

$$\begin{aligned} & \underset{\substack{c, p_0, p_1, \dots, p_N \in \mathbb{R} \\ \mathbf{Q}^{\pm, \ell} \in \mathbb{R}^{(M+1) \times (M+1)} \\ \mathbf{R}^{\pm, \ell} \in \mathbb{R}^{M \times M}}}{\text{minimize}} \quad c \quad \text{s.to} \quad p_N = \frac{1}{2^{N-1}}, \quad \mathbf{Q}^{\pm, \ell} \succeq \mathbf{0}, \quad \mathbf{R}^{\pm, \ell} \succeq \mathbf{0}, \\ & \text{and} \quad \sum_{i-j=m} Q_{i,j}^{\pm, \ell} + \alpha_\ell \sum_{i-j=m-1} R_{i,j}^{\pm, \ell} - \beta_\ell \sum_{i-j=m} R_{i,j}^{\pm, \ell} + \overline{\alpha_\ell} \sum_{i-j=m+1} R_{i,j}^{\pm, \ell} \\ & = \begin{cases} \frac{1}{2} \sigma_m c \pm \frac{1}{2} (\mathbf{Wp})_m, & m = 1 : M \\ \sigma_0 c \pm (\mathbf{Wp})_0, & m = 0 \end{cases}, \end{aligned} \quad (17)$$

where $\alpha_\ell = \frac{1}{2} \exp \left(\frac{i}{2} \arccos(a_\ell) + \frac{i}{2} \arccos(b_\ell) \right)$ and $\beta_\ell = \cos \left(\frac{1}{2} \arccos(a_\ell) - \frac{1}{2} \arccos(b_\ell) \right)$.

Figure 1 provides examples of Chebyshev polynomials of degree $N = 5$ for the union of $L = 3$ intervals which were computed by solving (17). In all cases, the Chebyshev polynomials equioscillate $N + 1 = 6$ times between $-w$ and $+w$ on K , as they should. However, they are not strict Chebyshev polynomials, since the number of equioscillation points on K is smaller than $N + L = 8$. We notice in (c) and (d) that some roots of the Chebyshev polynomials do not lie in the set K . We display in (e) and (f) the restricted Chebyshev polynomial for K , i.e. the monic polynomial of degree N with minimal max-norm on K which satisfies the additional constraint that all its roots lie in K . This constraint reads

$$P \text{ does not vanish on } (b_\ell, a_{\ell+1}), \quad \ell = 1 : L - 1. \quad (18)$$

We consider the semidefinite program (17) supplemented with the relaxed constraint

$$P \text{ does not change sign on } [b_\ell, a_{\ell+1}], \quad \ell = 1 : L - 1. \quad (19)$$

This is solved by selecting the smallest value (along with the corresponding minimizer) among the minima of 2^{L-1} semidefinite programs (17) indexed by $(\varepsilon_1, \dots, \varepsilon_{L-1}) \in \{\pm 1\}^{L-1}$, where the added constraint is the semidefinite characterization of the polynomial non-negativity condition

$$\varepsilon_\ell P(x) \geq 0 \quad \text{for all } x \in [b_\ell, a_{\ell+1}] \quad \text{and all } \ell = 1 : L - 1. \quad (20)$$

One checks whether the selected minimizer satisfies the original constraint (18). If it does, then the restricted Chebyshev polynomial has indeed been found, as in (e) and (f) of Fig. 1.

Remark Concerning the computation of the capacity of a union of intervals, we do not recommend using our semidefinite procedure or a Remez-type procedure to produce Chebyshev polynomials before invoking (4) to approximate the capacity. If one really wants to take such a route, it seems wiser to work with the numerically friendlier orthogonal polynomials

$$\mathcal{P}_N^K = \underset{P(x)=x^N+\dots}{\operatorname{argmin}} \|P\|_{L_2(K)}. \quad (21)$$

Indeed, we also have

$$\|\mathcal{P}_N^K\|_{L_2(K)}^{1/N} \xrightarrow[N \rightarrow \infty]{} \operatorname{cap}(K), \quad (22)$$

as a consequence of the inequalities

$$\begin{aligned} \frac{1}{N+1} \left[\min_{\ell=1:L} (b_\ell - a_\ell) \right]^{1/2} \|\mathcal{T}_N^K\|_K &\leq \|\mathcal{P}_N^K\|_{L_2(K)} \\ &\leq \left[\sum_{\ell=1}^L (b_\ell - a_\ell) \right]^{1/2} \|\mathcal{T}_N^K\|_K. \end{aligned} \quad (23)$$

3 Chebyshev Polynomials of the Second Kind

Still with K as in (5), but with an arbitrary (positive and continuous) weight function w , we are now targeting N th Chebyshev polynomials of the second kind for K with weight w , i.e.

$$\mathcal{U}_N^{K,w} \in \underset{P(x)=x^N+\dots}{\operatorname{argmin}} \left\| \frac{P}{w} \right\|_{L_1(K)}, \quad \text{where} \quad \left\| \frac{P}{w} \right\|_{L_1(K)} = \sum_{\ell=1}^L \int_{a_\ell}^{b_\ell} \frac{|P(t)|}{w(t)} dt. \quad (24)$$




Fig. 1 5th Chebyshev polynomials of the first kind for $K_1 = [-1, -\frac{1}{2}] \cup [-\frac{1}{5}, \frac{1}{5}] \cup [\frac{1}{2}, 1]$ and for $K_2 = [-1, -\frac{1}{2}] \cup [\frac{1}{10}, \frac{1}{5}] \cup [\frac{2}{3}, 1]$: the first two rows correspond to the unrestricted case, while restricted Chebyshev polynomials are shown in the last row; the first column corresponds to the unweighted case, while weighted Chebyshev polynomials with weight $w(x) = (1 + x^2)/(2 - x^2)$ are shown in the second column

Let us drop the superscript w and simply write \mathcal{U}_N^K for $\mathcal{U}_N^{K,w}$. Minimizing the L_1 -norm on K exactly seems out of reach, so instead we shall perform the minimization of a more tractable ersatz norm, which will be formally defined in Proposition 4. This

ersatz norm stems from a reformulation of the L_1 -norm on K , as described in the steps below. Given a polynomial P of degree at most N , we start by making two changes of variables to write

$$\begin{aligned} \left\| \frac{P}{w} \right\|_{L_1(K)} &= \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} \int_{-1}^1 \frac{|P_\ell(x)|}{w_\ell(x)} dx \\ &= \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} \int_0^\pi |P_\ell(\cos(\theta))| \frac{\sin(\theta)d\theta}{w_\ell(\cos(\theta))}, \end{aligned} \quad (25)$$

where P_ℓ and w_ℓ denote the functions $P_{|[a_\ell, b_\ell]}$ and $w_{|[a_\ell, b_\ell]}$ transplanted to $[-1, 1]$, for instance

$$P_\ell(x) = P \left(\frac{(b_\ell - a_\ell)x + a_\ell + b_\ell}{2} \right), \quad x \in [-1, 1]. \quad (26)$$

We continue by decomposing the signed measures $P_\ell(\cos(\theta)) \sin(\theta)/w_\ell(\cos(\theta))d\theta$ as differences of two non-negative measures, so that

$$\begin{aligned} \left\| \frac{P}{w} \right\|_{L_1(K)} &= \inf_{\mu_1^\pm, \dots, \mu_L^\pm} \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} \int_0^\pi d(\mu_\ell^+ + \mu_\ell^-) \text{ s.to } d(\mu_\ell^+ - \mu_\ell^-)(\theta) \\ &= P_\ell(\cos(\theta)) \frac{\sin(\theta)d\theta}{w_\ell(\cos(\theta))}, \end{aligned} \quad (27)$$

where the infimum is taken over all non-negative measures on $[0, \pi]$. As is well known, a minimization over non-negative measures can be reformulated as a minimization over their sequences of moments. There are several options to do so: here, emulating an approach already exploited in [6], see Section 3 there, we rely on the discrete trigonometric moment problem encapsulated in the following statement.

Proposition 3 *Given a sequence $\mathbf{y} \in \mathbb{R}^\mathbb{N}$, there exists a non-negative measure μ on $[0, \pi]$ such that*

$$\int_0^\pi \cos(k\theta) d\mu(\theta) = y_k, \quad k \geq 0, \quad (28)$$

if and only if the infinite Toeplitz matrix build from \mathbf{y} is positive semidefinite, i.e.

$$\mathbf{Toep}_\infty(\mathbf{y}) := \begin{bmatrix} y_0 & y_1 & y_2 & y_3 & \cdots \\ y_1 & y_0 & y_1 & y_2 & \\ y_2 & y_1 & y_0 & y_1 & \ddots \\ y_3 & y_2 & y_1 & \ddots & \ddots \\ \vdots & & \ddots & \ddots & \ddots \end{bmatrix} \succeq 0. \quad (29)$$

The latter means that all the finite sections of $\mathbf{Toep}_\infty(\mathbf{y})$ are positive semidefinite, i.e.

$$\mathbf{Toep}_d(\mathbf{y}) = \begin{bmatrix} y_0 & y_1 & \cdots & \cdots & y_d \\ y_1 & y_0 & y_1 & & \vdots \\ \vdots & y_1 & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & y_1 \\ y_d & \cdots & \cdots & y_1 & y_0 \end{bmatrix} \succeq 0 \quad \text{for all } d \geq 0. \quad (30)$$

With $\mathbf{y}^{1,\pm}, \dots, \mathbf{y}^{L,\pm} \in \mathbb{R}^{\mathbb{N}}$ representing the sequences of moments of $\mu_1^\pm, \dots, \mu_L^\pm$, the objective function in (27) just reads

$$\sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} \int_0^\pi d(\mu_\ell^+ + \mu_\ell^-) = \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+} + y_0^{\ell,-}).$$

As for the constraints in (27), with $\mathbf{W}^\ell \in \mathbb{R}^{(N+1) \times (N+1)}$ denoting the matrix of the linear map transforming the Chebyshev coefficients of P into the Chebyshev coefficients of the second kind of P_ℓ , so that

$$P_\ell = \sum_{n=0}^N (\mathbf{W}^\ell \mathbf{p})_n U_n, \quad (31)$$

they become, for all $\ell = 1 : L$ and all $k \geq 0$,

$$y_k^{\ell,+} - y_k^{\ell,-} = \sum_{n=0}^N (\mathbf{W}^\ell \mathbf{p})_n \int_0^\pi \cos(k\theta) U_n(\cos(\theta)) \frac{\sin(\theta) d\theta}{w_\ell(\cos(\theta))} = (\mathbf{J}^\ell \mathbf{W}^\ell \mathbf{p})_k, \quad (32)$$

where the infinite matrices $\mathbf{J}^\ell \in \mathbb{R}^{\mathbb{N} \times (N+1)}$ have entries

$$J_{k,n}^\ell = \int_0^\pi \frac{\cos(k\theta) \sin((n+1)\theta)}{w_\ell(\cos(\theta))} d\theta. \quad (33)$$

The finite matrices $\mathbf{J}^{\ell,d} \in \mathbb{R}^{(d+1) \times (N+1)}$, obtained by keeping the first $d+1$ rows of \mathbf{J}^ℓ , are to be precomputed numerically and can sometimes even be determined explicitly, e.g.

$$\text{when } w = 1, \quad J_{k,n}^{\ell,d} = \begin{cases} 0 & \text{if } k \text{ and } n \text{ have different parities,} \\ \frac{2(n+1)}{(n+1)^2 - k^2} & \text{if } k \text{ and } n \text{ have similar parities.} \end{cases} \quad (34)$$

Taking into account the constraints that the $\mathbf{y}^{\ell,\pm} \in \mathbb{R}^{\mathbb{N}}$ must be sequences of moments, we arrive at a semidefinite reformulation of the weighted L_1 -norm on K

given by

$$\left\| \frac{P}{w} \right\|_{L_1(K)} = \inf_{\mathbf{y}^{1,\pm}, \dots, \mathbf{y}^{L,\pm} \in \mathbb{R}^N} \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+} + y_0^{\ell,-})$$

s.to $\mathbf{y}^{\ell,+} - \mathbf{y}^{\ell,-} = \mathbf{J}^\ell \mathbf{W}^\ell \mathbf{p}$ and $\mathbf{Toep}_\infty(\mathbf{y}^{\ell,\pm}) \succeq \mathbf{0}$. (35)

This expression is not tractable due to the infinite dimensionality of the optimization variables and constraints, but truncating them to a level d leads to a tractable expression—the above-mentioned ersatz norm.

Proposition 4 *For each $d \geq N$, the expression*

$$\|P\|_d := \min_{\mathbf{y}^{1,\pm}, \dots, \mathbf{y}^{L,\pm} \in \mathbb{R}^{d+1}} \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+} + y_0^{\ell,-})$$

s.to $\mathbf{y}^{\ell,+} - \mathbf{y}^{\ell,-} = \mathbf{J}^{\ell,d} \mathbf{W}^\ell \mathbf{p}$ and $\mathbf{Toep}_d(\mathbf{y}^{\ell,\pm}) \succeq \mathbf{0}$ (36)

defines a norm on the space of polynomials of degree at most N . Moreover, one has

$$\dots \leq \|P\|_d \leq \|P\|_{d+1} \leq \dots \leq \left\| \frac{P}{w} \right\|_{L_1(K)} \quad \text{and} \quad \lim_{d \rightarrow \infty} \|P\|_d = \left\| \frac{P}{w} \right\|_{L_1(K)}. \quad (37)$$

Proof To justify that the expression in (36) defines a norm, we concentrate on the property $[\|P\|_d = 0] \implies [P = 0]$, as the other two norm properties are fairly clear. So, assuming that $\|P\|_d = 0$, there exist $\mathbf{y}^{1,\pm}, \dots, \mathbf{y}^{L,\pm} \in \mathbb{R}^{d+1}$ such that

$$\sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+} + y_0^{\ell,-}) = 0, \quad (38)$$

as well as, for all $\ell = 1 : L$,

$$\mathbf{y}^{\ell,+} - \mathbf{y}^{\ell,-} = \mathbf{J}^{\ell,d} \mathbf{W}^\ell \mathbf{p} \quad \text{and} \quad \mathbf{Toep}_d(\mathbf{y}^{\ell,\pm}) \succeq \mathbf{0}. \quad (39)$$

The semidefiniteness of the Toeplitz matrices implies that

$$|y_k^{\ell,\pm}| \leq y_0^{\ell,\pm} \quad \text{for all } k = 0 : d, \quad (40)$$

which, in view of (38), yields $\mathbf{y}^{\ell,\pm} = \mathbf{0}$. By the invertibility of the matrices \mathbf{W}^ℓ and the injectivity of the matrices $\mathbf{J}^{\ell,d}$ (easy to check from (33)), we derive that $\mathbf{p} = \mathbf{0}$, and in turn that $P = 0$, as desired.

Let us turn to the justification of (37). The chain of inequalities translates to the fact that the successive minimizations impose more and more constraints and, hence, produce larger and larger minima. It remains to prove that the limit of the sequence $(\|P\|_d)_{d \geq N}$ equals $\|P/w\|_{L_1(K)}$ (the limit exists, because the sequence is

non-decreasing and bounded above). For each $d \geq N$, as was done in (38) and (39), we consider minimizers of the problem (35)—they belong to \mathbb{R}^{d+1} but we pad them with zeros to create infinite sequences $\mathbf{y}^{1,\pm,d}, \dots, \mathbf{y}^{L,\pm,d}$ satisfying

$$\sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+d} + y_0^{\ell,-d}) = \|P\|_d, \quad (41)$$

as well as, for all $\ell = 1 : L$,

$$\mathbf{y}^{\ell,+d} - \mathbf{y}^{\ell,-d} = \mathbf{J}^\ell \mathbf{W}^\ell \mathbf{p} \quad \text{and} \quad \mathbf{Toep}_\infty(\mathbf{y}^{\ell,\pm,d}) \succeq \mathbf{0}. \quad (42)$$

The semidefiniteness of the Toeplitz matrices, together with (41), implies that, for all $k \geq 0$,

$$|y_k^{\ell,\pm,d}| \leq y_0^{\ell,\pm,d} \leq \frac{2}{b_\ell - a_\ell} \|P\|_d \leq \frac{2}{b_\ell - a_\ell} \left\| \frac{P}{w} \right\|_{L_1(K)}. \quad (43)$$

In other words, each sequence $(\mathbf{y}^{\ell,\pm,d})_{d \geq N}$, with entries in the sequence space ℓ_∞ , is bounded. The sequential compactness Banach–Alaoglu theorem guarantees the existence of convergent subsequences in the weak-star topology. With $(\mathbf{y}^{\ell,\pm,d_m})_{m \geq 0}$ denoting these subsequences and $\mathbf{y}^{\ell,\pm} \in \ell_\infty$ denoting their limits, the weak-star convergence implies that

$$y_k^{\ell,\pm,d_m} \xrightarrow[m \rightarrow \infty]{} y_k^{\ell,\pm} \quad \text{for all } k \geq 0. \quad (44)$$

Writing (42) for $d = d_m$ and passing to the limit reveals that the sequences $\mathbf{y}^{1,\pm}, \dots, \mathbf{y}^{L,\pm}$ are feasible for the problem (35). Hence

$$\begin{aligned} \left\| \frac{P}{w} \right\|_{L_1} &\leq \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+} + y_0^{\ell,-}) = \lim_{m \rightarrow \infty} \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+d_m} + y_0^{\ell,-d_m}) \\ &= \lim_{m \rightarrow \infty} \|P\|_{d_m} = \lim_{d \rightarrow \infty} \|P\|_d, \end{aligned} \quad (45)$$

where the last equality relied on the fact that the non-decreasing and bounded sequence $(\|P\|_d)_{d \geq N}$ is convergent. This concludes the justification of (37). \square

Given $d \geq N$, let us now consider ersatz N th Chebyshev polynomials of the second kind for K (a priori not guaranteed to be unique) defined by

$$\mathcal{V}_{N,d}^K \in \operatorname{argmin}_{P(x)=x^N+\dots} \|P\|_d. \quad (46)$$

It is possible to compute such a polynomial by solving the following semidefinite program:

$$\begin{aligned} & \underset{\substack{p_0, p_1, \dots, p_N \in \mathbb{R} \\ \mathbf{y}^{1,\pm}, \dots, \mathbf{y}^{L,\pm} \in \mathbb{R}^{d+1}}}{\text{minimize}} \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+} + y_0^{\ell,-}) \\ & \text{s.to } p_N = \frac{1}{2^{N-1}}, \quad \mathbf{y}^{\ell,+} - \mathbf{y}^{\ell,-} = \mathbf{J}^{\ell,d} \mathbf{W}^\ell \mathbf{p}, \quad \text{and} \quad \mathbf{Toep}_d(\mathbf{y}^{\ell,\pm}) \succeq \mathbf{0}. \end{aligned} \quad (47)$$

The qualitative result below ensures that, as d increases, the ersatz Chebyshev polynomials $\mathcal{V}_{N,d}^K$ approach genuine Chebyshev polynomials \mathcal{U}_N^K , which are themselves obtained by solving the following (unpractical) semidefinite program:

$$\begin{aligned} & \underset{\substack{p_0, p_1, \dots, p_N \in \mathbb{R} \\ \mathbf{y}^{1,\pm}, \dots, \mathbf{y}^{L,\pm} \in \mathbb{R}^N}}{\text{minimize}} \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+} + y_0^{\ell,-}) \\ & \text{s.to } p_N = \frac{1}{2^{N-1}}, \quad \mathbf{y}^{\ell,+} - \mathbf{y}^{\ell,-} = \mathbf{J}^\ell \mathbf{W}^\ell \mathbf{p}, \quad \text{and} \quad \mathbf{Toep}_\infty(\mathbf{y}^{\ell,\pm}) \succeq \mathbf{0}. \end{aligned} \quad (48)$$

Theorem 5 Any sequence $(\mathcal{V}_{N,d}^K)_{d \geq N}$ of minimizers of (46) admits a subsequence converging (with respect to any of the equivalent norms on the space of polynomials of degree at most N) to a minimizer \mathcal{U}_N^K of (24). Moreover, if (24) has a unique minimizer \mathcal{U}_N^K , then the whole sequence $(\mathcal{V}_{N,d}^K)_{d \geq N}$ converges to \mathcal{U}_N^K , i.e.

$$\mathcal{V}_{N,d}^K \xrightarrow[d \rightarrow \infty]{} \mathcal{U}_N^K. \quad (49)$$

Proof We first prove that the minima of (46) converge monotonically to the minimum of (24), i.e.

$$\begin{aligned} \dots & \leq \|\mathcal{V}_{N,d}^K\|_d \leq \|\mathcal{V}_{N,d+1}^K\|_{d+1} \leq \dots \leq \left\| \frac{\mathcal{U}_N^K}{w} \right\|_{L_1(K)} \quad \text{and} \\ & \lim_{d \rightarrow \infty} \|\mathcal{V}_{N,d}^K\|_d = \left\| \frac{\mathcal{U}_N^K}{w} \right\|_{L_1(K)}. \end{aligned} \quad (50)$$

The argument is quite similar to the proof of (37) in Proposition 4. The chain of inequalities holds because more and more constraints are imposed. Next, considering coefficients $p_0^d, p_1^d, \dots, p_N^d$ and infinite sequences $\mathbf{y}^{1,\pm,d}, \dots, \mathbf{y}^{L,\pm,d}$ satisfying

$$\sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+d} + y_0^{\ell,-d}) = \|\mathcal{V}_{N,d}^K\|_d, \quad (51)$$

as well as $p_N^d = 1/2^{N-1}$ and, for all $\ell = 1 : L$,

$$\mathbf{y}^{\ell,+,\pm,d} - \mathbf{y}^{\ell,-,\pm,d} = \mathbf{J}^\ell \mathbf{W}^\ell \mathbf{p}^d \quad \text{and} \quad \mathbf{Toep}_d(\mathbf{y}^{\ell,\pm,d}) \succeq \mathbf{0}, \quad (52)$$

the semidefiniteness of the Toeplitz matrices, together with (51), still implies that the sequences $(\mathbf{y}^{\ell,+,\pm,d})_{d \geq N}$ admit convergent subsequences in the weak-star topology, so we can write

$$y_k^{\ell,\pm,d_m} \xrightarrow[m \rightarrow \infty]{} y_k^{\ell,\pm} \quad \text{for all } k \geq 0. \quad (53)$$

We note that

$$\begin{aligned} \mathbf{p}^{d_m} &= (\mathbf{J}^{\ell,N} \mathbf{W}^\ell)^{-1} (\mathbf{y}_{\{0,\dots,N\}}^{\ell,+,\pm,d_m} - \mathbf{y}_{\{0,\dots,N\}}^{\ell,-,\pm,d_m}) \\ &\xrightarrow[m \rightarrow \infty]{} (\mathbf{J}^{\ell,N} \mathbf{W}^\ell)^{-1} (\mathbf{y}_{\{0,\dots,N\}}^{\ell,+} - \mathbf{y}_{\{0,\dots,N\}}^{\ell,-}) =: \mathbf{p}. \end{aligned} \quad (54)$$

It is easy to see that the coefficients $p_0, p_1, \dots, p_N \in \mathbb{R}$ thus defined, together with the sequences $\mathbf{y}^{1,\pm}, \dots, \mathbf{y}^{L,\pm} \in \mathbb{R}^N$, are feasible for the problem (48), which implies that

$$\begin{aligned} \left\| \frac{\mathcal{U}_N^K}{w} \right\|_{L_1} &\leq \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+} + y_0^{\ell,-}) = \lim_{m \rightarrow \infty} \sum_{\ell=1}^L \frac{b_\ell - a_\ell}{2} (y_0^{\ell,+,\pm,d_m} + y_0^{\ell,-,\pm,d_m}) \\ &= \lim_{m \rightarrow \infty} \|\mathcal{V}_{N,d_m}^K\|_{d_m} = \lim_{d \rightarrow \infty} \|\mathcal{V}_{N,d}^K\|_d. \end{aligned} \quad (55)$$

This concludes the justification of (50).

Let us now prove that the sequence $(\mathcal{V}_{N,d}^K)_{d \geq N}$ admits a subsequence converging to a minimizer \mathcal{U}_N^K of (24). This sequence is bounded (with respect to any of the equivalent norms, e.g. $\|\cdot\|_N$): indeed, as a consequence of (37) and (50), we have $\|\mathcal{V}_{N,d}^K\|_N \leq \|\mathcal{V}_{N,d}^K\|_d \leq \|\mathcal{U}_N^K/w\|_{L_1(K)}$. Therefore, there is a subsequence $(\mathcal{V}_{N,d_m}^K)_{m \geq 0}$ converging to some monic polynomial \mathcal{V}_N^K . Let us assume that \mathcal{V}_N^K is not one of the minimizers \mathcal{U}_N^K of (24), i.e. that $\|\mathcal{U}_N^K/w\|_{L_1(K)} < \|\mathcal{V}_N^K/w\|_{L_1(K)}$. In view of (37), we can choose d large enough so that

$$\begin{aligned} \left\| \mathcal{V}_N^K/w \right\|_{L_1(K)} &< \|\mathcal{V}_N^K\|_d + \varepsilon, \quad \text{where} \\ \varepsilon &:= \left\| \mathcal{V}_N^K/w \right\|_{L_1(K)} - \left\| \mathcal{U}_N^K/w \right\|_{L_1(K)} > 0. \end{aligned} \quad (56)$$

Let us observe that with d being fixed and by virtue of (37) and (50),

$$\|\mathcal{V}_N^K\|_d = \lim_{m \rightarrow \infty} \|\mathcal{V}_{N,d_m}^K\|_d \leq \lim_{m \rightarrow \infty} \|\mathcal{V}_{N,d_m}^K\|_{d_m} = \left\| \mathcal{U}_N^K/w \right\|_{L_1(K)}. \quad (57)$$

Combining (56) and (57) yields

$$\left\| \mathcal{V}_N^K / w \right\|_{L_1(K)} < \left\| \mathcal{U}_N^K / w \right\|_{L_1(K)} + \varepsilon = \left\| \mathcal{V}_N^K / w \right\|_{L_1(K)}, \quad (58)$$

which is of course a contradiction. This implies that \mathcal{V}_N^K is a minimizer of (24), as expected.

Finally, in case (24) has a unique minimizer \mathcal{U}_N^K , we can establish (49) by contradiction. Namely, if the sequence $(\mathcal{V}_{N,d}^K)_{d \geq N}$ did not converge to \mathcal{U}_N^K , then we could construct a subsequence $(\mathcal{V}_{N,d_m}^K)_{m \geq 0}$ converging to some monic polynomial $\mathcal{V}_N^K \neq \mathcal{U}_N^K$. Repeating the above arguments would imply that \mathcal{V}_N^K is a minimizer of (24), i.e. $\mathcal{V}_N^K = \mathcal{U}_N^K$, providing the required contradiction. \square

Theorem 5 does not indicate how to choose d a priori in order to reach a prescribed accuracy for the distance between $\mathcal{V}_{N,d}^K$ and \mathcal{U}_N^K . However, for a given d , we can assess a posteriori the distance between the ersatz minimum $\|\mathcal{V}_{N,d}^K\|_d$ and the genuine minimum $\|\mathcal{U}_N^K / w\|_{L_1(K)}$. Indeed, on the one hand, the semidefinite program (47) produces $\|\mathcal{V}_{N,d}^K\|_d$ while outputting $\mathcal{V}_{N,d}^K$; on the other hand, the weighted L_1 -norm $\|\mathcal{V}_{N,d}^K / w\|_{L_1(K)}$ can be computed once $\mathcal{V}_{N,d}^K$ has been output. These two facts provide lower and upper bounds for the unknown value $\|\mathcal{U}_N^K / w\|_{L_1(K)}$, as stated by the quantitative result below.

Proposition 6 *For any $d \geq N$, one has*

$$\|\mathcal{V}_{N,d}^K\|_d \leq \left\| \mathcal{U}_N^K / w \right\|_{L_1(K)} \leq \left\| \mathcal{V}_{N,d}^K / w \right\|_{L_1(K)}, \quad (59)$$

hence the weighted L_1 -norm of \mathcal{U}_N^K on K is approximated with a computable relative error of

$$\delta_{N,d}^K := 1 - \frac{\|\mathcal{V}_{N,d}^K\|_d}{\|\mathcal{V}_{N,d}^K / w\|_{L_1(K)}} \geq 0. \quad (60)$$

Proof By the definition (24) of the genuine Chebyshev polynomial of the second kind, we have

$$\|\mathcal{U}_N^K / w\|_{L_1(K)} \leq \|\mathcal{V}_{N,d}^K / w\|_{L_1(K)}, \quad (61)$$

and by the definition (46) of the ersatz Chebyshev polynomial of the second kind, together with (37), we have

$$\|\mathcal{V}_{N,d}^K\|_d \leq \|\mathcal{U}_N^K\|_d \leq \|\mathcal{U}_N^K / w\|_{L_1(K)}. \quad (62)$$

This establishes the bounds announced in (59). We also notice that the relative error satisfies

$$\delta_{N,d}^K = \frac{\|\mathcal{V}_{N,d}^K/w\|_{L_1(K)} - \|\mathcal{V}_{N,d}^K\|_d}{\|\mathcal{V}_{N,d}^K/w\|_{L_1(K)}} \xrightarrow{d \rightarrow \infty} 0, \quad (63)$$

since, according to (49) and (50), both $\|\mathcal{V}_{N,d}^K/w\|_{L_1(K)}$ and $\|\mathcal{V}_{N,d}^K\|_d$ converge to $\|\mathcal{U}_N^K/w\|_{L_1(K)}$ in case of uniqueness of \mathcal{U}_N^K . In case of non-uniqueness, (63) remains true at least for a subsequence. \square

Figure 2 shows ersatz Chebyshev polynomials of the second kind computed on the same examples as in Fig. 1. Notice that no ‘restricted’ ersatz Chebyshev polynomials of the second kind are displayed. This is because our experiments suggested that the polynomials $\mathcal{V}_{N,d}^K$ had simple roots all lying inside K . The corresponding statement for the polynomials \mathcal{U}_N^K , in case of uniqueness, can in fact be justified theoretically by the following observation.

Proposition 7 *Let \mathcal{U}_N^K be a weighted Chebyshev polynomial of the second kind for a finite union of closed intervals $K \subseteq [-1, 1]$. This polynomial is the unique minimizer of (24) if and only if it has N simple roots all lying inside K .*

Proof As a minimizer of (24), a Chebyshev polynomial of the second kind for K is characterized (see, e.g. [4, p. 84, Thm. 10.4]) by the condition

$$\int_K \frac{\operatorname{sgn}(\mathcal{U}_N^K(x)) P(x)}{w(x)} dx = 0$$

for all polynomials P of degree less than N . (64)

This implies that \mathcal{U}_N^K has N roots in $(-1, 1)$, as (64) would not hold for $P(x) = (x - \xi_1) \cdots (x - \xi_n)$ if \mathcal{U}_N^K had $n < N$ roots ξ_1, \dots, ξ_n in $(-1, 1)$. Moreover, if one of the roots was repeated, we would have $\mathcal{U}_N^K(x) = (x - \xi)^2 P(x)$ for some polynomial P of degree $N - 2$, but then (64) would not hold for this P either. Thus, the polynomial \mathcal{U}_N^K can be written, with distinct $\xi_1, \dots, \xi_N \in (-1, 1)$, as

$$\mathcal{U}_N^K(x) = (x - \xi_1) \cdots (x - \xi_i) \cdots (x - \xi_N). \quad (65)$$

Assume that \mathcal{U}_N^K is the unique Chebyshev polynomial of the second kind for K . If one of the ξ_i ’s does not lie inside K , i.e. if ξ_i belongs to one of the gaps $[b_\ell, a_{\ell+1}]$, then we can perturb ξ_i to $\tilde{\xi}_i$ while keeping it in $[b_\ell, a_{\ell+1}]$. Hence, the perturbed monic polynomial

$$\tilde{\mathcal{U}}_N^K(x) = (x - \xi_1) \cdots (x - \tilde{\xi}_i) \cdots (x - \xi_N). \quad (66)$$

still satisfies $\operatorname{sgn}(\tilde{\mathcal{U}}_N^K(x)) = \operatorname{sgn}(\mathcal{U}_N^K(x))$ for all $x \in K$. The condition (64) is then fulfilled by $\tilde{\mathcal{U}}_N^K$, too, so this monic polynomial is another minimizer of (24), which is impossible. We have, therefore, proved that the N simple roots of \mathcal{U}_N^K all lie inside K .




Fig. 2 Ersatz 5th Chebyshev polynomials of the second kind for $K_1 = [-1, -\frac{1}{2}] \cup [-\frac{1}{5}, \frac{1}{5}] \cup [\frac{1}{2}, 1]$ and for $K_2 = [-1, -\frac{1}{2}] \cup [\frac{1}{10}, \frac{1}{5}] \cup [\frac{2}{3}, 1]$: the first column corresponds to the unweighted case, while weighted ersatz Chebyshev polynomials with weight $w(x) = (1+x^2)/(2-x^2)$ are shown in the second column

Conversely, assume that \mathcal{U}_N^K has N simple roots all lying inside K and let us prove that \mathcal{U}_N^K is the unique minimizer of (24). Consider a monic polynomial $\tilde{\mathcal{U}}_N^K$ with $\|\tilde{\mathcal{U}}_N^K/w\|_{L_1(K)} = \|\mathcal{U}_N^K/w\|_{L_1(K)}$. In view of (64), we notice that

$$\int_K \frac{\operatorname{sgn}(\mathcal{U}_N^K(x))(\mathcal{U}_N^K(x) - \tilde{\mathcal{U}}_N^K(x))}{w(x)} dx = 0. \quad (67)$$

From here, it follows that

$$\begin{aligned} \left\| \frac{\mathcal{U}_N^K}{w} \right\|_{L_1(K)} &= \int_K \frac{|\mathcal{U}_N^K(x)|}{w(x)} dx = \int_K \frac{\operatorname{sgn}(\mathcal{U}_N^K(x))\mathcal{U}_N^K(x)}{w(x)} dx \\ &= \int_K \frac{\operatorname{sgn}(\mathcal{U}_N^K(x))\tilde{\mathcal{U}}_N^K(x)}{w(x)} dx \leq \int_K \frac{|\tilde{\mathcal{U}}_N^K(x)|}{w(x)} dx = \left\| \frac{\tilde{\mathcal{U}}_N^K}{w} \right\|_{L_1(K)}. \end{aligned} \quad (68)$$

The first and the last terms being equal, we must have equality all the way through, which means that $\text{sgn}(\tilde{\mathcal{U}}_N^K(x)) = \text{sgn}(\mathcal{U}_N^K(x))$ for all $x \in K$. Given that the polynomial \mathcal{U}_N^K vanishes at distinct points ξ_1, \dots, ξ_N inside K , the polynomial $\tilde{\mathcal{U}}_N^K$ must also vanish at ξ_1, \dots, ξ_N , and since both polynomials are monic, we must have $\tilde{\mathcal{U}}_N^K = \mathcal{U}_N^K$, proving the uniqueness. \square

References

1. Christiansen, J., Simon, B., Zinchenko, M.: Asymptotics of Chebyshev polynomials, I: subsets of \mathbb{R} . *Inventiones Mathematicae* **208**(1), 217–245 (2017)
2. Christiansen, J., Simon, B., Yuditskii, P., Zinchenko, M.: Asymptotics of Chebyshev polynomials, II: DCT subsets of \mathbb{R} . *Duke Mathematical Journal* **168**(2), 325–349 (2019)
3. CVX Research, Inc. CVX: MATLAB software for disciplined convex programming, version 2.1 (2017). <http://cvxr.com/cvx>
4. DeVore, R.A., Lorentz, G.G.: Constructive Approximation, vol. 303. Springer Science & Business Media, (1993)
5. Filip, S.-I.: A robust and scalable implementation of the Parks-McClellan algorithm for designing FIR filters. *ACM Transactions on Mathematical Software (TOMS)* **43**(1), 7 (2016)
6. Foucart, S., Lasserre, J.: Determining projection constants of univariate polynomial spaces. *Journal of Approximation Theory* **235**, 74–91 (2018)
7. Foucart, S., Powers, V.: Basc: constrained approximation by semidefinite programming. *IMA Journal of Numerical Analysis* **37**(2), 1066–1085 (2017)
8. Geronimo, J., Van Assche, W.: Orthogonal polynomials on several intervals via a polynomial mapping. *Transactions of the American Mathematical Society* **308**(2), 559–581 (1988)
9. Peherstorfer, F.: Orthogonal and extremal polynomials on several intervals. *Journal of Computational and Applied Mathematics* **48**(1-2), 187–205 (1993)
10. Ransford, T.: Potential Theory in the Complex Plane, vol. 28. Cambridge University Press, (1995)
11. Ransford, T., Rostand, J.: Computation of capacity. *Mathematics of Computation* **76**(259), 1499–1520 (2007)
12. Saff, E., Totik, V.: Logarithmic Potentials with External Fields, vol. 316. Springer Science & Business Media, (2013)
13. Trefethen, L.N. et al.: Chebfun Version 5, The Chebfun Development Team (2017). <http://www.chebfun.org>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.