



Mechanochemical C–H bond activation: Synthesis of the tuckover hydrides, $(C_5Me_5)_2Ln(\mu-H)(\mu-\eta^1:\eta^5-CH_2C_5Me_4)Ln(C_5Me_5)$ from solvent-free reactions of $(C_5Me_5)_2Ln(\mu-Ph)_2BPh_2$ with KC_5Me_5

David H. Woen, Jessica R.K. White, Joseph W. Ziller, William J. Evans*

Department of Chemistry, University of California, Irvine, CA, 92697-2025, United States

ARTICLE INFO

Article history:

Received 25 June 2019

Received in revised form

27 July 2019

Accepted 30 July 2019

Available online 2 August 2019

Keywords:

Lanthanide

Mechanochemistry

C–H activation

Reduction

ABSTRACT

Attempts to synthesize the sterically crowded complexes $(C_5Me_5)_3Ln$ of the smallest lanthanide metals, $Ln = Tm$ and Lu , from $(C_5Me_5)_2Ln(\mu-Ph)_2BPh_2$ and KC_5Me_5 via mechanochemical methods has revealed C–H bond activation chemistry: the tuckover hydride complexes, $(C_5Me_5)_2Ln(\mu-H)(\mu-\eta^1:\eta^5-CH_2C_5Me_4)Ln(C_5Me_5)$, were observed in low crystalline yields. In the case of $Ln = Yb$, attempts to synthesize the cationic precursor, $(C_5Me_5)_2Yb(\mu-Ph)_2BPh_2$, from the reaction of $(C_5Me_5)_2Yb(\eta^3-C_3H_5)$ and $[HNEt_3][BPh_4]$ led to the reduction of the metal and the formation of the $Yb(II)$ cationic complex, $(C_5Me_5)Yb(\mu-Ph)_2BPh_2$.



© 2019 Elsevier B.V. All rights reserved.

1. Introduction

For several decades after the $(C_5Me_5)^{1-}$ ligand was introduced into organometallic chemistry [1–3], it was assumed that three of these large ligands could not be put around a single metal center since the cone angle of $(C_5Me_5)^{1-}$ was estimated to be 142° [4,5]. However, this assumption was overturned by the discovery of $(C_5Me_5)_3Sm$, **1-Sm**, which formed from the reaction of the $Sm(II)$ complex, $(C_5Me_5)_2Sm$, with C_8H_8 , eq 1 [6]. X-ray crystallographic analysis of **1-Sm** revealed that the cone angle in this complex was 120° . This “reduced” cone angle was achieved because the ligands were positioned further away from the metal than in traditional $(C_5Me_5)^{1-}$ complexes of samarium: the average $Sm-C(\text{ring})$ distance was ca. 0.1 Å longer than those of all other $Sm(III)$ complexes previously characterized by X-ray crystallography [7]. Subsequently, $(C_5Me_5)_3M$ complexes were discovered with $M = La$ [8], Ce [9], Pr [9], Nd [10], Gd [11], Y [11], U [12,13], and Th [14], all of which also had long $M-C(C_5Me_5)$ distances compared to previously reported rare-earth cyclopentadienyl complexes [7].

Increased reactivity of the $(C_5Me_5)^{1-}$ ligand was observed due to the increased distance from the metal center. For example, with the


exception of the thorium complex, $(C_5Me_5)_3M$ complexes were capable of ring opening THF [9,15] to make complexes with $[O(CH_2)_4(C_5Me_5)]^{1-}$ ligands. They also displayed chemistry consistent with “ $(C_5Me_5)_2M(\eta^1-C_5Me_5)$ ” intermediates in ethylene polymerization, as well as one-electron reductive chemistry based on a $(C_5Me_5)^{1-}/(C_5Me_5)$ redox couple in reactions termed sterically-induced reduction [16]. In the case of yttrium, C–H bond activation was observed with aromatic solvents, benzene and toluene, to form phenyl and benzyl products, $(C_5Me_5)_2Y(C_6H_5)$ and $(C_5Me_5)_2Y(CH_2C_6H_5)$, respectively, eq 2 [11].




* Corresponding author.

E-mail address: wevans@uci.edu (W.J. Evans).

Most of the $(C_5Me_5)_3M$ complexes ($M = La, Ce, Pr, Nd, Gd, U, Th$) can be prepared from the reaction of the cationic complexes, $(C_5Me_5)_2M(\mu-Ph)_2BPh_2$, with one equiv of KC_5Me_5 in benzene or toluene. Since $(C_5Me_5)_3Y$, **1-Y**, readily reacts with benzene and toluene, **1-Y** was prepared by reacting the bridged hydride complex, $[(C_5Me_5)_2YH]_2$, with 1,2,3,4-tetramethylfulvene (TMF) in methylcyclohexane (MCH), eq 3 [11].



Recently, the sterically crowded $(C_5Me_5)_3Ln$, **1-Ln**, complexes of the smaller rare-earth metals, $Ln = Tb, Dy, Ho, Er$, and Y were successfully synthesized using mechanochemistry, eq 4 [17].




Specifically, the cationic metal precursors, $(C_5Me_5)_2Ln(\mu-Ph)_2BPh_2$, were reacted in the solid state with KC_5Me_5 by ball-milling with stainless steel spheres of 6 mm diameter in a disperser tube. This synthetic approach was advantageous since it avoided the benzene and toluene solvents previously shown to react with **1-Y**, eq 2.

We report here the use of this solvent-free mechanochemical method with the smallest metals in the lanthanide series, which led to the identification of mechanochemical C–H bond activation pathways. Analogs of the known tuckover hydride complexes $(C_5Me_5)_2Ln(\mu-H)(\mu-\eta^1:\eta^5-CH_2C_5Me_4)Ln(C_5Me_5)$ ($Ln = Y$ [18], La [19], Sm [20], Lu [21]) arising from C–H bond activation of $(C_5Me_5)^{1-}$ ligands are isolated with Tm and Lu . For Yb , we find that attempts to make the $(C_5Me_5)_2Ln(\mu-Ph)_2BPh_2$ precursor for this reaction leads to reduction and formation of the known $Yb(II)$ cationic complex, $(C_5Me_5)Yb(\mu-Ph)_2BPh_2$ [22].

2. Results and discussion

The cationic metal precursors, $(C_5Me_5)_2Ln(\mu-Ph)_2BPh_2$, **2-Ln** ($Ln = Tm, Lu$), were successfully prepared following the established procedure for analogs of the other rare-earth metals, by reaction of $LnCl_3$ with 2 equiv KC_5Me_5 to make $(C_5Me_5)_2Ln(\mu-Cl_2)K(THF)_x$, followed by conversion of the ate salt to $(C_5Me_5)_2Ln(\eta^3-C_3H_5)$, and protonolysis of the subsequent allyl complex with $[HNEt_3][BPh_4]$ [10,11]. Milling powders of pale orange **2-Tm** and colorless **2-Lu** with KC_5Me_5 using 6 mm diameter stainless steel balls in a BMT-20-S tube powered by an Ultra-Turrax Tube Drive gave a dark orange material with each metal. Extraction of the reaction product with pentane ($Ln = Tm$) or hexane ($Ln = Lu$) gave low yields of single crystals identified by X-ray crystallography as the tuck-over bridging hydride species, $(C_5Me_5)_2Ln(\mu-H)(\mu-\eta^1:\eta^5-CH_2C_5Me_4)Ln(C_5Me_5)$ ($Ln = Tm, Lu$), **3-Ln**, eq 5. Although the complexes were not isolated in synthetically useful yields, the result shows that C–H bond activation pathways are viable under mechanochemical conditions.



A small amount of crystals of **3-Tm** were grown from pentane in the triclinic space group, $P\bar{1}$, and the unit cell was found to be isomorphous to those of the previously reported **3-Lu** [21], **3-Sm** [20], and the heterobimetallic **3-La/Lu** analogs [18]. A low quantity of single crystals of the lutetium mechanochemical product were grown from a concentrated methylcyclohexane solution at $-35^\circ C$ and, in this case, **3-Lu** crystallized in the monoclinic space group, $P2_1/n$, Fig. 1. This unit cell is different that that of the previously reported structure [21] due to the presence of methylcyclohexane in the lattice. As shown in Table 1, the difference in unit cells did not significantly affect the structural parameters.

The tuckover hydride complexes, **3-Ln**, had previously been generated from the decomposition of the bridging-hydride complexes, $[(C_5Me_5)_2Ln(\mu-H)]_2$. The tuckover complexes presumably




Fig. 1. Thermal ellipsoid plot of **3-Lu** drawn at 50% probability level. A methylcyclohexane molecule and hydrogen atoms, except H1, are omitted for clarity.

formed by the loss of H₂ in a conventional C–H bond activation reaction involving sigma bond metathesis between the metal hydride and the methyl C–H bond [18–21]. Since no hydrides are present in the cationic precursor, **2-Ln**, a different mode of C–H bond activation may be occurring in this case. It is conceivable that the tuckover hydride complexes **3-Ln** arise from decomposition of the target molecule, (C₅Me₅)₃Ln, **1-Ln**. In the previous studies of mechanochemical synthesis of the structurally characterized complexes, (C₅Me₅)₃Ln (Ln = Ho and Er), **1-Ln**, according to eq 4 [17], **3-Ln** complexes were occasionally obtained as dark red-orange side products. X-ray crystallographic data on **3-Er** obtained from one of those reactions is given in the Supporting Information (Tables S6–10). (C₅Me₅)₃Ln complexes are known to react as if one of the (C₅Me₅)¹⁻ rings was a (η¹-C₅Me₅)¹⁻ ligand, i.e., "(C₅Me₅)₂Ln(η¹-C₅Me₅)" [9]. The (η¹-C₅Me₅)¹⁻ ligand would be capable of metalating a (C₅Me₅)¹⁻ ring to make C₅Me₅H and [C₅Me₄(CH₂)]²⁻, but the origin of the hydride ligand in **3-Ln** is more difficult to explain. Since it is now known that Lu(II) complexes exist [23] and that Ln(II) complexes can effect C–H bond activation to make hydride complexes [24,25], it is possible that the formation of **3-Ln** could proceed through (C₅Me₅)₃Ln-based sterically induced reduction [9,26,27] by a (C₅Me₅)¹⁻ ligand. However, no evidence of (C₅Me₅)₂, the byproduct of such a reduction, was observed. Since the mechanochemical reaction cannot be followed spectroscopically, there are no data with which to evaluate these possibilities.

For ytterbium, attempts to synthesize the cationic precursor, (C₅Me₅)₂Yb(μ-Ph)₂BPh₂, by the same literature procedures [10,11] successful for other rare-earth metals led to reduction and isolation of an Yb(II) complex. Following the previously reported procedures for the synthesis of (C₅Me₅)₂Ln(η³-C₃H₅) complexes [10,11,28], the reaction of (C₅Me₅)₂Yb(μ-Cl₂)K(THF)₂ with allyl magnesium chloride in toluene and subsequent extraction in hexane yielded the dark microcrystalline allyl complex, (C₅Me₅)₂Yb(η³-C₃H₅), **4**. Drying **4** further under reduced pressure at 45 °C over three days and placing a concentrated hexane solution of **4** in a –30 °C freezer gave black single crystals of **4** suitable for X-ray diffraction Fig. 2. The structural data for **4** are summarized in Table 2. These are consistent with other (C₅Me₅)₂Ln(η³-C₃H₅) complexes as shown [28].

Reaction of **4** with [HNET₃][BPh₄] in toluene resulted in the formation of the Yb(II) complex, (C₅Me₅)Yb(μ-Ph)₂BPh₂, **5** [22], eq 6, Fig. 3. Although this route to **5** is not synthetically useful, it shows the care that must be used with Yb(III) syntheses to avoid reduction to Yb(II) products.

The crystals of **5** were obtained from a concentrated toluene solution at –35 °C and formed in the P2₁/n space group with toluene in the lattice. Previously, a structure of **5** synthesized from




Fig. 2. Thermal ellipsoid plot of **4** drawn at the 50% probability level. Hydrogen atoms are omitted for clarity.

(C₅Me₅)₂Yb and [HNET₃][BPh₄] also formed crystals in the P2₁/n space group [22]. In that case, however, the complex was obtained from a concentrated benzene solution and had benzene in the lattice such that the complexes are not isomorphous. Nonetheless, both have two molecules of (C₅Me₅)Yb(μ-Ph)₂BPh₂ in the unit cell. As shown in Table 3, the difference in lattice solvent did not significantly affect the structural parameters.

It is interesting to note that the reduction reaction that formed **5** occurred by the addition of a Brønsted acid, [HNET₃][BPh₄]. However, formation of Yb(II) complexes from Yb(III) complexes is known to be facile due to the –1.15 V (vs NHE) Yb³⁺/Yb²⁺ reduction potential [30]. For example, the Yb(III) complex, [(C₅H₄Me)₂YbMe]₂, forms the Yb(II) complex, (C₅H₄Me)₂Yb(THF), upon exposure to UV light or by heating to 80 °C [31]. Andersen and coworkers have shown that ytterbium bipyridine (bipy) complexes, Cp^x₂Yb(bipy) (Cp^x = C₅Me₅, 1,3-(CMe₃)₂C₃H₃, C₅Me₄H; bipy = 2,2'-bipyridine) exist in temperature dependent equilibria as a mixture of Yb(III) (bipy)¹⁻ and Yb(II) neutral bipy complexes [32]. Since [BPh₄]¹⁻ is known to be a reductant [33–41], it is possible that this effects the reduction in eq 6. The ease in forming Yb(II) complexes suggests that if "(C₅Me₅)₃Yb" were formed by mechanochemistry, it might form the known (C₅Me₅)₂Yb [42].

3. Conclusion

Attempts to synthesize the sterically crowded complexes, (C₅Me₅)₃Ln, **1-Ln**, with the smaller lanthanide metals, Tm, Yb, and Lu, by mechanochemical reaction of (C₅Me₅)₂Ln(μ-Ph)₂BPh₂, **2-Ln**, and KC₅Me₅ have not been successful. The ball-milling reaction produces the tuckover hydride complexes, (C₅Me₅)₂Ln(μ-H)(μ-η¹:η⁵-CH₂C₅Me₄)Ln(C₅Me₅) (Ln = Tm, Lu), **3-Ln**, in low yields, but demonstrates that C–H bond activation reactivity can occur under these mechanochemical conditions. For ytterbium, efforts to prepare the (C₅Me₅)₂Ln(μ-Ph)₂BPh₂ precursor by reaction of (C₅Me₅)₂Yb(η³-C₃H₅) with [HNET₃][BPh₄] formed (C₅Me₅)Yb(μ-Ph)₂BPh₂, **5**, and revealed another facile reduction route for the

Table 1

Bond lengths [Å] and angles [°] of **3-Lu**, crystallized in P2₁/n with methylcyclohexane in the lattice and solvent free in P₁ [21].

	3-Lu (MCH) P2 ₁ /n	3-Lu P ₁
Lu(1)–Cnt1 ^a	2.321	2.333
Lu(1)–Cnt2	2.286	2.296
Lu(2)–Cnt3 ^b	2.320	2.334
Lu(2)–Cnt4	2.347	2.344
Lu(1)–H(1)	2.06(4)	2.01(5)
Lu(2)–H(1)	2.13(4)	2.09(5)
Lu(2)–C(6)	2.548(3)	2.556(5)
Lu(1)–C(26)	2.817(4)	2.815(5)
Cnt1–Lu(1)–C(26)	106.2	105.7
Cnt2–Lu(1)–C(26)	109.5	108.8
Cnt1–Lu(1)–Cnt2	136.1	136.7
Cnt3–Lu(2)–C(6)	107.1	110.7
Cnt4–Lu(2)–C(6)	109.6	110.7
Cnt3–Lu(2)–Cnt4	133.9	134.0

^a Cnt1 = C(1)–C(5).

^b Cnt3 = C(21)–C(25).

Table 2

Comparison of selected bond lengths [Å] and angles [°] of $\text{Cp}^*_2\text{Ln}(\eta^3\text{-C}_3\text{H}_5)$, **4**, with other $\text{Cp}^*_2\text{Ln}(\eta^3\text{-C}_3\text{H}_5)$ complexes ($\text{Cp}^* = \text{C}_5\text{Me}_5$). The eight coordinate ionic radii of trivalent Y, Yb and Lu are 1.019, 0.985, and 0.977 Å respectively [29].

	$\text{Cp}^*_2\text{Y}(\text{C}_3\text{H}_5)$ [28]	$\text{Cp}^*_2\text{Yb}(\text{C}_3\text{H}_5)$, 4	$\text{Cp}^*_2\text{Lu}(\text{C}_3\text{H}_5)$ [28]
Ln-Cnt1 ^a	2.381	2.328	2.318
Ln-Cnt2 ^b	2.362	2.313	2.301
Ln-C(allyl)	2.582(2)	2.555(4)	2.549(5)
	2.582(2)	2.559(4)	2.548(4)
	2.601(2)	2.562(4)	2.545(5)
C(allyl)-C(allyl)	1.392(3)	1.381(7)	1.382(8)
	1.391(3)	1.392(7)	1.383(9)
Cnt1-Ln-Cnt2	138.8	138.1	138.5
C(allyl)-C(allyl)-C(allyl)	125.9(2)	127.2(5)	127.7(5)

^a Cnt1 = C(1)–C(5).

^b Cnt2 = C(11)–C(15).




Fig. 3. Thermal ellipsoid plot of **5**. Hydrogen atoms, second $(\text{C}_5\text{Me}_5)\text{Yb}(\mu\text{-Ph})_2\text{BPh}_2$ unit, and two toluene molecules are omitted for clarity.

Table 3

Bond lengths [Å] and angles [°] comparison for $(\text{C}_5\text{Me}_5)\text{Yb}(\mu\text{-Ph})_2\text{BPh}_2$, **5**, crystallized in $\text{P2}_1/\text{n}$ with toluene (this study) or benzene [22] in the lattice.

	5(toluene)	5(benzene) [22]
Yb(1)-Cnt1 ^a	2.385	2.381 Å
Yb(1)-Cnt2 ^b	2.605	2.605 Å
Yb(1)-Cnt3 ^c	2.627	2.639 Å
Yb(2)-Cnt4 ^d	2.395	2.382 Å
Yb(2)-Cnt5 ^e	2.616	2.581 Å
Yb(2)-Cnt6 ^f	2.593	2.636 Å
Cnt1-Yb(1)-Cnt2	124.1	124.7°
Cnt1-Yb(1)-Cnt3	127.0	125.8°
Cnt2-Yb(1)-Cnt3	108.1	109.5°
Cnt4-Yb(2)-Cnt5	125.6	124.2°
Cnt4-Yb(2)-Cnt6	125.6	126.0°
Cnt5-Yb(2)-Cnt6	108.1	109.7°

^a Cnt1 = C(1)–C(5).

^b Cnt2 = C(11)–C(16).

^c Cnt3 = C(17)–C(22).

^d Cnt4 = C(35)–C(42) (See SI).

^e Cnt5 = C(45)–C(50) (See SI).

^f Cnt6 = C(51)–C(56) (See SI).

formation of Yb(II) complexes.

4. Experimental

All manipulations and syntheses described below were

conducted with the rigorous exclusion of air and water using standard Schlenk line and glovebox techniques under an argon or dinitrogen atmosphere. Solvents were sparged with UHP argon and dried by passage through columns containing Q-5 and molecular sieves prior to use. Deuterated NMR solvents were dried over NaK alloy, degassed by three freeze-pump-thaw cycles, and vacuum transferred before use. Allylmagnesium chloride and dioxane were purchased from Aldrich and used as obtained. $[\text{HNEt}_3][\text{BPh}_4]$ [43], $\text{K}(\text{C}_5\text{Me}_5)$ [44], and $(\text{C}_5\text{Me}_5)_2\text{Ln}(\mu\text{-Ph})_2\text{BPh}_2$ ($\text{Ln} = \text{Tm}$ [10], Lu [45]) were prepared according to literature procedures. $(\text{C}_5\text{Me}_5)_2\text{Er}(\mu\text{-H})(\mu\text{-}\eta^1\text{-}\eta^5\text{-CH}_2\text{C}_5\text{Me}_4)\text{Er}(\text{C}_5\text{Me}_5)$, **5-Er**, was prepared serendipitously using the literature procedure for $(\text{C}_5\text{Me}_5)_3\text{Er}$ [17]. ^1H and ^{13}C NMR spectra were obtained on a Bruker DRX500 MHz spectrometer with a TCI probe at 25 °C or a Bruker AVANCE600 MHz spectrometer with a BBO probe at 25 °C. Elemental analyses were conducted on a PerkinElmer 2400 Series II CHNS elemental analyzer. IR spectra were taken as compressed solids on a Thermo Scientific Nicolet iS5 spectrometer with an iD5 ATR attachment or with an Agilent Cary 630 FTIR spectrometer. Disperser milling was performed with an Ultra-Turrax Tube Drive and BMT-20-S tubes, both purchased from IKA.

$(\text{C}_5\text{Me}_5)_2\text{Yb}(\mu\text{-Cl}_2)\text{K}(\text{THF})_2$. In a nitrogen-filled glovebox, KC_5Me_5 (4.00 g, 22.9 mmol) was slowly added to a stirred colorless slurry of YbCl_3 (2.17 g, 7.78 mmol) in THF (100 mL). The reaction mixture initially turned deep blue then slowly changed color to dark maroon. After 2 d, the solution mixture was concentrated to

40 mL and centrifuged to remove insoluble material. The dark supernatant was filtered and dried under reduced pressure to give a dark maroon solid. This solid was washed using toluene (3x) and hexane (3x) over a glass filter frit, then collected and dried under reduced pressure to give the ate-salt complex, $(C_5Me_5)_2Yb(\mu-Cl)_2K(THF)_2$, (3.56 g, 5.10 mmol, 66%). 1H NMR (500 MHz, THF- d_8): δ 3.61 (t, 6H, THF), 3.27 (s, 30H, C_5Me_5), 1.77 (m, 6H, THF). ^{13}C NMR (600 MHz, THF- d_8): δ 129.47 (s, C_5Me_5), 128.71 (s, C_5Me_5), 67.71 (s, THF), 25.59 (s, THF), 11.34 (s, C_5Me_5). IR (cm^{-1}): 2903s, 2856s, 2726w, 1540 m, 1437 m, 1379 m, 1246w, 1023 m, 774s. Anal. Calcd for $C_{28}H_{46}Cl_2KO_2Yb$: C, 48.20; H, 6.65; Found: C, 44.75; H, 6.17. The found C/H ratio, $C_{28}H_{46}$, is consistent with the formula and suggests incomplete combustion [17,46–50].

$(C_5Me_5)_2Lu(\mu-H)(\mu-\eta^1-\eta^5-CH_2C_5Me_4)Lu(C_5Me_5)$, 3-Lu. In an argon-filled glovebox free of coordinating and aromatic solvents, a BMT-20-S tube was charged with $(C_5Me_5)_2Lu(\mu-Ph)_2BPh_2$ (340 mg, 0.46 mmol), KC_5Me_5 (135 mg, 0.766 mmol) and 40 stainless-steel balls (6 mm). The colorless mixture was milled for 40 min using an Ultra-Turrax Tube Drive (max speed setting), after which a pale orange mixture was obtained. Hexane (10 mL) was added to the mixture, followed by centrifugation to produce a dark red-orange solution which was collected in a vial. The solvent was removed under reduced pressure to give a low yield of dark orange oil. The oil was dissolved in minimal amount of methylcyclohexane and left at $-35^\circ C$ for over 7 days. A small quantity of orange single crystals of $(C_5Me_5)_2Lu(\mu-H)(\mu-\eta^1-\eta^5-CH_2C_5Me_4)Lu(C_5Me_5)$, **3-Lu** [21], was obtained after several days as confirmed by X-ray diffraction.

$(C_5Me_5)_2Tm(\mu-H)(\mu-\eta^1-\eta^5-CH_2C_5Me_4)Tm(C_5Me_5)$, 3-Tm. Similar to the procedure for **3-Lu**, in an argon-filled glovebox free of coordinating and aromatic solvents, $(C_5Me_5)_2Tm(\mu-Ph)_2BPh_2$ (200 mg, 0.26 mmol) and KC_5Me_5 (46 mg, 0.26 mmol) were milled, extracted, and dried to give a low yield of orange oil. Pentane (2 mL) was added to the oil, and the resulting orange solution was filtered and stored at $-35^\circ C$. A small amount of dark orange single crystals of $(C_5Me_5)_2Tm(\mu-H)(\mu-\eta^1-\eta^5-CH_2C_5Me_4)Tm(C_5Me_5)$, **3-Tm**, were obtained after over 7 days, as confirmed by matching unit cell using X-ray diffraction.

$(C_5Me_5)_2Yb(\eta^3-C_3H_5)$, 4. In a nitrogen-filled glovebox, allyl magnesium chloride (3.0 mL of a 2.0 M THF solution) was added to a stirred maroon slurry of $(C_5Me_5)_2Yb(\mu-Cl)_2K(THF)_2$ (3.56 g, 5.10 mmol) in toluene (50 mL). No obvious color change was observed. The reaction was allowed to stir for 4 h and then volatiles were removed under reduced pressure to yield a mixture of brown and white solids. Hexane (40 mL) and dioxane (1 mL) were added and the mixture was allowed to stir. After 4 h, the mixture was centrifuged, and the solvent was removed under reduced pressure to yield $(C_5Me_5)_2Yb(\eta^3-C_3H_5)$, **4**, as black crystalline solids. The solids were dried further under reduced pressure at $45^\circ C$ over three days to give black crystalline solids of **4** (1.20 g, 2.48 mmol, 49%). Black single crystals of **4** suitable for X-ray diffraction were grown from a concentrated hexane solution at $-30^\circ C$. 1H NMR (500 MHz, C_6D_6): δ 1.94 (s, 30H, C_5Me_5). The C_3H_5 resonances appear as broad signals at 6.16 ppm and 7.69 ppm. ^{13}C NMR (600 MHz, C_6D_6): δ 110.32 (s, C_5Me_5), 109.19 (s, C_5Me_5), 69.34 ($(CH_2)_2CH$), 66.31 ($(CH_2)_2CH$), 11.44 (s, C_5Me_5), 9.97 (s, C_5Me_5). No $(CH_2)_2CH$ resonances were observable. IR (cm^{-1}): 2903s, 2854s, 2718w, 1435 m, 1375 m, 1051s, 1020 m, 896 m. Anal. Calcd for $C_{21}H_{31}Yb$: C, 55.25; H, 6.84. The found values (54.26, 7.53), (54.21, 7.47), (55.88, 7.64), (55.92, 7.67), and (56.25, 7.62) from five different samples all have CH ratios of $C_{31}H_{34}$ (4 values) or $C_{31}H_{35}$ (1 value) which is close to the theoretical $C_{21}H_{31}$, but an exact match with the calculated values has not been obtainable [51].

$(C_5Me_5)_2Yb(\mu-Ph)_2BPh_2$, 5. In an argon-filled glovebox free of coordinating solvents, $[HNEt_3][BPh_4]$ (218 mg, 0.516 mmol) was slowly added to a stirred solution of $(C_5Me_5)_2Yb(\eta^3-C_3H_5)$ (250 mg,

0.516 mmol) in toluene (15 mL). The solution mixture changed from maroon to dark brown-green. The mixture was centrifuged to remove insoluble material and volatiles were removed from the solution under vacuum to yield crystalline solids of $(C_5Me_5)_2Yb(\mu-Ph)_2BPh_2$, **5** (280 mg, 0.447 mmol, 87%). Single crystals of **5** suitable for X-ray diffraction were grown from the toluene solution by pentane layer diffusion at $-30^\circ C$. 1H NMR (C_6D_6): δ 7.65 (s, BPh_4), 7.13 (m, BPh_4), 7.05 (m, BPh_4), 1.67 (s, C_5Me_5) ppm. This NMR matches with that of the previously reported complex [22].

Acknowledgments

This research was supported by the National Science Foundation under CHE-1855328. We also thank Mikey K. Wojnar and Austin J. Ryan for help with X-ray crystallography.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jorgchem.2019.120885>.

References

- [1] R.B. King, A. Fronzaglia, Organometallic chemistry of the transition metals. (XV). New olefinic and acetylenic derivatives of tungsten, *Inorg. Chem.* 5 (1966) 1837–1846, <https://doi.org/10.1021/ic50045a001>.
- [2] R.B. King, M.B. Bisnette, Organometallic chemistry of the transition metals (XXI). Some π -pentamethylcyclopentadienyl derivatives of various transition metals, *J. Organomet. Chem.* 8 (1967) 287–297, [https://doi.org/10.1016/s0022-328x\(00\)91042-8](https://doi.org/10.1016/s0022-328x(00)91042-8).
- [3] H. Brintzinger, J.E. Bercaw, Bis(pentamethylcyclopentadienyl)titanium(II). Isolation and reactions with hydrogen, nitrogen, and carbon monoxide, *J. Am. Chem. Soc.* 93 (1971) 2045–2046, <https://doi.org/10.1021/ja00737a033>.
- [4] C.E. Davies, I.M. Gardiner, J.C. Green, M.L.H. Green, N.J. Hazel, P.D. Grebenik, V.S.B. Mtetwa, K. Prout, Mono- η -cycloheptatrienyltitanium chemistry: synthesis, molecular and electronic structures, and reactivity of the complexes $[Ti(\eta-C_7H_7)L_2X]$ (L = tertiary phosphine, O- or N-donor ligand; X = Cl or alkyl), *J. Chem. Soc., Dalton Trans.* (1985) 669–683, <https://doi.org/10.1039/dt9850000669>.
- [5] T.D. Tilley, R.A. Andersen, Pentamethylcyclopentadienyl derivatives of the trivalent lanthanide elements neodymium, samarium, and ytterbium, *Inorg. Chem.* 20 (1981) 3267–3270, <https://doi.org/10.1021/ic50224a025>.
- [6] W.J. Evans, S.L. Gonzales, J.W. Ziller, Synthesis and X-ray crystal structure of the first tris(pentamethylcyclopentadienyl)metal complex: $(\eta^5-C_5Me_5)_3Sm$, *J. Am. Chem. Soc.* 113 (1991) 7423–7424, <https://doi.org/10.1021/ja00019a050>.
- [7] W.J. Evans, S.E. Foster, Structural trends in bis(pentamethylcyclopentadienyl) lanthanide and yttrium complexes, *J. Organomet. Chem.* (1992), [https://doi.org/10.1016/0022-328x\(92\)80131-G](https://doi.org/10.1016/0022-328x(92)80131-G).
- [8] W.J. Evans, B.L. Davis, J.W. Ziller, Synthesis and structure of tris(alkyl- and silyl-tetramethylcyclopentadienyl) complexes of lanthanum, *Inorg. Chem.* 40 (2001) 6341–6348, <https://doi.org/10.1021/ic010554i>.
- [9] W.J. Evans, J.M. Perotti, S.A. Kozimor, T.M. Champagne, B.L. Davis, G.W. Nyce, C.H. Fujimoto, R.D. Clark, M.A. Johnston, J.W. Ziller, Synthesis and comparative η^1 -alkyl and sterically induced reduction reactivity of $(C_5Me_5)_3Ln$ complexes of La, Ce, Pr, Nd, and Sm, *Organometallics* 24 (2005) 3916–3931, <https://doi.org/10.1021/om050402i>.
- [10] W.J. Evans, C.A. Seibel, J.W. Ziller, Unsolvated lanthanide metallocene cations $[(C_5Me_5)_2Ln][BPh_4]$: multiple syntheses, structural characterization, and reactivity including the formation of $(C_5Me_5)_3Nd$, *J. Am. Chem. Soc.* 120 (1998) 6745–6752, <https://doi.org/10.1021/ja980534o>.
- [11] W.J. Evans, B.L. Davis, T.M. Champagne, J.W. Ziller, C-H bond activation through steric crowding of normally inert ligands in the sterically crowded gadolinium and yttrium $(C_5Me_5)_3M$ complexes, *Proc. Natl. Acad. Sci.* 103 (2006) 12678–12683, <https://doi.org/10.1073/pnas.0602671103>.
- [12] W.J. Evans, K.J. Forrestal, J.W. Ziller, Activity of $[Sm(C_5Me_5)_3]$ in ethylene polymerization and synthesis of $[U(C_5Me_5)_3]$, the first tris(pentamethylcyclopentadienyl) 5f-element complex, *Angew. Chem. Int. Ed. Engl.* 36 (1997) 774–776, <https://doi.org/10.1002/anie.199707741>.
- [13] W.J. Evans, G.W. Nyce, K.J. Forrestal, J.W. Ziller, Multiple syntheses of $(C_5Me_5)_3U$, *Organometallics* 21 (2002) 1050–1055, <https://doi.org/10.1021/om010831t>.
- [14] R.R. Langeslay, G.P. Chen, C.J. Windorff, A.K. Chan, J.W. Ziller, F. Furche, W.J. Evans, Synthesis, structure, and reactivity of the sterically crowded Th^{3+} complex $(C_5Me_5)_3Th$ including formation of the thorium carbonyl, $[(C_5Me_5)_3Th(CO)][BPh_4]$, *J. Am. Chem. Soc.* 139 (2017) 3387–3398, <https://doi.org/10.1021/jacs.6b10826>.
- [15] T.M. Champagne, Ph.D. Dissertation, University of California, Irvine, 2006.

- [16] W.J. Evans, Perspectives in reductive lanthanide chemistry, *Coord. Chem. Rev.* (2000), [https://doi.org/10.1016/S0010-8545\(00\)00267-8](https://doi.org/10.1016/S0010-8545(00)00267-8).
- [17] D.H. Woen, C.M. Kotyk, T.J. Mueller, J.W. Ziller, W.J. Evans, Tris(pentamethylcyclopentadienyl) complexes of late lanthanides Tb, Dy, Ho, and Er: solution and mechanochemical syntheses and structural comparisons, *Organometallics* 36 (2017) 4558–4563, <https://doi.org/10.1021/acs.organomet.7b00385>.
- [18] M.E. Fieser, T.J. Mueller, J.E. Bates, J.W. Ziller, F. Furche, W.J. Evans, Differentiating chemically similar Lewis acid sites in heterobimetallic complexes: the rare-earth bridged hydride $(C_5Me_5)_2Ln(\mu-H)_2Ln'(C_5Me_5)_2$ and tuckover hydride $(C_5Me_5)_2Ln(\mu-H)(\mu-\eta^1-\eta^2-(CH)_2C_5Me_4)Ln'(C_5Me_5)_2$, *Organometallics* 33 (2014) 3882–3890, <https://doi.org/10.1021/om500624x>.
- [19] W.J. Evans, J.M. Perotti, J.W. Ziller, Trialkylboron/lanthanide metallocene hydride chemistry: polydentate bridging of $(HBEt_3)^{1-}$ to lanthanum, *Inorg. Chem.* 44 (2005) 5820–5825, <https://doi.org/10.1021/ic0501061>.
- [20] W.J. Evans, T.A. Ulibarri, J.W. Ziller, Reactivity of samarium complex $[(C_5Me_5)_2Sm(\mu-H)]_2$ in ether and arene solvents. X-ray crystal structures of the internally metalated complex $(C_5Me_5)_2Sm(\mu-H)(\mu-CH_2C_5Me_4)Sm(C_5Me_5)_2$, the benzyl complex $(C_5Me_5)_2Sm(CH_2C_6H_5)(THF)$, and the siloxide, *Organometallics* 10 (1991) 134–142, <https://doi.org/10.1021/om00047a040>.
- [21] W.J. Evans, T.M. Champagne, J.W. Ziller, Organolutetium vinyl and tuck-over complexes via C–H bond activation, *J. Am. Chem. Soc.* 128 (2006) 14270–14271, <https://doi.org/10.1021/ja0657881>.
- [22] W.J. Evans, T.M. Champagne, J.W. Ziller, Synthesis and reactivity of mono(pentamethylcyclopentadienyl) tetraphenylborate lanthanide complexes of ytterbium and samarium: tris(ring) precursors to $(C_5Me_5)_3Ln$ moieties, *Organometallics* 26 (2007) 1204–1211, <https://doi.org/10.1021/om060963s>.
- [23] M.R. MacDonald, J.E. Bates, J.W. Ziller, F. Furche, W.J. Evans, Completing the series of +2 ions for the lanthanide elements: synthesis of molecular complexes of Pr^{2+} , Gd^{2+} , Tb^{2+} , and Lu^{2+} , *J. Am. Chem. Soc.* 135 (2013) 9857–9868, <https://doi.org/10.1021/ja403753j>.
- [24] J.F. Corbey, D.H. Woen, C.T. Palumbo, M.E. Fieser, J.W. Ziller, F. Furche, W.J. Evans, Ligand effects in the synthesis of Ln^{2+} complexes by reduction of tris(cyclopentadienyl) precursors including C–H bond activation of an indenyl anion, *Organometallics* 34 (2015) 3909–3921, <https://doi.org/10.1021/acs.organomet.5b00500>.
- [25] F. Jaroschik, A. Momin, F. Nief, X.-F. Le Goff, G.B. Deacon, P.C. Junk, Dinitrogen reduction and C–H activation by the divalent organoneodymium complex $[(C_5H_2Bu_3)_2Nd(\mu-I)K([18]crown-6)]$, *Angew. Chem. Int. Ed.* 48 (2009) 1117–1121, <https://doi.org/10.1002/anie.200804934>.
- [26] W.J. Evans, K.J. Forrestal, J.W. Ziller, Reaction chemistry of sterically crowded tris(pentamethylcyclopentadienyl)samarium I, *J. Am. Chem. Soc.* 120 (1998) 9273–9282, <https://doi.org/10.1021/ja9809859>.
- [27] W.J. Evans, B.L. Davis, Chemistry of tris(pentamethylcyclopentadienyl) f-element complexes, $(C_5Me_5)_3M$, *Chem. Rev.* 102 (2002) 2119–2136, <https://doi.org/10.1021/cr010298r>.
- [28] W.J. Evans, S.A. Kozimor, J.C. Brady, B.L. Davis, G.W. Nyce, C.A. Seibel, J.W. Ziller, R.J. Doedens, Metalocene allyl reactivity in the presence of alkenes tethered to cyclopentadienyl ligands, *Organometallics* (2005), <https://doi.org/10.1021/om049286x>.
- [29] R.D. Shannon, Revised effective ionic radii and systematic studies of interatomic distances in halides and chalcogenides, *Acta Crystallogr. A* A32 (1976) 751–767, <https://doi.org/10.1107/S0567739476001551>.
- [30] L.J. Nugent, R.D. Baybarz, J.L. Burnett, Electron-transfer and f-d absorption bands of some lanthanide and actinide complexes and the standard (II–III) oxidation potential for each member of the lanthanide and actinide series, *J. Phys. Chem.* 77 (1973) 1528–1539.
- [31] H.A. Zinnen, J.J. Pluth, W.J. Evans, X-Ray crystallographic determination of the structure of bis(methyl-cyclopentadienyl)ytterbium tetrahydrofuranate and its ready formation by four new routes, *J. Chem. Soc., Chem. Commun.* (1980) 810, <https://doi.org/10.1039/c39800000810>.
- [32] M. Schultz, J.M. Boncella, D.J. Berg, T.D. Tilley, R.A. Andersen, Coordination of 2,2'-bipyridyl and 1,10-phenanthroline to substituted ytterbocenes: an experimental investigation of spin coupling in lanthanide complexes, *Organometallics* 21 (2002) 460–472, <https://doi.org/10.1021/om010661k>.
- [33] W.J. Evans, S.A. Kozimor, J.W. Ziller, $[(C_5Me_5)_2U][(\mu-Ph)_2BPh_2]$ as a four electron reductant, *Chem. Commun.* (2005) 4681, <https://doi.org/10.1039/b508612d>.
- [34] J.L.R. Williams, J.C. Doty, P.J. Grisdale, R. Searle, T.H. Regan, G.P. Happ, D.P. Maier, Boron photochemistry. I. Irradiation of sodium tetraarylborates in aqueous solution, *J. Am. Chem. Soc.* 89 (1967) 5153–5157, <https://doi.org/10.1021/ja00996a013>.
- [35] P.J. Grisdale, J.L.R. Williams, M.E. Glogowski, B.E. Babb, Boron photochemistry. Possible role of bridged intermediates in the photolysis of borate complexes, *J. Org. Chem.* 36 (1971) 544–549, <https://doi.org/10.1021/jo00803a012>.
- [36] J.J. Eisch, M.P. Boleslawski, K. Tamao, Bora-aromatic systems. 10. Photochemical generation of the diphenylborate(I) anion from metal tetraphenylborates(III) in aprotic media: repudiation of a contravening claim, *J. Org. Chem.* 54 (1989) 1627–1634, <https://doi.org/10.1021/jo00268a025>.
- [37] J.D. Wilkey, G.B. Schuster, Photochemistry of tetraarylborate salts (Ar_4B^-): formation of 2,5,7,7-tetraphenyl-7-boratabicyclo[4.1.0]hepta-2,4-diene (a boratanorcaradiene) by irradiation of (p-biphenyl)triphenyl borate, *J. Am. Chem. Soc.* 113 (1991) 2149–2155, <https://doi.org/10.1021/ja00006a037>.
- [38] S.H. Strauss, The search for larger and more weakly coordinating anions, *Chem. Rev.* 93 (1993) 927–942, <https://doi.org/10.1021/cr00019a005>.
- [39] J.J. Eisch, J.H. Shah, M.P. Boleslawski, Skeletal rearrangements of arylborane complexes mediated by redox reactions: thermal and photochemical oxidation by metal ions, *J. Organomet. Chem.* 464 (1994) 11–21, [https://doi.org/10.1016/0022-328x\(94\)87003-9](https://doi.org/10.1016/0022-328x(94)87003-9).
- [40] C.L. Crawford, M.J. Barnes, R.A. Peterson, W.R. Wilmarth, M.L. Hyder, Copper-catalyzed sodium tetraphenylborate, triphenylborane, diphenylborinic acid and phenylboronic acid decomposition kinetic studies in aqueous alkaline solutions, *J. Organomet. Chem.* 581 (1999) 194–206, [https://doi.org/10.1016/S0022-328x\(99\)00065-0](https://doi.org/10.1016/S0022-328x(99)00065-0).
- [41] P.K. Pal, S. Chowdhury, M.G.B. Drew, D. Datta, The electrooxidation of the tetraphenylborate ion revisited, *New J. Chem.* 26 (2002) 367–371, <https://doi.org/10.1039/b106356c>.
- [42] M. Schultz, C.J. Burns, D.J. Schwartz, R.A. Andersen, Solid-state structures of base-free ytterbocenes and inclusion compounds of bis(pentamethylcyclopentadienyl)ytterbium with neutral carboranes and toluene: the role of intermolecular contacts, *Organometallics* 19 (2000) 781–789, <https://doi.org/10.1021/om990821g>.
- [43] J.-C. Berthet, C. Villiers, J.-F. Le Maréchal, B. Delavaux-Nicot, M. Lance, M. Nierlich, J. Vigner, M. Ephritikhine, Anionic triscyclopentadienyluranium(III) hydrides, *J. Organomet. Chem.* 440 (1992) 53–65, [https://doi.org/10.1016/0022-328x\(92\)83484-y](https://doi.org/10.1016/0022-328x(92)83484-y).
- [44] W.J. Evans, S.A. Kozimor, J.W. Ziller, N. Kaltsoyannis, Structure, reactivity, and density functional theory analysis of the six-electron reductant, $[(C_5Me_5)_2U]_2(\mu-\eta^6-\eta^6-C_6H_6)$, synthesized via a new mode of $(C_5Me_5)_3M$ reactivity, *J. Am. Chem. Soc.* 126 (2004) 14533–14547, <https://doi.org/10.1021/ja0463886>.
- [45] B.M. Schmiede, J.W. Ziller, W.J. Evans, Reduction of dinitrogen with an yttrium metallocene hydride precursor, $[(C_5Me_5)_2YH]_2$, *Inorg. Chem.* 49 (2010) 10506–10511, <https://doi.org/10.1021/ic101558e>.
- [46] P.B. Hitchcock, M.F. Lappert, L. Maron, A.V. Protchenko, Lanthanum does form stable molecular compounds in the +2 oxidation state, *Angew. Chem. Int. Ed.* 47 (2008) 1488–1491, <https://doi.org/10.1002/anie.200704887>.
- [47] N.F. Chilton, C.A.P. Goodwin, D.P. Mills, R.E.P. Winpenny, The first near-linear bis(amide) f-block complex: a blueprint for a high temperature single molecule magnet, *Chem. Commun.* 51 (2015) 101–103, <https://doi.org/10.1039/c4cc08312a>.
- [48] C.A.P. Goodwin, K.C. Joslin, S.J. Lockyer, A. Formanuk, G.A. Morris, F. Ortu, I.J. Vitorica-Yrezabal, D.P. Mills, Homoleptic trigonal planar lanthanide complexes stabilized by superbulky silylamide ligands, *Organometallics* 34 (2015) 2314–2325, <https://doi.org/10.1021/om501123e>.
- [49] C.A.P. Goodwin, N.F. Chilton, G.F. Vettesse, E.M. Pineda, I.F. Crowe, J.W. Ziller, R.E.P. Winpenny, W.J. Evans, D.P. Mills, Physicochemical properties of near-linear lanthanide(II) bis(silylamide) complexes ($Ln = Sm, Eu, Tm, Yb$), *Inorg. Chem.* 55 (2016) 10057–10067, <https://doi.org/10.1021/acs.inorgchem.6b00808>.
- [50] C.T. Palumbo, D.P. Halter, V.K. Voora, G.P. Chen, A.K. Chan, M.E. Fieser, J.W. Ziller, W. Hieringer, F. Furche, K. Meyer, W.J. Evans, Metal versus ligand reduction in Ln^{3+} complexes of a mesitylene-anchored tris(aryloxide) ligand, *Inorg. Chem.* 57 (2018) 2823–2833, <https://doi.org/10.1021/acs.inorgchem.7b03236>.
- [51] F.P. Gabbaï, P.J. Chirik, D.E. Fogg, K. Meyer, D.J. Mindiola, L.L. Schafer, S.L. You, An editorial about elemental analysis, *Organometallics* (2016) 3255–3256, <https://doi.org/10.1021/acs.organomet.6b00720>.