1	Salinity stress increases the severity of ranavirus epidemics in amphibian populations
2	
3	Emily M. Hall ^{1*} , Jesse L. Brunner ¹ , Brandon Hutzenbiler ¹ , Erica J. Crespi ¹
4	¹ Washington State University, School of Biological Sciences, P.O. Box 644236, Pullman, WA
5	99164-4236
6	
7	
8	*Corresponding Author:
9	Emily M. Hall, Ph.D.
10	Present address: Department of Pathology, Microbiology, and Immunology, Vanderbilt
11	University School of Medicine, Nashville, Tennessee, USA
12	email: emily.m.hall@vanderbilt.edu
13	
14	
15	

Abstract

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

The stress-induced susceptibility hypothesis, which predicts chronic stress weakens immune defenses, was proposed to explain increasing infectious disease-related mass mortality and population declines. Previous work characterized wetland salinization as a chronic stressor to larval amphibian populations. Thus, we combined field observations with experimental exposures quantifying epidemiological parameters to test the role of salinity stress in the occurrence of ranavirus-associated mass mortality events. Despite ubiquitous pathogen presence (94%), populations exposed to salt runoff had slightly more frequent ranavirus related mass mortality events, more lethal infections, and 117-times greater pathogen environmental DNA. Experimental exposure to chronic elevated salinity (0.8-1.6 g/L Cl⁻) reduced tolerance to infection, causing greater mortality at lower doses. We found a strong negative relationship between splenocyte proliferation and corticosterone in ranavirus-infected larvae at a moderate elevation of salinity, supporting glucocorticoid-medicated immunosuppression, but not at high salinity. Salinity alone reduced proliferation further at similar corticosterone levels and infection intensities. Finally, larvae raised in elevated salinity had 10-times more intense infections and shed 5-times as much virus with similar viral decay rates, suggesting increased transmission. Our findings illustrate how a small change in habitat quality leads to more lethal infections and potentially greater transmission efficiency, increasing the severity of ranavirus epidemics.

1. Introduction

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Mass mortality events due to infectious diseases in animals are increasing in frequency across taxa [1], many of which involve multiple environmental stressors [2-5]. The stress-induced susceptibility hypothesis posits that infectious disease-associated population declines are caused by persistent stress that increases host susceptibility to infection [5-7]. The presumed mechanism is that chronic activation of the neuroendocrine stress axis (hypothalamus-pituitaryadrenal/interrenal axis) causes negative feedback, elevating glucocorticoid levels that directly signal immune cell apoptosis [8, 9]. Since the immunomodulatory effects of glucocorticoids depend on magnitude and timing [8, 10] and can be enhancing at acute elevated levels [9, 11], a key assumption of this hypothesis is that ecological change causes *chronic* stress, which is immunosuppressive. Other, glucocorticoid-independent mechanisms have been proposed, such as limited resources needed to fight infection [12], direct cytotoxicity (reviewed in [10]), or altered neuroendocrine-immune interactions causing homeostatic imbalance (e.g., endocrine disruptors [13]). A further challenge is translating individual-level effects to the population scale, since stressors can affect multiple, countervailing epidemiological factors [14]. Due to the complexity of host-pathogen-environment interactions, there are few examples in which causal links between ecological change, physiological stress, and the likelihood of pathogen-induced mass mortality in wildlife have been established [3, 15].

To investigate the effects of environmental change on pathogen-induced mass mortality events, we focused on amphibian populations affected by road salt runoff. In the United States, the yearly application of millions of tons of deicing salts (typically NaCl) has led to persistent salinization of freshwater systems (up to 25% of seawater) [16]. Salinization is also caused by increasing drought, mining activities, agriculture, and sea-level rise [17], and has been associated

with increased disease susceptibility in other systems (e.g., Streptococcus in freshwater tilapia [18]). Although some lineages have adapted to brackish water [19], even slight increases in salinity are energetically costly to evolutionarily naïve amphibians, causing physiological stress evidenced by developmental retardation, gill inflammation, and increased cell turnover [20-22]. At concentrations below the U.S. EPA's recommended level for aquatic life (230 mg/L Cl⁻), salinization of roadside habitats in Northeastern US (average of 150 mg/L Cl⁻ in our study system) [23] reduces amphibian embryonic survival [24, 25], larval growth and activity [26, 27], and elevates plasma corticosterone concentrations in adults [26]. The evidence of salinization's influence on amphibian disease dynamics, however, is mixed. Experimental salt exposure increased trematode, Ribeiroia ondatrae, infection intensity, but reduced the number of infections, potentially through negative effects of salinity on the trematode [28]. Yet in a similar study, trematode infection was more prevalent with salt exposure [29]. On the other hand, elevated salinity reduces survival of the fungal pathogen Batrachochytridum dendrobatidis (Bd) outside its host, thus appears to reduce transmission [30]. Predicting the net effects of stressors on host-pathogen interactions is not straight forward [14], but there is an urgent need to clarify the potential outcomes as secondary salinization is rapidly changing freshwater ecosystems globally [17, 31]. Here we tested whether salinization of wetlands from salt runoff increases the frequency or severity of ranavirus epidemics in larval wood frog (Lithobates sylvaticus) populations. Ranaviruses are widely distributed, multi-host, often lethal viruses in the genus *Ranavirus* (Family *Iridoviridae*) [32]. In a range-wide survey of wood frogs, ranavirus was more prevalent in regions with greater host population density and road density [33]. Wood frog larvae are highly susceptible to ranavirus infection compared to other ranids (e.g., bullfrogs) and epidemics

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

can vary from insubstantial mortality to the complete loss of a year class [34] with no clear explanation [35]. We predicted that if salinity stress plays a role increasing the severity or frequency of ranavirus epidemics, then more severe infections and greater mortality from infection would occur in populations affected by salt runoff. Following the stress-induced susceptibility hypothesis, we predicted the chronic stress of road salt exposure—which causes a reallocation of resources from body growth to coping with greater osmoregulatory demands [26]—will cause reduced immunocompetence through one or more of the aforementioned mechanisms. Since stress can reduce the probability of an epidemic through the loss of susceptible hosts [14], we would expect salinity stress to increase transmission, either by the increasing the viral shedding or environmental persistence, both of which would lead to a greater chance of mass mortality in populations in salinized wetlands. We combined field surveys to examine correlations with pathogen presence and mass mortality events and laboratory experiments to test causal mechanisms through which salinity stress could increase the likelihood of a die-off through effects on host susceptibility and viral transmission.

2. Methods

(a) Observations of ranavirus-related mass mortality events

We surveyed 18 ephemeral ponds with similar wood frog larval densities that spanned a range of proximities to roads that receive salt application [24, 36] in the mixed hardwood forests of Yale Myers Forest (YMF) in northeastern Connecticut, USA (Fig. 1a), where the primary disturbances are associated with road maintenance and rural housing. Roadside wetlands reach conductivity (a direct measure of salinity [37]) levels of 4000 μS/cm (~2 g/L Cl⁻) due to deicing salt runoff [24, 36], causing high levels of embryonic mortality in wood frogs [24, 38] thus decreased larval

density, which would reduce our ability to detect mortality events. Selected ponds less than 250 m from paved roads had higher salinity (conductivity range: 30-360 μS/cm; ~4-95 mg/L Cl⁻) than those further away (20-50 μ S/cm; ~2-5 mg/L Cl⁻; t-test: $t_{11.44} = 2.38$, P=0.036); other abiotic factors measured in each pond were not correlated with conductivity (Table S1). We conducted weekly surveys to observe carcasses from the mid-late larval period, from 3 June to 10 July 2013, when ranavirus-related mortality events are most common [36, 39]. Die-off events were defined as > 90% of larval amphibians found dead and ranavirus infections confirmed via qPCR. We also collected euthanized larvae (n=5) and pond water samples to determine ranavirus abundance in the community (250 ml filtered water, n=3/pond; hereafter "eDNA") at two points in larval development (see [35] for methods). We disinfected all equipment and waders with 10% bleach between ponds to prevent cross-pond contamination. All statistical analyses were performed in R version 3.4.3[40]. We applied an information theoretic approach [41] (see Supplemental Methods) to estimate the importance of pond characteristics in predicting the observation of die-offs (logistic regression) and Ranavirus eDNA concentrations early and late in larval development (linear regressions). (b) The progression of naturally acquired ranavirus infections We collected larvae (Gosner [42] stages 34-39, median = 37) from 11/18 surveyed sites that had no observed mortality at the time (numbered in Fig 1a, 10-13 per pond; 2-2236 m from a major road) within 3 days of the second eDNA sample, and monitored the progression of naturally acquired infections in the laboratory. Larvae were housed individually in containers of 525 ml of dechlorinated tap water (~ 200 μS/cm). With this design, we could examine lasting effects of road runoff exposure, which has been shown to continue even after returning larvae to freshwater [43], but we cannot rule out the confounding effect of exposure to a change in water chemistry

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

when brought to the laboratory. We analyzed survival to 17 d post metamorphic climax by salinity using a mixed effects Cox proportional hazard (Cox PH) model with site as random variable (*coxme* function in *coxme* package [44]). We compared log₁₀ ranavirus liver titers at death/euthanasia using a linear mixed model with site as a random variable (*lmer* in the *lmerTest* package).

(c) Effect of elevated salinity on laboratory-raised larvae

For this and subsequent experiments, we reared animals throughout their lives in freshwater or water with added road salt (NaCl collected from a Department of Transportation salt-shed (Union, CT) to achieve desired conductivities. We collected wood frog eggs from a pond in Poughkeepsie, NY, where ranavirus die-offs had not been observed (150 μS/cm conductivity) and transported them to Washington State University where they were housed at 15° C with 15L/9D light cycle. Eggs were rinsed with clean water and divided into three treatments: freshwater (dechlorinated water with a conductivity ~200 μS/cm due to calcium ions), average salinity (1500 μS/cm; ~0.76 g/L Cl²), and near maximum salinity (3000 μS/cm; ~1.6 g/L Cl²) for wood frog ponds in this region [24, 26]). While the highest salinity treatment is more saline than those in our field survey, previous surveys have detected larvae surviving in ponds across the full range of treatment levels [24, 45]. We euthanized 10 larvae and screened for ranavirus before the exposure experiment begun; none amplified.

First, we estimated the effects of road salt exposure on mortality from infection by conducting a dose-response experiment using the salinity treatments above crossed with four doses of an FV3 ranavirus: mock exposure, low, medium and high ranavirus doses (0, 3×10³, 3×10⁴, 3×10⁵ plaque forming units per ml [pfu/ml], respectively; N=20/dose×salinity level; see Supplementary Methods). Larvae were distributed into treatment groups to standardize

developmental stage (same median: Gosner 34, and range: 30-38). We monitored survival until metamorphosis and compared case mortality (qPCR positives only) from infection in individually housed larvae using a logistic regression (*glm* with family bionmial, link = logit; in the package *stats* [40]) and a parametric survival analysis (*survreg* in the package *survival* [46]), with development stage at exposure as a covariate in both analyses.

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

We then exposed another sample of larvae from the same cohort (N=20/salinity level, Gosner stages 33-36) to a moderate dose $(3\times10^4 \text{ pfu/ml})$ or mock exposure to determine the effect of elevated salinity on infection intensity in gastrointestinal (GI) tissue, corticosterone and immune response, and shedding rate. At 6 days post-inoculation (dpi) we euthanized animals, dissected spleens (see below) and froze carcasses in liquid nitrogen. We also filtered 150 ml of water from each container to collect eDNA (following methods of [35]) to estimate viral shedding rate. Ranavirus eDNA is positively related to infectiousness in wood frog tadpoles [47], but this method does not measure concentrations of viable virions. We extracted DNA from dissected GI tissue and filters and assayed both using qPCR assays (N=20). We measured corticosterone concentrations of dissected interrenal glands (N=10/salinity×ranavirus level), where this hormone is synthesized [48], using ether lipid extractions and enzyme immunoassays. We measured splenocyte proliferation as an indicator of immune system activity by marking mitotic cells with a polycolonal antibody for phospho-histone H3 (PH3) using immunofluorohistochemistry methods. Prior research showed that elevated glucocorticoid signaling inhibits proliferation and stimulates apoptosis of splenocytes in amphibians [48]. See Supplemental Methods for details of each of these procedures.

We analyzed log₁₀ GI and eDNA ranavirus titers and log₁₀ interrenal corticosterone concentrations per mg of tissue by water treatments using univariate linear regressions for each

response (lm analyzed using Anova in the package car [49]). We compared prevalence of infection using a X^2 test of proportions. We analyzed the number of PH3 ir-cells (n=6-10/salinity× ranavirus levels) using a Poisson regression (glm in the package stats), with main effects of ranavirus and salinity, their interaction, and the area of the spleen section as a covariate. We also used Poisson regression with a three-way interaction term between corticosterone or ranavirus titer, salinity and ranavirus exposure to determine if the relationships between these variables varied among treatments.

In a third sample of larvae from this cohort, we tested whether salinity affects contact rates to increase direct transmission [50], because elevated salinity reduces larval activity [26, 28]. We counted the number of contacts between group-housed larvae (16 replicates of 10 L aquaria, 6 larvae each) that were either mock or ranavirus-exposed (high dose, same as above) in either freshwater or 3000 μS/cm saltwater. After 7 days of acclimation, we tallied the number of contacts during a 10 min-period 1 d before exposure, 2 dpi, and 5 dpi (at same time of day). We analyzed contact rate between water and ranavirus treatments using a linear mixed model with time and tank as random variables (*lmer* in the package *lmerTest* [51]).

To estimate the effects of salinity on viral persistence in the environment, we collected Gosner stages 33-38 wood frog larvae from YMF (Site 4 in Fig. 1a) when a die-off was occurring (ensuring they had high ranavirus titers). We set up 8 replicate 10 L aquaria containing 6 larvae in either freshwater or high salinity (concentrations as above). Carcasses were removed daily with a disinfected net until day 7 when we removed all larvae. Water was sampled (150 ml; filtered and quantified as above) in each aquarium at eight timepoints. We analyzed ranavirus eDNA titer using a linear regression before larvae were removed with the number of larvae

remaining at d7 and timepoint as covariates, and after larvae were removed to compare decay rate between water treatments.

200

201

3. Results

- 202 *(a) Observations of ranavirus-related mass mortality events*
- We observed ranavirus-related die-offs in seven (39%) of 18 ponds (Fig. 1a), five of which had
- elevated salinity (measured as conductivity) and were within 250 m of a paved road, and in three
- of which we observed ranavirus related die-offs in 2011 (EMH pers. obs.) and 2014 [36].
- Distance to road was in all of the best-supported models, and salinity was in next best model that
- 207 did not include distance to roads (Table S2 & S3), which were the only two variables in the best-
- 208 fit models that predicted the probability of a die-off. Specifically, the probability declined with
- distance (Fig. 1b; $\beta_{sqrt(meters)} = -0.132 \pm 0.058$, z = -2.272, P = 0.023) and marginally increased
- with salinity ($\beta_{log10(conductivity)} = 4.300 \pm 2.333$, z = 1.843, P = 0.065). Our eDNA surveys revealed
- 211 that ranavirus was ubiquitous (17/18 ponds) and ranavirus eDNA concentrations were positively
- related to salinity levels during early development (N=18; $\beta_{log10(\mu s/cm)}$ = 1.93 ± 0.690, $t_{2,15}$ =
- 2.798, P = 0.014; Fig. 1c), and during late development and once die-offs began. Specifically,
- 214 ponds with elevated salinity near major roads had ~117 times higher concentrations of ranavirus
- eDNA in pond water when larvae were nearing metamorphosis, which for some ponds coincided
- with or after observed die-offs (N=17; $\beta_{sqrt(meters)} = -0.045 \pm 0.017$, $t_{2,14} = -2.591$, P = 0.021; β
- 217 $log10(us/cm) = 1.93 \pm 0.823, t_{2,14} = 2.351, P = 0.034).$
- 218 *(b)* The progression of naturally acquired ranavirus infections
- All of the larvae collected had natural ranavirus infections at death or euthanasia, but larvae
- 220 collected from ponds with higher salinity died at a faster rate from infection (Cox PH with pond

- as a random effect: $\beta_{log10(\mu s/cm)}$ = 4.239 ± 1.943, z = 1.94, P =0.029; Fig. 1d) and had greater larval
- 222 titers at death or euthanasia (n=96; 3-11/site; $\beta_{log10(us/cm)} = 4.596 \pm 1.911$, $t_{1, 9.759} = 2.405$, P =
- 223 0.038; Fig. 1e). Because salinity and eDNA concentrations were correlated, survival was also
- 224 predicted by viral eDNA ($\beta_{log10(copies/ml)} = 1.535 \pm 0.540$, z = 2.82, P = 0.005).
- 225 (c) Effects of elevated salinity on laboratory-raised larvae
- Larvae had greater odds of dying from infection (positives only) with increasing virus dose and
- developmental stage at exposure ($\beta_{log10\ RV\ dose} = 0.899 \pm 0.160$, z = 5.637, P < 0.001; $\beta_{stage} =$
- 228 0.271 \pm 0.082, z = 3.310, P < 0.001), and with higher salinity ($\beta_{salinity} = 3.737 \pm 1.439 \times 10^{-4}$, z =
- 229 2.598, P = 0.009; Fig. 2). Exposure to high salinity increased the odds of mortality by 2.85-fold
- relative to freshwater across ranavirus doses, and reduced the estimated LD₅₀ (dose lethal to
- 50%) by nearly 4-fold compared with freshwater (from 30,200 pfu/mL in freshwater to 7,762
- pfu/mL). Larvae in the high salt treatment also died faster from infections (parametric survival
- analysis: $\beta_{salinity} = -1.23 \pm 0.45 \text{ x } 10^{-3}, z = -2.84, P = 0.005; \beta_{log 10 RV dose} = -0.851 \pm 0.124, z = -0.005; \beta_{log 10 RV dose} = -0.005; \beta_{log 10$
- 234 6.88, P < 0.001; $\beta_{stage} = -0.179 \pm 0.044$, z = -4.08, P < 0.001). In the absence of ranavirus
- exposure, salinity had a marginal effect on mortality ($\beta_{salinity} = 8.636 \pm 5.157 \times 10^{-4}$, z = 1.675, P
- = 0.094), and salinity did not affect the proportion infected (P = 0.62).
- Larvae exposed to ranavirus had elevated resting interrenal corticosterone concentrations
- 6 dpi compared to unexposed larvae (df = 1, F = 23.7, P < 0.001). Those reared in elevated
- salinity exhibited marginally elevated corticosterone (ANOVA; df = 1, F = 3.576, P = 0.064;
- Fig. 3a), but the interaction with ranavirus exposure was not significant (P = 0.27). However, we
- 241 detected main and interactive effects of salinity and ranavirus-exposure on proliferating
- splenocyte counts (GLM, Salinity: $X^2 = 16.483$, df = 1, P < 0.001; Ranavirus: $X^2 = 5.672$, df= 1,
- P = 0.017; Interaction: $X^2 = 4.210$, df = 1, P = 0.040; Fig. 3b). Salinity also increased GI viral

titers of infected larvae (df = 1, F = 7.97, P = 0.01; Fig. 3c). Splenocyte proliferation was negatively related to corticosterone levels (β_{CORT} = -0.724 ± 0.099, z = -7.31, P < 0.001), but this relationship depended on salinity and ranavirus exposure (three-way interaction: $X^2 = 8.343$, df = 1, P = 0.004, CORT-salinity interaction: $X^2 = 8.594$, df=1, P = 0.003, ranavirus-salinity interaction: $X^2 = 3.188$, df = 1, P = 0.074; Fig. 3d). Specifically, salt exposed animals had steeper slopes in the CORT-proliferation relationship compared to those raised in freshwater, except in high salt and ranavirus exposed larvae, which had a flatter slope with a lower intercept. Splenocyte proliferation was also negatively related to infection intensity, but this relationship depended on exposure to salinity (GI titer-salinity interaction: $X^2 = 6.248$, df=1, P = 0.012; GI titer: $X^2 = 7.202$, df = 1, P = 0.007; Fig. S1). Further, greater salinity exposure slowed developmental rates (Kruskal-Wallis: $X^2=7.532$, df=2, P=0.023), and splenocyte proliferation and developmental rate were negatively associated in elevated salinity but not in freshwater (Developmental rate-salinity interaction: $X^2 = 9.041$, df=1, P = 0.003, Development: P = 0.7374, Salinity: P = 0.329, RV exposure = 0.329, Fig. S1).

Our experiments designed to measure effects of salinity on routes of transmission showed that salinity treatments tended to have more extreme shedders (quantile regression ANOVA of 90^{th} from 50^{th} percentile: F = 4.879, P = 0.029; whiskers shown in Fig. 4). When we sum the ranavirus eDNA concentrations across individuals to estimate a population-level response, larvae in the high salinity treatment shed 5.5-times more virus than the freshwater group (20,417 compared to 3,715 copies total). Shed viral eDNA concentrations were correlated with GI titers $(\beta_{log10~GI~titer} = 0.954 \pm 0.075$, $t_{1,58} = 12.737$, P < 0.001). However, elevated salinity did not affect contact rates between larvae with or without ranavirus exposure ($t_{21.18} = -1.175~P = 0.253$).

We also observed an increase in viral shedding in high salinity conditions in naturally infected larvae collected from a pond during a die-off. eDNA concentrations were more than an order of magnitude greater in the high salinity treatment compared to freshwater ($\beta_{salinity} = 1.411 \pm 0.220$, $t_{1,6} = 6.407$, P < 0.001), and remained higher for 12 d after we removed larvae, though there were no differences in decay rates between salinity treatments ($t_{2,21} = 1.598$, P = 0.125, Fig. 4).

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

266

267

268

269

270

271

4. Discussion

Since first proposed to explain infectious disease-linked amphibian population declines [7], the stress-induced susceptibility hypothesis has received support linking environmental change to physiological stress and greater mortality from infection (e.g., agricultural runoff, but not natural stressors; see review [52]). Yet few studies causally link immunosuppression observed in the laboratory to prevalence of infection on the landscape [53], and none we know of link to mass mortality events. Since salinization is a persistent, energetically demanding stressor [22, 26] we hypothesized reduced immune function via glucocorticoid-dependent and -independent pathways at the individual-level; and immunocompromised populations with increased transmission and mortality from infection would pass the tipping point of more frequent and severe ranavirus epidemics (Fig. S2). Though our field observations were necessarily biased towards robust populations in order to observe mass mortality (since higher salinity roadside ponds had much lower larval densities we selected roadside populations with lower than typical salinity [26, 36]), we discovered a correlation between the probability of observed ranavirus-associated die-offs and proximity to roads and a weak correlation with salinity. This observation was corroborated with >100 times greater concentration of viral eDNA—which is tightly associated with larval

titers [35]—and greater mortality and viral loads in larvae with naturally acquired infections from more saline wetlands. Perhaps at the maximum tolerable salinity levels, embryonic mortality reduces the probability of an epidemic through the loss of susceptible individuals [14], thus future work is needed to more strongly link salt stress and mass mortality events through broader surveys. Altogether, these correlations set the stage for experimentally testing whether salinity stress plays a role in the severity and outcome of infections in wood frog larvae.

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

Our experimental exposures at environmentally relevant road salt concentrations supported several components of the stress-induced susceptibility hypothesis and revealed additional nuances. Larvae reared in elevated salinity were more likely to die from virus exposure and exhibited suppressed immune function. Previous work also found that ranavirus infection induces a stress response [54] similar to other infections [55]. However, contrary to our expectations for the involvement of the HPI axis, salinity only marginally increased resting glucocorticoid levels. Although salinity and ranavirus exposure did not interact to predict resting interrenal corticosterone concentrations, we observed a strong negative relationship between splenocyte proliferation and corticosterone that depended on salinity and ranavirus exposure. Thus, glucocorticoid-mediated immunosuppression explains in part reduced tolerance (i.e., greater infection intensity) due to salt exposure. Splenocyte proliferation is negatively affected by glucocorticoid signaling in experimental studies in anurans [48] and is an essential acute inflammatory response that ranavirus is likely capable of inhibiting [56], offering a potential explanation to the negative relationship we found between infection loads and proliferation. Additional work is needed to determine if this splenocyte response coincides with the inhibition of other antiviral responses that occurs in other vertebrates via increased GR signaling [57] to cause greater viral loads observed here. While the increased ranavirus titers and ranavirusinduced mortality of field-caught larvae is consistent with these experimental responses, future work is still needed to determine whether wild amphibian populations exposed to deicing salt also experience a rise in glucocorticoid levels and immunosuppression similar to laboratory findings to cause more frequent mortality events.

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

Our findings also show that elevated salinity suppressed splenocyte proliferation more than what could be explained by corticosterone levels. Specifically, we observed a dosedependent reduction of splenocyte proliferation with elevated salinity in ranavirus-unexposed larvae, and lower proliferation across corticosterone levels at high salinity in ranavirus-exposed larvae. These patterns suggest that elevated salinity is suppressing immunity through glucocorticoid-independent mechanisms. Since we observed reduced feeding activity in a prior study [26], salinity may affect resource acquisition or cause a delay in developmental processes that affect susceptibility to ranavirus infection. Our findings support evidence of a trade-off between the significant energetic demands of osmoregulation [19] and mounting an immune response, specifically, salt exposed larvae exhibited lower splenocyte proliferation when maintaining a comparable developmental rate, whereas the freshwater individuals exhibited no such trade-off. Similarly, migratory waterbirds experienced a trade-off between mounting an immune response and maintaining osmotic homeostasis in response to elevated salinity [58]. Additional experiments are also needed to test whether elevated internal osmolality is cytotoxic to the spleen, or whether salt changes interrenal function in ranavirus-infected larvae. This finding highlights that the stress-induced susceptibility hypothesis operates via mechanisms beyond glucocorticoid-mediated immunosuppression.

At the population level, these individual-level effects of salinity on ranavirus infection appear to drive more severe epidemics through increased disease-induced mortality rate

combined with potentially greater transmission efficiency. The increase in cumulative viral shedding we observed in high salinity would be expected to increase rates of transmission through the water [47], which corroborates our finding of greater pathogen eDNA levels in higher salinity ponds. Contact rates were not affected by salinity, but we would expect the >10fold more intense infections are more transmissible given a contact. Elevated salinity could also increase water-borne transmission through its negative effect survival of zooplankton [59], which consume infectious ranaviruses in the water [60]; or the accumulation of infected carcasses resulting from greater mortality could increase ranavirus transmission either by contact or via necrophagy [61]. Thus, small changes in habitat quality can affect multiple mechanisms that concertedly push host-pathogen systems beyond the tipping point towards a rapid, mass mortality event [1]. Such interactions are increasingly recognized as critical to understanding and perhaps mitigating the impact of infectious disease on host populations [62]. Recent work with integral projection models of dose-dependent host-parasite interactions show that small changes in host resistance or tolerance, akin to those we observed with salinity, can lead to large changes in the population-level outcomes [63].

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

Mass mortality events caused by infectious diseases are often associated with environmental changes [1, 2, 5], yet, we are just beginning to clarify generalizable patterns and mechanisms to explain the rise of emerging infectious diseases in wildlife [64]. In addition to salinity, numerous environmental changes are known to act on the neuroendocrine stress axis in amphibians (e.g., low pH [65], see review [52]). Although some cause increased mortality from infection (e.g., atrazine [66]; see review [67]), further research is needed to determine whether these individual-level effects manifest as more common or severe epidemics. Our research suggests that the mechanisms relating stressors to disease outcomes at the individual and

population level will be complex, and experiments need to be designed to test multiple hypotheses. Considering salinization has been proposed as a management strategy to provide refuge from another amphibian disease, Bd [30], our study suggests a holistic approach is required to successfully mitigate the impact of stress and disease on amphibian populations. **Data Availability** Raw data from the environmental survey and laboratory experiments are available on Dryad (doi:10.5061/dryad.ffbg79cr5). **Author Contribution** E.M.H., J.L.B., and E.J.C. contributed to study design, data analysis, and writing the manuscript. E.M.H. and B.H. contributed to data collection. **Competing Interests** The authors declare no competing interests.

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

374 References

- 1. Fey S.B., Siepielski A.M., Nusslé S., Cervantes-Yoshida K., Hwan J.L., Huber E.R., Fey
- 376 M.J., Catenazzi A., Carlson S.M. 2015 Recent shifts in the occurrence, cause, and magnitude of
- animal mass mortality events. Proceedings of the National Academy of Sciences 112(4), 1083-
- 378 1088.
- Daszak P., Cunningham A., Hyatt A. 2001 Anthropogenic environmental change and the
- emergence of infectious diseases in wildlife. *Acta tropica* **78**(2), 103-116.
- 381 3. Brearley G., Rhodes J., Bradley A., Baxter G., Seabrook L., Lunney D., Liu Y.,
- 382 McAlpine C. 2013 Wildlife disease prevalence in human-modified landscapes. *Biological*
- 383 *reviews of the Cambridge Philosophical Society* **88**(2), 427-442. (doi:10.1111/brv.12009).
- 384 4. Marcogliese D.J., Pietrock M. 2011 Combined effects of parasites and contaminants on
- animal health: parasites do matter. *Trends in parasitology* **27**(3), 123-130.
- 386 5. Patz J.A., Graczyk T.K., Geller N., Vittor A.Y. 2000 Effects of environmental change on
- emerging parasitic diseases. *International journal for parasitology* **30**(12), 1395-1405.
- Smith K.F., Sax D.F., Lafferty K.D. 2006 Evidence for the role of infectious disease in
- species extinction and endangerment. Conservation biology **20**(5), 1349-1357.
- 390 7. Carey C., Cohen N., Rollins-Smith L. 1999 Amphibian declines: an immunological
- perspective. Developmental and Comparative Immunology 23(6), 459-472. (doi:10.1016/s0145-
- 392 305x(99)00028-2).
- 8. Sapolsky R.M., Romero L.M., Munck A.U. 2000 How do glucocorticoids influence
- stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions.
- 395 Endocrine Reviews **21**(1), 55-89. (doi:10.1210/er.21.1.55).
- 396 9. Dhabhar F.S., Mcewen B.S. 1997 Acute stress enhances while chronic stress suppresses
- cell-mediated immunityin vivo: A potential role for leukocyte trafficking. *Brain, behavior, and immunity* **11**(4), 286-306.
- 399 10. Martin L.B., Hopkins W.A., Mydlarz L.D., Rohr J.R. 2010 The effects of anthropogenic
- 400 global changes on immune functions and disease resistance. *Annals of the New York Academy of*
- 401 *Sciences* **1195**(1), 129-148.
- 402 11. Bauer M.E., Perks P., Lightman S.L., Shanks N. 2001 Restraint stress is associated with
- 403 changes in glucocorticoid immunoregulation. *Physiology & behavior* **73**(4), 525-532.
- 404 12. McEwen B.S., Wingfield J.C. 2003 The concept of allostasis in biology and biomedicine.
- 405 *Hormones and Behavior* **43**(1), 2-15. (doi:10.1016/s0018-506x(02)00024-7).
- 406 13. Ahmed S.A. 2000 The immune system as a potential target for environmental estrogens
- 407 (endocrine disrupters): a new emerging field. *Toxicology* **150**(1-3), 191-206.
- 408 14. Lafferty K.D., Holt R.D. 2003 How should environmental stress affect the population
- 409 dynamics of disease? *Ecol Lett* **6**(7), 654-664.
- 410 15. Plowright R.K., Sokolow S.H., Gorman M.E., Daszak P., Foley J.E. 2008 Causal
- 411 inference in disease ecology: investigating ecological drivers of disease emergence. Frontiers in
- 412 Ecology and the Environment 6(8), 420-429.
- 413 16. Jackson R.B., Jobbagy E.G. 2005 From icy roads to salty streams. Proceedings of the
- National Academy of Sciences of the United States of America 102(41), 14487-14488.
- 415 (doi:10.1073/pnas.0507389102).
- 416 17. Herbert E.R., Boon P., Burgin A.J., Neubauer S.C., Franklin R.B., Ardón M.,
- Hopfensperger K.N., Lamers L.P., Gell P. 2015 A global perspective on wetland salinization:
- ecological consequences of a growing threat to freshwater wetlands. *Ecosphere* **6**(10), 1-43.

- 419 18. Chang P., Plumb J. 1996 Effects of salinity on Streptococcus infection of Nile tilapia,
- 420 Oreochromis niloticus. *Journal of Applied Aquaculture* **6**(1), 39-45.
- 421 19. Hopkins G.R., Brodie J.E.D. 2015 Occurrence of Amphibians in Saline Habitats: A
- Review and Evolutionary Perspective. *Herpetological Monographs* **29**(1), 1-27.
- 423 20. Gomez-Mestre I., Tejedo M., Ramayo E., Estepa J. 2004 Developmental alterations and
- osmoregulatory physiology of a larval anuran under osmotic stress. *Physiological and*
- 425 *Biochemical Zoology* **77**(2), 267-274. (doi:10.1086/378143).
- 426 21. Uchiyama M., Yoshizawa H. 1992 Salinity tolerance and structure of external and
- internal gills in tadpoles of the crab-eating frog, Rana cancrivora. *Cell and Tissue Research*
- 428 **267**(1), 35-44. (doi:10.1007/bf00318689).
- 429 22. Bernabò I., Bonacci A., Coscarelli F., Tripepi M., Brunelli E. 2013 Effects of salinity
- 430 stress on Bufo balearicus and Bufo bufo tadpoles: Tolerance, morphological gill alterations and
- Na+/K+-ATPase localization. *Aquatic toxicology* **132**, 119-133.
- 432 23. Findlay S.E.G., Kelly V.R. 2011 Emerging indirect and long-term road salt effects on
- ecosystems. In Year in Ecology and Conservation Biology (pp. 58-68.
- 434 24. Brady S.P. 2013 Microgeographic maladaptive performance and deme depression in
- response to roads and runoff. *PeerJ* 1, e163. (doi:10.7717/peerj.163).
- 436 25. Karraker N.E., Gibbs J.P. 2011 Road deicing salt irreversibly disrupts osmoregulation of
- salamander egg clutches. *Environmental Pollution* **159**(3), 833-835.
- 438 (doi:10.1016/j.envpol.2010.11.019).
- 439 26. Hall E.M., Brady S.P., Mattheus N.M., Earley R.L., Diamond M., Crespi E.J. 2017
- Physiological consequences of exposure to salinized roadside ponds on wood frog larvae and
- adults. *Biological Conservation* **209**, 98-106. (doi:10.1016/j.biocon.2017.02.013).
- 442 27. Squires Z.E., Bailey P.C., Reina R.D., Wong B.B. 2008 Environmental deterioration
- increases tadpole vulnerability to predation. *Biol Lett* **4**(4), 392-394.
- 444 (doi:10.1098/rsbl.2008.0144).
- 445 28. Milotic D., Milotic M., Koprivnikar J. 2017 Effects of road salt on larval amphibian
- susceptibility to parasitism through behavior and immunocompetence. Aquatic Toxicology 189,
- 447 42-49.
- 448 29. Buss N., Hua J. 2018 Parasite susceptibility in an amphibian host is modified by
- salinization and predators. *Environmental Pollution* **236**, 754-763.
- 450 (doi:https://doi.org/10.1016/j.envpol.2018.01.060).
- 451 30. Stockwell M., Clulow J., Mahony M. 2015 Evidence of a salt refuge: chytrid infection
- loads are suppressed in hosts exposed to salt. *Oecologia* 177(3), 901-910.
- 453 31. Hintz W.D., Relyea R.A. 2019 A review of the species, community, and ecosystem
- impacts of road salt salinisation in fresh waters. Freshwater biology **64**(6), 1081-1097.
- 455 32. Miller D., Gray M., Storfer A. 2011 Ecopathology of ranaviruses infecting amphibians.
- 456 *Viruses* **3**(11), 2351-2373.
- 457 33. Crespi E.J., Rissler L.J., Mattheus N.M., Engbrecht K., Duncan S.I., Seaborn T., Hall
- 458 E.M., Peterson J.D., Brunner J.L. 2015 Geophysiology of Wood Frogs: Landscape Patterns of
- 459 Prevalence of Disease and Circulating Hormone Concentrations across the Eastern Range.
- 460 *Integrative and Comparative Biology* **55**(4), 602-617. (doi:10.1093/icb/icv096).
- 461 34. Brunner J.L., Barnett K.E., Gosier C.J., McNulty S.A., Rubbo M.J., Kolozsvary M.B.
- 2011 Ranavirus infection in die-offs of vernal pool amphibians in New York, USA. Herpetol Rev
- 463 *Herpetological Review* **42**(1), 76-79.

- 464 35. Hall E.M., Crespi E.J., Goldberg C.S., Brunner J.L. 2016 Evaluating environmental
- DNA-based quantification of ranavirus infection in wood frog populations. *Molecular Ecology*
- 466 Resources **16**(2), 423-433. (doi:10.1111/1755-0998.12461).
- 467 36. Hall E.M., Goldberg C.S., Brunner J.L., Crespi E.J. 2018 Seasonal dynamics and
- 468 potential drivers of ranavirus epidemics in wood frog populations. *Oecologia*.
- 469 (doi:10.1007/s00442-018-4274-4).
- 470 37. Granato G.E., Smith K.P. 1999 Estimating concentrations of road-salt constituents in
- 471 highway-runoff from measurements of specific conductance, US Department of the Interior, US
- 472 Geological Survey.
- 473 38. Dananay K.L., Krynak K.L., Krynak T.J., Benard M.F. 2015 Legacy of road salt:
- 474 Apparent positive larval effects counteracted by negative postmetamorphic effects in wood frogs.
- 475 Environ Toxicol Chem **34**(10), 2417-2424. (doi:10.1002/etc.3082).
- 476 39. Green D.E., Converse K.A., Schrader A.K. 2002 Epizootiology of sixty-four amphibian
- 477 morbidity and mortality events in the USA, 1996-2001. In *Domestic Animal/Wildlife Interface*:
- 478 *Issue for Disease Control, Conservation, Sustainable Food Production, and Emerging Diseases* 479 (pp. 323-339.
- 480 40. R Core Team. 2015 R: A language and environment for statistical computing. R
- 481 Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.
- 482 41. Burnham K.P., Anderson D.R. 2004 Multimodel inference: understanding AIC and BIC
- in model selection. *Sociological methods & research* **33**(2), 261-304.
- 484 42. Gosner K.L. 1960 A Simplified Table for Staging Anuran Embryos and Larvae with
- 485 Notes on Identification. *Herpetologica* **16**(3), 183-190. (doi:10.2307/3890061).
- 486 43. Wu C.-S., Gomez-Mestre I., Kam Y.-C. 2012 Irreversibility of a bad start: early exposure
- 487 to osmotic stress limits growth and adaptive developmental plasticity. *Oecologia* **169**(1), 15-22.
- 488 44. Therneau T.M., Therneau M.T.M. 2018 Package 'coxme'. Mixed effects cox models R
- 489 package version 2.
- 490 45. Karraker N.E., Gibbs J.P., Vonesh J.R. 2008 Impacts of road deicing salt on the
- demography of vernal pool-breeding amphibians. *Ecological Applications* **18**(3), 724-734.
- 492 (doi:10.1890/07-1644.1).
- 493 46. Therneau T.M., Lumley T. 2015 Package 'survival'. R Top Doc 128.
- 494 47. Araujo A., Kirschman L., Warne R.W. 2016 Behavioural phenotypes predict disease
- susceptibility and infectiousness. *Biol Lett* **12**(8). (doi:10.1098/rsbl.2016.0480).
- 496 48. Rollins-Smith L.A. 1998 Metamorphosis and the amphibian immune system.
- 497 *Immunological Reviews* **166**, 221-230. (doi:10.1111/j.1600-065X.1998.tb01265.x).
- 498 49. Fox J., Weisberg S., Adler D., Bates D., Baud-Bovy G., Ellison S., Firth D., Friendly M.,
- 499 Gorjanc G., Graves S. 2012 Package 'car'. Vienna: R Foundation for Statistical Computing.
- 500 50. Brunner J.L., Schock D.M., Collins J.P. 2007 Transmission dynamics of the amphibian
- ranavirus Ambystoma tigrinum virus. *Dis Aquat Organ* 77(2), 87-95. (doi:10.3354/dao01845).
- 502 51. Kuznetsova A., Brockhoff P.B., Christensen R.H.B. 2017 lmerTest package: tests in
- 503 linear mixed effects models. *Journal of Statistical Software* **82**(13).
- 504 52. Rollins-Smith L.A. 2017 Amphibian immunity–stress, disease, and climate change.
- 505 Developmental & Comparative Immunology **66**, 111-119.
- 506 53. Rohr J.R., Schotthoefer A.M., Raffel T.R., Carrick H.J., Halstead N., Hoverman J.T.,
- Johnson C.M., Johnson L.B., Lieske C., Piwoni M.D. 2008 Agrochemicals increase trematode
- infections in a declining amphibian species. *Nature* **455**(7217), 1235-1239.

- 509 54. Warne R.W., Crespi E.J., Brunner J.L. 2011 Escape from the pond: stress and
- developmental responses to ranavirus infection in wood frog tadpoles. Functional Ecology 25(1),
- 511 139-146. (doi:10.1111/j.1365-2435.2010.01793.x).
- 512 55. O'Dwyer K., Dargent F., Forbes M.R., Koprivnikar J. 2019 Parasite infection leads to
- widespread glucocorticoid hormone increases in vertebrate hosts: A meta-analysis. *Journal of*
- 514 Animal Ecology.
- 515 56. De Jesús Andino F., Chen G., Li Z., Grayfer L., Robert J. 2012 Susceptibility of Xenopus
- laevis tadpoles to infection by the ranavirus Frog-Virus 3 correlates with a reduced and delayed
- innate immune response in comparison with adult frogs. *Virology* **432**(2), 435-443.
- 518 (doi:https://doi.org/10.1016/j.virol.2012.07.001).
- 519 57. Sheridan J.F., Dobbs C., Jung J., Chu X., Konstantinos A., Padgett D., Glaser R. 1998
- 520 Stress-Induced Neuroendocrine Modulation of Viral Pathogenesis and Immunity. *Annals of the*
- 521 *New York Academy of Sciences* **840**(1), 803-808.
- 522 58. Gutiérrez J.S., Abad-Gómez J.M., Villegas A., Sánchez-Guzmán J.M., Masero J.A. 2013
- Effects of salinity on the immune response of an 'osmotic generalist' bird. *Oecologia* **171**(1), 61-
- 524 69. (doi:10.1007/s00442-012-2405-x).
- 525 59. Van Meter R.J., Swan C.M. 2014 Road salts as environmental constraints in urban pond
- food webs. *PloS one* **9**(2), e90168.
- 527 60. Johnson A., Brunner J. 2014 Persistence of an amphibian ranavirus in aquatic
- 528 communities. Diseases of Aquatic Organisms 111(2), 129-138.
- 529 61. Le Sage M.J., Towey B.D., Brunner J.L. 2019 Do scavengers prevent or promote disease
- transmission? The effect of invertebrate scavenging on Ranavirus transmission. Functional
- 531 *Ecology* **33**(7), 1342-1350.
- 62. Kock R.A., Orynbayev M., Robinson S., Zuther S., Singh N.J., Beauvais W., Morgan
- E.R., Kerimbayev A., Khomenko S., Martineau H.M. 2018 Saigas on the brink:
- Multidisciplinary analysis of the factors influencing mass mortality events. *Science advances*
- 535 **4**(1), eaao2314.
- 536 63. Wilber M.Q., Knapp R.A., Toothman M., Briggs C.J. 2017 Resistance, tolerance and
- environmental transmission dynamics determine host extinction risk in a load-dependent
- 538 amphibian disease. *Ecology letters* **20**(9), 1169-1181.
- 539 64. Hing S., Narayan E.J., Thompson R.A., Godfrey S.S. 2016 The relationship between
- 540 physiological stress and wildlife disease: consequences for health and conservation. Wildlife
- 541 *Research* **43**(1), 51-60.
- 542 65. Chambers D.L., Wojdak J.M., Du P., Belden L.K. 2013 Pond Acidification May Explain
- 543 Differences in Corticosterone among Salamander Populations. *Physiological and Biochemical*
- 544 Zoology **86**(2), 224-232. (doi:10.1086/669917).
- 545 66. Forson D.D., Storfer A. 2006 Atrazine increases ranavirus susceptibility in the tiger
- salamander, Ambystoma tigrinum. Ecological Applications 16(6), 2325-2332.
- 547 67. Blaustein A., Urbina J., Snyder P., Reynolds E., Dang T., Hoverman J., Han B., Olson
- D., Searle C., Hambalek N. 2018 Effects of Emerging Infectious Diseases on Amphibians: A
- Review of Experimental Studies. *Diversity* **10**(3), 81.

551 Acknowledgements

We would like to thank the anonymous reviewers for helpful comments. This research was approved by the Animal Care and Use Committee (Protocol #04366-002) of Washington State University. Collections were approved by the Connecticut Department of Energy and Environmental Protection (Scientific collections permit #1114003) and Yale Myers Forest, and New York Department of Environmental Conservation (Collection Permit #1481). This work was supported by an EPA STAR Fellowship, Elling Endowment at Washington State University, and the Sigma-Xi Grants in Aid of Research to EMH; NSF-DEB 1139199 to JLB, NSF-DEB 1754474 to JLB & EJC and an Undergraduate Summer Research Minigrant from the College of Arts and Sciences at Washington State University to BH. J. Cundiff grew and cultured ranavirus isolates, R. Reeve completed the microscopy and imaging, and A. Storfer, J. Owen, M. Lambert, M. Atwood, and D. Skelly provided key project advice. We thank Yale Myers Forest for research sites and usage of facilities, J. Schwarz for collection and shipment of wood frog eggs used in lab experiments, and C. Goldberg for extracting eDNA samples, and her funding from Department of Defense Environmental Security Technology Certification Program. This publication was developed under STAR Fellowship Assistance Agreement no. 91767901 awarded by the U.S. Environmental Protection Agency (EPA). It has not been formally reviewed by EPA. The views expressed in this publication are solely those of E.M. Hall and co-authors, and EPA does not endorse any products or commercial services mentioned in this publication.

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

Figure Captions


Figure 1. a, Field survey of ranavirus occurrence and associated die-offs. Left: Map indicating locations of the 18 ephemeral ponds monitored weekly in June and July for ranavirus-associated die-offs of wood frog larvae in the vicinity of Yale Myers Forest in Northeastern CT, USA (top-left subset), a 32 km² managed mixed-hardwood forest. Numbered sites on map refer to the ponds from which larvae were collected to observe the progression of naturally acquired ranavirus infections (11 ponds, panels d & e). Die-offs of > 90% were observed in seven ponds while die-offs were not observed in the remaining eleven ponds. b, Proximity to major roads was correlated with the probability of observed die-offs. c, Log₁₀ Ranavirus eDNA concentration by conductivity (measure of salinity) during early larval development. (Mean ±SEM) d, Average days to death of naturally infected animals in captivity after collection with a line of predicted survival from Cox proportional hazards analysis; e, Average larval ranavirus (RV) titer of infected animals collected in the field and monitored for mortality in the laboratory by pond salinity (11 sites numbered in a; N=96; 3-11/site). Lines and shaded areas are the logistic (b) or linear (c, e) regression lines and the 95% confidence envelope. In all panels, black and gray points indicate ponds where die-offs were and were not observed, respectively.

Figure 2. Tolerance to ranavirus exposure of naive larvae reared in road salt-treated water in the laboratory. Salinity exposures reflect the average (1500 μ S/cm) and maximum (3000 μ S/cm) conductivities of roadside ponds surveyed in this region [24]. We standardized ranavirus exposures to the range of susceptible developmental stages for this species (Gosner 30-38; [54]). Points represent the proportion dead with a positive ranavirus titer (N=20 per dose/group were exposed), and lines and 95% confidence envelope (shaded area) reflect the best-fit logistic model.


Figure 3. Stress and immune responses of naïve larvae reared in road salt-treated water in a laboratory experiment at 6 d post exposure (N=20 per salinity/exposure group; exposed to 10⁵ pfu/ml: RV+; or mock exposed: RV-). **a**, LS mean interrenal corticosterone (CORT) levels (±SEM, N=10) accounting for developmental stage effects **b**, LS mean proliferating splenocyte counts (±SEM, N=6-10) accounting for spleen size and **c**, positive Log₁₀ ranavirus titers in gastrointestinal (GI) tissue (N=6-8). In a-c, dark and light bars represent ranavirus-exposed and unexposed, respectively. **d**, The relationship between proliferating splenocyte count and corticosterone levels across salinity (divided plots corresponding to x-axis of panels a-c) and ranavirus treatments (N=4-9). Lines represent best fit Poisson regressions with white points and solid lines representing RV unexposed, and black points and dotted lines representing ranavirus exposed.

Figure 4. Effects of road salt on the shedding rate and persistence of ranavirus from infected larvae from two laboratory experiments. Left: Concentration of ranavirus DNA shed into housing water (eDNA) in points and box plots showing median, lower and upper quantiles of shed DNA, whiskers extend to 10th and 90th quantiles, representing the increase in "supershedders" in the salinity treatments (N=20). Right: Average (±SEM, N=4 aquaria) Log₁₀ ranavirus eDNA concentration in aquaria water after 6 ranavirus-infected wood frog larvae were

introduced. Larvae were removed at 7 d (vertical dotted line), and eDNA sampling continued to
examine decay rates.



620 621 Figuire 1



622623 Figure 2624





Figure 3



629 Figure 4