

Tight paths in convex geometric hypergraphs

Zoltán Füredi* Tao Jiang[†] Alexandr Kostochka[‡] Dhruv Mubayi[§]
Jacques Verstraëte[¶]

Received 24 June 2018; Published 28 February 2020

Abstract: In this paper, we prove a theorem on tight paths in convex geometric hypergraphs, which is asymptotically sharp in infinitely many cases. Our geometric theorem is a common generalization of early results of Hopf and Pannwitz [12], Sutherland [19], Kupitz and Perles [16] for convex geometric graphs, as well as the classical Erdős-Gallai Theorem [6] for graphs. As a consequence, we obtain the first substantial improvement on the Turán problem for tight paths in uniform hypergraphs.

Key words and phrases: Paths, convex geometric hypergraphs, Turan number

1 Introduction

In this paper, we address extremal questions for tight paths in uniform hypergraphs and in convex geometric hypergraphs. For $k \geq 1$ and $r \geq 2$, a *tight k -path* is an r -uniform hypergraph (or simply r -graph) $P_k^r = \{v_i v_{i+1} \dots v_{i+r-1} : 0 \leq i < k\}$. Let $\text{ex}(n, P_k^r)$ denote the maximum number of edges in an n -vertex r -graph not containing a tight k -path. It appears to be difficult to determine $\text{ex}(n, P_k^r)$ in general, and even the asymptotics as $n \rightarrow \infty$ are not known. The following is a special case of a conjecture of Kalai [9] on tight trees, generalizing the well-known Erdős-Sós Conjecture [7]:

*Supported by grant K116769 from the National Research, Development and Innovation Office NKFIH and by the Simons Foundation Collaboration grant 317487.

[†]Supported by National Science Foundation award DMS-1400249.

[‡]Supported by NSF grant DMS1600592, by Arnold O. Beckman Award RB20003 of the University of Illinois at Urbana-Champaign and by grants 18-01-00353A and 19-01-00682 of the Russian Foundation for Basic Research.

[§]Supported by NSF grants DMS-1300138 and DMS-1763317.

[¶]Supported by NSF grants DMS-1556524 and DMS-1800332.

Conjecture 1.1 (Kalai). For $n \geq r \geq 2$ and $k \geq 1$, $\text{ex}(n, P_k^r) \leq \frac{k-1}{r} \binom{n}{r-1}$.

A construction based on combinatorial designs shows this conjecture if true is tight – the existence of designs was established by Keevash [13] and also more recently by Glock, Kühn, Lo and Osthus [10]. It is straightforward to see that any n -vertex r -graph H that does not contain a tight k -path has at most $(k-1) \binom{n}{r-1}$ edges. Patkós [18] gave an improvement over this bound in the case $k < 3r/4$. In the special case $k = 4$ and $r = 3$, it is shown in [8] that $\text{ex}(n, P_4^3) = \binom{n}{2}$ for all $n \geq 5$. In this paper, we give the first non-trivial upper bound on $\text{ex}(n, P_k^r)$ valid for all k and r :

Theorem 1.2. For $n \geq 1$, $r \geq 2$, and $k \geq 1$,

$$\text{ex}(n, P_k^r) \leq \begin{cases} \frac{k-1}{2} \binom{n}{r-1} & \text{if } r \text{ is even} \\ \frac{1}{2} (k + \lfloor \frac{k-1}{r} \rfloor) \binom{n}{r-1} & \text{if } r \text{ is odd} \end{cases}$$

The case $r = 2$ of this result is the well-known Erdős-Gallai Theorem [6] on paths in graphs. We prove Theorem 1.2 by introducing a novel method for extremal problems for paths in convex geometric hypergraphs.

Convex geometric hypergraphs. A *convex geometric hypergraph* (or cgh for short) is an r -graph whose vertex set is a set Ω_n of n vertices in strictly convex position in the plane, and whose edges are viewed as convex r -gons with vertices from Ω_n . Given an r -uniform cgh F , let $\text{ex}_\circ(n, F)$ denote the maximum number of edges in an n -vertex r -uniform cgh that does not contain F . Extremal problems for convex geometric graphs (or cggs for short) have been studied extensively, going back to theorems in the 1930's on disjoint line segments in the plane. We refer the reader to the papers of Braß, Károlyi and Valtr [3], Capovileas and Pach [5] and the references therein for many related extremal problems on convex geometric graphs and to Aronov, Dujmović, Morin, Ooms and da Silveira [1], Braß [2], Brass, Rote and Swanepoel [4], and Pach and Pinchasi [17] for problems in convex geometric hypergraphs, and their connections to important problems in discrete geometry, as well as the triangle-removal problem (see Aronov, Dujmović, Morin, Ooms and da Silveira [1] and Gowers and Long [11]).

Concerning results on convex geometric graphs, let M_k denote the cgg consisting of k pairwise disjoint line segments. Generalizing results of Hopf and Pannwitz [12] and Sutherland [19], Kupitz [15] and Kupitz and Perles [16] showed that for $n \geq k \geq 2$,

$$\text{ex}_\circ(n, M_k) \leq (k-1)n.$$

Perles proved the following even stronger theorem. Define a k -zigzag P_k to be a k -path $v_0 v_1 \dots v_k$ with vertices in Ω_n such that in a fixed cyclic ordering of Ω_n , the vertices appear in the order $v_0, v_2, v_4, \dots, v_5, v_3, v_1, v_0$ (see the left picture in Figure 1).

Theorem 1.3 (Perles). For $n, k \geq 1$, $\text{ex}_\circ(n, P_k) \leq (k-1)n/2$.

The bound in Theorem 1.3 is tight when k divides n since any disjoint union of cliques of order k does not contain any path with k edges. In particular, since P_{2k-1} contains M_k , Theorem 1.3 implies $\text{ex}_\circ(n, M_k) \leq \text{ex}_\circ(n, P_{2k-1}) \leq (k-1)n$. It appears to be challenging to determine for all k and r the exact value of the extremal function or the extremal cghs without k -zigzag (see Keller and Perles [14] for a discussion of extremal constructions in the case $r = 2$).

In this paper, we generalize Theorem 1.3 to convex geometric hypergraphs, and use its proof technique to prove Theorem 1.2. We let $\prec$ denote a fixed cyclic ordering of the vertices of Ω_n , and let $[u, v] = \{w \in \Omega_n : u \prec w \prec v\}$ denote a *segment* of Ω_n . If $I_1, I_2, \dots \subset \Omega_n$, then we write $I_1 \prec I_2 \prec \dots$ if all vertices of I_j are followed in the ordering $\prec$ by all vertices of I_{j+1} for $j \geq 1$. We use the following definition of a path in a convex geometric hypergraph:

Definition 1.4 (Zigzag paths). For $k \geq 1$ and even $r \geq 2$, a tight k -path $v_0 v_1 \dots v_{k+r-2}$ with vertices in Ω_n is a k -zigzag, denoted P_k^r , if there exist disjoint segments $I_0 \prec I_1 \prec \dots \prec I_{r-1}$ of Ω_n such that $\{v_i : i \equiv j \pmod{r}\} \subseteq I_j$ for $0 \leq j < r$ and

- (i) if j is even, then $v_j \prec v_{j+r} \prec v_{j+2r} \prec \dots$.
- (ii) if j is odd, then $v_j \succ v_{j+r} \succ v_{j+2r} \succ \dots$.

In words, the vertices of the zigzag with subscripts congruent to $j \pmod{r}$ appear in increasing order of subscripts if j is even, followed by the vertices with subscripts congruent to $j+1 \pmod{r}$ in decreasing order of subscripts with respect to the cyclic ordering $\prec$. In the case of graphs, a k -zigzag is simply $P_k^2 = P_k$ from Theorem 1.3. We give examples of zigzag paths P_6^2 and P_5^4 in Figure 1 below (the last edge of each path is indicated in bold).




Figure 1: Zigzag paths

The following result generalizes Theorem 1.3 to r -uniform cghs when r is even:

Theorem 1.5. Let $n, k \geq 1$, and let $r \geq 2$ be even. Then

$$\text{ex}_{\odot}(n, P_k^r) \leq \frac{(k-1)(r-1)}{r} \binom{n}{r-1}.$$

This theorem is asymptotically sharp in infinitely many cases, and is a common generalization Theorem 1.3 and the Erdős-Gallai Theorem [6]. The proof of Theorem 1.5 is also the basis for our proof of Theorem 1.2.

Organization. This paper is organized as follows. In Section 2, we give a method for extending a k -zigzag in an r -uniform cgh to a $(k+1)$ -zigzag. This is used in the short proof of Theorem 1.5 in Section 3. In Section 4, we give constructions of dense cghs without k -zigzags, and in Section 5, we prove Theorem 1.2 using the proof technique of Theorem 1.5.

Notation. We let Ω_n denote a generic set of n points in strictly convex position in the plane, and let $\prec$ denote a cyclic ordering of Ω_n . For $u, v \in \Omega_n$, we write $[u, v] = \{w : u \prec w \prec v\}$; this is the set of vertices in the segment of Ω_n from u to v (including u and v) in the ordering $\prec$. For $u, v \in \Omega_n$, let $\ell(u, v) = \min\{|[u, v]| - 1, |[v, u]| - 1\}$. In other words, $\ell(u, v)$ is the number of sides in a shortest segment of Ω_n between u and v . Throughout this paper, cghs have vertex set in Ω_n with cyclic ordering $\prec$. For an r -uniform cgh F , let $\text{ex}_{\odot}(n, F)$ denote the maximum number of edges in an r -uniform cgh on Ω_n that does not contain an ordered substructure isomorphic to F . We write $V(H)$ for the vertex set of a hypergraph H , and represent the edges as unordered lists of vertices. We identify a hypergraph H with its edge-set, denoting by $|H|$ the number of edges in H . For $v \in V(H)$, the *neighborhood* of v is $N(v) = \bigcup_{e \in H} (e \setminus \{v\})$. Let ∂H denote the *shadow* of an r -graph H , namely $\{e \setminus \{x\} : x \in e \in H\}$.

2 Extending zigzags

2.1 Extending zigzags in graphs

We start with a short proof of Theorem 1.3 for zigzags of odd length, along the lines of Perles' proof, which gives an idea of the proof of Theorem 1.5.

Proposition 2.1. *Let $k \geq 0$. If G is an n -vertex cgg with no $(2k+1)$ -zigzag, then $|G| \leq kn$.*

Proof. Proceed by induction on k ; for $k = 0$, the statement is clear. Suppose $k \geq 1$ and G is an n -vertex cgg with no $(2k+1)$ -zigzag. For $v \in V(G)$, let $f(v)$ be the first vertex of $N(v)$ after v in the ordering $\prec$. Let $E = \{vf(v) : v \in V(G)\}$. If $v_0v_1 \dots v_{2k-1}$ is a $(2k-1)$ -zigzag in $F = G \setminus E$, then $f(v_0)v_0 \dots v_{2k-1}f(v_{2k-1})$ is a $(2k+1)$ -zigzag in G . So F has no $(2k-1)$ -zigzag, and $|F| \leq (k-1)n$ by induction. Since $|E| \leq n$, $|G| = |F| + |E| \leq kn$. $\square$

A key point is that a zigzag $v_0v_1 \dots v_k$ can be extended to a $(k+1)$ -zigzag $v_0v_1 \dots v_kv$ if v is adjacent to v_k and $v \in [v_k, v_{k-1}]$ if k is even, whereas $v \in [v_{k-1}, v_k]$ if k is odd (the reader may find it helpful to refer to Figure 1). In the next section, we generalize these ideas to uniform cghs.

2.2 Extending zigzags in hypergraphs

Fixing an even $r \geq 2$, we write v_k as shorthand for $(v_{k-1}, v_k, \dots, v_{k+r-2})$. We use this as notation for the ordering of the last edge of a k -zigzag:

Definition 2.2. The *end* of a k -zigzag $v_0v_1 \dots v_{k+r-2}$ is $v_k = (v_{k-1}, v_k, \dots, v_{k+r-2})$. Let $I(v_k) = [v_{k-1}, v_k]$ if k is odd and $I(v_k) = [v_{k+r-2}, v_{k-1}]$ if k is even, and

$$X(v_k) = \{v \in I(v_k) : vv_kv_{k+1} \dots v_{k+r-2} \in H\}$$

Referring to Figure 1, in the picture on the left $X(v_6)$ is the set of v in the segment from v_6 to v_5 clockwise such that v_6v is an edge. In the picture on the right, $X(v_5)$ is the set of v in the segment from v_4 to v_5 clockwise such that $v_5v_6v_7v$ is an edge. In the next proposition, we see that any vertex in $X(v_k)$ can be used to “extend” a k -zigzag ending in v_k to a $(k+1)$ -zigzag:

Proposition 2.3. *Let $v_k \in V(H)^r$ be the end of a k -zigzag P in H . Then for any $v_{k+r-1} \in X(v)$, $P \cup \{v_kv_{k+1} \dots v_{k+r-1}\}$ is a $(k+1)$ -zigzag ending in v_{k+1} .*

Proof. Let $P = v_0v_1 \dots v_{k+r-2}$ and let $I_0 \prec I_1 \prec \dots \prec I_{r-1}$ be the segments in Definition 1.4. Let $v_{k-1} \in I_j$, so $j \equiv k-1 \pmod{r}$. If k is odd, then j is even, and the vertices of $I_j \cup I_{j+1}$ appear in the order $v_j \prec \dots \prec v_{k-1} \prec v_k \prec \dots \prec v_{j+1}$ by Definition 1.4(i). Then for any $v_{k+r-1} \in X(v_k)$, $e = v_kv_{k+1} \dots v_{k+r-2}v_{k+r-1} \in H$ and adding e to P and v_{k+r-1} to I_j before v_{k-1} in the clockwise orientation, we obtain a $(k+1)$ -zigzag. Similarly, if k is even, then j is odd so the vertices of $I_{j-1} \cup I_j$ appear in the order $v_{j-1} \prec \dots \prec v_{k+r-2} \prec v_{k-1} \prec \dots \prec v_j$ by Definition 1.4(ii), and we add e to P and v_{k+r-1} after v_{k-1} in I_j in the clockwise orientation. $\square$

Definition 2.4. Let $S_k(H)$ be the set of ends $v_k \in V(H)^r$ of k -zigzags in H , and

$$T_k(H) = \{v_k \in S_k(H) : X(v_k) = \emptyset\}.$$

Informally, $T_k(H)$ is the set of ends of k -zigzags which cannot be “extended” to $(k+1)$ -zigzags. The two key propositions for the proof of Theorem 1.5 are as follows.

Proposition 2.5. *For $v_k \in S_k(H) \setminus T_k(H)$, let $v_{k+r-1} \in X(v_k)$ be as close as possible to v_{k-1} in the segment $I(v_k)$. Then $f(v_k) = v_{k+1}$ is an injection from $S_k(H) \setminus T_k(H)$ to $S_{k+1}(H)$. In particular,*

$$|S_{k+1}(H)| \geq |S_k(H) \setminus T_k(H)|. \quad (2.1)$$

Proof. By Proposition 2.3, $f(v_k) \in S_{k+1}(H)$. Furthermore, $f(v_k) = f(w_k)$ implies $v_{k+1} = w_{k+1}$, which gives $v_i = w_i$ for $k \leq i \leq k+r-1$. If $v_{k-1} \neq w_{k-1}$, then either w_{k-1} is closer to v_{k-1} than w_{k+r-1} in $I(v_k)$, or v_{k-1} is closer to w_{k-1} than v_{k+r-1} in $I(v_k)$. These contradictions imply $v_{k-1} = w_{k-1}$, and so $v_k = w_k$ and f is an injection. $\square$

Proposition 2.6. *For $v_k \in T_k(H)$, the map $g(v_k) = (v_k, v_{k+1}, \dots, v_{k+r-2})$ is an injection from $T_k(H)$ to cyclically ordered elements of ∂H . In particular,*

$$|T_k(H)| \leq (r-1)|\partial H|. \quad (2.2)$$

Proof. If $g(v_k) = g(w_k)$, then $w_i = v_i$ for $k \leq i \leq k+r-2$. Suppose $v_{k-1} \neq w_{k-1}$. Then either $v_{k+r-2} \prec w_{k-1} \prec v_{k-1}$, and $v_{k-1} \in X(w_k)$, or $v_{k-1} \prec w_{k-1} \prec v_k$, and $w_{k-1} \in X(v_k)$. In either case, $v_k \notin T_k$ or $w_k \notin T_k$, a contradiction. So $v_{k-1} = w_{k-1}$, which implies $v_k = w_k$. $\square$

3 Proof of Theorem 1.5 on zigzags

The following theorem implies Theorem 1.5, since if H is an n -vertex r -uniform cgh not containing a k -zigzag, then $S_k(H) = \emptyset$, and we always have $|\partial H| \leq \binom{n}{r-1}$.

Theorem 3.1. *Let $k \geq 1$ and let $r \geq 2$ be even. Then for any r -uniform cgh H ,*

$$|S_k(H)| \geq r|H| - (r-1)(k-1)|\partial H|. \quad (3.1)$$

Proof. We prove (3.1) by induction on k . Let $k = 1$ and $e \in H$. By Definition 1.4(i), there are r possible orderings of the vertices of e giving a 1-zigzag: having chosen the first vertex, the ordering of the remaining vertices of e is determined. Therefore $|S_1(H)| \geq r|H|$. For the induction step, suppose $k \geq 1$ and (3.1) holds. By (2.1) and (2.2),

$$\begin{aligned} |S_{k+1}(H)| &\geq |S_k(H) \setminus T_k(H)| \geq r|H| - (r-1)(k-1)|\partial H| - |T_k(H)| \\ &\geq r|H| - (r-1)k|\partial H|. \end{aligned}$$

This proves (3.1). □

4 Stack-free constructions

Let $k \geq 1$ and let $r \geq 2$ be even. A k -stack, denoted M_k^r , consists of edges $\{v_{ir}, v_{ir+1}, \dots, v_{ir+r-1} : 0 \leq i < k\}$ where $v_0 v_1 \dots v_{(k-1)r-1}$ is an r -uniform zigzag path; in other words we pick every r th edge from a zigzag path $P_{(k-1)r+1}^r$. An example for $r = 4$ and $k = 7$ is shown below, where the extreme points on the perimeter form Ω_{28} .

There is a simple construction of an r -uniform cgh with no k -stack when k is odd with $(k-1)(r-1)\binom{n}{r-1} + O(n^{r-2})$ edges. If $k \geq 3$ is odd, let H be the cgh consisting of r -sets e from Ω_n such that $\ell(u, v) \leq k-1$ for some $u, v \in e$. It is straightforward to see that $|H| = (r-1)(k-1)\binom{n}{r-1} + O(n^{r-2})$, and H contains no k -stack since the “middle” edge e in the stack – drawn in bold in Figure 2 – has $\ell(u, v) \geq k$ for all $u, v \in e$.

In this section, we extend this construction to all values of k , thereby proving the following theorem, which may be of independent interest. In particular, this construction does not contain $P_{(k-1)r+1}^r$, and shows Theorem 1.5 is asymptotically tight for zigzags of length $1 \pmod{r}$.

Theorem 4.1. *Let $k \geq 1$ and $r \geq 2$ be even. Then*

$$\text{ex}_{\cup}(n, M_k^r) = (k-1)(r-1)\binom{n}{r-1} + O(n^{r-2}).$$

Proof. We have $\text{ex}_{\cup}(n, M_k^r) \leq (k-1)(r-1)\binom{n}{r-1}$ from Theorem 1.5. The main part of the proof is the construction of an r -uniform cgh with $(k-1)(r-1)\binom{n}{r-1} + O(n^{r-2})$ edges that does not contain a k -stack.




Figure 2: Stack

It will be convenient to let $\Omega_n = \{0, 1, 2, \dots, n-1\}$ in cyclic order, and view our edges as ordered r -tuples $(v_0, v_1, \dots, v_{r-1})$ where $0 \leq v_0 < v_1 < \dots < v_{r-1} \leq n-1$.

Our construction $H = H(n, r, k)$ has the form $H = \bigcup_{j=0}^{k-1} H_j$, where

- (i) $H_0 = \{(v_0, v_1, v_2, \dots, v_{r-1}) : v_0 = 0\}$,
- (ii) $H_j = \bigcup_{h=0}^{r-1} \{(v_0, v_1, \dots, v_{r-1}) \notin H_0 : \ell(v_h, v_{h+1}) = j\}$ for $1 \leq j \leq k-2$,
- (iii) $H_{k-1} = \bigcup_{h=1}^{r/2-1} \{(v_0, v_1, \dots, v_{r-1}) \notin H_0 : \ell(v_{2h-1}, v_{2h}) \in \{k-1, k\}\}$.

Claim 1. $|H| = (k-1)(r-1) \binom{n}{r-1} + O(n^{r-2})$.

Proof. By definition, $|H_0| = \binom{n-1}{r-1}$, and $H_0 \cap \bigcup_{j=1}^{k-1} H_j = \emptyset$. For any $j : 1 \leq j \leq k-2$, as $n \rightarrow \infty$,

$$|H_j| = (n-1) \binom{n-j-1}{r-2} + O(n^{r-2}) = (r-1) \binom{n}{r-1} + O(n^{r-2})$$

and also

$$|H_{k-1}| = 2(r/2-1) \binom{n-1}{r-1} + O(n^{r-2}) = (r-2) \binom{n}{r-1} + O(n^{r-2}).$$

If $1 \leq i < j \leq k-1$, $|H_i \cap H_j| = O(n^{r-2})$. By inclusion-exclusion,

$$|H| \geq |H_0| + \sum_{j=1}^{k-1} |H_j| - \sum_{i < j} |H_i \cap H_j| = (k-1)(r-1) \binom{n}{r-1} + O(n^{r-2}).$$

This proves the claim. □

Claim 2. $M_k^r \not\subseteq H$.

Proof. Suppose H contains a k -stack. The key is to consider the “middle” two edges of the stack, say e and f . Then the vertex 0 is in at most one of e and f . If v_0 is the first vertex of e and w_0 is the first vertex of f after 0 in the clockwise direction, then without loss of generality we may assume $v_0 < w_0$. Now consider the pairs w_1w_2, w_3w_4 up to $w_{r-1}w_r$ which are in f . We claim all of these pairs have length at least $k+1$, contradicting the definition of H , since f would then not be a member of H . To see the claim, fix $h : 1 \leq h < r/2$. Notice that there are $k/2$ edges of the stack (excluding f) which contain a pair of vertices in the segment $[w_{2h-1}, w_{2h}]$, and these pairs are vertex disjoint. However, then $\ell(w_{2h-1}, w_{2h}) \geq 2(k/2 + 1) - 1 = k + 1$, and this holds for $1 \leq h < r/2$. $\square$

5 Proof of Theorem 1.2 on tight paths

Proof for r even. Let H be an n -vertex r -graph with no tight k -path, where r is even. We aim to prove the following, which gives Theorem 1.2 for r even:

$$|H| \leq \frac{k-1}{2} |\partial H|. \quad (5.1)$$

We follow the approach used to prove Theorem 1.5 on a carefully chosen subgraph G of H . This subgraph is defined via a random partition of $V(H)$: let $s = r/2$ and let $\chi : V(G) \rightarrow \{0, 1, \dots, s-1\}$ be a random s -coloring of the vertices of H such that $P(\chi(v) = i) = 1/s$ for $0 \leq i \leq s-1$ and each vertex $v \in V(H)$, and such that vertices are colored independently. Let $B_i = \{v \in V(H) : \chi(v) = i\}$, and define the following (random) subgraph of H :

$$G = \{e \in H : |e \cap B_i| = 2 \text{ for } 0 \leq i \leq s-1\}.$$

In other words, each edge of G has two vertices in each of the sets B_i . For $0 \leq i \leq s-1$, let

$$\partial_i G = \{e \in \partial G : |e \cap B_i| = 1\} \subset \{e \in \partial H : |e \cap B_i| = 1, |e \cap B_j| = 2 \text{ for } j \neq i\}.$$

Then we have the following expected values:

$$E(|G|) = \frac{r!}{2^s s^r} |H| \quad \text{and} \quad E(|\partial_i G|) \leq \frac{(r-1)!}{2^{s-1} s^{r-1}} |\partial H|. \quad (5.2)$$

The next step is to introduce some geometric structure on G . Let $\prec$ denote a cyclic ordering of the vertices of each of $B_0, B_1, \dots, B_{s-1}$.

Definition 5.1 (Good paths). We call a tight path $v_0v_1 \dots v_{k+r-2}$ in G *good* if

- (i) for $0 \leq j < k+r-2$, $v_j, v_{j+1} \in B_i$ whenever $j \equiv 2i \pmod{r}$.
- (ii) the cyclic order in B_i is always $v_j \prec v_{j+r} \prec v_{j+2r} \prec \dots \prec v_{j+1+2r} \prec v_{j+1+r} \prec v_{j+1}$.

An r -uniform good path with k edges is shown in Figure 3, for $r = 6$ and $k = 4$.

We now follow the ideas in Section 2. By Definition 5.1(i), $v_j \in B_i$ if and only if $i = h(j) = \lfloor j/2 \rfloor \pmod{s}$. Let $i = h(k-1)$, so that $v_{k-1} \in B_i$. We write $[u, v] = \{w \in B_i : u \prec w \prec v\}$. Define $I(v_k) = [v_{k-1}, v_k] \subseteq B_i$ if k is odd and $I(v_k) = [v_{k+r-2}, v_{k-1}] \subseteq B_i$ if k is even, and

$$X(v_k) = \{v \in I(v_k) : vv_kv_{k+1} \dots v_{k+r-2} \in H\}$$




Figure 3: Good paths

Note that the definition of $X(v_k)$ is identical to that in Section 2 but with respect to the ordering $\prec$ of B_i , where $i = h(k-1)$, and in particular, $I(v_k), X(v_k) \subseteq B_i$. In Figure 3, $X(v_4)$ consists of all vertices $v \in B_1$ clockwise from v_8 to v_3 such that $v_4 v_5 v_6 v_7 v_8 v \in G$. Let $S_k(G)$ be the set of the ends of good k -paths in G , and let $T_k(G) = \{v_k \in S_k(H) : X(v_k) = \emptyset\}$.

Claim 1. For $k \geq 1$, if $i = h(k-1)$, then

$$|T_k(G)| \leq 2^{s-1} |\partial_i G|. \quad (5.3)$$

Proof. If $v_k \in S_k(G)$, then $v_{k-1} \in B_i$ since $i = h(k-1)$. For $v_k \in T_k(G)$, define $g(v_k) = (v_k, v_{k+1}, \dots, v_{k+r-2})$. Then $v_k v_{k+1} \dots v_{k+r-2} \in \partial_i G$ and $(v_k, v_{k+1}, \dots, v_{k+r-2})$ is uniquely determined by specifying the order of the pairs $\{v_k, v_{k+1}, \dots, v_{k+r-2}\} \cap B_j$ for each $j \neq i$. Therefore $g(v_k)$ injectively maps elements of $T_k(G)$ to ordered elements of $\partial_i G$, where each element of $\partial_i G$ is ordered in 2^{s-1} ways. We conclude $|T_k(G)| \leq 2^{s-1} |\partial_i G|$. $\square$

Claim 2. For $k \geq 1$,

$$|S_k(G)| \geq 2^s |G| - 2^{s-1} \sum_{i=0}^{k-2} |\partial_{h(i)} G|. \quad (5.4)$$

Proof. For $k = 1$, we observe for $e \in G$, there are two ways to label the pair $e \cap B_i$ for each $i \in [s]$, and therefore $|S_1(G)| \geq 2^s |G|$. Suppose (5.4) holds for some $k \geq 1$. Then we copy the proofs of Propositions 2.3 and 2.5 to obtain $|S_{k+1}(G)| \geq |S_k(G) \setminus T_k(G)|$. By the induction hypothesis (5.4) and Claim 1,

$$\begin{aligned} |S_{k+1}(G)| &\geq |S_k(G) \setminus T_k(G)| \geq 2^s |G| - 2^{s-1} \sum_{i=0}^{k-2} |\partial_{h(i)} G| - |T_k(G)| \\ &\geq 2^s |G| - 2^{s-1} \sum_{i=0}^{k-1} |\partial_{h(i)} G|. \end{aligned}$$

This completes the induction step and proves (5.4). $\square$

Proof of (5.1). Finally we prove (5.1). Taking expectations on both sides of (5.4), and using (5.2) and the linearity of expectation:

$$\mathbb{E}(|S_k(G)|) \geq 2^s \mathbb{E}(|G|) - 2^{s-1} \sum_{i=0}^{k-2} \mathbb{E}(|\partial_{h(i)} G|) \geq \frac{r!}{s^r} |H| - \frac{(r-1)!(k-1)}{s^{r-1}} |\partial H|. \quad (5.5)$$

Since $G \subseteq H$ has no tight k -path, $S_k(G) = \emptyset$. Using this in (5.5), we obtain (5.1). $\square$

Proof for r odd. Let H be an n -vertex r -graph containing no tight k -path. We aim to show

$$|H| \leq \frac{1}{2} \left(k + \left\lfloor \frac{k-1}{r} \right\rfloor \right) |\partial H|. \quad (5.6)$$

To prove (5.6), we reduce the case r is odd to the case r is even, and apply (5.1) from the last proof. Form the $(r+1)$ -graph H^+ by adding a set X of vertices to $V(H)$, and let $H^+ = \{\{x\} \cup e : x \in X, e \in H\}$. It is convenient to let $\phi(\ell) = \lceil (\ell+r)/(r+1) \rceil$ for $\ell \geq 1$.

It is straightforward to check that if $P = v_0 v_1 \dots v_{\ell+r-1}$ is a tight ℓ -path in H^+ , then $|V(P) \cap X| \leq \phi(\ell)$. In addition, the sequence of vertices $v_i \in V(P) \setminus X$ in increasing order of subscripts forms a tight path in H of length at least $\ell+1 - \phi(\ell)$. Setting $\ell = k + \lfloor (k-1)/r \rfloor + 1$, we have $\ell+1 - \phi(\ell) = k$, and therefore H^+ has no tight ℓ -path. By (5.1) applied to H^+ ,

$$|H^+| \leq \frac{\ell-1}{2} |\partial H^+|.$$

Since $|H^+| = |X||H|$ and $|\partial H^+| = |X||\partial H| + |H|$, we find

$$|X||H| \leq \frac{\ell-1}{2} |X||\partial H| + \frac{\ell-1}{2} |H|.$$

Choosing $|X| > (\ell-1)|H|/2$ and dividing by $|X|$, we obtain $|H| \leq (\ell-1)|\partial H|/2$. Since $(\ell-1)/2 = (k + \lfloor (k-1)/r \rfloor)/2$, this proves (5.6). $\square$

6 Concluding remarks

- It turns out using Steiner systems with arbitrarily large block sizes [10, 13]) that for each fixed $k, r \geq 2$, both of the following limits exist:

$$z(k, r) := \lim_{n \rightarrow \infty} \frac{\text{ex}_{\odot}(n, P_k^r)}{\binom{n}{r-1}} \quad \text{and} \quad p(k, r) := \lim_{n \rightarrow \infty} \frac{\text{ex}(n, P_k^r)}{\binom{n}{r-1}}.$$

The first limit is determined by Theorem 1.5 and the construction in Section 4 for $k \equiv 1 \pmod{r}$, and for $r \geq 4$ the problem is wide open in all remaining cases, even for $k = 2$.

- For $k \leq r+1$, an improvement over Theorem 1.2 is possible, slightly improving the results of Patkós [18]: we prove by induction on r that if $r \geq k-1$, the

$$\text{ex}(n, P_k^r) \leq \frac{k^2}{2r} \binom{n}{r-1}.$$

If $r = k - 1$, this follows from Theorem 1.2. Suppose $r \geq k$ and we have proved the bound for $(r - 1)$ -graphs. Let H be an r -graph with no tight k -path and pick a vertex $v \in V(H)$ contained in at least $r|H|/n$ edges of H . Consider the link hypergraph $H_v = \{e \in \partial H : e \cup \{v\} \in H\}$. Then H_v has no tight k -path, otherwise adding v to each edge we get a tight k -path in H . By induction,

$$\frac{r|H|}{n} \leq |H_v| \leq \frac{k^2}{2(r-1)} \binom{n-1}{r-2} \leq \frac{k^2}{2n} \binom{n}{r-1}$$

and this implies $|H| \leq \frac{k^2}{2r} \binom{n}{r-1}$, as required.

- It is possible when $r \geq 3$ is odd to obtain a very slight improvement over Theorem 1.2, namely

$$\text{ex}(n, P_k^r) \leq \frac{1}{r} (\sqrt{a} + \sqrt{b})^2 \binom{n}{r-1}$$

where $a = \lfloor (k-1)/r \rfloor$ and $b = (r-1)(k-1-a)/2$ and n is sufficiently large. For the purpose of comparison, we obtain

$$p(k, r) \leq k \cdot \left(\frac{1}{2} + \frac{\sqrt{2}-1}{r} + c \right)$$

where $c = O(r^{-2})$. For $r = 3$, we find that the upper bound is at most $\frac{1}{9}(3 + \sqrt{8})k \cdot \binom{n}{2}$.

- The proof in Section 5 shows that if $s = r/2$, n is a multiple of s , and G is an n -vertex r -graph such that $V(G)$ is partitioned into sets $B_0, B_1, \dots, B_{s-1}$ with $|B_i| = n/s$ and $|e \cap B_i| = 2$ for $0 \leq i < s$ and every edge $e \in G$, then $|G| \leq 2^{s-1}(k-1)(n/r)^{r-1}$, and this is asymptotically tight if $k \equiv 1 \pmod{r}$. Indeed, let $B_0, B_1, \dots, B_{s-1}$ be disjoint sets of size n/s , and let $A_i \subset B_i$ have size $(k-1)/r$. Then let G_i consist of all r -sets with one vertex in A_i , one vertex in $B_i \setminus A_i$, and two vertices in each $B_j \setminus A_j$ for $0 \leq j < s, j \neq i$. Let $G = \bigcup_{i=0}^{s-1} G_i$. Then $|e \cap f| \leq r-2$ for $e \in G_i$ and $f \in G_j$ with $i \neq j$, so if G contains a tight k -path, then the tight k -path is contained in some G_i . However, A_i is a transversal of each G_i , so G_i cannot contain a tight k -path. Therefore G has no tight k -path, and furthermore

$$|G| = \sum_{i=0}^{s-1} |G_i| = s \frac{k-1}{r} \frac{n}{s} \left(\frac{n/s}{2} \right)^{s-1} + O(n^{r-2}) = 2^{s-1}(k-1) \left(\frac{n}{r} \right)^{r-1} + O(n^{r-2}).$$

- In forthcoming work, we consider extremal problems for various other analogs of paths and matchings in the setting of convex geometric hypergraphs, having considered only zigzag paths and stacks of even uniformity in this paper.

Acknowledgments

The last author would like to thank Gil Kalai for stimulating discussions occurring at the Oberwolfach Combinatorics Conference, 2017. This research was partly conducted during AIM SQuaRes (Structured Quartet Research Ensembles) workshops, and we gratefully acknowledge the support of AIM. The authors are also grateful to a referee for carefully reading the paper and providing many valuable comments that improved the presentation.

References

- [1] B. Aronov, V. Dujmović, P. Morin, A. Ooms, L. da Silveira, More Turán-type theorems for triangles in convex point sets, *Electron. J. Combin.* 26 (2019), no. 1, Paper 1.8, 26 pp. [2](#)
- [2] P. Braß, Turán-type extremal problems for convex geometric hypergraphs, *Contemporary Mathematics*, 342, 25–34, 2004. [2](#)
- [3] P. Braß, G. Károlyi, P. Valtr, A Turán-type extremal theory of convex geometric graphs, *Goodman-Pollack Festschrift*, Springer 2003, 277–302. [2](#)
- [4] P. Braß, G. Rote, K. Swanepoel, Triangles of extremal area or perimeter in a finite planar point set, *Discrete Comp. Geom.*, 26 (1), 51–58, 2001. [2](#)
- [5] V. Capovileas, J. Pach, A Turán-type theorem for chords of a convex polygon, *J. Combin. Theory Ser. B*, 56, 9–15. [2](#)
- [6] P. Erdős and T. Gallai, On maximal paths and circuits of graphs, *Acta Math. Acad. Sci. Hungar.* **10** (1959), 337–356. [1](#), [2](#), [3](#)
- [7] P. Erdős, Extremal problems in graph theory M. Fiedler (Ed.), *Theory of Graphs and its Applications*, Academic Press (1965), pp. 29–36. [1](#)
- [8] Z. Füredi, T. Jiang, A. Kostochka, D. Mubayi, J. Verstraëte, Hypergraphs not containing a tight tree with a bounded trunk, *SIAM J. Discrete Math.* 33 (2019), no. 2, 862–873. [2](#)
- [9] P. Frankl, Z. Füredi, Exact solution of some Turán-type problems. *J. Comb. Theory Ser. A* 45 (2), (1987), 226–262. [1](#)
- [10] S. Glock, D. Kühn, A. Lo, D. Osthus, The existence of designs via iterative absorption, Preprint: <https://arxiv.org/abs/1611.06827>, 63 pages. [2](#), [10](#)
- [11] W. T. Gowers, E. Long, The length of an s -increasing sequence of r -tuples, Preprint: <https://arxiv.org/abs/1609.08688>. [2](#)
- [12] H. Hopf and E. Pannwitz: Aufgabe Nr. 167, *Jahresbericht d. Deutsch. Math. Verein.* 43 (1934), 114. [1](#), [2](#)
- [13] P. Keevash, The existence of designs II, Preprint: <https://arxiv.org/abs/1802.05900>, 39 pages. [2](#), [10](#)
- [14] C. Keller, M. Perles, On convex geometric graphs with no $k + 1$ pairwise disjoint edges. *Graphs Combin.* 32 (2016), no. 6, 2497–2514. [2](#)
- [15] Y. S. Kupitz, On Pairs of Disjoint Segments in Convex Position in the Plane, *Annals Discrete Math.*, 20 (1984), 203–208. [2](#)
- [16] Y. S. Kupitz, M. Perles, Extremal theory for convex matchings in convex geometric graphs, *Discrete Comput Geom.* 15, (1996), 195–220. [1](#), [2](#)

- [17] J. Pach, R. Pinchasi, How many unit equilateral triangles can be generated by n points in general position? *Amer. Math. Monthly* 110 (2003), 100–106. [2](#)
- [18] B. Patkós, A note on traces of set families, *Moscow Journal of Combinatorics and Number Theory* 2 (2012), 47–55. See detailed calculations in: On tight paths of fixed length, <http://www.renyi.hu/~patkos/tight-paths-fixed.pdf>. Posted on 2 Dec 2017, 3 pp. [2](#), [10](#)
- [19] J. W. Sutherland, Lösung der Aufgabe 167, *Jahresbericht Deutsch. Math.-Verein.* 45 (1935), 33–35. [1](#), [2](#)

AUTHORS

Zoltán Füredi
 Alfréd Rényi Institute of Mathematics
 Hungarian Academy of Sciences
 Reáltanoda utca 13-15.
 H-1053, Budapest, Hungary.
zfuredi@gmail.com

Tao Jiang
 Department of Mathematics
 Miami University
 Oxford, OH 45056, USA.
jiangt@miamioh.edu

Alexandr Kostochka
 University of Illinois at Urbana–Champaign
 Urbana, IL 61801
 and Sobolev Institute of Mathematics
 Novosibirsk 630090, Russia.
kostochk@math.uiuc.edu

Dhruv Mubayi
 Department of Mathematics, Statistics
 and Computer Science
 University of Illinois at Chicago
 Chicago, IL 60607.
mubayi@uic.edu

ZOLTÁN FÜREDI, TAO JIANG, ALEXANDR KOSTOCHKA, DHARUV MUBAYI AND JACQUES VERSTRAËTE

Jacques Verstraëte
Department of Mathematics
University of California at San Diego
9500 Gilman Drive, La Jolla, California 92093-0112, USA.
jverstra@math.ucsd.edu.