

MULTIPOINT DISTRIBUTION OF PERIODIC TASEP

JINHO BAIK AND ZHIPENG LIU

1. INTRODUCTION

The models in the KPZ universality class are expected to have the 1:2:3 scaling for the height fluctuations, spatial correlations, and time correlations as time $t \rightarrow \infty$. This means that the scaled two-dimensional fluctuation field

$$(1.1) \quad h_t(\gamma, \tau) := \frac{H(c_1\gamma(\tau t)^{2/3}, \tau t) - (c_2(\tau t) + c_3(\tau t)^{2/3})}{c_4(\tau t)^{1/3}}$$

of the height function $H(\ell, t)$, where ℓ is the spatial variable and t is time, is believed to converge to a universal field which depends only on the initial condition.¹ Here c_1, c_2, c_3, c_4 are model-dependent constants. Determining the limiting two-dimensional fluctuation field

$$(1.2) \quad (\gamma, \tau) \mapsto h(\gamma, \tau) := \lim_{t \rightarrow \infty} h_t(\gamma, \tau)$$

is an outstanding question.

By now there are several results for the one-point distribution. The one-point distribution of $h(\gamma, \tau)$ for fixed (γ, τ) is given by random matrix distributions (Tracy-Widom distributions) or their generalizations. The convergence is proved for a quite long list of models including PNG, TASEP, ASEP, q -TASEP, random tilings, last passage percolations, directed polymers, the KPZ equation, and so on. See, for example, [1, 2, 6, 26, 41], and the review article [11]. These models were studied using various integrable methods under standard initial conditions. See also the recent papers [13, 37] for general initial conditions.

The spatial one-dimensional process, $\gamma \mapsto h(\gamma, \tau)$ for fixed τ , is also well understood. This process is given by the Airy process and its variations. However, the convergence is proved rigorously only for a smaller number of models. It was proved for the determinantal models like PNG, TASEP, last passage percolation,² but not yet for other integrable models such as ASEP, q -TASEP, finite-temperature directed polymers, and the KPZ equation.

The two-dimensional fluctuation field, $(\gamma, \tau) \mapsto h(\gamma, \tau)$, on the other hand, is less well understood. The joint distribution is known only for the two-point distribution. In 2015, Johansson [28] considered the zero temperature Brownian semidiscrete

Received by the editors October 18, 2017, and, in revised form, October 26, 2018.

2010 *Mathematics Subject Classification*. Primary 60K35; Secondary 82C22.

The first author was supported in part by NSF grants DMS-1361782, DMS-1664531, and DMS-1664692, and the Simons Fellows program. The work was done in part when the second author was at Courant Institute, New York University.

¹For some initial conditions, such as the stationary initial condition, one may need to translate the space in the characteristic direction.

²See, for example, [3, 8–10, 25, 27, 34] for special initial conditions. See the recent paper [31] for general initial conditions for TASEP.

directed polymer and computed the limit of the two-point (in time and location) distribution.³ The limit is obtained in terms of rather complicated series involving the determinants of matrices whose entries contain the Airy kernel. The formula is simplified more recently in terms of a contour integral of a Fredholm determinant in [29] in which the author also extended his work to the directed last passage percolation model with geometric weights. Two other papers studied qualitative behaviors of the temporal correlations. Using a variational problem involving two independent Airy processes, Ferrari and Spohn [20] proved in 2016 the power law of the covariance in the time direction in the large- and small-time limits, $\tau_1/\tau_2 \rightarrow 0$ and $\tau_1/\tau_2 \rightarrow 1$. Here, τ_1 and τ_2 denote the scaled time parameters. De Nardis and Le Doussal [14] extended this work further and also, augmented by other physics arguments, computed the similar limits of the two-time distribution when one of the arguments is large. It is yet to be seen if one can deduce these results from the formula of Johansson.

The objective of this paper is to study the two-dimensional fluctuation field of *spatially periodic* KPZ models. Specifically, we evaluate the multipoint distribution of the periodic TASEP (totally asymmetric simple exclusion process) and compute a large-time limit in a certain critical regime.

We denote by L the period and by N the number of particles per period. Set $\rho = N/L$, the average density of particles. The periodic TASEP (of period L and density ρ) is defined by the occupation function $\eta_j(t)$ satisfying the spatial periodicity:

$$(1.3) \quad \eta_j(t) = \eta_{j+L}(t), \quad j \in \mathbb{Z}, \quad t \geq 0.$$

Apart from this condition, the particles follow the usual TASEP rules.

Consider the limit as $t, L, N \rightarrow \infty$ with fixed $\rho = N/L$. Since the spatial fluctuations of the usual infinite TASEP is $O(t^{2/3})$, all of the particles in the periodic TASEP are correlated when $t^{2/3} = O(L)$. We say that the periodic TASEP is in the *relaxation time scale* if

$$(1.4) \quad t = O(L^{3/2}).$$

If $t \ll L^{3/2}$, we expect that the system size has negligible effect and, therefore, the system follows the KPZ dynamics. See, for example, [5]. On the other hand, if $t \gg L^{3/2}$, then the system is basically in a finite system, and hence we expect the stationary dynamics. See, for example, [15]. Therefore, in the relaxation time scale, we predict that the KPZ dynamics and the stationary dynamics are both present.

Even though the periodic TASEP is as natural as the infinite TASEP, the one-point distribution was obtained only recently. Over the last two years, in a physics paper [36] and, independently, in mathematics papers [4, 30], the authors evaluated the one-point function of the height function in finite time and computed the large-time limit in the relaxation time scale. The one-point function follows the KPZ scaling $O(t^{1/3})$ but the limiting distribution is different from that of the infinite TASEP.⁴ This result was obtained for the three initial conditions of periodic step, flat, and stationary. Some earlier related studies can be found in physics papers

³There are non-rigorous physics papers for the two-time distribution of directed polymers [16–18]. However, another physics paper [14] indicates that the formulas in these papers are not correct.

⁴The formulas obtained in [4, 30] and [36] are similar, but different. It is yet to be checked that these formulas are the same.

[15, 21–24, 32, 33, 35], including results on the large deviation and spectral properties of the system.

In this paper, we extend the analysis of the papers [4, 30] and compute the multipoint (in time and location) distribution of the periodic TASEP with a special initial condition called the periodic step initial condition. Here we allow any number of points unlike the previous work of Johansson on the infinite TASEP. It appears that the periodicity of the model simplifies the algebraic computation compared with the infinite TASEP. In a separate paper we will consider flat and stationary initial conditions. The main results are the following:

- (1) For arbitrary initial conditions, we evaluate finite-time joint distribution functions of the periodic TASEP at multiple points in the space-time coordinates in terms of a multiple integral involving a determinant of size N . See Theorem 3.1 and Corollary 3.3.
- (2) For the periodic step initial condition, we simplify the determinant to a Fredholm determinant. See Theorem 4.6 and Corollary 4.7.
- (3) We compute the large-time limit of the multipoint (in the space-time coordinates) distribution in the relaxation time scale for the periodic TASEP with the periodic step initial condition. See Theorem 2.1.

One way of studying the usual infinite TASEP is the following. First, one computes the transition probability using the coordinate Bethe ansatz method. This means that we solve the Kolmogorov forward equation explicitly after replacing it (which contains complicated interactions between the particles) by the free evolution equation with certain boundary conditions. In [40], Schütz obtained the transition probability of the infinite TASEP. Second, one evaluates the marginal or joint distribution by taking a sum of the transition probabilities. It is important that the resulting expression should be suitable for the asymptotic analysis. This is achieved typically by obtaining a Fredholm determinant formula. In [38], Rákos and Schütz rederived the famous finite-time Fredholm determinant formula of Johansson [26] for the one-point distribution in the case of the step initial condition using this procedure. Subsequently, Sasamoto [39] and Borodin, Ferrari, Prähofer, and Sasamoto [9] obtained a Fredholm determinant formula for the joint distribution of multiple points with equal time. This was further extended by Borodin and Ferrari [7] to the points in spatial directions of each other. However, it was not extended to the case when the points are temporal directions of each other. The third step is to analyze the finite-time formula asymptotically using the method of steepest-descent. See [7, 9, 26, 39] and also a more recent paper [31]. In the KPZ 1:2:3 scaling limit, the above algebraic formulas give only the spatial process $\gamma \mapsto h(\gamma, \tau)$.

We applied the above procedure to the one-point distribution of the periodic TASEP in [4]. We obtained a formula for the transition probability, which is a periodic analogue of the formula of Schütz. Using that, we computed the finite-time one-point distribution for an arbitrary initial condition. The distribution was given by an integral of a determinant of size N . We then simplified the determinant to a Fredholm determinant for the cases of the step and flat initial conditions. The resulting expression was suitable for the asymptotic analysis. A similar computation for the stationary initial condition was carried out in [30].

In this paper, we extend the analysis of [4, 30] to multipoint distributions. For general initial conditions, we evaluate the joint distribution by taking a multiple

sum of the transition probabilities obtained in [4]. The computation can be reduced to an evaluation of a sum involving only two arbitrary points in the space-time coordinates (with different time coordinates). The main technical result of this paper, presented in Proposition 3.4, is the evaluation of this sum in a compact form. The key point, compared with the infinite TASEP [7, 9, 31], is that the points do not need to be restricted to the spatial directions.⁵ The final formula is suitable for the large-time asymptotic analysis in relaxation time scale.

If we take the period L to infinity while keeping other parameters fixed, the periodic TASEP becomes the infinite TASEP. Moreover, it is easy to check (see Section 8 below) that the joint distributions of the periodic TASEP and the infinite TASEP are equal even for fixed L if L is large enough compared with the times. Hence, the finite-time joint distribution formula obtained in this paper (Theorem 3.1 and Corollary 3.3) in fact gives a formula of the joint distribution of the infinite TASEP; see the equations (8.7) and (8.8). This formula contains an auxiliary parameter L which has no meaning in the infinite TASEP. From this observation, we find that if we take the large-time limit of our formula in the subrelaxation time scale, $t \ll L^{3/2}$, then the limit, if it exists, is the joint distribution of the two-dimensional process $h(\gamma, \tau)$ in (1.2). However, it is not clear at this moment if our formula is suitable for the asymptotic analysis in the subrelaxation time scale; the kernel of the operator in the Fredholm determinant does not seem to converge in the subrelaxation time scale while it converges in the relaxation time scale. The question of computing the limit in the subrelaxation time scale, and hence the multipoint distribution of the infinite TASEP, will be left as a later project.

This paper is organized as follows. We state the limit theorem in Section 2. The finite-time formula for general initial conditions is in Section 3. Its simplification for the periodic step initial condition is obtained in Section 4. In Section 5, we prove Proposition 3.4, the key algebraic computation. The asymptotic analysis of the formula obtained in Section 4 is carried out in Section 6, proving the result in Section 2. We discuss some properties of the limit of the joint distribution in Section 7. In Section 8 we show that the finite-time formulas obtained in Sections 3 and 4 are also valid for infinite TASEP for all large enough L .

2. LIMIT THEOREM FOR MULTIPONT DISTRIBUTION

2.1. Limit theorem. Consider the periodic TASEP of period L with N particles per period. We set $\rho = N/L$, the average particle density. We assume that the particles move to the right. Let $\eta_j(t)$ be the occupation function of periodic TASEP: $\eta_j(t) = 1$ if the site j is occupied at time t , otherwise $\eta_j(t) = 0$, and it satisfies the periodicity $\eta_j(t) = \eta_{j+L}(t)$. We consider the *periodic step initial condition* defined by

$$(2.1) \quad \eta_j(0) = \begin{cases} 1 & \text{for } -N+1 \leq j \leq 0, \\ 0 & \text{for } 1 \leq j \leq L-N, \end{cases}$$

and $\eta_{j+L}(0) = \eta_j(0)$.

⁵In the large-time limit, we add a certain restriction when the rescaled times are equal. See Theorem 2.1. The outcome of the above computation is that we find the joint distribution in terms of a multiple integral involving a determinant of size N . For the periodic step initial condition, we simplify the determinant further to a Fredholm determinant.




FIGURE 1. The pictures represent the density profile at times $t = 0$, $t = L$, and $t = 10L$ when $\rho = 1/2$. The horizontal axis is scaled down by L .




FIGURE 2. The pictures represent the limiting height function at times $t = 0.5nL$ for $n = 0, 1, 2, \dots$. The left picture is when $\rho = 1/2$ and the right picture is when $\rho = 2/5$. Both the horizontal axis (location) and the vertical axis (height) are scaled down by L .

We state the results in terms of the height function

$$(2.2) \quad h(\mathbf{p}), \quad \text{where } \mathbf{p} = \ell \mathbf{e}_1 + t \mathbf{e}_2 = (\ell, t) \in \mathbb{Z} \times \mathbb{R}_{\geq 0}.$$

Here $\mathbf{e}_1 = (1, 0)$ and $\mathbf{e}_2 = (0, 1)$ are the unit coordinate vectors in the spatial and time directions, respectively. The height function is defined by

$$(2.3) \quad h(\ell \mathbf{e}_1 + t \mathbf{e}_2) = \begin{cases} 2J_0(t) + \sum_{j=1}^{\ell} (1 - 2\eta_j(t)), & \ell \geq 1, \\ 2J_0(t), & \ell = 0, \\ 2J_0(t) - \sum_{j=\ell+1}^0 (1 - 2\eta_j(t)), & \ell \leq -1, \end{cases}$$

where $J_0(t)$ counts the number of particles jumping through the bond from 0 to 1 during the time interval $[0, t]$. The periodicity implies that

$$(2.4) \quad h((\ell + nL) \mathbf{e}_1 + t \mathbf{e}_2) = h(\ell \mathbf{e}_1 + t \mathbf{e}_2) + n(L - 2N)$$

for integers n .

See Figure 1 for the evolution of the density profile and Figure 2 for the limiting height function. Note that the step initial condition (2.1) generates shocks.⁶. By solving the Burgers' equation in a periodic domain, one could derive the explicit formulas of the density profile, the limiting height function and the shock location. These computations were done in [5].

We represent the space-time position in new coordinates. Let

$$(2.5) \quad \mathbf{e}_c := (1 - 2\rho) \mathbf{e}_1 + \mathbf{e}_2$$

⁶These shocks are generated when faster particles from the lower density region enter the higher density region and are forced to slow down. See, for example, [12, 19] for the study of similar behaviors in infinite TASEP.




FIGURE 3. Illustration of the points \mathbf{p}_j , $j = 1, \dots, m$, in the region \mathbf{R} .

be a vector parallel to the characteristic directions. If we represent $\mathbf{p} = \ell \mathbf{e}_1 + t \mathbf{e}_c$ in terms of \mathbf{e}_1 and \mathbf{e}_c , then

$$(2.6) \quad \mathbf{p} = s \mathbf{e}_1 + t \mathbf{e}_c, \quad \text{where } s = \ell - t(1 - 2\rho).$$

Consider the region

$$(2.7) \quad \begin{aligned} \mathbf{R} := & \{\ell \mathbf{e}_1 + t \mathbf{e}_2 \in \mathbb{Z} \times \mathbb{R}_{\geq 0} : 0 \leq \ell - (1 - 2\rho)t \leq L\} \\ = & \{s \mathbf{e}_1 + t \mathbf{e}_c \in \mathbb{Z} \times \mathbb{R}_{\geq 0} : 0 \leq s \leq L\}. \end{aligned}$$

See Figure 3. Due to the periodicity, the height function in \mathbf{R} determines the height function in the whole space-time plane.

The following theorem is the main asymptotic result. We take the limit as follows. We take $L, N \rightarrow \infty$ in such a way that the average density $\rho = N/L$ is fixed, or more generally ρ stays in a compact subset of the interval $(0, 1)$. We consider m distinct points $\mathbf{p}_i = s_i \mathbf{e}_1 + t_i \mathbf{e}_c$ in the space-time plane such that their temporal coordinate $t_j \rightarrow \infty$ and they satisfy the relaxation time scale $t_j = O(L^{3/2})$. The relative distances of the coordinates are scaled as in the 1:2:3 KPZ prediction: $t_i - t_j = O(L^{3/2}) = O(t_i)$, $s_i = O(L) = O(t_i^{2/3})$, and the height at each point is scaled by $O(L^{1/2}) = O(t_i^{1/3})$.

Theorem 2.1 (Limit of multipoint joint distribution for periodic TASEP). *Fix two constants c_1 and c_2 satisfying $0 < c_1 < c_2 < 1$. Let $N = N_L$ be a sequence of integers such that $c_1 L \leq N \leq c_2 L$ for all sufficiently large L . Consider the periodic TASEP of period L and average particle density $\rho = \rho_L = N/L$. Assume the periodic step initial condition (2.1). Let m be a positive integer. Fix m points $\mathbf{p}_j = (\gamma_j, \tau_j)$, $j = 1, \dots, m$, in the region*

$$(2.8) \quad \mathbf{R} := \{(\gamma, \tau) \in \mathbb{R} \times \mathbb{R}_{>0} : 0 \leq \gamma \leq 1\}.$$

Assume that

$$(2.9) \quad \tau_1 < \tau_2 < \dots < \tau_m.$$

Let $\mathbf{p}_j = s_j \mathbf{e}_1 + t_j \mathbf{e}_c$ be m points⁷ in the region \mathbf{R} shown in Figure 3, where $\mathbf{e}_1 = (1, 0)$ and $\mathbf{e}_c = (1 - 2\rho, 1)$, with

$$(2.10) \quad s_j = \gamma_j L, \quad t_j = \tau_j \frac{L^{3/2}}{\sqrt{\rho(1 - \rho)}}.$$

Then, for arbitrary fixed $x_1, \dots, x_m \in \mathbb{R}$,

$$(2.11) \quad \lim_{L \rightarrow \infty} \mathbb{P} \left(\bigcap_{j=1}^m \left\{ \frac{\mathbb{h}(\mathbf{p}_j) - (1 - 2\rho)s_j - (1 - 2\rho + 2\rho^2)t_j}{-2\rho^{1/2}(1 - \rho)^{1/2}L^{1/2}} \leq x_j \right\} \right) = \mathbb{F}(x_1, \dots, x_m; \mathbf{p}_1, \dots, \mathbf{p}_m),$$

where the function \mathbb{F} is defined in (2.15). The convergence is locally uniform in x_j, τ_j , and γ_j . If $\tau_i = \tau_{i+1}$ for some i , then (2.11) still holds if we assume that $x_i < x_{i+1}$.

Remark 2.2. Suppose that we have arbitrary m distinct points $\mathbf{p}_j = (\gamma_j, \tau_j)$ in \mathbf{R} . Then we may rearrange them so that $0 < \tau_1 \leq \dots \leq \tau_m$. If τ_j are all different, we can apply the above theorem since the result holds for arbitrarily ordered γ_j . If some of the τ_j are equal, then we may rearrange the points further so that x_j are ordered with those τ_j , and use the theorem if x_j are distinct. The only case which is not covered by the above theorem is when some of the τ_j are equal and the corresponding x_j are also equal.

The case when $m = 1$ was essentially obtained in our previous paper [4] (and also [36]). In that paper, we considered the location of a tagged particle instead of the height function, but it is straightforward to translate the result to the height function.

Remark 2.3. We will check that $\mathbb{F}(x_1, \dots, x_m; \mathbf{p}_1, \dots, \mathbf{p}_m)$ is periodic with respect to each of the space coordinates γ_j in Subsection 2.2. By this spatial periodicity, we can remove the restrictions $0 \leq \gamma_j \leq 1$ in the above theorem.

Remark 2.4. Since we expect the KPZ dynamics in the subrelaxation scale $t_j \ll L^{3/2}$, we expect that the $\tau_j \rightarrow 0$ limit of the above result should give rise to a result for the usual infinite TASEP. Concretely, we expect that the limit

$$(2.12) \quad \lim_{\tau \rightarrow 0} \mathbb{F}((\tau_1 \tau)^{1/3} x_1, \dots, (\tau_m \tau)^{1/3} x_m; (\gamma_1 (\tau_1 \tau)^{2/3}, \tau_1 \tau), \dots, (\gamma_m (\tau_m \tau)^{2/3}, \tau_m \tau))$$

exists and it is the limit of the multitime, multilocation joint distribution of the height function $\mathfrak{H}(s, t)$ of the usual TASEP with step initial condition,

$$(2.13) \quad \lim_{T \rightarrow \infty} \mathbb{P} \left(\bigcap_{j=1}^m \left\{ \frac{\mathfrak{H}(\gamma_j \tau_j^{2/3} T^{2/3}, 2\tau_j T) - \tau_j T}{-\tau_j^{1/3} T^{1/3}} \leq x_j \right\} \right).$$

In particular, we expect that when $m = 1$, (2.12) is $F_{GUE}(x_1 + \gamma_1^2/4)$, the Tracy-Widom GUE distribution; we expect that when $\tau_1 = \dots = \tau_m$, (2.12) is equal to the corresponding joint distribution of the Airy₂ process [34] $A_2(\gamma/2) - \gamma^2/4$;

⁷Since \mathbf{p}_j should have an integer value for its spatial coordinate, to be precise, we need to take the integer part of $s_j + t_j(1 - 2\rho)$ for the spatial coordinate. This small distinction does not change the result since the limits are uniform in the parameters γ_j, τ_j . Therefore, we suppress the integer value notation throughout this paper.

and when $m = 2$, (2.12) is expected to match the two-time distribution $F_{\text{two-time}}(\gamma_1/2, x_1 + \gamma_1^2/4; \gamma_2/2, x_2 + \gamma_2^2/4; \tau_1^{1/3}(\tau_2 - \tau_1)^{-1/3})$ obtained by Johansson [28, 29]. See also Section 8.

2.2. Formula of the limit of the joint distribution.

2.2.1. Definition of \mathbb{F} .

Definition 2.5. Fix a positive integer m . Let $\mathbf{p}_j = (\gamma_j, \tau_j)$ for each $j = 1, \dots, m$, where $\gamma_j \in \mathbb{R}$ and

$$(2.14) \quad 0 < \tau_1 < \tau_2 < \dots < \tau_m.$$

Define, for $x_1, \dots, x_m \in \mathbb{R}$,

$$(2.15) \quad \mathbb{F}(x_1, \dots, x_m; p_1, \dots, p_m) = \oint \dots \oint C(\mathbf{z}) D(\mathbf{z}) \frac{dz_m}{2\pi i z_m} \dots \frac{dz_1}{2\pi i z_1},$$

where $\mathbf{z} = (z_1, \dots, z_m)$ and the contours are nested circles in the complex plane satisfying $0 < |z_m| < \dots < |z_1| < 1$. Set $\mathbf{x} = (x_1, \dots, x_m)$, $\boldsymbol{\tau} = (\tau_1, \dots, \tau_m)$, and $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_m)$. The function $C(\mathbf{z}) = C(\mathbf{z}; \mathbf{x}, \boldsymbol{\tau})$ is defined by (2.21) and it depends on \mathbf{x} and $\boldsymbol{\tau}$ but not on $\boldsymbol{\gamma}$. The function $D(\mathbf{z}) = D(\mathbf{z}; \mathbf{x}, \boldsymbol{\tau}, \boldsymbol{\gamma})$ depends on all \mathbf{x} , $\boldsymbol{\tau}$, and $\boldsymbol{\gamma}$, and it is given by the Fredholm determinant $D(\mathbf{z}) = \det(1 - \mathbf{K}_1 \mathbf{K}_2)$ defined in (2.38).

The functions in the above definition satisfy the following properties. The proofs of (P1), (P3), and (P4) are scattered in this section while (P2) is proved later in Lemma 7.1.

- (P1) For each i , $C(\mathbf{z})$ is a meromorphic function of z_i in the disk $|z_i| < 1$. It has simple poles at $z_i = z_{i+1}$ for $i = 1, \dots, m-1$.
- (P2) For each i , $D(\mathbf{z})$ is analytic in the punctured disk $0 < |z_i| < 1$.
- (P3) For each i , $D(\mathbf{z})$ does not change if we replace γ_i by $\gamma_i + 1$. Therefore, \mathbb{F} is periodic, with period 1, in the parameter γ_i for each i .
- (P4) If $\tau_i = \tau_{i+1}$, the function \mathbb{F} is still well-defined for $x_i < x_{i+1}$.

Remark 2.6. It is not easy to check directly from the formula that \mathbb{F} defines a joint distribution function. Nonetheless, we may check them indirectly. From the fact that \mathbb{F} is a limit of a sequence of joint distribution functions, $0 \leq \mathbb{F} \leq 1$ and \mathbb{F} is a non-decreasing function of x_k for each k . It also follows from the fact that the joint distribution can be majorized by a marginal distribution that \mathbb{F} converges to 0 if any coordinate $x_k \rightarrow -\infty$ since it was shown in (4.10) of [4] that the $m = 1$ case is indeed a distribution function; see Lemma 7.5 below. The most difficult property to prove is the consistency which \mathbb{F} should satisfy as a coordinate $x_k \rightarrow +\infty$. We prove this property in Section 7 by finding a probabilistic interpretation of the formula of \mathbb{F} when the z_i -contours are not nested; see Theorem 7.3 and Proposition 7.4.

2.2.2. Definition of $C(\mathbf{z})$. Let $\log z$ be the principal branch of the logarithm function with cut $\mathbb{R}_{\leq 0}$. Let $\text{Li}_s(z)$ be the polylogarithm function defined by

$$(2.16) \quad \text{Li}_s(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^s} \quad \text{for } |z| < 1 \text{ and } s \in \mathbb{C}.$$

It has an analytic continuation using the formula

$$(2.17) \quad \text{Li}_s(z) = \frac{z}{\Gamma(s)} \int_0^{\infty} \frac{x^{s-1}}{e^x - z} dx \quad \text{for } z \in \mathbb{C} \setminus [1, \infty) \text{ if } \Re(s) > 0.$$

Set

$$(2.18) \quad A_1(z) = -\frac{1}{\sqrt{2\pi}} \text{Li}_{3/2}(z), \quad A_2(z) = -\frac{1}{\sqrt{2\pi}} \text{Li}_{5/2}(z).$$

For $0 < |z|, |z'| < 1$, set

$$(2.19) \quad B(z, z') = \frac{zz'}{2} \iint \frac{\eta \xi \log(-\xi + \eta)}{(e^{-\xi^2/2} - z)(e^{-\eta^2/2} - z')} \frac{d\xi}{2\pi i} \frac{d\eta}{2\pi i} = \frac{1}{4\pi} \sum_{k, k' \geq 1} \frac{z^k (z')^{k'}}{(k + k') \sqrt{kk'}},$$

where the integral contours are the vertical lines $\Re \xi = \mathfrak{a}$ and $\Re \eta = \mathfrak{b}$ with constants \mathfrak{a} and \mathfrak{b} satisfying $-\sqrt{-\log |z|} < \mathfrak{a} < 0 < \mathfrak{b} < \sqrt{-\log |z'|}$. The equality of the double integral and the series is easy to check (see (9.27)–(9.30) in [4] for a similar calculation). Note that $B(z, z') = B(z', z)$. When $z = z'$, we can also check that

$$(2.20) \quad B(z) := B(z, z) = \frac{1}{4\pi} \int_0^z \frac{(\text{Li}_{1/2}(y))^2}{y} dy.$$

Definition 2.7. Define

$$(2.21) \quad C(\mathbf{z}) := \left[\prod_{\ell=1}^m \frac{z_\ell}{z_\ell - z_{\ell+1}} \right] \left[\prod_{\ell=1}^m \frac{e^{x_\ell A_1(z_\ell) + \tau_\ell A_2(z_\ell)}}{e^{x_\ell A_1(z_{\ell+1}) + \tau_\ell A_2(z_{\ell+1})}} e^{2B(z_\ell) - 2B(z_{\ell+1}, z_\ell)} \right],$$

where we set $z_{m+1} := 0$.

Since A_1, A_2, B are analytic inside the unit circle, it is clear from the definition that $C(\mathbf{z})$ satisfies property (P1) in Subsubsection 2.2.1.

2.2.3. Definition of $D(\mathbf{z})$. The function $D(\mathbf{z})$ is given by a Fredholm determinant. Before we describe the operator and the space, we first introduce a few functions.

For $|z| < 1$, define the function

$$(2.22) \quad h(\zeta, z) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\zeta} \text{Li}_{1/2}(ze^{(\zeta^2 - y^2)/2}) dy \quad \text{for } \Re(\zeta) < 0$$

and

$$(2.23) \quad h(\zeta, z) = -\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{-\zeta} \text{Li}_{1/2}(ze^{(\zeta^2 - y^2)/2}) dy \quad \text{for } \Re(\zeta) > 0.$$

The integration contour lies in the half-plane $\Re(y) < 0$, and is given by the union of the interval $(-\infty, \Re(\pm\zeta)]$ on the real axis and the line segment from $\Re(\pm\zeta)$ to $\pm\zeta$. Since $|ze^{(\zeta^2 - y^2)/2}| < 1$ on the integration contour, $\text{Li}_{1/2}(ze^{(\zeta^2 - y^2)/2})$ is well-defined. Thus, we find that the integrals are well-defined using $\text{Li}_{1/2}(\omega) \sim \omega$ as $\omega \rightarrow 0$.

Observe the symmetry

$$(2.24) \quad h(\zeta, z) = h(-\zeta, z) \quad \text{for } \Re(\zeta) < 0.$$

We also have

$$(2.25) \quad h(\zeta, z) = \int_{-i\infty}^{i\infty} \frac{\log(1 - ze^{\omega^2/2})}{\omega - \zeta} \frac{d\omega}{2\pi i} \quad \text{for } \Re(\zeta) < 0.$$

This identity can be obtained by the power series expansion and using the fact that $\frac{1}{\sqrt{2\pi}} \int_{-\infty}^u e^{-\omega^2/2} d\omega = \int_{-i\infty}^{i\infty} \frac{e^{(-u^2 + \omega^2)/2}}{\omega - u} \frac{d\omega}{2\pi i}$ for u with $\arg(u) \in (3\pi/4, 5\pi/4)$;




FIGURE 4. The pictures represent the roots of the equation $e^{-\zeta^2/2} = z$ (dots) and the contours $\Re(\zeta^2) = -2 \log |z|$ (solid curves) for $z = 0.05e^i, 0.4e^i, 0.8e^i$, from the left to the right.

see (4.8) of [4]. From (2.24) and (2.25), we find that

$$(2.26) \quad h(\zeta, z) = O(\zeta^{-1}) \quad \text{as } \zeta \rightarrow \infty \text{ in the region } \left| \arg(\zeta) \pm \frac{\pi}{2} \right| > \epsilon$$

for any fixed z satisfying $|z| < 1$.

Let \mathbf{x} , $\boldsymbol{\tau}$, and $\boldsymbol{\gamma}$ be the parameters in Definition 2.5. We set

$$(2.27) \quad f_i(\zeta) := \begin{cases} e^{-\frac{1}{3}(\tau_i - \tau_{i-1})\zeta^3 + \frac{1}{2}(\gamma_i - \gamma_{i-1})\zeta^2 + (x_i - x_{i-1})\zeta} & \text{for } \Re(\zeta) < 0, \\ e^{\frac{1}{3}(\tau_i - \tau_{i-1})\zeta^3 - \frac{1}{2}(\gamma_i - \gamma_{i-1})\zeta^2 - (x_i - x_{i-1})\zeta} & \text{for } \Re(\zeta) > 0 \end{cases}$$

for $i = 1, \dots, m$, where we set $\tau_0 = \gamma_0 = x_0 = 0$.

Now we describe the space and the operators. For a non-zero complex number z , consider the roots ζ of the equation $e^{-\zeta^2/2} = z$. The roots are on the contour $\Re(\zeta^2) = -2 \log |z|$. It is easy to check that if $0 < |z| < 1$, the contour $\Re(\zeta^2) = -2 \log |z|$ consists of two disjoint components, one in $\Re(\zeta) > 0$ and the other in $\Re(\zeta) < 0$. See Figure 4. The asymptotes of the contours are the straight lines of slope ± 1 . For $0 < |z| < 1$, we define the discrete sets

$$(2.28) \quad \begin{aligned} L_z &:= \{\zeta \in \mathbb{C} : e^{-\zeta^2/2} = z\} \cap \{\Re(\zeta) < 0\}, \\ R_z &:= \{\zeta \in \mathbb{C} : e^{-\zeta^2/2} = z\} \cap \{\Re(\zeta) > 0\}. \end{aligned}$$

For distinct complex numbers z_1, \dots, z_m satisfying $0 < |z_i| < 1$, define the sets

$$(2.29) \quad S_1 := L_{z_1} \cup R_{z_2} \cup L_{z_3} \cup \dots \cup \begin{cases} R_{z_m} & \text{if } m \text{ is even,} \\ L_{z_m} & \text{if } m \text{ is odd,} \end{cases}$$

and

$$(2.30) \quad S_2 := R_{z_1} \cup L_{z_2} \cup R_{z_3} \cup \dots \cup \begin{cases} L_{z_m} & \text{if } m \text{ is even,} \\ R_{z_m} & \text{if } m \text{ is odd.} \end{cases}$$

See Figure 5. Now we define two operators

$$(2.31) \quad K_1 : \ell^2(S_2) \rightarrow \ell^2(S_1), \quad K_2 : \ell^2(S_1) \rightarrow \ell^2(S_2)$$

by kernels as follows. If

$$(2.32) \quad \zeta \in (L_{z_i} \cup R_{z_i}) \cap S_1 \text{ and } \zeta' \in (L_{z_j} \cup R_{z_j}) \cap S_2$$




FIGURE 5. Example of S_1 (black dots) and S_2 (white dots) when $m = 3$. The level sets are shown for visual convenience.

for some $i, j \in \{1, \dots, m\}$, then we set

$$(2.33) \quad K_1(\zeta, \zeta') = (\delta_i(j) + \delta_i(j + (-1)^i)) \frac{f_i(\zeta) e^{2h(\zeta, z_i) - h(\zeta, z_{i-(-1)^i}) - h(\zeta', z_{j-(-1)^j})}}{\zeta(\zeta - \zeta')} Q_1(j).$$

Similarly, if

$$(2.34) \quad \zeta \in (L_{z_i} \cup R_{z_i}) \cap S_2 \text{ and } \zeta' \in (L_{z_j} \cup R_{z_j}) \cap S_1$$

for some $i, j \in \{1, \dots, m\}$, then we set

$$(2.35) \quad K_2(\zeta, \zeta') = (\delta_i(j) + \delta_i(j - (-1)^i)) \frac{f_i(\zeta) e^{2h(\zeta, z_i) - h(\zeta, z_{i+(-1)^i}) - h(\zeta', z_{j+(-1)^j})}}{\zeta(\zeta - \zeta')} Q_2(j).$$

Here the delta function $\delta_i(k) = 1$ if $k = i$ or 0 otherwise. We also set $z_0 = z_{m+1} = 0$ so that

$$(2.36) \quad e^{-h(\zeta, z_0)} = e^{-h(\zeta, z_{m+1})} = 1,$$

and the functions $Q_1(j)$ and $Q_2(j)$ are defined by

$$(2.37) \quad Q_1(j) = 1 - \frac{z_{j-(-1)^j}}{z_j} \text{ and } Q_2(j) = 1 - \frac{z_{j+(-1)^j}}{z_j}.$$

Definition 2.8. Define

$$(2.38) \quad D(\mathbf{z}) := \det(1 - K_1 K_2)$$

for $\mathbf{z} = (z_1, \dots, z_m)$, where $0 < |z_i| < 1$ and z_i are distinct.

In this definition, we temporarily assumed that z_i are distinct in order to ensure that the term $\zeta - \zeta'$ in the denominators in (2.33) and (2.35) does not vanish. However, as we stated in (P2) in Subsubsection 2.2.1, $D(\mathbf{z})$ is still well-defined when the z_i are equal. See Lemma 7.1.

The definition of L_z and R_z implies that $|\arg(\zeta)| \rightarrow 3\pi/4$ as $|\zeta| \rightarrow \infty$ along $\zeta \in L_z$ and $|\arg(\zeta)| \rightarrow \pi/4$ as $|\zeta| \rightarrow \infty$ along $\zeta \in R_z$. Hence, due to the cubic term ζ^3 in (2.27), $f_i(\zeta) \rightarrow 0$ super exponentially as $|\zeta| \rightarrow \infty$ on the set $L_z \cup R_z$ if $\tau_1 < \dots < \tau_m$. Hence, using the property (2.26) of h , we see that the kernels decay super-exponentially fast as $|\zeta|, |\zeta'| \rightarrow \infty$ on the spaces. Therefore, the Fredholm determinant is well-defined if $\tau_1 < \dots < \tau_m$.

We now check property (P4). If $\tau_i = \tau_{i+1}$, the exponent of f_i has no cubic term ζ^3 . The quadratic term contributes to $O(1)$ since $|e^{-\zeta^2/2}| = |z_i|$ for $\zeta \in L_{z_i} \cup R_{z_i}$, and hence $|e^{c\zeta^2}| = O(1)$. On the other hand, the linear term in the exponent of f_i has a negative real part if $x_i < x_{i+1}$. Hence, if $\tau_i = \tau_{i+1}$ and $x_i < x_{i+1}$, then $|f_i(\zeta)| \rightarrow 0$ exponentially as $\zeta \rightarrow \infty$ along $\zeta \in S_1 \cup S_2$ and hence the kernel decays exponentially fast as $|\zeta|, |\zeta'| \rightarrow \infty$ on the spaces. Therefore, the Fredholm determinant is still well-defined if $\tau_i = \tau_{i+1}$ and $x_i < x_{i+1}$. This proves (P4).

2.3. Matrix kernel formula of K_1 and K_2 . Due to the delta functions, $K_1(\zeta, \zeta') \neq 0$ only when

$$(2.39) \quad \zeta \in L_{z_{2\ell-1}} \cup R_{z_{2\ell}} \text{ and } \zeta' \in R_{z_{2\ell-1}} \cup L_{z_{2\ell}}$$

for some integer ℓ , and similarly $K_2(\zeta, \zeta') \neq 0$ only when

$$(2.40) \quad \zeta \in L_{z_{2\ell}} \cup R_{z_{2\ell+1}} \text{ and } \zeta' \in R_{z_{2\ell}} \cup L_{z_{2\ell+1}}$$

for some integer ℓ . Thus, if we represent the kernels as $m \times m$ matrix kernels, then they have 2×2 block structures.

For example, consider the case when $m = 5$. Let us use ξ_i and η_i to represent variables in L_{z_i} and R_{z_i} , respectively:

$$(2.41) \quad \xi_i \in L_{z_i}, \quad \eta_i \in R_{z_i}.$$

The matrix kernels are given by

$$(2.42) \quad K_1 = \begin{bmatrix} k(\xi_1, \eta_1) & k(\xi_1, \xi_2) & & & \\ k(\eta_2, \eta_1) & k(\eta_2, \xi_2) & & & \\ & & k(\xi_3, \eta_3) & k(\xi_3, \xi_4) & \\ & & k(\eta_4, \eta_3) & k(\eta_4, \xi_4) & \\ & & & & k(\xi_5, \eta_5) \end{bmatrix}$$

and

$$(2.43) \quad K_2 = \begin{bmatrix} k(\eta_1, \xi_1) & & & & \\ & k(\xi_2, \eta_2) & k(\xi_2, \xi_3) & & \\ & k(\eta_3, \eta_2) & k(\eta_3, \xi_3) & & \\ & & & k(\xi_4, \eta_4) & k(\xi_4, \xi_5) \\ & & & k(\eta_5, \eta_4) & k(\eta_5, \xi_5) \end{bmatrix},$$

where the empty entries are zeros and the function k is given below. When m is odd, the structure is similar. On the other hand, when m is even, K_1 consists only of 2×2 blocks and K_2 contains an additional non-zero 1×1 block at the bottom right corner.

We now define k . For $1 \leq i \leq m-1$, writing

$$(2.44) \quad \xi = \xi_i, \quad \eta = \eta_i, \quad \xi' = \xi_{i+1}, \quad \eta' = \eta_{i+1},$$

we define

$$(2.45)$$

$$\begin{bmatrix} k(\xi, \eta) & k(\xi, \xi') \\ k(\eta', \eta) & k(\eta', \xi') \end{bmatrix} = \begin{bmatrix} f_i(\xi) & \\ & f_{i+1}(\eta') \end{bmatrix} \begin{bmatrix} \frac{e^{2h(\xi, z_i)}}{\xi e^{h(\xi, z_{i+1})}} & \\ & \frac{e^{2h(\eta', z_{i+1})}}{\eta' e^{h(\eta', z_i)}} \end{bmatrix} \begin{bmatrix} \frac{1}{\xi - \eta} & \frac{1}{\xi - \xi'} \\ \frac{1}{\eta' - \eta} & \frac{1}{\eta' - \xi'} \end{bmatrix} \\ \times \begin{bmatrix} \frac{1}{e^{h(\eta, z_{i+1})}} & \\ & \frac{1}{e^{h(\xi', z_i)}} \end{bmatrix} \begin{bmatrix} 1 - \frac{z_{i+1}}{z_i} & \\ & 1 - \frac{z_i}{z_{i+1}} \end{bmatrix}.$$

This means that

$$(2.46) \quad k(\xi_i, \eta_i) = f_i(\xi_i) \frac{e^{2h(\xi_i, z_i)}}{\xi_i e^{h(\xi_i, z_{i+1})}} \frac{1}{\xi_i - \eta_i} \frac{1}{e^{h(\eta_i, z_{i+1})}} \left(1 - \frac{z_{i+1}}{z_i}\right),$$

$$(2.47) \quad k(\xi_i, \xi_{i+1}) = f_i(\xi_i) \frac{e^{2h(\xi_i, z_i)}}{\xi_i e^{h(\xi_i, z_{i+1})}} \frac{1}{\xi_i - \xi_{i+1}} \frac{1}{e^{h(\xi_{i+1}, z_i)}} \left(1 - \frac{z_i}{z_{i+1}}\right),$$

and so on. The term $k(\xi_m, \eta_m)$ is defined by the $(1, 1)$ entry of (2.45) with $i = m$, where we set $z_{m+1} = 0$. The term $k(\eta_1, \xi_1)$ is defined by the $(2, 2)$ entry of (2.45) with $i = 0$, where we set $z_0 = 0$.

2.4. Series formulas for $D(\mathbf{z})$. We present two series formulas for the function $D(\mathbf{z})$. The first one (2.52) is the series expansion of Fredholm determinant using the block structure of the matrix kernel. The second formula (2.53) is obtained after evaluating the finite determinants in (2.52) explicitly.

To simplify formulas, we introduce the following notation.

Definition 2.9 (Notational conventions). For complex vectors $\mathbf{W} = (w_1, \dots, w_n)$ and $\mathbf{W}' = (w'_1, \dots, w'_{n'})$, we set

$$(2.48) \quad \Delta(\mathbf{W}) = \prod_{i < j} (w_j - w_i) = \det \left[w_i^{j-1} \right], \quad \Delta(\mathbf{W}; \mathbf{W}') = \prod_{\substack{1 \leq i \leq n \\ 1 \leq i' \leq n'}} (w_i - w'_{i'}).$$

For a function h of single variable, we write

$$(2.49) \quad h(\mathbf{W}) = \prod_{i=1}^n h(w_i).$$

We also use the notation

$$(2.50) \quad \Delta(S; S') = \prod_{\substack{s \in S \\ s' \in S'}} (s - s'), \quad f(S) = \prod_{s \in S} f(s)$$

for finite sets S and S' .

The next lemma follows from a general result whose proof is given in Subsection 4.3 below.

Lemma 2.10 (Series formulas for $D(\mathbf{z})$). *We have*

$$(2.51) \quad D(\mathbf{z}) = \sum_{\mathbf{n} \in (\mathbb{Z}_{\geq 0})^m} \frac{1}{(\mathbf{n}!)^2} D_{\mathbf{n}}(\mathbf{z})$$

with $\mathbf{n}! = \prod_{\ell=1}^m n_{\ell}!$ for $\mathbf{n} = (n_1, \dots, n_m)$, where $D_{\mathbf{n}}(\mathbf{z})$ can be expressed in the following two ways.

(i) We have, for $\mathbf{n} = (n_1, \dots, n_m)$,

$$(2.52) \quad D_{\mathbf{n}}(\mathbf{z}) = (-1)^{|\mathbf{n}|} \sum_{\substack{\mathbf{U}^{(\ell)} \in (\mathbf{L}_{z_\ell})^{n_\ell} \\ \mathbf{V}^{(\ell)} \in (\mathbf{R}_{z_\ell})^{n_\ell} \\ \ell=1, \dots, m}} \det [\mathbf{K}_1(\zeta_i, \zeta'_j)]_{i,j=1}^{|\mathbf{n}|} \det [\mathbf{K}_2(\zeta'_i, \zeta_j)]_{i,j=1}^{|\mathbf{n}|},$$

where $\mathbf{U} = (\mathbf{U}^{(1)}, \dots, \mathbf{U}^{(m)})$, $\mathbf{V} = (\mathbf{V}^{(1)}, \dots, \mathbf{V}^{(m)})$ with $\mathbf{U}^{(\ell)} = (\mathbf{u}_1^{(\ell)}, \dots, \mathbf{u}_{n_\ell}^{(\ell)})$, $\mathbf{V}^{(\ell)} = (\mathbf{v}_1^{(\ell)}, \dots, \mathbf{v}_{n_\ell}^{(\ell)})$, and where

$$(2.53) \quad \zeta_i = \begin{cases} \mathbf{u}_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with odd integer } \ell, \\ \mathbf{v}_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with even integer } \ell, \end{cases}$$

and

$$(2.54) \quad \zeta'_i = \begin{cases} \mathbf{v}_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with odd integer } \ell, \\ \mathbf{u}_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with even integer } \ell. \end{cases}$$

Here, we set $n_0 = 0$.

(ii) We also have

$$(2.55) \quad D_{\mathbf{n}}(\mathbf{z}) = \sum_{\substack{\mathbf{U}^{(\ell)} \in (\mathbf{L}_{z_\ell})^{n_\ell} \\ \mathbf{V}^{(\ell)} \in (\mathbf{R}_{z_\ell})^{n_\ell} \\ \ell=1, \dots, m}} d_{\mathbf{n}, \mathbf{z}}(\mathbf{U}, \mathbf{V})$$

with

$$(2.56) \quad d_{\mathbf{n}, \mathbf{z}}(\mathbf{U}, \mathbf{V}) := \left[\prod_{\ell=1}^m \frac{\Delta(\mathbf{U}^{(\ell)})^2 \Delta(\mathbf{V}^{(\ell)})^2}{\Delta(\mathbf{U}^{(\ell)}; \mathbf{V}^{(\ell)})^2} \hat{f}_\ell(\mathbf{U}^{(\ell)}) \hat{f}_\ell(\mathbf{V}^{(\ell)}) \right] \times \left[\prod_{\ell=2}^m \frac{\Delta(\mathbf{U}^{(\ell)}; \mathbf{V}^{(\ell-1)}) \Delta(\mathbf{V}^{(\ell)}; \mathbf{U}^{(\ell-1)}) e^{-h(\mathbf{V}^{(\ell)}, z_{\ell-1}) - h(\mathbf{V}^{(\ell-1)}, z_\ell)}}{\Delta(\mathbf{U}^{(\ell)}; \mathbf{U}^{(\ell-1)}) \Delta(\mathbf{V}^{(\ell)}; \mathbf{V}^{(\ell-1)}) e^{h(\mathbf{U}^{(\ell)}, z_{\ell-1}) + h(\mathbf{U}^{(\ell-1)}, z_\ell)}} \right. \\ \left. \left(1 - \frac{z_{\ell-1}}{z_\ell} \right)^{n_\ell} \left(1 - \frac{z_\ell}{z_{\ell-1}} \right)^{n_{\ell-1}} \right],$$

where

$$(2.57) \quad \hat{f}_\ell(\zeta) := \frac{1}{\zeta} f_\ell(\zeta) e^{2h(\zeta, z_\ell)}.$$

Recall (2.22), (2.23), and (2.27) for the definition of h and f_j .

Property (P3) in Subsubsection 2.2.1 follows easily from (2.56). Note that γ_i only appears in the factor $\hat{f}_\ell(\mathbf{U}^{(\ell)}) \hat{f}_\ell(\mathbf{V}^{(\ell)})$ for $\ell = i$ or $i+1$. If we replace γ_i by γ_i+1 , then $f_i(\zeta)$ and $f_{i+1}(\zeta)$ are changed by $z_i^{-1} f_i(\zeta)$ and $z_{i+1} f_{i+1}(\zeta)$ if $\Re(\zeta) < 0$, or by $z_i f_i(\zeta)$ and $z_{i+1}^{-1} f_{i+1}(\zeta)$ if $\Re(\zeta) > 0$. But $\mathbf{U}^{(\ell)}$ has the same number of components as $\mathbf{V}^{(\ell)}$ for each ℓ . Therefore $\hat{f}_\ell(\mathbf{U}^{(\ell)}) \hat{f}_\ell(\mathbf{V}^{(\ell)})$ does not change. We can also check (P3) from the original Fredholm determinant formula.

The analyticity property (P2) is proved in Lemma 7.1 later using the series formula.

3. JOINT DISTRIBUTION FUNCTION FOR THE GENERAL INITIAL CONDITION

We obtain the limit theorem of the previous section from a finite-time formula of the joint distribution. In this section, we describe a formula of the finite-time joint distribution for an arbitrary initial condition. We simplify the formula further in the next section for the case of the periodic step initial condition.

We state the results in terms of particle locations instead of the height function used in the previous section. It is easy to convert one to another; see (6.5). The particle locations are denoted by $\mathbf{x}_i(t)$, where

$$(3.1) \quad \cdots < \mathbf{x}_0(t) < \mathbf{x}_1(t) < \mathbf{x}_2(t) < \cdots.$$

Due to the periodicity of the system, we have $\mathbf{x}_i(t) = \mathbf{x}_{i+nN}(t) - nL$ for all integers n .

The periodic TASEP can be described if we keep track of N consecutive particles, say $\mathbf{x}_1(t) < \cdots < \mathbf{x}_N(t)$. If we focus only on these particles, they follow the usual TASEP rules plus the extra condition that $\mathbf{x}_N(t) < \mathbf{x}_1(t) + L$ for all t . Define the configuration space

$$(3.2) \quad \mathcal{X}_N(L) = \{(x_1, x_2, \dots, x_N) \in \mathbb{Z}^N : x_1 < x_2 < \cdots < x_N < x_1 + L\}.$$

We call the process of the N particles *TASEP in $\mathcal{X}_N(L)$* . We use the same notation $\mathbf{x}_i(t)$, $i = 1, \dots, N$, to denote the particle locations in the TASEP in $\mathcal{X}_N(L)$. We state the result for the TASEP in $\mathcal{X}_N(L)$ first and then for the periodic TASEP as a corollary.

For $z \in \mathbb{C}$, consider the polynomial of degree L given by

$$(3.3) \quad q_z(w) = w^N(w+1)^{L-N} - z^L.$$

Denote the set of the roots by

$$(3.4) \quad \mathcal{R}_z = \{w \in \mathbb{C} : q_z(w) = 0\}.$$

The roots are on the level set $|w^N(w+1)^{L-N}| = |z|^L$. It is straightforward to check the following properties of the level set. Set

$$(3.5) \quad \mathbf{r}_0 := \rho^\rho(1-\rho)^{1-\rho},$$

where, as before, $\rho = N/L$. The level set becomes larger as $|z|$ increases; see Figure 6. If $0 < |z| < \mathbf{r}_0$, the level set consists of two closed contours, one in $\Re(w) < -\rho$ enclosing the point $w = -1$ and the other in $\Re(w) > -\rho$ enclosing the point $w = 0$. When $|z| = \mathbf{r}_0$, the level set has a self-intersection at $w = -\rho$. If $|z| > \mathbf{r}_0$, then the level set is a connected closed contour. Now consider the set of roots \mathcal{R}_z . Note that if $z \neq 0$, then $-1, 0 \notin \mathcal{R}_z$. It is also easy to check that if a non-zero z satisfies $z^L \neq \mathbf{r}_0^L$, then the roots of $q_z(w)$ are all simple. On the other hand, if $z^L = \mathbf{r}_0^L$, then there is a double root at $w = -\rho$ and the remaining $L-2$ roots are simple. For the results in this section, we take z to be any non-zero complex number. But in the next section, we restrict $0 < |z| < \mathbf{r}_0$.

Theorem 3.1 (Joint distribution of TASEP in $\mathcal{X}_N(L)$ for the general initial condition). *Consider the TASEP in $\mathcal{X}_N(L)$. Let $Y = (y_1, \dots, y_N) \in \mathcal{X}_N(L)$ and assume that $(\mathbf{x}_1(0), \dots, \mathbf{x}_N(0)) = Y$. Fix a positive integer m . Let $(k_1, t_1), \dots, (k_m, t_m)$ be m distinct points in $\{1, \dots, N\} \times [0, \infty)$. Assume that $0 \leq t_1 \leq \cdots \leq t_m$. Let*




FIGURE 6. The pictures represent the roots of the equation $w^N(w+1)^{L-N} = z^L$ and the contours $|w^N(w+1)^{L-N}| = |z|^L$ with $N = 8, L = 24$ for three different values of z . The value of $|z|$ increases from the left picture to the right picture. The middle picture is when $z = \rho^\rho(1-\rho)^{1-\rho}$, where $\rho = N/L = 1/3$.

$a_i \in \mathbb{Z}$ for $1 \leq i \leq m$. Then

$$(3.6) \quad \mathbb{P}_Y(\mathbf{x}_{k_1}(t_1) \geq a_1, \dots, \mathbf{x}_{k_m}(t_m) \geq a_m) = \oint \cdots \oint \mathcal{C}(\mathbf{z}, \mathbf{k}) \mathcal{D}_Y(\mathbf{z}, \mathbf{k}, \mathbf{a}, \mathbf{t}) \frac{dz_m}{2\pi i z_m} \cdots \frac{dz_1}{2\pi i z_1},$$

where the contours are nested circles in the complex plane satisfying $0 < |z_m| < \cdots < |z_1|$. Here $\mathbf{z} = (z_1, \dots, z_m)$, $\mathbf{k} = (k_1, \dots, k_m)$, $\mathbf{a} = (a_1, \dots, a_m)$, and $\mathbf{t} = (t_1, \dots, t_m)$. The functions in the integrand are

$$(3.7) \quad \mathcal{C}(\mathbf{z}, \mathbf{k}) = (-1)^{(k_m-1)(N+1)} z_1^{(k_1-1)L} \prod_{\ell=2}^m \left[z_\ell^{(k_\ell-k_{\ell-1})L} \left(\left(\frac{z_\ell}{z_{\ell-1}} \right)^L - 1 \right)^{N-1} \right]$$

and

$$(3.8) \quad \mathcal{D}_Y(\mathbf{z}, \mathbf{k}, \mathbf{a}, \mathbf{t}) = \det \left[\sum_{\substack{w_1 \in \mathcal{R}_{z_1} \\ \vdots \\ w_m \in \mathcal{R}_{z_m}}} \frac{w_1^i (w_1+1)^{y_i-i} w_m^{-j}}{\prod_{\ell=2}^m (w_\ell - w_{\ell-1})} \prod_{\ell=1}^m \mathcal{G}_\ell(w_\ell) \right]_{i,j=1}^N$$

with

$$(3.9) \quad \mathcal{G}_\ell(w) = \frac{w(w+1)}{L(w+\rho)} \frac{w^{-k_\ell}(w+1)^{-a_\ell+k_\ell} e^{t_\ell w}}{w^{-k_{\ell-1}}(w+1)^{-a_{\ell-1}+k_{\ell-1}} e^{t_{\ell-1} w}},$$

where we set $t_0 = k_0 = a_0 = 0$.

Remark 3.2. The limiting joint distribution \mathbb{F} in the previous section was not defined for all parameters: when $\tau_i = \tau_{i+1}$, we need to put the restriction $x_i < x_{i+1}$. See property (P4) in Subsubsection 2.2.1. The finite-time joint distribution does not require such restrictions. The sums in the entries of the determinant $\mathcal{D}_Y(\mathbf{z}, \mathbf{k}, \mathbf{a}, \mathbf{t})$ are over finite sets, and hence there is no issue with the convergence. Therefore, the right-hand side of (3.6) is well-defined for all real numbers t_i and integers a_i and k_i .

Corollary 3.3 (Joint distribution of periodic TASEP for the general initial condition). *Consider the periodic TASEP with a general initial condition determined by*

$Y = (y_1, \dots, y_N) \in \mathcal{X}_N(L)$ and its periodic translations; $\mathbf{x}_{j+nN}(0) = y_j + nL$ for all $n \in \mathbb{Z}$ and $j = 1, \dots, N$. Then (3.6) holds for all $k_i \in \mathbb{Z}$ without the restriction that $k_i \in \{1, \dots, N\}$.

Proof. The particles in the periodic TASEP satisfy $\mathbf{x}_{j+nN}(t) = \mathbf{x}_j(t) + nL$ for every integer n . Hence if k_ℓ is not between 1 and N , we may translate it. This amounts to changing k_ℓ to $k_\ell + nN$ and a_ℓ to $a_\ell + nL$ for some integer n . Hence it is enough to show that the right-hand side of (3.6) is invariant under these changes. Under these changes, the term $\mathcal{C}(\mathbf{z}, \mathbf{k})$ is multiplied by the factor $z_\ell^{nNL} z_{\ell+1}^{-nNL}$ if $1 \leq \ell \leq m-1$ and by z_m^{nNL} if $\ell = m$. On the other hand, $\mathcal{G}_\ell(w_\ell)$ produces the multiplicative factor $w_\ell^{-nN} (w_\ell + 1)^{-nL+nN}$ which is z_ℓ^{-nL} by (3.4). Taking this factor outside the determinant (3.8), we cancel out the factor z_ℓ^{nNL} from $\mathcal{C}(\mathbf{z}, \mathbf{k})$. Similarly $\mathcal{G}_{\ell+1}(w_{\ell+1})$ produces a factor which cancel out $z_{\ell+1}^{-nNL}$ if $1 \leq \ell \leq m-1$. \square

Before we prove the theorem, let us comment on the analytic property of the integrand in the formula (3.6). The function $\mathcal{C}(\mathbf{z}, \mathbf{k})$ is clearly analytic in each $z_\ell \neq 0$. Consider the function $\mathcal{D}_Y(\mathbf{z}, \mathbf{k}, \mathbf{a}, \mathbf{t})$. Note that

$$(3.10) \quad \frac{d}{dw} q_z(w) = \frac{L(w + \rho)}{w(w + 1)} w^N (w + 1)^{L-N}.$$

Hence, if $F(w)$ is an analytic function of w in $\mathbb{C} \setminus \{-1, 0\}$ and

$$f(w) = F(w) w^N (w + 1)^{L-N},$$

then

$$(3.11) \quad \begin{aligned} & \sum_{w \in \mathcal{R}_z} F(w) \frac{w(w + 1)}{L(w + \rho)} \\ &= \frac{1}{2\pi i} \oint_{|w|=r} \frac{f(w)}{q_z(w)} dw - \frac{1}{2\pi i} \oint_{|w+1|=\epsilon_1} \frac{f(w)}{q_z(w)} dw - \frac{1}{2\pi i} \oint_{|w|=\epsilon_2} \frac{f(w)}{q_z(w)} dw \end{aligned}$$

for any $\epsilon_1, \epsilon_2 > 0$ and $r > \max\{\epsilon_1 + 1, \epsilon_2\}$ such that all roots of $q_z(w)$ lie in the region $\{w : \epsilon_2 < |w| < r, |w + 1| > \epsilon_1\}$. Note that $q_z(w)$ is an entire function of z for each w . Since we may take r arbitrarily large and ϵ_1, ϵ_2 arbitrarily small and positive, the right-hand side of (3.11) defines an analytic function of $z \neq 0$. Now the entries of the determinant in (3.8) are of the form

$$(3.12) \quad \sum_{\substack{w_1 \in \mathcal{R}_{z_1} \\ \vdots \\ w_m \in \mathcal{R}_{z_m}}} F(w_1, \dots, w_m) \prod_{\ell=1}^m \frac{w_\ell(w_\ell + 1)}{L(w_\ell + \rho)}$$

for a function $F(w_1, \dots, w_m)$ which is analytic in each variable in $\mathbb{C} \setminus \{-1, 0\}$ as long as $w_\ell \neq w_{\ell-1}$ for all $\ell = 2, \dots, m$. The last condition is due to the factor $\prod_{\ell=2}^m (w_\ell - w_{\ell-1})$ in the denominator. Note that if $w_\ell = w_{\ell-1}$, then $z_\ell = z_{\ell-1}$. Hence by using (3.11) m times, each entry of (3.8), and hence $\mathcal{D}_Y(\mathbf{z}, \mathbf{k}, \mathbf{a}, \mathbf{t})$, is an analytic function of each $z_\ell \neq 0$ in the region where all z_ℓ are distinct.

When $m = 1$, the product in (3.7) is set to be 1 and the formula (3.6) in this case was obtained in Proposition 6.1 in [4]. For $m \geq 2$, as we mentioned in the Introduction, we prove (3.6) by taking a multiple sum of the transition probability. The main new technical result is a summation formula and we summarize it in Proposition 3.4 below.

The transition probability was obtained in Proposition 5.1 of [4]. Denote by $\mathbb{P}_X(X'; t)$ the transition probability from $X = (x_1, \dots, x_N) \in \mathcal{X}_N(L)$ to $X' = (x'_1, \dots, x'_N) \in \mathcal{X}_N(L)$ in time t ,

$$(3.13) \quad \mathbb{P}_X(X'; t) = \oint \det \left[\frac{1}{L} \sum_{w \in \mathcal{R}_z} \frac{w^{j-i+1} (w+1)^{-x'_i+x_j+i-j} e^{tw}}{w+\rho} \right]_{i,j=1}^N \frac{dz}{2\pi iz},$$

where the integral is over any simple closed contour in $|z| > 0$ which contains 0 inside. The integrand is an analytic function of z for $z \neq 0$ by using (3.11).

Proof of Theorem 3.1. It is enough to consider $m \geq 2$. It is also sufficient to consider the case when the times are distinct, $t_1 < \dots < t_m$, because both sides of (3.6) are continuous functions of t_1, \dots, t_m . Note that (3.8) involves only finite sums.

Denoting by $X^{(\ell)} = (x_1^{(\ell)}, \dots, x_N^{(\ell)})$ the configuration of the particles at time t_ℓ , the joint distribution function on the left-hand side of (3.6) is equal to

$$(3.14) \quad \sum_{\substack{X^{(\ell)} \in \mathcal{X}_N(L) \cap \{x_{k_\ell}^{(\ell)} \geq a_\ell\} \\ \ell=1, \dots, m}} \mathbb{P}_Y(X^{(1)}; t_1) \mathbb{P}_{X^{(1)}}(X^{(2)}, t_2 - t_1) \cdots \mathbb{P}_{X^{(m-1)}}(X^{(m)}; t_m - t_{m-1}).$$

Applying the Cauchy-Binet formula to (3.13), we have

$$(3.15) \quad \mathbb{P}_X(X'; t) = \oint \sum_{W \in (\mathcal{R}_z)^N} \mathcal{L}_X(W) \mathcal{R}_{X'}(W) \mathcal{Q}(W; t) \frac{dz}{2\pi iz},$$

where, for $W = (w_1, \dots, w_N) \in \mathbb{C}$,

$$(3.16) \quad \mathcal{L}_X(W) = \det \left[w_i^j (w_i + 1)^{x_j - j} \right]_{i,j=1}^N, \quad \mathcal{R}_{X'}(W) = \det \left[w_i^{-j} (w_i + 1)^{-x'_j + j} \right]_{i,j=1}^N,$$

and

$$(3.17) \quad \mathcal{Q}(W; t) = \frac{1}{N!L^N} \prod_{i=1}^N \frac{w_i e^{tw_i}}{w_i + \rho}.$$

Here the factor $N!$ in the denominator comes from the Cauchy-Binet formula; it will eventually disappear since we will apply the Cauchy-Binet identity backward again at the end of the proof.

We insert (3.15) into (3.14) and interchange the order of the sums and the integrals. Assuming that the series converges absolutely so that the interchange is possible, the joint distribution is equal to

$$(3.18) \quad \oint \frac{dz_1}{2\pi iz_1} \cdots \oint \frac{dz_m}{2\pi iz_m} \sum_{\substack{W^{(1)} \in (\mathcal{R}_{z_1})^N \\ \vdots \\ W^{(m)} \in (\mathcal{R}_{z_m})^N}} \mathcal{P}(W^{(1)}, \dots, W^{(m)}) \prod_{\ell=1}^m \mathcal{Q}(W^{(\ell)}; t_\ell - t_{\ell-1}),$$

where $W^{(\ell)} = (w_1^{(\ell)}, \dots, w_N^{(\ell)})$ and

$$(3.19) \quad \begin{aligned} & \mathcal{P}(W^{(1)}, \dots, W^{(m)}) \\ &= \mathcal{L}_Y(W^{(1)}) \left[\prod_{\ell=1}^{m-1} \mathcal{H}_{k_\ell, a_\ell}(W^{(\ell)}; W^{(\ell+1)}) \right] \left[\sum_{\substack{X \in \mathcal{X}_N(L) \\ x_{k_m} \geq a_m}} \mathcal{R}_X(W^{(m)}) \right]. \end{aligned}$$

Here we set

$$(3.20) \quad \mathcal{H}_{k,a}(W; W') := \sum_{X \in \mathcal{X}_N(L) \cap \{x_k \geq a\}} \mathcal{R}_X(W) \mathcal{L}_X(W')$$

for a pair of complex vectors $W = (w_1, \dots, w_N)$ and $W' = (w'_1, \dots, w'_N)$. Let us now show that it is possible to exchange the sums and integrals if we take the z_i -contours properly. We first consider the convergence of (3.20) and the sum in (3.19). Note that, shifting the summation variable X to $X - (b, \dots, b)$,

$$(3.21) \quad \begin{aligned} & \sum_{X \in \mathcal{X}_N(L) \cap \{x_k = b\}} \mathcal{R}_X(W) \mathcal{L}_X(W') \\ &= \left[\sum_{Y \in \mathcal{X}_N(X) \cap \{y_k = 0\}} \mathcal{R}_Y(W) \mathcal{L}_Y(W') \right] \left(\prod_{j=1}^N \frac{w'_j + 1}{w_j + 1} \right)^b. \end{aligned}$$

The right-hand side of (3.20) is the sum of the above formula over $b \geq a$. Hence (3.20) converges absolutely and the convergence is uniform for w_i, w'_i if $\prod_{j=1}^N |\frac{w'_j + 1}{w_j + 1}|$ is in a compact subset of $[0, 1)$. Similarly, the sum of $\mathcal{R}_X(W^{(m)})$ in (3.19) converges if $\prod_{j=1}^N |w_j^{(m)} + 1| > 1$. Therefore, (3.19) converges absolutely if the intermediate variables $W^{(\ell)} = (w_1^{(\ell)}, \dots, w_N^{(\ell)})$ satisfy

$$(3.22) \quad \prod_{j=1}^N |w_j^{(1)} + 1| > \prod_{j=1}^N |w_j^{(2)} + 1| > \dots > \prod_{j=1}^N |w_j^{(m)} + 1| > 1.$$

We now show that it is possible to choose the contours of z_i so that (3.22) is achieved. Since $W^{(\ell)} \in (R_{z_\ell})^N$, $w_j^{(\ell)}$ satisfies the equation $|w^N(w+1)^{L-N}| = |z_\ell^L|$. Hence $|w_j^{(\ell)}| = |z_j| + O(1)$ as $|z_j| \rightarrow \infty$. Therefore, if we take the contours $|z_\ell| = r_\ell$ where $r_1 > \dots > r_m > 0$ and $r_\ell - r_{\ell+1}$ are large enough (where $r_{m+1} := 0$), then (3.22) is satisfied. Thus, (3.20) and the sum in (3.19) converge absolutely. It is easy to see that the convergences are uniform. Hence we can exchange the sums and integrals, and therefore, the joint distribution is indeed given by (3.18) if we take the contours of z_i to be large nested circles.

We simplify (3.18). The terms $\mathcal{H}_{k_\ell, a_\ell}(W^{(\ell)}; W^{(\ell+1)})$ are evaluated in Proposition 3.4 below. Note that since the z_i -contours are the large nested circles, we have (3.22), and hence the assumptions in Proposition 3.4 are satisfied. On the other hand, the sum of $\mathcal{R}_X(W^{(m)})$ in (3.18) was computed in [4]. Lemma 6.1 in [4]

implies that for $W = (w_1, \dots, w_N) \in \mathcal{R}_z^N$,

$$(3.23) \quad \begin{aligned} \sum_{\substack{X \in \mathcal{X}_N(L) \\ x_k = a}} \mathcal{R}_X(W) &= (-1)^{(k-1)(N+1)} z^{(k-1)L} \\ &\times \left[1 - \prod_{j=1}^N (w_j + 1)^{-1} \right] \left[\prod_{j=1}^N w_j^{-k} (w_j + 1)^{-a+k+1} \right] \det [w_j^{-i}]_{i,j=1}^N. \end{aligned}$$

Hence, from the geometric series, for $W = (w_1, \dots, w_N) \in \mathcal{R}_z^N$,

$$(3.24) \quad \begin{aligned} \sum_{\substack{X \in \mathcal{X}_N(L) \\ x_k \geq a}} \mathcal{R}_X(W) \\ &= (-1)^{(k-1)(N+1)} z^{(k-1)L} \left[\prod_{j=1}^N w_j^{-k} (w_j + 1)^{-a+k+1} \right] \det [w_j^{-i}]_{i,j=1}^N \end{aligned}$$

if $\prod_{j=1}^N |w_j + 1| > 1$. The last condition is satisfied for $W = W^{(m)}$. We thus find that (3.19) is equal to an explicit factor times a product of $m-1$ Cauchy determinants times a Vandermonde determinant. By using the Cauchy-Binet identity m times, we obtain (3.6) assuming that the z_i -contours are large nested circles.

Finally, using the analyticity of the integrand on the right-hand side of (3.6), which was discussed before the start of this proof, we can deform the contours of z_i to any nested circles, not necessarily large circles. This completes the proof. \square

The main technical part of this section is the following summation formula. We prove it in Section 5.

Proposition 3.4. *Let z and z' be two non-zero complex numbers satisfying $z^L \neq (z')^L$. Let $W = (w_1, \dots, w_N) \in (\mathcal{R}_z)^N$ and $W' = (w'_1, \dots, w'_N) \in (\mathcal{R}_{z'})^N$. Suppose that $\prod_{j=1}^N |w'_j + 1| < \prod_{j=1}^N |w_j + 1|$. Consider $\mathcal{H}_{k,a}(W; W')$ defined in (3.20). Then for any $1 \leq k \leq N$ and integer a ,*

$$(3.25) \quad \begin{aligned} \mathcal{H}_{k,a}(W; W') \\ &= \left(\frac{z}{z'} \right)^{(k-1)L} \left(1 - \left(\frac{z'}{z} \right)^L \right)^{N-1} \left[\prod_{j=1}^N \frac{w_j^{-k} (w_j + 1)^{-a+k+1}}{(w'_j)^{-k} (w'_j + 1)^{-a+k}} \right] \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i,i'=1}^N. \end{aligned}$$

4. PERIODIC STEP INITIAL CONDITION

We now assume the following periodic step condition:

$$(4.1) \quad \mathbf{x}_{i+nN}(0) = i - N + nL \text{ for } 1 \leq i \leq N \text{ and } n \in \mathbb{Z}.$$

In the previous section, we obtained a formula for general initial conditions. In this section, we find a simpler formula for the periodic step initial condition which is suitable for the asymptotic analysis. We express $\mathcal{D}_Y(\mathbf{z}, \mathbf{k}, \mathbf{a}, \mathbf{t})$ as a Fredholm determinant times a simple factor. The result is described in terms of two functions $C(\mathbf{z})$ and $D(\mathbf{z})$. We first define them and then state the result.

Throughout this section, we fix a positive integer m , and fix parameters $k_1, \dots, k_m, a_1, \dots, a_m, t_1, \dots, t_m$ as in the previous section.

4.1. Definitions. Recall the function $q_z(w) = w^N(w+1)^{L-N} - z^L$ for complex z in (3.3) and the set of its roots

$$(4.2) \quad \mathcal{R}_z = \{w \in \mathbb{C} : q_z(w) = 0\}$$

in (3.4). Set

$$(4.3) \quad \mathbb{r}_0 := \rho^\rho(1-\rho)^{1-\rho}, \quad \rho = N/L,$$

as in (3.5). We discussed in the previous section that if $0 < |z| < \mathbb{r}_0$, then the contour $|q_z(w)| = 0$ consists of two closed contours, one in $\Re(w) < -\rho$ enclosing the point $w = -1$ and the other in $\Re(w) > -\rho$ enclosing the point $w = 0$. Now, for $0 < |z| < \mathbb{r}_0$, set

$$(4.4) \quad L_z = \{w \in \mathcal{R}_z : \Re(w) < -\rho\}, \quad R_z = \{w \in \mathcal{R}_z : \Re(w) > -\rho\}.$$

It is not difficult to check that

$$(4.5) \quad |L_z| = L - N, \quad |R_z| = N.$$

See the left picture in Figure 6 in Section 3. (Note that if $z = 0$, then the roots are $w = -1$ with multiplicity $L - N$ and $w = 0$ with multiplicity N .) From the definitions, we have

$$(4.6) \quad \mathcal{R}_z = L_z \cup R_z.$$

In Theorem 3.1, we took the contours of z_i as nested circles of arbitrary sizes. In this section, we assume that the circles satisfy

$$(4.7) \quad 0 < |z_m| < \cdots < |z_1| < \mathbb{r}_0.$$

Hence L_{z_i} and R_{z_i} are all well-defined.

We define two functions $C(\mathbf{z})$ and $D(\mathbf{z})$ of $\mathbf{z} = (z_1, \dots, z_m)$, both of which depend on the parameters k_i, t_i, a_i . The first one is the following. Recall the notational convention introduced in Definition 2.9. For example, $\Delta(R_z; L_z) = \prod_{v \in R_z} \prod_{u \in L_z} (v - u)$.

Definition 4.1. Define

$$(4.8) \quad \begin{aligned} C(\mathbf{z}) &= \left[\prod_{\ell=1}^m \frac{E_\ell(z_\ell)}{E_{\ell-1}(z_\ell)} \right] \left[\prod_{\ell=1}^m \frac{\prod_{u \in L_{z_\ell}} (-u)^N \prod_{v \in R_{z_\ell}} (v+1)^{L-N}}{\Delta(R_{z_\ell}; L_{z_\ell})} \right] \\ &\quad \times \left[\prod_{\ell=2}^m \frac{z_{\ell-1}^L}{z_{\ell-1}^L - z_\ell^L} \right] \left[\prod_{\ell=2}^m \frac{\Delta(R_{z_\ell}; L_{z_{\ell-1}})}{\prod_{u \in L_{z_{\ell-1}}} (-u)^N \prod_{v \in R_{z_\ell}} (v+1)^{L-N}} \right], \end{aligned}$$

where

$$(4.9) \quad E_i(z) := \prod_{u \in L_z} (-u)^{k_i - N - 1} \prod_{v \in R_z} (v+1)^{-a_i + k_i - N} e^{t_i v}$$

for $i = 1, \dots, m$, and $E_0(z) := 1$.

It is easy to see that all terms in $C(\mathbf{z})$ other than $\prod_{\ell=2}^m \frac{z_{\ell-1}^L}{z_{\ell-1}^L - z_\ell^L}$ are analytic for z_1, \dots, z_m within the disk $\{z; |z| < \mathbb{r}_0\}$. Hence $C(\mathbf{z})$ is analytic in the disk except for the simple poles when $z_{\ell-1}^L = z_\ell^L$, $\ell = 2, \dots, m$.

We now define $D(\mathbf{z})$. It is given by a Fredholm determinant. Set

$$(4.10) \quad \begin{aligned} F_i(w) &:= w^{-k_i + N + 1} (w+1)^{-a_i + k_i - N} e^{t_i w} \quad \text{for } i = 1, \dots, m, \\ F_0(w) &:= 1. \end{aligned}$$




FIGURE 7. Example of S_1 and S_2 when $m = 3$. The black dots are S_1 and the white dots are S_2 . The level sets are shown for visual convenience.

Define

$$(4.11) \quad f_i(w) = \begin{cases} \frac{F_i(w)}{F_{i-1}(w)} & \text{for } \Re(w) < -\rho, \\ \frac{F_{i-1}(w)}{F_i(w)} & \text{for } \Re(w) > -\rho. \end{cases}$$

Also, set

$$(4.12) \quad J(w) = \frac{w(w+1)}{L(w+\rho)}.$$

Define, for $0 < |z| < r_0$,

$$(4.13) \quad l_z(w) = \frac{1}{(w+1)^{L-N}} \prod_{u \in L_z} (w-u), \quad r_z(w) = \frac{1}{w^N} \prod_{u \in R_z} (w-u).$$

Note that $l_z(w)r_z(w) = \frac{q_z(w)}{(w+1)^{L-N}w^N} = \frac{w^N(w+1)^{L-N}-z^L}{w^N(w+1)^{L-N}}$. Set

$$(4.14) \quad H_z(w) := \begin{cases} l_z(w) & \text{for } \Re(w) > -\rho, \\ r_z(w) & \text{for } \Re(w) < -\rho. \end{cases}$$

When $z = 0$, we define $l_z(w) = r_z(w) = 1$ and hence $H_z(w) = 1$.

Define two sets

$$(4.15) \quad S_1 := L_{z_1} \cup R_{z_2} \cup L_{z_3} \cup \dots \cup \begin{cases} L_{z_m} & \text{if } m \text{ is odd,} \\ R_{z_m} & \text{if } m \text{ is even,} \end{cases}$$

and

$$(4.16) \quad S_2 := R_{z_1} \cup L_{z_2} \cup R_{z_3} \cup \dots \cup \begin{cases} R_{z_m} & \text{if } m \text{ is odd,} \\ L_{z_m} & \text{if } m \text{ is even.} \end{cases}$$

See Figure 7. We define two operators

$$(4.17) \quad K_1 : \ell^2(S_2) \rightarrow \ell^2(S_1), \quad K_2 : \ell^2(S_1) \rightarrow \ell^2(S_2)$$

by kernels. If $w \in \mathcal{R}_{z_i} \cap S_1$ and $w' \in \mathcal{R}_{z_j} \cap S_2$ for some $i, j \in \{1, \dots, m\}$, we set

$$(4.18) \quad K_1(w, w') = (\delta_i(j) + \delta_i(j + (-1)^i)) \frac{J(w) f_i(w) (H_{z_i}(w))^2}{H_{z_{i-(-1)^i}}(w) H_{z_{j-(-1)^j}}(w') (w - w')} Q_1(j).$$

Similarly, if $w \in \mathcal{R}_{z_i} \cap S_2$ and $w' \in \mathcal{R}_{z_j} \cap S_1$ for some $i, j \in \{1, \dots, m\}$, we set

$$(4.19) \quad K_2(w, w') = (\delta_i(j) + \delta_i(j - (-1)^i)) \frac{J(w) f_i(w) (H_{z_i}(w))^2}{H_{z_{i+(-1)^i}}(w) H_{z_{j+(-1)^j}}(w') (w - w')} Q_2(j).$$

Here we set $z_0 = z_{m+1} = 0$. We also set

$$(4.20) \quad Q_1(j) := 1 - \left(\frac{z_{j-(-1)^j}}{z_j} \right)^L, \quad Q_2(j) := 1 - \left(\frac{z_{j+(-1)^j}}{z_j} \right)^L.$$

Definition 4.2. Define

$$(4.21) \quad D(\mathbf{z}) = \det(I - K_1 K_2).$$

Remark 4.3. The matrix kernels for K_1 and K_2 have block structures similar to the infinite-time case discussed in Subsection 2.3. The only change is that \mathbf{k} is replaced by \mathbf{k} which is given as follows. For

$$(4.22) \quad u \in L_{z_i}, \quad v \in R_{z_i}, \quad u' \in L_{z_{i+1}}, \quad v' \in R_{z_{i+1}},$$

we have

$$(4.23) \quad \begin{bmatrix} k(u, v) & k(u, u') \\ k(v', v) & k(v', u') \end{bmatrix} = \begin{bmatrix} \frac{F_i(u)}{F_{i-1}(u)} & \frac{u(u+1)r_{z_i}(u)^2}{L(u+\rho)r_{z_{i+1}}(u)} \\ \frac{F_i(v')}{F_{i+1}(v')} & \frac{v'(v'+1)l_{z_{i+1}}(v')^2}{L(v'+\rho)l_{z_i}(v')} \end{bmatrix} \times \begin{bmatrix} \frac{1}{u-v} & \frac{1}{u-u'} \\ \frac{1}{v'-v} & \frac{1}{v'-u'} \end{bmatrix} \begin{bmatrix} \frac{1}{l_{z_{i+1}}(v)} & \frac{1}{r_{z_i}(u')} \\ \frac{1}{r_{z_i}(u')} & \frac{1}{r_{z_{i+1}}(v')} \end{bmatrix} \begin{bmatrix} 1 - \frac{z_{i+1}^L}{z_i^L} & \\ & 1 - \frac{z_i^L}{z_{i+1}^L} \end{bmatrix}.$$

As in Subsection 2.4, the above Fredholm determinant also has two series formulas. The proof of the following lemma is given in Subsection 4.3.

Lemma 4.4 (Series formulas of $D(\mathbf{z})$). *We have*

$$(4.24) \quad D(\mathbf{z}) = \sum_{\mathbf{n} \in (\mathbb{Z}_{\geq 0})^m} \frac{1}{(\mathbf{n}!)^2} D_{\mathbf{n}}(\mathbf{z})$$

with $\mathbf{n}! = \prod_{\ell=1}^m n_{\ell}!$ for $\mathbf{n} = (n_1, \dots, n_m)$ and $D_{\mathbf{n}}(\mathbf{z})$ can be expressed in the following two ways. Here we set $D_{\mathbf{n}}(\mathbf{z}) = 0$ if one of n_{ℓ} is larger than N .

(a) We have

$$(4.25) \quad D_{\mathbf{n}}(z_1, \dots, z_m) = (-1)^{|\mathbf{n}|} \sum_{\substack{U^{(\ell)} \in (L_{z_\ell})^{n_\ell} \\ V^{(\ell)} \in (R_{z_\ell})^{n_\ell} \\ l=1, \dots, m}} \det [K_1(w_i, w'_j)]_{i,j=1}^{|\mathbf{n}|} \det [K_2(w'_i, w_j)]_{i,j=1}^{|\mathbf{n}|},$$

where $U = (U^{(1)}, \dots, U^{(m)})$, $V = (V^{(1)}, \dots, V^{(m)})$ with $U^{(\ell)} = (u_1^{(\ell)}, \dots, u_{n_\ell}^{(\ell)})$, $V^{(\ell)} = (v_1^{(\ell)}, \dots, v_{n_\ell}^{(\ell)})$, and where

$$(4.26) \quad w_i = \begin{cases} u_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with odd integer } \ell, \\ v_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with even integer } \ell, \end{cases}$$

and

$$(4.27) \quad w'_i = \begin{cases} v_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with odd integer } \ell, \\ u_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with even integer } \ell. \end{cases}$$

(b) We have

$$(4.28) \quad D_{\mathbf{n}}(\mathbf{z}) = \sum_{\substack{U^{(\ell)} \in (L_{z_\ell})^{n_\ell} \\ V^{(\ell)} \in (R_{z_\ell})^{n_\ell} \\ \ell=1, \dots, m}} \left[\prod_{\ell=1}^m \frac{(\Delta(U^{(\ell)}))^2 (\Delta(V^{(\ell)}))^2}{(\Delta(U^{(\ell)}; V^{(\ell)}))^2} \hat{f}_\ell(U^{(\ell)}) \hat{f}_\ell(V^{(\ell)}) \right] \times \left[\prod_{\ell=2}^m \frac{\Delta(U^{(\ell)}; V^{(\ell-1)}) \Delta(V^{(\ell)}; U^{(\ell-1)}) (1 - \frac{z_\ell^L}{z_{\ell-1}^L})^{n_{\ell-1}} (1 - \frac{z_{\ell-1}^L}{z_\ell^L})^{n_\ell}}{\Delta(U^{(\ell)}; U^{(\ell-1)}) \Delta(V^{(\ell)}; V^{(\ell-1)}) r_{z_{\ell-1}}(U^{(\ell)}) r_{z_\ell}(U^{(\ell-1)}) l_{z_{\ell-1}}(V^{(\ell)}) l_{z_\ell}(V^{(\ell-1)})} \right],$$

where

$$(4.29) \quad \hat{f}_\ell(w) := J(w) f_\ell(w) (H_{z_\ell}(w))^2 = \begin{cases} J(w) \frac{F_\ell(w)}{F_{\ell-1}(w)} (r_{z_\ell}(w))^2 & \text{for } w \in L_{z_\ell}, \\ J(w) \frac{F_{\ell-1}(w)}{F_\ell(w)} (l_{z_\ell}(w))^2 & \text{for } w \in R_{z_\ell}. \end{cases}$$

Remark 4.5. From (4.28), we can check that $D_{\mathbf{n}}(\mathbf{z})$ is analytic for each z_ℓ in $0 < |z_\ell| < \mathfrak{r}_0$, $1 \leq \ell \leq m$, just like $D(\mathbf{z})$ of Section 2. The proof for $D(\mathbf{z})$ is in Lemma 7.1. The proof for $D_{\mathbf{n}}(\mathbf{z})$ is similar, and we skip it.

4.2. Result and proof.

Theorem 4.6 (Joint distribution of TASEP in $\mathcal{X}_N(L)$ for the step initial condition). *Consider the TASEP in $\mathcal{X}_N(L)$ with the step initial condition $\mathbf{x}_i(0) = i - N$, $1 \leq i \leq N$. Set $\rho = N/L$. Fix a positive integer m . Let $(k_1, t_1), \dots, (k_m, t_m)$ be m distinct points in $\{1, \dots, N\} \times [0, \infty)$. Assume that $0 \leq t_1 \leq \dots \leq t_m$. Let $a_i \in \mathbb{Z}$ for $1 \leq i \leq m$. Then*

$$(4.30) \quad \mathbb{P}(\mathbf{x}_{k_1}(t_1) \geq a_1, \dots, \mathbf{x}_{k_m}(t_m) \geq a_m) = \oint \dots \oint C(\mathbf{z}) D(\mathbf{z}) \frac{dz_m}{2\pi i z_m} \dots \frac{dz_1}{2\pi i z_1},$$

where $\mathbf{z} = (z_1, \dots, z_m)$ and the contours are nested circles satisfying

$$(4.31) \quad 0 < |z_m| < \dots < |z_1| < \mathfrak{r}_0$$

with $\mathfrak{r}_0 = \rho^\rho(1 - \rho)^{1-\rho}$. The functions $C(\mathbf{z})$ and $D(\mathbf{z})$ are defined in (4.8) and (4.21), respectively.

Recall that $C(\mathbf{z})$ is analytic in $|z_\ell| < \mathfrak{r}_0$ except for the poles when $z_{\ell-1}^L = z_\ell^L$, and $D(\mathbf{z})$ is analytic in $0 < |z_\ell| < \mathfrak{r}_0$. We point out that Remark 3.2 still applies to the above theorem; the Fredholm determinant expansion involves only finite sums.

Corollary 4.7 (Joint distribution of periodic TASEP for the periodic step initial condition). *Consider the periodic TASEP with the periodic step initial condition $\mathbf{x}_{i+nN}(0) = i - N + nL$ for $1 \leq i \leq N$ and $n \in \mathbb{Z}$. Then (4.30) holds for all integer indices k_1, \dots, k_m without the restriction that they are between 1 and N .*

Proof. As in the proof of Corollary 3.3, it is enough to show that the formulas are invariant under the changes $k_i \mapsto k_i \pm N$ and $a_i \mapsto a_i \pm L$ for some i . This can be checked easily for $C(\mathbf{z})$ using the identity $\prod_{u \in \mathbf{L}_z} (-u)^N = \prod_{v \in \mathbf{R}_z} (v+1)^{L-N}$, which is easy to prove; see (4.52) below. For $D(\mathbf{z})$, we use the fact that $u^N(u+1)^{L-N} = z_\ell^L$ for $u \in \mathbf{L}_{z_\ell}$ and $v_i^N(v_i+1)^{L-N} = z_i^L$ for $v \in \mathbf{R}_{z_\ell}$ (plus the special structure of K_1 and K_2). \square

Proof of Theorem 4.6. When $m = 1$, the result was obtained in Theorem 7.4 of [4]. We assume $m \geq 2$. In Theorem 3.1, $\mathcal{C}(\mathbf{z}, \mathbf{k})$ does not depend on the initial condition. Let us denote $\mathcal{D}_Y(\mathbf{z}, \mathbf{k}, \mathbf{a}, \mathbf{t})$ by $\mathcal{D}_{\text{step}}$ when $Y = (1 - N, \dots, 1, 0)$, the step initial condition. We need to show that $\mathcal{D}_{\text{step}} = D(\mathbf{z}) \frac{C(\mathbf{z})}{\mathcal{C}(\mathbf{z}, \mathbf{k})}$.

Inserting the initial condition $y_i = i - N$, rewriting \mathcal{G}_ℓ in terms of F_ℓ and \mathbf{J} in (4.10) and (4.12), and reversing the rows, we get

$$(4.32) \quad \mathcal{D}_{\text{step}} = (-1)^{N(N-1)/2} \det \left[\sum \frac{w_1^{-i} w_m^{-j}}{\prod_{\ell=2}^m (w_\ell - w_{\ell-1})} \prod_{\ell=1}^m \mathbf{J}(w_\ell) \frac{F_\ell(w_\ell)}{F_{\ell-1}(w_\ell)} \right]_{i,j=1}^N.$$

The sum is over all $w_1 \in \mathcal{R}_{z_1}, \dots, w_m \in \mathcal{R}_{z_m}$. Using the Cauchy-Binet identity m times,

$$(4.33) \quad \mathcal{D}_{\text{step}} = \frac{(-1)^{N(N-1)/2}}{(N!)^m} \sum_{\substack{W^{(\ell)} \in (\mathcal{R}_{z_\ell})^N \\ \ell=1, \dots, m}} \mathcal{E}(\mathbf{W}) \prod_{\ell=1}^m \mathbf{J}(W^{(\ell)}) \frac{F_\ell(W^{(\ell)})}{F_{\ell-1}(W^{(\ell)})},$$

where $W^{(\ell)} = (w_1^{(\ell)}, \dots, w_N^{(\ell)})$ with $w_i^{(\ell)} \in \mathcal{R}_{z_\ell}$ for each i , $\mathbf{W} = (W^{(1)}, \dots, W^{(m)})$, and

$$(4.34) \quad \mathcal{E}(\mathbf{W}) = \det \left[(w_i^{(1)})^{-j} \right] \det \left[\frac{1}{w_i^{(2)} - w_j^{(1)}} \right] \dots \det \left[\frac{1}{w_i^{(m)} - w_j^{(m-1)}} \right] \det \left[(w_i^{(m)})^{-j} \right].$$

Here all matrices are indexed by $1 \leq i, j \leq N$. Note that in (4.33), we use the notational convention such as $F_\ell(W^{(\ell)}) = \prod_{i=1}^N F_\ell(w_i^{(\ell)})$ mentioned in Definition 2.9. Evaluating the Vandermonde determinants and the Cauchy determinants, $\mathcal{D}_{\text{step}}$ is

equal to (recall the notation (2.48))

$$(4.35) \quad \frac{(-1)^{mN(N-1)/2}}{(N!)^m} \sum_{\substack{W^{(\ell)} \in (\mathcal{R}_{z_\ell})^N \\ \ell=1, \dots, m}} \frac{\prod_{\ell=1}^m \Delta(W^{(\ell)})^2}{\prod_{\ell=2}^m \Delta(W^{(\ell)}; W^{(\ell-1)})} \\ \times \left[\prod_{i=1}^N (w_i^{(1)} w_i^{(m)})^{-N} \right] \prod_{\ell=1}^m J(W^{(\ell)}) \frac{F_\ell(W^{(\ell)})}{F_{\ell-1}(W^{(\ell)})}.$$

Note that for each ℓ , we may assume that the coordinates of the vector $W^{(\ell)}$ are all distinct since otherwise the summand is zero due to $\Delta(W^{(\ell)})$. Also note that the summand is a symmetric function of the coordinates of $W^{(\ell)}$ for each ℓ . Hence instead of taking the sum over the vectors $W^{(\ell)} \in (\mathcal{R}_{z_\ell})^N$, we can take a sum over the subsets $\tilde{W}^{(\ell)} \subset \mathcal{R}_{z_\ell}$ of size N : $\mathcal{D}_{\text{step}}$ is equal to

$$(4.36) \quad (-1)^{mN(N-1)/2} \\ \times \sum_{\substack{\tilde{W}^{(\ell)} \subset \mathcal{R}_{z_\ell} \\ |\tilde{W}^{(\ell)}|=N \\ \ell=1, \dots, m}} \frac{\prod_{\ell=1}^m \Delta(\tilde{W}^{(\ell)})^2}{\prod_{\ell=2}^m \Delta(\tilde{W}^{(\ell)}; \tilde{W}^{(\ell-1)})} \left[\prod_{i=1}^N (w_i^{(1)} w_i^{(m)})^{-N} \right] \prod_{\ell=1}^m J(\tilde{W}^{(\ell)}) \frac{F_\ell(\tilde{W}^{(\ell)})}{F_{\ell-1}(\tilde{W}^{(\ell)})},$$

where $w_i^{(1)}$ are the elements of $\tilde{W}^{(1)}$ and $w_i^{(m)}$ are the elements of $\tilde{W}^{(m)}$.

We now change the sum as follows. Since \mathcal{R}_{z_ℓ} is the disjoint union of L_{z_ℓ} and R_{z_ℓ} , some elements of the set $\tilde{W}^{(\ell)}$ are in L_{z_ℓ} and the rest are in R_{z_ℓ} . (Recall that $|L_{z_\ell}| = L - N$ and $|R_{z_\ell}| = N$.) Let $\tilde{U}^{(\ell)} = \tilde{W}^{(\ell)} \cap L_{z_\ell}$ and $\tilde{V}^{(\ell)} = R_{z_\ell} \setminus \tilde{W}^{(\ell)}$. Observe that since $|\tilde{W}^{(\ell)}| = |R_{z_\ell}| (= N)$, we have $|\tilde{U}^{(\ell)}| = |\tilde{V}^{(\ell)}|$. Call this last number n_ℓ . We thus find that the sum in (4.36) can be replaced by the sums

$$(4.37) \quad \sum_{\substack{n_\ell=0, \dots, N \\ \ell=1, \dots, m}} \sum_{\substack{\tilde{U}^{(\ell)} \subset L_{z_\ell}, \tilde{V}^{(\ell)} \subset R_{z_\ell} \\ |\tilde{U}^{(\ell)}|=|\tilde{V}^{(\ell)}|=n_\ell \\ \ell=1, \dots, m}} \dots$$

We now express the summand in terms of $\tilde{U}^{(\ell)}$ and $\tilde{V}^{(\ell)}$ instead of $\tilde{W}^{(\ell)}$. First, for any function h ,

$$(4.38) \quad h(\tilde{W}^{(\ell)}) = h(\tilde{U}^{(\ell)}) \frac{h(R_{z_\ell})}{h(\tilde{V}^{(\ell)})}.$$

Now consider $\Delta(\tilde{W}^{(\ell)})^2$. We suppress the dependence on ℓ in the next few sentences to make the notation light. Tentatively set $\tilde{S} = R_z \setminus \tilde{V}$ so that $R_z = \tilde{V} \cup \tilde{S}$. Note that $\tilde{W} = \tilde{U} \cup \tilde{S}$, a disjoint union. We thus have

$$(4.39) \quad \Delta(\tilde{W})^2 = \Delta(\tilde{U})^2 \Delta(\tilde{S})^2 \Delta(\tilde{U}; \tilde{S})^2, \quad \Delta(R_z)^2 = \Delta(\tilde{V})^2 \Delta(\tilde{S})^2 \Delta(\tilde{V}; \tilde{S})^2.$$

Let

$$(4.40) \quad q_{z,R}(w) := \prod_{v \in R_z} (w - v) = w^N r_z(w).$$

Then,

$$(4.41) \quad q_{z,R}(\tilde{U}) = \Delta(\tilde{U}; \tilde{V}) \Delta(\tilde{U}; \tilde{S}), \quad q'_{z,R}(\tilde{V}) = (-1)^{N(N-1)/2} \Delta(\tilde{V})^2 \Delta(\tilde{V}; \tilde{S}).$$

It is also direct to see that

$$(4.42) \quad \Delta(R_z)^2 = (-1)^{N(N-1)/2} q'_{z,R}(R_z).$$

From these, after canceling out all terms involving \tilde{S} and inserting the dependence on ℓ , we find that

$$(4.43) \quad \Delta(\tilde{W}^{(\ell)})^2 = (-1)^{N(N-1)/2} \frac{\Delta(\tilde{U}^{(\ell)})^2 \Delta(\tilde{V}^{(\ell)})^2 (q'_{z_\ell,R}(\tilde{U}^{(\ell)}))^2}{\Delta(\tilde{U}^{(\ell)}; \tilde{V}^{(\ell)})^2 (q'_{z_\ell,R}(\tilde{V}^{(\ell)}))^2} q'_{z_\ell,R}(R_{z_\ell}).$$

(This computation was also given in (7.48) and (7.50) of [4].) Similarly,

$$(4.44) \quad \frac{\Delta(\tilde{W}^{(\ell)}; \tilde{W}^{(\ell-1)})}{\Delta(R_{z_\ell}; R_{z_{\ell-1}})} = \frac{\Delta(\tilde{U}^{(\ell)}; \tilde{U}^{(\ell-1)}) \Delta(\tilde{V}^{(\ell)}; \tilde{V}^{(\ell-1)}) q'_{z_{\ell-1},R}(\tilde{U}^{(\ell)}) q'_{z_\ell,R}(\tilde{U}^{(\ell-1)})}{\Delta(\tilde{U}^{(\ell)}; \tilde{V}^{(\ell-1)}) \Delta(\tilde{V}^{(\ell)}; \tilde{U}^{(\ell-1)}) q'_{z_{\ell-1},R}(\tilde{V}^{(\ell)}) q'_{z_\ell,R}(\tilde{V}^{(\ell-1)})}.$$

We express the summands in (4.36) using (4.38), (4.43), and (4.44). We then change the subsets $\tilde{U}^{(\ell)} \subset L_{z_\ell}$ and $\tilde{V}^{(\ell)} \subset R_{z_\ell}$ to vectors $U^{(\ell)} \in (L_{z_\ell})^{n_\ell}$ and $V^{(\ell)} \in (R_{z_\ell})^{n_\ell}$. This has the effect of introducing the factors $\frac{1}{(n_\ell!)^2}$. We thus obtain

$$(4.45) \quad \mathcal{D}_{\text{step}} = \mathcal{B}(\mathbf{z}) \left[\sum_{\mathbf{n} \in (\mathbb{Z}_{\geq 0})^m} \frac{1}{(\mathbf{n}!)^2} \hat{D}_{\mathbf{n}}(\mathbf{z}) \right]$$

with

$$(4.46) \quad \mathcal{B}(\mathbf{z}) = \left[\frac{\prod_{\ell=1}^m q'_{z_\ell,R}(R_{z_\ell})}{\prod_{\ell=2}^m \Delta(R_{z_\ell}; R_{z_{\ell-1}})} \right] \left[\prod_{v \in R_{z_1}} v^{-N} \right] \left[\prod_{v \in R_{z_m}} v^{-N} \right] \left[\prod_{\ell=1}^m J(R_{z_\ell}) \frac{F_\ell(R_{z_\ell})}{F_{\ell-1}(R_{z_\ell})} \right]$$

and

$$(4.47) \quad \begin{aligned} \hat{D}_{\mathbf{n}}(\mathbf{z}) &= \sum_{\substack{U^{(\ell)} \in (L_{z_\ell})^{n_\ell} \\ V^{(\ell)} \in (R_{z_\ell})^{n_\ell} \\ \ell=1, \dots, m}} \left[\prod_{\ell=1}^m \frac{(\Delta(U^{(\ell)}))^2 (\Delta(V^{(\ell)}))^2 (q'_{z_\ell,R}(U^{(\ell)}))^2 J(U^{(\ell)}) F_\ell(U^{(\ell)}) F_{\ell-1}(V^{(\ell)})}{(\Delta(U^{(\ell)}; V^{(\ell)}))^2 (q'_{z_\ell,R}(V^{(\ell)}))^2 J(V^{(\ell)}) F_\ell(V^{(\ell)}) F_{\ell-1}(U^{(\ell)})} \right] \\ &\quad \times \left[\prod_{i=1}^{n_1} \left(\frac{u_i^{(1)}}{v_i^{(1)}} \right)^{-N} \right] \left[\prod_{i=1}^{n_m} \left(\frac{u_i^{(m)}}{v_i^{(m)}} \right)^{-N} \right] \\ &\quad \times \left[\prod_{\ell=2}^m \frac{\Delta(U^{(\ell)}; V^{(\ell-1)}) \Delta(V^{(\ell)}; U^{(\ell-1)}) q'_{z_{\ell-1},R}(V^{(\ell)}) q'_{z_\ell,R}(V^{(\ell-1)})}{\Delta(U^{(\ell)}; U^{(\ell-1)}) \Delta(V^{(\ell)}; V^{(\ell-1)}) q'_{z_{\ell-1},R}(U^{(\ell)}) q'_{z_\ell,R}(U^{(\ell-1)})} \right]. \end{aligned}$$

We rewrite $(q'_{z_\ell,R}(V^{(\ell)}))^2$ using the identity, $q'_{z,R}(v) = \frac{v^N}{J(v)l_z(v)}$, which we prove later in (4.51). We also use $q_{z,R}(w) = w^N r_z(w)$ (see (4.40)). Furthermore, we re-express $r_{z_\ell}(V^{(\ell)})$ in terms of $l_z(V^{(\ell)})$ using the identity

$$(4.48) \quad r_z(v) = \frac{1}{l_z(v)} \left(1 - \frac{z^L}{(z')^L} \right) \quad \text{for } v \in R_{z'}$$

which follows from the fact that $r_z(w)l_z(w) = 1 - \frac{z^L}{w^N(w+1)^{L-N}}$. Then, using the notation (4.29), we find that $\hat{D}_{\mathbf{n}}(\mathbf{z}) = D_{\mathbf{n}}(\mathbf{z})$, given by (4.28).

Thus, the theorem is proved if we show that $\mathcal{C}(\mathbf{z}, \mathbf{k})\mathcal{B}(\mathbf{z}) = C(\mathbf{z})$. Before we prove it, we make the following observations.

- For any $v' \in R_{z'}$, we have $0 = q_{z'}(v') = (v')^N(v' + 1)^{L-N} - (z')^L$. Hence, for another complex number z ,

$$(4.49) \quad z'^L - z^L = (v')^N(v' + 1)^{L-N} - z^L = q_z(v') = \prod_{u \in L_z} (v' - u) \prod_{v \in R_z} (v' - v).$$

- As a special case of the above identity, taking $z' = 0$ and $v' = 0$, we obtain

$$(4.50) \quad z^L = (-1)^{N-1} \prod_{u \in L_z} (-u) \prod_{v \in R_z} v.$$

- Setting $q_{z,L}(w) = \prod_{u \in L_z} (w - u)$, we have $q_{z,R}(w)q_{z,L}(w) = q_z(w) = w^N(w + 1)^{L-N} - z^L$, and hence,

$$(4.51) \quad q'_{z,R}(v) = \frac{L(v + \rho)}{v(v + 1)} \frac{v^N(v + 1)^{L-N}}{q_{z,L}(v)} = \frac{v^N}{J(v)l_z(v)} \quad \text{for } v \in R_z.$$

- Since $z^{NL} = \prod_{u \in L_z} (-u)^N \prod_{v \in R_z} v^N$ from equality (4.50) and $z^{NL} = \prod_{v \in R_z} v^N(v + 1)^{L-N}$ by using the definition of R_z , we find that

$$(4.52) \quad \prod_{u \in L_z} (-u)^N = \prod_{v \in R_z} (v + 1)^{L-N}.$$

We now prove that $\mathcal{C}(\mathbf{z}, \mathbf{k})\mathcal{B}(\mathbf{z}) = C(\mathbf{z})$. Consider $\mathcal{C}(\mathbf{z}, \mathbf{k})$ defined in (3.7). Using (4.49),

$$\begin{aligned} & (-1)^{(k_m-1)(N+1)} z_1^{(k_1-1)L} \prod_{\ell=2}^m z_\ell^{(k_\ell-k_{\ell-1})L} \\ &= \left[\prod_{u \in L_{z_1}} (-u)^{k_1-1} \prod_{v \in R_{z_1}} v^{(k_1-1)} \right] \cdot \prod_{\ell=2}^m \left[\prod_{u \in L_{z_\ell}} (-u)^{k_\ell-k_{\ell-1}} \prod_{v \in R_{z_\ell}} v^{(k_\ell-k_{\ell-1})} \right]. \end{aligned}$$

Using (4.50) and (4.49),

$$\begin{aligned} & \prod_{\ell=2}^m \left(\left(\frac{z_\ell}{z_{\ell-1}} \right)^L - 1 \right)^N \\ &= \prod_{\ell=2}^m \frac{\Delta(R_{z_\ell}; L_{z_{\ell-1}}) \Delta(R_{z_\ell}; R_{z_{\ell-1}})}{\prod_{u \in L_{z_{\ell-1}}} (-u)^N \prod_{v \in R_{z_{\ell-1}}} v^N} \\ &= \left[\prod_{\ell=2}^m \Delta(R_{z_\ell}; R_{z_{\ell-1}}) \right] \left[\prod_{\ell=2}^m \frac{\Delta(R_{z_\ell}; L_{z_{\ell-1}})}{\prod_{u \in L_{z_{\ell-1}}} (-u)^N \prod_{v \in R_{z_\ell}} (v + 1)^{L-N}} \right] \\ & \quad \times \left[\prod_{\ell=2}^m \prod_{v \in R_{z_\ell}} (v + 1)^{L-N} \right] \left[\prod_{\ell=1}^{m-1} \prod_{v \in R_{z_\ell}} v^{-N} \right]. \end{aligned}$$

Now consider $\mathcal{B}(\mathbf{z})$. Using (4.51), the fact that $\prod_{v \in R_z} q_{z,L}(v) = \Delta(R_z; L_z)$, and (4.52), we see that

$$\begin{aligned} q'_{z,R}(R_z)J(R_z) &= \frac{\prod_{v \in R_z} (v+1)^{L-N}}{\Delta(R_z; L_z)} \left[\prod_{v \in R_z} v^N \right] \\ &= \frac{\left[\prod_{v \in R_z} (v+1)^{L-N} \right] \left[\prod_{u \in L_z} (-u)^N \right]}{\Delta(R_z; L_z)} \left[\prod_{v \in R_z} v^N \right] \left[\prod_{v \in R_z} (v+1)^{-L+N} \right]. \end{aligned}$$

This implies that

$$\begin{aligned} \prod_{\ell=1}^m q'_{z_\ell,R}(R_{z_\ell})J(R_{z_\ell}) &= \prod_{\ell=1}^m \frac{\left[\prod_{v \in R_{z_\ell}} (v+1)^{L-N} \right] \left[\prod_{u \in L_{z_\ell}} (-u)^N \right]}{\Delta(R_{z_\ell}; L_{z_\ell})} \\ &\quad \left[\prod_{\ell=1}^m \prod_{v \in R_{z_\ell}} v^N \right] \left[\prod_{\ell=2}^m \prod_{v \in R_{z_\ell}} (v+1)^{-L+N} \right] \left[\prod_{u \in L_{z_1}} (-u)^{-N} \right]. \end{aligned}$$

From these calculations, we find that $\mathcal{C}(\mathbf{z}, \mathbf{k})\mathcal{B}(\mathbf{z}) = C(\mathbf{z})$. □

4.3. Equivalence of the Fredholm determinant and series formulas. We presented three formulas of $D(\mathbf{z})$ in Subsection 4.1; see Lemma 4.4. One of them is a Fredholm determinant (4.21) and the other two are series formulas (4.24) with (4.25) and (4.28). In this subsection, we prove the equivalence of these formulas. The proof is general, and the same argument also gives a proof of Lemma 2.10, which states the equivalence of three formulas of $D(\mathbf{z})$, a limit of $D(\mathbf{z})$.

First, we prove Lemma 4.4(a) and Lemma 2.10(i). These are special cases of the next general result which follows from the series definition of Fredholm determinant together with a block structure of the operator.

Lemma 4.8. *Let $\Sigma_1, \dots, \Sigma_m$ be disjoint sets in \mathbb{C} and let $\mathcal{H} = L^2(\Sigma_1 \cup \dots \cup \Sigma_m, \mu)$ for some measure μ . Let $\Sigma'_1, \dots, \Sigma'_m$ be another collection of disjoint sets in \mathbb{C} and let $\mathcal{H}' = L^2(\Sigma'_1 \cup \dots \cup \Sigma'_m, \mu')$ for some measure μ' . Let A be an operator from \mathcal{H}' to \mathcal{H} and B an operator from \mathcal{H} to \mathcal{H}' , both of which are defined by kernels. Suppose the following block structures:*

- $A(w, w') = 0$ unless there is an index i such that $w \in \Sigma_{2i-1} \cup \Sigma_{2i}$ and $w' \in \Sigma'_{2i-1} \cup \Sigma'_{2i}$,
- $B(w', w) = 0$ unless there is an index i such that $w' \in \Sigma'_{2i} \cup \Sigma'_{2i+1}$ and $w \in \Sigma_{2i} \cup \Sigma_{2i+1}$.

Assume that the Fredholm determinant $\det(1 - AB)$ is well-defined and is equal to the usual Fredholm determinant series expansion. Then

$$(4.53) \quad \det(1 - AB) = \sum_{\mathbf{n} \in (\mathbb{Z}_{\geq 0})^m} \frac{(-1)^{|\mathbf{n}|}}{(\mathbf{n}!)^2} \int_{\Sigma_1^{n_1} \times \cdots \times \Sigma_m^{n_m}} \int_{(\Sigma'_1)^{n_1} \times \cdots \times (\Sigma'_m)^{n_m}} \det [A(w_i, w'_j)]_{i,j=1}^{|\mathbf{n}|} \det [B(w'_i, w_j)]_{i,j=1}^{|\mathbf{n}|} \prod_{i=1}^{|\mathbf{n}|} d\mu'(w'_i) \prod_{i=1}^{|\mathbf{n}|} d\mu(w_i),$$

where $\mathbf{n} = (n_1, \dots, n_m)$.

Proof. From the standard Fredholm determinant series expansion,

$$\det(1 - AB) = \sum_{\mathbf{n} \in (\mathbb{Z}_{\geq 0})^m} \frac{(-1)^{|\mathbf{n}|}}{\mathbf{n}!} \int_{\Sigma_1^{n_1} \times \cdots \times \Sigma_m^{n_m}} \det [(AB)(w_i, w_j)]_{i,j=1}^{|\mathbf{n}|} \prod_{i=1}^{|\mathbf{n}|} d\mu(w_i),$$

where among $|\mathbf{n}|$ variables w_i , the first n_1 are in Σ_1 , the next n_2 are in Σ_2 , and so on. By the Cauchy-Binet formula, for given \mathbf{n} ,

$$\begin{aligned} & \det [(AB)(w_i, w_j)]_{i,j=1}^{|\mathbf{n}|} \\ &= \frac{1}{|\mathbf{n}|!} \int_{(\Sigma'_1 \cup \cdots \cup \Sigma'_m)^{|\mathbf{n}|}} \det [A(w_i, w'_j)]_{i,j=1}^{|\mathbf{n}|} \det [B(w'_i, w_j)]_{i,j=1}^{|\mathbf{n}|} \prod_{i=1}^{|\mathbf{n}|} d\mu'(w'_i) \\ &= \sum_{\substack{\mathbf{n}' \in (\mathbb{Z}_{\geq 0})^m \\ |\mathbf{n}'| = |\mathbf{n}|}} \frac{1}{(\mathbf{n}')!} \int_{(\Sigma'_1)^{n'_1} \times \cdots \times (\Sigma'_m)^{n'_m}} \det [A(w_i, w'_j)]_{i,j=1}^{|\mathbf{n}|} \\ & \quad \times \det [B(w'_i, w_j)]_{i,j=1}^{|\mathbf{n}|} \prod_{i=1}^{|\mathbf{n}|} d\mu'(w'_i). \end{aligned}$$

From the structure of the kernels, we find that the matrix $A(w_i, w'_j)$ has a natural block structure of block sizes $n_1 + n_2, n_3 + n_4$, and so on for the rows and of sizes $n'_1 + n'_2, n'_3 + n'_4$, and so on for the columns. The matrix has non-zero entries only in the diagonal blocks, which have sizes $(n_1 + n_2) \times (n'_1 + n'_2), (n_3 + n_4) \times (n'_3 + n'_4), \dots$. Similarly, the matrix $B(w_i, w'_j)$ has non-zero entries only on the diagonal blocks, which have sizes $n_1 \times n'_1, (n_2 + n_3) \times (n'_2 + n'_3), (n_4 + n_5) \times (n'_4 + n'_5), \dots$. From these structures, we find that the determinant of $A(w_i, w'_j)$ times the determinant of $B(w_i, w'_j)$ is zero unless $\mathbf{n}' = \mathbf{n}$. Hence

$$\begin{aligned} & \det [(AB)(w_i, w_j)]_{i,j=1}^{|\mathbf{n}|} \\ &= \frac{1}{\mathbf{n}!} \int_{(\Sigma'_1)^{n_1} \times \cdots \times (\Sigma'_m)^{n_m}} \det [A(w_i, w'_j)]_{i,j=1}^{|\mathbf{n}|} \det [B(w'_i, w_j)]_{i,j=1}^{|\mathbf{n}|} \prod_{i=1}^{|\mathbf{n}|} d\mu'(w'_i). \end{aligned}$$

This implies (4.53). \square

Now, we prove Lemma 4.4(b) by showing that (4.25) and (4.28) are the same. We prove Lemma 2.10(ii) similarly. For this purpose, we use the following general result.

Lemma 4.9. *Let $U = (u_1, \dots, u_m)$, $V = (v_1, \dots, v_m)$, $U' = (u'_1, \dots, u'_n)$, and $V' = (v'_1, \dots, v'_n)$ be four complex vectors. Then for any single-variable functions \mathcal{F} , \mathcal{G} , \mathcal{I} , and \mathcal{J} ,*

$$(4.54) \quad \begin{aligned} & \frac{\Delta(U)\Delta(V)\Delta(U')\Delta(V')\Delta(U'; V)\Delta(V'; U)}{\Delta(U; V)\Delta(U'; V')\Delta(U'; U)\Delta(V'; V)} \mathcal{F}(U)\mathcal{G}(V)\mathcal{I}(U')\mathcal{J}(V') \\ & = (-1)^{\binom{n+m}{2}+n+mn} \det \begin{bmatrix} \left[\frac{\mathcal{F}(u_i)\mathcal{G}(v_j)}{u_i-v_j} \right]_{m \times m} & \left[\frac{\mathcal{F}(u_i)\mathcal{I}(u'_j)}{u_i-u'_j} \right]_{m \times n} \\ \left[\frac{\mathcal{J}(v'_i)\mathcal{G}(v_j)}{v'_i-v_j} \right]_{n \times m} & \left[\frac{\mathcal{J}(v'_i)\mathcal{I}(u'_j)}{v'_i-u'_j} \right]_{n \times n} \end{bmatrix}. \end{aligned}$$

Proof. It follows directly from the Cauchy determinant formula. \square

Consider (4.28). We can write it as

$$(4.55) \quad D_{\mathbf{n}}(\mathbf{z}) = \sum_{\substack{U^{(\ell)} \in (L_{z_\ell})^{n_\ell} \\ V^{(\ell)} \in (R_{z_\ell})^{n_\ell} \\ l=1, \dots, m}} \tilde{D}_0 \cdots \tilde{D}_m,$$

where

$$(4.56) \quad \tilde{D}_0 = \frac{\Delta(U^{(1)})\Delta(V^{(1)})}{\Delta(U^{(1)}; V^{(1)})} \hat{f}_1(V^{(1)}), \quad \tilde{D}_m = \frac{\Delta(U^{(m)})\Delta(V^{(m)})}{\Delta(U^{(m)}; V^{(m)})} \hat{f}_m(U^{(m)}),$$

and

$$(4.57) \quad \begin{aligned} \tilde{D}_\ell &= \frac{\Delta(U^{(\ell)})\Delta(V^{(\ell)})\Delta(U^{(\ell+1)})\Delta(V^{(\ell+1)})\Delta(U^{(\ell+1)}; V^{(\ell)})\Delta(V^{(\ell+1)}; U^{(\ell)})}{\Delta(U^{(\ell)}; V^{(\ell)})\Delta(U^{(\ell+1)}; V^{(\ell+1)})\Delta(U^{(\ell+1)}; U^{(\ell)})\Delta(V^{(\ell+1)}; V^{(\ell)})} \\ &\times \frac{\hat{f}_\ell(U^{(\ell)})(1 - \frac{z_{\ell+1}^L}{z_\ell^L})^{n_\ell}(1 - \frac{z_\ell^L}{z_{\ell+1}^L})^{n_{\ell+1}} \hat{f}_\ell(V^{(\ell+1)})}{r_{z_{\ell+1}}(U^{(\ell)})l_{z_{\ell+1}}(V^{(\ell)})r_{z_\ell}(U^{(\ell+1)})l_{z_\ell}(V^{(\ell+1)})} \end{aligned}$$

for $1 \leq \ell \leq m-1$. For $1 \leq \ell \leq m-1$, we apply Lemma 4.9 with $m = n_\ell$, $n = n_{\ell+1}$, $\mathcal{F}(u) = \frac{\hat{f}_\ell(u)}{r_{z_{\ell+1}}(u)}$, $\mathcal{G}(v) = \frac{1}{l_{z_{\ell+1}}(v)}(1 - \frac{z_{\ell+1}^L}{z_\ell^L})$, $\mathcal{I}(u') = \frac{1}{r_{z_\ell}(u')}(1 - \frac{z_\ell^L}{z_{\ell+1}^L})$, and $\mathcal{J}(v') = \frac{\hat{f}_{\ell+1}(v')}{l_{z_\ell}(v')}$. Then, recalling (4.29),

$$\begin{aligned} \tilde{D}_\ell &= (-1)^{\binom{n_\ell+n_{\ell+1}}{2}+n_{\ell+1}+n_\ell n_{\ell+1}} \\ &\times \det \begin{bmatrix} \left[K_1(u_i^{(\ell)}, v_j^{(\ell)}) \right]_{n_\ell \times n_\ell} & \left[K_1(u_i^{(\ell)}, u_j^{(\ell+1)}) \right]_{n_\ell \times n_{\ell+1}} \\ \left[K_1(v_i^{(\ell+1)}, v_j^{(\ell)}) \right]_{n_{\ell+1} \times n_\ell} & \left[K_1(v_i^{(\ell+1)}, u_j^{(\ell+1)}) \right]_{n_{\ell+1} \times n_{\ell+1}} \end{bmatrix} \end{aligned}$$

if ℓ is odd, and

$$\begin{aligned} \tilde{D}_\ell &= (-1)^{\binom{n_\ell+n_{\ell+1}}{2}+n_{\ell+1}+n_\ell n_{\ell+1}} \\ &\times \det \begin{bmatrix} \left[K_2(u_i^{(\ell)}, v_j^{(\ell)}) \right]_{n_\ell \times n_\ell} & \left[K_2(u_i^{(\ell)}, u_j^{(\ell+1)}) \right]_{n_\ell \times n_{\ell+1}} \\ \left[K_2(v_i^{(\ell+1)}, v_j^{(\ell)}) \right]_{n_{\ell+1} \times n_\ell} & \left[K_2(v_i^{(\ell+1)}, u_j^{(\ell+1)}) \right]_{n_{\ell+1} \times n_{\ell+1}} \end{bmatrix} \end{aligned}$$

if ℓ is even. On the other hand, using the Cauchy determinant formula,

$$(4.58) \quad \tilde{D}_0 = (-1)^{\binom{n_1}{2} + n_1} \det \left[K_2(v_i^{(1)}, u_j^{(1)}) \right]_{i,j=1}^{n_1}$$

and

$$(4.59) \quad \tilde{D}_m = \begin{cases} (-1)^{\binom{n_m}{2}} \det \left[K_1(u_i^{(m)}, v_j^{(m)}) \right]_{i,j=1}^{n_m} & \text{if } m \text{ is even,} \\ (-1)^{\binom{n_m}{2}} \det \left[K_2(u_i^{(m)}, v_j^{(m)}) \right]_{i,j=1}^{n_m} & \text{if } m \text{ is odd.} \end{cases}$$

The formula (4.25) follows by combining the product of \tilde{D}_ℓ for odd ℓ as a single determinant of a block diagonal matrix, and combining the product \tilde{D}_ℓ for even indices as another single determinant, we obtain (4.25). This proves Lemma 4.4(b). The proof of Lemma 2.10(ii) is similar.

5. PROOF OF PROPOSITION 3.4

As we mentioned before, Proposition 3.4 is the key technical result of this paper. We prove it in this section.

Let z and z' be two non-zero complex numbers satisfying $z^L \neq (z')^L$. Let $W = (w_1, \dots, w_N) \in (\mathcal{R}_z)^N$ and $W' = (w'_1, \dots, w'_N) \in (\mathcal{R}_{z'})^N$ be two complex vectors satisfying $\prod_{j=1}^N |w'_j + 1| < \prod_{j=1}^N |w_j + 1|$. Let

$$(5.1) \quad \mathcal{L}_Y(W) = \det \left[w_i^j (w_i + 1)^{y_j - j} \right]_{i,j=1}^N, \quad \mathcal{R}_X(W) = \det \left[w_i^{-j} (w_i + 1)^{-x_j + j} \right]_{i,j=1}^N.$$

The goal is to evaluate

$$(5.2) \quad \mathcal{H}_{k,a}(W; W') = \sum_{X \in \mathcal{X}_N(L) \cap \{x_k \geq a\}} \mathcal{R}_X(W) \mathcal{L}_X(W')$$

and show that it is equal to the right-hand side of (3.25).

We first reduce the general k case to the $k = 1$ case.

Lemma 5.1. *Under the same assumptions as in Proposition 3.4,*

$$(5.3) \quad \mathcal{H}_{k,a}(W; W') = \left(\frac{z}{z'} \right)^{(k-1)L} \left[\prod_{j=1}^N \left(\frac{(w_j + 1)w'_j}{w_j(w'_j + 1)} \right)^{k-1} \right] \mathcal{H}_{1,a}(W; W').$$

Proof. The sum in (5.2) is over the discrete variables $x_1 < \dots < x_N < x_1 + L$ with $x_k \geq a$. The first condition is equivalent to $x_k < \dots < x_N < x_1 + L < \dots < x_{k-1} + L < x_k + L$. Hence, if we set $x'_j = x_{j+k-1}$ for $j = 1, \dots, N-k+1$ and $x'_j = x_{j+k-1-N} + L$ for $j = N-k+2, \dots, N$, then the sum is over $x'_1 < \dots < x'_N < x'_1 + L$ with $x'_1 \geq a$. Consider

$$\mathcal{R}_X(W) = \det \left[w_i^{-j} (w_i + 1)^{-x_j + j} \right]_{i,j=1}^N.$$

If we move the first $k-1$ columns to the end of the matrix and use the variables x'_j , then $\mathcal{R}_X(W)$ is equal to $(-1)^{(k-1)(N-1)}$ times the determinant of the matrix whose (i,j) -entry is $w_i^{-j-k+1} (w_i + 1)^{-x'_j + j + k - 1}$ for the first $N-k+1$ columns and is $w_i^{-j-k+1+N} (w_i + 1)^{-x'_j + L + j + k - 1 - N}$ for the remaining $k-1$ columns. Since $w_i^N (w_i + 1)^{L-N} = z^L$ for $w_i \in \mathcal{R}_z$, the entries of the last $k-1$ columns are z^L times $w_i^{-j-k+1} (w_i + 1)^{-x'_j + j + k - 1}$. Thus, the row i of the matrix has the common

multiplicative factor $w_i^{-k+1}(w_i+1)^{k-1}$. Factoring out z^L from the last $k-1$ columns and also the common row factors, we find that, setting $X' = (x'_1, \dots, x'_N)$,

$$(5.4) \quad \mathcal{R}_X(W) = (-1)^{(k-1)(N-1)} z^{(k-1)L} \left[\prod_{j=1}^N \left(\frac{w_j+1}{w_j} \right)^{k-1} \right] \mathcal{R}_{X'}(W).$$

Similarly,

$$(5.5) \quad \mathcal{L}_X(W') = (-1)^{(k-1)(N-1)} (z')^{-(k-1)L} \left[\prod_{j=1}^N \left(\frac{w'_j+1}{w'_j} \right)^{-k+1} \right] \mathcal{L}_{X'}(W').$$

Hence the sum (5.2) is a certain explicit constant times the sum

$$\sum_{X' \in \mathcal{X}_N(L) \cap \{x'_1 \geq a\}} \mathcal{R}_{X'}(W) \mathcal{L}_{X'}(W')$$

which is $\mathcal{H}_{1,a}(W; W')$. Checking the multiplicative constant factor explicitly, we obtain the lemma. \square

It is thus enough to prove Proposition 3.4 for $k = 1$. Set

$$(5.6) \quad H_a(W; W') := \mathcal{H}_{1,a}(W; W') - \mathcal{H}_{1,a+1}(W; W') = \sum_{\substack{X \in \mathcal{X}_N(L) \\ x_1=a}} \mathcal{R}_X(W) \mathcal{L}_X(W').$$

We prove the following result in this section.

Proposition 5.2. *Let $z, z' \in \mathbb{C} \setminus \{0\}$ such that $z^L \neq (z')^L$. Then for every $W = (w_1, \dots, w_N) \in (\mathcal{R}_z)^N$, $W' = (w'_1, \dots, w'_N) \in (\mathcal{R}_{z'})^N$, and integer a ,*

$$(5.7) \quad H_a(W; W') = - \left(\left(\frac{z'}{z} \right)^L - 1 \right)^{N-1} \times \left(\prod_{j=1}^N \frac{w'_j(w'_j+1)^{a-1}}{w_j(w_j+1)^{a-2}} - \prod_{j=1}^N \frac{w'_j(w'_j+1)^a}{w_j(w_j+1)^{a-1}} \right) \det \left[\frac{1}{w'_{i'} - w_i} \right]_{i,i'=1}^N.$$

Note that the sum (5.6) is over the finite set $a < x_2 < \dots < x_N < a + L$. Since it is a finite sum, there is no issue of convergence, and hence we do not need to assume that $\prod_{j=1}^N |w'_j+1| < \prod_{j=1}^N |w_j+1|$. Now, if we assume this extra condition, then the sum of $H_b(W; W')$ over $b = a, a+1, a+2, \dots$ converges and is equal to $\mathcal{H}_{1,a}(W; W')$, and the resulting formula is

$$(5.8) \quad \mathcal{H}_{1,a}(W; W') = - \left(\left(\frac{z'}{z} \right)^L - 1 \right)^{N-1} \left[\prod_{j=1}^N \frac{w'_j(w'_j+1)^{a-1}}{w_j(w_j+1)^{a-2}} \right] \det \left[\frac{1}{w'_{i'} - w_i} \right]_{i,i'=1}^N.$$

This is precisely Proposition 3.4 when $k = 1$. Hence, by Lemma 5.1, Proposition 3.4 is obtained if we prove Proposition 5.2. The rest of this section is devoted to the proof of Proposition 5.2.

The proof is split into four steps. The condition that $W \in (\mathcal{R}_z)^N$, $W' \in (\mathcal{R}_{z'})^N$ is used only in step 4. Steps 1, 2, and 3 apply to any complex vectors.

5.1. Step 1. We expand $\mathcal{R}_X(W)$ and $\mathcal{L}_X(W')$ as sums and interchange the order of summation.

Let S_N be the symmetric group of order N . Let $\text{sgn}(\sigma)$ denote the sign of permutation $\sigma \in S_N$. Expanding the determinant of $\mathcal{R}_X(W)$ and $\mathcal{L}_X(W')$, we have

$$(5.9) \quad H_a(W; W') = \sum_{\sigma, \sigma' \in S_N} \text{sgn}(\sigma \sigma') \left[\prod_{j=1}^N \left(\frac{w'_{\sigma'(j)}}{w_{\sigma(j)}} \right)^j \right] S_{\sigma, \sigma'}(W; W'),$$

where

$$(5.10) \quad S_{\sigma, \sigma'}(W; W') = \sum_{\substack{X \in \mathcal{X}_N(L) \\ x_1 = a}} \prod_{j=1}^N \left(\frac{w'_{\sigma'(j)} + 1}{w_{\sigma(j)} + 1} \right)^{x_j - j}.$$

We rewrite the last sum using the following formula.

Lemma 5.3. *For complex numbers f_j , set*

$$(5.11) \quad F_{m,n} = \prod_{j=m}^n f_j \quad \text{for } 1 \leq m \leq n.$$

Then

$$(5.12) \quad \sum_{\substack{X \in \mathcal{X}_N(L) \\ x_1 = a}} \prod_{j=1}^N (f_j)^{x_j - j} = \sum_{k=0}^{N-1} \sum_{1 < s_1 < \dots < s_k \leq N} \frac{(F_{1,s_1-1})^{a-1}}{\prod_{j=2}^{s_1-1} (1 - F_{j,s_1-1})} \\ \times \prod_{i=1}^k \frac{(F_{s_i, s_{i+1}-1})^{a+L-N-1}}{(1 - (F_{s_i, s_{i+1}-1})^{-1}) \prod_{j=s_i+1}^{s_{i+1}-1} (1 - F_{j, s_{i+1}-1})},$$

where we set $s_{k+1} = N + 1$. When $k = 0$, the summand is $\frac{(F_{1,N})^{a-1}}{\prod_{j=2}^N (1 - F_{j,N})}$.

Proof. The sum is over $a < x_2 < \dots < x_N < a + L$. We evaluate the repeated sums in the following order: x_N, x_{N-1}, \dots, x_2 . For x_j the summation range is $x_{j-1} + 1 \leq x_j \leq a + L - N + j - 1$. Hence the sum is over

$$\sum_{x_2=a+1}^{a+L-N+1} \dots \sum_{x_{N-1}=x_{N-2}+1}^{a+L-2} \sum_{x_N=x_{N-1}+1}^{a+L-1}.$$

For the sum over x_j , we use the simple identity

$$(5.13) \quad \sum_{x_j=x_{j-1}+1}^{a+L-N+j-1} A^{x_j - j} = \frac{A^{x_{j-1} - (j-1)}}{1 - A} + \frac{A^{a+L-N-1}}{1 - A^{-1}}.$$

Note that the first term of the right-hand side involves x_{j-1} and the second term does not. Applying (5.13) repeatedly, the left-hand side of (5.12) is equal to a sum of 2^{N-1} terms. Each term is a product of $N - 1$ terms, some of which are from the first term of (5.13) and the rest are from the second term. We combine the summand into groups depending on how many times the second term $\frac{A^{a+L-N-1}}{1 - A^{-1}}$ is used. This number is represented by k : $0 \leq k \leq N - 1$. For given $1 \leq k \leq N - 1$, we denote by $1 \leq s_1 < \dots < s_k < N$ the indices j such that we had chosen the second term when we take the sum over x_j (5.13). The result then follows after a simple algebra. \square

We apply the lemma to $S_{\sigma, \sigma'}(W; W')$ and exchange the summation orders. Then,

$$(5.14) \quad H_a(W; W') = \sum_{k=0}^{N-1} \sum_{1 < s_1 < \dots < s_k \leq N} \sum_{\sigma, \sigma' \in S_N} \operatorname{sgn}(\sigma \sigma') \prod_{i=0}^k U_i,$$

where

$$(5.15) \quad U_0 = \frac{P_1(s_1 - 1) (Q_1(s_1 - 1))^{a-1}}{\prod_{j=2}^{s_1-1} (1 - Q_j(s_1 - 1))}$$

and, for $1 \leq i \leq k$,

$$(5.16) \quad U_i = \frac{P_{s_i}(s_{i+1} - 1) (Q_{s_i}(s_{i+1} - 1))^{a+L-N-1}}{\left(1 - (Q_{s_i}(s_{i+1} - 1))^{-1}\right) \prod_{j=s_i+1}^{s_{i+1}-1} (1 - Q_j(s_{i+1} - 1))}$$

with

$$(5.17) \quad Q_m(n) := \prod_{j=m}^n \frac{w'_{\sigma'(j)} + 1}{w_{\sigma(j)} + 1}, \quad P_m(n) := \prod_{j=m}^n \left(\frac{w'_{\sigma'(j)}}{w_{\sigma(j)}} \right)^j$$

for $1 \leq m \leq n \leq N$. Here we set $s_{k+1} := N + 1$.

We now rewrite the last two sums in (5.14) for fixed k . Given s_1, \dots, s_k and σ, σ' , let $I_i = \sigma(\{s_i, \dots, s_{i+1} - 1\})$ and $I'_i = \sigma'(\{s_i, \dots, s_{i+1} - 1\})$ for $0 \leq i \leq k$, where we set $s_0 := 1$. With this notation,

$$(5.18) \quad Q_{s_i}(s_{i+1} - 1) = \frac{\prod_{j' \in I'_i} (w'_{j'} + 1)}{\prod_{j \in I_i} (w_j + 1)}.$$

Consider the restriction of the permutation σ on $\{s_i, \dots, s_{i+1} - 1\}$. This gives rise to a bijection σ_i from $\{1, \dots, |I_i|\}$ to I_i by setting $\sigma_i(\ell) = \sigma(s_i + \ell - 1)$. Similarly we obtain a bijection σ'_i from σ . Then we have

$$(5.19) \quad Q_{s_i+j}(s_{i+1} - 1) = \prod_{\ell=j}^{|I_i|} \left(\frac{w'_{\sigma'_i(\ell)} + 1}{w_{\sigma_i(\ell)} + 1} \right)$$

for $j = 1, \dots, |I_i|$, and

$$(5.20) \quad P_{s_i}(s_{i+1} - 1) = \left[\prod_{j=1}^{|I_i|} \left(\frac{w'_{\sigma'_i(j)} + 1}{w_{\sigma_i(j)} + 1} \right)^j \right] \left(\frac{\prod_{j' \in I'_i} w'_{j'}}{\prod_{j \in I_i} w_j} \right)^{|I_0| + \dots + |I_{i-1}|}.$$

We now reorganize the summations over s_1, \dots, s_k and σ, σ' in (5.14) as follows. We first decide on two partitions I_0, \dots, I_k and I'_0, \dots, I'_k of $\{1, \dots, N\}$ satisfying $|I_i| = |I'_i| \neq 0$ for all i . Then for each I_i , we consider a bijection σ_i from $\{1, \dots, |I_i|\}$ to I_i and a bijection σ'_i from $\{1, \dots, |I_i|\}$ to I_i . The collection of $\sigma_0, \dots, \sigma_k$ is equivalent to a permutation σ . Note that the sign becomes

$$(5.21) \quad \operatorname{sgn}(\sigma) = (-1)^{\#(I_0, \dots, I_k)} \prod_{i=0}^k \operatorname{sgn}(\sigma_i),$$

where

$$(5.22) \quad \#(I_0, \dots, I_k) := |\{(m, n) \in I_i \times I_j : m < n, i > j\}|.$$

It is easy to see that summing over the partitions and then over the bijections σ_i and σ'_i is equivalent to summing over s_1, \dots, s_k and σ, σ' in (5.14). Hence we obtain

(5.23)

$$H_a(W; W') = \sum_{k=0}^{N-1} \sum_{\mathcal{I}, \mathcal{I}'} (-1)^{\#(\mathcal{I}) + \#(\mathcal{I}')} \left[\prod_{i=1}^k \left(\frac{\prod_{j' \in I'_i} w'_{j'}}{\prod_{j \in I_i} w_j} \right)^{|I_0| + \dots + |I_{i-1}|} \right] \tilde{\mathcal{G}}(I_0, I'_0) \prod_{i=1}^k \mathcal{G}(I_i, I'_i),$$

where the second sum is over two partitions $\mathcal{I} = (I_0, \dots, I_k)$ and $\mathcal{I}' = (I'_0, \dots, I'_k)$ of $\{1, \dots, N\}$ such that $|I_i| = |I'_i| \neq 0$ for all $0 \leq i \leq k$. The function $\mathcal{G}(I; I')$ on two subsets of $\{1, \dots, N\}$ of equal cardinality is defined by

(5.24)

$$\mathcal{G}(I, I') = \left[\sum_{\substack{\sigma: \{1, \dots, |I|\} \rightarrow I \\ \sigma': \{1, \dots, |I'|\} \rightarrow I'}} \text{sgn}(\sigma\sigma') \frac{P(\sigma, \sigma')}{\prod_{j=2}^{|I|} (1 - Q_j(\sigma, \sigma'))} \right] \frac{Q(I, I')^{a+L-N-1}}{1 - Q(I, I')^{-1}}.$$

Here the sum is over bijections σ and σ' , and

$$(5.25) \quad \begin{aligned} P(\sigma, \sigma') &= \prod_{\ell=1}^{|I|} \left(\frac{w'_{\sigma'(\ell)}}{w_{\sigma(\ell)}} \right)^\ell, & Q_j(\sigma, \sigma') &= \prod_{\ell=j}^{|I|} \left(\frac{w'_{\sigma'(\ell)} + 1}{w_{\sigma(\ell)} + 1} \right), \\ Q(I, I') &= \frac{\prod_{\ell' \in I'} (w'_{\ell'} + 1)}{\prod_{\ell \in I} (w_\ell + 1)}. \end{aligned}$$

Note that $Q_1(\sigma, \sigma')$ does not depend on σ, σ' and is equal to $Q(I, I')$. On the other hand,

$$(5.26) \quad \tilde{\mathcal{G}}(I, I') = \left[\sum_{\substack{\sigma: \{1, \dots, |I|\} \rightarrow I \\ \sigma': \{1, \dots, |I'|\} \rightarrow I'}} \text{sgn}(\sigma\sigma') \frac{P(\sigma, \sigma')}{\prod_{j=2}^{|I|} (1 - Q_j(\sigma, \sigma'))} \right] Q(I, I')^{a-1}.$$

5.2. Step 2. We simplify $\tilde{\mathcal{G}}(I, I')$ and $\mathcal{G}(I, I')$. They have a common sum. We claim that this sum is equal to

$$(5.27) \quad \left(\frac{\prod_{i' \in I'} w'_{i'}}{\prod_{i \in I} w_i} \right) \left(\prod_{i \in I} (w_i + 1) - \prod_{i' \in I'} (w'_{i'} + 1) \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in I, i' \in I'}.$$

This implies that

$$(5.28) \quad \begin{aligned} \tilde{\mathcal{G}}(I, I') &= \left(\frac{\prod_{i' \in I'} (w'_{i'} + 1)}{\prod_{i \in I} (w_i + 1)} \right)^{a-1} \left(\frac{\prod_{i' \in I'} w'_{i'}}{\prod_{i \in I} w_i} \right) \\ &\times \left(\prod_{i \in I} (w_i + 1) - \prod_{i' \in I'} (w'_{i'} + 1) \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in I, i' \in I'} \end{aligned}$$

and

(5.29)

$$\mathcal{G}(I, I') = - \frac{\left(\prod_{i' \in I'} (w'_{i'} + 1) \right)^{a+L-N}}{\left(\prod_{i \in I} (w_i + 1) \right)^{a+L-N-1}} \left(\frac{\prod_{i' \in I'} w'_{i'}}{\prod_{i \in I} w_i} \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in I, i' \in I'}.$$

We now prove (5.27). This follows from the next lemma. This lemma assumes that $I = I' = \{1, \dots, n\}$ but the general case is obtained if we relabel the indices.

Lemma 5.4. *For complex numbers w_i and w'_i , $i = 1, \dots, n$,*

$$(5.30) \quad \begin{aligned} & \sum_{\sigma, \sigma' \in S_n} \frac{\operatorname{sgn}(\sigma \sigma') \prod_{i=1}^n \left(\frac{w'_{\sigma'(i)}}{w_{\sigma(i)}} \right)^{i-1}}{\prod_{j=2}^n \left(1 - \prod_{i=j}^n \frac{w'_{\sigma'(i)} + 1}{w_{\sigma(i)} + 1} \right)} \\ &= \left(\prod_{j=1}^n (w_j + 1) - \prod_{j=1}^n (w'_j + 1) \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i, i'=1}^n. \end{aligned}$$

Proof. We use an induction in n . It is direct to check that the identity holds for $n = 1$.

Let $n \geq 2$ and assume that the identity holds for index $n - 1$. We now prove the identity for index n . We consider the sum according to different values of $\sigma(1)$ and $\sigma'(1)$. Setting $\sigma(1) = l$ and $\sigma'(1) = l'$, and then renaming the shifted version of the rest of σ and σ' by again σ and σ' , respectively, the left-hand side of (5.30) is equal to

$$\begin{aligned} & \sum_{\ell, \ell'=1}^n (-1)^{\ell+\ell'} \frac{\frac{w_\ell}{w'_{\ell'}} \prod_{k=1}^n \left(\frac{w'_k}{w_k} \right)}{1 - \frac{w_\ell + 1}{w'_{\ell'} + 1} \prod_{k=1}^n \left(\frac{w'_k + 1}{w_k + 1} \right)} \\ & \times \left[\sum_{\substack{\sigma: \{1, \dots, n-1\} \rightarrow \{1, \dots, n\} \setminus \{\ell\} \\ \sigma': \{1, \dots, n-1\} \rightarrow \{1, \dots, n\} \setminus \{\ell'\}}} \frac{\operatorname{sgn}(\sigma) \operatorname{sgn}(\sigma') \prod_{i=2}^{n-1} \left(\frac{w'_{\sigma'(i)}}{w_{\sigma(i)}} \right)^{i-1}}{\prod_{j=2}^{n-1} \left(1 - \prod_{i=j}^{n-1} \frac{w'_{\sigma'(i)} + 1}{w_{\sigma(i)} + 1} \right)} \right]. \end{aligned}$$

From the induction hypothesis, the bracket term is equal to

$$\left(\prod_{\substack{j=1 \\ j \neq \ell}}^n (w_j + 1) - \prod_{\substack{j=1 \\ j \neq \ell'}}^n (w'_j + 1) \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{\substack{1 \leq i, i' \leq n \\ i \neq \ell, i' \neq \ell'}}.$$

Hence the sum is equal to

$$\sum_{\ell, \ell'=1}^n (-1)^{\ell+\ell'} \left(\prod_{\substack{1 \leq k' \leq n \\ k' \neq \ell'}} w'_{k'} \right) \left(\prod_{\substack{1 \leq k \leq n \\ k \neq \ell}} \frac{w_k + 1}{w_k} \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{\substack{1 \leq i, i' \leq n \\ i \neq \ell, i' \neq \ell'}}.$$

This is the same as

$$\left(\prod_{k=1}^n \frac{w'_k (w_k + 1)}{w_k} \right) \left(\sum_{i, i'=1}^n (-1)^{\ell+\ell'} \frac{w_\ell}{(w_\ell + 1) w'_{\ell'}} \det \left[\frac{1}{w_i - w'_{i'}} \right]_{\substack{1 \leq i, i' \leq n \\ i \neq \ell, i' \neq \ell'}} \right).$$

Lemma 5.5 below implies that the sum in the second parentheses is equal to

$$\left(\prod_{k=1}^n \frac{w_k}{w'_k (w_k + 1)} \right) \left(\prod_{i=1}^n (w_i + 1) - \prod_{i=1}^n (w'_i + 1) \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i, i'=1}^n.$$

This completes the proof. \square

Lemma 5.5. *For distinct complex numbers x_1, \dots, x_n and y_1, \dots, y_n , let C be the $n \times n$ Cauchy matrix with entries $\frac{1}{x_i - y_j}$. Let $C_{\ell,k}$ be the matrix obtained from C by removing row ℓ and column k . Define the functions*

$$(5.31) \quad A(z) = \prod_{i=1}^n (z - x_i), \quad B(z) = \prod_{i=1}^n (z - y_i).$$

Then

$$(5.32) \quad \sum_{\ell,k=1}^n (-1)^{\ell+k} \frac{x_\ell}{(x_\ell + 1)y_k} \det [C_{\ell,k}] = \frac{A(0)}{B(0)} \left(1 - \frac{B(-1)}{A(-1)} \right) \det [C].$$

Proof. Recall the Cauchy determinant formula,

$$\det [C] = \frac{\prod_{i < j} (x_i - x_j)(y_j - y_i)}{\prod_{i,j} (x_i - y_j)}.$$

Note that $C_{\ell,k}$ is also a Cauchy matrix. Hence we find that

$$(5.33) \quad \frac{\det [C_{\ell,k}]}{\det [C]} = (-1)^{\ell+k+1} \frac{B(x_\ell)A(y_k)}{(x_\ell - y_k)A'(x_\ell)B'(y_k)}.$$

Let f and g be meromorphic functions with finitely many poles and consider the double integral

$$(5.34) \quad \frac{1}{(2\pi i)^2} \oint_{|\xi|=r_2} \oint_{|z|=r_1} f(z)g(\xi) \frac{B(z)A(\xi)}{(z - \xi)A(z)B(\xi)} dz d\xi.$$

Here we take $r_1 < r_2$ large so that the poles of $\frac{f(z)}{A(z)}$ are inside $|z| < r_1$ and the poles of $\frac{g(\xi)}{B(\xi)}$ are inside $|\xi| < r_2$. Consider $f(z) = \frac{z}{z+1}$ and $g(\xi) = \frac{1}{\xi}$. By changing the order of integration and noting that the integrand is $O(\xi^{-2})$ as $\xi \rightarrow \infty$ for fixed z , the double integral is zero by taking the ξ contour to infinity. On the other hand, we also evaluate the double integral by residue calculus. For given ξ , the integral in z is equal to

$$\frac{A(\xi)B(-1)}{\xi(\xi + 1)B(\xi)A(-1)} + \sum_{\ell=1}^n \frac{x_\ell A(\xi)B(x_\ell)}{(x_\ell + 1)\xi(x_\ell - \xi)B(\xi)A'(x_\ell)}.$$

The first term is $O(\xi^{-2})$ and hence the integral with respect to ξ is zero. The integral of the second term is, by residue calculus,

$$\left[\sum_{\ell=1}^n \frac{B(x_\ell)}{(x_\ell + 1)A'(x_\ell)} \right] \frac{A(0)}{B(0)} + \sum_{\ell,k=1}^n \frac{x_\ell B(x_\ell)A(y_k)}{(x_\ell + 1)y_k(x_\ell - y_k)A'(x_\ell)B'(y_k)}.$$

Since the double integral is zero, the above expression is zero. Now the sum inside the bracket can be simplified to $1 - \frac{B(-1)}{A(-1)}$ by considering the integral of $\frac{B(z)}{(z+1)A(z)}$. Therefore, we obtain

$$\sum_{\ell,k=1}^n \frac{x_\ell B(x_\ell)A(y_k)}{(x_\ell + 1)y_k(x_\ell - y_k)A'(x_\ell)B'(y_k)} = - \left(1 - \frac{B(-1)}{A(-1)} \right) \frac{A(0)}{B(0)}.$$

Using (5.33), we obtain the lemma. \square

Remark 5.6. Note that, using the rank one property and Hadamard's formula for the inverse of a matrix,

$$\begin{aligned}
 (5.35) \quad \det \left[\frac{1}{x_i - y_j} + f(x_i)g(y_j) \right]_{i,j=1}^n &= \det[C] \left(1 + \sum_{\ell,k=1}^n g(y_k)(C^{-1})_{k,\ell} f(x_\ell) \right) \\
 &= \det[C] + \sum_{\ell,k=1}^n (-1)^{\ell+k} \det[C_{\ell,k}] f(x_\ell)g(y_k).
 \end{aligned}$$

Hence, the above lemma implies that

$$(5.36) \quad \det \left[\frac{1}{x_i - y_j} + \frac{x_i}{(x_i + 1)y_j} \right]_{i,j=1}^n = \left(1 + \frac{A(0)}{B(0)} - \frac{A(0)B(-1)}{B(0)A(-1)} \right) \det \left[\frac{1}{x_i - y_j} \right]_{i,j=1}^n.$$

We will need the following variation of the above lemma in the next subsection.

Lemma 5.7. *Using the same notation as in Lemma 5.5,*

$$(5.37) \quad \det \left[\frac{1}{x_i - y_j} + \frac{u}{x_j} \right]_{i,j=1}^n = \left(1 + u \left(1 - \frac{B(0)}{A(0)} \right) \right) \det[C].$$

Proof. The proof is similar to the previous lemma using $f(z) = \frac{u}{z}$ and $g(\xi) = 1$ in (5.34) instead. We also use (5.35). \square

5.3. Step 3. We insert the formulas (5.28) and (5.29) into (5.23), and then reorganize the sum as follows. For the partitions $\mathcal{I} = (I_0, \dots, I_k)$ and $\mathcal{I}' = (I'_0, \dots, I'_k)$, we consider the first parts I_0 and I'_0 separately. Set $\tilde{\mathcal{I}} = (I_1, \dots, I_k)$ and $\tilde{\mathcal{I}}' = (I'_1, \dots, I'_k)$. Note that

$$\#(\mathcal{I}) = \#(I_0, I_0^c) + \#(\tilde{\mathcal{I}}), \quad \#(\mathcal{I}') = \#(I'_0, (I'_0)^c) + \#(\tilde{\mathcal{I}}'),$$

where $I_0^c = \{1, \dots, N\} \setminus I_0$ and $(I'_0)^c = \{1, \dots, N\} \setminus I'_0$. Then (5.23) can be written as

$$\begin{aligned}
 (5.38) \quad H_a(W; W') &= \sum_{\substack{I_0, I'_0 \subseteq \{1, \dots, N\} \\ |I_0| = |I'_0| \neq 0}} (-1)^{\#(I_0, I_0^c) + \#(I'_0, (I'_0)^c)} \tilde{\mathcal{G}}(I_0, I'_0) \\
 &\quad \times \left[\frac{\left(\prod_{i' \in (I'_0)^c} (w'_{i'} + 1) \right)^{a+L-N}}{\left(\prod_{i \in I_0^c} (w_i + 1) \right)^{a+L-N-1}} \right] S(I_0, I'_0),
 \end{aligned}$$

where

$$\begin{aligned}
 (5.39) \quad S(I_0, I'_0) &= \sum_{k=1}^{N-|I_0|} \sum_{\tilde{\mathcal{I}}, \tilde{\mathcal{I}}'} (-1)^{k + \#(\tilde{\mathcal{I}}) + \#(\tilde{\mathcal{I}}')} \\
 &\quad \times \prod_{j=1}^k \left[\left(\frac{\prod_{i' \in I'_j} w'_{i'}}{\prod_{i \in I_j} w_i} \right)^{|I_0| + \dots + |I_{j-1}| + 1} \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in I_j, i' \in I'_j} \right].
 \end{aligned}$$

Here the second sum in (5.39) is over all partitions $\tilde{\mathcal{I}} = (I_1, \dots, I_k)$ of $\{1, \dots, N\} \setminus I_0$ and partitions $\tilde{\mathcal{I}}' = (I'_1, \dots, I'_k)$ of $\{1, \dots, N\} \setminus I'_0$ satisfying $|I_i| = |I'_i| \neq 0$ for all

$1 \leq i \leq k$. When $|I_0| = N$, we understand that $S(I_0, I'_0) = 1$. The following lemma simplifies $S(I_0, I'_0)$.

Lemma 5.8. *Let $n \geq 1$. Let w_i and w'_i , $1 \leq i \leq n$, be complex numbers. Then*

$$(5.40) \quad \begin{aligned} & \sum_{k=1}^n \sum_{\mathcal{J}, \mathcal{J}'} (-1)^{k+\#(\mathcal{J})+\#(\mathcal{J}')} \prod_{j=1}^k \left(\frac{\prod_{i' \in J'_j} w'_{i'}}{\prod_{i \in J_j} w_i} \right)^{1+\sum_{\ell=1}^{j-1} |J_\ell|} \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in J_j, i' \in J'_j} \\ &= \left[\prod_{j=1}^n \left(\frac{w'_j}{w_j} \right)^n \right] \det \left[\frac{1}{-w_i + w'_{i'}} \right]_{1 \leq i, i' \leq n}. \end{aligned}$$

Here the second sum is over all partitions $\mathcal{J} = (J_1, \dots, J_k)$ and $\mathcal{J}' = (J'_1, \dots, J'_k)$ of $\{1, \dots, n\}$ such that $|J_i| = |J'_i| \neq 0$ for all $1 \leq i \leq k$.

Proof. We use an induction in n . When $n = 1$ the identity is trivial. Now we prove that the identity holds for index n assuming that it holds for all indices less than n . We fix $J_1 = J$ and $J'_1 = J'$ with $|J| = |J'| \neq 0$ and then apply the induction hypothesis on the remaining sum with index $n - |J|$. Then, the left-hand side of (5.40) is equal to

$$(5.41) \quad \begin{aligned} & - \sum_{\substack{J, J' \subseteq \{1, \dots, n\} \\ |J| = |J'| \neq 0}} (-1)^{\#(J, J^c) + \#(J', (J')^c)} \left(\frac{\prod_{i' \in J'} w'_{i'}}{\prod_{i \in J} w_i} \right) \left(\frac{\prod_{i' \in (J')^c} w'_{i'}}{\prod_{i \in J^c} w_i} \right)^n \\ & \quad \times \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in J, i' \in J'} \det \left[\frac{1}{-w_i + w'_{i'}} \right]_{i \in J^c, i' \in (J')^c}. \end{aligned}$$

Here the minus sign comes from $(-1)^k$, where k , the number of parts of partition \mathcal{J} , is reduced by 1 when we apply the induction hypothesis since we remove J_1 in the counting. Now, note that the right-hand side of (5.40) is the summand of (5.41) when $|J| = |J'| = 0$. Hence, after dividing by $\prod_{j=1}^n \left(\frac{w'_j}{w_j} \right)^n$, we find that (5.40) is obtained if we show that

$$\begin{aligned} & \sum_{\substack{J, J' \subseteq \{1, \dots, n\} \\ |J| = |J'|}} (-1)^{\#(J, J^c) + \#(J', (J')^c)} \left(\frac{\prod_{i' \in J'} w'_{i'}}{\prod_{i \in J} w_i} \right)^{1-n} \\ & \quad \times \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in J, i' \in J'} \det \left[\frac{1}{-w_i + w'_{i'}} \right]_{i \in J^c, i' \in (J')^c} \end{aligned}$$

is equal to 0. Here the sum is over all $J, J' \subseteq \{1, \dots, n\}$ with $|J| = |J'|$, including the case when $|J| = |J'| = 0$. By Lemma 5.9 below, this sum is equal to

$$\det \left[\frac{(w_i)^{n-1}}{(w'_{i'})^{n-1}} \frac{1}{w_i - w'_{i'}} + \frac{1}{-w_i + w'_{i'}} \right]_{i, i'=1}^n.$$

Hence it is enough to show that

$$\det \left[\frac{(w'_{i'})^{n-1} - w_i^{n-1}}{w'_{i'} - w_i} \right]_{i, i'=1}^n = 0.$$

Using $\frac{a^{n-1} - b^{n-1}}{a-b} = a^{n-2} + a^{n-3}b + \cdots + b^{n-2}$, this last determinant is a sum of the determinants of the form

$$\det [(w'_{i'})^{\alpha_i} w_i^{n-2-\alpha_i}]_{i,i'=1}^n = \det [(w'_{i'})^{\alpha_i}]_{i,i'=1}^n \prod_{i=1}^n w_i^{n-2-\alpha_i}$$

for some $\alpha_i \in \{0, 1, \dots, n-2\}$. The last determinant is zero since at least two rows are equal. This completes the proof. \square

Lemma 5.9. *For two $n \times n$ matrices A and B ,*

$$(5.42) \quad \begin{aligned} & \sum_{\substack{J, J' \subseteq \{1, \dots, n\} \\ |J|=|J'|}} (-1)^{\#(J, J^c) + \#(J', (J')^c)} \det [A(i, i')]_{i \in J, i' \in J'} \det [B(i, i')]_{i \in J^c, i' \in (J')^c} \\ & = \det [A + B], \end{aligned}$$

where the sum is over all subsets J and J' of equal size, including the case when $J = J' = \emptyset$.

Proof. It is direct to check by expanding all determinants by sums using definition. \square

We apply Lemma 5.8 to $S(I_0, I'_0)$. Note that the power of the products of $w'_{i'}$ and w_i in (5.39) is $|I_0| + |I_1| + \cdots + |I_k| + 1$ while the corresponding products have power $|J_1| + \cdots + |J_k| + 1$ in the lemma. We find that

$$(5.43) \quad S(I_0, I'_0) = \left(\frac{\prod_{i' \in (I'_0)^c} w'_{i'}}{\prod_{i \in I_0^c} w_i} \right)^N \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in I_0^c, i' \in (I'_0)^c}.$$

Using this formula and also the formula (5.28) for $\tilde{G}(I_0, I'_0)$, the equation (5.38) becomes

$$(5.44) \quad \begin{aligned} H_a(W; W') &= \sum_{\substack{I_0, I'_0 \subseteq \{1, \dots, N\} \\ |I_0|=|I'_0| \neq 0}} (-1)^{\#(I_0, I_0^c) + \#(I'_0, (I'_0)^c)} \\ &\times \left[\frac{\prod_{i' \in I'_0} w'_{i'} (w'_{i'} + 1)^a}{\prod_{i \in I_0} w_i (w_i + 1)^{a-1}} \right] \left[\frac{\prod_{i' \in (I'_0)^c} (w'_{i'})^N (w'_{i'} + 1)^{a+L-N}}{\prod_{i \in I_0^c} w_i^N (w_i + 1)^{a+L-N-1}} \right] \\ &\times \left(\frac{\prod_{i \in I_0} (w_i + 1)}{\prod_{i' \in I'_0} (w'_{i'} + 1)} - 1 \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in I_0, i' \in I'_0} \det \left[\frac{1}{-w_i + w'_{i'}} \right]_{i \in I_0^c, i' \in (I'_0)^c}, \end{aligned}$$

where we added the zero partitions $|I_0| = |I'_0| = 0$ in the sum since the summand is zero in that case.

5.4. Step 4. We evaluate the sum (5.44). So far w_i and $w'_{i'}$ were any complex numbers. We now use the assumption that $w_i \in \mathcal{R}_z$ and $w'_{i'} \in \mathcal{R}_{z'}$. This means that

$$w_i^N (w_i + 1)^{L-N} = z^L, \quad (w'_{i'})^N (w'_{i'} + 1)^{L-N} = (z')^L$$

for all i and i' , and hence the second square bracket in (5.44) can be simplified. We then separate the formula of $H_a(W; W')$ into two terms using the two terms

in the big parentheses. The second term, which comes from the term 1 in the big parenthesis, can be written as

$$\left[\prod_{j=1}^N \frac{(w'_j + 1)^a}{(w_j + 1)^{a-1}} \right] \left\{ \sum_{\substack{I_0, I'_0 \subseteq \{1, \dots, N\} \\ |I_0| = |I'_0|}} (-1)^{\#(I_0, I'_0) + \#(I'_0, (I'_0)^c)} \left[\frac{\prod_{i' \in I'_0} w'_{i'}}{\prod_{i \in I_0} w_i} \right] \right. \\ \left. \times \left[\frac{\prod_{i' \in (I'_0)^c} (z')^L}{\prod_{i \in I'_0} z^L} \right] \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i \in I_0, i' \in I'_0} \det \left[\frac{1}{-w_i + w'_{i'}} \right]_{i \in I'_0, i' \in (I'_0)^c} \right\}.$$

The sum is, after inserting the terms in the products into the determinants, of the form in Lemma 5.9 above, and hence is equal to

$$\det \left[\frac{w'_{i'}}{w_i(w_i - w'_{i'})} - \left(\frac{z'}{z} \right)^L \frac{1}{w_i - w'_{i'}} \right]_{1 \leq i, i' \leq N}.$$

The first term is also similar, and we obtain

$$(5.45) \quad H_a(W; W') = \left[\prod_{j=1}^N \frac{(w'_j + 1)^a}{(w_j + 1)^{a-1}} \right] (D_1 - D_2),$$

where, for $k = 1, 2$,

$$(5.46) \quad D_k = \det \left[\frac{\frac{f_k(w'_{i'})}{f_k(w_i)}}{w_i - w'_{i'}} - \frac{q}{w_i - w'_{i'}} \right]_{i, i'=1}^N$$

with

$$(5.47) \quad f_1(w) = \frac{w}{w+1}, \quad f_2(w) = w, \quad q = \left(\frac{z'}{z} \right)^L.$$

The entries of the determinants D_k are of the form

$$\frac{\frac{f(w'_{i'})}{f(w_i)} - 1}{w_i - w'_{i'}} + \frac{1 - q}{w_i - w'_{i'}} = (1 - q) \left(\frac{1}{w_i - w'_{i'}} + \frac{f(w'_{i'}) - f(w_i)}{(1 - q)f(w_i)(w_i - w'_{i'})} \right),$$

where f represents either f_1 or f_2 . For $f(w) = f_2(w) = w$, the last term is $-\frac{1}{(1-q)w_i}$ and

$$(5.48) \quad D_2 = (1 - q)^N \det \left[\frac{1}{w_i - w'_{i'}} - \frac{1}{(1 - q)w_i} \right] \\ = (1 - q)^N \left(1 - \frac{1}{1 - q} \left(1 - \prod_{i=1}^N \frac{w'_{i'}}{w_i} \right) \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i, i'=1}^N,$$

where we used Lemma 5.7 for the second equality. On the other hand,

$$D_1 = (1 - q)^N \det \left[\frac{1}{w_i - w'_{i'}} - \frac{1}{(1 - q)w_i(w'_{i'} + 1)} \right].$$

We may evaluate this determinant by finding a variation of Lemma 5.7. Alternatively, we set $w_i = \frac{1}{x_i - 1}$ and $w'_{i'} = \frac{1}{y_{i'} - 1}$, and use Lemma 5.7 to obtain

$$(5.49) \quad \begin{aligned} D_1 &= \frac{(1-q)^N}{\prod_{i=1}^N (w_i w'_i)} \det \left[\frac{1}{y_{i'} - x_i} - \frac{1}{(1-q)y_{i'}} \right] \\ &= (1-q)^N \left(1 - \frac{1}{1-q} \left(1 - \prod_{j=1}^N \frac{(w_j + 1)w'_j}{w_j(w'_j + 1)} \right) \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i,i'=1}^N. \end{aligned}$$

From (5.45), (5.48), and (5.49), we obtain

$$H_a(W; W') = \left(1 - \left(\frac{z'}{z} \right)^L \right)^{N-1} \left[\prod_{j=1}^N \frac{w'_j}{w_j} \right] \left(\prod_{j=1}^N \frac{(w'_j + 1)^{a-1}}{(w_j + 1)^{a-2}} - \prod_{j=1}^N \frac{(w'_j + 1)^a}{(w_j + 1)^{a-1}} \right) \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i,i'=1}^N.$$

This completes the proof of Proposition 5.2.

6. PROOF OF THEOREM 2.1

In this section, we prove Theorem 2.1 starting from Corollary 4.7.

6.1. Parameters. We evaluate the limit of

$$(6.1) \quad \mathbb{P}(\mathbb{h}(\mathbf{p}_1) \geq b_1, \dots, \mathbb{h}(\mathbf{p}_m) \geq b_m)$$

as $L \rightarrow \infty$, where $\mathbf{p}_j = s_j \mathbf{e}_1 + t_j \mathbf{e}_c = \ell_j \mathbf{e}_1 + t_j \mathbf{e}_2$ with $s_j = \ell_j - (1-2\rho)t_j$,

$$(6.2) \quad t_j = \tau_j \frac{L^{3/2}}{\sqrt{\rho(1-\rho)}} + O(L),$$

$$(6.3) \quad \ell_j = (1-2\rho)t_j + \gamma_j L + O(L^{1/2}),$$

and

$$(6.4) \quad b_j = 2\rho(1-\rho)t_j + (1-2\rho)\ell_j - 2x_j \rho^{1/2} (1-\rho)^{1/2} L^{1/2}$$

for fixed $0 < \tau_1 < \dots < \tau_m$, $\gamma_j \in [0, 1]$, and $x_j \in \mathbb{R}$. The analysis needs small changes if $\tau_i = \tau_{i+1}$ and $x_i < x_{i+1}$ for some i . We will comment on such changes in the footnotes throughout the analysis.

It is tedious but easy to check in the analysis that the convergence is uniform for the parameters τ_j , γ_j , and x_j in compact sets. In order to keep the writing light, we do not keep track of the uniformity.

The height function and the particle location have the following relation:

$$(6.5) \quad \mathbb{h}(\ell \mathbf{e}_1 + t \mathbf{e}_2) \geq b \text{ if and only if } \mathbb{x}_{N-\frac{b-\ell}{2}+1}(t) \geq \ell + 1$$

for all $b - 2\ell \in 2\mathbb{Z}$ satisfying $b \geq \mathbb{h}(\ell \mathbf{e}_1)$. Hence,

$$(6.6) \quad \mathbb{P}(\mathbb{h}(\mathbf{p}_1) \geq b_1, \dots, \mathbb{h}(\mathbf{p}_m) \geq b_m) = \mathbb{P}(\mathbb{x}_{k_1}(t_1) \geq a_1, \dots, \mathbb{x}_{k_m}(t_m) \geq a_m),$$

where t_j are the same as above,

$$(6.7) \quad a_j = \ell_j + 1,$$

and

$$(6.8) \quad k_j = N - \frac{b_j - \ell_j}{2} + 1.$$

From Corollary 4.7, we thus need to evaluate the limit of

$$(6.9) \quad \mathbb{P}(\mathbb{x}_{k_1}(t_1) \geq a_1, \dots, \mathbb{x}_{k_m}(t_m) \geq a_m) = \oint \cdots \oint C(\mathbf{Z}) D(\mathbf{Z}) \frac{dZ_1}{2\pi i Z_1} \cdots \frac{dZ_m}{2\pi i Z_m},$$

where the contours are nest circles satisfying $0 < |Z_m| < \cdots < |Z_1| < \mathbb{r}_0$. Here we use the notation Z_j instead of z_j . The notation z_j will be reserved for the rescaled parameter of Z_j in the asymptotic analysis.

In [4] and [30], we analyzed the case when $m = 1$. The case when $m \geq 2$ is similar and we follow the same strategy. The results and analysis of the above two papers are used heavily in this section.

We change the variables Z_j to z_j defined by

$$(6.10) \quad z_j = (-1)^N \frac{Z_j^L}{\mathbb{r}_0^L}.$$

Then $\frac{dZ_j}{Z_j} = \frac{dz_j}{Lz_j}$, but the simple closed circle of Z_j becomes a circle with multiplicity L of z_j . Note that $C(\mathbf{Z})$ and $D(\mathbf{Z})$ depend on Z_j through the set of the roots of the equation $w^N(w+1)^{L-N} = Z_j^L$, which is unchanged if Z_j is changed to $Z_j e^{2\pi i k/L}$ for integer k . Hence the above integral is the same as

$$(6.11) \quad \mathbb{P}(\mathbb{x}_{k_1}(t_1) \geq a_1, \dots, \mathbb{x}_{k_m}(t_m) \geq a_m) = \oint \cdots \oint C(\mathbf{Z}) D(\mathbf{Z}) \frac{dz_1}{2\pi i z_1} \cdots \frac{dz_m}{2\pi i z_m},$$

where the integrals are nested simple circles such that $0 < |z_m| < \cdots < |z_1| < 1$ on the contours, and for given z_j , Z_j is any one of the L roots of the equation $Z_j^L = (-1)^L \mathbb{r}_0^L z_j$.

Remark 6.1. We note that the analysis in this section does not depend on the fact that $|z_i|$ are ordered in a particular way. It is easy to check in each step of the analysis that we only require that $|z_i|$ are distinct. Hence, if we fix $0 < r_m < \cdots < r_1 < 1$, and $|z_j| = r_{\sigma(j)}$ for a permutation σ of $1, \dots, m$, then the asymptotics and error estimates of $C(\mathbf{Z})$, $D(\mathbf{Z})$ and the integral on the right-hand side of (6.11) still hold (with different constants ϵ, c, C in the error estimates). This fact is used in an important way in the next section; see the proof of Theorem 7.3.

6.2. Asymptotics of $C(\mathbf{Z})$.

Lemma 6.2 ([30]). *Let $N = N_L$ be a sequence of integers such that $\rho = \rho_L := \frac{N}{L}$ stays in a compact subset of $(0, 1)$ for all L . Fix $\epsilon \in (0, 1/2)$. Fix a complex number z such that $0 < |z| < 1$ and let Z satisfy $Z^L = (-1)^N \mathbb{r}_0^L z$. Assume that for fixed $\tau > 0$, $\gamma \in [0, 1)$, and $x \in \mathbb{R}$,*

$$(6.12) \quad t = \frac{\tau}{\sqrt{\rho(1-\rho)}} L^{3/2} + O(L), \quad a = \ell + 1, \quad k = N - \frac{b - \ell}{2} + 1,$$

where

$$(6.13) \quad \ell = (1-2\rho)t + \gamma L + O(L^{1/2}), \quad b = 2\rho(1-2\rho)t + (1-2\rho)\ell - 2x\rho^{1/2}(1-\rho)^{1/2}L^{1/2}.$$

Recall the definition of $E(Z)$ in (4.9):

$$(6.14) \quad E(Z) = E(Z; a, k, t) := \prod_{u \in L_Z} (-u)^{k-N-1} \prod_{v \in R_Z} (v+1)^{-a+k-N} e^{tv}.$$

Then

$$(6.15) \quad E(Z) = e^{xA_1(z) + \tau A_2(z)} \left(1 + O(N^{\epsilon-1/2}) \right),$$

where $A_1(z)$ and $A_2(z)$ are polylogarithm functions defined in (2.18).

Proof. This follows from (4.25) and (4.26) of Section 4.3 of [30]: we have $E(Z; a, k, t) = \mathcal{C}_{N,2}^{(2)}(Z; k-N, a)$ in terms of the notation used in [30]. (Note that the limit there is $e^{\tau^{1/3} x A_1(z) + \tau A_2(z)}$, but in this paper, we use the scale of the height function so that $\tau^{1/3} x$ is changed to x .) It is easy to check that the conditions (4.1), (4.2), and (4.3) in [30] are satisfied, and hence we obtain the above lemma. The analysis was based on the integral representation of $\log E(Z)$ (see (4.28) of [30]) and applying the method of the steepest-descent. Indeed, by the residue theorem ($w = 0$ is the case we need),

$$(6.16) \quad \log \left(\prod_{u \in L_Z} (w-u) \right) = (L-N) \log(w+1) + \frac{LZ^L}{2\pi i} \oint \frac{(u+\rho) \log(w-u)}{u(u+1)q_Z(u)} du$$

for $\Re(w) > -\rho$, where the contour is a simple closed curve in $\Re(u) < -\rho$ which contains L_Z inside. We can find a similar formula for the product of v . \square

Lemma 6.3. *Let N , L , and ϵ be as in the previous lemma. Fix complex numbers z_1, z_2 such that $0 < |z_1|, |z_2| < 1$. Then for Z_1, Z_2 satisfying $Z_1^L = (-1)^N \mathbb{r}_0^L z_1$, $Z_2^L = (-1)^N \mathbb{r}_0^L z_2$,*

$$(6.17) \quad \frac{\prod_{v \in R_{Z_1}} (v+1)^{L-N} \prod_{u \in L_{Z_2}} (-u)^N}{\Delta(R_{Z_1}; L_{Z_2})} = e^{2B(z_1, z_2)} (1 + O(N^{\epsilon-1/2})),$$

where $B(z_1, z_2)$ is defined in (2.19).

Proof. The case when $Z_1 = Z_2$ was obtained in [4]: take the square of (8.17) in Lemma 8.2, whose proof was given in Section 9.2. The case of general Z_1 and Z_2 is almost the same, which we outline here. From the residue theorem,

$$(6.18) \quad \begin{aligned} & (L-N) \sum_{v \in R_{Z_1}} \log(v+1) + N \sum_{u \in L_{Z_2}} \log(-u) - \sum_{v \in R_{Z_1}} \sum_{u \in L_{Z_2}} \log(v-u) \\ &= Z_1^L Z_2^L \int_{-\rho_a - i\infty}^{-\rho_a + i\infty} \int_{-\rho_b - i\infty}^{-\rho_b + i\infty} \log \left(\frac{N^{1/2}(v-u)}{\rho \sqrt{1-\rho}} \right) \frac{L(v+\rho)}{v(v+1)q_{Z_1}(v)} \\ & \quad \times \frac{L(u+\rho)}{u(u+1)q_{Z_2}(u)} \frac{du}{2\pi i} \frac{dv}{2\pi i}, \end{aligned}$$

where the contours for u and v are two vertical lines $\Re(u) = -\rho_a := -\rho + \alpha \rho \sqrt{1-\rho} N^{-1/2}$ and $\Re(v) = -\rho_b := -\rho + \beta \rho \sqrt{1-\rho} N^{-1/2}$ with constants α and β satisfying $-\sqrt{-\log|z_2|} < \alpha < 0 < \beta < \sqrt{-\log|z_1|}$. This formula is similar to (9.26) in [4] and the proof is also similar. We divide the double integral (6.18) into two parts: $|u + \rho_a|, |v + \rho_a| \leq N^{\epsilon/3}$ and the rest. It is direct to check that the

formula (6.18) where the integral is restricted to $|u + \rho_a|, |v + \rho_a| \leq N^{\epsilon/3}$ is equal to

$$(6.19) \quad z_1 z_2 \int_{\Re \xi = a} \int_{\Re \zeta = b} \frac{\xi \zeta \log(\zeta - \xi)}{(e^{-\xi^2/2} - z_2)(e^{-\zeta^2/2} - z_1)} \frac{d\xi}{2\pi i} \frac{d\zeta}{2\pi i} \left(1 + O(N^{\epsilon-1/2})\right),$$

where we used the change of variables $u = -\rho + \xi \rho \sqrt{1-\rho} N^{-1/2}$ and $v = -\rho + \zeta \rho \sqrt{1-\rho} N^{-1/2}$. On the other hand, if $|u + \rho_a| \geq N^{\epsilon/3}$ or $|v + \rho_b| \geq N^{\epsilon/3}$, the integrand decays exponentially fast and hence the integral in these regions is exponentially small. See Section 9.2 of [4] for more discussions. From (2.19), we obtain (6.17). \square

Recalling the definitions (4.8) and (2.21), and using Lemma 6.2 and Lemma 6.3, we find that

$$(6.20) \quad C(\mathbf{Z}) = C(\mathbf{z}) (1 + O(N^{\epsilon-1/2})).$$

6.3. Analysis of $D(\mathbf{Z})$. We can obtain the limit of $D(\mathbf{Z})$ using either the Fredholm determinant or its series expansion (4.25). Both are suitable for the asymptotic analysis. Here we use the series expansion. From (4.25), we have

$$(6.21) \quad D(\mathbf{Z}) = \sum_{\mathbf{n} \in (\mathbb{Z}_{\geq 0})^m} \frac{1}{(\mathbf{n}!)^2} D_{\mathbf{n}}(\mathbf{Z})$$

with

$$(6.22) \quad D_{\mathbf{n}}(\mathbf{Z}) = (-1)^{|\mathbf{n}|} \sum_{\substack{U^{(\ell)} \in (L_{Z_\ell})^{n_\ell} \\ V^{(\ell)} \in (R_{Z_\ell})^{n_\ell} \\ l=1, \dots, m}} \det [K_1(w_i, w'_j)]_{i,j=1}^{|\mathbf{n}|} \det [K_2(w'_i, w_j)]_{i,j=1}^{|\mathbf{n}|},$$

where $U = (U^{(1)}, \dots, U^{(m)})$, $V = (V^{(1)}, \dots, V^{(m)})$ with $U^{(\ell)} = (u_1^{(\ell)}, \dots, u_{n_\ell}^{(\ell)})$, $V^{(\ell)} = (v_1^{(\ell)}, \dots, v_{n_\ell}^{(\ell)})$, and where

$$(6.23) \quad w_i = \begin{cases} u_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with odd integer } \ell, \\ v_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with even integer } \ell, \end{cases}$$

and

$$(6.24) \quad w'_i = \begin{cases} v_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with odd integer } \ell, \\ u_k^{(\ell)} & \text{if } i = n_1 + \dots + n_{\ell-1} + k \text{ for some } k \leq n_\ell \text{ with even integer } \ell. \end{cases}$$

We prove the convergence of this series to the series (2.51) with (2.52).

6.3.1. Strategy. To be able to cite easily later, let us state the following simple fact.

Lemma 6.4. *Suppose that*

- (A) *for each fixed \mathbf{n} , $D_{\mathbf{n}}(\mathbf{Z}) \rightarrow D_{\mathbf{n}}(\mathbf{z})$ as $L \rightarrow \infty$, and*
- (B) *there is a constant $C > 0$ such that $|D_{\mathbf{n}}(\mathbf{Z})| \leq C^{|\mathbf{n}|}$ for all \mathbf{n} and for all large enough L .*

Then $D(\mathbf{Z}) \rightarrow D(\mathbf{z})$ as $L \rightarrow \infty$.

Proof. It follows from the dominated convergence theorem. \square

We are going to show that conditions (A) and (B) are satisfied. To be precise, we need to show that (A) and (B) hold locally uniformly in \mathbf{z} so that the conclusion holds locally uniformly. The local uniformity is easy to check throughout the proof. To make the presentation light, we do not state the local uniformity explicitly and, instead, state only the pointwise convergence for each \mathbf{z} in the rest of this section.

Let us discuss the strategy of verifying the conditions (A) and (B). Suppose for a moment that $N/L = \rho$ is fixed. If $Z^L = (-1)^N \mathbf{r}_0^L z$ for a fixed z , then the contour $|w^N(w+1)|^{L-N} = |Z|^L$, on which the roots of $q_Z(w)$ lie, converges to the self-crossing contour $|w^\rho(w+1)^{1-\rho}| = \mathbf{r}_0$ as $L, N \rightarrow \infty$ since $|Z| \rightarrow \mathbf{r}_0$ (see Figure 6). The point of self-intersection is $w = -\rho$. For large L, N and the parameters satisfying the conditions of Theorem 2.1, it turns out that the main contribution to the sum $D_{\mathbf{n}}(\mathbf{Z})$ comes from the points $U^{(\ell)}$ and $V^{(\ell)}$ near the self-crossing point $w = -\rho$. As $L \rightarrow \infty$, $q_Z(w)$ has more and more roots. We scale the roots near the point $w = -\rho$ in such a way that the distances between the scaled roots are $O(1)$; this is achieved if⁸ we take $N^{1/2}(w+\rho) \mapsto w$. Under this scale, for each w on the set $L_Z \cup R_Z$ (which depends on L and N ; see (4.4)) in a neighborhood of the point $-\rho$, there is a unique point ζ on the set $L_z \cup R_z$ (which is independent of L and N ; see (2.28)), and vice versa. See Lemma 6.5 for the precise statement. We show that K_1 and K_2 converge to K_1 and K_2 pointwise for the points near $-\rho$. We then estimate the kernels when one of the arguments is away from $-\rho$. These two calculations are enough to prove conditions (A) and (B). See Lemma 6.6 for the precise statement. The fact that we only assume that $N/L = \rho$ is in a compact subset of $(0, 1)$ for all L does not change the analysis.

The following lemma is from [4].

Lemma 6.5 (Lemma 8.1 of [4]). *Let $0 < \epsilon < 1/2$. Fix a complex number z satisfying $0 < |z| < 1$ and let Z be a complex number satisfying $Z^L = (-1)^N \mathbf{r}_0^L z$. Recall the definitions of the sets L_Z and R_Z in (4.4), and the sets L_z and R_z in (2.28). Let $\mathcal{M}_{N,L}$ be the map from $L_Z \cap \{w : |w+\rho| \leq \rho\sqrt{1-\rho}N^{\epsilon/4-1/2}\}$ to L_z defined by*

$$(6.25) \quad \mathcal{M}_{N,L}(w) = \xi, \quad \text{where } \xi \in L_z \text{ and } \left| \xi - \frac{(w+\rho)N^{1/2}}{\rho\sqrt{1-\rho}} \right| \leq N^{3\epsilon/4-1/2} \log N.$$

Then for all large enough N the following hold:

- (a) $\mathcal{M}_{N,L}$ is well-defined.
- (b) $\mathcal{M}_{N,L}$ is injective.
- (c) Setting $L_z^{(c)} := L_z \cap \{\xi : |\xi| \leq c\}$ for $c > 0$, we have

$$(6.26) \quad L_z^{(N^{\epsilon/4}-1)} \subseteq I(\mathcal{M}_{N,L}) \subseteq L_z^{(N^{\epsilon/4}+1)},$$

where $I(\mathcal{M}_{N,L}) := \mathcal{M}_{N,L}(L_Z \cap \{w : |w+\rho| \leq \rho\sqrt{1-\rho}N^{\epsilon/4-1/2}\})$, the image of the map $\mathcal{M}_{N,L}$.

If we define the mapping $\mathcal{M}_{N,R}$ in the same way but replace L_Z and L_z by R_Z and R_z , respectively, the same results hold for $\mathcal{M}_{N,R}$.

Before we go further, we conjugate the kernels which leaves the determinants in (6.22) unchanged. We use the conjugated kernels in the rest of this subsection.

⁸The roots are less dense near the self-crossing point $w = -\rho$ than elsewhere. A typical distance between two neighboring roots is $O(N^{-1})$, but near $w = -\rho$ this distance is $O(N^{-1/2})$.

This makes the necessary convergences possible. For $w \in \mathcal{R}_{z_i} \cap S_1$ and $w' \in \mathcal{R}_{z_j} \cap S_2$, we change (4.18) to

$$(6.27) \quad \tilde{K}_1(w, w') = -(\delta_i(j) + \delta_i(j + (-1)^i)) \frac{J(w)\sqrt{f_i(w)}\sqrt{f_j(w')}(H_{z_i}(w))^2}{H_{z_{i-(-1)^i}}(w)H_{z_{j-(-1)^j}}(w')(w-w')} Q_1(j),$$

and for $w \in \mathcal{R}_{z_i} \cap S_2$ and $w' \in \mathcal{R}_{z_j} \cap S_1$, we change (4.19) to

$$(6.28) \quad \tilde{K}_2(w, w') = -(\delta_i(j) + \delta_i(j - (-1)^i)) \frac{J(w)\sqrt{f_i(w)}\sqrt{f_j(w')}(H_{z_i}(w))^2}{H_{z_{i+(-1)^i}}(w)H_{z_{j+(-1)^j}}(w')(w-w')} Q_2(j),$$

where we set (note the change from $f_j(w)$ in (4.11))

$$(6.29) \quad f_j(w) = \begin{cases} \frac{F_j(w)F_{j-1}(-\rho)}{F_{j-1}(w)F_j(-\rho)} & \text{for } \Re(w) < -\rho, \\ \frac{F_{j-1}(w)F_j(-\rho)}{F_j(w)F_{j-1}(-\rho)} & \text{for } \Re(w) > -\rho \end{cases}$$

with (see (4.10))

$$(6.30) \quad F_i(w) := w^{-k_i+N+1}(w+1)^{-a_i+k_i-N}e^{t_i w}.$$

We multiplied by -1 to remove a minus sign in the limit. The square-root function is defined as follows: for a complex number $w = re^{i\theta}$ with $r \geq 0$ and $-\pi < \theta \leq \pi$, we set $\sqrt{w} = r^{1/2}e^{i\theta/2}$. Note that \sqrt{w} is not continuous when w is a negative real number, and hence $\tilde{K}_1(w, w')$ and $\tilde{K}_2(w, w')$ may be discontinuous for some w and w' . However, the product of $\det[\tilde{K}_1(w_i, w'_j)]_{i,j=1}^{|\mathbf{n}|}$ and $\det[\tilde{K}_2(w'_i, w_j)]_{i,j=1}^{|\mathbf{n}|}$ is still continuous at the branch cuts since each of $\sqrt{f_i(w)}$ is multiplied twice. We also note that the change from f_j to f_j has the effect of conjugating the matrices in the determinants $\det[K_1(w_i, w'_j)]_{i,j=1}^{|\mathbf{n}|}$ and $\det[K_2(w'_i, w_j)]_{i,j=1}^{|\mathbf{n}|}$ and multiplying both by $(-1)^{|\mathbf{n}|}$. Hence $D_{\mathbf{n}}(\mathbf{Z})$ in (6.22) is unchanged if we replace K_1 and K_2 by \tilde{K}_1 and \tilde{K}_2 .

We also conjugate the limiting kernels. For $\zeta \in (\mathcal{L}_{z_i} \cup \mathcal{R}_{z_i}) \cap S_1$ and $\zeta' \in (\mathcal{L}_{z_j} \cup \mathcal{R}_{z_j}) \cap S_2$ for some i, j , we change (2.33) to

$$(6.31) \quad \tilde{K}_1(\zeta, \zeta') = (\delta_i(j) + \delta_i(j + (-1)^i)) \times \frac{\sqrt{f_i(\zeta)}\sqrt{f_j(\zeta')}e^{2h(\zeta, z_i) - h(\zeta, z_{i-(-1)^i}) - h(\zeta', z_{j-(-1)^j})}}{\zeta(\zeta - \zeta')} Q_1(j),$$

and for $\zeta \in (\mathcal{L}_{z_i} \cup \mathcal{R}_{z_i}) \cap S_2$ and $\zeta' \in (\mathcal{L}_{z_j} \cup \mathcal{R}_{z_j}) \cap S_1$, we change (2.35) to

$$(6.32) \quad \tilde{K}_2(\zeta, \zeta') = (\delta_i(j) + \delta_i(j - (-1)^i)) \times \frac{\sqrt{f_i(\zeta)}\sqrt{f_j(\zeta')}e^{2h(\zeta, z_i) - h(\zeta, z_{i+(-1)^i}) - h(\zeta', z_{j+(-1)^j})}}{\zeta(\zeta - \zeta')} Q_2(j).$$

Recall the definition of f_i in (2.27).

The next lemma shows how we prove the strategy mentioned above.

Lemma 6.6. *Fix $0 < \epsilon < 1/(1 + 2m)$. Let*

$$(6.33) \quad \Omega = \Omega_N := \left\{ w \in \mathbb{C} : |w + \rho| \leq \frac{\rho\sqrt{1 - \rho}}{N^{1/2 - \epsilon/4}} \right\}$$

be a disk centered at $-\rho$. Suppose the following:

(i) *We have*

$$(6.34) \quad \begin{aligned} |\tilde{K}_1(w, w')| &= |\tilde{K}_1(\zeta, \zeta')| + O(N^{\epsilon-1/2} \log N), \\ |\tilde{K}_2(w', w)| &= |\tilde{K}_2(\zeta', \zeta)| + O(N^{\epsilon-1/2} \log N) \end{aligned}$$

as $L \rightarrow \infty$, uniformly for $w \in S_1 \cap \Omega$ and $w' \in S_2 \cap \Omega$, where $\zeta \in S_1$ and $\zeta' \in S_2$ are the unique points corresponding to w and w' under either the map $\mathcal{M}_{N,L}$ or $\mathcal{M}_{N,R}$ in Lemma 6.5.

(ii) *For each \mathbf{n} ,*

$$\begin{aligned} \det \left[\tilde{K}_1(w_i, w'_j) \right]_{i,j=1}^{|\mathbf{n}|} &\rightarrow \det \left[\tilde{K}_1(\zeta_i, \zeta'_j) \right]_{i,j=1}^{|\mathbf{n}|}, \\ \det \left[\tilde{K}_2(w'_i, w_j) \right]_{i,j=1}^{|\mathbf{n}|} &\rightarrow \det \left[\tilde{K}_2(\zeta'_i, \zeta_j) \right]_{i,j=1}^{|\mathbf{n}|} \end{aligned}$$

as $L \rightarrow \infty$, where for $w_i \in S_1 \cap \Omega$ and $w'_i \in S_2 \cap \Omega$, $\zeta_i \in S_1$ and $\zeta'_i \in S_2$ are the unique points corresponding to w and w' under either the map $\mathcal{M}_{N,L}$ or $\mathcal{M}_{N,R}$ in Lemma 6.5.

(iii) *There are positive constants c and α such that*

$$|\tilde{K}_1(w, w')| = O(e^{-cN^\alpha}), \quad |\tilde{K}_2(w', w)| = O(e^{-cN^\alpha})$$

as $L \rightarrow \infty$, uniformly for $w \in S_1 \cap \Omega^c$ and $w' \in S_2$, and also for $w' \in S_2 \cap \Omega^c$ and $w \in S_1$.

Then conditions (A) and (B) in Lemma 6.4 hold, and therefore, $D(\mathbf{Z}) \rightarrow D(\mathbf{z})$.

If the absolute values in (i) are removed, then (i) will imply (ii). However, due to the discontinuity of the branch cuts of the square-root functions, $\tilde{K}_1(w, w')$ may converge to $-\tilde{K}_1(\zeta, \zeta')$ if the points are at the branch cuts. Nevertheless, the branch cuts do not affect the determinants as we discussed before. To emphasize this point, we state (ii) separately.

Proof. It is direct to check that due to the term f_i (see (2.27)) the kernels⁹ $\tilde{K}_1(\zeta, \zeta') = O(e^{-c_1|\zeta|^3})$ for some positive constant c_1 as $|\zeta| \rightarrow \infty$ along $\zeta \in S_1$ uniformly for $\zeta' \in S_2$, and also $\tilde{K}_1(\zeta, \zeta') = O(e^{-c_1|\zeta'|^3})$ as $|\zeta'| \rightarrow \infty$ along $\zeta' \in S_2$ uniformly for $\zeta \in S_1$. There are similar estimates for $\tilde{K}_2(\zeta', \zeta)$. This implies, in particular, that $\tilde{K}_1(\zeta, \zeta')$ and $\tilde{K}_2(\zeta', \zeta)$ are bounded for $\zeta \in S_1$ and $\zeta' \in S_2$.

Since S_1 and S_2 have $O(N^m)$ number of points, assumption (iii) implies that

$$(6.35) \quad \sum_{w' \in S_2} |\tilde{K}_1(w, w')|^2 = O(e^{-\frac{1}{2}cN^{\alpha\epsilon}}), \quad \sum_{w' \in S_2} |\tilde{K}_2(w', w)|^2 = O(e^{-\frac{1}{2}cN^{\alpha\epsilon}})$$

⁹If $\tau_i = \tau_{i+1}$ and $x_i < x_{i+1}$ for some i , then we have $O(e^{-c_1|\zeta|})$ and $O(e^{-c_1|\zeta'|})$, which are enough for the analysis.

uniformly for $w \in S_1 \cap \Omega^c$ and, similarly,

$$(6.36) \quad \sum_{w \in S_1} |\tilde{K}_1(w, w')|^2 = O(e^{-\frac{1}{2}cN^{\alpha\epsilon}}), \quad \sum_{w \in S_1} |\tilde{K}_2(w', w)|^2 = O(e^{-\frac{1}{2}cN^{\alpha\epsilon}})$$

uniformly for $w' \in S_2 \cap \Omega^c$.

We now show that there is a positive constant C_1 such that

$$(6.37) \quad \sum_{w \in S_1} \sqrt{\sum_{w' \in S_2} |\tilde{K}_1(w, w')|^2} \leq C_1$$

for all large enough L . The inequality is obtained if

$$\begin{aligned} & \sum_{w \in S_1 \cap \Omega^c} \sqrt{\sum_{w' \in S_2} |\tilde{K}_1(w, w')|^2}, \\ & \sum_{w \in S_1 \cap \Omega} \sqrt{\sum_{w' \in S_2 \cap \Omega^c} |\tilde{K}_1(w, w')|^2}, \\ & \sum_{w \in S_1 \cap \Omega} \sqrt{\sum_{w' \in S_2 \cap \Omega} |\tilde{K}_1(w, w')|^2} \end{aligned}$$

are all $O(1)$. The first two terms are bounded from assumption (iii) and the fact that S_1 and S_2 have $O(N^m)$ points. For the third term, we use assumption (i). It is direct to check that there are $O(N^{m\epsilon})$ number of points in $S_1 \cap \Omega$ and $S_2 \cap \Omega$. Hence the third term is bounded by

$$2 \sum_{\zeta \in S_1} \sqrt{\sum_{\zeta' \in S_2} |\tilde{K}_1(\zeta, \zeta')|^2} + O(N^{(m+1/2)\epsilon-1/2} \log N).$$

This is bounded since $|\tilde{K}_1(\zeta, \zeta')|$ decays fast as $|\zeta| \rightarrow \infty$ or $|\zeta'| \rightarrow \infty$ on S_1, S_2 and $\epsilon < 1/(1+2m)$. Hence we proved (6.37). Similarly, we have

$$(6.38) \quad \sum_{w' \in S_2} \sqrt{\sum_{w \in S_1} |\tilde{K}_2(w', w)|^2} \leq C_1.$$

We now show that (B) in Lemma 6.4 holds. Consider the formula (6.22) of $D_{\mathbf{n}}(\mathbf{Z})$. As we mentioned before, we change K_1 and K_2 to \tilde{K}_1 and \tilde{K}_2 without changing the determinants. From Hadamard's inequality, for all different w'_j ,

(6.39)

$$\left| \det \left[\tilde{K}_1(w_i, w'_j) \right]_{i,j=1}^{|\mathbf{n}|} \right| \leq \prod_{i=1}^{|\mathbf{n}|} \sqrt{\sum_{j=1}^{|\mathbf{n}|} |\tilde{K}_1(w_i, w'_j)|^2} \leq \prod_{i=1}^{|\mathbf{n}|} \sqrt{\sum_{w' \in S_2} |\tilde{K}_1(w_i, w')|^2}$$

and, similarly, for all different w_j ,

(6.40)

$$\left| \det \left[\tilde{K}_2(w'_i, w_j) \right]_{i,j=1}^{|\mathbf{n}|} \right| \leq \prod_{i=1}^{|\mathbf{n}|} \sqrt{\sum_{j=1}^{|\mathbf{n}|} |\tilde{K}_2(w'_i, w_j)|^2} \leq \prod_{i=1}^{|\mathbf{n}|} \sqrt{\sum_{w \in S_1} |\tilde{K}_2(w'_i, w)|^2}.$$

Hence,

$$\begin{aligned} |D_{\mathbf{n}}(\mathbf{Z})| &\leq \sum_{\substack{w_i \in S_1 \\ w'_i \in S_2 \\ i=1, \dots, |\mathbf{n}|}} \left(\prod_{i=1}^{|\mathbf{n}|} \sqrt{\sum_{w' \in S_2} |\tilde{K}_1(w_i, w')|^2} \right) \left(\prod_{i=1}^{|\mathbf{n}|} \sqrt{\sum_{w \in S_1} |\tilde{K}_2(w'_i, w)|^2} \right) \\ &= \left(\sum_{w \in S_1} \sqrt{\sum_{w' \in S_2} |\tilde{K}_1(w, w')|^2} \right)^{|\mathbf{n}|} \left(\sum_{w' \in S_2} \sqrt{\sum_{w \in S_1} |\tilde{K}_2(w', w)|^2} \right)^{|\mathbf{n}|}. \end{aligned}$$

Using (6.37) and (6.38), we obtain (B) with $C = C_1^2$.

We now prove that (A) in Lemma 6.4 holds. Fix \mathbf{n} . We divide the sum in the formula of $D_{\mathbf{n}}(\mathbf{Z})$ into two parts: the part that all $u_j^{(\ell)}, v_j^{(\ell)}$ are in Ω and the rest. By assumption (ii) and Lemma 6.5(c), the first part converges, as $L \rightarrow \infty$, to

$$(-1)^{|\mathbf{n}|} \sum_{\substack{\mathbf{U}^{(\ell)} \in (\mathbf{L}_{z_\ell})^{n_\ell} \\ \mathbf{V}^{(\ell)} \in (\mathbf{R}_{z_\ell})^{n_\ell} \\ \ell=1, \dots, m}} \det \left[\tilde{K}_1(\zeta_i, \zeta'_j) \right]_{i,j=1}^{|\mathbf{n}|} \det \left[\tilde{K}_2(\zeta'_i, \zeta_j) \right]_{i,j=1}^{|\mathbf{n}|}$$

which is equal to $D_{\mathbf{n}}(\mathbf{z})$ in (2.52). On the other hand, for the second part, note that (6.37) and (6.38) imply, in particular, that there is a positive constant C_2 such that

$$(6.41) \quad \sqrt{\sum_{w' \in S_2} |\tilde{K}_1(w, w')|^2} \leq C_2, \quad \sqrt{\sum_{w \in S_1} |\tilde{K}_2(w', w)|^2} \leq C_2$$

uniformly for $w \in S_1$ for the first inequality, and for $w' \in S_2$ for the second inequality. Now, by Hadamard's inequality (see (6.39) and (6.40)) and the estimates (6.35), (6.36), and (6.41), we find that for the second part,

$$(6.42) \quad \left| \det \left[\tilde{K}_1(w_i, w'_j) \right]_{i,j=1}^{|\mathbf{n}|} \det \left[\tilde{K}_2(w'_i, w_j) \right]_{i,j=1}^{|\mathbf{n}|} \right| \leq C_2^{2|\mathbf{n}|} C_3 e^{-cN^\alpha}$$

for a positive constant C_3 , since one of the variables is in Ω^c . Since there are only $O(N^m)$ points in S_1 and S_2 , we find that the second part converges to zero. Hence, we obtain (A). \square

6.3.2. Asymptotics of r_Z , l_Z , and f_j . In the remainder of this section, we verify assumptions (i), (ii), and (iii) of Lemma 6.6. The kernels contain $l_Z(w)$, $r_Z(w)$, and f_j . We first find the asymptotics of these functions.

The following asymptotic result was proved in [4].

Lemma 6.7 ([4]). *Let z be a complex number satisfying $0 < |z| < 1$. Let Z be a complex number such that $Z^L = (-1)^N \mathbf{r}_0^L z$. For a complex number $\hat{\omega}$, set*

$$(6.43) \quad w = -\rho + \frac{\rho\sqrt{1-\rho}}{N^{1/2}} \hat{\omega}.$$

There is a positive constant C such that the following hold:

(a) *If $\Re(\hat{\omega}) > c$ for some $c > 0$, then, uniformly in $\hat{\omega}$,*

$$(6.44) \quad l_Z(w) = e^{h(\hat{\omega}, z)} (1 + O(N^{\epsilon/2-1} \log N)) \quad \text{for } |\hat{\omega}| \leq N^{\epsilon/4}$$

and

$$(6.45) \quad e^{-CN^{-\epsilon/4}} \leq |l_Z(w)| \leq e^{CN^{-\epsilon/4}} \quad \text{for } |\hat{\omega}| \geq N^{\epsilon/4}.$$

(b) If $\Re(\hat{\omega}) < -c$ for some $c < 0$, then, uniformly in $\hat{\omega}$,

$$(6.46) \quad r_Z(w) = e^{h(\hat{\omega}, z)} (1 + O(N^{\epsilon/2-1} \log N)) \quad \text{for } |\hat{\omega}| \leq N^{\epsilon/4}$$

and

$$(6.47) \quad e^{-CN^{-\epsilon/4}} \leq |r_Z(w)| \leq e^{CN^{-\epsilon/4}} \quad \text{for } |\hat{\omega}| \geq N^{\epsilon/4}.$$

The errors are uniform for z in a compact subset of $0 < |z| < 1$. The function h is defined in (2.22) and (2.23).

Proof. The case when $|\hat{\omega}| \leq N^{\epsilon/4}$ is in Lemma 8.2(a) and (b) in [4], where we used the notation $q_{Z,L}(w) = (w+1)^{L-N} l_Z(w)$, $q_{Z,R}(w) = w^N r_Z(w)$, $h_L(\hat{\omega}, z) = h(\hat{\omega}, z)$ for $\hat{\omega} < 0$, and $h_R(\hat{\omega}, z) = h(\hat{\omega}, z)$ for $\hat{\omega} > 0$.

The upper bounds of $l_Z(w)$ and $r_Z(w)$ when $|\hat{\omega}| \geq N^{\epsilon/4}$ were computed in the proof of Lemma 8.4(c) of the same paper: (9.57) shows $|r_Z(w)| = \left| \frac{q_{Z,R}(w)}{w^N} \right| = O(e^{CN^{-\epsilon/4}})$. (There is a typo in that equation: the denominator on the left-hand side should be w^N instead of w^{L-N} .) This upper bound was obtained from an upper bound of $\log |r_Z(w)|$. Indeed, in the analysis between (9.52) and (9.57) in [4], we first write (there is another typo in (9.52): $(L-N) \log(-u)$ should be $N \log(-u)$)

$$\begin{aligned} \log |r_Z(w)| &= \Re \left[\sum_{v \in R_z} \log(-w+v) - N \log(-w) \right] \\ &= \Re \left[-LZ^L \int_{-\rho+i\mathbb{R}} \log \left(\frac{v-w}{-\rho-w} \right) \frac{(v+\rho)}{v(v+1)q_Z(v)} \frac{dv}{2\pi i} \right]. \end{aligned}$$

After estimating the integral in an elementary way, we obtain $|\log |r_Z(w)|| \leq CN^{-\epsilon/4}$. This implies both the upper and the lower bounds, $e^{-N^{-\epsilon/4}} \leq |r_Z(w)| \leq e^{N^{-\epsilon/4}}$. The estimate of $l_Z(w)$ is similar. \square

Now we consider $f_j(w)$. The following result applies to $F_j(w)$ (recall (6.30)), and hence f_j , for w near $-\rho$.

Lemma 6.8 ([30]). *Assume the same conditions for k, t, a as in Lemma 6.2. Set*

$$(6.48) \quad g(w) = w^{-k+N+1} (w+1)^{-a+k-N} e^{tw}.$$

Then, for

$$(6.49) \quad w = -\rho + \frac{\rho\sqrt{1-\rho}}{N^{1/2}} \hat{\omega}$$

with $\hat{\omega} = O(N^{\epsilon/4})$, we have

$$(6.50) \quad \frac{g(w)}{g(-\rho)} = e^{x\hat{\omega} + \frac{1}{2}\gamma\hat{\omega}^2 - \frac{1}{3}\tau\hat{\omega}^3} (1 + O(N^{\epsilon-1/2})).$$

Proof. If we set $k' = k - N + 1$ and $\ell' = a + 2$, then $g(w)$ is the same as $\tilde{g}_2(w)$ in (4.41) of [30] with k' and ℓ' instead of k and ℓ (and $\tau^{1/3}x$ replaced by x). Since $\ell' = a + 2 = \ell + 3$ satisfies condition (6.13), we see that $\frac{g(w)}{g(-\rho)}$ is equal to $g_2(w)$ in (4.40) of [30] with $j = 0$. The asymptotics of $g_2(w)$ was obtained in (4.46) of [30], which is the same as (6.50), under the conditions on t , k' , and ℓ' satisfying (4.1),

(4.2), and (4.3) of [30]. From (6.12) and (6.13), we find that these conditions are satisfied, and hence we obtain the lemma. \square

The next lemma is about $\mathbf{f}_j(w)$ when w is away from $-\rho$.

Lemma 6.9. *Suppose $x_j \in \mathbb{R}$, $\gamma_j \in [0, 1]$, and $\tau_j \in \mathbb{R}_{\geq 0}$ are fixed constants, and assume that¹⁰ $\tau_1 < \dots < \tau_m$. Let t_j , ℓ_j , and b_j satisfy (6.2), (6.3), and (6.4). Let*

$$(6.51) \quad a_j = \ell_j + 1, \quad k_j = N - \frac{b_j - \ell_j}{2} + 1$$

as defined in (6.7) and (6.8). Then, for $w \in \mathbf{L}_Z \cup \mathbf{R}_Z$ satisfying $|w + \rho| \geq \rho\sqrt{1 - \rho}N^{\epsilon/4 - 1/2}$,

$$(6.52) \quad \mathbf{f}_j(w) = O(e^{-cN^{3\epsilon/4}}).$$

The errors are uniform for z in a compact subset of $0 < |z| < 1$.

Proof. The proof is similar to the first part of the proof of Lemma 8.4(c) in [4], which is given in Section 9.4 of the same paper. We prove the case when $w \in \mathbf{L}_Z$. The case when $w \in \mathbf{R}_Z$ is similar. Recall from (6.29) and (6.30) that for $w \in \mathbf{L}_Z$, $\mathbf{f}_j(w) = \frac{F_j(w)F_{j-1}(-\rho)}{F_{j-1}(w)F_j(-\rho)}$ and

$$\frac{F_j(w)}{F_{j-1}(w)} = w^{k_{j-1} - k_j} (w + 1)^{-a_j + k_j + a_{j-1} - k_{j-1}} e^{(t_j - t_{j-1})w}.$$

We start with the following Claim. This is similar to a claim in Section 9.4 part (c).

Claim. *Suppose m , n , and ℓ are positive integers and $\rho \in (0, 1)$ satisfying*

$$(6.53) \quad -\frac{m}{\rho} + \frac{n}{1 - \rho} + \ell \geq 0.$$

Then the function

$$(6.54) \quad |w^m(w + 1)^n e^{\ell w}|$$

increases as $\Re(w)$ increases along any fixed contour

$$(6.55) \quad |w^\rho(w + 1)^{1-\rho}| = \text{const.}$$

If (6.53) is not satisfied, then (6.54) increases as $\Re(w)$ increases along the part of the contour (6.55) satisfying

$$(6.56) \quad |w + \rho|^2 \geq -\frac{\rho(1 - \rho)}{\ell} \left(-\frac{m}{\rho} + \frac{n}{1 - \rho} + \ell \right).$$

Proof of Claim. Set

$$c = -\frac{\rho n - (1 - \rho)m}{\rho \ell}.$$

We have $c \leq 1 - \rho$ from condition (6.53). Note that

$$|w^m(w + 1)^n e^{\ell w}| = \text{const} \cdot \left| (w + 1)^{n - \frac{1-\rho}{\rho}m} e^{\ell w} \right| = \text{const} \cdot |(w + 1)^{-c} e^w|^\ell$$

by using the condition that w is on the contour (6.55). It is direct to check by parameterizing the contour and taking the derivatives that the function $|(w + 1)^{-c} e^w|$ increases as $\Re(w)$ increases along the contour (6.55); see the Claim in Section 9.4

¹⁰If $\tau_{j-1} = \tau_j$ and $x_{j-1} < x_j$, then the errors change to $O(e^{-cN^{\epsilon/4}})$. In the proof, this change comes when we conclude the error term from (6.62).

part (c) of [4]. If (6.53) is not satisfied, then $c > 1 - \rho$, and in this case, again it is direct to check that the function $|(w + 1)^{-c}e^w|$ increases as $\Re(w)$ increases along the contour (6.55) if w is restricted to $|w + \rho|^2 \geq \rho(c - 1 + \rho)$. The last condition is (6.56). \square

We continue the proof of Lemma 6.9. We prove that the function $\left| \frac{F_j(w)}{F_{j-1}(w)} \right|$ increases as $\Re(w)$ increases along parts of the contour

$$(6.57) \quad |w^\rho(w + 1)^{1-\rho}| = \mathfrak{r}_0|z|^{1/L}$$

with the restriction

$$(6.58) \quad |w + \rho| \geq \rho\sqrt{1 - \rho}N^{\epsilon/4-1/2}.$$

Recall that $L_Z \cup R_Z$ is a discrete subset of the contour (6.57). By the Claim, it is sufficient to show that

$$(6.59) \quad \begin{aligned} & \left(\rho\sqrt{1 - \rho}N^{\epsilon/4-1/2} \right)^2 \\ & \geq -\frac{\rho(1 - \rho)}{t_j - t_{j-1}} \left(-\frac{-k_j + k_{j-1}}{\rho} + \frac{-a_j + k_j + a_{j-1} - k_{j-1}}{1 - \rho} + t_j - t_{j-1} \right). \end{aligned}$$

From (6.51), and (6.2), (6.3), and (6.4), the right-hand side is equal to

$$\begin{aligned} & -\frac{1}{2\rho(1 - \rho)(t_j - t_{j-1})} \\ & \times \left((b_j - b_{j-1}) - 2\rho(1 - \rho)(t_j - t_{j-1}) - (1 - 2\rho)(\ell_j - \ell_{j-1}) + O(t_j^{2/3}) \right). \end{aligned}$$

This is $O(N^{-1})$. Thus (6.59) holds for sufficiently large N , and we obtain the monotonicity of $\left| \frac{F_j(w)}{F_{j-1}(w)} \right|$.

The monotonicity implies that

$$(6.60) \quad \left| \frac{F_j(w)}{F_{j-1}(w)} \right| \leq \left| \frac{F_j(u_c)}{F_{j-1}(u_c)} \right|$$

with u_c on the contour satisfying

$$(6.61) \quad u_c = -\rho + \rho\sqrt{1 - \rho}\xi_c N^{-1/2}$$

for a complex number ξ_c satisfying $|\xi_c| = N^{\epsilon/4}$. Now by Lemma 6.8,

$$(6.62) \quad \begin{aligned} & \left| \frac{F_j(u_c)F_{j-1}(-\rho)}{F_{j-1}(u_c)F_j(-\rho)} \right| \\ & = \left| e^{(x_j - x_{j-1})\xi_c + \frac{1}{2}(\gamma_j - \gamma_{j-1})\xi_c^2 - \frac{1}{3}(\tau_j - \tau_{j-1})\xi_c^3} \right| \left(1 + O(N^{\epsilon-1/2}) \right). \end{aligned}$$

The point u_c is on the contour (6.57) and the contour is close to the contour $|w^\rho(w + 1)^{1-\rho}| = \mathfrak{r}_0$ exponentially, which is self-intersecting at $w = -\rho$ like an x. This implies that $\arg(\xi_c)$ converges to either $\frac{3\pi}{4}$ or $-\frac{3\pi}{4}$. Since $\tau_j - \tau_{j-1}$ is positive, we find that the right-hand side of (6.62) decays fast and is of order $e^{-cN^{3\epsilon/4}}$. This proves (6.52) when $w \in L_Z$. The case when $w \in R_Z$ is similar. \square

From Lemmas 6.8 and 6.9, we obtain the following asymptotics of $\mathbf{f}_j(w)$.

Lemma 6.10. *Recall $\mathbf{f}_j(w)$ introduced in (6.29) and $\mathbf{f}_j(w)$ defined in (2.27). Assume the same conditions on the parameters as in Lemma 6.9. Then¹¹ there is a positive constant c such that for $w \in \mathbf{L}_Z \cup \mathbf{R}_Z$,*

$$(6.63) \quad \mathbf{f}_j(w) = \begin{cases} \mathbf{f}_j(\hat{\omega})(1 + O(N^{\epsilon-1/2})) & \text{if } |\hat{\omega}| \leq N^{\epsilon/4}, \\ O(e^{-cN^{3\epsilon/4}}) & \text{if } |\hat{\omega}| \geq N^{\epsilon/4}, \end{cases}$$

where for given w , $\hat{\omega}$ is defined by the relation

$$(6.64) \quad w = -\rho + \frac{\rho\sqrt{1-\rho}}{N^{1/2}}\hat{\omega}.$$

In particular, $\mathbf{f}_j(w)$ is bounded uniformly for $w \in \mathbf{L}_Z \cup \mathbf{R}_Z$ as $N \rightarrow \infty$.

6.3.3. *Verification of conditions (i) and (ii) of Lemma 6.6.* Let $w \in \mathcal{R}_{Z_i} \cap \mathbf{S}_1$ and $w' \in \mathcal{R}_{Z_j} \cap \mathbf{S}_2$. Let ζ and ζ' be the image of w and w' under either the map $\mathcal{M}_{N,\mathbf{L}}$ or $\mathcal{M}_{N,\mathbf{R}}$ in Lemma 6.5, depending on whether the point is on \mathbf{L}_z or \mathbf{R}_z . We also set

$$(6.65) \quad \hat{\omega} := \frac{N^{1/2}}{\rho\sqrt{1-\rho}}(w + \rho), \quad \hat{\omega}' := \frac{N^{1/2}}{\rho\sqrt{1-\rho}}(w' + \rho).$$

Then, by Lemma 6.5,

$$(6.66) \quad |\hat{\omega} - \zeta|, \quad |\hat{\omega}' - \zeta'| \leq N^{3\epsilon/4-1/2} \log N.$$

We have

$$(6.67) \quad \begin{aligned} \tilde{K}_1(w, w') &= -(\delta_i(j) + \delta_i(j + (-1)^i)) \\ &\times \frac{J(w)\sqrt{f_i(w)}\sqrt{f_j(w')}(H_{Z_i}(w))^2}{H_{Z_{i-(-1)^i}}(w)H_{Z_{j-(-1)^j}}(w')(w - w')} Q_1(j). \end{aligned}$$

Clearly, $Q_1(j) = Q_1(j)$. Assume that $w, w' \in \Omega$. Then

$$(6.68) \quad J(w) = -\frac{\rho\sqrt{1-\rho}}{\hat{\omega}N^{1/2}}(1 + O(N^{\epsilon/4-1/2})).$$

The overall minus sign in $\tilde{K}_1(w, w')$ cancels the minus sign from $J(w)$. For other factors in (6.67), we use Lemma 6.7 and Lemma 6.10. Here, we recall from (4.14) that $H_z(w) = 1_z(w)$ for $\Re(w) < -\rho$ and $H_z(w) = r_z(w)$ for $\Re(w) > -\rho$. We obtain

$$(6.69) \quad |\tilde{K}_1(w, w')| = |\tilde{K}_1(\hat{\omega}, \hat{\omega}')|(1 + O(N^{\epsilon-1/2} \log N)).$$

We then take the approximate $\hat{\omega}$ and $\hat{\omega}'$ by ζ and ζ' using (6.66). Since the derivatives of $\tilde{K}_1(\hat{\omega}, \hat{\omega}')$ are bounded (which is straightforward to check), we find that

$$(6.70) \quad |\tilde{K}_1(w, w')| = |\tilde{K}_1(\zeta, \zeta')|(1 + O(N^{\epsilon-1/2} \log N)) + O(N^{3\epsilon/4-1/2} \log N).$$

Recall that $|\tilde{K}_1(\zeta, \zeta')|$ are bounded; see the first paragraph of the proof of Lemma 6.6. Therefore, we obtain the first equation of (6.34). The estimate of $|\tilde{K}_2(w, w')|$ is similar, and we obtain (i) of Lemma 6.6. Part (ii) of the lemma is similar.

¹¹If $\tau_{j-1} = \tau_j$ and $x_{j-1} < x_j$, then the error changes to $O(e^{-cN^{\epsilon/4}})$ for $|\hat{\omega}| \geq N^{\epsilon/4}$.

6.3.4. *Verification of condition (iii) of Lemma 6.6.* Consider $|\tilde{K}_1(w, w')|$ when w or w' is in Ω^c . Here $w \in S_1$ and $w' \in S_2$. We estimate each of the factors on the right-hand side of (6.67). We have the following. The estimates are all uniform in w or w' in the domain specified. The positive constants C , C' , and c may be different from line to line.

- (1) $N^{1/2}|w + \rho| \geq C$ for $w \in S_1 \cup S_2$. This follows from Lemma 6.5.
- (2) $N^{1/2}|w - w'| \geq C$ for $w \in S_1$ and $w' \in S_2$.
- (3) $|\mathbf{J}(w)| \leq CN^{-1/2}$ for $w \in S_1 \cup S_2$. Recall (4.12) for the definition of $\mathbf{J}(w)$.
- (4) $|\mathbf{f}_i(w)| \leq C$ for $w \in S_1 \cup S_2$ by Lemma 6.10.
- (5) $|\mathbf{f}_i(w)| \leq Ce^{-cN^{3\epsilon/4}}$ for $w \in \Omega^c \cap (S_1 \cup S_2)$ by Lemma 6.10.
- (6) $C \leq |H_{Z_i}(w)| \leq C'$ for $w \in S_1 \cup S_2$ by Lemma 6.7. For the upper bound, we also use the decay property of $\mathbf{h}(\hat{\omega}, z) = O(\hat{\omega}^{-1})$ from (2.26). Recall (4.14) for the definition of H_Z .
- (7) $|\mathbf{Q}_1(i)| \leq C$ and $|\mathbf{Q}_2(i)| \leq C$.

By combining these facts we obtain¹² that $|\tilde{K}_1(w, w')| = O(e^{-cN^{3\epsilon/4}})$ and $|\tilde{K}_2(w', w)| = O(e^{-cN^{3\epsilon/4}})$ if w or w' is in Ω^c . Hence we obtain (iii) of Lemma 6.6. This completes the proof of Theorem 2.1.

7. PROPERTIES OF THE LIMIT OF THE JOINT DISTRIBUTION

In this section, we discuss a few properties of the function

$$(7.1) \quad \mathbb{F}(x_1, \dots, x_m; p_1, \dots, p_m)$$

introduced in Section 2, which is a limit of the joint distribution. In order to emphasize that this is a function of m variables with m parameters, let us use the notation

$$(7.2) \quad \mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p}) = \mathbb{F}(x_1, \dots, x_m; p_1, \dots, p_m),$$

where

$$(7.3) \quad \mathbf{x} = (x_1, \dots, x_m), \quad \mathbf{p} = (p_1, \dots, p_m).$$

We also use the notation

$$(7.4) \quad \mathbf{x}^{[k]} = (x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_m), \quad \mathbf{p}^{[k]} = (p_1, \dots, p_{k-1}, p_{k+1}, \dots, p_m)$$

for vectors of size $m - 1$ with x_k and p_k removed, respectively.

Recall the formula

$$(7.5) \quad \mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p}) = \oint \dots \oint \mathbf{C}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) \mathbf{D}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) \frac{dz_m}{2\pi i z_m} \dots \frac{dz_1}{2\pi i z_1},$$

where $\mathbf{z} = (z_1, \dots, z_m)$, and the contours are nest circles satisfying

$$(7.6) \quad 0 < |z_m| < \dots < |z_1| < 1.$$

We wrote $\mathbf{C}(\mathbf{z})$ and $\mathbf{D}(\mathbf{z})$ by $\mathbf{C}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p})$ and $\mathbf{D}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p})$ to emphasize that they are functions of m variables z_1, \dots, z_m and depend on \mathbf{x} and \mathbf{p} . Recall from property (P1) in Subsubsection 2.2.1, which we proved in Subsubsection 2.2.2, that for each i , $\mathbf{C}^{(m)}(\mathbf{z})$ is a meromorphic function of z_i in the disk $|z_i| < 1$ and the only simple poles are $z_i = z_{i+1}$ for $i = 1, \dots, m-1$. Part (i) of the next lemma shows the analytic property of $\mathbf{D}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p})$ and proves property (P2) in Subsubsection 2.2.1. Part (ii) is used later.

¹²If $\tau_{j-1} = \tau_j$ and $x_{j-1} < x_j$, then the error terms in (5) and the kernels are $O(e^{-cN^{\epsilon/4}})$.

Lemma 7.1. (i) $D^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p})$ is analytic in each of the variables z_k in the deleted disk $0 < |z_k| < 1$.
(ii) For $1 \leq k \leq m-1$,

$$(7.7) \quad \lim_{z_k \rightarrow z_{k+1}} D^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) = D^{(m-1)}(\mathbf{z}^{[k]}; \mathbf{x}^{[k]}, \mathbf{p}^{[k]}),$$

where $\mathbf{z}^{[k]} = (z_1, \dots, z_{k-1}, z_{k+1}, \dots, z_m)$.

Proof. Recall the series formula (2.55):

$$(7.8) \quad D^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) = \sum_{\mathbf{n} \in (\mathbb{Z}_{\geq 0})^m} \frac{1}{(\mathbf{n}!)^2} D_{\mathbf{n}}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}), \quad D_{\mathbf{n}}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) = \sum \mathbf{d}_{\mathbf{n}, \mathbf{z}}^{(m)}(\mathbf{U}, \mathbf{V}; \mathbf{x}, \mathbf{p}),$$

where the last sum is over $\mathbf{U}^{(\ell)} \in (\mathcal{L}_{z_\ell})^{n_\ell}$ and $\mathbf{V}^{(\ell)} \in (\mathcal{R}_{z_\ell})^{n_\ell}$, $\ell = 1, \dots, m$. Here (recall the notational convention in Definition 2.9),

$$(7.9) \quad \begin{aligned} \mathbf{d}_{\mathbf{n}, \mathbf{z}}^{(m)}(\mathbf{U}, \mathbf{V}; \mathbf{x}, \mathbf{p}) &= \left[\prod_{\ell=1}^m \frac{\Delta(\mathbf{U}^{(\ell)})^2 \Delta(\mathbf{V}^{(\ell)})^2}{\Delta(\mathbf{U}^{(\ell)}; \mathbf{V}^{(\ell)})^2} \hat{f}_\ell(\mathbf{U}^{(\ell)}) \hat{f}_\ell(\mathbf{V}^{(\ell)}) \right] \\ &\times \left[\prod_{\ell=2}^m \frac{\Delta(\mathbf{U}^{(\ell)}; \mathbf{V}^{(\ell-1)}) \Delta(\mathbf{V}^{(\ell)}; \mathbf{U}^{(\ell-1)}) e^{-h(\mathbf{V}^{(\ell)}, z_{\ell-1}) - h(\mathbf{V}^{(\ell-1)}, z_\ell)}}{\Delta(\mathbf{U}^{(\ell)}; \mathbf{U}^{(\ell-1)}) \Delta(\mathbf{V}^{(\ell)}; \mathbf{V}^{(\ell-1)}) e^{h(\mathbf{U}^{(\ell)}, z_{\ell-1}) + h(\mathbf{U}^{(\ell-1)}, z_\ell)}} \right. \\ &\quad \left. \left(1 - \frac{z_{\ell-1}}{z_\ell}\right)^{n_\ell} \left(1 - \frac{z_\ell}{z_{\ell-1}}\right)^{n_{\ell-1}} \right], \end{aligned}$$

where $\mathbf{U} = (\mathbf{U}^{(1)}, \dots, \mathbf{U}^{(m)})$, $\mathbf{V} = (\mathbf{V}^{(1)}, \dots, \mathbf{V}^{(m)})$ with $\mathbf{U}^{(\ell)} = (u_1^{(\ell)}, \dots, u_{n_\ell}^{(\ell)})$, $\mathbf{V}^{(\ell)} = (v_1^{(\ell)}, \dots, v_{n_\ell}^{(\ell)})$. Note that we may take the components of $\mathbf{U}^{(\ell)}$ and $\mathbf{V}^{(\ell)}$ to be all distinct due to the factors $\Delta(\mathbf{U}^{(\ell)})$ and $\Delta(\mathbf{V}^{(\ell)})$. Recall that $\hat{f}_\ell(\zeta) := \frac{1}{\zeta} f_\ell(\zeta) e^{2h(\zeta, z_\ell)}$, where h and f_j are defined in (2.22), (2.23), and (2.27).

The points $u_j^{(\ell)}$ and $v_j^{(\ell)}$ are roots of the equation $e^{-\zeta^2/2} = z_\ell$, and hence they depend on z_ℓ analytically (if we order them properly). Note that the only denominators in (7.9) which can vanish are $\Delta(\mathbf{U}^{(\ell)}; \mathbf{U}^{(\ell-1)})$ and $\Delta(\mathbf{V}^{(\ell)}; \mathbf{V}^{(\ell-1)})$. They vanish only when $z_{\ell-1} = z_\ell$. Hence the only possible poles of $D^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p})$ are $z_k = z_{k+1}$ for $k = 1, \dots, m-1$. Hence (ii) implies (i).

We now prove (ii). We may assume that z_1, \dots, z_m are all distinct and take the limit as $z_k \rightarrow z_{k+1}$; otherwise, we may take successive limits. Let us consider which terms in (7.9) vanish when $z_k = z_{k+1}$. Clearly, $(1 - \frac{z_k}{z_{k+1}})^{n_{k+1}} (1 - \frac{z_{k+1}}{z_k})^{n_k}$ vanishes. On the other hand, when $z_k = z_{k+1}$, $\mathbf{U}^{(k)}$ and $\mathbf{U}^{(k+1)}$ are from the same set. If there is a non-zero overlap between the pair of vectors, then $\Delta(\mathbf{U}^{(k+1)}; \mathbf{U}^{(k)})$, which is in the denominator in (7.9), vanishes. Similarly, $\Delta(\mathbf{V}^{(k+1)}; \mathbf{V}^{(k)})$ may also vanish. Hence $\mathbf{d}_{\mathbf{n}, \mathbf{z}}^{(m)}(\mathbf{U}, \mathbf{V}; \mathbf{x}, \mathbf{p})$ is equal to

$$(7.10) \quad \frac{(z_{k+1} - z_k)^{n_{k+1} + n_k}}{\Delta(\mathbf{U}^{(k+1)}; \mathbf{U}^{(k)}) \Delta(\mathbf{V}^{(k+1)}; \mathbf{V}^{(k)})}$$

times a non-vanishing factor as $z_k \rightarrow z_{k+1}$. Now, note that if $u_i^{(k)} \rightarrow u_j^{(k+1)}$ for some i and j , then from $e^{-(u_j^{(k+1)})^2/2} = z_{k+1}$,

$$(7.11) \quad \lim_{u_i^{(k)} \rightarrow u_j^{(k+1)}} \frac{z_{k+1} - z_k}{u_j^{(k+1)} - u_i^{(k)}} = \frac{dz_{k+1}}{du_j^{(k+1)}} = -u_j^{(k+1)} z_{k+1}.$$

Since $\Delta(\mathbf{U}^{(k+1)}; \mathbf{U}^{(k)})$ has at most $\min\{n_{k+1}, n_k\}$ vanishing factors $u_j^{(k+1)} - u_i^{(k)}$ (because the components of $\mathbf{U}^{(k)}$ are all distinct, and so are $\mathbf{U}^{(k+1)}$) and, similarly, $\Delta(\mathbf{V}^{(k+1)}; \mathbf{V}^{(k)})$ has at most $\min\{n_{k+1}, n_k\}$ vanishing factors $v_j^{(k+1)} - v_i^{(k)}$, we find, by looking at the degree of the numerator, that (7.10) is non-zero only if (a) $n_k = n_{k+1}$, (b) $\mathbf{U}^{(k)}$ converges to a vector whose components are a permutation of the components of $\mathbf{U}^{(k+1)}$, and (c) $\mathbf{V}^{(k)}$ converges to a vector whose components are a permutation of the components of $\mathbf{V}^{(k+1)}$. When $n_k = n_{k+1}$, $\mathbf{U}^{(k)} \rightarrow \mathbf{U}^{(k+1)}$, and $\mathbf{V}^{(k)} \rightarrow \mathbf{V}^{(k+1)}$, the term (7.10) converges to

$$\frac{z_{k+1}^{2n_{k+1}} \prod_{i=1}^{n_{k+1}} u_i^{(k+1)} v_i^{(k+1)}}{\Delta(\mathbf{U}^{(k+1)})^2 \Delta(\mathbf{V}^{(k+1)})^2}.$$

Hence, using the fact that (see (2.27))

$$f_k(w) f_{k+1}(w) = \begin{cases} e^{-\frac{1}{3}(\tau_{k+1}-\tau_{k-1})\zeta^3 + \frac{1}{2}(\gamma_{k+1}-\gamma_{k-1})\zeta^2 + (x_{k+1}-x_{k-1})\zeta} & \text{for } \Re(w) < 0, \\ e^{\frac{1}{3}(\tau_{k+1}-\tau_{k-1})\zeta^3 - \frac{1}{2}(\gamma_{k+1}-\gamma_{k-1})\zeta^2 - (x_{k+1}-x_{k-1})\zeta} & \text{for } \Re(w) > 0, \end{cases}$$

we find that $d_{\mathbf{n}, \mathbf{z}}^{(m)}(\mathbf{U}, \mathbf{V}; \mathbf{x}, \mathbf{p})$ converges to

$$d_{\mathbf{n}^{[k]}, \mathbf{z}^{[k]}}^{(m-1)}(\mathbf{U}^{[k]}, \mathbf{V}^{[k]}; \mathbf{x}^{[k]}, \mathbf{p}^{[k]}),$$

where $\mathbf{n}^{[k]}, \mathbf{U}^{[k]}, \mathbf{V}^{[k]}$ are the same as $\mathbf{n}, \mathbf{U}, \mathbf{V}$ with $n_k, \mathbf{U}^{(k)}, \mathbf{V}^{(k)}$ removed. We obtain the same limit if $n_k = n_{k+1}$, $\mathbf{U}^{(k)}$ converges to a vector whose components are a permutation of the components of $\mathbf{U}^{(k+1)}$, and $\mathbf{V}^{(k)}$ converges to a vector whose components are a permutation of the components of $\mathbf{V}^{(k+1)}$. Hence,

$$\lim_{z_k \rightarrow z_{k+1}} D_{\mathbf{n}}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) = (n_k!)^2 \sum d_{\mathbf{n}^{[k]}, \mathbf{z}^{[k]}}^{(m-1)}(\mathbf{U}^{[k]}, \mathbf{V}^{[k]}; \mathbf{x}^{[k]}, \mathbf{p}^{[k]}),$$

where the sum is over $\mathbf{U}^{(\ell)} \in (\mathbf{L}_{z_\ell})^{n_\ell}$ and $\mathbf{V}^{(\ell)} \in (\mathbf{R}_{z_\ell})^{n_\ell}$, $\ell = 1, \dots, k-1, k+1, \dots, m$. This implies (7.7). This completes the proof of (ii), and hence (i), too. \square

The next lemma shows what happens if we interchange two consecutive contours in the formula (7.5) of $\mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p})$. Although we do not state it explicitly, we can also obtain an analogous formula for the finite-time joint distribution from a similar computation.

Lemma 7.2. *For every $1 \leq k \leq m-1$,*

$$(7.12) \quad \begin{aligned} \mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p}) &= \mathbb{F}^{(m-1)}(\mathbf{x}^{[k]}; \mathbf{p}^{[k]}) \\ &+ \oint \cdots \oint C^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) D^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) \frac{dz_m}{2\pi i z_m} \cdots \frac{dz_1}{2\pi i z_1}, \end{aligned}$$

where the contours are nested circles satisfying

$$(7.13) \quad 0 < |z_m| < \cdots < |z_{k+1}| < |z_{k-1}| < \cdots < |z_1| < 1 \text{ and } 0 < |z_k| < |z_{k+1}|.$$

Proof. We start with the formula (7.5). We fix all other contours and deform the z_k -contour so that $|z_k|$ is smaller than $|z_{k+1}|$. Then the integral (7.5) is equal to a term due to the residue plus the same integral with the contours changed to satisfy

$$0 < |z_m| < \cdots < |z_{k+2}| < |z_k| < |z_{k+1}| < |z_{k-1}| < \cdots < |z_1| < 1.$$

Since the integrand has poles only at $z_i = z_{i+1}$, $i = 1, \dots, m-1$, it is analytic at $z_k = z_i$ for $i \geq k+2$. Hence the conditions that $|z_k| > |z_i|$, $i \geq k+2$, are not necessary, and we can take the contours as (7.13).

It remains to show that the residue term is $\mathbb{F}^{(m-1)}(\mathbf{x}^{[k]}; \mathbf{p}^{[k]})$. It is direct to check from the definition (2.21) that

$$(7.14) \quad \lim_{z_k \rightarrow z_{k+1}} (z_k - z_{k+1}) C^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) = z_{k+1} C^{(m-1)}(\mathbf{z}^{[k]}; \mathbf{x}^{[k]}, \mathbf{p}^{[k]}),$$

where we set $\mathbf{z}^{[k]} = (z_1, \dots, z_{k-1}, z_{k+1}, \dots, z_m)$. On the other hand, from Lemma 7.1(ii), $D^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p})$ converges to $D^{(m)}(\mathbf{z}^{[k]}; \mathbf{x}^{[k]}, \mathbf{p}^{[k]})$. Thus, noting $\frac{dz_k}{2\pi i z_k}$ in (7.5), the residue term is equal to $\mathbb{F}^{(m-1)}(\mathbf{x}^{[k]}; \mathbf{p}^{[k]})$. \square

The multiple integral in (7.12) has a natural probabilistic interpretation. The following theorem gives an interpretation for more general choices of contours.

Theorem 7.3. *Assume the same conditions as Theorem 2.1. Let E_j^\pm be the events defined by*

$$(7.15) \quad E_j^- = \left\{ \frac{\ln(\mathbf{p}_j) - (1 - 2\rho)s_j - (1 - 2\rho + 2\rho^2)t_j}{-2\rho^{1/2}(1 - \rho)^{1/2}L^{1/2}} \leq x_j \right\}$$

and

$$(7.16) \quad E_j^+ = \left\{ \frac{\ln(\mathbf{p}_j) - (1 - 2\rho)s_j - (1 - 2\rho + 2\rho^2)t_j}{-2\rho^{1/2}(1 - \rho)^{1/2}L^{1/2}} > x_j \right\}.$$

Then

$$(7.17)$$

$$\lim_{L \rightarrow \infty} \mathbb{P}(E_1^\pm \cap \dots \cap E_{m-1}^\pm \cap E_m^-) = (-1)^\# \oint \dots \oint C(\mathbf{z}) D(\mathbf{z}) \frac{dz_m}{2\pi i z_m} \dots \frac{dz_1}{2\pi i z_1},$$

where $\#$ denotes the number of appearances of $+$ in $E_1^\pm \cap \dots \cap E_{m-1}^\pm$, and the contours are circles of radii between 0 and 1 such that for each $1 \leq j \leq m-1$,

$$(7.18) \quad |z_j| > |z_{j+1}| \text{ if we have } E_j^- \text{ and } |z_j| < |z_{j+1}| \text{ if we have } E_j^+.$$

Proof. Theorem 2.1 is the special case when we take E_j^- for all j . The general case follows from the same asymptotic analysis starting with a different finite-time formula. We first change Theorem 3.1. Let $\tilde{E}_j^+ := \{\mathbf{x}_{k_j}(t_j) \geq a_j\}$ and $\tilde{E}_j^- := \{\mathbf{x}_{k_j}(t_j) < a_j\}$ be events of TASEP in $\mathcal{X}_N(L)$. Then we claim that

$$(7.19) \quad \begin{aligned} & \mathbb{P}_Y \left(\tilde{E}_1^\pm \cap \dots \cap \tilde{E}_{m-1}^\pm \cap \tilde{E}_m^- \right) \\ &= (-1)^\# \oint \dots \oint C(\mathbf{z}, \mathbf{k}) D_Y(\mathbf{z}, \mathbf{k}, \tilde{\mathbf{a}}, \mathbf{t}) \frac{dz_m}{2\pi i z_m} \dots \frac{dz_1}{2\pi i z_1}, \end{aligned}$$

where the contours are circles of radii between 0 and 1 such that for each $1 \leq j \leq m-1$, $|z_j| > |z_{j+1}|$ if we have \tilde{E}_j^- and $|z_j| < |z_{j+1}|$ if we have \tilde{E}_j^+ . Here $\#$ denotes the number of appearances of $+$ in $\tilde{E}_1^\pm \cap \dots \cap \tilde{E}_{m-1}^\pm$ and $\tilde{\mathbf{a}} = (\tilde{a}_1, \dots, \tilde{a}_m)$ with $\tilde{a}_j = a_j$ or $a_j - 1$ depending on whether we have \tilde{E}_j^- or \tilde{E}_j^+ , respectively. The identity (7.19) follows from the same proof of Theorem 3.1 with a small change. The contour condition is used in the proof when we apply Proposition 3.4. Now, in Proposition 3.4, if we replace the assumption¹³ $\prod_{j=1}^N |w'_j + 1| < \prod_{j=1}^N |w_j + 1|$

¹³See the paragraph including the equation (3.22) for a discussion about how the condition $\prod_{j=1}^N |w'_j + 1| < \prod_{j=1}^N |w_j + 1|$ is related to the condition $|z'| < |z|$.

by $\prod_{j=1}^N |w'_j + 1| > \prod_{j=1}^N |w_j + 1|$, then the conclusion changes to

$$(7.20) \quad \sum_{X \in \mathcal{X}_N(L) \cap \{x_k < a\}} \mathcal{R}_X(W) \mathcal{L}_X(W') = - \left(\frac{z}{z'} \right)^{(k-1)L} \left(1 - \left(\frac{z'}{z} \right)^L \right)^{N-1} \times \left[\prod_{j=1}^N \frac{w_j^{-k} (w_j + 1)^{-a+k+2}}{(w'_j)^{-k} (w'_j + 1)^{-a+k+1}} \right] \det \left[\frac{1}{w_i - w'_{i'}} \right]_{i,i'=1}^N.$$

Note the change in the summation domain from $x_k \geq a$ to $x_k < a$. This identity follows easily from Proposition 5.2 and the geometric series formula. The probability of the event $\tilde{E}_1^\pm \cap \dots \cap \tilde{E}_{m-1}^\pm \cap \tilde{E}_m^-$ is then obtained from the same calculation as before using, for each i , either Proposition 3.4 or (7.20) depending on whether we have \tilde{E}_i^+ or \tilde{E}_i^- .

For the periodic step initial condition, Theorem 4.6 showed that $\mathcal{C}(\mathbf{z}, \mathbf{k})$ times $\mathcal{D}_Y(\mathbf{z}, \mathbf{k}, \tilde{\mathbf{a}}, \mathbf{t})$ is equal to $C(\mathbf{z})D(\mathbf{z})$. This proof does not depend on how the z_i -contours are nested. Hence we obtain

$$(7.21) \quad \mathbb{P} \left(\tilde{E}_1^\pm \cap \dots \cap \tilde{E}_{m-1}^\pm \cap \tilde{E}_m^- \right) = (-1)^\# \oint \dots \oint C(\mathbf{z}) D(\mathbf{z}) \frac{dz_m}{2\pi i z_m} \dots \frac{dz_1}{2\pi i z_1},$$

where $C(\mathbf{z})$ and $D(\mathbf{z})$ are the same as that in Theorem 4.6 except that the parameter \mathbf{a} is replaced by $\tilde{\mathbf{a}}$. The contours are the same as that in (7.19). Now, the asymptotic analysis follows from Section 6. Recall that the analysis of Section 6 does not depend on the ordering of $|z_i|$; see Remark 6.1. \square

We now prove the consistency of $\mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p})$ when one of the variables tends to positive infinity.

Proposition 7.4 (Consistency). *We have*

$$(7.22) \quad \lim_{x_k \rightarrow +\infty} \mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p}) = \mathbb{F}^{(m-1)}(\mathbf{x}^{[k]}; \mathbf{p}^{[k]}).$$

Proof. We consider the case when $k = m$ first and then the case when $k < m$.

(a) Assume $k = m$. When $m = 1$, we showed in Section 4 of [4] that $\mathbb{F}^{(1)}(x_1; p_1)$ is a distribution function. Hence,

$$(7.23) \quad \lim_{x_1 \rightarrow +\infty} \mathbb{F}^{(1)}(x_1; p_1) = 1.$$

Now we assume that $m \geq 2$ and take $x_m \rightarrow +\infty$. Recall the formula (7.5) in which the z_m -contour is the smallest contour. From the definition (see (2.21)),

$$\begin{aligned} \mathbb{C}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) &= \mathbb{C}^{(m-1)}(\mathbf{z}^{[m]}; \mathbf{x}^{[m]}, \mathbf{p}^{[m]}) \frac{z_{m-1}}{z_{m-1} - z_m} \\ &\times \frac{e^{x_m A_1(z_m) + \tau_m A_2(z_m)}}{e^{x_{m-1} A_1(z_m) + \tau_{m-1} A_2(z_m)}} e^{2B(z_m) - 2B(z_m, z_{m-1})}. \end{aligned}$$

We choose the contour for z_m given by $z_m = \frac{1}{x_m^2} e^{i\theta}$, $\theta \in [0, 2\pi]$. Since $A_i(z) = O(z)$, $B(z) = O(z)$, and $B(z, w) = O(z)$ as $z \rightarrow 0$, we have

$$(7.24) \quad \mathbb{C}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) = \mathbb{C}^{(m-1)}(\mathbf{z}^{[m]}; \mathbf{x}^{[m]}, \mathbf{p}^{[m]}) + O(x_m^{-1})$$

as $x_m \rightarrow \infty$.

We now show that

$$(7.25) \quad \mathbb{D}^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) = \mathbb{D}^{(m-1)}(\mathbf{z}^{[m]}; \mathbf{x}^{[m]}, \mathbf{p}^{[m]}) + O(e^{-cx_m})$$

as $x_m \rightarrow \infty$, where we had chosen $|z_m| = \frac{1}{x_m^2}$. From the series formula (2.51),

$$(7.26) \quad D^{(m)}(\mathbf{z}; \mathbf{x}, \mathbf{p}) - D^{(m-1)}(\mathbf{z}^{[m]}; \mathbf{x}^{[m]}, \mathbf{p}^{[m]}) = \sum_{n_m \geq 1} \sum_{n_1, \dots, n_{m-1} \geq 0} \frac{1}{(\mathbf{n}!)^2} D_{\mathbf{n}}(\mathbf{z}).$$

We need to show that the sum is exponentially small. From (2.55) and (2.56),

$$(7.27) \quad D_{\mathbf{n}}(\mathbf{z}) = \sum_{\substack{U^{(\ell)} \in (L_{z_\ell})^{n_\ell} \\ V^{(\ell)} \in (R_{z_\ell})^{n_\ell} \\ \ell=1, \dots, m-1}} \left[\sum_{\substack{U^{(m)} \in (L_{z_m})^{n_m} \\ V^{(m)} \in (R_{z_m})^{n_m}}} d_{\mathbf{n}, \mathbf{z}}(U, V) \right],$$

where for each U and V , $d_{\mathbf{n}, \mathbf{z}}(U, V)$ is equal to

$$(7.28) \quad \begin{aligned} & \frac{\Delta(U^{(m)})^2 \Delta(V^{(m)})^2}{\Delta(U^{(m)}; V^{(m)})^2} \hat{f}_m(U^{(m)}) \hat{f}_m(V^{(m)}) \\ & \times \frac{\Delta(U^{(m)}; V^{(m-1)}) \Delta(V^{(m)}; U^{(m-1)}) e^{-h(V^{(m)}, z_{m-1}) - h(V^{(m-1)}, z_m)}}{\Delta(U^{(m)}; U^{(m-1)}) \Delta(V^{(m)}; V^{(m-1)}) e^{h(U^{(m)}, z_{m-1}) + h(U^{(m-1)}, z_m)}} \\ & \times \left(1 - \frac{z_{m-1}}{z_m}\right)^{n_m} \left(1 - \frac{z_m}{z_{m-1}}\right)^{n_{m-1}} \end{aligned}$$

times

$$(7.29) \quad d_{\mathbf{n}^{[m]}, \mathbf{z}^{[m]}}(U^{[m]}, V^{[m]}),$$

a factor which does not depend on z_m , and hence also on $U^{(m)}$ and $V^{(m)}$ (and x_m and \mathbf{p}_m). The term (7.29) is the same as $d_{\mathbf{n}, \mathbf{z}}(U, V)$ when m is replaced by $m-1$. Since $n_m \geq 1$ in the sum (7.26), the inside sum in (7.27) is not over an empty set. We show that for each U and V , (7.28) is

$$(7.30) \quad O(e^{-cx_m|U^{(m)}|-cx_m|V^{(m)}|})$$

for a constant $c > 0$, where $|U^{(m)}|$ is the sum of the absolute values of $U^{(m)}$, and $|V^{(m)}|$ is similarly defined. This proves (7.25).

To show the decay of (7.28), we first note that every component u of $U^{(m)}$ is a solution of the equation $e^{-u^2/2} = z_m$ satisfying $\Re(u) < 0$. As $|z_m| = \frac{1}{x_m^2} \rightarrow 0$, $\Re(u^2) \rightarrow \infty$. Since $\Re(u^2) = (\Re(u))^2 - (\Im(u))^2$, this implies that $\Re(u) \rightarrow -\infty$, and hence $|u| \rightarrow \infty$. Similarly, every component v of $V^{(m)}$ satisfies $\Re(v) \rightarrow \infty$ and $|v| \rightarrow \infty$. It is also easy to check (see Figure 4) that the solutions of the equation $e^{-\zeta^2/2} = z$ lie in the sectors $-\pi/4 < \arg(\zeta) < \pi/4$ or $3\pi/4 < \arg(\zeta) < 5\pi/4$ for any $0 < |z| < 1$. Hence $|u| \leq \sqrt{2}\Re(u)$, $|v| \leq \sqrt{2}\Re(v)$, and $\sqrt{2}|u-v| \geq |u| + |v|$. We now consider each term in (7.28). Considering the degrees,

$$\frac{\Delta(U^{(m)})^2 \Delta(V^{(m)})^2}{\Delta(U^{(m)}; V^{(m)})^2} = O(1), \quad \frac{\Delta(U^{(m)}; V^{(m-1)}) \Delta(V^{(m)}; U^{(m-1)})}{\Delta(U^{(m)}; U^{(m-1)}) \Delta(V^{(m)}; V^{(m-1)})} = O(1).$$

From the formula of h in (2.25), and using (2.24),

$$h(V^{(m)}, z_{m-1}), h(U^{(m)}, z_{m-1}), h(V^{(m-1)}, z_m), h(U^{(m-1)}, z_m) = O(1).$$

Recall from (2.57) that $\hat{f}_m(\zeta) = \frac{1}{\zeta} f_\ell(\zeta) e^{2h(\zeta, z_m)}$. As above,

$$h(V^{(m)}, z_m), h(U^{(m)}, z_m) = O(1).$$

On the other hand, from the definition (2.27),

$$|\mathbf{f}_\ell(u)| \leq e^{-cx_m|u|}, \quad |\mathbf{f}_\ell(v)| \leq e^{-cx_m|v|}$$

for the components u and v of $\mathbf{U}^{(m)}$ and $\mathbf{V}^{(m)}$, implying that

$$|\hat{\mathbf{f}}_m(\mathbf{U}^{(m)})\hat{\mathbf{f}}_m(\mathbf{V}^{(m)})| = O(e^{-cx_m|\mathbf{U}^{(m)}|-cx_m|\mathbf{V}^{(m)}|}).$$

This term dominates the factor

$$\left(1 - \frac{z_{m-1}}{z_m}\right)^{n_m} = O(x_m^{2n_m}).$$

Combining together, we obtain the decay (7.30) of (7.28), and hence we obtain (7.25).

We insert (7.24) and (7.25) in (7.5), and integrating over θ , we obtain

$$\mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p}) = \mathbb{F}^{(m-1)}(\mathbf{x}^{[m]}, \mathbf{p}^{[m]}) + O(x_m^{-1})$$

for all large x_m . Hence we proved (7.22) for $k = m$.

(b) Assume that $k < m$. Let us denote the integral in (7.17) with the contours (7.18) by

$$\tilde{\mathbb{F}}^{(m)}(x_1^\pm, \dots, x_{m-1}^\pm, x_m^-; \mathbf{p}).$$

Note that

$$\tilde{\mathbb{F}}^{(m)}(x_1^-, \dots, x_{m-1}^-, x_m^-; \mathbf{p}) = \mathbb{F}^{(m)}(x_1, \dots, x_{m-1}, x_m; \mathbf{p}).$$

Fix k such that $1 \leq k \leq m-1$. By Lemma 7.2, we obtain (7.22) if we show that

$$(7.31) \quad \lim_{x_k \rightarrow +\infty} \tilde{\mathbb{F}}^{(m)}(x_1^-, \dots, x_{k-1}^-, x_k^+, x_{k+1}^-, \dots, x_m^-; \mathbf{p}) = 0.$$

Now, from the joint probability function interpretation stated in Theorem 7.3,

$$(7.32) \quad \tilde{\mathbb{F}}^{(m)}(\dots, x_{i-1}^\pm, x_i^\pm, x_{i+1}^\pm, \dots; \mathbf{p}) \leq \tilde{\mathbb{F}}^{(m-1)}(\dots, x_{i-1}^\pm, x_{i+1}^\pm, \dots; \mathbf{p}^{[i]})$$

for any $1 \leq i \leq m$ (for any choice of \pm -sign for x_i , $1 \leq i \leq m-1$, and the choice of $-$ sign for x_m). Using (7.32) $m-1$ times, we find that

$$(7.33) \quad \tilde{\mathbb{F}}^{(m)}(x_1^-, \dots, x_{k-1}^-, x_k^+, x_{k+1}^-, \dots, x_m^-; \mathbf{p}) \leq \tilde{\mathbb{F}}^{(1)}(x_k^+; p_k) = 1 - \mathbb{F}^{(1)}(x_k; p_k).$$

The one-point function $\mathbb{F}^{(1)}(x_k; p_k)$ converges to 1 as $x_k \rightarrow +\infty$ from (7.23). Hence we obtain (7.31), and this completes the proof of (7.22) when $1 \leq k \leq m-1$. \square

In the opposite direction, we have the following result.

Lemma 7.5. *We have*

$$(7.34) \quad \lim_{x_k \rightarrow -\infty} \mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p}) = 0.$$

Proof. Since a joint probability is smaller than a marginal distribution and $\mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p})$ is a limit of joint probabilities,

$$(7.35) \quad \mathbb{F}^{(m)}(\mathbf{x}; \mathbf{p}) \leq \mathbb{F}^{(1)}(x_k; p_k).$$

As mentioned before, the function $\mathbb{F}^{(1)}(x_k; p_k) = \mathbb{F}(x_k; p_k)$ is shown to be a distribution function in Section 4 of [4]. This implies the lemma. \square

8. INFINITE TASEP

If we take $L \rightarrow \infty$ while keeping all other parameters fixed, the periodic TASEP becomes the infinite TASEP with N particles. In terms of the joint distribution, this is still true if L is fixed but large enough.

Lemma 8.1. *Consider the infinite TASEP on \mathbb{Z} with N particles and let $\tilde{x}_i(t)$ denote the location of the i th particle (from left to right) at time t . Assume that the infinite TASEP has the initial condition given by $\tilde{x}_i(0) = y_i$, where $y_1 < \dots < y_N$. Also consider the TASEP in $\mathcal{X}_N(L)$ and denote by $\mathbf{x}_i(t)$ the location of the i th particle. Assume that*

$$(8.1) \quad L > y_N - y_1$$

and let $\mathbf{x}_i(t)$ have the same initial condition given by $\mathbf{x}_i(0) = y_i$. Fix a positive integer m . Let k_1, \dots, k_m be integers in $\{1, \dots, N\}$, let a_1, \dots, a_m be integers, and let t_1, \dots, t_m be positive real numbers. Then for any integer L satisfying (in addition to (8.1))

$$(8.2) \quad L > \max\{a_1 - k_1, \dots, a_m - k_m\} - y_1 + N + 1,$$

we have

$$(8.3) \quad \mathbb{P}(\tilde{x}_{k_1}(t_1) \leq a_1, \dots, \tilde{x}_{k_m}(t_m) \leq a_m) = \mathbb{P}(\mathbf{x}_{k_1}(t_1) \leq a_1, \dots, \mathbf{x}_{k_m}(t_m) \leq a_m).$$

*Proof.*¹⁴ We first observe that the particles $\mathbf{x}_i(t)$ are in the configuration space $\mathcal{X}_N(L)$, while the particles $\tilde{x}_i(t)$ are in the configuration space $W_N := \{(x_1, \dots, x_N) \in \mathbb{Z}^N : x_1 < \dots < x_N\}$. The only difference between these two configuration spaces is that $\mathcal{X}_N(L)$ has an extra restriction $x_N \leq x_1 + L - 1$. Therefore, if this restriction does not take an effect before time t , i.e., $\mathbf{x}_N(s) < \mathbf{x}_1(s) + L - 1$ for all $0 < s < t$, then the dynamics of the TASEP on $\mathcal{X}_N(L)$ is the same as that of the infinite TASEP (with the same initial condition) before time t . Furthermore, if we focus on the i th particle in the TASEP in $\mathcal{X}_N(L)$, there exists a smallest random time T_i such that the dynamics of this particle are the same in both the TASEP $\mathcal{X}_N(L)$ and the infinite TASEP before time T_i . The times T_i are determined inductively as follows. First, T_N is the smallest time such that $\mathbf{x}_N(t) = \mathbf{x}_1(t) + L - 1$. Next, T_{N-1} is the smallest time that satisfies $t \geq T_N$ and $\mathbf{x}_{N-1}(t) = \mathbf{x}_N(t) - 1$. For general index $1 \leq i \leq N-1$, T_i is the smallest time that satisfies $t \geq T_{i+1}$ and $\mathbf{x}_i(t) = \mathbf{x}_{i+1}(t) - 1$. Note that $T_1 \geq T_2 \geq \dots \geq T_N$ and for $1 \leq i \leq N-1$,

$$(8.4) \quad \mathbf{x}_i(T_i) = \mathbf{x}_{i+1}(T_i) - 1 \geq \mathbf{x}_{i+1}(T_{i+1}) - 1.$$

The same consideration shows that if we consider m particles $\mathbf{x}_{k_1}(t_1), \dots, \mathbf{x}_{k_m}(t_m)$ of the TASEP in $\mathcal{X}_N(L)$ at possibly different times, their joint distribution is the same as that of the infinite TASEP if $t_i < T_{k_i}$ for all i . Therefore, we obtain (8.3) if we show that under the condition (8.2), the event that $\mathbf{x}_{k_i}(t_i) \leq a_i$ for all $1 \leq i \leq m$ is a subset of the event that $t_i < T_{k_i}$ for all $1 \leq i \leq m$. Now, suppose that $t_i \geq T_{k_i}$ for some i . Then, writing $\ell = k_i$ and using (8.4),

$$\mathbf{x}_\ell(t_i) \geq \mathbf{x}_\ell(T_\ell) \geq \mathbf{x}_{\ell+1}(T_{\ell+1}) - 1 \geq \dots \geq \mathbf{x}_N(T_N) - (N - \ell).$$

¹⁴This lemma can be seen easily from the directed last passage percolation interpretation of the TASEP.

Since $x_N(\tau_N) = \mathbf{x}_1(T_N) + L - 1$ and $\mathbf{x}_1(T_N) \geq \mathbf{x}_1(0) = y_1$, this implies that (recall that $\ell = k_i$)

$$\mathbf{x}_\ell(t_i) \geq y_1 + L - 1 - (N - \ell) > a_\ell$$

using the condition (8.2). Hence we are not in the event that $\mathbf{x}_{k_i}(t_i) \leq a_i$ for all $1 \leq i \leq m$. This completes the proof. \square

The above result implies, using the inclusion-exclusion principle,

$$(8.5) \quad \mathbb{P}(\tilde{x}_{k_1}(t_1) \geq a_1, \dots, \tilde{x}_{k_m}(t_m) \geq a_m) = \mathbb{P}(\mathbf{x}_{k_1}(t_1) \geq a_1, \dots, \mathbf{x}_{k_m}(t_m) \geq a_m)$$

for L satisfying

$$(8.6) \quad L > \max\{a_1 - k_1, \dots, a_m - k_m, y_N - N\} - y_1 + N.$$

Therefore, Theorem 3.1 implies that

$$(8.7) \quad \mathbb{P}(\tilde{x}_{k_1}(t_1) \geq a_1, \dots, \tilde{x}_{k_m}(t_m) \geq a_m) = \text{the right-hand side of (3.6)}$$

for any L satisfying (8.6). In particular, for the initial condition $\tilde{x}_i(0) = i - N$, $i = 1, \dots, N$, by Theorem 4.6, we find that

$$(8.8) \quad \mathbb{P}(\tilde{x}_{k_1}(t_1) \geq a_1, \dots, \tilde{x}_{k_m}(t_m) \geq a_m) = \text{the right-hand side of (4.30)}$$

for any integer L satisfying

$$(8.9) \quad L \geq 2N + \max\{a_1 - k_1, \dots, a_m - k_m, -N\}.$$

Note that since the particles move only to the right, the above joint probability is the same as that of the infinite TASEP (with infinitely many particles) with the step initial condition. Hence we obtained a formula for the finite-time joint distribution in multiple times and locations of the infinite TASEP with the step initial condition. Actually we have infinitely many formulas, one for each L satisfying (8.9). Since the infinite TASEP does not involve the parameter L , all these formulas should give an equal value for all L satisfying (8.9).

Now, if we want to compute the large-time limit of the joint distribution of the infinite TASEP under the KPZ scaling, we need to take $a_i = O(t)$. The above restriction on L implies that $L \geq O(t)$. This implies that $t \ll L^{3/2}$, which corresponds to the *subrelaxation* time scale. Hence the large-time limit of the joint distribution of the infinite TASEP is equal to the large-time limit, if it exists, of the joint distribution of the periodic TASEP in the subrelaxation time scale. However, it is not immediately clear if the formula (4.30) is suitable for the subrelaxation time scale when $m \geq 2$. In particular, the kernels $K_1(w, w')$ and $K_2(w, w')$ do not seem to converge. We leave the analysis of the multipoint distribution of the infinite TASEP as a future project.

REFERENCES

- [1] Gideon Amir, Ivan Corwin, and Jeremy Quastel, *Probability distribution of the free energy of the continuum directed random polymer in 1 + 1 dimensions*, Comm. Pure Appl. Math. **64** (2011), no. 4, 466–537, DOI 10.1002/cpa.20347. MR2796514
- [2] Jinho Baik, Percy Deift, and Kurt Johansson, *On the distribution of the length of the longest increasing subsequence of random permutations*, J. Amer. Math. Soc. **12** (1999), no. 4, 1119–1178, DOI 10.1090/S0894-0347-99-00307-0. MR1682248
- [3] Jinho Baik, Patrik L. Ferrari, and Sandrine Péché, *Limit process of stationary TASEP near the characteristic line*, Comm. Pure Appl. Math. **63** (2010), no. 8, 1017–1070, DOI 10.1002/cpa.20316. MR2642384

- [4] Jinho Baik and Zhipeng Liu, *Fluctuations of TASEP on a ring in relaxation time scale*, Comm. Pure Appl. Math. **71** (2018), no. 4, 747–813, DOI 10.1002/cpa.21702. MR3772401
- [5] Jinho Baik and Zhipeng Liu, *TASEP on a ring in sub-relaxation time scale*, J. Stat. Phys. **165** (2016), no. 6, 1051–1085, DOI 10.1007/s10955-016-1665-y. MR3575637
- [6] Alexei Borodin, Ivan Corwin, and Patrik Ferrari, *Free energy fluctuations for directed polymers in random media in $1 + 1$ dimension*, Comm. Pure Appl. Math. **67** (2014), no. 7, 1129–1214, DOI 10.1002/cpa.21520. MR3207195
- [7] Alexei Borodin and Patrik L. Ferrari, *Large time asymptotics of growth models on space-like paths. I. PushASEP*, Electron. J. Probab. **13** (2008), no. 50, 1380–1418, DOI 10.1214/EJP.v13-541. MR2438811
- [8] Alexei Borodin, Patrik L. Ferrari, and Michael Prähofer, *Fluctuations in the discrete TASEP with periodic initial configurations and the Airy₁ process*, Int. Math. Res. Pap. IMRP **1** (2007), Art. ID rpm002, 47. MR2334008
- [9] Alexei Borodin, Patrik L. Ferrari, Michael Prähofer, and Tomohiro Sasamoto, *Fluctuation properties of the TASEP with periodic initial configuration*, J. Stat. Phys. **129** (2007), no. 5–6, 1055–1080, DOI 10.1007/s10955-007-9383-0. MR2363389
- [10] Alexei Borodin, Patrik L. Ferrari, and Tomohiro Sasamoto, *Transition between Airy₁ and Airy₂ processes and TASEP fluctuations*, Comm. Pure Appl. Math. **61** (2008), no. 11, 1603–1629, DOI 10.1002/cpa.20234. MR2444377
- [11] Ivan Corwin, *The Kardar-Parisi-Zhang equation and universality class*, Random Matrices Theory Appl. **1** (2012), no. 1, 1130001, 76, DOI 10.1142/S2010326311300014. MR2930377
- [12] Ivan Corwin, Patrik L. Ferrari, and Sandrine Péché, *Limit processes for TASEP with shocks and rarefaction fans*, J. Stat. Phys. **140** (2010), no. 2, 232–267, DOI 10.1007/s10955-010-9995-7. MR2659279
- [13] Ivan Corwin, Zhipeng Liu, and Dong Wang, *Fluctuations of TASEP and LPP with general initial data*, Ann. Appl. Probab. **26** (2016), no. 4, 2030–2082, DOI 10.1214/15-AAP1139. MR354389
- [14] Jacopo de Nardis and Pierre Le Doussal, *Tail of the two-time height distribution for KPZ growth in one dimension*, J. Stat. Mech. Theory Exp. **5** (2017), 053212, 72, DOI 10.1088/1742-5468/aa6bce. MR3664392
- [15] Bernard Derrida and Joel L. Lebowitz, *Exact large deviation function in the asymmetric exclusion process*, Phys. Rev. Lett. **80** (1998), no. 2, 209–213, DOI 10.1103/PhysRevLett.80.209. MR1604439
- [16] Victor Dotsenko, *Two-time free energy distribution function in $(1 + 1)$ directed polymers*, J. Stat. Mech. Theory Exp. **6** (2013), P06017, 23, DOI 10.1088/1742-5468/2013/06/p06017. MR3090739
- [17] Victor Dotsenko, *Two-time distribution function in one-dimensional random directed polymers*, J. Phys. A **48** (2015), no. 49, 495001, 18, DOI 10.1088/1751-8113/48/49/495001. MR3434822
- [18] Victor Dotsenko, *On two-time distribution functions in $(1 + 1)$ random directed polymers*, J. Phys. A **49** (2016), no. 27, 27LT01, 8, DOI 10.1088/1751-8113/49/27/27LT01. MR3512099
- [19] Patrik L. Ferrari and Peter Nejjar, *Anomalous shock fluctuations in TASEP and last passage percolation models*, Probab. Theory Related Fields **161** (2015), no. 1–2, 61–109, DOI 10.1007/s00440-013-0544-6. MR3304747
- [20] Patrik L. Ferrari and Herbert Spohn, *On time correlations for KPZ growth in one dimension*, SIGMA Symmetry Integrability Geom. Methods Appl. **12** (2016), Paper No. 074, 23, DOI 10.3842/SIGMA.2016.074. MR3529743
- [21] O. Golinelli and K. Mallick, *Bethe ansatz calculation of the spectral gap of the asymmetric exclusion process*, J. Phys. A **37** (2004), no. 10, 3321–3331, DOI 10.1088/0305-4470/37/10/001. MR2039850
- [22] O. Golinelli and K. Mallick, *Spectral gap of the totally asymmetric exclusion process at arbitrary filling*, J. Phys. A **38** (2005), no. 7, 1419–1425, DOI 10.1088/0305-4470/38/7/001. MR2117259
- [23] Shamik Gupta, Satya N. Majumdar, Claude Godrèche, and Mustansir Barma, *Tagged particle correlations in the asymmetric simple exclusion process: finite-size effects*, Phys. Rev. E (3) **76** (2007), no. 2, 021112, 17, DOI 10.1103/PhysRevE.76.021112. MR2365530
- [24] L.-H. Gwa and H. Spohn, *Bethe solution for the dynamical-scaling exponent of the noisy Burgers equation*, Phys. Rev. A **46** (1992), 844–854.

- [25] T. Imamura and T. Sasamoto, *Fluctuations of the one-dimensional polynuclear growth model with external sources*, Nuclear Phys. B **699** (2004), no. 3, 503–544, DOI 10.1016/j.nuclphysb.2004.07.030. MR2098552
- [26] Kurt Johansson, *Shape fluctuations and random matrices*, Comm. Math. Phys. **209** (2000), no. 2, 437–476, DOI 10.1007/s002200050027. MR1737991
- [27] Kurt Johansson, *Discrete polynuclear growth and determinantal processes*, Comm. Math. Phys. **242** (2003), no. 1-2, 277–329, DOI 10.1007/s00220-003-0945-y. MR2018275
- [28] Kurt Johansson, *Two time distribution in Brownian directed percolation*, Comm. Math. Phys. **351** (2017), no. 2, 441–492, DOI 10.1007/s00220-016-2660-5. MR3613511
- [29] K. Johansson, *The two-time distribution in geometric last-passage percolation*, 2018, arXiv:1802.00729.
- [30] Zhipeng Liu, *Height fluctuations of stationary TASEP on a ring in relaxation time scale* (English, with English and French summaries), Ann. Inst. Henri Poincaré Probab. Stat. **54** (2018), no. 2, 1031–1057, DOI 10.1214/17-AIHP831. MR3795076
- [31] K. Matetski, J. Quastel, and D. Remenik, *The KPZ fixed point*, arXiv:1701.00018.
- [32] V. S. Poghosyan and V. B. Priezzhev, *Determinant solution for the TASEP with particle-dependent hopping probabilities on a ring*, Markov Process. Related Fields **14** (2008), no. 2, 233–254. MR2437530
- [33] A. M. Povolotsky and V. B. Priezzhev, *Determinant solution for the totally asymmetric exclusion process with parallel update. II. Ring geometry*, J. Stat. Mech. Theory Exp. **8** (2007), P08018, 27, DOI 10.1088/1742-5468/2007/08/p08018. MR2338272
- [34] Michael Prähofer and Herbert Spohn, *Scale invariance of the PNG droplet and the Airy process*, J. Statist. Phys. **108** (2002), no. 5-6, 1071–1106, DOI 10.1023/A:1019791415147. Dedicated to David Ruelle and Yasha Sinai on the occasion of their 65th birthdays. MR1933446
- [35] V. Priezzhev, *Exact nonstationary probabilities in the asymmetric exclusion process on a ring*, Phys. Rev. Lett. **91** (2003), no. 5, 050601.
- [36] S. Prolhac, *Finite-time fluctuations for the totally asymmetric exclusion process*, Phys. Rev. Lett. **116** (2016), 090601.
- [37] J. Quastel and D. Remenik, *How flat is flat in random interface growth?* arXiv:1606.09228.
- [38] A. Rákos and G. M. Schütz, *Current distribution and random matrix ensembles for an integrable asymmetric fragmentation process*, J. Stat. Phys. **118** (2005), no. 3-4, 511–530, DOI 10.1007/s10955-004-8819-z. MR2123646
- [39] T. Sasamoto, *Spatial correlations of the 1D KPZ surface on a flat substrate*, J. Phys. A **38** (2005), no. 33, L549–L556, DOI 10.1088/0305-4470/38/33/L01. MR2165697
- [40] Gunter M. Schütz, *Exact solution of the master equation for the asymmetric exclusion process*, J. Statist. Phys. **88** (1997), no. 1-2, 427–445, DOI 10.1007/BF02508478. MR1468391
- [41] Craig A. Tracy and Harold Widom, *Asymptotics in ASEP with step initial condition*, Comm. Math. Phys. **290** (2009), no. 1, 129–154, DOI 10.1007/s00220-009-0761-0. MR2520510

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN 48109
Email address: `baik@umich.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, LAWRENCE, KANSAS 66045
Email address: `zhipeng@ku.edu`