



A Paradiifferential Approach for Well-Posedness of the Muskat Problem

HUY Q. NGUYEN & BENOÎT PAUSADER

Communicated by P. CONSTANTIN

Abstract

We study the Muskat problem for one fluid or two fluids, with or without viscosity jump, with or without rigid boundaries, and in arbitrary space dimension d of the interface. The Muskat problem is scaling invariant in the Sobolev space $H^{s_c}(\mathbb{R}^d)$ where $s_c = 1 + \frac{d}{2}$. Employing a paradiifferential approach, we prove local well-posedness for large data in any subcritical Sobolev spaces $H^s(\mathbb{R}^d)$, $s > s_c$. Moreover, the rigid boundaries are only required to be Lipschitz and can have arbitrarily large variation. The Rayleigh–Taylor stability condition is assumed for the case of two fluids with viscosity jump but is proved to be automatically satisfied for the case of one fluid. The starting point of this work is a reformulation solely in terms of the Dirichlet–Neumann operator. The key elements of proofs are new paralinearization and contraction results for the Dirichlet–Neumann operator in rough domains.

1. Introduction

1.1. The Muskat problem

In its full generality, the Muskat problem describes the dynamics of two immiscible fluids in a porous medium with different densities ρ^\pm and different viscosities μ^\pm . Let us denote the interface between the two fluids by Σ and assume that it is the graph of a time-dependent function $\eta(x, t)$, that is

$$\Sigma_t = \left\{ (x, \eta(t, x)) : x \in \mathbb{R}^d \right\}. \quad (1.1)$$

The associated time-dependent fluid domains are then given by


$$\Omega_t^+ = \left\{ (x, y) \in \mathbb{R}^d \times \mathbb{R} : \eta(t, x) < y < \underline{b}^+(x) \right\} \quad (1.2)$$

and

$$\Omega_t^- = \{(x, y) \in \mathbb{R}^d \times \mathbb{R} : \underline{b}^-(x) < y < \eta(t, x)\} \quad (1.3)$$

where \underline{b}^\pm are the parametrizations of the rigid boundaries

$$\Gamma^\pm = \{(x, \underline{b}^\pm(x)) : x \in \mathbb{R}^d\}. \quad (1.4)$$



(a) The one-phase problem

(b) The two-phase problem

The incompressible fluid velocity u^\pm in each region is governed by Darcy's law:

$$\mu^\pm u^\pm + \nabla_{x,y} p^\pm = -(0, \rho^\pm) \quad \text{in } \Omega_t^\pm, \quad (1.5)$$

and

$$\operatorname{div}_{x,y} u^\pm = 0 \quad \text{in } \Omega_t^\pm. \quad (1.6)$$

Note that we have normalized gravity to 1 in (1.5).

At the interface Σ , the normal velocity is continuous:

$$u^+ \cdot n = u^- \cdot n \quad \text{on } \Sigma_t, \quad (1.7)$$

where $n = \frac{1}{\sqrt{1+|\nabla \eta|^2}}(-\nabla \eta, 1)$ is the upward pointing unit normal to Σ_t . Then, the interface moves with the fluid:

$$\partial_t \eta = \sqrt{1+|\nabla \eta|^2} u^- \cdot n|_{\Sigma_t}. \quad (1.8)$$

By neglecting the effect of surface tension, the pressure is continuous at the interface:

$$p^+ = p^- \quad \text{on } \Sigma_t. \quad (1.9)$$

Finally, at the two rigid boundaries, the no-penetration boundary conditions are imposed:

$$u^\pm \cdot v^\pm = 0 \quad \text{on } \Gamma^\pm, \quad (1.10)$$

where $v^\pm = \pm \frac{1}{\sqrt{1+|\nabla \underline{b}^\pm|^2}}(-\nabla \underline{b}^\pm, 1)$ denotes the outward pointing unit normal to Γ^\pm . We will also consider the case that at least one of Γ^\pm is empty (infinite depth); (1.10) is then replaced by the vanishing of u at infinity.

We shall refer to the system (1.2)–(1.10) as the two-phase Muskat problem. When the top phase corresponds to vacuum, that is $\mu^+ = \rho^+ = 0$, the two-phase Muskat problem reduces to the one-phase Muskat problem and (1.9) becomes

$$p^- = 0 \quad \text{on } \Sigma_t. \quad (1.11)$$

1.2. Presentation of the Main Results

It turns out that the Muskat problem can be recast as a *quasilinear* evolution problem of the interface η only (see for example [6, 25, 28, 39, 50]). Moreover, in the case of infinite bottom, if $\eta(t, x)$ is a solution then so is

$$\eta_\lambda(t, x) := \lambda^{-1} \eta(\lambda t, \lambda x), \quad \lambda > 0,$$

and thus the Sobolev space $H^{1+\frac{d}{2}}(\mathbb{R}^d)$ is scaling invariant. Our main results assert that the Muskat problem in arbitrary dimension is locally well-posed for large data in *all subcritical Sobolev spaces* $H^s(\mathbb{R})$, $s > 1 + \frac{d}{2}$, either in the case of one fluid or the case of two fluids with or without viscosity jump, and when the bottom is either empty or is the graph of a Lipschitz function with arbitrarily large variation. We state here an informal version of our main results and refer to Theorems 2.3 and 2.4 for precise statements.

Theorem 1.1. (Informal version) *Let $d \geq 1$ and $s > 1 + \frac{d}{2}$.*

- (i) *(The one-phase problem) Consider $\rho^- > \rho^+ = 0$ and $\mu^- > \mu^+ = 0$. Assume either that the depth is infinite or that the bottom is the graph of a Lipschitz function that does not touch the surface. Then the one-phase Muskat problem is locally well posed in $H^s(\mathbb{R}^d)$.*
- (ii) *(The two-phase problem) Consider $\rho^- > \rho^+ > 0$ and $\mu^\pm > 0$. Assume that the upper and lower boundaries are either empty or graphs of Lipschitz functions that do not touch the interface. The two-phase Muskat problem is locally well posed in $H^s(\mathbb{R}^d)$ in the sense that any initial data in $H^s(\mathbb{R}^d)$ satisfying the Rayleigh–Taylor condition leads to a unique solution in $L^\infty([0, T]; H^s(\mathbb{R}^d))$ for some $T > 0$.*

The starting point of our analysis is the fact that the Muskat problem has a very simple reformulation in terms of the Dirichlet–Neumann map G (see the definition (2.2) below); most strikingly, in the case of one fluid, it is equivalent to

$$\partial_t \eta + G(\eta) \eta = 0 \tag{1.12}$$

(see Proposition 2.1). This makes it clear that

- Any precise result on the continuity of the Dirichlet–Neumann map leads to direct application for the Muskat problem. This is especially relevant in view of the recent intensive work in the context of water-waves [2, 3, 7, 31, 47].
- The Muskat problem is the natural parabolic analog of the water-wave problem and as such is a useful toy-model to understand some of the outstanding challenges for the water-wave problem.

The second point above applies to the study of possible splash singularities, see [14, 15]. Another problem is the question of optimal low-regularity well-posedness for quasilinear problems. This seems a rather formidable problem for water-waves since the mechanism of dispersion is harder to properly pin down in the quasilinear case (see [1–4, 32, 46, 54, 61, 62]), but becomes much more tractable in the case of the Muskat problem due to its parabolicity. This is the question we consider here.

The Muskat problem exists in many incarnations: with or without viscosity jump, with or without surface tension, with or without bottom, with or without permeability jump, in $2d$ or $3d$, when the interface are graphs or curves. Our main objective is to provide a flexible approach that covers many aspects at the same time and provides almost sharp well-posedness results. The main questions that we do not address here are

- The case of surface tension or jump in permeability (see for example [11, 41, 50, 51, 56]). This can also be covered by the paradifferential formalism, but we decided to leave it for another work in order to highlight the centrality of the Dirichlet–Neumann operator.
- The case the interface is not a graph. We believe that so long as the interface is a graph over *some* smooth reference surface, the approach here may be adapted, but this would require substantial additional technicalities.
- The case of beaches when the bottom and the interface meet. This is again a difficult problem (see for example [30]).
- The case of critical regularity. This is a delicate issue, especially for large data, or in the presence of corners. We believe that the approach outlined here could lead to interesting new insights into this question, but the estimates we provide would need to be significantly refined.

Finally, let us stress the fact that in our quasilinear case, there is a significant difference between small and large data, even for local existence. Indeed, the solution is created through some scheme which amounts to decomposing

$$\partial_t \eta = \mathcal{D}\eta + \Pi,$$

where $\partial_t - \mathcal{D}$ can be more or less explicitly integrated, while Π contains the perturbative terms. There are two ways the terms can be perturbative in an expansion

- (1) because they are small and at the same level of regularity,
- (2) because they are more regular.

The first possibility allows us, in the case of small data, to bypass the precise understanding of the terms entailing derivative losses, so long as they are compatible with the regularity of solutions to $(\partial_t - \mathcal{D})\eta = 0$. In our case, when considering large data, we need to extract the terms corresponding to the loss of derivatives in (1.12) and this is where the *paradifferential calculus approach* is particularly useful.

1.3. Prior Results

The Muskat problem was introduced in [53] and has recently been the subject of intense study, both numerically and analytically. Interestingly, the Muskat problem is mathematically analogous to the Hele-Shaw problem [43, 44, 57] for viscous flows between two closely spaced parallel plates. We will mostly discuss the issue of well-posedness and refer to [14, 15, 37] for interesting results on singularity formation and to [36, 40] for recent reviews on the Muskat problem. In the case of small data and infinite depth, global strong solutions have been constructed in subcritical

spaces [13, 17–22, 25, 28, 59] and in critical spaces [39]. We note in particular that [28] allows for interfaces with large slope and that [18, 39] allow for viscosity jump. Global weak solutions were obtained in [20, 29, 38]. We also refer to [26, 27, 34, 42].

As noted earlier there is a significant difference between small and large data for this quasilinear problem. We now discuss in detail the issue of local well-posedness for large data. Early results on local well-posedness for large data in Sobolev spaces date back to [17, 33, 63] and [8, 9]. CÓRDOBA and GANCEDO [25] introduced the contour dynamic formulation for the Muskat problem without viscosity jump and with infinite depth, and proved local well-posedness in $H^{d+2}(\mathbb{R}^d)$, $d = 1, 2$; here the interface is the graph of a function. In [23, 24], CÓRDOBA, CÓRDOBA and GANCEDO extended this result to the case of viscosity jump and nongraph interfaces satisfying the arc-chord and the Rayleigh–Taylor conditions. Note that in the case with viscosity jump, one needs to invert a highly nonlocal equation to obtain the vorticity as an operator of the interface. Using an ALE (Arbitrary Lagrangian–Eulerian) approach, CHENG et al. [18] proved local well-posedness for the one-phase problem with flat bottom when the initial surface $\eta \in H^2(\mathbb{T})$ which allows for unbounded curvature. This result was then extended by MATIOC [51] to the case of viscosity jump (but no bottom). For the case of constant viscosity, using nonlinear lower bounds, the authors in [21] obtained local well-posedness for $\eta \in W^{2,p}(\mathbb{R})$ with $p \in (1, \infty]$. Note that $W^{2,1}(\mathbb{R})$ is scaling invariant yet requires 1/2 more derivative compared to $H^{3/2}(\mathbb{R})$. By rewriting the problem as an abstract parabolic equation in a suitable functional setting, MATIOC [50] sharpened the local well-posedness theory to $\eta \in H^{3/2+\varepsilon}(\mathbb{R})$ for the case of constant viscosity and infinite depth. This covers all subcritical (L^2 -based) Sobolev spaces for the given one-dimensional setting. We also note the recent work of ALAZARD–LAZAR [5] which extends this result by allowing non L^2 -data.

Our Theorem 1.1 thus confirms local-wellposedness for large data in all subcritical Sobolev spaces for a rather general setting allowing for viscosity jump, large bottom variations and higher dimensions. A notable feature of our approach is that it is entirely phrased in terms of the Dirichlet–Neumann operator and as a result, once this operator is properly understood, there is no significant difficulty in passing from constant viscosity to viscosity jump. Furthermore, we obtain an explicit quasilinear parabolic form [see (2.19) and (2.21)] of the Muskat problem by extracting the elliptic and the transport part in the nonlinearity.

1.4. Organization of the Paper

In Section 2, we reformulate the Muskat problem in terms of the Dirichlet–Neumann operator and present the main results of the paper. In Section 3, we properly define the Dirichlet–Neumann operator in our setting and obtain preliminary low-regularity bounds which are then used to obtain paralinearization and contraction estimates in higher norms via a paradifferential approach. These are key technical ingredients for the proof of the main results which are given in Section 4. “Appendix A” gathers trace theorems for homogeneous Sobolev spaces; “Appendix B” is devoted to the proof of (2.8) and (2.9); finally, a review of the paradifferential calculus machinery is presented in “Appendix C”.

2. Reformulation and Main Results

2.1. Reformulation

In order to state our reformulation for the Muskat problem, let us define the Dirichlet–Neumann operators $G^\pm(\eta)$ associated to Ω^\pm . For a given function f , if ϕ^\pm solves

$$\begin{cases} \Delta_{x,y}\phi^\pm = 0 & \text{in } \Omega^\pm, \\ \phi^\pm = f & \text{on } \Sigma, \\ \frac{\partial\phi^\pm}{\partial v^\pm} = 0 & \text{on } \Gamma^\pm \end{cases} \quad (2.1)$$

then

$$G^\pm(\eta)f := \sqrt{1 + |\nabla\eta|^2} \frac{\partial\phi^\pm}{\partial n}. \quad (2.2)$$

The Dirichlet–Neumann operator will be studied in detail in Section 3. We can now restate the Muskat problem in terms of G^\pm .

Proposition 2.1. *Let $d \geq 1$.*

(i) *If (u, p, η) solve the one-phase Muskat problem then $\eta : \mathbb{R}^d \rightarrow \mathbb{R}$ obeys the equation*

$$\partial_t \eta + \kappa G^-(\eta)\eta = 0, \quad \kappa = \frac{\rho^-}{\mu^-}. \quad (2.3)$$

Conversely, if η is a solution of (2.3) then the one-phase Muskat problem has a solution which admits η as the free surface.

(ii) *If (u^\pm, p^\pm, η) is a solution of the two-phase Muskat problem then*

$$\partial_t \eta = -\frac{1}{\mu^-} G^-(\eta) f^-, \quad (2.4)$$

where $f^\pm := p^\pm|_\Sigma + \rho^\pm \eta$ satisfy

$$\begin{cases} f^+ - f^- = (\rho^+ - \rho^-)\eta, \\ \frac{1}{\mu^+} G^+(\eta) f^+ - \frac{1}{\mu^-} G^-(\eta) f^- = 0. \end{cases} \quad (2.5)$$

Conversely, if η is a solution of (2.4) with f^\pm solution of (2.5) then the two-phase Muskat problem has a solution which admits η as the free interface.

We refer to [6, 16] for similar reformulations and derivation of a number of interesting properties.

Proof. (i) Assume first that (u^-, p^-, η) solve the one-phase Muskat problem. Setting $q = p^- + \rho^- y$, then q solves the elliptic problem

$$\Delta_{x,y}q = 0 \quad \text{in } \Omega_t^-, \quad q = \rho^- \eta \quad \text{on } \Sigma_t, \quad \partial_v q = 0 \quad \text{on } \Gamma^-. \quad (2.6)$$

Since $\sqrt{1 + |\nabla\eta|^2} u^- \cdot n|_{\Sigma_t} = -G^-(\eta)(\rho^- \eta)$, (2.3) follows from (1.8) and (1.5).

Conversely, if η satisfies (2.3) then the pressure $p^- = q - \rho^- y$ is obtained by solving (2.6), and the velocity is determined from the Darcy's law (1.5).

(ii) As before, (2.4) follows from (1.8) and (1.5) for Ω^- . The jump of f in (2.5) is a consequence of the continuity (1.9) of the pressure. Lastly, the jump of Dirichlet–Neumann operators is exactly the continuity (1.7) of the normal velocity. Conversely, if η is known then (u^\pm, p^\pm) can be easily determined. \square

Remark 2.2. For a given function $\eta \in W^{1,\infty}(\mathbb{R}^d) \cap H^{\frac{1}{2}}(\mathbb{R}^d)$, we prove in Proposition 4.8 below that there exists a unique pair f^\pm solving (2.5) in a variational sense.

2.2. Main Results

The Rayleigh–Taylor stability condition requires that the pressure is increasing in the normal direction when crossing the interface from the top fluid to the bottom fluid. More precisely,

$$RT(x, t) = (\nabla_{x,y} p^+ - \nabla_{x,y} p^-) \cdot n|_{\Sigma_t} > 0. \quad (2.7)$$

In terms of η and f^\pm , we have

$$RT(x, t) = (1 + |\nabla \eta|^2)^{\frac{1}{2}} (\llbracket \rho \rrbracket - \llbracket B \rrbracket), \quad (2.8)$$

where

$$\llbracket \rho \rrbracket = \rho^- - \rho^+, \quad \llbracket B \rrbracket = B^- - B^+$$

and

$$B^\pm = \frac{\nabla \eta \cdot \nabla f^\pm + G^\pm(\eta) f^\pm}{1 + |\nabla \eta|^2}.$$

Using the Darcy law (1.5) we can write that

$$RT(x, t) = \llbracket \rho \rrbracket (1 + |\nabla \eta|^2)^{-\frac{1}{2}} + \llbracket \mu \rrbracket u \cdot n, \quad \llbracket \mu \rrbracket = (\mu^- - \mu^+). \quad (2.9)$$

See Appendix B for the proof of (2.8) and (2.9). Let us denote

$$Z^s(T) = L^\infty([0, T]; H^s(\mathbb{R}^d)) \cap L^2([0, T]; H^{s+\frac{1}{2}}(\mathbb{R}^d)). \quad (2.10)$$

For the one-phase problem, we prove local well-posedness *without* assuming the Rayleigh–Taylor stability condition which in fact *always holds*, even in finite depth (see Remark 2.6).

Theorem 2.3. *Let $\mu^- > 0$ and $\rho^- > 0$. Let $s > 1 + \frac{d}{2}$ with $d \geq 1$. Consider either $\Gamma^- = \emptyset$ or $b^- \in \dot{W}^{1,\infty}(\mathbb{R}^d)$. Let $\eta_0 \in H^s(\mathbb{R}^d)$ satisfy*

$$\text{dist}(\eta_0, \Gamma^-) \geq 2h > 0. \quad (2.11)$$

Then, there exist a positive time T depending only on (s, κ) , h and $\|\eta_0\|_{H^s(\mathbb{R}^d)}$, and a unique solution $\eta \in Z^s(T)$ to equation (2.3) such that $\eta|_{t=0} = \eta_0$ and

$$\text{dist}(\eta(t), \Gamma^-) \geq h \quad \forall t \in [0, T]. \quad (2.12)$$

Furthermore, the L^2 norm of η in nonincreasing in time.

As for the two-phase problem, we prove local well-posedness in the stable regime ($\rho^+ < \rho^-$) for large data satisfying the Rayleigh–Taylor stability condition.

Theorem 2.4. *Let $\mu^\pm > 0$ and $\rho^- > \rho^+ > 0$. Let $s > 1 + \frac{d}{2}$ with $d \geq 1$. Consider any combination of $\Gamma^\pm = \emptyset$ and $\underline{b}^\pm \in \dot{W}^{1,\infty}(\mathbb{R}^d)$. Let $\eta_0 \in H^s(\mathbb{R}^d)$ satisfy*

$$\text{dist}(\eta_0, \Gamma^\pm) \geq 2h > 0, \quad (2.13)$$

$$\inf_{x \in \mathbb{R}^d} \text{RT}(x, 0) > 2\alpha > 0. \quad (2.14)$$

Then, there exist a positive time T depending only on $(s, \mu^\pm, [\![\rho]\!])$, (h, α) and $\|\eta_0\|_{H^s(\mathbb{R}^d)}$, and a unique solution $\eta \in Z^s(T)$ to Eqs. (2.4)–(2.5) such that $\eta|_{t=0} = \eta_0$,

$$\text{dist}(\eta(t), \Gamma^\pm) \geq h \quad \forall t \in [0, T], \quad (2.15)$$

$$\inf_{(x,t) \in \mathbb{R}^d \times [0, T]} \text{RT}(x, t) > \alpha. \quad (2.16)$$

Furthermore, the L^2 norm of η is nonincreasing in time.

Several remarks on our main results are in order.

Remark 2.5. The solutions constructed in Theorems 2.3 and 2.4 are unique in $L_t^\infty H_x^s$ and the solution maps are locally Lipschitz in $L_t^\infty H_x^s$ with respect to the topology of $L_t^\infty H_x^{s-1}$. The proof of Theorem 2.4 also provides the following estimate for f^\pm :

$$\|f^\pm\|_{\tilde{H}_\pm^s(\mathbb{R}^d)} \leq \mathcal{F}(\|\eta\|_{H^s(\mathbb{R}^d)}) \|\eta\|_{H^s(\mathbb{R}^d)}, \quad (2.17)$$

where the space $\tilde{H}_\pm^s(\mathbb{R}^d)$ is defined by (3.24). Modulo some minor modifications, our proofs work equally for the periodic case.

Remark 2.6. The Rayleigh–Taylor (RT) condition is ubiquitous in free boundary problems. For irrotational water-waves (one fluid), WU [61] proved that this condition is automatically satisfied if there is no bottom. In the presence of a bottom that is the graph of a function, LANNEZ [47] proved this condition assuming that the second fundamental form of the bottom is sufficiently small, covering the case of flat bottoms. In the context of the Muskat problem, there are various scenarios for the stable regime $\rho^+ < \rho^-$. When the interface is a general curve/surface, the RT condition was assumed in [8, 23, 24]. On the other hand, when the interface is a graph, we see from (2.9) that this condition always holds if there is no viscosity jump but need not be so otherwise. In particular, for the one-phase problem, the local well-posedness result in [18] assumes the RT condition for flat bottoms. However, we prove in Proposition 4.3 that the RT condition holds in the one-phase case so long as the bottom is either empty or is the graph of a Lipschitz function which can be unbounded and have large variation.

Remark 2.7. When the surface tension effect is taken into account, well-posedness holds without the Rayleigh–Taylor condition. It turns out that $H^{1+\frac{d}{2}}(\mathbb{R}^d)$ is also the scaling invariant Sobolev space for the Muskat problem with surface tension. Local well-posedness for all subcritical data in $H^s(\mathbb{R}^d)$, $s > 1 + \frac{d}{2}$, is established

in [55]. Furthermore, at the same level of regularity, Flynn and Nguyen [35] proves that solutions constructed in Theorems 2.3 and 2.4 are limits of solutions to the problem with surface tension as surface tension vanishes.

2.3. Strategy of Proof

Let us briefly explain our strategy for a priori estimates. The main step consists in obtaining a precise paralinearization for the Dirichlet–Neumann operator $G^\pm(\eta) f$ when $\eta \in H^s(\mathbb{R}^d)$, $s > 1 + \frac{d}{2}$ and f has the maximal regularity $H^s(\mathbb{R}^d)$. We prove in Theorem 3.18 that

$$G^-(\eta) f = T_\lambda(f - T_B \eta) - T_V \cdot \nabla \eta + R^-(\eta) f, \quad (2.18)$$

where (B, V) are explicit functions (see (3.54)), λ is an elliptic first-order symbol (see (3.50)) and the remainder $R^-(\eta) f$ obeys

$$\|R^-(\eta) f\|_{H^{s-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s})(1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|f\|_{\tilde{H}^s_-},$$

provided that $0 < \delta < \min(s - 1 - \frac{d}{2}, \frac{1}{2})$. Here we note that the term $f - T_B \eta$ comes from the consideration of Alinhac’s good unknown.

(1) For the one-phase problem (2.3), taking $f = \eta$ yields

$$\partial_t \eta = -\kappa T_{\lambda(1-B)} \eta + \kappa T_V \cdot \nabla \eta + R_1, \quad (2.19)$$

where

$$\|R_1\|_{H^{s-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s})(1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|\eta\|_{H^s}. \quad (2.20)$$

We observe that the transport term $T_V \cdot \nabla \eta$ is harmless for energy estimates and the term $-\kappa T_{\lambda(1-B)} \eta$ would give the parabolicity if $1 - B > 0$. Then this latter term entails a gain of $\frac{1}{2}$ derivative when measured in L_t^2 , compensating the loss of $\frac{1}{2}$ derivative in the remainder R_1 . Moreover, the fact that the highest order term $\|\eta\|_{H^{s+\frac{1}{2}-\delta}}$ in (2.20) appears linearly with a gain of δ derivative gives room to choose the time T as a small parameter. We thus obtain a closed a priori estimate in $L_t^\infty H_x^s \cap L_t^2 H_x^{s+\frac{1}{2}}$. Finally, we prove in Proposition 4.3 that the stability condition $1 - B > 0$ is automatically satisfied.

(2) As for the two-phase problem (2.4)–(2.5), we apply the paralinearization (2.18) and obtain a reduced equation similar to (2.19):

$$\partial_t \eta = -\frac{1}{\mu^+ + \mu^-} T_{\lambda([\rho] - [B])} \eta + \frac{1}{\mu^+ + \mu^-} T_{[V]} \cdot \nabla \eta + R_2, \quad (2.21)$$

where R_2 obeys the same bound (2.20) as R_1 . Consequently, the parabolicity holds if $[\rho] - [B] > 0$ and in view of (2.8), this is equivalent to $RT > 0$. This shows a remarkable link between Alinhac’s good unknown and the Rayleigh–Taylor stability condition.

Finally, we remark that the contraction estimate for the solutions requires a fine contraction estimate for the Dirichlet–Neumann operator, see Theorem 3.24.

3. The Dirichlet–Neumann Operator: Continuity, Paralinearization and Contraction Estimates

This section is devoted to the study of the Dirichlet–Neumann operator. For the two-phase problem (2.4), the function f^- obtained from solving (2.5) is only determined up to additive constants and we need to define $G^-(\eta)f$ for f belonging to a suitable homogeneous space. Since f is the trace of a harmonic function (see (2.1)) with bounded gradient in L^2 , the trace theory recently developed in [49] is perfectly suited for this purpose, allowing us to take f in a “screened” homogeneous Sobolev space (see (3.5)) tailored to the bottom. This is the content of Section 3.1, where we obtain existence and modest regularity of the variational solution to the appropriate Dirichlet problem.

Next, in Section 3.2, we obtain a precise paralinearization for $G^-(\eta)f$ by extracting all the first order symbols. This is done when η has subcritical regularity $H^s(\mathbb{R}^d)$, $s > 1 + \frac{d}{2}$, and f has the maximal regularity $H^s(\mathbb{R}^d)$. The error estimate is precise enough to obtain closed a priori estimates afterwards. Finally, in Section 3.3 we prove a contraction estimate for $G^-(\eta_1) - G^-(\eta_2)$, showing a gain of derivative for $\eta_1 - \eta_2$ which will be crucial for the contraction estimate of solutions.

3.1. Definition and Continuity

We study the Dirichlet–Neumann problem associated to the fluid domain Ω^- underneath the free interface $\Sigma = \{(x, \eta(x)) : x \in \mathbb{R}^d\}$. Here and in what follows, the time variable is frozen. We say that a function $g : \mathbb{R}^d \rightarrow \mathbb{R}$ is Lipschitz, $g \in \dot{W}^{1,\infty}(\mathbb{R}^d)$, if $\nabla g \in L^\infty(\mathbb{R}^d)$. As for the bottom Γ^- , we assume that either

- $\Gamma^- = \emptyset$ or
- $\Gamma^- = \{(x, \underline{b}^-(x)) : x \in \mathbb{R}^d\}$ where $\underline{b}^- \in \dot{W}^{1,\infty}(\mathbb{R})$ satisfies

$$\text{dist}(\Sigma, \Gamma^-) > h > 0. \quad (3.1)$$

In either case, $\text{dist}(\Sigma, \Gamma^-) > h > 0$. Consider the elliptic problem

$$\begin{cases} \Delta_{x,y}\phi = 0 & \text{in } \Omega^-, \\ \phi = f & \text{on } \Sigma, \\ \frac{\partial \phi}{\partial \nu^-} = 0 & \text{on } \Gamma^-, \end{cases} \quad (3.2)$$

where in the case of infinite depth ($\Gamma^- = \emptyset$), the Neumann condition is replaced by the vanishing of $\nabla_{x,y}\phi$ as $y \rightarrow -\infty$

$$\lim_{y \rightarrow -\infty} \nabla_{x,y}\phi = 0. \quad (3.3)$$

The Dirichlet–Neumann operator associated to Ω^- is formally defined by

$$G^-(\eta)f = \sqrt{1 + |\nabla \eta|^2} \frac{\partial \phi}{\partial n}, \quad (3.4)$$

where we recall that n is the upward-pointing unit normal to Σ . Similarly, if ϕ solves the elliptic problem (3.2) with $(\Omega^-, \Gamma^-, v^-)$ replaced by $(\Omega^+, \Gamma^+, v^+)$ then we define

$$G^+(\eta)f = \sqrt{1 + |\nabla \eta|^2} \frac{\partial \phi}{\partial n}.$$

Note that n is inward-pointing for Ω^+ . In the rest of this section, we only state results for $G^-(\eta)$ since corresponding results for $G^+(\eta)$ are completely parallel.

The Dirichlet data f for (3.2) will be taken in the following “screened” fractional Sobolev space [49]

$$\begin{aligned} \tilde{H}_\Theta^{\frac{1}{2}}(\mathbb{R}^d) = \left\{ f \in \mathcal{S}'(\mathbb{R}^d) \cap L_{loc}^2(\mathbb{R}^d) : \right. \\ \left. \int_{\mathbb{R}^d} \int_{B_{\mathbb{R}^d}(0, \Theta(x))} \frac{|f(x+k) - f(x)|^2}{|k|^{d+1}} dk dx < \infty \right\} / \mathbb{R}, \quad (3.5) \end{aligned}$$

where $\Theta : \mathbb{R}^d \rightarrow (0, \infty]$ is a given lower semi-continuous function. We will choose for an arbitrary number $a \in (0, 1)$ that

$$\Theta(x) = \begin{cases} \infty & \text{when } \Gamma^- = \emptyset, \\ \mathfrak{d}(x) := a(\eta(x) - \underline{b}^-(x)) & \text{when } \underline{b}^- \in \dot{W}^{1,\infty}(\mathbb{R}^d). \end{cases} \quad (3.6)$$

In view of assumption (3.1),

$$\mathfrak{d} \geq ah. \quad (3.7)$$

We also define the slightly-homogeneous Sobolev spaces

$$H^{1,\sigma}(\mathbb{R}^d) = \{f \in \mathcal{S}'(\mathbb{R}^d) \cap L_{loc}^2(\mathbb{R}^d) : \nabla f \in H^{\sigma-1}(\mathbb{R}^d)\} / \mathbb{R}. \quad (3.8)$$

Remark 3.1. According to Theorem 2.2(b) in [60], $f \in \tilde{H}_1^{\frac{1}{2}}(\mathbb{R}^d)$ ($\Theta \equiv 1$) if and only if $f \in \mathcal{S}'(\mathbb{R}^d) \cap L_{loc}^2(\mathbb{R}^d)$ and \hat{f} is locally L^2 in the complement of the origin such that

$$\int_{\mathbb{R}^d} \min(|\xi|, |\xi|^2) |\hat{f}(\xi)|^2 d\xi < \infty; \quad (3.9)$$

moreover, $\|f\|_{\tilde{H}_1^{\frac{1}{2}}(\mathbb{R}^d)}^2$ is bounded above and below by a multiple of (3.9) so that

$$\tilde{H}_1^{\frac{1}{2}}(\mathbb{R}^d) = \dot{H}^{1,\frac{1}{2}}(\mathbb{R}^d). \quad (3.10)$$

On the other hand, $f \in \tilde{H}_\infty^{\frac{1}{2}}(\mathbb{R}^d)$ ($\Theta \equiv \infty$) if and only if $f \in L_{loc}^2(\mathbb{R}^d)$ and \hat{f} is locally L^2 in the complement of the origin, with

$$\int_{\mathbb{R}^d} |\xi| |\hat{f}(\xi)|^2 d\xi < \infty; \quad (3.11)$$

moreover, $\|f\|_{\tilde{H}_\infty^{\frac{1}{2}}(\mathbb{R}^d)}^2$ is a constant multiple of (3.11). Thus, we have the continuous embeddings

$$\dot{H}^{\frac{1}{2}}(\mathbb{R}^d) = \tilde{H}_\infty^{\frac{1}{2}}(\mathbb{R}^d) \subset \tilde{H}_\Theta^{\frac{1}{2}}(\mathbb{R}^d) \subset \tilde{H}_1^{\frac{1}{2}}(\mathbb{R}^d) = H^{1,\frac{1}{2}}(\mathbb{R}^d) \quad (3.12)$$

upon recalling the lower bound (3.7) for \mathfrak{d} . In addition, under condition (3.1),

$$\text{if } \underline{\eta}, \underline{b}^- \in W^{1,\infty}(\mathbb{R}^d) \text{ then } \widetilde{H}_0^{\frac{1}{2}}(\mathbb{R}^d) = \widetilde{H}_1^{\frac{1}{2}}(\mathbb{R}^d). \quad (3.13)$$

See Theorem 3.13 [49]. To accommodate unbounded bottoms, we have only assumed that $\underline{b}^- \in \dot{W}^{1,\infty}(\mathbb{R}^d)$ and thus (3.13) is not applicable. Nevertheless, we have the following proposition:

Proposition 3.2. *Assume that $\sigma_1, \sigma_2 : \mathbb{R}^d \rightarrow (0, \infty]$ satisfy*

$$\inf_{x \in \mathbb{R}^d} \{\sigma_1(x), \sigma_2(x)\} > h > 0, \quad \|\sigma_1 - \sigma_2\|_{L^\infty(\mathbb{R}^d)} \leq M < \infty. \quad (3.14)$$

Then there exists $C = C(d, h, M)$ such that

$$\|f\|_{\widetilde{H}_{\sigma_1}^{\frac{1}{2}}(\mathbb{R}^d)} \leq C \|f\|_{\widetilde{H}_{\sigma_2}^{\frac{1}{2}}(\mathbb{R}^d)} \quad \forall f \in \widetilde{H}_{\sigma_2}^{\frac{1}{2}}(\mathbb{R}^d). \quad (3.15)$$

It follows that for any two surfaces η_1 and η_2 in $L^\infty(\mathbb{R}^d)$ satisfying (3.1), the screened Sobolev space $\widetilde{H}_0^{\frac{1}{2}}(\mathbb{R}^d)$, \mathfrak{d} given by (3.6), is independent of η_j . The proof of Proposition 3.2 is given in Appendix A.4.

We will solve (3.2) in the homogeneous Sobolev space $\dot{H}^1(\Omega^-)$ where

$$\dot{H}^1(U) = \{u \in L^2_{loc}(U) : \nabla u \in L^2(U)\} / \mathbb{R} \quad (3.16)$$

for $U \subset \mathbb{R}^N$ connected. Here, the norm of $\dot{H}^1(U)$ is given by $\|u\|_{\dot{H}^1(U)} = \|\nabla u\|_{L^2(U)}$.

Proposition 3.3. *The vector space $\dot{H}^1(U)$ equipped with the norm $\|u\|_{\dot{H}^1(U)} = \|\nabla u\|_{L^2(U)}$ is complete.*

Proof. Suppose that u_n is a Cauchy sequence in $\dot{H}^1(U)$. Then $\nabla u_n \rightarrow F$ in $L^2(U)$. We claim that $F = \nabla u$ for some $u \in L^2_{loc}(U)$. Indeed, for any bounded domain $V \subset U$, the sequence $u_n - |V|^{-1} \int_V u_n$ is bounded in $L^2(V)$, according to the Poincaré inequality, hence weakly converges in $L^2(V)$. By a diagonal process, we can find $u \in L^2_{loc}(U)$ and a subsequence $n_k \rightarrow \infty$ such that

$$u_{n_k} - |V|^{-1} \int_V u_{n_k} \rightharpoonup u \quad \text{in } L^2(V)$$

for any bounded $V \subset U$. Let $\varphi \in C_c^\infty(U)$ be a test vector field with $\text{supp } \varphi \subset V \Subset U$. We have

$$\begin{aligned} \int_U \varphi \cdot \nabla u_n \, dx &= \int_V \varphi \nabla (u_n - |V|^{-1} \int_V u_n \, dy) \, dx \\ &= - \int_V (u_n - |V|^{-1} \int_V u_n \, dy) \, \text{div } \varphi \, dx \rightarrow - \int_V u \, \text{div } \varphi \, dx. \end{aligned}$$

Thus,

$$\int_U F \cdot \varphi \, dx = \lim_{n \rightarrow \infty} \int_U \nabla u_n \cdot \varphi \, dx = - \int_U u \, \text{div } \varphi \, dx$$

for any test vector field φ . This proves that $F = \nabla u$ and thus finishes the proof. \square

We refer to ‘‘Appendix A’’ of the present paper for a summary of trace theory, taken from [49], when U is an infinite strip-like domain or a Lipschitz half space.

Proposition 3.4. *Consider the finite-depth case with $\underline{b}^- \in \dot{W}^{1,\infty}(\mathbb{R}^d)$. If $\eta \in \dot{W}^{1,\infty}(\mathbb{R}^d)$ then for every $f \in \widetilde{H}_0^{\frac{1}{2}}(\mathbb{R}^d)$, there exists a unique variational solution $\phi \in \dot{H}^1(\Omega^-)$ to (3.2). Moreover, ϕ satisfies*

$$\|\nabla_{x,y}\phi\|_{L^2(\Omega^-)} \leq \mathcal{F}(\|\nabla\eta\|_{L^\infty}) \|f\|_{\widetilde{H}_0^{\frac{1}{2}}(\mathbb{R}^d)} \quad (3.17)$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on h and $\|\nabla \underline{b}^-\|_{L^\infty(\mathbb{R}^d)}$.

Proof. By virtue of Theorem A.2, there exists $\underline{f} \in \dot{H}^1(\Omega^-)$ such that $\text{Tr}(\underline{f})(x, \eta(x)) = f(x)$, $\text{Tr}(\underline{f})(x, \underline{b}^-(x)) = f(x)$, and

$$\|\nabla_{x,y}\underline{f}\|_{L^2(\Omega^-)} \leq \mathcal{F}(\|\nabla\eta\|_{L^\infty}) \|f\|_{\widetilde{H}_0^{\frac{1}{2}}(\mathbb{R}^d)} \quad (3.18)$$

where \mathcal{F} depends only on h and $\|\nabla \underline{b}^-\|_{L^\infty(\mathbb{R}^d)}$. Set

$$H_{0,*}^1(\Omega^-) = \{u \in \dot{H}^1(\Omega^-) : \text{Tr}(u)|_\Sigma = 0\}$$

endowed with the norm of $\dot{H}^1(\Omega^-)$. We then define ϕ solution to (3.2) to be

$$\phi(x, y) = \underline{f}(x, y) + u(x, y), \quad (3.19)$$

where $u \in H_{0,*}^1(\Omega^-)$ is the unique solution to the variational problem

$$\int_{\Omega^-} \nabla_{x,y} u \cdot \nabla_{x,y} \varphi \, dx \, dy = - \int_{\Omega^-} \nabla_{x,y} \underline{f} \cdot \nabla_{x,y} \varphi \, dx \, dy \quad \forall \varphi \in H_{0,*}^1(\Omega^-). \quad (3.20)$$

The existence and uniqueness of u is guaranteed by the Lax–Milgram theorem upon using the bound (3.18). Setting $\varphi = u$ in (3.20) and recalling the definition (3.19) of ϕ we obtain the estimate (3.17). It follows from (3.20) that

$$\int_{\Omega^-} \nabla_{x,y} \phi \cdot \nabla_{x,y} \varphi \, dx \, dy = 0 \quad \forall \varphi \in H_{0,*}^1(\Omega^-).$$

Thus, if ϕ is smooth then ϕ solves (3.2) in the classical sense upon integrating by parts. Finally, it is easy to see that the solution ϕ constructed by (3.19) and (3.20) is independent of the choice of $\underline{f} \in \dot{H}^1(\Omega^-)$ that has trace f on Σ . \square

Remark 3.5. As the functions \underline{b}^\pm are fixed, we shall omit the dependence on $\|\nabla \underline{b}^\pm\|_{L^\infty(\mathbb{R}^d)}$.

Proposition 3.6. *Consider the infinite-depth case $\Gamma^- = \emptyset$. If $\eta \in \dot{W}^{1,\infty}(\mathbb{R}^d)$ then for every $f \in \widetilde{H}_\infty^{\frac{1}{2}}(\mathbb{R}^d)$ there exists a unique variational solution $\phi \in \dot{H}^1(\Omega^-)$ to (3.2). Moreover, ϕ satisfies*

$$\|\nabla_{x,y}\phi\|_{L^2(\Omega^-)} \leq \mathcal{F}(\|\nabla\eta\|_{L^\infty}) \|f\|_{\widetilde{H}_\infty^{\frac{1}{2}}(\mathbb{R}^d)} \quad (3.21)$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on h .

Proof. The proof follows along the same lines as in the proof of Proposition 3.4 upon using the trace Theorem A.3 and the lifting Theorem A.4 for the half space $U = \Omega^-$. The fact that $\nabla\phi \in L^2(\Omega^-)$ gives a sense to the boundary condition (3.3). \square

Notation 3.7. We denote

$$\tilde{H}_-^{\frac{1}{2}} = \begin{cases} \tilde{H}_\infty^{\frac{1}{2}}(\mathbb{R}^d) & \text{if } \Gamma^- = \emptyset, \\ \tilde{H}_{\mathfrak{d}'}^{\frac{1}{2}}(\mathbb{R}^d) & \text{if } \underline{b}^- \in \dot{W}^{1,\infty}(\mathbb{R}^d), \end{cases} \quad (3.22)$$

and

$$\tilde{H}_+^{\frac{1}{2}} = \begin{cases} \tilde{H}_\infty^{\frac{1}{2}}(\mathbb{R}^d) & \text{if } \Gamma^+ = \emptyset, \\ \tilde{H}_{\mathfrak{d}'}^{\frac{1}{2}}(\mathbb{R}^d) & \text{if } \underline{b}^+ \in \dot{W}^{1,\infty}(\mathbb{R}^d), \quad \mathfrak{d}'(x) := a(\eta(x) - \underline{b}^+(x)), \quad a \in (0, 1). \end{cases} \quad (3.23)$$

For $s > \frac{1}{2}$, we denote

$$\tilde{H}_\pm^s = \tilde{H}_\pm^{\frac{1}{2}} \cap H^{1,s}(\mathbb{R}^d). \quad (3.24)$$

Proposition 3.8. *If $\eta \in \dot{W}^{1,\infty}(\mathbb{R}^d)$ then the Dirichlet–Neumann operator is continuous from $\tilde{H}_-^{\frac{1}{2}}$ to $H^{-\frac{1}{2}}(\mathbb{R}^d)$. Moreover, there exists a constant $C > 0$ depending only on h such that*

$$\|G^-(\eta)f\|_{H^{-\frac{1}{2}}} \leq \mathcal{F}(\|\nabla\eta\|_{L^\infty})\|f\|_{\tilde{H}_-^{\frac{1}{2}}}. \quad (3.25)$$

Proof. Let ϕ solve (3.2). By virtue of Propositions 3.4 and 3.6, we have $\phi \in \dot{H}^1(\Omega^-)$ and $\Delta\phi = 0$. According to Theorem A.5, the trace

$$\sqrt{1 + |\nabla\eta|^2} \frac{\partial\phi}{\partial n}|_\Sigma = G^-(\eta)f$$

is well-defined in $H^{-\frac{1}{2}}(\mathbb{R}^d)$ and

$$\|G^-(\eta)f\|_{H^{-\frac{1}{2}}} \leq Ch^{-\frac{1}{2}}(1 + \|\nabla\eta\|_{L^\infty})\|\nabla\phi\|_{L^2(\Omega_h^-)}$$

where C is an absolute constant and $\Omega_h^- = \{(x, y) \in \mathbb{R}^{d+1} : \eta(x) - h < y < \eta(x)\}$. Thus, (3.25) follows from (3.17) and (3.21). \square

To propagate higher Sobolev regularity for ϕ and hence for $G^-(\eta)f$, following [3,47] we straighten the boundary as follows: set

$$\begin{cases} \Omega_1^- = \{(x, y) : x \in \mathbb{R}^d, \eta(x) - h < y < \eta(x)\}, \\ \Omega_2^- = \{(x, y) \in \Omega^- : y \leq \eta(x) - h\} \end{cases} \quad (3.26)$$

and

$$\begin{cases} \widetilde{\Omega_1^-} = \mathbb{R}^d \times (-1, 0), \\ \widetilde{\Omega_2^-} = \{(x, z) \in \mathbb{R}^d \times (-\infty, -1] : (x, z + 1 + \eta(x) - h) \in \Omega_2^-\}, \\ \widetilde{\Omega^-} = \widetilde{\Omega_1^-} \cup \widetilde{\Omega_2^-}. \end{cases} \quad (3.27)$$

Define

$$\begin{cases} \varrho(x, z) = (1+z)e^{\tau z \langle D_x \rangle} \eta(x) - z \{ e^{-(1+z)\tau \langle D_x \rangle} \eta(x) - h \} & \text{if } (x, z) \in \widetilde{\Omega_1^-}, \\ \varrho(x, z) = z + 1 + \eta(x) - h & \text{if } (x, z) \in \widetilde{\Omega_2^-}, \end{cases} \quad (3.28)$$

where $\tau > 0$ will be chosen in the next lemma.

Lemma 3.9. *Assume $\eta \in B_{\infty,1}^1(\mathbb{R}^d)$.*

(1) *There exists a constant $C > 0$ independent of τ such that*

$$\|\nabla_{x,z} \varrho\|_{L^\infty(\widetilde{\Omega}^-)} \leq 1 + C \|\eta\|_{B_{\infty,1}^1}$$

for all $(x, z) \in \widetilde{\Omega}^-$.

(2) *There exists $K > 0$ such that if*

$$\tau \|\eta\|_{B_{\infty,1}^1} \leq \frac{h}{2K} \quad (3.29)$$

then $\min(1, \frac{h}{2}) \leq \partial_z \varrho \leq K \|\eta\|_{B_{\infty,1}^1}$ and thus the mappings $(x, z) \in \widetilde{\Omega}_k^- \mapsto (x, \varrho(x, z)) \in \Omega_k^-$, $k = 1, 2$ are Lipschitz diffeomorphisms.

Lemma 3.9 follows from straightforward calculations which we omit. Note that $H^s(\mathbb{R}^d) \subset B_{\infty,1}^1(\mathbb{R}^d)$ for any $s > 1 + \frac{d}{2}$. A direct calculation shows that if $f : \Omega^- \rightarrow \mathbb{R}$ then $\tilde{f}(x, z) = f(x, \varrho(x, z))$ satisfies

$$\operatorname{div}_{x,z}(\mathcal{A} \nabla_{x,z} \tilde{f})(x, z) = \partial_z \varrho (\Delta_{x,y} f)(x, \varrho(x, z)) \quad (3.30)$$

with

$$\mathcal{A} = \begin{bmatrix} \partial_z \varrho \cdot Id & -\nabla_x \varrho \\ -(\nabla_x \varrho)^T & \frac{1 + |\nabla_x \varrho|^2}{\partial_z \varrho} \end{bmatrix}. \quad (3.31)$$

In order to study functions inside the domain, we introduce adapted functional spaces. Given $\mu \in \mathbb{R}$ we define the interpolation spaces

$$\begin{aligned} X^\mu(I) &= C_z^0(I; H^\mu(\mathbb{R}^d)) \cap L_z^2\left(I; H^{\mu+\frac{1}{2}}(\mathbb{R}^d)\right), \\ Y^\mu(I) &= L_z^1(I; H^\mu(\mathbb{R}^d)) + L_z^2\left(I; H^{\mu-\frac{1}{2}}(\mathbb{R}^d)\right). \end{aligned} \quad (3.32)$$

We prove the following useful inequalities:

Lemma 3.10. *Let s_0, s_1 and s_2 be real numbers, and let $J \subset \mathbb{R}$.*

(1) *If*

$$\begin{cases} s_0 \leq \min\{s_1 + 1, s_2 + 1\}, \\ s_1 + s_2 > s_0 + \frac{d}{2} - 1, \\ s_1 + s_2 + 1 > 0, \end{cases} \quad (3.33)$$

then

$$\|u_1 u_2\|_{Y^{s_0}(J)} \lesssim \|u_1\|_{X^{s_1}(J)} \|u_2\|_{X^{s_2}(J)}. \quad (3.34)$$

(2) If

$$\begin{cases} s_0 \leq \min\{s_1, s_2\}, \\ s_1 + s_2 > s_0 + \frac{d}{2}, \\ s_1 + s_2 > 0, \end{cases} \quad (3.35)$$

then

$$\|u_1 u_2\|_{Y^{s_0}(J)} \lesssim \|u_1\|_{Y^{s_1}(J)} \|u_2\|_{X^{s_2}(J)}. \quad (3.36)$$

In fact, we have

$$\begin{aligned} \|T_{u_2} u_1\|_{Y^{s_0}(J)} &\lesssim \|u_1\|_{Y^{s_1}(J)} \|u_2\|_{L^\infty(J; H^{s_2})} \quad \text{if } s_0 \leq s_1, \\ s_1 + s_2 &> s_0 + \frac{d}{2}, \end{aligned} \quad (3.37)$$

$$\begin{aligned} \|T_{u_1} u_2\|_{Y^{s_0}(J)} &\lesssim \|u_1\|_{Y^{s_1}(J)} \|u_2\|_{L^\infty(J; H^{s_2})} \quad \text{if } s_0 \leq s_2, \\ s_1 + s_2 &> s_0 + \frac{d}{2}, \end{aligned} \quad (3.38)$$

$$\begin{aligned} \|R(u_1, u_2)\|_{L^1(J; H^{s_0})} &\lesssim \|u_1\|_{Y^{s_1}(J)} \|u_2\|_{X^{s_2}(J)} \quad \text{if } s_1 + s_2 > 0, \\ s_1 + s_2 &> s_0 + \frac{d}{2}. \end{aligned} \quad (3.39)$$

The proof of Lemma 3.10 is given in ‘‘Appendix D’’.

Lemma 3.11. *Let $\eta \in H^s(\mathbb{R}^d)$ with $s > 1 + \frac{d}{2}$ and $\tau > 0$ such that (3.29) holds. If*

$$\begin{aligned} \nabla_{x,z} \tilde{f} &\in L^2([-1, 0]; L^2(\mathbb{R}^d)), \\ \operatorname{div}_{x,z}(\mathcal{A} \nabla_{x,z} \tilde{f}) &\in L^2([-1, 0]; H^{-1}(\mathbb{R}^d)) \end{aligned}$$

then $\nabla_{x,z} \tilde{f} \in X^{-\frac{1}{2}}([-1, 0])$ and

$$\begin{aligned} \|\nabla_{x,z} \tilde{f}\|_{X^{-\frac{1}{2}}([-1, 0])} &\leq \mathcal{F}(\|\eta\|_{H^s})(\|\nabla_{x,z} \tilde{f}\|_{L^2([-1, 0]; L^2)} \\ &\quad + \|\operatorname{div}_{x,z}(\mathcal{A} \nabla_{x,z} \tilde{f})\|_{L^2([-1, 0]; H^{-1})}). \end{aligned} \quad (3.40)$$

Proof. By the definition of $X^{-\frac{1}{2}}([-1, 0])$, it suffices to prove that $\nabla_{x,z} \tilde{f} \in C([-1, 0]; H^{-\frac{1}{2}}(\mathbb{R}^d))$ with norm bounded by the right-hand side of (3.40). By virtue of the interpolation Theorem A.6, $\nabla_x \tilde{f} \in C([-1, 0]; H^{-\frac{1}{2}}(\mathbb{R}^d))$ and

$$\begin{aligned} \|\nabla_x \tilde{f}\|_{C([-1, 0]; H^{-\frac{1}{2}})} &\lesssim \|\nabla_x \tilde{f}\|_{L^2([-1, 0]; L^2)} + \|\partial_z \nabla_x \tilde{f}\|_{L^2([-1, 0]; H^{-1})} \\ &\lesssim \|\nabla_{x,z} \tilde{f}\|_{L^2([-1, 0]; L^2)}. \end{aligned} \quad (3.41)$$

Thus, it remains to prove that $\partial_z \tilde{f} \in C([-1, 0]; H^{-\frac{1}{2}}(\mathbb{R}^d))$. Setting $\Xi(x, z) = -\nabla_x \varrho \cdot \nabla_x \tilde{f} + \frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho} \partial_z \tilde{f}$ we find that $\partial_z \Xi$ is a divergence

$$\partial_z \Xi = -\operatorname{div}_x(\partial_z \varrho \nabla_x \tilde{f} - \nabla_x \varrho \partial_z \tilde{f}) + \operatorname{div}_{x,z}(\mathcal{A} \nabla_{x,z} \tilde{f}).$$

Consequently,

$$\begin{aligned} \|\partial_z \Xi\|_{L^2([-1,0]; H^{-1})} &\leq \mathcal{F}(\|\eta\|_{H^s})(\|\nabla_{x,z} \tilde{f}\|_{L^2([-1,0]; L^2)} \\ &\quad + \|\operatorname{div}_{x,z}(\mathcal{A} \nabla_{x,z} \tilde{f})\|_{L^2([-1,0]; H^{-1})}). \end{aligned}$$

On the other hand, using Lemma 3.9, it is easy to see that

$$\|\Xi\|_{L^2([-1,0]; L^2)} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\nabla_{x,z} \tilde{f}\|_{L^2([-1,0]; L^2)}.$$

Then, applying Theorem A.6 we obtain that $\Xi \in C([-1, 0]; H^{-\frac{1}{2}}(\mathbb{R}^d))$ and

$$\begin{aligned} \|\Xi\|_{C([-1,0]; H^{-\frac{1}{2}})} &\leq \mathcal{F}(\|\eta\|_{H^s})(\|\nabla_{x,z} \tilde{f}\|_{L^2([-1,0]; L^2)} \\ &\quad + \|\operatorname{div}_{x,z}(\mathcal{A} \nabla_{x,z} \tilde{f})\|_{L^2([-1,0]; H^{-1})}). \end{aligned}$$

Now from the definition of Ξ we have

$$\partial_z \tilde{f} = \frac{\partial_z \varrho}{1 + |\nabla_x \varrho|^2} (\Xi(x, z) + \nabla_x \varrho \cdot \nabla_x \tilde{f}).$$

For $s > 1 + \frac{d}{2} \geq \frac{3}{2}$, using the product rule (C.12) and the nonlinear estimate (C.13) gives

$$\begin{aligned} &\left\| \frac{\partial_z \varrho}{1 + |\nabla_x \varrho|^2} \Xi \right\|_{C([-1,0]; H^{-\frac{1}{2}})} \\ &\lesssim \left(\left\| \frac{\partial_z \varrho}{1 + |\nabla_x \varrho|^2} - h \right\|_{C([-1,0]; H^{s-1})} + h \right) \|\Xi\|_{C([-1,0]; H^{-\frac{1}{2}})} \\ &\leq \mathcal{F}(\|\eta\|_{H^s}) (\|\nabla_{x,z} \tilde{f}\|_{L^2([-1,0]; L^2)} + \|\operatorname{div}_{x,z}(\mathcal{A} \nabla_{x,z} \tilde{f})\|_{L^2([-1,0]; H^{-1})}) \end{aligned}$$

and

$$\begin{aligned} &\left\| \frac{\partial_z \varrho \nabla_x \varrho}{1 + |\nabla_x \varrho|^2} \cdot \nabla_x \tilde{f} \right\|_{C([-1,0]; H^{-\frac{1}{2}})} \\ &\lesssim \left\| \frac{\partial_z \varrho \nabla_x \varrho}{1 + |\nabla_x \varrho|^2} \right\|_{C([-1,0]; H^{s-1})} \|\nabla_x \tilde{f}\|_{C([-1,0]; H^{-\frac{1}{2}})} \\ &\leq \mathcal{F}(\|\eta\|_{H^s}) \|\nabla_{x,z} \tilde{f}\|_{L^2([-1,0]; L^2)}, \end{aligned}$$

where (3.41) was used in the last estimate. This finishes the proof. \square

Denote $v(x, z) = \phi(x, \varrho(x, z)) : \mathbb{R}^d \times [-1, 0] \rightarrow \mathbb{R}$ where ϕ is the solution of (3.2). Then v satisfies $v|_{z=0} = f$ and

$$\operatorname{div}_{x,z}(\mathcal{A} \nabla_{x,z} v) = 0, \tag{3.42}$$

while, by the chain rule,

$$G^-(\eta) f = \left(\frac{1 + |\nabla_x \varrho|^2}{\partial_z \varrho} \partial_z v - \nabla_x \varrho \cdot \nabla_x v \right)_{|z=0}. \tag{3.43}$$

Expanding (3.42) yields

$$(\partial_z^2 + \alpha \Delta_x + \beta \cdot \nabla_x \partial_z - \gamma \partial_z)v = 0 \quad \text{in } \mathbb{R}^d \times [-1, 0], \quad (3.44)$$

where

$$\alpha = \frac{(\partial_z \rho)^2}{1 + |\nabla_x \rho|^2}, \quad \beta = -2 \frac{\partial_z \rho \nabla_x \rho}{1 + |\nabla_x \rho|^2}, \quad \gamma = \frac{1}{\partial_z \rho} (\partial_z^2 \rho + \alpha \Delta_x \rho + \beta \cdot \nabla_x \partial_z \rho). \quad (3.45)$$

Note that the restriction to $z \in [-1, 0]$ guarantees that ρ is smooth in z . We have the following Sobolev estimates for the inhomogeneous version of (3.44):

Proposition 3.12. ([3, Proposition 3.16]) *Let $d \geq 1$, $s > 1 + \frac{d}{2}$ and $\frac{1}{2} \leq \sigma \leq s$. Consider $f \in H^{1,\sigma}(\mathbb{R}^d)$ and $\eta \in H^s(\mathbb{R}^d)$ satisfying $\text{dist}(\eta, \Gamma^-) \geq h > 0$. Assume that $F_0 \in Y^{\sigma-1}([z_1, 0])$, $z_1 \in (-1, 0)$ and v a solution of*

$$(\partial_z^2 + \alpha \Delta_x + \beta \cdot \nabla_x \partial_z - \gamma \partial_z)v = F_0 \quad \text{in } \mathbb{R}^d \times [-1, 0] \quad (3.46)$$

with $v|_{z=0} = f$. If $z_0 \in (z_1, 0)$ and

$$\nabla_{x,z} v \in X^{-\frac{1}{2}}([z_0, 0]) \quad (3.47)$$

then $\nabla_{x,z} v \in X^{\sigma-1}([z_0, 0])$ and

$$\begin{aligned} \|\nabla_{x,z} v\|_{X^{\sigma-1}([z_0, 0])} &\leq \mathcal{F}(\|\eta\|_{H^s}) \\ &\left(\|\nabla_x f\|_{H^{\sigma-1}} + \|F_0\|_{Y^{\sigma-1}([z_1, 0])} + \|\nabla_{x,z} v\|_{X^{-\frac{1}{2}}([z_0, 0])} \right) \end{aligned}$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (σ, s, h, z_0, z_1) .

Remark 3.13. In fact, Proposition 3.16 in [3] assumes $f \in H^\sigma(\mathbb{R}^d)$. This comes from estimating v solving

$$\partial_z v + T_A v = -w \quad z \in [z_0, 0], \quad v|_{z=0} = f$$

where $w \in X^\sigma([z_0, 0])$ and $A \in \Gamma_\varepsilon^1$, $\varepsilon \in (0, \max\{\frac{1}{2}, s - 1 - \frac{d}{2}\})$, is given by (3.61) below. To obtain estimates involving only $\|\nabla_x f\|_{H^{\sigma-1}}$, it suffices to differentiate this equation in x and apply Proposition C.6 to control $T_{\nabla_x A} v$.

In the rest of this subsection, we fix $s > 1 + \frac{d}{2}$. For $v(x, z) = \phi(x, \rho(x, z))$, ϕ solution of (3.2), Lemma 3.11 combined with (3.42) yields

$$\begin{aligned} \|\nabla_{x,z} v\|_{X^{-\frac{1}{2}}([-1, 0])} &\leq \mathcal{F}(\|\eta\|_{H^s}) \|\nabla_{x,z} v\|_{L^2([-1, 0]; L^2(\mathbb{R}^d))} \\ &\leq \mathcal{F}(\|\eta\|_{H^s}) \|\nabla_{x,y} \phi\|_{L^2(\Omega^-)}. \end{aligned}$$

In conjunction with (3.17) and (3.21), this implies

$$\|\nabla_{x,z} v\|_{X^{-\frac{1}{2}}([-1, 0])} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^{\frac{1}{2}}}. \quad (3.48)$$

This verifies condition (3.47) of Proposition 3.12 from which the estimate for $\|\nabla_{x,z} v\|_{X^{\sigma-1}}$, $\sigma \in [\frac{1}{2}, s]$, follows. Using this and the product rule (C.12) one can easily deduce the continuity of $G^-(\eta)$ in higher Sobolev norms.

Theorem 3.14. ([3, Theorem 3.12]) Let $d \geq 1$, $s > 1 + \frac{d}{2}$ and $\frac{1}{2} \leq \sigma \leq s$. Consider $f \in \tilde{H}_-^\sigma$ and $\eta \in H^s(\mathbb{R}^d)$ with $\text{dist}(\eta, \Gamma^-) \geq h > 0$. Then we have $G^-(\eta)f \in H^{\sigma-1}(\mathbb{R}^d)$, together with the estimate

$$\|G^-(\eta)f\|_{H^{\sigma-1}} \leq \mathcal{F}(\|\eta\|_{H^s})\|f\|_{\tilde{H}_-^\sigma} \quad (3.49)$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (s, σ, h) .

Remark 3.15. Theorem 3.14 was proved in [3] for $f \in H^\sigma(\mathbb{R}^d)$. The two-phase Muskat problem involves $G^-(\eta)f^-$ where f^- is obtained from system (2.5). In particular, f^- is only determined up to an additive constant.

3.2. Paralinearization with Tame Error Estimate

The principal symbol of the Dirichlet–Neumann operator is given by

$$\lambda(x, \xi) = \sqrt{(1 + |\nabla \eta(x)|^2)|\xi|^2 - (\nabla \eta(x) \cdot \xi)^2}. \quad (3.50)$$

Note that when $d = 1$, (3.50) reduces to $\lambda(x, \xi) = |\xi|$.

We first recall a paralinearization result from [3].

Theorem 3.16. ([3, Proposition 3.13]) Let $r > 1 + \frac{d}{2}$ with $d \geq 1$, and let $\delta \in (0, \frac{1}{2}]$ satisfy $\delta < r - 1 - \frac{d}{2}$. Let $\sigma \in [\frac{1}{2}, r - \delta]$. If $\eta \in H^r(\mathbb{R}^d)$ and $f \in \tilde{H}_-^\sigma$ with $\text{dist}(\eta, \Gamma^-) > h > 0$, then we have

$$G^-(\eta)f = T_\lambda f + R_0^-(\eta)f, \quad (3.51)$$

$$\|R_0^-(\eta)f\|_{H^{\sigma-1+\delta}} \leq \mathcal{F}(\|\eta\|_{H^r})\|f\|_{\tilde{H}_-^\sigma} \quad (3.52)$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (r, σ, δ, h) .

Remark 3.17. In the statement of Proposition 3.1.3 in [3], $\sigma \in [\frac{1}{2}, r - \frac{1}{2}]$. However, its proof (see page 116) allows for $\sigma \in [\frac{1}{2}, r - \delta]$.

Our goal in this subsection is to prove the next theorem, which isolates the main term in the Dirichlet–Neumann operator as an operator and which will be the key ingredient for obtaining a priori estimates for the Muskat problem in any subcritical Sobolev regularity.

Theorem 3.18. Let $s > 1 + \frac{d}{2}$ with $d \geq 1$, and let $\delta \in (0, \frac{1}{2}]$ satisfy $\delta < s - 1 - \frac{d}{2}$. For any $\sigma \in [\frac{1}{2}, s]$, if $\eta \in H^{s+\frac{1}{2}-\delta}(\mathbb{R}^d)$ and $f \in \tilde{H}_-^\sigma$ satisfies $\text{dist}(\eta, \Gamma^-) > h > 0$ then

$$G^-(\eta)f = T_\lambda(f - T_B\eta) - T_V \cdot \nabla \eta + R^-(\eta)f, \quad (3.53)$$

where

$$B = \frac{\nabla \eta \cdot \nabla f + G^-(\eta)f}{1 + |\nabla \eta|^2}, \quad V = \nabla f - B \nabla \eta \quad (3.54)$$

and the remainder $R^-(\eta)f$ satisfies

$$\|R^-(\eta)f\|_{H^{\sigma-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s})(1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}})\|f\|_{\tilde{H}_-^\sigma}. \quad (3.55)$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (s, σ, δ, h) . In fact, $B = (\partial_y \phi)|_{y=\eta(x)}$ and $V = (\nabla_x \phi)|_{y=\eta(x)}$ where ϕ is the solution of (3.2).

Remark 3.19. For $s > 1 + \frac{d}{2}$, we can apply Theorem 3.16 with $r = s + \frac{1}{2}$, $\sigma = s$ and $\delta = \frac{1}{2}$ (the maximal value allowed) to have

$$\|R_0^-(\eta)f\|_{H^{s-\frac{1}{2}}} \leq \mathcal{F}\left(\|\eta\|_{H^{s+\frac{1}{2}}}\right) \|f\|_{\tilde{H}_-^s}. \quad (3.56)$$

Both (3.56) and (3.55) provide a gain of $\frac{1}{2}$ derivative for f . The improvement of (3.55) is in that (1) there is a gain of δ derivative for η ; (2) the highest norm $\|\eta\|_{H^{s+\frac{1}{2}-\delta}}$ of η appears linearly. For the sake of a priori estimates, (1) gives room to choose the time of existence T as a small parameter; (2) is required to gain $\frac{1}{2}$ derivative using the parabolicity when measured in L^2 in time.

We fix $s > 1 + \frac{d}{2}$ in the rest of this subsection. Setting

$$Q = \partial_z^2 + \alpha \Delta_x + \beta \cdot \nabla_x \partial_z, \quad (3.57)$$

we can rewrite (3.44) as

$$Qv = \frac{\partial_z v}{\partial_z \varrho} Q\varrho. \quad (3.58)$$

The coefficients of Q can easily be controlled using (3.45), (3.28) and Lemma 3.9:

$$\begin{aligned} & \|\alpha - h^2\|_{X^{r-1}([-1,0])} + \|\beta\|_{X^{r-1}([-1,0])} + \|\gamma\|_{X^{r-2}([-1,0])} \\ & \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^r} \quad \forall r \geq s \end{aligned} \quad (3.59)$$

since it follows from (3.28) that

$$\|(\nabla_x \varrho, \partial_z \varrho - h)\|_{X^{\mu-1}([-1,0])} \leq C \|\eta\|_{H^\mu} \quad \forall \mu \in \mathbb{R}. \quad (3.60)$$

We start with a factorization of Q by paradifferential operators and a remainder:

Lemma 3.20. *With the symbols*

$$\begin{aligned} a &= \frac{1}{2} \left(-i\beta \cdot \xi - \sqrt{4\alpha|\xi|^2 - (\beta \cdot \xi)^2} \right), \\ A &= \frac{1}{2} \left(-i\beta \cdot \xi + \sqrt{4\alpha|\xi|^2 - (\beta \cdot \xi)^2} \right), \end{aligned} \quad (3.61)$$

we define R_Q by

$$R_Q g = Qg - (\partial_z - T_a)(\partial_z - T_A)g. \quad (3.62)$$

Let $0 < \delta \leq 1$ satisfy $\delta < s - 1 - \frac{d}{2}$. If θ satisfies

$$\theta \leq s - \delta, \quad \theta + s > 1 + \delta, \quad (3.63)$$

then, for any $z_0 \in (-1, 0)$ we have

$$\|R_Q g\|_{L^2([z_0, 0]; H^\theta)} \leq \mathcal{F}(\|\eta\|_{H^s}) \left(1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}} \right) \|\nabla_{x,z} g\|_{X^\theta([z_0, 0])}. \quad (3.64)$$

On the other hand, if θ satisfies

$$\theta \leq s, \quad \theta + s > 1 + \delta, \quad (3.65)$$

$$\|R_Q g\|_{Y^\theta([z_0, 0])} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\nabla_{x,z} g\|_{X^{\theta-\delta}([z_0, 0])}. \quad (3.66)$$

Proof. From (3.61) we have that $a + A = -i\beta \cdot \xi$, $aA = -\alpha|\xi|^2$, and hence

$$\partial_z^2 + T_\alpha \Delta_x + T_\beta \cdot \nabla_x \partial_z = (\partial_z - T_a)(\partial_z - T_A)g - (T_a T_A - T_\alpha \Delta_x)g + T_{\partial_z A}g.$$

It follows that

$$R_Q g = (\alpha - T_\alpha) \Delta_x g + (\beta - T_\beta) \cdot \nabla_x \partial_z g - (T_a T_A - T_\alpha \Delta_x)g + T_{\partial_z A}g, \quad (3.67)$$

Proof of (3.64). Assuming (3.63), we claim that

$$\begin{aligned} & \|(\alpha - T_\alpha) \Delta_x g + (\beta - T_\beta) \cdot \nabla_x \partial_z g\|_{L^2([z_0, 0]; H^\theta)} \\ & \leq \mathcal{F}(\|\eta\|_{H^s}) (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|\nabla_{x,z} g\|_{L^\infty([z_0, 0]; H^\theta)}. \end{aligned} \quad (3.68)$$

Using (C.11), (3.59) and (3.63), we have

$$\begin{aligned} \|(\beta - \beta) \nabla_x \partial_z g\|_{L^2 H^\theta} & \lesssim \|\beta\|_{L^2 H^{s-\delta}} \|\nabla_x \partial_z g\|_{L^\infty H^{\theta-1}} \lesssim \|\beta\|_{L^2 H^{s-\delta}} \|\partial_z g\|_{L^\infty H^\theta} \\ & \leq \mathcal{F}(\|\eta\|_{H^s}) (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|\partial_z g\|_{L^\infty H^\theta}. \end{aligned}$$

As for $(\alpha - T_\alpha) \Delta_x g$, we write

$$(\alpha - T_\alpha) \Delta_x g = (\alpha - h^2 - T_{\alpha-h^2}) \Delta_x g + h^2 (\text{Id} - T_1) \Delta_x g.$$

The first term can be estimated as above and in view of (C.2), $\text{Id} - T_1 = \text{Id} - \Psi(D)$ is a smoothing operator so that

$$\|(\text{Id} - T_1) \Delta_x g\|_{L^2 H^\theta} \leq C \|\nabla_x g\|_{L^2 H^\theta}.$$

We thus obtain (3.68).

Next it is readily seen that A and a satisfy (see (C.1))

$$M_\delta^1(a; [-1, 0]) + M_\delta^1(A; [-1, 0]) \leq \mathcal{F}(\|\eta\|_{H^s}) \quad (3.69)$$

$$M_{\frac{1}{2}}^1(a; [-1, 0]) + M_{\frac{1}{2}}^1(A; [-1, 0]) \leq \mathcal{F}(\|\eta\|_{H^s}) \left(1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}\right) \quad (3.70)$$

Consequently, by Theorem C.4 (ii), $T_a T_A - T_\alpha \Delta_x$ is of order $\frac{3}{2}$ and

$$\|(T_a T_A - T_\alpha \Delta_x)g\|_{L^2 H^\theta} \leq \mathcal{F}(\|\eta\|_{H^s}) \left(1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}\right) \|\nabla_x g\|_{L^2 H^{\theta+\frac{1}{2}}}, \quad (3.71)$$

where Remark C.5 has been used. Now in view of the seminorm bounds

$$\|M_0^1(\partial_z A)\|_{L^2([-1, 0])} \leq \mathcal{F}(\|\eta\|_{H^s}) \left(1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}\right),$$

Theorem C.4 (i) combined with Remark C.5 gives

$$\|T_{\partial_z A} g\|_{L^2 H^\theta} \leq \mathcal{F}(\|\eta\|_{H^s}) \left(1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}\right) \|\nabla_x g\|_{L^\infty H^\theta}. \quad (3.72)$$

From (3.68), (3.71) and (3.72), the proof of (3.64) is complete.

Proof of (3.66). Assume (3.65). Using (C.8), (C.9) and (3.59), we have

$$\begin{aligned} \|T_{\nabla_x \partial_z g} \cdot \beta\|_{L^2 H^{\theta-\frac{1}{2}}} & \lesssim \|\nabla_x \partial_z g\|_{L^\infty H^{\theta-1-\delta}} \|\beta\|_{L^2 H^{s-\frac{1}{2}}} \\ & \leq \mathcal{F}(\|\eta\|_{H^s}) \|\partial_z g\|_{L^\infty H^{\theta-\delta}} \end{aligned}$$

and

$$\begin{aligned}\|R(\nabla_x \partial_z g, \beta)\|_{L^1 H^\theta} &\lesssim \|\beta\|_{L^2 H^{s-\frac{1}{2}}} \|\nabla_x \partial_z g\|_{L^2 H^{\theta-\frac{1}{2}-\delta}} \\ &\leq \mathcal{F}(\|\eta\|_{H^s}) \|\partial_z g\|_{L^2 H^{\theta+\frac{1}{2}-\delta}}.\end{aligned}$$

The term $(T_\alpha - \alpha) \Delta_x g$ can be treated similarly. Next using (3.69) and Theorem C.4 (ii) we find that $T_a T_A - T_\alpha \Delta_x$ is of order $2 - \delta$ and

$$\|(T_a T_A - T_\alpha \Delta_x)g\|_{L^2 H^{\theta-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\nabla_x g\|_{L^2 H^{\theta+\frac{1}{2}-\delta}}. \quad (3.73)$$

As for $T_{\partial_z} A g$, we note that since

$$\|(\alpha - h^2, \beta)\|_{L^\infty([-1,0]; H^{s-1})} + \|(\partial_z \alpha, \partial_z \beta)\|_{L^\infty([-1,0]; H^{s-2})} \leq \mathcal{F}(\|\eta\|_{H^s})$$

and $H^{s-2}(\mathbb{R}^d) \subset C_*^{-1+\delta}(\mathbb{R}^d)$, we have

$$M_{-1+\delta}^1(\partial_z A) \leq \mathcal{F}(\|\eta\|_{H^s}).$$

By virtue of Proposition C.6, $T_{\partial_z} A$ is of order $2 - \delta$ and

$$\|T_{\partial_z} A g\|_{L^2 H^{\theta-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\nabla_x g\|_{L^2 H^{\theta+\frac{1}{2}-\delta}}.$$

This completes the proof of (3.66). \square

We can now start analyzing (3.58). We fix $\sigma \in [\frac{1}{2}, s]$ and apply Proposition 3.12 to have

$$\|\nabla_{x,z} v\|_{X^{\sigma-1}([z_0,0])} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma} \quad (3.74)$$

for all $z_0 \in (-1, 0)$.

Next we introduce

$$\mathfrak{b} = \frac{\partial_z v}{\partial_z \varrho}, \quad u = v - T_{\mathfrak{b}} \varrho. \quad (3.75)$$

We note that $\mathfrak{b}|_{z=0} = (\partial_y \phi)(x, \eta(x)) = B$ given by (3.54), and $u|_{z=0} = f - T_B \eta$. The new variable u is known as the “good unknown” à la Alinhac. Fixing $\delta \in (0, \frac{1}{2}]$ satisfying $\delta < s - 1 - \frac{d}{2}$, for all $\sigma \in [\frac{1}{2}, s]$ we have

$$\sigma + s > 2 + \delta. \quad (3.76)$$

Lemma 3.21. *Let $z_0 \in (-1, 0)$. For $\sigma \in [\frac{1}{2}, s]$ we have*

$$\|\mathfrak{b}\|_{L^\infty([z_0,0]; H^{\sigma-1})} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma}, \quad (3.77)$$

$$\|\nabla_{x,z} \mathfrak{b}\|_{L^2([z_0,0]; H^{\sigma-\frac{3}{2}})} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma} \quad \text{if } \sigma + s > \frac{5}{2}, \quad (3.78)$$

$$\|\nabla_{x,z} \mathfrak{b}\|_{L^\infty([z_0,0]; H^{\sigma-2})} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma} \quad \text{if } \sigma + s > 3, \quad (3.79)$$

$$\|\partial_z \mathfrak{b}\|_{Y^{\sigma-1}([z_0,0])} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma}. \quad (3.80)$$

Proof. (3.77) follows from (3.60), (3.74) and the product rule (C.12) with $s_0 = s_1 = \sigma - 1$ and $s_2 = s - 1$ since $\sigma - 1 + s - 1 > 0$ in view of (3.76).

The estimate (3.78) for $\nabla_x b$ can be proved similarly upon using (C.12) with $s_0 = s_1 = \sigma - \frac{3}{2}$, $s_2 = s - 1$ and noting that guarantees $s_1 + s_2 = \sigma + s - \frac{5}{2} > 0$. As for $\partial_z b$ we use (3.44) to have the formula for $\partial_z^2 v$, then apply (C.12) as in the estimate for $\nabla_x b$.

The proof of (3.79) is similar to (3.78): we apply (C.12) with $s_0 = s_1 = \sigma - 2$, $s_2 = s - 1$ and note that $s_1 + s_2 > 0$ if $\sigma + s > 3$.

Let us prove (3.80). We first compute using (3.44) that

$$\begin{aligned}\partial_z b &= -\partial_z v \frac{\partial_z^2 \varrho}{(\partial_z \varrho)^2} + \frac{\partial_z^2 v}{\partial_z \varrho} \\ &= -\partial_z v \frac{\partial_z^2 \varrho}{(\partial_z \varrho)^2} - \frac{\alpha}{\partial_z \varrho} \Delta_x v - \frac{\beta}{\partial_z \varrho} \cdot \nabla_x \partial_z v + \partial_z v \frac{\gamma}{\partial_z \varrho}.\end{aligned}$$

For the first and last terms, we apply (3.34) with $s_0 = \sigma - 1$, $s_1 = \sigma - 1$ and $s_2 = s - 2$, giving

$$\|\partial_z v \frac{\partial_z^2 \varrho}{(\partial_z \varrho)^2}\|_{X^{\sigma-1}} \lesssim \|\partial_z v\|_{X^{\sigma-1}} \|\frac{\partial_z^2 \varrho}{(\partial_z \varrho)^2}\|_{X^{s-2}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma}.$$

Next for the second and third terms, applying (3.34) with $s_0 = \sigma - 1$, $s_1 = s - 1$ and $s_2 = \sigma - 2$ yields

$$\|\frac{\beta}{\partial_z \varrho} \cdot \nabla_x \partial_z v\|_{Y^{\sigma-1}} \lesssim \|\frac{\beta}{\partial_z \varrho}\|_{X^{s-1}} \|\nabla_x \partial_z v\|_{X^{\sigma-2}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma}.$$

This finishes the proof of (3.80). \square

We can now state our main technical estimate.

Lemma 3.22. *For any $z_0 \in (-1, 0)$ and $\sigma \in [\frac{1}{2}, s]$, we have*

$$(\partial_z - T_a)[(\partial_z - T_A)v - T_b(\partial_z - T_A)\varrho] = F_2, \quad (3.81)$$

$$\|F_2\|_{Y^{\sigma-\frac{1}{2}}([z_0, 0])} \leq \mathcal{F}(\|\eta\|_{H^s}) (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|f\|_{\tilde{H}_-^\sigma}. \quad (3.82)$$

Remark 3.23. The direct consideration of the good unknown $u = v - T_b \varrho$ in [2, 31] consists in obtaining good estimates for

$$(\partial_z - T_a)(\partial_z - T_A)u.$$

In our setting, even when $\sigma = s$, estimating this in $Y^{s-\frac{1}{2}}$ demands an estimate for $\|\partial_z^2 b\|_{X^{s-3}}$. However, in one space dimension, the low regularity $s > \frac{3}{2}$ makes it challenging to prove that $\partial_z^3 v \in X^{s-3}$, where $\partial_z^3 v$ appears when differentiating b twice in z . Lemma 3.22 avoids this issue.

Proof of Lemma 3.22. Using (3.58) and Lemma 3.20 with $\theta = \sigma - 1$, we see that

$$\begin{aligned} (\partial_z - T_a)(\partial_z - T_A)v + R_Q v &= T_b(\partial_z - T_a)(\partial_z - T_A)\varrho \\ &\quad + (\mathfrak{b}Q\varrho - T_bQ\varrho) + T_bR_Q\varrho \end{aligned}$$

which gives

$$F_2 = [T_b, (\partial_z - T_a)](\partial_z - T_A)\varrho - R_Q v + T_bR_Q\varrho + (\mathfrak{b}Q\varrho - T_bQ\varrho).$$

It is readily checked that

$$\|Q\varrho\|_{L^2 H^{s-1-\delta}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^{s+\frac{1}{2}-\delta}}$$

which in conjunction with (C.11) and (3.77) yields

$$\begin{aligned} \|\mathfrak{b}Q\varrho - T_bQ\varrho\|_{L^2 H^{\sigma-1}} &\lesssim \|\mathfrak{b}\|_{L^\infty H^{\sigma-1}} \|Q\varrho\|_{L^2 H^{s-1-\delta}} \\ &\leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^{s+\frac{1}{2}-\delta}} \|f\|_{\tilde{H}_-^\sigma}, \end{aligned}$$

where we have used (3.76) in the first inequality.

In view of (3.76), (3.64) can be applied with $\theta = \sigma - 1$, implying the control of $R_Q v$. As for $T_bR_Q\varrho$ we apply (C.8), (3.64) with $\theta = s - 1$, and (3.77) to have

$$\begin{aligned} \|T_bR_Q\varrho\|_{L^2 H^{\sigma-1}} &\lesssim \|\mathfrak{b}\|_{L^\infty H^{\sigma-1}} \|R_Q\varrho\|_{L^2 H^{s-1}} \\ &\leq \mathcal{F}(\|\eta\|_{H^s}) (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|f\|_{\tilde{H}_-^\sigma}. \end{aligned}$$

Regarding the commutator in F_2 , we write

$$[T_b, (\partial_z - T_a)](\partial_z - T_A)\varrho = -T_{\partial_z} \mathfrak{b}(\partial_z - T_A)\varrho - [T_b, T_a](\partial_z - T_A)\varrho.$$

For $T_{\partial_z} \mathfrak{b}(\partial_z - T_A)\varrho$ we distinguish two cases.

Cases 1 $\sigma \in [s - \delta, s]$. Then, $\sigma + s \geq 2s - \delta > \frac{5}{2}$ and (3.78) can be applied. Noting in addition that $\sigma - 1 \leq s - 1 \leq s - \frac{1}{2} - \delta$, (C.8) yields

$$\begin{aligned} \|T_{\partial_z} \mathfrak{b}(\partial_z - T_A)\varrho\|_{L^2 H^{\sigma-1}} &\lesssim \|\partial_z \mathfrak{b}\|_{L^2 H^{\sigma-\frac{3}{2}}} \|(\partial_z - T_A)\varrho - h\|_{L^\infty H^{s-\frac{1}{2}-\delta}} \\ &\leq \mathcal{F}(\|\eta\|_{H^s}) (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|f\|_{\tilde{H}_-^\sigma}. \end{aligned}$$

Cases 2 $\sigma \in [\frac{1}{2}, s - \delta]$. In view of (3.80), applying (3.38) we obtain

$$\begin{aligned} \|T_{\partial_z} \mathfrak{b}(\partial_z - T_A)\varrho\|_{Y^{\sigma-\frac{1}{2}}} &\lesssim \|\partial_z \mathfrak{b}\|_{Y^{\sigma-1}} \|(\partial_z - T_A)\varrho - h\|_{L^\infty H^{s-\frac{1}{2}-\delta}} \\ &\leq \mathcal{F}(\|\eta\|_{H^s}) (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|f\|_{\tilde{H}_-^\sigma}. \end{aligned}$$

To treat the commutator $[T_a, T_b]$ we again distinguish two cases.

Case 1 $\sigma \in (s - \delta, s]$. Then we have $\nu := \sigma + \delta - s \in (0, \frac{1}{2}]$ since $\delta \leq \frac{1}{2}$ and $\sigma \leq s$. In addition, $\sigma - 1 - \frac{d}{2} > \nu$ and $s - 1 - \frac{d}{2} > \nu$, implying that $\mathfrak{b} \in H^{\sigma-1} \subset W^{\nu, \infty}$ and $a \in \Gamma_\nu^1$ uniformly in z . Consequently, by virtue of Theorem C.4, $[T_a, T_b]$ is of order $1 - \nu$ and

$$\|[T_b, T_a](\partial_z - T_A)\varrho\|_{L^2 H^{\sigma-1}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma} \|(\partial_z - T_A)\varrho - h\|_{L^2 H^{\sigma-\nu}}$$

$$\begin{aligned}
&= \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma} \|(\partial_z - T_A)\rho - h\|_{L^2 H^{s-\delta}} \\
&\leq \mathcal{F}(\|\eta\|_{H^s}) (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|f\|_{\tilde{H}_-^\sigma},
\end{aligned}$$

where we have used (3.77) in the first inequality.

Case 2 $\sigma \in [\frac{1}{2}, s - \delta]$. In this case we do not use the structure of the commutator but directly estimate using Theorem C.4 (i) and (C.8):

$$\begin{aligned}
\|T_a T_b (\partial_z - T_A) \varrho\|_{L^2 H^{\sigma-1}} &\leq \mathcal{F}(\|\eta\|_{H^s}) \|T_b (\partial_z - T_A) \varrho\|_{L^2 H^\sigma} \\
&\leq \mathcal{F}(\|\eta\|_{H^s}) \|\mathfrak{b}\|_{L^\infty H^{\sigma-1}} \|(\partial_z - T_A) \varrho - h\|_{L^2 H^{s-\delta}} \\
&\leq \mathcal{F}(\|\eta\|_{H^s}) (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \|f\|_{\tilde{H}_-^\sigma},
\end{aligned}$$

where in the second inequality we have used the fact that $\sigma \leq s - \delta$. The term $T_b T_a (\partial_z - T_A) \varrho$ can be controlled similarly. This completes the proof of Lemma 3.22. \square

Proof of Theorem 3.18. The proof proceeds in two steps.

Step 1 Let us fix $-1 < z_0 < z_1 < 0$ and introduce a cut-off χ satisfying $\chi(z) = 1$ for $z > z_1$ and $= 0$ for $z < z_0$. Set

$$w = \chi(z) [(\partial_z - T_A)v - T_b (\partial_z - T_A)\varrho].$$

It follows from (3.81) that

$$(\partial_z - T_a)w = F_3 := \chi(z)F_2 + \chi'(z)[(\partial_z - T_A)v - T_b (\partial_z - T_A)\varrho]. \quad (3.83)$$

By virtue of (3.82), (3.74), (3.77) and (3.69) we have

$$\|F_3\|_{Y^{\sigma-\frac{1}{2}}([z_0, 0])} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma} (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}). \quad (3.84)$$

Next we note that

$$\operatorname{Re}(-a) \geq \frac{1}{\mathcal{F}(\|\eta\|_{H^s})} |\xi|.$$

Since $w(z_0) = 0$, applying Proposition C.12 to equation (3.83) with the aid of (3.84) we obtain

$$\begin{aligned}
\|w\|_{X^{\sigma-\frac{1}{2}}([z_0, 0])} &\leq \mathcal{F}(\|\eta\|_{H^s}) \|F_3\|_{Y^{\sigma-\frac{1}{2}}([z_0, 0])} \\
&\leq \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma} (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}).
\end{aligned} \quad (3.85)$$

Step 2 Starting from (3.43) and using Bony's decomposition, we find that

$$\begin{aligned}
G^-(\eta) f &= T_{\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho}} (T_A v - T_b T_A \varrho) - \left(T_{\nabla_x \varrho} \nabla_x v - T_{\partial_z v} T_{\frac{\nabla_x \varrho}{\partial_z \varrho}} \nabla_x \varrho \right) \\
&\quad - \left(T_{\nabla_x v} - T_{\partial_z v} T_{\frac{\nabla_x \varrho}{\partial_z \varrho}} \right) \nabla_x \varrho \\
&\quad + \left\{ T_{\partial_z v} \left(\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho} \right) + T_{\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho}} T_b \partial_z \varrho - 2 T_{\partial_z v} T_{\frac{\nabla_x \varrho}{\partial_z \varrho}} \nabla_x \varrho \right\} \\
&\quad + T_{\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho}} w - R(\nabla_x \varrho, \nabla_x v) + R\left(\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho}, \partial_z v\right),
\end{aligned}$$

where the right-hand side is evaluated at $z = 0$. We will see that this gives (3.53) by estimating each term one by one.

Using Theorem C.4, (3.69), (3.70) and (3.74), we first observe that

$$R^1 = \left\{ T_{\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho}} (T_A v - T_b T_A \varrho) - (T_{\nabla_x \varrho} \nabla_x v - T_{\partial_z v} T_{\frac{\nabla_x \varrho}{\partial_z \varrho}} \nabla_x \varrho) \right\} \\ - T_{\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho}} A - i \xi \cdot \nabla_x \varrho \underbrace{(v - T_b \rho)}_{=u}$$

satisfies estimates as in (3.55). Using the formula (3.50) and (3.61), we see that

$$A(x, \xi) \frac{1 + |\nabla_x \varrho|^2}{\partial_z \varrho} - i \xi \cdot \nabla_x \varrho|_{z=0} = \lambda, \quad (3.86)$$

and this gives the first main term in (3.53). Similarly, we obtain that

$$R^2 = - \left(T_{\nabla_x v} - T_{\partial_z v} T_{\frac{\nabla_x \varrho}{\partial_z \varrho}} \right) \nabla_x \varrho + T_{\nabla_x v} - b \nabla_x \varrho \nabla_x \varrho \\ = \left(T_{\partial_z v} T_{\frac{\nabla_x \varrho}{\partial_z \varrho}} - T_{\frac{\partial_z v \nabla_x \varrho}{\partial_z \varrho}} \right) \nabla_x \varrho$$

is acceptable, and since

$$\nabla_x v - b \nabla_x \varrho|_{z=0} = \nabla f - B \nabla \eta = V,$$

we obtain the second main estimate in (3.53).

We claim that all the other terms are remainders. Next we paralinearize the function $F(m, n) = \frac{1+|m|^2}{n+h} - h^{-1}$ where $m \in \mathbb{R}^d$ and $n \in \mathbb{R}$. Clearly $F(0, 0) = 0$, $\nabla_m F = \frac{2m}{n+h}$, and $\partial_n F = -\frac{1+m^2}{(n+h)^2}$. Applying Theorem C.11 with $\mu = s - \frac{1}{2} - \delta$ and $\tau = \delta$ yields

$$\frac{1 + |\nabla_x \varrho|^2}{\partial_z \varrho} - h^{-1} = F(\nabla_x \varrho, \partial_z \varrho - h) = T_{\frac{\nabla_x \varrho}{\partial_z \varrho}} \cdot \nabla_x \varrho - T_{\frac{1+|\nabla_x \varrho|^2}{(\partial_z \varrho)^2}} (\partial_z \varrho - h) \\ + R_F(\nabla_x \varrho, \partial_z \varrho - h^{-1})$$

with

$$\|R_F(\nabla_x \varrho, \partial_z \varrho - h^{-1})\|_{H^{s-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^{s+\frac{1}{2}-\delta}}.$$

Then by virtue of Theorem C.4 (ii) with $\rho = \delta$ we obtain that

$$R^3 = T_{\partial_z v} \left(\frac{1 + |\nabla_x \varrho|^2}{\partial_z \varrho} - h^{-1} \right) - 2 T_{\partial_z v} T_{\frac{\nabla_x \varrho}{\partial_z \varrho}} \cdot \nabla_x \varrho + T_{\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho}} T_{\frac{\partial_z v}{\partial_z \varrho}} (\partial_z \varrho - h)$$

is acceptable, as in (3.55). The next term follows from (3.85). Finally, by (3.74), (C.9) we get

$$\begin{aligned} \|R(\nabla_x \varrho, \nabla_x v)\|_{L^\infty H^{\sigma-\frac{1}{2}}} &\lesssim \|\nabla_x \varrho\|_{L^\infty H^{s-\frac{1}{2}-\delta}} \|\nabla_x v\|_{L^\infty H^{\sigma-1}} \\ &\lesssim \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma} \|\eta\|_{H^{s+\frac{1}{2}-\delta}}, \end{aligned}$$

and similarly, since $\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho} - \frac{1}{h} \in L^\infty H^{s-\frac{1}{2}-\delta} \subset L^\infty C_*^{\frac{1}{2}}$, it follows from (C.10) that

$$\begin{aligned} &\|R\left(\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho}, \nabla_x v\right)\|_{L^\infty H^{\sigma-\frac{1}{2}}} \\ &\lesssim \left(\|\frac{1+|\nabla_x \varrho|^2}{\partial_z \varrho} - \frac{1}{h}\|_{L^\infty H^{s-\frac{1}{2}-\delta}} + \frac{1}{h}\right) \|\nabla_x v\|_{L^\infty H^{\sigma-1}} \\ &\lesssim \mathcal{F}(\|\eta\|_{H^s}) \|f\|_{\tilde{H}_-^\sigma} (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}). \end{aligned}$$

The proof of Theorem 3.18 is complete. \square

3.3. Contraction Estimates

In order to obtain uniqueness and stability estimates for the Muskat problem, we need contraction estimates for the Dirichlet–Neumann operator associated to two different surfaces η_1 and η_2 . Since we always assume in this subsection that $\eta_j \in L^\infty(\mathbb{R}^d)$ and $\text{dist}(\eta_j, \Gamma^-) > h > 0$, Proposition 3.2 guarantees that the spaces \tilde{H}_\pm^s , defined by (3.22)–(3.23)–(3.24), are independent of η_j . We have the following results:

Theorem 3.24. *Let $s > 1 + \frac{d}{2}$ with $d \geq 1$. Let $\delta \in (0, \frac{1}{2}]$ satisfy $\delta < s - 1 - \frac{d}{2}$. Consider $f \in \tilde{H}_-^s$ and $\eta_1, \eta_2 \in H^s(\mathbb{R}^d)$ with $\text{dist}(\eta_j, \Gamma^-) > 4h > 0$ for $j = 1, 2$. Then for any $\sigma \in [\frac{1}{2} + \delta, s]$, we have*

$$G^-(\eta_1)f - G^-(\eta_2)f = -T_{\lambda_2 B_2}(\eta_1 - \eta_2) - T_{V_2} \cdot \nabla(\eta_1 - \eta_2) + R_2^-(\eta_1, \eta_2)f \quad (3.87)$$

where

$$\|R_2^-(\eta_1, \eta_2)f\|_{H^{\sigma-1}} \leq \mathcal{F}(\|(\eta_1, \eta_2)\|_{H^s}) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s} \quad (3.88)$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (s, σ, h, δ) .

Corollary 3.25. *Let $s > 1 + \frac{d}{2}$ with $d \geq 1$. Consider $f \in \tilde{H}_-^s$ and $\eta_1, \eta_2 \in H^s(\mathbb{R}^d)$ with $\text{dist}(\eta_j, \Gamma^-) > 4h > 0$ for $j = 1, 2$. Then for all $\sigma \in [\frac{1}{2}, s]$, we have*

$$\|G^-(\eta_1)f - G^-(\eta_2)f\|_{H^{\sigma-1}} \leq \mathcal{F}(\|(\eta_1, \eta_2)\|_{H^s}) \|\eta_1 - \eta_2\|_{H^\sigma} \|f\|_{\tilde{H}_-^s} \quad (3.89)$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (s, σ, h) .

Remark 3.26. The following contraction estimate was obtained in Theorem 5.2 in [3]:

$$\|G^-(\eta_1)f - G(\eta_2)f\|_{H^{r-2}} \leq \mathcal{F}(\|(\eta_1, \eta_2)\|_{H^r}) \|\eta_1 - \eta_2\|_{H^{r-1}} \|f\|_{H^{r-\frac{1}{2}}}, \quad (3.90)$$

where $r > \frac{3}{2} + \frac{d}{2}$ ($\frac{1}{2}$ derivative above scaling). It was also noted in [3] (see Remark 5.3 therein) that the authors were unable to obtain a similar estimate in higher norms. Applying 3.89 with $s = r - \frac{1}{2}$ gives such estimates.

From now on, to simplify notation, we let

$$N_s = \|\eta_1\|_{H^s} + \|\eta_2\|_{H^s}.$$

The rest of this section is devoted to the proof of Theorem 3.24. We follow similar steps as to those in the previous section, the main novelty coming from the two different domains. To define $G(\eta_j)f$ we call ϕ_j solution to (3.2) with surface η_j and Dirichlet data f . For the sake of contraction estimates, we shall use a diffeomorphism different from the one defined by (3.26)–(3.27)–(3.28). Assume $\text{dist}(\eta_j, \Gamma^-) > 4h > 0$. There exists $\eta_* : \mathbb{R}^d \rightarrow \mathbb{R}$ such that

$$\|\eta_* + \frac{3h}{2}\|_{H^{s+100}} \leq CN_s \quad (3.91)$$

and

$$\underline{b}^-(x) + 2h < \eta_*(x) < \eta_j(x) - h \quad (3.92)$$

when the depth is finite and $\eta_*(x) < \eta_j(x) - h$ when $\Gamma^- = \emptyset$. One can take η_* to be a mollification of $\min(\eta_1(x), \eta_2(x)) - \frac{3h}{2}$. Then we divide Ω_j^- into

$$\begin{cases} \Omega_{j,1}^- = \{(x, y) : x \in \mathbb{R}^d, \eta_*(x) < y < \eta_j(x)\}, \\ \Omega_{j,2}^- = \{(x, y) \in \Omega_j^- : y \leq \eta_*(x)\} \end{cases} \quad (3.93)$$

and set $\widetilde{\Omega}^- = \widetilde{\Omega}_1^- \cup \widetilde{\Omega}_2^-$ where

$$\begin{cases} \widetilde{\Omega}_1^- = \mathbb{R}^d \times (-1, 0), \\ \widetilde{\Omega}_2^- = \begin{cases} \mathbb{R}^d \times (-\infty, -1] & \text{if } \Gamma^- = \emptyset, \\ \{(x, z) \in \mathbb{R}^d \times (-\infty, -1] : z + 1 + \eta_*(x) > \underline{b}^-(x)\} & \text{if } \underline{b}^- \in W^{1,\infty}. \end{cases} \end{cases} \quad (3.94)$$

Note that $\Omega_{1,2}^- = \Omega_{2,2}^-$ and the sets $\widetilde{\Omega}_1^-$ and $\widetilde{\Omega}_2^-$ are independent of η_j . Define

$$\begin{cases} \varrho_j(x, z) = (1 + z)e^{\tau z \langle D_x \rangle} \eta_j(x) - z\eta_*(x) & \text{for } (x, z) \in \widetilde{\Omega}_1^-, \\ \varrho_j(x, z) = z + 1 + \eta_*(x) & \text{for } (x, z) \in \widetilde{\Omega}_2^-. \end{cases} \quad (3.95)$$

In particular, $\varrho_1 = \varrho_2$ in $\widetilde{\Omega}_2^-$. For $\tau > 0$ sufficiently small, it is easy to check that the mappings $(x, z) \in \widetilde{\Omega}^- \mapsto (x, \varrho_j(x, z)) \in \Omega_j^-$ and $(x, z) \in \Omega_1^- \mapsto (x, \varrho_j(x, z)) \in \Omega_{j,1}^-$ are Lipschitz diffeomorphisms, where the latter is smooth in z . Letting also $\varrho_\delta = \varrho_1 - \varrho_2$, we observe as in (3.60) that

$$\begin{aligned} \min(\partial_z \varrho_1, \partial_z \varrho_2) &\geq \min\left(1, \frac{h}{2}\right), \\ \|\left(\nabla_x \varrho_j, \partial_z \varrho_j - \frac{3h}{2}\right)\|_{X^{s-1}([-1, 0])} &\lesssim \|\eta_1\|_{H^s} + \|\eta_2\|_{H^s}, \\ \|\varrho_\delta\|_{X^\sigma((-\infty, 0])} &\lesssim \|\eta_1 - \eta_2\|_{H^\sigma} \end{aligned} \quad (3.96)$$

if $\tau > 0$ is chosen small enough (depending on $\|\eta_1\|_{B_{\infty,1}^1} + \|\eta_2\|_{B_{\infty,1}^1}$).

As in (3.44),

$$v_j(x, z) := \phi_j(x, \varrho_j(x, z))$$

solves

$$\mathcal{L}_j v_j := (\partial_z^2 + \alpha_j \Delta_x + \beta_j \cdot \nabla_x \partial_z - \gamma_j \partial_z) v_j = 0 \quad \text{in } \mathbb{R}^d \times [-1, 0], \quad (3.97)$$

with $(\alpha_j, \beta_j, \gamma_j)$ defined in terms of ϱ_j as in (3.45) and satisfies¹

$$\|\nabla_{x,z} v_j\|_{X^{s-1}([-1, 0])} \lesssim \mathcal{F}(N_s) \|f\|_{\tilde{H}_-^s}. \quad (3.98)$$

The difference

$$v = v_1 - v_2$$

then solves

$$\begin{aligned} \mathcal{L}_1 v &= F := -(\alpha_1 - \alpha_2) \Delta_x v_2 - (\beta_1 - \beta_2) \cdot \nabla_x \partial_z v_2 \\ &\quad + (\gamma_1 - \gamma_2) \partial_z v_2 \quad \text{in } \mathbb{R}^d \times [-1, 0]. \end{aligned} \quad (3.99)$$

As before, we start with an estimate for v in the low norm $X^{-\frac{1}{2}}([-1, 0])$.

Lemma 3.27.

$$\|\nabla_{x,z} v\|_{X^{-\frac{1}{2}}([-1, 0])} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\frac{1}{2}}} \|f\|_{\tilde{H}_-^s}. \quad (3.100)$$

Proof. We first recall the variational characterization (3.20)

$$\int_{\Omega_j^-} \nabla_{x,y} \phi_j \cdot \nabla_{x,y} \varphi \, dx \, dy = 0 \quad \forall \varphi \in H_{0,*}^1(\Omega_j^-) = \{\varphi \in \dot{H}^1(\Omega_j^-) : \varphi|_{\Sigma_j} = 0\}. \quad (3.101)$$

In the fixed domain $\widetilde{\Omega}^-$, this becomes

$$\begin{aligned} \int_{\widetilde{\Omega}^-} \mathcal{A}_j \nabla_{x,z} v_j \cdot \nabla_{x,z} \theta \, dx \, dy &= 0 \quad \forall \theta \in H_{0,*}^1(\widetilde{\Omega}^-) \\ &= \{\varphi \in \dot{H}^1(\widetilde{\Omega}^-) : \varphi|_{z=0} = 0\}, \end{aligned}$$

where

$$\mathcal{A}_j = \begin{bmatrix} \partial_z \varrho_j \cdot Id & -\nabla_x \varrho_j \\ -(\nabla_x \varrho_j)^T & \frac{1+|\nabla_x \varrho_j|^2}{\partial_z \varrho_j} \end{bmatrix}.$$

Consequently,

¹ A priori, Proposition 3.12 would only give a bound in $X^{s-1}([z_0, 0])$ for some $z_0 > -1$. However, one can first apply this with ϱ_j replaced by $\varrho_{j,*}$ which is equal to ϱ_j for $-1 \leq z \leq 0$ and smooth for $-2 \leq z \leq 0$ to obtain a bound on $[-1, 0]$.

$$\begin{aligned} & \int_{\widetilde{\Omega}^-} \mathcal{A}_1 \nabla_{x,z} v \cdot \nabla_{x,z} \theta \, dx \, dy \\ &= \int_{\widetilde{\Omega}^-} (\mathcal{A}_2 - \mathcal{A}_1) \nabla_{x,z} v_2 \cdot \nabla_{x,z} \theta \, dx \, dy \quad \forall \theta \in H_{0,*}^1(\widetilde{\Omega}^-). \end{aligned} \quad (3.102)$$

Since $v|_{z=0} = 0$, we have $v \in H_{0,*}^1(\widetilde{\Omega}^-)$. Inserting $\theta = v$ into (3.102) yields

$$\begin{aligned} \int_{\widetilde{\Omega}^-} \mathcal{A}_1 \nabla_{x,z} v \cdot \nabla_{x,z} v \, dx \, dy &= \int_{\widetilde{\Omega}^-} (\mathcal{A}_2 - \mathcal{A}_1) \nabla_{x,z} v_2 \cdot \nabla_{x,z} v \, dx \, dy \\ &= \int_{\mathbb{R}^d \times (-1,0)} (\mathcal{A}_2 - \mathcal{A}_1) \nabla_{x,z} v_2 \cdot \nabla_{x,z} v \, dx \, dy, \end{aligned} \quad (3.103)$$

where we used the fact that $\mathcal{A}_2 = \mathcal{A}_1$ in $\widetilde{\Omega}_2^-$, which in turn comes from the fact that $\varrho_1 = \varrho_2$ in $\widetilde{\Omega}_2^-$. In view of (3.96) and (3.98),

$$\begin{aligned} \|\mathcal{A}_1 - \mathcal{A}_2\|_{L^2(\mathbb{R}^d \times (-1,0))} &\leq \mathcal{F}(\|\eta_1, \eta_2\|_{B_{\infty,1}^1}) \|\eta_1 - \eta_2\|_{H^{\frac{1}{2}}}, \\ \|\nabla_{x,z} v_2\|_{L^\infty(\mathbb{R}^d \times (-1,0))} &\leq \mathcal{F}(N_s) \|f\|_{\tilde{H}_-^s} \end{aligned} \quad (3.104)$$

and since (see (3.96))

$$\langle \mathcal{A}_1 \nabla_{x,z} v \cdot \nabla_{x,z} v \rangle \geq \min \left(\partial_z \varrho_1, \frac{1}{\partial_z \varrho_1} \right) |\nabla_{x,z} v|^2 \geq \frac{1}{\mathcal{F}(\|\eta_1\|_{B_{\infty,1}^1})} |\nabla_{x,z} v|^2$$

pointwise in $\widetilde{\Omega}^-$, we obtain that

$$\|\nabla_{x,z} v\|_{L^2(\widetilde{\Omega}^-)} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\frac{1}{2}}} \|f\|_{\tilde{H}_-^s}. \quad (3.105)$$

Since $\mathbb{R}^d \times (-1,0) \subset \widetilde{\Omega}^-$, this yields

$$\|\nabla_{x,z} v\|_{L^2((-1,0); L^2(\mathbb{R}^d))} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\frac{1}{2}}} \|f\|_{\tilde{H}_-^s}. \quad (3.106)$$

According to Theorem A.6,

$$\|\nabla_x v\|_{X^{-\frac{1}{2}}([-1,0])} \lesssim \|\nabla_{x,z} v\|_{L^2((-1,0); L^2)}. \quad (3.107)$$

As for $\|\partial_z v\|_{X^{-\frac{1}{2}}([-1,0])}$ it remains to estimate $\|\partial_z v\|_{C([-1,0]; H^{-\frac{1}{2}})}$. Setting

$$\Xi_j(x, z) = -\nabla_x \varrho_j \cdot \nabla_x v_j + \frac{1 + |\nabla_x \varrho_j|^2}{\partial_z \varrho_j} \partial_z v_j,$$

it follows from the equation $\operatorname{div}_{x,z}(\mathcal{A}_j \nabla_{x,z} v_j) = 0$ that

$$\partial_z \Xi_j = -\operatorname{div}_x(\partial_z \varrho_j \nabla_x v_j - \nabla_x \varrho_j \partial_z v_j).$$

Hence $\Xi = \Xi_1 - \Xi_2$ is a divergence

$$\partial_z \Xi = -\operatorname{div}_x(\partial_z \varrho_\delta \nabla_x v_1 + \partial_z \varrho_2 \nabla_x v - \nabla_x \varrho_\delta \partial_z v_1 - \nabla_x \varrho_2 \partial_z v),$$

and using (3.96) and (3.98), we obtain the bounds

$$\begin{aligned} \|\Xi\|_{L_z^2 L^2} &\leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\frac{1}{2}}} \|f\|_{\tilde{H}_-^s}, \\ \|\partial_z \Xi\|_{L_z^2 H^{-1}} &\leq \|\partial_z \varrho_2 \nabla_x v_1 + \partial_z \varrho_2 \nabla_x v - \nabla_x \varrho_2 \partial_z v_1 - \nabla_x \varrho_2 \partial_z v\|_{L_z^2 L^2}, \quad (3.108) \\ &\leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\frac{1}{2}}} \|f\|_{\tilde{H}_-^s}. \end{aligned}$$

Theorem A.6 then yields

$$\|\Xi\|_{C([-1,0]; H^{-\frac{1}{2}})} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\frac{1}{2}}} \|f\|_{\tilde{H}_-^s}.$$

Finally, by writting

$$\begin{aligned} \partial_z v &= \frac{\partial_z \varrho_2}{1 + |\nabla_x \varrho_2|^2} \left\{ \Xi + \nabla_x \varrho_2 \cdot \nabla_x v + \frac{(1 + |\nabla_x \varrho_1|^2)}{\partial_z \varrho_1 \partial_z \varrho_2} \partial_z \varrho_2 \right. \\ &\quad \left. + \left(\nabla_x v_1 - \frac{\partial_z v_1 \nabla_x (\varrho_1 + \varrho_2)}{\partial_z \varrho_1} \right) \nabla_x \varrho_2 \right\} \end{aligned}$$

we deduce that

$$\|\partial_z v\|_{C([-1,0]; H^{-\frac{1}{2}})} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\frac{1}{2}}} \|f\|_{\tilde{H}_-^s}.$$

This completes the proof of Lemma 3.27. \square

This low-regularity bound can easily be upgraded to a bound with no loss of regularity in $\eta_1 - \eta_2$ with the aid of the next lemma. We shall use frequently the fact that for $s > 1 + \frac{d}{2}$, $\sigma \in [\frac{1}{2}, s]$ and $\delta < s - 1 - \frac{d}{2}$, we have

$$\sigma + s > 2 + \delta. \quad (3.109)$$

Lemma 3.28. *For any $\sigma \in [\frac{1}{2}, s]$, we have*

$$\begin{aligned} \|\alpha_1 - \alpha_2\|_{X^{\sigma-1}([-1,0])} + \|\beta_1 - \beta_2\|_{X^{\sigma-1}([-1,0])} \\ \leq \mathcal{F}(\|\eta\|_{H^\sigma}) \|\eta_1 - \eta_2\|_{H^\sigma}, \quad (3.110) \end{aligned}$$

$$\|\gamma_1 - \gamma_2\|_{Y^{\sigma-1}([-1,0])} \leq \mathcal{F}(\|\eta\|_{H^\sigma}) \|\eta_1 - \eta_2\|_{H^\sigma}. \quad (3.111)$$

Proof. From the definition of α and β we see that they are nonlinear functions of $\nabla_{x,z} \varrho$ which is bounded in $L_z^\infty H_x^{s-1}$. By the product rule (C.12), we have that multiplication with H^{s-1} is a continuous linear operator from H^ν to H^ν for any $\nu \in [-\frac{1}{2}, s]$. Thus, (3.110) follows easily.

For $\gamma_1 - \gamma_2$, let us consider the typical term $\frac{\alpha_1}{\partial_z \varrho_1} \Delta_x \varrho_1 - \frac{\alpha_2}{\partial_z \varrho_2} \Delta_x \varrho_2$ which in turns contains the following typical terms

$$E_1 = \frac{\alpha_1 - \alpha_2}{\partial_z \varrho_1} \Delta_x \varrho_1, \quad E_2 = \frac{\alpha_2}{\partial_z \varrho_2} \Delta_x \varrho_2.$$

In view of (3.109), using (3.34) with $s_0 = \sigma - 1$, $s_1 = \sigma - 1$ and $s_2 = s - 2$ we obtain

$$\|E_1\|_{Y^{\sigma-1}} \lesssim \left\| \frac{\alpha_1 - \alpha_2}{\partial_z \mathcal{Q}_1} \right\|_{X^{\sigma-1}} \|\Delta_x \mathcal{Q}_1\|_{X^{s-2}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta_1 - \eta_2\|_{H^\sigma}.$$

On the other hand, applying (3.34) with $s_0 = \sigma - 1$, $s_1 = s - 1$ and $s_2 = \sigma - 2$, we bound E_2 as

$$\|E_2\|_{Y^{\sigma-1}} \lesssim \left\| \frac{\alpha_2}{\partial_z \mathcal{Q}_2} \right\|_{X^{s-1}} \|\Delta_x \mathcal{Q}_\delta\|_{X^{\sigma-2}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta_1 - \eta_2\|_{H^\sigma}.$$

The other terms can be treated similarly; this finishes the proof of (3.111). \square

Lemma 3.29. *For any $\sigma \in [\frac{1}{2}, s]$, we have*

$$\|\nabla_{x,z} v\|_{X^{\sigma-1}([z_0, 0])} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^\sigma} \|f\|_{\tilde{H}_-^s}. \quad (3.112)$$

Proof. We claim that for $z_1 \in (-1, 0)$, and F as in (3.99), there exists \mathcal{F} such that

$$\|F\|_{Y^{\sigma-1}([z_1, 0])} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^\sigma} \|f\|_{\tilde{H}_-^s}. \quad (3.113)$$

We first apply (3.36) with $s_0 = \sigma - 1$, $s_1 = \sigma - 1$ and $s_2 = s - 1$:

$$\|(\gamma_1 - \gamma_2) \partial_z v_2\|_{Y^{\sigma-1}} \lesssim \|\gamma_1 - \gamma_2\|_{Y^{\sigma-1}} \|\partial_z v_2\|_{X^{s-1}} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^\sigma} \|f\|_{\tilde{H}_-^s},$$

where we have used (3.111) in the second inequality. As for $(\alpha_1 - \alpha_2) \Delta_x v_2$ we apply (3.34) with $s_0 = \sigma - 1$, $s_1 = \sigma - 1$ and $s_2 = s - 2$:

$$\|(\alpha_1 - \alpha_2) \Delta_x v_2\|_{Y^{\sigma-1}} \lesssim \|\alpha_1 - \alpha_2\|_{X^{\sigma-1}} \|\Delta_x v_2\|_{X^{s-2}} \lesssim \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^\sigma} \|f\|_{\tilde{H}_-^s}.$$

Proceeding similarly for $(\beta_1 - \beta_2) \cdot \nabla_x \partial_z v_2$, we obtain (3.113). Since $v|_{z=0} = 0$, Proposition 3.12 gives that

$$\|\nabla_{x,z} v\|_{X^{\sigma-1}([z_0, 0])} \leq \mathcal{F}(N_s) (\|F\|_{Y^{\sigma-1}([z_1, 0])} + \|\nabla_{x,z} v\|_{X^{-\frac{1}{2}}([-1, 0])}) \quad (3.114)$$

for $-1 < z_1 < z_0 < 0$. Combining this with (3.113), Lemma 3.27 and the condition $\sigma \geq \frac{1}{2}$, we finish the proof. \square

Proof of Corollary 3.25. Corollary 3.25 can be deduced from Theorem 3.89, Theorem C.4 (i) and the fact that B_2 and V_2 are in L_x^∞ . Here we give a short proof using Lemma 3.29. In view of (3.43), we find that typical terms in $G^-(\eta_1) f - G^-(\eta_2) f$ are $\nabla_x (\mathcal{Q}_1 - \mathcal{Q}_2) \cdot \nabla_x v_1|_{z=0}$ and $\nabla_x \mathcal{Q}_2 \cdot \nabla_x (v_1 - v_2)|_{z=0}$. Using (C.12) and (3.109), we have at $z = 0$ that

$$\begin{aligned} \|\nabla_x (\mathcal{Q}_1 - \mathcal{Q}_2) \cdot \nabla_x v_1\|_{H^{\sigma-1}} &\lesssim \|\nabla_x (\mathcal{Q}_1 - \mathcal{Q}_2)\|_{H^{\sigma-1}} \|\nabla_x v_1\|_{H^{s-1}} \\ &\leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^\sigma} \|f\|_{\tilde{H}_-^s}, \\ \|\nabla_x \mathcal{Q}_2 \cdot \nabla_x (v_1 - v_2)\|_{H^{\sigma-1}} &\lesssim \|\nabla_x \mathcal{Q}_2\|_{H^{s-1}} \|\nabla_x v\|_{H^{\sigma-1}} \\ &\leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^\sigma} \|f\|_{\tilde{H}_-^s}, \end{aligned}$$

where we have applied Lemma 3.29 in the last inequality. This finishes the proof of Corollary 3.25. \square

Let us turn to the proof of Theorem 3.24. Fixing $\sigma \in [\frac{1}{2} + \delta, s]$, we have $\sigma - \delta \in [\frac{1}{2}, s - \delta]$, and hence Lemma 3.29 yields the contraction estimate

$$\|\nabla_{x,z} v\|_{X^{\sigma-1-\delta}([z_0, 0])} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}. \quad (3.115)$$

We first prove a technical analog of Lemma 3.22.

Lemma 3.30. *With notation similar to Lemma 3.20, letting $b_2 = \frac{\partial_z v_2}{\partial_z \varrho_2}$ we have*

$$\begin{aligned} (\partial_z - T_{a_1}) [(\partial_z - T_{A_1})v - T_{b_2}(\partial_z - T_{A_1})\varrho_\delta] &= R_\delta^1, \\ \|R_\delta^1\|_{Y^{\sigma-1}([z_0, 0])} &\leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s} \end{aligned} \quad (3.116)$$

for any $z_0 \in (-1, 0)$.

Proof. Applying (3.77), (3.78) and (3.79) with $\sigma = s$ (note that $s + s > 3$) we obtain that b_2 satisfies

$$\|b_2\|_{L^\infty([z_0, 0]; H^{s-1})} + \|\nabla_{x,z} b_2\|_{X^{s-2}([z_0, 0])} \leq \mathcal{F}(N_s) \|f\|_{\tilde{H}_-^s}. \quad (3.117)$$

We also recall from (3.98) that

$$\|\nabla_{x,z} v_j\|_{X^{s-1}([-1, 0])} \lesssim \mathcal{F}(N_s) \|f\|_{\tilde{H}_-^s}.$$

Set $Q_j = \partial_z^2 + \alpha_j \Delta_x + \beta_j \cdot \nabla_x \partial_z$. Using (C.12), (3.96) and (3.109) we obtain the bounds

$$\begin{aligned} &\|(\nabla_x \varrho_1, \nabla_x \varrho_2)\|_{X^{s-1}} + \left\| \left(\partial_z \varrho_1 - \frac{3}{2}h, \partial_z \varrho_2 - \frac{3}{2}h \right) \right\|_{X^{s-1}} \\ &+ \|(Q_1 \varrho_1, \gamma_1)\|_{X^{s-2}} \leq \mathcal{F}(N_s), \\ &\|\nabla_{x,z} Q_\delta\|_{X^{\sigma-1-\delta}} + \|(\alpha_1 - \alpha_2, \beta_1 - \beta_2)\|_{X^{\sigma-1-\delta}} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}}. \end{aligned}$$

On the other hand, we claim that

$$\|Q_1 \varrho_\delta\|_{Y^{\sigma-1-\delta}} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}}. \quad (3.118)$$

Indeed, from the definition of ϱ_δ we have

$$\|\partial_z^2 \varrho_\delta\|_{L^2 H^{\sigma-\frac{3}{2}-\delta}} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}};$$

on the other hand, by (3.34),

$$\|\alpha_1 \Delta_x \varrho_\delta\|_{Y^{\sigma-1-\delta}} \lesssim (\|\alpha_1 - h^2\|_{X^{s-1}} + 1) \|\Delta_x \varrho_\delta\|_{X^{\sigma-2-\delta}} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}}$$

and similarly for $\beta_j \cdot \nabla_x \partial_z \varrho_\delta$.

Step 1 From (3.97) and the definition (3.45) of γ we have

$$\begin{aligned} Q_1 v - (\gamma_1 - \gamma_2) \partial_z v_2 &= \gamma_1 \partial_z v - (Q_1 - Q_2) v_2, \\ \gamma_1 - \gamma_2 - \frac{1}{\partial_z \varrho_2} Q_1 \varrho_\delta &= \frac{1}{\partial_z \varrho_2} (Q_1 - Q_2) \varrho_2 + \frac{Q_1 \varrho_1}{\partial_z \varrho_1 \partial_z \varrho_2} \partial_z \varrho_\delta. \end{aligned}$$

It follows that

$$Q_1 v - \mathfrak{b}_2 Q_1 \varrho_\delta = R'_\delta, \quad (3.119)$$

$$R'_\delta = \gamma_1 \partial_z v - (Q_1 - Q_2)v_2 + \mathfrak{b}_2(Q_1 - Q_2)\varrho_2 + \frac{\mathfrak{b}_2}{\partial_z Q_1} Q_1 \varrho_1 \partial_z \varrho_\delta. \quad (3.120)$$

We claim that

$$\|R'_\delta\|_{Y^{\sigma-1}([z_0, 0])} \leqq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}. \quad (3.121)$$

We first apply (3.34) with $s_0 = \sigma - 1$, $s_1 = s - 2$, $s_2 = \sigma - 1 - \delta$, giving

$$\|\gamma_1 \partial_z v\|_{Y^{\sigma-1}} \lesssim \|\gamma_1\|_{X^{s-2}} \|\partial_z v\|_{X^{\sigma-1-\delta}} \leqq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}. \quad (3.122)$$

By the same argument we can control $(Q_1 - Q_2)\varrho_2$ and $(Q_1 - Q_2)v_2$ in $Y^{\sigma-1}$. For example, when distributing derivatives in the term $(\alpha_1 - \alpha_2)\Delta_x v_2$ in $(Q_1 - Q_2)v_2$, we see that $\alpha_1 - \alpha_2$ and $\Delta_x v_2$ respectively play the role of $\partial_z v$ and γ_1 in the product $\gamma_1 \partial_z v$.

For products we note that (3.36) gives

$$\|ab\|_{Y^{\sigma-1}} \lesssim \|a\|_{X^{s-1}} \|b\|_{Y^{\sigma-1}}.$$

Applying this for $(a, b) = (\mathfrak{b}_2, (Q_1 - Q_2)\varrho_2)$ we obtain the control of $\mathfrak{b}_2(Q_1 - Q_2)\varrho_2$. As for the last term in R'_δ , we take $a = \frac{\mathfrak{b}_2}{\partial_z Q_1}$ and $b = Q_1 \varrho_1 \partial_z \varrho_\delta$ where applying (3.34) again yields

$$\|Q_1 \varrho_1 \partial_z \varrho_\delta\|_{Y^{\sigma-1}} \lesssim \|Q_1 \varrho_1\|_{X^{s-2}} \|\partial_z \varrho_\delta\|_{X^{\sigma-1-\delta}} \leqq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}.$$

We thus conclude the proof of (3.121).

Step 2 Using (3.62), we factorize $Q_1 v$ and $Q_1 \varrho_\delta$ in (3.119). Then we obtain the first equation in (3.116) with

$$\begin{aligned} R_\delta^1 &= (\partial_z - T_{a_1}) [(\partial_z - T_{A_1})v - T_{\mathfrak{b}_2}(\partial_z - T_{A_1})\varrho_\delta] \\ &= R'_\delta + T_{Q_1 \varrho_\delta} \mathfrak{b}_2 + R(\mathfrak{b}_2, Q_1 \varrho_\delta) - R_{Q_1} v + T_{\mathfrak{b}_2} R_{Q_1} \varrho_\delta \\ &\quad + [T_{\mathfrak{b}_2}, \partial_z - T_{a_1}] (\partial_z - T_{A_1}) \varrho_\delta. \end{aligned}$$

In view of (3.109), applying (3.39) and (3.118) gives

$$\|R(\mathfrak{b}_2, Q_1 \varrho_\delta)\|_{L^1 H^{\sigma-1}} \lesssim \|Q_1 \varrho_\delta\|_{Y^{\sigma-1-\delta}} \|\mathfrak{b}_2\|_{X^{s-1}} \leqq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}.$$

On the other hand, by (3.38) and (3.118),

$$\|T_{Q_1 \varrho_\delta} \mathfrak{b}_2\|_{Y^{\sigma-1}} \lesssim \|Q_1 \varrho_\delta\|_{Y^{\sigma-1-\delta}} \|\mathfrak{b}_2\|_{L^\infty H^{s-1}} \leqq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}.$$

Next we compute

$$[T_{\mathfrak{b}_2}, \partial_z - T_{a_1}] (\partial_z - T_{A_1}) \varrho_\delta = -T_{\partial_z \mathfrak{b}_2} (\partial_z - T_{A_1}) \varrho_\delta - [T_{\mathfrak{b}_2}, T_{a_1}] (\partial_z - T_{A_1}) \varrho_\delta.$$

In view of (3.117) for $\|\partial_z \mathfrak{b}_2\|_{X^{s-2}}$, we have

$$\begin{aligned} \|T_{\partial_z \mathfrak{b}_2} (\partial_z - T_{A_1}) \varrho_\delta\|_{L^2 H^{\sigma-\frac{3}{2}}} &\lesssim \|\partial_z \mathfrak{b}_2\|_{L^2 H^{\sigma-\frac{3}{2}}} \|(\partial_z - T_{A_1}) \varrho_\delta\|_{L^\infty H^{\sigma-1-\delta}} \\ &\leqq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}. \end{aligned}$$

On the other hand, the commutator $[T_{b_2}, T_{a_1}](\partial_z - T_{A_1})\varrho_\delta$ can be controlled in $L^2 H^{\sigma - \frac{3}{2}}$ upon using Theorem C.4 (ii).

It remains to control terms involving R_{Q_1} . Using (3.109) we see that $\theta = \sigma - 1$ and δ satisfy (3.63). Consequently, the estimate (3.66) in Lemma 3.20 can be applied, giving

$$\|R_Q g\|_{Y^{\sigma-1}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\nabla_{x,z} g\|_{X^{\sigma-1-\delta}}.$$

Applying this with $g = v$ and taking into account (3.115) we deduce that $R_{Q_1} v$ is controllable. Finally, with $g = \varrho_\delta$ and with the aid of (3.37), we have

$$\|T_{b_2} R_{Q_1} \varrho_\delta\|_{Y^{\sigma-1}} \lesssim \|b_2\|_{L^\infty H^{s-1}} \|R_{Q_1} \varrho_\delta\|_{Y^{\sigma-1}} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}.$$

The proof of (3.116) is complete. \square

Proof of Theorem 3.24. As in the proof of Theorem 3.18, let us fix $-1 < z_0 < z_1 < 0$ and introduce a cut-off χ satisfying $\chi(z) = 1$ for $z > z_1$ and $\chi(z) = 0$ for $z < z_0$. It follows from Lemma 3.30 that

$$w_\delta := \chi(z) [(\partial_z - T_{A_1})v - T_{b_2}(\partial_z - T_{A_1})\varrho_\delta]$$

satisfies

$$(\partial_z - T_{a_1})w_\delta = R_\delta^2 := \chi(z)R_\delta^1 + \chi'(z)[(\partial_z - T_{A_1})v - T_{b_2}(\partial_z - T_{A_1})\varrho].$$

Applying Theorem C.4 (i) and (3.115) we have

$$\begin{aligned} & \|(\partial_z - T_{A_1})v\|_{L^2([z_0, 0]; H^{\sigma - \frac{3}{2}})} \\ & \leq \mathcal{F}(N_s) \|\nabla_{x,z} v\|_{L^2([z_0, 0]; H^{\sigma - \frac{3}{2}})} \\ & \leq \mathcal{F}(N_s) \|\nabla_{x,z} v\|_{L^2([z_0, 0]; H^{\sigma - \frac{1}{2} - \delta})} \\ & \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}. \end{aligned}$$

In addition, (C.8) together with (3.117) implies

$$\begin{aligned} & \|T_{b_2}(\partial_z - T_{A_1})\varrho_\delta\|_{L^2([z_0, 0]; H^{\sigma - \frac{3}{2}})} \\ & \lesssim \|b_2\|_{L^\infty([z_0, 0]; H^{s-1})} \|(\partial_z - T_{A_1})\varrho_\delta\|_{L^2([z_0, 0]; H^{\sigma - \frac{3}{2}})} \\ & \leq \mathcal{F}(N_s) \|f\|_{\tilde{H}_-^s} \|\nabla_{x,z} \varrho_\delta\|_{L^2([z_0, 0]; H^{\sigma - \frac{3}{2}})} \\ & \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}. \end{aligned}$$

Thus, R_δ^2 satisfies similar estimates as R_δ^1 in (3.116). Since $w_\delta(z_0) = 0$, applying Proposition C.12 yields

$$\|w_\delta\|_{X^{\sigma-1}([z_0, 0])} \leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}. \quad (3.123)$$

In the rest of this proof, functions of (x, z) are evaluated at $z = 0$. Besides, we write $g_1 \simeq g_2$ to signify that g_1 and g_2 agree up to acceptable errors,

$$\|g_1 - g_2\|_{H^{\sigma-1}} \lesssim \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}} \|f\|_{\tilde{H}_-^s}.$$

Set

$$p_j = \frac{1 + |\nabla_x \varrho_j|^2}{\partial_z \varrho_j} = \frac{\partial_z \varrho_j}{\alpha_j}, \quad j = 1, 2$$

so that, by (3.96),

$$\begin{aligned} \|p_1\|_{H^{s-1}} + \|p_2\|_{H^{s-1}} + \|\nabla_{x,z} \varrho_1\|_{H^{s-1}} + \|\nabla_{x,z} \varrho_2\|_{H^{s-1}} &\leq \mathcal{F}(N_s) \\ \|p_1 - p_2\|_{H^{\sigma-1-\delta}} + \|\nabla_{x,z} \varrho_\delta\|_{H^{\sigma-1-\delta}} &\leq \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}}. \end{aligned} \quad (3.124)$$

Using (3.43) and the fact that $v|_{z=0} \equiv 0$, we write

$$\begin{aligned} G^-(\eta_1)f - G^-(\eta_2)f &= (p_1 - p_2)\partial_z v_2 + p_1\partial_z v - \nabla_x \varrho_\delta \cdot \nabla_x v_2 \\ &= (p_1 - p_2)\partial_z v_2 - \nabla_x \varrho_\delta \cdot \nabla_x v_2 \\ &\quad + p_1 [T_{A_1}v + T_{b_2}(\partial_z - T_{A_1})\varrho_\delta] + p_1 w_\delta, \end{aligned} \quad (3.125)$$

where $T_{A_1}v = 0$ (at $z = 0$). Using this, (3.98), (3.123), (3.124) with Theorem C.4 (i), (C.8), (C.11), we find that

$$G^-(\eta_1)f - G^-(\eta_2)f \simeq T_{\partial_z v_2}(p_1 - p_2) - T_{\nabla_x v_2} \nabla_x \varrho_\delta + T_{p_1} T_{b_2}(\partial_z - T_{A_1})\varrho_\delta,$$

where by virtue of Theorem C.4 (ii),

$$T_{p_1} T_{b_2}(\partial_z - T_{A_1})\varrho_\delta \simeq T_{b_2} T_{p_1}(\partial_z - T_{A_1})\varrho_\delta.$$

Next applying Theorem C.11 we find that

$$T_{\partial_z v_2}(p_1 - p_2) \simeq T_{2b_2 \nabla_x \varrho_2} \cdot \nabla_x \varrho_\delta - T_{p_2 b_2} \partial_z \varrho_\delta.$$

We thus arrive at

$$\begin{aligned} G^-(\eta_1)f - G^-(\eta_2)f &\simeq -T_{b_2}(T_{p_1 A_1} - T_{\nabla_x \varrho_2} \nabla_x) \varrho_\delta + T_{b_2 \nabla_x \varrho_2 - \nabla_x v_2} \nabla_x \varrho_\delta \\ &\quad + T_{b_2} T_{p_1 - p_2} \partial_z \varrho_\delta. \end{aligned}$$

By (3.86), we have

$$\begin{aligned} p_1 A_1 - p_2 A_2 &= \lambda_1 - \lambda_2 - i\xi \cdot \nabla_x \varrho_\delta = \tilde{A} \cdot \nabla_x \varrho_\delta, \\ M_\delta^0(\nabla_x \varrho_\delta) + M_\delta^1(\tilde{A}) &\lesssim \mathcal{F}(N_s). \end{aligned}$$

Theorem C.4 (i) and (C.8) yield that

$$\begin{aligned} \|T_{p_1 - p_2} \partial_z \varrho_\delta\|_{H^{\sigma-1}} &\lesssim \|\partial_z \varrho_\delta\|_{H^{s-1}} \|p_1 - p_2\|_{H^{\sigma-1-\delta}} \\ &\lesssim \mathcal{F}(N_s) \|\eta_1 - \eta_2\|_{H^{\sigma-\delta}}, \\ \|T_{p_1 A_1 - p_2 A_2} \varrho_\delta\|_{H^{\sigma-1}} &\leq \mathcal{F}(N_s) \|\varrho_\delta\|_{H^{\sigma-\delta}} + \|T_{\nabla_x \varrho_\delta} T_{\tilde{A}} \varrho_\delta\|_{H^{\sigma-1}} \\ &\lesssim \mathcal{F}(N_s) \|\varrho_\delta\|_{H^{\sigma-\delta}}. \end{aligned}$$

We conclude that

$$G^-(\eta_1)f - G^-(\eta_2)f \simeq -T_{\lambda_2 B_2}(\eta_1 - \eta_2) + T_{V_2} \cdot \nabla(\eta_1 - \eta_2)$$

which finishes the proof of Theorem 3.24. \square

For future reference, let us end this subsection by providing a variant of Corollary 3.25.

Proposition 3.31. *Let $s > 1 + \frac{d}{2}$ with $d \geq 1$. Consider $\eta_1, \eta_2 \in H^s(\mathbb{R}^d)$ with $\text{dist}(\eta_j, \Gamma^-) > 4h > 0$ for $j = 1, 2$. For all $\sigma \in [\frac{1}{2}, s]$, there exists $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (s, σ, h) such that*

$$\|G^-(\eta_1)f - G^-(\eta_2)f\|_{H^{\sigma-1}} \leq \mathcal{F}(\|(\eta_1, \eta_2)\|_{H^s})\|\eta_1 - \eta_2\|_{H^s}\|f\|_{\tilde{H}_-^\sigma}. \quad (3.126)$$

Proof. We shall use the notation in the proof of Theorem 3.24. In our setting, we can strengthen (3.96) to

$$\|\varrho_\delta\|_{X^s} \lesssim \|\eta_1 - \eta_2\|_{H^s}, \quad (3.127)$$

but as $f \in \tilde{H}_-^\sigma$, in place of (3.98) we have

$$\|\nabla_{x,z} v_j\|_{X^{\sigma-1}([-1, 0])} \lesssim \mathcal{F}(N_s)\|f\|_{\tilde{H}_-^\sigma}, \quad N_s = \|(\eta_1, \eta_2)\|_{H^s}. \quad (3.128)$$

Recall that $v = v_1 - v_2$ solves (3.99). Upon using the product rule (C.12) and proceeding as in the proof of Corollary 3.25, (3.126) follows easily from the following estimate for v

$$\|\nabla_{x,z} v\|_{X^{\sigma-1}([z_0, 0])} \leq \mathcal{F}(N_s)\|\eta_1 - \eta_2\|_{H^s}\|f\|_{\tilde{H}_-^\sigma}, \quad z_0 \in (-1, 0). \quad (3.129)$$

To prove (3.129), we apply Proposition 3.12 to have

$$\|\nabla_{x,z} v\|_{X^{\sigma-1}([z_0, 0])} \leq \mathcal{F}(N_s) \left(\|F\|_{Y^{\sigma-1}([z_1, 0])} + \|\nabla_{x,z} v\|_{X^{-\frac{1}{2}}([-1, 0])} \right),$$

where $-1 < z_1 < z_0 < 0$. Let us estimate each term on the right-hand side. We claim that

$$\|\nabla_{x,z} v\|_{X^{-\frac{1}{2}}([-1, 0])} \leq \mathcal{F}(N_s)\|\eta_1 - \eta_2\|_{H^s}\|f\|_{\tilde{H}_-^\sigma}.$$

This follows along the same lines as the proof of Lemma 3.27, except that for the right-hand side of (3.103), in place of (3.104) we estimate

$$\begin{aligned} \|\mathcal{A}_1 - \mathcal{A}_2\|_{L^\infty(\mathbb{R}^d \times (-1, 0))} &\lesssim \|\mathcal{A}_1 - \mathcal{A}_2\|_{L^\infty((-1, 0); H^{s-1})} \lesssim \mathcal{F}(N_s)\|\eta_1 - \eta_2\|_{H^s}, \\ \|\nabla_{x,z} v_2\|_{L^2(\mathbb{R}^d \times (-1, 0))} &\lesssim \|\nabla_{x,z} v_2\|_{L^2((-1, 0); H^{\sigma-\frac{1}{2}})} \lesssim \mathcal{F}(N_s)\|f\|_{\tilde{H}_-^\sigma}. \end{aligned}$$

It remains to estimate $\|F\|_{Y^{s-\frac{3}{2}}([z_1, 0])}$. The first term in F can be bounded using (3.34) as

$$\begin{aligned} \|(\alpha_1 - \alpha_2)\Delta_x v_2\|_{Y^{\sigma-1}} &\lesssim \|\alpha_1 - \alpha_2\|_{X^{s-1}}\|\Delta_x v_2\|_{X^{\sigma-2}} \\ &\lesssim \mathcal{F}(\|\eta\|_{H^s})\|\eta_1 - \eta_2\|_{H^s}\|f\|_{\tilde{H}_-^\sigma}. \end{aligned}$$

Applying (3.34) again gives

$$\begin{aligned} \|(\gamma_1 - \gamma_2)\partial_z v_2\|_{Y^{\sigma-1}} &\lesssim \|\gamma_1 - \gamma_2\|_{X^{s-2}}\|\partial_z v_2\|_{X^{\sigma-1}} \\ &\lesssim \mathcal{F}(\|\eta\|_{H^s})\|\eta_1 - \eta_2\|_{H^s}\|f\|_{\tilde{H}_-^\sigma}. \end{aligned}$$

The proof is complete. \square

4. Proof of the Main Theorems

4.1. Proof of Theorem 2.3

Let us fix $s > 1 + \frac{d}{2}$ and consider either $\Gamma^- = \emptyset$ or $\underline{b}^- \in \dot{W}^{1,\infty}(\mathbb{R}^d)$. This section is organized as follows. First, we assume that η is a solution of (2.3) on $[0, T]$ such that

$$\eta \in C([0, T]; H^s) \cap L^2([0, T]; H^{s+\frac{1}{2}}), \quad (4.1)$$

$$\text{dist}(\eta(t), \Gamma^-) \geq h \quad \forall t \in [0, T], \quad (4.2)$$

$$\inf_{x \in \mathbb{R}^d} (1 - B(x, t)) \geq \alpha > 0 \quad \forall t \in [0, T], \quad (4.3)$$

where B is given by (3.54) with $f = \eta$. Under these assumptions, a priori estimates are derived in Sections 4.1.1, 4.1.2, 4.1.3 and 4.1.4. These estimates will be used for solutions of (4.32) which is similar to (2.3) with $\partial_t \rightarrow \partial_t - \varepsilon \Delta$. This modification only improves the equation and for simplicity, we perform the analysis on solutions of the simpler equation (2.3). Finally, the proof of Theorem 2.3 is given in Section 4.1.5.

4.1.1. Paradifferential reduction We first apply Theorem 3.18 with $f = \eta$ and $\sigma = s$ to have

$$\partial_t \eta = -\kappa T_\lambda(\eta - T_B \eta) + \kappa T_V \cdot \nabla \eta - \kappa R^-(\eta) \eta \quad (4.4)$$

where $R^-(\eta) \eta$ obeys the estimate (3.55) with $\sigma = s$

$$\|R^-(\eta) \eta\|_{H^{s-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^s} (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \quad (4.5)$$

where $\delta \in (0, \frac{1}{2}]$ satisfies $\delta < s - 1 - \frac{d}{2}$. Recall that V and B can be expressed in terms of η by virtue of the formulas (3.54) with $f = \eta$. Note that

$$M_\delta^1(\lambda) \leq \mathcal{F}(\|\eta\|_{H^s}), \quad \|B\|_{W^{\delta,\infty}} \leq C \|B\|_{H^{s-1}} \leq \mathcal{F}(\|\eta\|_{H^s}). \quad (4.6)$$

Owing to Theorem C.4 (ii), $T_\lambda T_B - T_\lambda B$ is of order $1 - \delta$ and

$$\|(T_\lambda T_B - T_\lambda B) \eta\|_{H^{s-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^{s+\frac{1}{2}-\delta}}. \quad (4.7)$$

Combining (4.4), (4.5) and (4.7) we arrive at the following paradifferential reduction for the one-phase Muskat problem (2.3).

Proposition 4.1. *For $\delta \in (0, \frac{1}{2}]$ satisfying $\delta < s - 1 - \frac{d}{2}$, there exists $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (s, δ, h, κ) such that*

$$\partial_t \eta - \kappa T_V \cdot \nabla \eta + \kappa T_{\lambda(1-B)} \eta = f \quad (4.8)$$

with f satisfying

$$\|f\|_{H^{s-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^s} (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}). \quad (4.9)$$

4.1.2. Parabolicity In equation (4.8), $T_V \cdot \nabla \eta$ is an advection term. We now prove that $T_{\kappa\lambda(1-B)}$ is an elliptic operator, showing that (4.8) is a first-order drift-diffusion equation.

Lemma 4.2. *For any $t \in [0, T]$ we have*

$$\sup_{x \in \mathbb{R}^d} B(x, t) < 1. \quad (4.10)$$

In view of the formula (3.54) for B , Lemma 4.2 is a direct consequence of the following surprising upper bound.

Proposition 4.3. *Assume that either $\Gamma^- = \emptyset$ or $\underline{b}^- \in \dot{W}^{1,\infty}(\mathbb{R}^d)$. If $f \in H^s(\mathbb{R}^d)$ with $s > 1 + \frac{d}{2}$, $d \geq 1$, then there exists $c_0 > 0$ such that*

$$G^-(f)f(x) < 1 - c_0 \quad \forall x \in \mathbb{R}^d. \quad (4.11)$$

Proof. Let Ω^- denote the fluid domain with the top boundary $\Sigma = \{y = f(x)\}$ and the bottom Γ^- .

1. *Finite depth* According to Proposition 3.4, there exists a unique solution $\phi \in \dot{H}^1(\Omega^-)$ to the problem

$$\begin{cases} \Delta_{x,y}\phi = 0 & \text{in } \Omega^-, \\ \phi = f & \text{on } \Sigma, \\ \frac{\partial \phi}{\partial \nu^-} = 0 & \text{on } \Gamma^- \end{cases} \quad (4.12)$$

in the sense of (3.20)

$$\int_{\Omega^-} \nabla_{x,y}\phi \cdot \nabla_{x,y}\varphi \, dx \, dy = 0 \quad \forall \varphi \in H_{0,*}^1(\Omega^-). \quad (4.13)$$

Inserting $\varphi = \min\{\phi - \inf_{\mathbb{R}^d} f, 0\} \in H_{0,*}^1(\Omega^-)$ into (4.13) we obtain the minimum principle

$$\inf_{\Omega^-} \phi \geq \inf_{\mathbb{R}^d} f. \quad (4.14)$$

Consequently,

$$\psi(x, y) := \phi(x, y) - y \geq \inf_{\mathbb{R}^d} f - y > 0 \quad (4.15)$$

for $(x, y) \in \Omega^-$ with $y \leq k := (\inf_{\mathbb{R}^d} f) - 1$. We claim that ψ is also nonnegative elsewhere:

$$\psi(x, y) \geq 0 \quad \text{in } \Omega_{\tilde{b}} := \{(x, y) \in \mathbb{R}^d \times \mathbb{R} : \tilde{b}(x) < y < f(x)\}, \quad (4.16)$$

where $\tilde{b}(x) = \max\{\underline{b}^-(x), k\}$ is a Lipschitz and bounded function, $\tilde{b} \in W^{1,\infty}(\mathbb{R}^d)$. Let $\chi : \mathbb{R}^d \rightarrow \mathbb{R}^+$ be a compactly function that equals 1 in $B(0, 1)$ and vanishes outside $B(0, 2)$. Then consider the test functions $\varphi_n = \psi^-(x, y)\chi(\frac{x}{n}) \leq 0$ where $\psi^- = \min\{\psi, 0\}$ and $n \geq 1$. By (4.15), $\text{supp } \psi^- \subset \overline{\Omega_{\tilde{b}}}$ and thus

$$\text{supp } \varphi_n \subset \overline{\Omega_{\tilde{b},n}}, \quad \Omega_{\tilde{b},n} := \Omega_{\tilde{b}} \cap \{|x| < 2n\}$$

which gives $\varphi_n \in H_{0,*}^1(\Omega^-)$. Replacing ϕ with $\psi + y$ and φ with φ_n in (4.13) gives

$$\begin{aligned} \int_{\Omega^-} \nabla_{x,y} \psi \cdot \nabla_{x,y} \varphi_n \, dx \, dy &= - \int_{\Omega_{\tilde{b}} \cap \{|x| < 2n\}} \partial_y \varphi_n \, dx \, dy \\ &= - \int_{\{|x| < 2n\}} \int_{\tilde{b}(x)}^{f(x)} \partial_y \varphi_n(x, y) \, dy \, dx \\ &= \int_{\{|x| < 2n\}} \varphi_n(x, \tilde{b}(x)) \, dx \leqq 0. \end{aligned}$$

On the other hand,

$$\begin{aligned} \int_{\Omega^-} \nabla_{x,y} \psi \cdot \nabla_{x,y} \varphi_n \, dx \, dy &= \int_{U_n} |\nabla_{x,y} \psi|^2 \chi\left(\frac{x}{n}\right) \, dx \, dy \\ &\quad + \frac{1}{n} \int_{U_n} \psi \nabla_x \psi \cdot (\nabla_x \chi)\left(\frac{x}{n}\right) \, dx \, dy, \end{aligned}$$

where $U_n = \{(x, y) \in \Omega_{\tilde{b},n} : \psi(x, y) < 0\}$. Thus,

$$\int_{U_n} |\nabla_{x,y} \psi|^2 \chi\left(\frac{x}{n}\right) \, dx \, dy \leqq \frac{-1}{n} \int_{U_n} \psi \nabla_x \psi \cdot (\nabla_x \chi)\left(\frac{x}{n}\right) \, dx \, dy \quad \forall n \geqq 1. \quad (4.17)$$

Since

$$\psi(x, y) = - \int_y^{f(x)} \partial_y \phi(x, y') \, dy' + f(x) - y$$

and $f(x) - y \geqq 0$, we deduce that if $\psi(x, y) < 0$ then

$$\begin{aligned} |\psi(x, y)| &= \left| \int_y^{f(x)} \partial_y \phi(x, y') \, dy' \right| - |f(x) - y| \\ &\leqq L^{\frac{1}{2}} \left(\int_y^{f(x)} |\partial_y \phi(x, y')|^2 \, dy' \right)^{\frac{1}{2}} \end{aligned}$$

where $L = \|f - \tilde{b}\|_{L^\infty(\mathbb{R}^d)} < \infty$. In particular, for $(x, y) \in U_n \subset \Omega_{\tilde{b},n}$ we have

$$\begin{aligned} \int_{U_n} |\psi(x, y)|^2 \, dx \, dy &\leqq L \int_{\Omega_{\tilde{b},n}} \int_y^{f(x)} |\partial_y \phi(x, y')|^2 \, dy' \, dx \, dy \\ &\leqq L^2 \int_{\Omega_{\tilde{b},n}} |\partial_y \phi(x, y')|^2 \, dx \, dy'. \end{aligned}$$

This combined with the fact that $\nabla_x \psi = \nabla_x \phi$ yields

$$\begin{aligned} &\left| \frac{1}{n} \int_{U_n} \psi \nabla_x \psi \cdot (\nabla_x \chi)\left(\frac{x}{n}\right) \, dx \, dy \right| \\ &\leqq \frac{L \|\nabla \chi\|_{L^\infty(\mathbb{R}^d)}}{n} \|\nabla_x \phi\|_{L^2(\Omega_{\tilde{b},n})} \|\partial_y \phi\|_{L^2(\Omega_{\tilde{b},n})} \\ &\leqq \frac{L \|\nabla \chi\|_{L^\infty(\mathbb{R}^d)}}{n} \|\nabla_{x,y} \phi\|_{L^2(\Omega^-)}^2 \end{aligned}$$

which tends to 0 as $n \rightarrow \infty$. Then applying the monotone convergence theorem to (4.17), we arrive that

$$\int_{\{(x,y) \in \Omega_b^- : \psi(x,y) < 0\}} |\nabla_{x,y} \psi|^2 dx dy \leq 0$$

which proves that $\psi \geq 0$ in Ω_b^- as claimed in (4.16). Combining (4.15) and (4.16) we conclude that $\psi \geq 0$ in Ω^- .

Now by virtue of Proposition 3.12, $\phi \in C_b^1(\Omega_h^-)$ where

$$\Omega_h^- = \{(x, y) \in \mathbb{R}^d \times \mathbb{R} : \eta(x) - h < y < \eta(x)\}$$

for some $h > 0$. Consequently, $\psi \in C_b^1(\Omega_h^-)$, $\psi \geq 0$ everywhere in Ω_h^- and $\psi|_{\Sigma} = 0$. The infimum of ψ over Ω_h^- is thus 0 and is attained at any points of Σ ; moreover, $\psi < 0$ in Ω_h^- thanks to the strong maximum principle. On one hand, it follows from Theorem 3.14 that $G^-(f)f \in H^{s-1}(\mathbb{R}^d)$ with $s-1 > \frac{d}{2}$, implying that $G(f)f(x) < \frac{1}{2}$ for all $|x| \geq M$ for M sufficiently large. On the other hand, letting $V = \Omega_h^- \cap \{|x| < 2M\}$, we can apply Hopf's lemma (see [58]) to the $C^{1,\alpha}$ boundary $\Sigma \cap \{|x| < 2M\}$ of V to have $\frac{\partial \psi}{\partial n} < 0$ for $|x| \leq M$, where n is the upward-pointing normal to Σ . Hence,

$$\frac{\partial \phi}{\partial n} < \frac{\partial y}{\partial n} = \frac{1}{\sqrt{1 + |\nabla f|^2}}$$

which yields $G^-(f)f(x) = \sqrt{1 + |\nabla f|^2} \frac{\partial \phi}{\partial n} < 1$ for all $|x| \leq M$. By the continuity of $G^-(f)f$ on $\{|x| \leq M\}$, we conclude that $G^-(f)f(x) \leq 1 - c_0$ for some $c_0 > 0$ and for all $x \in \mathbb{R}^d$.

2. Infinite depth The proof for this case is in fact simpler. We first let $\phi \in \dot{H}^1(\Omega^-)$ be the solution in the sense of Proposition 3.6 to the problem

$$\begin{cases} \Delta_{x,y} \phi = 0 & \text{in } \Omega^-, \\ \phi = f & \text{on } \Sigma, \\ \nabla_{x,y} \phi \rightarrow 0 & \text{as } y \rightarrow -\infty; \end{cases} \quad (4.18)$$

that is, (4.13) holds. The minimum principle (4.14) remains valid, implying (4.15). Then we can proceed as in the previous case upon replacing Ω_b^- with $\{(x, y) \in \mathbb{R}^d \times \mathbb{R} : \eta(x) - k < y < \eta(x)\}$. \square

Remark 4.4. The proof of Proposition 4.3 is simpler for the periodic case $x \in \mathbb{T}^d$. Indeed, when $x \in \mathbb{T}^d$ we have $\psi^- \in \dot{H}^1(\Omega^-)$ and thus localization in x by $\chi(\frac{x}{n})$ is not needed.

Remark 4.5. The one-phase problem (2.3) dissipates the energy $E(t) = \frac{1}{2} \|\eta(t)\|_{L^2}^2$ since

$$\frac{1}{2} \|\eta(t)\|_{L^2}^2 - \frac{1}{2} \|\eta(0)\|_{L^2}^2 = -\kappa \int_0^t (G^-(\eta)\eta, \eta)_{L^2(\mathbb{R}^d)} dr \leq 0.$$

By virtue of the upper bound (4.11), if $\eta(t)$ remains nonnegative on $[0, T]$ then the energy dissipation over $[0, T]$ is bounded by the L^1 norm of η :

$$\kappa \int_0^T (G^-(\eta)\eta, \eta)_{L^2(\mathbb{R}^d)} dr \leq \kappa \int_0^T \|\eta(r)\|_{L^1(\mathbb{R}^d)} dr.$$

Note that this bound is linear, while the energy is quadratic. In the case of constant viscosity, the same bound was proved in [20] without the sign condition on η .

4.1.3. A priori estimates for η Denote $\langle D_x \rangle = (1 + |D_x|^2)^{\frac{1}{2}}$ and set $\eta_s = \langle D_x \rangle^s$. Conjugating the paradifferential equation (4.8) with $\langle D_x \rangle$ gives

$$\partial_t \eta_s = \kappa T_V \cdot \nabla \eta_s - \kappa T_{\lambda(1-B)} \eta_s + f_1 \quad (4.19)$$

where

$$f_1 = \kappa [\langle D_x \rangle^s, T_V \cdot \nabla] \eta - \kappa [\langle D_x \rangle^s, T_{\lambda(1-B)}] \eta + \langle D_x \rangle^s f.$$

As in (4.6) we have $\|V\|_{W^{\delta, \infty}} \leq \mathcal{F}(\|\eta\|_{H^s})$. This combined with (4.9), (4.6) and Theorem C.4 (iii) implies that $[\langle D_x \rangle^s, T_V \cdot \nabla]$ and $[\langle D_x \rangle^s, T_{\lambda(1-B)}] \eta$ are of order $s + 1 - \delta$, and that

$$\|f_1\|_{H^{-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^s} (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}). \quad (4.20)$$

Taking the L^2 inner product of (4.19) with η_s gives

$$\frac{1}{2} \frac{d}{dt} \|\eta_s\|_{L^2}^2 = \kappa (T_V \cdot \nabla \eta_s, \eta_s)_{L^2} - \kappa (T_{\lambda(1-B)} \eta_s, \eta_s)_{L^2} + (f_1, \eta_s)_{L^2}. \quad (4.21)$$

We have

$$\kappa (T_V \cdot \nabla \eta_s, \eta_s)_{L^2} = \frac{\kappa}{2} \left((T_V \cdot \nabla + (T_V \cdot \nabla)^*) \eta_s, \eta_s \right)_{L^2}$$

where by virtue of Theorem C.4 (iii), $T_V \cdot \nabla + (T_V \cdot \nabla)^*$ is of order $1 - \delta$ and

$$\|((T_V \cdot \nabla)^* + T_V \cdot \nabla) \eta_s\|_{H^{-\frac{1}{2}+\delta}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta_s\|_{H^{\frac{1}{2}}}.$$

Consequently,

$$\kappa |(T_V \cdot \nabla \eta_s, \eta_s)_{L^2}| \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta_s\|_{H^{\frac{1}{2}}} \|\eta_s\|_{H^{\frac{1}{2}-\delta}}. \quad (4.22)$$

Next we write

$$\begin{aligned} (T_{\lambda(1-B)} \eta_s, \eta_s)_{L^2} &= \left(T_{\sqrt{\omega}} \eta_s, T_{\sqrt{\omega}} \eta_s \right)_{L^2} + \left(T_{\sqrt{\omega}} \eta_s, ((T_{\sqrt{\omega}})^* - T_{\sqrt{\omega}}) \eta_s \right)_{L^2} \\ &\quad + \left((T_{\omega} - T_{\sqrt{\omega}} T_{\sqrt{\omega}}) \eta_s, \eta_s \right)_{L^2} \end{aligned} \quad (4.23)$$

where $\omega = \lambda(1 - B)$. In view of (3.50) and (4.3) we have $\omega \geq \alpha |\xi|$, hence

$$M_{\delta}^{\frac{1}{2}}(\sqrt{\omega}) \leq \mathcal{F} \left(\|\eta\|_{H^s}, \frac{1}{\alpha} \right).$$

According to Theorem C.4 (ii) and (iii), $(T_{\sqrt{\omega}})^* - T_{\sqrt{\omega}}$ and $T_{\omega} - T_{\sqrt{\omega}}T_{\sqrt{\omega}}$ are of order $\frac{1}{2} - \delta$ and $1 - \delta$ respectively. Thus

$$\begin{aligned} \left| \left(T_{\sqrt{\omega}}\eta_s, ((T_{\sqrt{\omega}})^* - T_{\sqrt{\omega}})\eta_s \right)_{L^2} \right| &\leq \|T_{\sqrt{\omega}}\eta_s\|_{H^{-\delta}} \|((T_{\sqrt{\omega}})^* - T_{\sqrt{\omega}})\eta_s\|_{H^\delta} \\ &\leq \mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right) \|\eta_s\|_{H^{\frac{1}{2}-\delta}} \|\eta_s\|_{H^{\frac{1}{2}}} \end{aligned} \quad (4.24)$$

and

$$\begin{aligned} \left| \left((T_{\omega} - T_{\sqrt{\omega}}T_{\sqrt{\omega}})\eta_s, \eta_s \right)_{L^2} \right| &\leq \|(T_{\omega} - T_{\sqrt{\omega}}T_{\sqrt{\omega}})\eta_s\|_{H^{-\frac{1}{2}+\delta}} \|\eta_s\|_{H^{\frac{1}{2}-\delta}} \\ &\leq \mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right) \|\eta_s\|_{H^{\frac{1}{2}}} \|\eta_s\|_{H^{\frac{1}{2}-\delta}}. \end{aligned} \quad (4.25)$$

In addition, Theorem C.4 (iii) gives that $T_{\sqrt{\omega}^{-1}}T_{\sqrt{\omega}} - \text{Id}$ is of order $-\delta$ and that

$$\begin{aligned} \|\eta_s\|_{H^{\frac{1}{2}}} &\leq \|T_{\sqrt{\omega}^{-1}}T_{\sqrt{\omega}}\eta_s\|_{H^{\frac{1}{2}}} + \mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right) \|\eta_s\|_{H^{\frac{1}{2}-\delta}} \\ &\leq \mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right) \left(\|T_{\sqrt{\omega}}\eta_s\|_{L^2} + \|\eta_s\|_{H^{\frac{1}{2}-\delta}} \right), \end{aligned}$$

whence

$$\|T_{\sqrt{\omega}}\eta_s\|_{L^2}^2 \geq \frac{1}{\mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right)} \|\eta_s\|_{H^{\frac{1}{2}}}^2 - \|\eta_s\|_{H^{\frac{1}{2}-\delta}}^2. \quad (4.26)$$

Combining (4.23), (4.20), (4.24), (4.25), and (4.26) leads to

$$\begin{aligned} -\kappa(T_{\lambda(1-B)}\eta_s, \eta_s)_{L^2} &\leq -\frac{1}{\mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right)} \|\eta_s\|_{H^{\frac{1}{2}}}^2 \\ &\quad + \mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right) \|\eta_s\|_{H^{\frac{1}{2}}} \|\eta_s\|_{H^{\frac{1}{2}-\delta}}. \end{aligned} \quad (4.27)$$

Putting together (4.21), (4.22), and (4.27) we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\eta_s\|_{L^2}^2 &\leq -\frac{1}{\mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right)} \|\eta_s\|_{H^{\frac{1}{2}}}^2 + \mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right) \\ &\quad \left(\|\eta_s\|_{H^{\frac{1}{2}}} \|\eta_s\|_{H^{\frac{1}{2}-\delta}} + \|\eta_s\|_{H^{\frac{1}{2}}} \|\eta_s\|_{L^2} \right). \end{aligned} \quad (4.28)$$

We assume without loss of generality that $\delta \leq \frac{1}{2}$. The gain of δ derivative gives room to interpolate

$$\|\eta_s\|_{H^{\frac{1}{2}}} \|\eta_s\|_{H^{\frac{1}{2}-\delta}} \leq C \|\eta_s\|_{H^{\frac{1}{2}}}^{1+\mu} \|\eta_s\|_{L^2}^{1-\mu}$$

for some $\mu \in (0, 1)$. Applying Young's inequality yields

$$\frac{1}{2} \frac{d}{dt} \|\eta_s\|_{L^2}^2 \leq -\frac{1}{\mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right)} \|\eta_s\|_{H^{\frac{1}{2}}}^2 + \mathcal{F}\left(\|\eta\|_{H^s}, \frac{1}{\alpha}\right) \|\eta_s\|_{L^2}^2,$$

where \mathcal{F} depends only on (s, h, κ) . Finally, using Grönwall's lemma we obtain the following a priori estimate for η :

Proposition 4.6. *Let $s > 1 + \frac{d}{2}$. Assume that η is a solution of (2.3) on $[0, T]$ with the properties (4.1), (4.2) and (4.3). Then there exists an increasing function $\mathcal{F} : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (s, h, κ) such that*

$$\begin{aligned} & \|\eta\|_{L^\infty([0, T]; H^s)} + \|\eta\|_{L^2([0, T]; H^{s+\frac{1}{2}})} \\ & \leq \mathcal{F}\left(\|\eta(0)\|_{H^s} + T\mathcal{F}\left(\|\eta\|_{L^\infty([0, T]; H^s)}, \mathfrak{a}^{-1}\right), \mathfrak{a}^{-1}\right). \end{aligned} \quad (4.29)$$

In order to close (4.29), we prove a priori estimates for \mathfrak{a} and h in the next subsection.

4.1.4. A priori estimates for the parabolicity and the depth Using (2.3) (or the approximate equation (4.32) below) and Theorem 3.14 (with $\sigma = \frac{1}{2}$) we first observe that

$$\begin{aligned} \|\eta(t) - \eta(0)\|_{H^{-\frac{1}{2}}} & \leq \int_0^t \|\partial_t \eta(\tau)\|_{H^{-\frac{1}{2}}} d\tau \\ & \leq t\mathcal{F}\left(\|\eta\|_{L^\infty([0, t]; H^s)}\right) \|\eta\|_{L^\infty([0, t]; H^{\frac{3}{2}})} \lesssim t\mathcal{F}(\|\eta\|_{L^\infty([0, t]; H^s)}). \end{aligned}$$

Hence, by interpolation, for $s^- = s - \frac{\delta}{2} > 1 + \frac{d}{2} + \frac{\delta}{2}$,

$$\|\eta(t) - \eta(0)\|_{H^{s^-}} \lesssim (\|\eta(t)\|_{H^s} + \|\eta(0)\|_{H^s})^{1-\theta} \|\eta(t) - \eta(0)\|_{H^{-\frac{1}{2}}}^\theta, \quad \theta \in (0, 1)$$

and similarly, by virtue of Theorem 3.14 and Corollary 3.25 (note that $s^- > s - \frac{1}{2} - \delta$),

$$\begin{aligned} \|G(\eta(t))(\eta(t) - \eta(0))\|_{H^{s^-}} & \leq \mathcal{F}(\|\eta\|_{L^\infty H^s}) \|\eta(t) - \eta(0)\|_{H^{s^-}}, \\ \|G(\eta(t) - G(\eta(0)))\eta(0)\|_{H^{s^-}} & \leq \mathcal{F}(\|\eta\|_{L^\infty H^s}) \|\eta(0)\|_{H^s} \|\eta(t) - \eta(0)\|_{H^{s^-}}. \end{aligned}$$

Thus, there exists $\theta > 0$ such that

$$\begin{aligned} & \|G(\eta(t))\eta(t) - G(\eta(0))\eta(0)\|_{L^\infty} + \|\nabla_x \eta(t) - \nabla_x \eta(0)\|_{L^\infty} \\ & + \|\eta(t) - \eta(0)\|_{L^\infty} \leq t^\theta \mathcal{F}(\|\eta\|_{L^\infty([0, t]; H^s)}). \end{aligned}$$

Recalling the definition in (3.54), we deduce that

$$\inf_{(x, t) \in \mathbb{R}^d \times [0, T]} (1 - B(x, t)) \geq \inf_{x \in \mathbb{R}^d} (1 - B(x, 0)) - T^\theta \mathcal{F}_1(\|\eta\|_{L^\infty([0, T]; H^s)}) \quad (4.30)$$

and that

$$\inf_{t \in [0, T]} \text{dist}(\eta(t), \Gamma^-) \geq \text{dist}(\eta_0, \Gamma^-) - T^\theta \mathcal{F}_1(\|\eta\|_{L^\infty([0, T]; H^s)}), \quad (4.31)$$

where $\theta \in (0, 1)$ and $\mathcal{F}_1 : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depends only on (s, h, κ) .

4.1.5. Proof of Theorem 2.3 Having the a priori estimates (4.29), (4.30) and (4.31) in hand, we turn to prove the existence of H^s solutions of (2.3). By a contraction mapping argument, we can prove that for each $\varepsilon \in (0, 1)$ the parabolic approximation

$$\partial_t \eta_\varepsilon = -\kappa G(\eta_\varepsilon) \eta_\varepsilon + \varepsilon \Delta \eta_\varepsilon, \quad \eta_\varepsilon|_{t=0} = \eta_0, \quad (4.32)$$

has a unique solution η_ε in the complete metric space

$$\begin{aligned} E_h(T_\varepsilon) &= \{v \in C([0, T_\varepsilon]; H^s) \cap L^2([0, T_\varepsilon]; H^{s+1}) : \text{dist}(v(t), \Gamma^-) \\ &\geqq h \quad \forall t \in [0, T_\varepsilon]\} \end{aligned} \quad (4.33)$$

provided that T_ε is sufficiently small and that $\text{dist}(\eta_0, \Gamma^-) \geqq 2h > 0$. Let us note that the dissipation term $\varepsilon \Delta \eta_\varepsilon$ in (4.32) has higher order than the term $-\kappa G(\eta_\varepsilon) \eta_\varepsilon$ so that the parabolicity coming from $-\kappa G(\eta_\varepsilon) \eta_\varepsilon$ is not needed in the definition of E_h .

On the other hand, since $\eta_0 \in H^s(\mathbb{R}^d)$ with $s > 1 + \frac{d}{2}$, applying the upper bound in Proposition 4.3 with $f = \eta_0$ we obtain that

$$\inf_{x \in \mathbb{R}^d} (1 - B_\varepsilon(x, 0)) \geqq 2\alpha \quad (4.34)$$

for some constant $\alpha > 0$ independent of ε . It then follows from the a priori estimates (4.29), (4.30), (4.31) and a continuity argument that there exists a positive time T such that $T < T_\varepsilon$ for all $\varepsilon \in (0, 1)$. Moreover, on $[0, T]$, we have the uniform bounds

$$\inf_{(x,t) \in \mathbb{R}^d \times [0, T]} (1 - B_\varepsilon(x, t)) \geqq \alpha, \quad (4.35)$$

$$\text{dist}(\eta_\varepsilon(T), \Gamma^-) \geqq h, \quad (4.36)$$

$$\|\eta_\varepsilon\|_{L^\infty([0, T]; H^s)} + \|\eta_\varepsilon\|_{L^2([0, T]; H^{s+\frac{1}{2}})} \leqq \mathcal{F}(\|\eta(0)\|_{H^s}, \alpha^{-1}), \quad (4.37)$$

where \mathcal{F} depends only on (s, h, κ) . In addition, the L^2 norm of η_ε is nonincreasing in time since

$$(G(\eta_\varepsilon) \eta_\varepsilon, \eta_\varepsilon)_{L^2(\mathbb{R}^d)} \geqq 0, \quad (\Delta \eta_\varepsilon, \eta_\varepsilon)_{L^2(\mathbb{R}^d)} = -\|\nabla \eta_\varepsilon\|_{L^2(\mathbb{R}^d)}^2 \leqq 0.$$

Next we show that for any sequence $\varepsilon_n \rightarrow 0$, the solution sequence $\eta_n \equiv \eta_{\varepsilon_n}$ is Cauchy in the space

$$Z^{s-1}(T) = L^\infty([0, T]; H^{s-1}) \cap L^2([0, T]; H^{s-\frac{1}{2}}).$$

Fix $\delta \in (0, \frac{1}{2}]$ satisfying $\delta < s - 1 - \frac{d}{2}$. We introduce the difference $\eta_\delta = \eta_m - \eta_n$ and claim that it satisfies a nice equation:

$$\partial_t \eta_\delta = -\kappa (T_{\mathfrak{P}} \eta_\delta - T_V \cdot \nabla_x \eta_\delta) + \mathcal{R}_1 + \mathcal{R}_2, \quad (4.38)$$

where

$$\mathfrak{P} = \frac{1}{2}(\lambda_n(1 - B_n) + \lambda_m(1 - B_m)), \quad V = \frac{1}{2}(V_n + V_m),$$

and the remainder terms satisfy

$$\begin{aligned}\|\mathcal{R}_1(t)\|_{H^{s-\frac{3}{2}}} &\leq \mathcal{F}(\|\eta_n\|_{L^\infty H^s} + \|\eta_m\|_{L^\infty H^s}) \|\eta_\delta(t)\|_{H^{s-\frac{1}{2}-\delta}}, \\ \|\mathcal{R}_2(t)\|_{H^{s-\frac{3}{2}}} &\leq (\varepsilon_m + \varepsilon_n) \left(\|\eta_m(t)\|_{H^{s+\frac{1}{2}}} + \|\eta_n(t)\|_{H^{s+\frac{1}{2}}} \right)\end{aligned}$$

Indeed, taking the difference in (4.32), we obtain

$$\partial_t \eta_\delta = -\kappa \left(G^-(\eta_m) \eta_\delta + [G^-(\eta_m) - G^-(\eta_n)] \eta_n \right) + \varepsilon_m \Delta \eta_m - \varepsilon_n \Delta \eta_n.$$

and we can directly set $\mathcal{R}_2 := \varepsilon_m \Delta \eta_m - \varepsilon_n \Delta \eta_n$. For the remaining terms, we apply Theorem 3.16 (with $\sigma = s - \frac{1}{2} - \delta$) and Theorem 3.24 (with $\sigma = s - \frac{1}{2}$) to get

$$\begin{aligned}G^-(\eta_m) \eta_\delta + [G^-(\eta_m) - G^-(\eta_n)] \eta_n &= T_{\lambda_m} \eta_\delta - T_{\lambda_n B_n} \eta_\delta - T_{V_n} \cdot \nabla_x \eta_\delta \\ &\quad + R_0^-(\eta_m) \eta_\delta + R_2^-(\eta_m, \eta_n) \eta_n.\end{aligned}$$

Note that the remainder R_0^- and R_2^- lead to acceptable terms as in \mathcal{R}_1 , but we also have

$$\begin{aligned}G^-(\eta_n) \eta_\delta + [G^-(\eta_n) - G^-(\eta_m)] \eta_m &= T_{\lambda_n} \eta_\delta - T_{\lambda_m B_m} \eta_\delta - T_{V_m} \cdot \nabla_x \eta_\delta \\ &\quad + R_0^-(\eta_n) \eta_\delta + R_2^-(\eta_n, \eta_m) \eta_n.\end{aligned}$$

Thus, by taking the average of the above two identities, we arrive at (4.38).

Now, H^{s-1} energy estimates using (4.38) give that

$$\begin{aligned}\frac{1}{2} \frac{d}{dt} \|\eta_\delta(t)\|_{H^{s-1}}^2 &= -\kappa \left(\langle D_x \rangle^{s-\frac{3}{2}} T_{\mathfrak{B}} \eta_\delta, \langle D_x \rangle^{s-\frac{1}{2}} \eta_\delta \right)_{L^2} \\ &\quad + \kappa \left(\langle D_x \rangle^{s-\frac{3}{2}} T_V \cdot \nabla \eta_\delta, \langle D_x \rangle^{s-\frac{1}{2}} \eta_\delta \right)_{L^2} \\ &\quad + \left(\langle D_x \rangle^{s-\frac{3}{2}} \mathcal{R}_1, \langle D_x \rangle^{s-\frac{1}{2}} \eta_\delta \right)_{L^2} \\ &\quad + \left(\langle D_x \rangle^{s-\frac{3}{2}} \mathcal{R}_2, \langle D_x \rangle^{s-\frac{1}{2}} \eta_\delta \right)_{L^2} \\ &:= I_1 + I_2 + I_3 + I_4.\end{aligned}$$

We can now estimate each term one by one. First, it follows from the above estimates for \mathcal{R}_1 and \mathcal{R}_2 that

$$\begin{aligned}|I_4| &\leq \|\mathcal{R}_2\|_{H^{s-\frac{3}{2}}} \|\eta_\delta\|_{H^{s-\frac{1}{2}}} \\ &\leq (\varepsilon_m + \varepsilon_n) \left(\|\eta_m\|_{H^{s+\frac{1}{2}}}^2 + \|\eta_n\|_{H^{s+\frac{1}{2}}}^2 + \|\eta_\delta\|_{H^{s-\frac{1}{2}}}^2 \right), \\ |I_3| &\leq \|\mathcal{R}_1\|_{H^{s-\frac{3}{2}}} \|\eta_\delta\|_{H^{s-\frac{1}{2}}} \\ &\lesssim \mathcal{F}(\|\eta_n\|_{L^\infty H^s} + \|\eta_m\|_{L^\infty H^s}) \|\eta_\delta\|_{H^{s-\frac{1}{2}-\delta}} \|\eta_\delta\|_{H^{s-\frac{1}{2}}} \\ &\lesssim \mathcal{F}(\|\eta_n\|_{L^\infty H^s} + \|\eta_m\|_{L^\infty H^s}) \|\eta_\delta\|_{H^{s-\frac{1}{2}}}^{1+\theta} \|\eta_\delta\|_{H^{s-1}}^{1-\theta} \\ &\leq \varepsilon_* \|\eta_\delta\|_{H^{s-\frac{1}{2}}}^2 + C_{\varepsilon_*} \mathcal{F}(\|\eta_n\|_{L^\infty H^s} + \|\eta_m\|_{L^\infty H^s}) \|\eta_\delta\|_{H^{s-1}}^2,\end{aligned}$$

where $\varepsilon_* > 0$ is arbitrary. Proceeding as in (4.22), we see that

$$\begin{aligned} |I_2| &\leq \mathcal{F}(\|\eta_n\|_{L^\infty H^s} + \|\eta_m\|_{L^\infty H^s}) \|\eta_\delta\|_{H^{s+\frac{1}{2}}} \|\eta_\delta\|_{H^{s-\frac{1}{2}+\delta}} \\ &\leq \varepsilon_* \|\eta_\delta\|_{H^{s-\frac{1}{2}}}^2 + C_{\varepsilon_*} \mathcal{F}(\|\eta_n\|_{L^\infty H^s} + \|\eta_m\|_{L^\infty H^s}) \|\eta_\delta\|_{H^{s-1}}^2 \end{aligned}$$

and finally, as in (4.27),

$$\begin{aligned} I_1 &\leq -\frac{1}{\mathcal{F}(\|\eta_n\|_{L^\infty H^s}, \frac{1}{\alpha_n})} \|\eta_\delta\|_{H^{s-\frac{1}{2}}}^2 \\ &\quad + \mathcal{F}(\|\eta_n\|_{L^\infty H^s}, \frac{1}{\alpha_n}) \|\eta_\delta\|_{H^{s-\frac{1}{2}}} \|\eta_\delta\|_{H^{s-\frac{1}{2}+\delta}} \\ &\leq -\left(\frac{1}{\mathcal{F}(\|\eta_n\|_{L^\infty H^s}, \frac{1}{\alpha_n})} - \varepsilon_*\right) \|\eta_\delta\|_{H^{s-\frac{1}{2}}}^2 \\ &\quad + C_{\varepsilon_*} \mathcal{F}(\|\eta_n\|_{L^\infty H^s}, \frac{1}{\alpha_n}) \|\eta_\delta\|_{H^{s-1}}^2. \end{aligned}$$

Adding all the above estimates yields

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\eta_\delta(t)\|_{H^{s-1}}^2 &\leq -\left(\frac{1}{\mathcal{F}(\|\eta_n\|_{L^\infty H^s}, \frac{1}{\alpha_n})} - 3\varepsilon_* - \varepsilon_m - \varepsilon_n\right) \|\eta_\delta(t)\|_{H^{s-\frac{1}{2}}}^2 \\ &\quad + C_{\varepsilon_*} \mathcal{F}(\|\eta_n\|_{L^\infty H^s} + \|\eta_m\|_{L^\infty H^s}) \|\eta_\delta(t)\|_{H^{s-1}}^2 \\ &\quad + (\varepsilon_m + \varepsilon_n) \left(\|\eta_m(t)\|_{H^{s+\frac{1}{2}}}^2 + \|\eta_n(t)\|_{H^{s+\frac{1}{2}}}^2 \right). \end{aligned}$$

By virtue of the uniform bounds for η_n , there exists $c_* > 0$ such that

$$\frac{1}{\mathcal{F}(\|\eta_n\|_{L^\infty([0,T];H^s)}, \frac{1}{\alpha_n})} \geq c_* \quad \forall n \in \mathbb{N}.$$

Choosing $\varepsilon_* = \frac{c_*}{10}$ and taking m and n sufficiently large so that $\varepsilon_m, \varepsilon_n \leq \frac{c_*}{10}$, we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\eta_\delta(t)\|_{H^{s-1}}^2 &\leq -\frac{c_*}{2} \|\eta_\delta\|_{H^{s-\frac{1}{2}}}^2 + C \|\eta_\delta\|_{H^{s-1}}^2 \\ &\quad + (\varepsilon_m + \varepsilon_n) \left(\|\eta_m\|_{H^{s+\frac{1}{2}}}^2 + \|\eta_n\|_{H^{s+\frac{1}{2}}}^2 \right). \end{aligned} \tag{4.39}$$

Ignoring the first term on the right-hand side, then integrating in time we obtain

$$\|\eta_\delta(t)\|_{H^{s-1}}^2 \leq C \int_0^t \|\eta_\delta(\tau)\|_{H^{s-1}}^2 d\tau + C(\varepsilon_m + \varepsilon_n) \left(\|\eta_m\|_{H^{s+\frac{1}{2}}}^2 + \|\eta_n\|_{H^{s+\frac{1}{2}}}^2 \right).$$

In view of (4.37), the sequence $\|\eta_n\|_{L^2([0,T];H^{s+\frac{1}{2}})}^2$ is bounded, whence Grönwall's lemma implies

$$\|\eta_\delta\|_{L^\infty([0,T];H^{s-1})}^2 \leq C'(\varepsilon_m + \varepsilon_n) \exp(C'T) \tag{4.40}$$

We then integrate (4.39) in time and use (4.40) to get the dissipation estimate

$$\|\eta_\delta\|_{L^2([0,T];H^{s-\frac{1}{2}})}^2 \leq C''(\varepsilon_m + \varepsilon_n) \exp(C''T). \quad (4.41)$$

It follows from (4.40) and (4.41) that η_n is a Cauchy sequence in $Z^{s-1}(T)$. Therefore, there exists $\eta \in Z^s(T)$ such that $\eta_n \rightarrow \eta$ in $Z^{s-1}(T)$. By virtue of Theorem 3.14 and Corollary 3.25, $G^-(\eta_n)\eta_n \rightarrow G^-(\eta)\eta$ in H^{s-1} and thus η is a solution of (2.3) in $Z^s(T)$.

Repeating the above proof of the fact that η_n is a Cauchy sequence in $Z^{s-1}(T)$, we obtain the following stability estimate.

Proposition 4.7. *Let η_1 and η_2 be two solutions of (2.3) in $Z^s(T)$ defined by (2.10) with $s > 1 + \frac{d}{2}$ and*

$$\begin{aligned} \text{dist}(\eta_j(t), \Gamma^-) &\geq h \quad \forall t \in [0, T], \\ \inf_{x \in \mathbb{R}^d} (1 - B_j(x, t)) &\geq \alpha > 0 \quad \forall t \in [0, T]. \end{aligned}$$

Then,

$$\|\eta_1 - \eta_2\|_{Z^{s-1}(T)} \leq \mathcal{F}(\|(\eta_1, \eta_2)\|_{L^\infty([0, T]; H^s)}) \|(\eta_1 - \eta_2)|_{t=0}\|_{H^{s-1}} \quad (4.42)$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on (s, h, α, κ) .

Finally, uniqueness and continuous dependence on initial data for the solution $\eta \in Z^{s-1}(T)$ constructed above follow at once from Proposition 4.7.

4.2. Proof of Theorem 2.4

We now consider the two-phase problem (2.4)–(2.5). We assume throughout that either $\Gamma^\pm = \emptyset$ or $\underline{b}^\pm \in \dot{W}^{1,\infty}(\mathbb{R}^d)$.

4.2.1. Well-posedness of the elliptic problem (2.5)

Proposition 4.8. *Let $\eta \in W^{1,\infty}(\mathbb{R}^d) \cap H^{\frac{1}{2}}(\mathbb{R}^d)$. Then there exists a unique solution $f^\pm \in \tilde{H}_{\Theta_\pm}^{\frac{1}{2}}(\mathbb{R}^d)$,*

$$\Theta_\pm(x) := \begin{cases} \infty & \text{if } \Gamma^\pm = \emptyset, \\ \frac{\mp(\eta^\pm - \underline{b}^\pm(x))}{2(\|\nabla_x \eta\|_{L^\infty(\mathbb{R}^d)} + \|\underline{b}^\pm\|_{L^\infty(\mathbb{R}^d)})} & \text{if } \underline{b}^\pm \in \dot{W}^{1,\infty}(\mathbb{R}^d), \end{cases}$$

to the system (2.5). Moreover, f^\pm satisfy

$$\begin{aligned} \|f^\pm\|_{\tilde{H}_{\Theta_\pm}^{\frac{1}{2}}(\mathbb{R}^d)} &\leq C(1 + \|\eta\|_{W^{1,\infty}(\mathbb{R}^d)})(1 + \|(\nabla_x \eta, \nabla_x \underline{b}^\pm)\|_{L^\infty(\mathbb{R}^d)} \|\rho\| \|\eta\|_{H^{\frac{1}{2}}(\mathbb{R}^d)}), \end{aligned} \quad (4.43)$$

where the constant C depends only on (μ^\pm, h) .

Proof. Observe that $f^\pm = q^\pm|_\Sigma$ where q^\pm solve the two-phase elliptic problem

$$\begin{cases} \Delta q^\pm = 0 & \text{in } \Omega^\pm, \\ q^+ - q^- = -[\![\rho]\!] \eta, \quad \frac{\partial_n q^+}{\mu^+} - \frac{\partial_n q^-}{\mu^-} = 0 & \text{on } \Sigma, \\ \partial_{\nu^\pm} q^\pm = 0 & \text{on } \Gamma^\pm. \end{cases} \quad (4.44)$$

Here the Neumann boundary conditions need to be modified as in (3.3) when Γ^+ or Γ^- is empty. Thus, it remains to prove the unique solvability of (4.44). To remove the jump of q at the interface, let us fix a cut-off $\chi \in C^\infty(\mathbb{R})$ satisfying $\chi(z) = 1$ for $|z| < \frac{1}{2}$, $\chi(z) = 0$ for $|z| > 1$, and set

$$\underline{\varrho}(x, z) = -\frac{[\![\rho]\!]}{2} \chi(z) e^{-|z|} \langle D_x \rangle \eta(x), \quad (x, z) \in \mathbb{R}^d \times \mathbb{R}$$

and

$$\theta(x, y) = \underline{\varrho} \left(x, \frac{y - \eta(x)}{h} \right), \quad (x, y) \in \mathbb{R}^d \times \mathbb{R}.$$

Then $\theta(x, \eta(x)) = -\frac{[\![\rho]\!]}{2} \eta(x)$ and θ vanishes near Γ^\pm . Moreover, we have

$$\|\theta\|_{H^1(\Omega)} \leq C[\![\rho]\!](1 + \|\eta\|_{W^{1,\infty}}) \|\eta\|_{H^{\frac{1}{2}}}, \quad \Omega = \Omega^+ \cup \Omega^-. \quad (4.45)$$

We then need to prove that there exists a unique solution $r \in \dot{H}^1(\Omega)$ to the problem

$$\begin{cases} -\Delta r = \pm \Delta \theta & \text{in } \Omega^\pm, \\ \frac{\partial_n r}{\mu^+} - \frac{\partial_n r}{\mu^-} = -\partial_n \theta \left(\frac{1}{\mu^+} + \frac{1}{\mu^-} \right) & \text{on } \Sigma, \\ \partial_{\nu^\pm} r = 0 & \text{on } \Gamma^\pm. \end{cases} \quad (4.46)$$

The pair $q^\pm := r|_{\Omega^\pm} \pm \theta$ is the unique solution of (4.44). For a smooth solution r and for any smooth test function $\phi : \Omega \rightarrow \mathbb{R}$ we have after integrating by parts that

$$\int_{\Omega^+} \nabla r \cdot \nabla \phi \, dx + \int_{\Sigma} \partial_n r \phi \, dS = - \int_{\Omega^+} \nabla \theta \cdot \nabla \phi \, dx - \int_{\Sigma} \partial_n \theta \phi \, dS$$

and

$$\int_{\Omega^-} \nabla r \cdot \nabla \phi \, dx - \int_{\Sigma} \partial_n r \phi \, dS = \int_{\Omega^-} \nabla \theta \cdot \nabla \phi \, dx - \int_{\Sigma} \partial_n \theta \phi \, dS.$$

Multiplying the first equation by $\frac{1}{\mu^+}$ and the second one by $\frac{1}{\mu^-}$ then adding and using the jump conditions in (4.46) we obtain

$$\begin{aligned} & \frac{1}{\mu^+} \int_{\Omega^+} \nabla r \cdot \nabla \phi \, dx + \frac{1}{\mu^-} \int_{\Omega^-} \nabla r \cdot \nabla \phi \, dx \\ &= -\frac{1}{\mu^+} \int_{\Omega^+} \nabla \theta \cdot \nabla \phi \, dx + \frac{1}{\mu^-} \int_{\Omega^-} \nabla \theta \cdot \nabla \phi \, dx. \end{aligned} \quad (4.47)$$

Conversely, if r is a sufficiently smooth function that verifies (4.47), then upon integrating by parts we can show that r solves (4.46). Therefore, $r \in \dot{H}^1(\Omega)$ is a variational solution of (4.46) if the weak formulation (4.47) is satisfied for all test functions $\phi \in \dot{H}^1(\Omega)$. By virtue of the estimate (4.45) and Proposition 3.3, the Lax–Milgram theorem guarantees the existence of a unique variational solution r ; moreover, the variational bound

$$\|\nabla r\|_{L^2(\Omega)} \leq C[\rho](1 + \|\eta\|_{W^{1,\infty}})\|\eta\|_{H^{\frac{1}{2}}}$$

holds for some constant C depending only on (μ^\pm, h) . This combined with (4.45) implies that $q^\pm = r|_{\Omega^\pm} \pm \theta$ satisfy the same bound

$$\|q^\pm\|_{\dot{H}^1(\Omega^\pm)} \leq C[\rho](1 + \|\eta\|_{W^{1,\infty}})\|\eta\|_{H^{\frac{1}{2}}}.$$

Finally, (4.43) follows from this and the trace inequalities (A.2) and (A.5). \square

Since it is always possible to find $a \in (0, 1)$ such that $\Theta_\pm(x) \geq \mp a(\eta^\pm(x) - \underline{b}^\pm(x))$, we have $f^\pm \in \widetilde{H}_\pm^{\frac{1}{2}}$.

Remark 4.9. (1) In fact, the proof of Proposition 4.8 shows that for $\eta \in W^{1,\infty}(\mathbb{R}^d)$ and $g \in H^{\frac{1}{2}}(\mathbb{R}^d)$, there exists a unique variational solution $f^\pm \in \widetilde{H}_{\Theta_\pm}^{\frac{1}{2}}(\mathbb{R}^d)$ to the system

$$\begin{cases} f^+ - f^- = g, \\ \frac{1}{\mu^+} G^+(\eta) f^+ - \frac{1}{\mu^-} G^-(\eta) f^- = 0. \end{cases} \quad (4.48)$$

In addition, there exists a constant C depending only on (μ^\pm, h) such that

$$\|f^\pm\|_{\widetilde{H}_{\Theta_\pm}^{\frac{1}{2}}(\mathbb{R}^d)} \leq C(1 + \|\eta\|_{W^{1,\infty}(\mathbb{R}^d)})(1 + \|(\nabla_x \eta, \nabla_x \underline{b}^\pm)\|_{L^\infty(\mathbb{R}^d)})\|g\|_{H^{\frac{1}{2}}(\mathbb{R}^d)}. \quad (4.49)$$

(2) If $\Gamma^+ = \Gamma^- = \emptyset$, then it suffices to assume $\eta \in \dot{W}^{1,\infty}(\mathbb{R}^d)$ and $g \in \dot{H}^{\frac{1}{2}}(\mathbb{R}^d)$ since localization away from Γ^\pm is not needed and one can choose

$$\underline{\theta}(x, z) = \frac{1}{2}e^{-|z||D_x|}g(x), \quad (x, z) \in \mathbb{R}^d \times \mathbb{R}.$$

Then, $\theta(x, y) := \underline{\theta}(x, y - \eta(x))$ satisfies

$$\|\theta\|_{\dot{H}^1(\Omega)} = \|\theta\|_{\dot{H}^1(\mathbb{R}^d)} \leq C(1 + \|\nabla_x \eta\|_{L^\infty(\mathbb{R}^d)})\|g\|_{\dot{H}^{\frac{1}{2}}(\mathbb{R}^d)}.$$

Consequently,

$$\|f^\pm\|_{\widetilde{H}_\infty^{\frac{1}{2}}(\mathbb{R}^d)} \leq C(1 + \|\nabla_x \eta\|_{L^\infty(\mathbb{R}^d)})^2\|g\|_{\dot{H}^{\frac{1}{2}}(\mathbb{R}^d)}. \quad (4.50)$$

4.2.2. A priori higher regularity estimate for f^\pm The weak regularity bound can then be bootstrapped to regularity of the surface η .

Proposition 4.10. *Let f^\pm be the solution of (2.5) as given by Proposition 4.8. If $\eta \in H^s(\mathbb{R}^d)$ with $s > 1 + \frac{d}{2}$ then for any $r \in [\frac{1}{2}, s]$, we have*

$$\|f^\pm\|_{\tilde{H}_\pm^r} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^r}. \quad (4.51)$$

Proof. Fix $\delta \in (0, \min(s - 1 - \frac{d}{2}, 1))$. We first claim that whenever $f^\pm \in \tilde{H}^\sigma$ with $\sigma \in [\frac{1}{2}, s - \delta]$ we have $f^\pm \in \tilde{H}^{\sigma+\delta}$ and

$$\|f^\pm\|_{\tilde{H}^{\sigma+\delta}} \leq \mathcal{F}(\|\eta\|_{H^s}) (\|f^\pm\|_{\tilde{H}^\sigma} + \|\eta\|_{H^{\sigma+\delta}}). \quad (4.52)$$

Indeed, applying Theorem 3.16 we have

$$\begin{aligned} G^\pm(\eta) f^\pm &= \mp T_\lambda f^\pm + R_0^\pm(\eta) f^\pm, \\ \|R_0^\pm(\eta) f^\pm\|_{H^{\sigma-1+\delta}} &\leq \mathcal{F}(\|\eta\|_{H^s}) \|f^\pm\|_{\tilde{H}_\pm^\sigma}. \end{aligned}$$

Plugging this into the second equation in (2.5) we obtain

$$\left\| \frac{1}{\mu^-} T_\lambda f^- + \frac{1}{\mu^+} T_\lambda f^+ \right\|_{H^{\sigma-1+\delta}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|f^\pm\|_{\tilde{H}_\pm^\sigma}.$$

However, $f^+ = f^- - [\rho]$, from the first equation in (2.5), hence

$$\|T_\lambda f^-\|_{H^{\sigma-1+\delta}} \leq \mathcal{F}(\|\eta\|_{H^s}) (\|f^\pm\|_{\tilde{H}_\pm^\sigma} + \|\eta\|_{H^{\sigma+\delta}}). \quad (4.53)$$

On the other hand, by virtue of Theorem C.4 (ii) we have

$$\|\Psi(D)g\|_{H^\nu} = \|T_1 g\|_{H^\nu} \leq \mathcal{F}(\|\eta\|_{H^s}) (\|T_\lambda g\|_{H^{\nu-1}} + \|g\|_{H^{1,\nu-\delta}}) \quad \forall \nu \in \mathbb{R}.$$

Combining this with the inequality

$$\|g\|_{\tilde{H}_\pm^\nu} \lesssim \|g\|_{\tilde{H}_\pm^{\frac{1}{2}}} + \|\psi(D)g\|_{H^\nu} \quad \forall \nu \geq \frac{1}{2}$$

yields

$$\|g\|_{\tilde{H}_\pm^\nu} \lesssim \|g\|_{\tilde{H}_\pm^{\frac{1}{2}}} + \mathcal{F}(\|\eta\|_{H^s}) (\|T_\lambda g\|_{H^{\nu-1}} + \|g\|_{H^{1,\nu-\delta}}) \quad \forall \nu \geq \frac{1}{2}.$$

Applying this with $g = f^-$ and $\nu = \sigma + \delta \geq \frac{1}{2} + \delta$ we deduce in view of (4.53) and (4.43) that

$$\|f^-\|_{\tilde{H}_\pm^{\sigma+\delta}} \leq \mathcal{F}(\|\eta\|_{H^s}) (\|f^\pm\|_{\tilde{H}_\pm^\sigma} + \|\eta\|_{H^{\sigma+\delta}})$$

for all $\sigma \in [\frac{1}{2}, s - \delta]$. Clearly, this implies (4.52).

Then, because (4.51) holds for $r = \frac{1}{2}$, an induction argument using (4.52) shows that (4.51) holds for any $r \in [\frac{1}{2}, s]$. \square

4.2.3. Paradifferential reduction

Proposition 4.11. *Let $s > 1 + \frac{d}{2}$ and let $\delta \in (0, s - 1 - \frac{d}{2})$ and $\delta \leq 1$. If $\eta \in H^s(\mathbb{R}^d)$ and $f^\pm \in \tilde{H}_\pm^s$ solve the system (2.4)–(2.5), then we have*

$$\begin{aligned} \partial_t \eta &= -\frac{1}{\mu^-} G^-(\eta) f^- \\ &= -\frac{1}{\mu^+ + \mu^-} T_\lambda \left([\![\rho]\!] \eta - T_{[\![B]\!]} \eta \right) + \frac{1}{\mu^+ + \mu^-} T_{[\![V]\!]} \cdot \nabla \eta + R(\eta) \end{aligned} \quad (4.54)$$

where $[\![B]\!] = B^- - B^+$, $[\![V]\!] = V^- - V^+$, B^\pm and V^\pm are given by

$$B^\pm = \frac{\nabla \eta \cdot \nabla f^\pm + G^\pm(\eta) f^\pm}{1 + |\nabla \eta|^2}, \quad V^\pm = \nabla f^\pm - B^\pm \nabla \eta, \quad (4.55)$$

and $R(\eta)$ obeys the bound

$$\|R(\eta)\|_{H^{s-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^s} (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}) \quad (4.56)$$

for some $\mathcal{F} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ depending only on $(s, \mu^\pm, [\![\rho]\!], h)$.

Proof. We first apply the paralinearization in Theorem 3.18 to $G^\pm(\eta) f^\pm$ with $\sigma = s$, to have

$$\begin{aligned} G^-(\eta) f^- &= T_\lambda(f^- - T_{B^-} \eta) - T_{V^-} \cdot \nabla \eta + R^-(\eta) f^-, \\ G^+(\eta) f^+ &= -T_\lambda(f^+ - T_{B^+} \eta) + T_{V^+} \cdot \nabla \eta + R^+(\eta) f^+, \end{aligned} \quad (4.57)$$

where, using Proposition 4.10,

$$\|R^\pm(\eta) f^\pm\|_{H^{s-\frac{1}{2}}} \leq \mathcal{F}(\|\eta\|_{H^s}) \|\eta\|_{H^s} (1 + \|\eta\|_{H^{s+\frac{1}{2}-\delta}}). \quad (4.58)$$

In view of the second equation in (2.5), (4.57) yields

$$\begin{aligned} T_\lambda \left(\frac{f^+}{\mu^+} + \frac{f^-}{\mu^-} \right) &= T_\lambda T_{\frac{B^+}{\mu^+} + \frac{B^-}{\mu^-}} \eta + T_{\frac{V^+}{\mu^+} + \frac{V^-}{\mu^-}} \cdot \nabla \eta + \frac{1}{\mu^+} R^+(\eta) f^+ \\ &\quad - \frac{1}{\mu^-} R^-(\eta) f^-. \end{aligned} \quad (4.59)$$

Since $f^+ = f^- - [\![\rho]\!] \eta$, (4.59) implies

$$\begin{aligned} T_\lambda f^- &= \frac{[\![\rho]\!] \mu^-}{\mu^+ + \mu^-} T_\lambda \eta + \frac{1}{\mu^+ + \mu^-} T_\lambda T_{\mu^+ B^- + \mu^- B^+} \eta \\ &\quad + \frac{1}{\mu^+ + \mu^-} T_{\mu^+ V^- + \mu^- V^+} \cdot \nabla \eta \\ &\quad + \frac{1}{\mu^+ + \mu^-} (\mu^- R^+(\eta) f^+ - \mu^+ R^-(\eta) f^-). \end{aligned}$$

Plugging this into the first equation in (4.57), we arrive at (4.54) with

$$R(\eta) = -\frac{1}{\mu^+ + \mu^-} (R^+(\eta) f^+ - \frac{\mu^+}{\mu^-} R^-(\eta) f^-).$$

In view of (4.58), this finishes the proof. \square

When the top fluid is vacuum, equation (4.54) reduces to equation (4.8) previously obtained for the one-phase problem. Remarkably, (4.54) together with the fact that

$$T_\lambda (\llbracket \rho \rrbracket \eta - T_{\llbracket B \rrbracket} \eta) \sim T_{\lambda(\llbracket \rho \rrbracket - \llbracket B \rrbracket)} \eta$$

shows that the two-phase Muskat problem is parabolic so long as the Rayleigh–Taylor condition $RT = \sqrt{1 + |\nabla \eta|^2}(\llbracket \rho \rrbracket - \llbracket B \rrbracket) > 0$ holds. In addition, for constant viscosity, $\llbracket \mu \rrbracket = 0$, by using (2.8) and (2.9) we find that the parabolic term becomes explicit

$$T_{\lambda(\llbracket \rho \rrbracket - \llbracket B \rrbracket)} \eta = T_{\lambda \llbracket \rho \rrbracket (1 + |\nabla_x \eta|^2)^{-1}} \eta. \quad (4.60)$$

4.2.4. Proof of Theorem 2.4 We observe that the paradifferential equation (4.54) has the same form as equation (4.8) for the one-phase problem. In particular, an H^s energy estimate on $[0, T]$ can be obtained as in Section 4.1.3 provided that $RT(x, t) > 0$ for all $(x, t) \in \mathbb{R}^d \times [0, T]$. This stability condition can be propagated as in Section 4.1.4 if it is assumed to hold at initial time. In the rest of this subsection, we only sketch the approximation scheme that preserves the aforementioned a priori estimates.

For each $\varepsilon \in (0, 1)$, consider the approximate problem

$$\partial_t \eta_\varepsilon = -\frac{1}{\mu^-} G^-(\eta_\varepsilon) f_\varepsilon^- + \varepsilon \Delta \eta_\varepsilon, \quad (4.61)$$

where f_ε^\pm solves (2.5):

$$\begin{cases} f_\varepsilon^+ - f_\varepsilon^- = (\rho^+ - \rho^-) \eta_\varepsilon, \\ \frac{1}{\mu^+} G^+(\eta_\varepsilon) f_\varepsilon^+ - \frac{1}{\mu^-} G^-(\eta_\varepsilon) f_\varepsilon^- = 0. \end{cases} \quad (4.62)$$

Proposition 4.12. *Let $s > 1 + \frac{d}{2}$ with $d \geq 1$. For each $\eta_0 \in H^s(\mathbb{R}^d)$ with $\text{dist}(\eta_0, \Gamma^\pm) > 2h > 0$, there exist $T_\varepsilon = T_\varepsilon(\|\eta_0\|_{H^s}, h, s, \mu^\pm, \llbracket \rho \rrbracket) > 0$ and a unique solution η_ε to (4.61)–(4.62) on $[0, T_\varepsilon]$ such that $\eta_\varepsilon|_{t=0} = \eta_0$,*

$$\eta_\varepsilon \in C([0, T_\varepsilon]; H^s(\mathbb{R}^d)) \cap L^2([0, T_\varepsilon]; H^{s+1}(\mathbb{R}^d)),$$

and

$$\text{dist}(\eta_\varepsilon(t), \Gamma^\pm) \geq h \quad \forall t \in [0, T_\varepsilon].$$

Proof. The unique existence of η_ε can be obtained via a contraction mapping argument in the space

$$\begin{aligned} E'_h(T_\varepsilon) = \{v \in L^\infty([0, T_\varepsilon]; H^s) \cap L^2([0, T_\varepsilon]; H^{s+1}) : \text{dist}(v(t), \Gamma^\pm) \\ \geq h \text{ almost everywhere } t \in [0, T_\varepsilon]\} \end{aligned} \quad (4.63)$$

provided that T_ε is sufficiently small. Note the similarity between E'_h and E_h defined by (4.33). \square

Acknowledgements. The work of HQN was partially supported by NSF Grant DMS-1907776. BP was partially supported by NSF Grant DMS-1700282. We would like to thank the reviewer for his/her positive and insightful comments on the manuscript.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Appendix A. Traces for Homogeneous Sobolev Spaces

A.1. Infinite Strip-Like Domains

Let η_1 and η_2 be two Lipschitz functions on \mathbb{R}^d , $\eta_j \in \dot{W}^{1,\infty}(\mathbb{R}^d)$, such that $\eta_1 > \eta_2$. Set

$$L = \|\nabla \eta_1\|_{L^\infty} + \|\nabla \eta_2\|_{L^\infty}, \quad \Theta(x) = \frac{\eta_1(x) - \eta_2(x)}{2L}.$$

Consider the infinite strip-like domain

$$U = \{(x, y) \in \mathbb{R}^{d+1} : \eta_2(x) < y < \eta_1(x)\}. \quad (\text{A.1})$$

We record in this Appendix the trace theory in [49] (see also [60]) for $\dot{H}^1(U)$ where

$$\dot{H}^1(U) = \{u \in L^2_{loc}(U) : \nabla u \in L^2(U)\} / \mathbb{R}.$$

Theorem A.1. ([49, Theorem 5.1]) *There exists a unique linear operator*

$$Tr : \dot{H}^1(U) \rightarrow L^2_{loc}(\mathbb{R}^d)$$

such that the following hold:

- (1) $Tr(u) = u|_{\partial U}$ for all $u \in \dot{H}^1(U) \cap C(\overline{U})$.
- (2) There exists a positive constant $C = C(d)$ such that for all $u \in \dot{H}^1(U)$, the functions $g_j = Tr(u)(\cdot, \eta_j(\cdot))$ are in $\widetilde{H}_\Theta^{\frac{1}{2}}(\mathbb{R}^d)$ and satisfy

$$\|g_j\|_{\widetilde{H}_\Theta^{\frac{1}{2}}(\mathbb{R}^d)} \leq C(1 + L) \|u\|_{\dot{H}^1(U)}, \quad (\text{A.2})$$

$$\int_{\mathbb{R}^d} \frac{|g_1(x) - g_2(x)|^2}{\eta_1(x) - \eta_2(x)} dx \leq C \int_U |\partial_y u(x, y)|^2 dx dy. \quad (\text{A.3})$$

Recall that the space $\widetilde{H}_\Theta^{\frac{1}{2}}(\mathbb{R}^d)$ is defined by (3.5).

Theorem A.2. ([49, Theorem 5.4]) *Suppose that g_1 and g_2 are in $\widetilde{H}_{a(\eta_1 - \eta_2)}^{\frac{1}{2}}(\mathbb{R}^d)$ for some $a \in (0, 1)$ such that*

$$\int_{\mathbb{R}^d} \frac{|g_1(x) - g_2(x)|^2}{\eta_1(x) - \eta_2(x)} dx < \infty.$$

Then there exists $u \in \dot{H}^1(U)$ such that $Tr(u)(\cdot, \eta_j(\cdot)) = g_j$ and

$$\begin{aligned} \|u\|_{\dot{H}^1(U)}^2 &\leq C(1+L)^2 \int_{\mathbb{R}^d} \frac{|g_1(x) - g_2(x)|^2}{|\eta_1(x) - \eta_2(x)|} dx \\ &\quad + C(1+L)^2 \left(\|g_1\|_{\tilde{H}_{a(\eta_1-\eta_2)}^{\frac{1}{2}}(\mathbb{R}^d)}^2 + \|g_2\|_{\tilde{H}_{a(\eta_1-\eta_2)}^{\frac{1}{2}}(\mathbb{R}^d)}^2 \right) \end{aligned} \quad (\text{A.4})$$

where $C = C(d)$.

A.2. Lipschitz Half Spaces

For a Lipschitz function η on \mathbb{R}^d we consider the associated half-space

$$U = \{(x, y) \in \mathbb{R}^{n+1} : y < \eta(x)\}.$$

Theorem A.3. *There exists a unique linear operator*

$$Tr : \dot{H}^1(U) \rightarrow L^2_{loc}(\mathbb{R}^d)$$

such that the following hold:

- (1) $Tr(u) = u|_{\partial U}$ for all $u \in \dot{H}^1(U) \cap C(\overline{U})$.
- (2) There exists a positive constant $C = C(d)$ such that for all $u \in \dot{H}^1(U)$, the function $g = Tr(u)(\cdot, \eta(\cdot))$ is in $\tilde{H}_{\infty}^{\frac{1}{2}}(\mathbb{R}^d)$ and satisfies

$$\|g\|_{\tilde{H}_{\infty}^{\frac{1}{2}}(\mathbb{R}^d)} \leq C(1 + \|\nabla \eta\|_{L^{\infty}(\mathbb{R}^d)}) \|u\|_{\dot{H}^1(U)}. \quad (\text{A.5})$$

Theorem A.4. *For each $g \in \tilde{H}_{\infty}^{\frac{1}{2}}(\mathbb{R}^d)$, there exists $u \in \dot{H}^1(U)$ such that $Tr(u)(\cdot, \eta(\cdot)) = g$ and*

$$\|u\|_{\dot{H}^1(U)} \leq C(1 + \|\nabla \eta\|_{L^{\infty}(\mathbb{R}^d)}) \|g\|_{\tilde{H}_{\infty}^{\frac{1}{2}}(\mathbb{R}^d)} \quad (\text{A.6})$$

where $C = C(d)$.

A.3. Trace of Normalized Normal Derivative

We consider the infinite strip-like domain

$$U_h = \{(x, y) \in \mathbb{R}^d \times \mathbb{R} : \eta(x) - h < y < \eta(x)\}$$

of width h underneath the graph of η

$$\Sigma = \{(x, \eta(x)) : x \in \mathbb{R}^d\}.$$

Note that $n = \frac{1}{\sqrt{1+|\nabla \eta|}}(-\nabla \eta, 1)$ is the upward pointing unit normal to Σ .

Our goal is to show that for any function u in the homogeneous maximal domain of the Laplace operator Δ

$$E(U_h) = \{u \in \dot{H}^1(U_h) : \Delta_{x,y}u \in L^2(U_h)\},$$

the trace

$$\sqrt{1 + |\nabla \eta|^2} \frac{\partial u}{\partial n}|_{\Sigma} = \partial_y u(x, \eta(x)) - (\nabla_x u)(x, \eta(x)) \cdot \nabla \eta(x)$$

makes sense in $H^{-\frac{1}{2}}(\mathbb{R}^d)$.

Theorem A.5. *Assume that $\eta \in \dot{W}^{1,\infty}(\mathbb{R}^d)$. There exists a unique linear operator $\mathcal{N} : E(U_h) \rightarrow H^{-\frac{1}{2}}(\mathbb{R}^d)$ such that the following hold:*

- (1) $\mathcal{N}(u) = \sqrt{1 + |\nabla \eta|^2} \frac{\partial u}{\partial n}|_{\Sigma}$ if $u \in \dot{H}^1(U_h) \cap C^1(\overline{U}_h)$.
- (2) There exists an absolute constant $C > 0$ such that

$$\begin{aligned} \|\mathcal{N}(u)\|_{H^{-\frac{1}{2}}(\mathbb{R}^d)} &\leq Ch^{-\frac{1}{2}}(1 + \|\nabla \eta\|_{L^\infty}) \|\nabla_{x,y} u\|_{L^2(U_h)} \\ &\quad + Ch^{-\frac{1}{2}} \|\Delta_{x,y} u\|_{L^2(U_h)} \end{aligned}$$

for all $u \in E(U_h)$.

Proof. Set $S = \mathbb{R}^d \times (-1, 0)$ and introduce $\theta(x, z) = \eta(x) + zh$ for $(x, z) \in S$. It is clear that $(x, z) \mapsto (x, \theta(x, z))$ is a diffeomorphism from S onto U_h . If $f : U_h \rightarrow \mathbb{R}$ we denote $\tilde{f}(x, z) = f(x, \theta(x, z))$. It follows that

$$\begin{aligned} (\partial_y f)(x, \theta(x, z)) &= \frac{1}{h} \partial_z \tilde{f}(x, z), \\ (\nabla_x f)(x, \theta(x, z)) &= \left(\nabla_x - \frac{\nabla \eta}{h} \partial_z \right) \tilde{f}(x, z) := \Lambda \tilde{f}. \end{aligned}$$

For $u \in E(U)$ we have that $f = \Delta u$ satisfies

$$\tilde{f} = \operatorname{div}_{x,z}(A \nabla_{x,z} \tilde{u}) \tag{A.7}$$

with

$$A = \begin{bmatrix} Id & -\frac{\nabla \eta}{h} \\ -\frac{(\nabla \eta)^T}{h} & \frac{1 + |\nabla \eta|^2}{h^2} \end{bmatrix}. \tag{A.8}$$

Equivalently, we have

$$\operatorname{div}_x \Lambda \tilde{u} + \partial_z \left(-\frac{\nabla \eta}{h} \nabla_x \tilde{u} + \frac{1 + |\nabla \eta|^2}{h^2} \partial_z \tilde{u} \right) = \tilde{f}. \tag{A.9}$$

Let us consider the quantity

$$\begin{aligned} &(\partial_y u)(x, \theta(x, z)) - (\nabla_x u)(x, \theta(x, z)) \cdot \nabla \eta(x) \\ &= \frac{1}{h} \partial_z \tilde{u}(x, z) - \nabla \eta(x, z) \cdot \left(\nabla_x - \frac{\nabla \eta}{h} \partial_z \right) \tilde{u}(x, z) \\ &= -\frac{\nabla \eta(x)}{h} \cdot \nabla_x \tilde{u}(x, z) + \frac{1 + |\nabla \eta(x, z)|^2}{h^2} \partial_z \tilde{u}(x, z) \\ &:= \Xi(x, z). \end{aligned}$$

Using the first expression we deduce easily that

$$\|\Xi\|_{L^2((-1,0); L^2(\mathbb{R}^d))} \leq h^{-\frac{1}{2}} (1 + \|\nabla \eta\|_{L^\infty}) \|\nabla_{x,y} u\|_{L^2(U_h)}. \quad (\text{A.10})$$

Moreover, (A.9) implies

$$\partial_z \Xi = \tilde{f} - \operatorname{div}_x \Lambda \tilde{u}. \quad (\text{A.11})$$

Note that if $u \in C^1(\overline{U})$ then

$$\partial_y u(x, \eta(x)) - (\nabla_x u)(x, \eta(x)) \cdot \nabla \eta(x) = \Xi(x, 0).$$

We shall appeal to Theorem A.6 below to prove that the trace $\Xi|_{z=0}$ is well-defined in $H^{-\frac{1}{2}}(\mathbb{R}^d)$ for $u \in \dot{H}^1(U_h)$. To this end, we use (A.11) to have

$$\|\partial_z \Xi\|_{L^2((-1,0); H^{-1}(\mathbb{R}^d))} \leq \|\tilde{f}\|_{L^2((-1,0); H^{-1}(\mathbb{R}^d))} + \|\Lambda \tilde{u}\|_{L^2((-1,0); L^2(\mathbb{R}^d))}$$

where

$$\|\Lambda \tilde{u}\|_{L^2((-1,0); L^2(\mathbb{R}^d))} \leq h^{-\frac{1}{2}} \|\nabla_x u\|_{L^2(U_h)}.$$

On the other hand,

$$\|\tilde{f}\|_{L^2((-1,0); H^{-1}(\mathbb{R}^d))} \leq \|\tilde{f}\|_{L^2(S)} \leq h^{-\frac{1}{2}} \|f\|_{L^2(U_h)},$$

hence

$$\|\partial_z \Xi\|_{L^2((-1,0); H^{-1}(\mathbb{R}^d))} \leq h^{-\frac{1}{2}} (\|f\|_{L^2(U_h)} + \|\nabla_x u\|_{L^2(U_h)}). \quad (\text{A.12})$$

Combining (A.10) and (A.12) we conclude by virtue of Theorem A.6 that $\Xi \in C([-1, 0]; L^2(\mathbb{R}^d))$ and

$$\|\Xi\|_{C([-1,0]; L^2(\mathbb{R}^d))} \leq C h^{-\frac{1}{2}} (1 + \|\nabla \eta\|_{L^\infty}) \|\nabla_{x,y} u\|_{L^2(U_h)} + C h^{-\frac{1}{2}} \|f\|_{L^2(U_h)} \quad (\text{A.13})$$

for some absolute constant $C > 0$. \square

Theorem A.6. ([48, Theorem 3.1]) *Let $s \in \mathbb{R}$ and I be a closed (bounded or unbounded) interval in \mathbb{R} . Let $u \in L_z^2(I, H^{s+\frac{1}{2}}(\mathbb{R}^d))$ such that $\partial_z u \in L_z^2(I, H^{s-\frac{1}{2}}(\mathbb{R}^d))$. Then $u \in BC(I, H^s(\mathbb{R}^d))$ and there exists an absolute constant $C > 0$ such that*

$$\sup_{z \in I} \|u(z, \cdot)\|_{H^s(\mathbb{R}^d)} \leq C \left(\|u\|_{L^2(I, H^{s+\frac{1}{2}}(\mathbb{R}^d))} + \|\partial_z u\|_{L^2(I, H^{s-\frac{1}{2}}(\mathbb{R}^d))} \right).$$

A.4. Proof of Proposition 3.2

Assuming (3.14), let us prove (3.15). By comparing σ_1 with $\min\{\sigma_1, \sigma_2\} + M$ it suffices to prove the following claim. If $\sigma \geq 2h > 0$ for some $h > 0$, then for any $M > 0$, there exists $C = C(d, h, M)$ such that

$$\|f\|_{\tilde{H}_{\sigma+M}^{\frac{1}{2}}} \leq C \|f\|_{\tilde{H}_{\sigma}^{\frac{1}{2}}}. \quad (\text{A.14})$$

By iteration, (A.14) will follow from the same estimate with M replaced by $\delta = \frac{h}{10}$. To prove this, we first note that

$$\begin{aligned} \|f\|_{\tilde{H}_{\sigma+\delta}^{\frac{1}{2}}}^2 &= \|f\|_{\tilde{H}_{\sigma}^{\frac{1}{2}}}^2 + \int_{x \in \mathbb{R}^d} \int_{\{\sigma(x) \leq |k| \leq \sigma(x) + \delta\}} \\ &\quad \frac{|f(x+k) - f(x)|^2}{|k|^{d+1}} dk dx := \|f\|_{\tilde{H}_{\sigma}^{\frac{1}{2}}}^2 + J. \end{aligned}$$

Letting $\theta(x) = 1 - \frac{h}{2\sigma(x)}$, we have $\theta \in [\frac{1}{2}, 1]$ and

$$|y\theta(x)| \leq \sigma(x) - \frac{h}{4} \quad \text{when} \quad |y| \leq \sigma(x) + \delta. \quad (\text{A.15})$$

Then, for $|u| \leq h/4$, we decompose $J \leq 2J_1 + 2J_2$ where

$$\begin{aligned} J_1 &= \int_{x \in \mathbb{R}^d} \int_{\{\sigma(x) \leq |k| \leq \sigma(x) + \delta\}} \frac{|f(x+k) - f(x + \theta k + u)|^2}{|k|^{d+1}} dk dx \\ J_2 &= \int_{x \in \mathbb{R}^d} \int_{\{\sigma(x) \leq |k| \leq \sigma(x) + \delta\}} \frac{|f(x + \theta k + u) - f(x)|^2}{|k|^{d+1}} dk dx. \end{aligned}$$

We can easily dispense with J_2 by changing variable $k \mapsto z = \theta k + u$ and using (A.15):

$$J_2 \lesssim \int_{x \in \mathbb{R}^d} \int_{\{|z| \leq \sigma(x)\}} \frac{|f(x+z) - f(x)|^2}{|z|^{d+1}} dx dz = \|f\|_{H_{\sigma}^s}^2$$

uniformly in $|u| \leq h/4$. To estimate J_1 , we average over $|u| \leq h/4$:

$$\begin{aligned} J_1 &\lesssim h^{-d} \int_{u \in B(0, h/4)} \int_{x \in \mathbb{R}^d} \int_{\{\sigma(x) \leq |k| \leq \sigma(x) + \delta\}} \\ &\quad \frac{|f(x+k) - f(x+k + (\theta-1)k + u)|^2}{|k|^{d+1}} dk dx du. \end{aligned}$$

Since $|(\theta - 1)k + u| \leq h$, by the changes of variables $u \mapsto (\theta - 1)k + u$ and then $k \mapsto k + x$, we get

$$\begin{aligned}
J_1 &\lesssim h^{-d} \int_{u \in B(0, h)} \int_{x, k \in \mathbb{R}^d} \\
&\quad \frac{|f(x + k) - f(x + k + u)|^2}{|k|^{d+1}} 1_{\{\sigma(x) \leq |k| \leq \sigma(x) + \delta\}} dk dx du \\
&\lesssim h^{-d} \int_{z \in \mathbb{R}^d} \int_{u \in B(0, h)} |f(z) - f(z + u)|^2 \\
&\quad \left\{ \int_{x \in \mathbb{R}^d} \frac{1}{|z - x|^{d+1}} 1_{\{\sigma(x) \leq |z - x| \leq \sigma(x) + \delta\}} dx \right\} du dz \\
&\lesssim h \int_{z \in \mathbb{R}^d} \int_{u \in B(0, h)} \frac{|f(z) - f(z + u)|^2}{|u|^{d+1}} du dz \\
&\lesssim h \|f\|_{H_\sigma^{\frac{1}{2}}}^2,
\end{aligned}$$

which finishes the proof.

Appendix B. Proof of (2.8) and (2.9)

Since $p^+(x, \eta(x)) = p^-(x, \eta(x))$ we have

$$\nabla_x p^+ - \nabla_x p^- = (\partial_y p^- - \partial_y p^+) \nabla \eta \quad \text{on } \Sigma = \{y = \eta(x)\}.$$

Then

$$\begin{aligned}
\sqrt{1 + |\nabla \eta|^2} RT &= -(\nabla_x p^+ - \nabla_x p^-)|_\Sigma \cdot \nabla \eta + (\partial_y p^+ - \partial_y p^-)|_\Sigma \\
&= (\partial_y p^+ - \partial_y p^-)|_\Sigma (1 + |\nabla \eta|^2).
\end{aligned}$$

Finally, using the fact that

$$\partial_y p^\pm|_\Sigma = \partial_y q^\pm|_\Sigma - \rho^\pm = B^\pm - \rho^\pm,$$

we obtain

$$RT = \sqrt{1 + |\nabla \eta|^2} [(\rho^- - \rho^+) - (B^- - B^+)],$$

which proves (2.8). As for (2.9), we use the Darcy law (1.5) and the continuity (1.7) of $u \cdot n$ to have

$$\mu^\pm u \cdot n + \nabla p^\pm = -(0, \rho^\pm) \cdot n = -\rho^\pm (1 + |\nabla \eta|^2)^{-\frac{1}{2}},$$

yielding

$$RT = (\nabla p^+ - \nabla p^-) \cdot n = (\mu^- - \mu^+) u \cdot n + (\rho^- - \rho^+) \rho^\pm (1 + |\nabla \eta|^2)^{-\frac{1}{2}}.$$

Appendix C. Paradifferential Calculus

This section is devoted to a review of basic features of Bony's paradifferential calculus (see for example [3, 12, 45, 52]).

Definition C.1. 1. (Symbols) Given $\rho \in [0, \infty)$ and $m \in \mathbb{R}$, $\Gamma_\rho^m(\mathbb{R}^d)$ denotes the space of locally bounded functions $a(x, \xi)$ on $\mathbb{R}^d \times (\mathbb{R}^d \setminus 0)$, which are C^∞ with respect to ξ for $\xi \neq 0$ and such that, for all $\alpha \in \mathbb{N}^d$ and all $\xi \neq 0$, the function $x \mapsto \partial_\xi^\alpha a(x, \xi)$ belongs to $W^{\rho, \infty}(\mathbb{R}^d)$ and there exists a constant C_α such that

$$\forall |\xi| \geq \frac{1}{2}, \quad \|\partial_\xi^\alpha a(\cdot, \xi)\|_{W^{\rho, \infty}(\mathbb{R}^d)} \leq C_\alpha (1 + |\xi|)^{m - |\alpha|}.$$

Let $a \in \Gamma_\rho^m(\mathbb{R}^d)$, we define the semi-norm

$$M_\rho^m(a) = \sup_{|\alpha| \leq 2(d+2)+\rho} \sup_{|\xi| \geq \frac{1}{2}} \|(1 + |\xi|)^{|\alpha|-m} \partial_\xi^\alpha a(\cdot, \xi)\|_{W^{\rho, \infty}(\mathbb{R}^d)}. \quad (\text{C.1})$$

2. (Paradifferential operators) Given a symbol a , we define the paradifferential operator T_a by

$$\widehat{T_a u}(\xi) = (2\pi)^{-d} \int \chi(\xi - \eta, \eta) \widehat{a}(\xi - \eta, \eta) \Psi(\eta) \widehat{u}(\eta) d\eta, \quad (\text{C.2})$$

where $\widehat{a}(\theta, \xi) = \int e^{-ix \cdot \theta} a(x, \xi) dx$ is the Fourier transform of a with respect to the first variable; χ and Ψ are two fixed C^∞ functions such that

$$\Psi(\eta) = 0 \quad \text{for } |\eta| \leq \frac{1}{5}, \quad \Psi(\eta) = 1 \quad \text{for } |\eta| \geq \frac{1}{4}, \quad (\text{C.3})$$

and $\chi(\theta, \eta)$ satisfies, for $0 < \varepsilon_1 < \varepsilon_2$ small enough,

$$\chi(\theta, \eta) = 1 \quad \text{if } |\theta| \leq \varepsilon_1 |\eta|, \quad \chi(\theta, \eta) = 0 \quad \text{if } |\theta| \geq \varepsilon_2 |\eta|,$$

and such that

$$\forall (\theta, \eta), \quad |\partial_\theta^\alpha \partial_\eta^\beta \chi(\theta, \eta)| \leq C_{\alpha, \beta} (1 + |\eta|)^{-|\alpha|-|\beta|}.$$

Remark C.2. The cut-off χ can be appropriately chosen so that when $a = a(x)$, the paradifferential operator $T_a u$ becomes the usual paraproduct.

Definition C.3. Let $m \in \mathbb{R}$. An operator T is said to be of order m if, for all $\mu \in \mathbb{R}$, it is bounded from H^μ to $H^{\mu-m}$.

Symbolic calculus for paradifferential operators is summarized in the following theorem:

Theorem C.4. (Symbolic calculus) *Let $m \in \mathbb{R}$ and $\rho \in [0, 1)$.*

(i) *If $a \in \Gamma_0^m(\mathbb{R}^d)$, then T_a is of order m . Moreover, for all $\mu \in \mathbb{R}$ there exists a constant K such that*

$$\|T_a\|_{H^\mu \rightarrow H^{\mu-m}} \leq K M_0^m(a). \quad (\text{C.4})$$

(ii) If $a \in \Gamma_\rho^m(\mathbb{R}^d)$, $b \in \Gamma_\rho^{m'}(\mathbb{R}^d)$ then $T_a T_b - T_{ab}$ is of order $m + m' - \rho$. Moreover, for all $\mu \in \mathbb{R}$ there exists a constant K such that

$$\|T_a T_b - T_{ab}\|_{H^\mu \rightarrow H^{\mu-m-m'+\rho}} \leq K(M_\rho^m(a) M_0^{m'}(b) + M_0^m(a) M_\rho^{m'}(b)). \quad (\text{C.5})$$

(iii) Let $a \in \Gamma_\rho^m(\mathbb{R}^d)$. Denote by $(T_a)^*$ the adjoint operator of T_a and by \bar{a} the complex conjugate of a . Then $(T_a)^* - T_{\bar{a}}$ is of order $m - \rho$ where Moreover, for all μ there exists a constant K such that

$$\|(T_a)^* - T_{\bar{a}}\|_{H^\mu \rightarrow H^{\mu-m+\rho}} \leq K M_\rho^m(a). \quad (\text{C.6})$$

Remark C.5. In the definition (C.2) of paradifferential operators, the cut-off Ψ removes the low frequency part of u . In particular, when $a \in \Gamma_0^m$ we have

$$\|T_a u\|_{H^\sigma} \leq C M_0^m(a) \|\nabla u\|_{H^{\sigma+m-1}}.$$

To handle symbols of negative Zygmund regularity, we shall appeal to the following.

Proposition C.6. ([3, Proposition 2.12]) Let $m \in \mathbb{R}$ and $\rho < 0$. We denote by $\dot{\Gamma}_\rho^m(\mathbb{R}^d)$ the class of symbols $a(x, \xi)$ that are homogeneous of order m in ξ , smooth in $\xi \in \mathbb{R}^d \setminus \{0\}$ and such that

$$M_\rho^m(a) = \sup_{|\alpha| \leq 2(d+2)+\rho} \sup_{|\xi| \geq \frac{1}{2}} \||\xi|^{|\alpha|-m} \partial_\xi^\alpha a(\cdot, \xi)\|_{C_*^\rho(\mathbb{R}^d)} < \infty.$$

If $a \in \dot{\Gamma}_\rho^m$ then the operator T_a defined by (C.2) is of order $m - \rho$.

Notation C.7. If a and u depend on a parameter $z \in J \subset \mathbb{R}$ we denote

$$(T_a u)(z) = T_{a(z)} u(z), \quad M_\rho^m(a; J) = \sup_{z \in J} M_\rho^m(a(z)).$$

If $M_\rho^m(a; J)$ is finite we write $a \in \Gamma_\rho^m(\mathbb{R}^d \times J)$.

Definition C.8. Given two functions a , u defined on \mathbb{R}^d the Bony's remainder is defined by

$$R(a, u) = au - T_a u - T_u a.$$

We gather here several useful product and paraproduct rules.

Theorem C.9. Let s_0, s_1 and s_2 be real numbers.

(1) For any $s \in \mathbb{R}$,

$$\|T_a u\|_{H^s} \leq C \|a\|_{L^\infty} \|u\|_{H^s}. \quad (\text{C.7})$$

(2) If $s_0 \leq s_2$ and $s_0 < s_1 + s_2 - \frac{d}{2}$, then

$$\|T_a u\|_{H^{s_0}} \leq C \|a\|_{H^{s_1}} \|u\|_{H^{s_2}}. \quad (\text{C.8})$$

(3) If $s_1 + s_2 > 0$ then

$$\|R(a, u)\|_{H^{s_1+s_2-\frac{d}{2}}(\mathbb{R}^d)} \leq C \|a\|_{H^{s_1}(\mathbb{R}^d)} \|u\|_{H^{s_2}(\mathbb{R}^d)}, \quad (\text{C.9})$$

$$\|R(a, u)\|_{H^{s_1+s_2}(\mathbb{R}^d)} \leq C \|a\|_{C_*^{s_1}(\mathbb{R}^d)} \|u\|_{H^{s_2}(\mathbb{R}^d)}. \quad (\text{C.10})$$

(4) If $s_1 + s_2 > 0$, $s_0 \leq s_1$ and $s_0 < s_1 + s_2 - \frac{d}{2}$ then

$$\|au - T_a u\|_{H^{s_0}} \leq C \|a\|_{H^{s_1}} \|u\|_{H^{s_2}}. \quad (\text{C.11})$$

(5) If $s_1 + s_2 > 0$, $s_0 \leq s_1$, $s_0 \leq s_2$ and $s_0 < s_1 + s_2 - \frac{d}{2}$ then

$$\|u_1 u_2\|_{H^{s_0}} \leq C \|u_1\|_{H^{s_1}} \|u_2\|_{H^{s_2}}. \quad (\text{C.12})$$

Theorem C.10. Consider $F \in C^\infty(\mathbb{C}^N)$ such that $F(0) = 0$.

(i) For $s > \frac{d}{2}$, there exists a non-decreasing function $\mathcal{F}: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that, for any $U \in H^s(\mathbb{R}^d)^N$,

$$\|F(U)\|_{H^s} \leq \mathcal{F}(\|U\|_{L^\infty}) \|U\|_{H^s}. \quad (\text{C.13})$$

(ii) For $s > 0$, there exists an increasing function $\mathcal{F}: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that, for any $U \in C_*^s(\mathbb{R}^d)^N$,

$$\|F(U)\|_{C_*^s} \leq \mathcal{F}(\|U\|_{L^\infty}) \|U\|_{C_*^s}. \quad (\text{C.14})$$

Theorem C.11. ([10, Theorem 2.92] and [52, Theorem 5.2.4]) (Paralinearization for nonlinear functions) Let μ , τ be positive real numbers and let $F \in C^\infty(\mathbb{C}^N)$ be a scalar function satisfying $F(0) = 0$. If $U = (u_j)_{j=1}^N$ with $u_j \in H^\mu(\mathbb{R}^d) \cap C_*^\tau(\mathbb{R}^d)$ then we have

$$F(U) = \sum_{j=1}^N T_{\partial_j F(U)} u_j + R_F(U) \quad (\text{C.15})$$

with

$$\|R_F(U)\|_{H^{\mu+\tau}} \leq \mathcal{F}(\|U\|_{L^\infty}) \|U\|_{C_*^\tau} \|U\|_{H^\mu}.$$

Recall the definitions (3.32). The next proposition provides parabolic estimates for elliptic paradifferential operators.

Proposition C.12. ([3, Proposition 2.18]) Let $r \in \mathbb{R}$, $\varrho \in (0, 1)$, $J = [z_0, z_1] \subset \mathbb{R}$ and let $p \in \Gamma_\varrho^1(\mathbb{R}^d \times J)$ satisfying

$$\operatorname{Re} p(z; x, \xi) \geq c|\xi|,$$

for some positive constant c . Then for any $f \in Y^r(J)$ and $w_0 \in H^r(\mathbb{R}^d)$, there exists $w \in X^r(J)$ solution of the parabolic evolution equation

$$\partial_z w + T_p w = f, \quad w|_{z=z_0} = w_0, \quad (\text{C.16})$$

satisfying

$$\|w\|_{X^r(J)} \leq \mathcal{F}\left(\mathcal{M}_\varrho^1(p), \frac{1}{c}\right) (\|w_0\|_{H^r} + \|f\|_{Y^r(J)})$$

for some increasing function \mathcal{F} depending only on r and ϱ . Furthermore, this solution is unique in $X^s(J)$ for any $s \in \mathbb{R}$.

Appendix D. Proof of Lemma 3.10

(1) Assuming (3.33), we prove (3.34). We decompose $u_1 u_2 = T_{u_1} u_2 + T_{u_2} u_1 + R(u_1, u_2)$. Since $s_0 \leq s_2 + 1$ and $s_1 + s_2 > s_0 + \frac{d}{2} - 1$, (C.8) gives

$$\|T_{u_1} u_2\|_{L^2 H^{s_0 - \frac{1}{2}}} \lesssim \|u_1\|_{L^\infty H^{s_1}} \|u_2\|_{L^2 H^{s_2 + \frac{1}{2}}} \lesssim \|u_1\|_{X^{s_1}} \|u_2\|_{X^{s_2}}.$$

The paraproduct $T_{u_2} u_1$ can be estimated similarly. As for $R(u_1, u_2)$ we use (C.9) and the conditions $s_1 + s_2 + 1 > 0$ and $s_1 + s_2 > s_0 + \frac{d}{2} - 1$:

$$\|R(u_1, u_2)\|_{L^1 H^{s_0}} \lesssim \|u_1\|_{L^2 H^{s_1 + \frac{1}{2}}} \|u_2\|_{L^2 H^{s_2 + \frac{1}{2}}} \lesssim \|u_1\|_{X^{s_1}} \|u_2\|_{X^{s_2}}.$$

(2) Assuming (3.35), we prove (3.36). Again, we decompose $u_1 u_2 = T_{u_1} u_2 + T_{u_2} u_1 + R(u_1, u_2)$. If $u_1 = a + b$ with $a \in L^1 H^{s_1}$ and $b \in L^2 H^{s_1 - \frac{1}{2}}$ then

$$u_1 u_2 = T_a u_2 + T_b u_2 + T_{u_2} a + T_{u_2} b + R(a, u_2) + R(b, u_2).$$

Using the conditions $s_0 \leq s_2$ and $s_1 + s_2 > s_0 + \frac{d}{2}$, we can apply (C.8) to get

$$\begin{aligned} \|T_a u_2\|_{L^1 H^{s_0}} &\lesssim \|a\|_{L^1 H^{s_1}} \|u_2\|_{L^\infty H^{s_2}}, \\ \|T_b u_2\|_{L^2 H^{s_0 - \frac{1}{2}}} &\lesssim \|b\|_{L^2 H^{s_1 - \frac{1}{2}}} \|u_2\|_{L^\infty H^{s_2}}. \end{aligned}$$

Next from the conditions $s_0 \leq s_1$ and $s_1 + s_2 > s_0 + \frac{d}{2}$, (C.8) yields

$$\begin{aligned} \|T_{u_2} a\|_{L^1 H^{s_0}} &\lesssim \|u_2\|_{L^\infty H^{s_2}} \|a\|_{L^1 H^{s_1}}, \\ \|T_{u_2} b\|_{L^2 H^{s_0 - \frac{1}{2}}} &\lesssim \|u_2\|_{L^\infty H^{s_2}} \|b\|_{L^2 H^{s_1 - \frac{1}{2}}}. \end{aligned}$$

Finally, under the conditions $s_1 + s_2 > 0$ and $s_1 + s_2 > s_0 + \frac{d}{2}$, using (C.9) we obtain

$$\begin{aligned} \|R(a, u_2)\|_{L^1 H^{s_0}} &\lesssim \|a\|_{L^1 H^{s_1}} \|u_2\|_{L^\infty H^{s_2}}, \\ \|R(b, u_2)\|_{L^1 H^{s_0}} &\lesssim \|b\|_{L^2 H^{s_1 - \frac{1}{2}}} \|u_2\|_{L^2 H^{s_2 + \frac{1}{2}}}. \end{aligned}$$

We have proved that

$$\|u_1 u_2\|_{Y^{s_0}} \lesssim (\|a\|_{L^1 H^{s_1}} + \|b\|_{L^2 H^{s_1 - \frac{1}{2}}}) \|u_2\|_{X^{s_2}}$$

for any decomposition $u_1 = a + b$. Therefore, (3.36) follows. We have also proved (3.37), (3.38) and (3.39).

References

1. AI, A.: Low regularity solutions for gravity water waves, preprint, [arXiv:1712.07821](https://arxiv.org/abs/1712.07821), 2017
2. ALAZARD, T., BURQ, N., ZUILY, C.: On the water waves equations with surface tension. *Duke Math. J.* **158**(3), 413–499, 2011
3. ALAZARD, T., BURQ, N., ZUILY, C.: On the Cauchy problem for gravity water waves. *Invent. Math.* **198**(1), 71–163, 2014
4. ALAZARD, T., BURQ, N., ZUILY, C.: *Strichartz Estimates and the Cauchy Problem for the Gravity Water Waves Equations. Memoirs of the AMS*, Vol. 256, 2018
5. ALAZARD, T., LAZAR, O.: Paralinearization of the Muskat equation and application to the Cauchy problem, preprint [arXiv:1907.02138](https://arxiv.org/abs/1907.02138), 2019
6. ALAZARD, T., MEUNIER, N., SMETS, D.: Lyapounov functions, identities and the Cauchy problem for the Hele-Shaw equation, preprint [arXiv:1907.03691](https://arxiv.org/abs/1907.03691), 2019
7. ALAZARD, T., MÉTIVIER, G.: Paralinearization of the Dirichlet to Neumann operator, and regularity of diamond waves. *Commun. Partial Differ. Equ.* **34**(10–12), 1632–1704, 2009
8. AMBROSE, D.M.: Well-posedness of two-phase Hele-Shaw flow without surface tension. *Eur. J. Appl. Math.* **15**(5), 597–607, 2004
9. AMBROSE, D.M.: Well-posedness of two-phase Darcy flow in 3D. *Q. Appl. Math.* **65**(1), 189–203, 2007
10. BAHOURI, H., CHEMIN, J-Y, DANCHIN, R.: *Fourier analysis and nonlinear partial differential equations*, volume 343 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, 2011
11. BERSELLI, L.C., CÓRDOBA, D., GRANERO-BELINCHÓN, R.: Local solvability and turning for the inhomogeneous Muskat problem. *Interfaces Free Bound.* **16**(2), 175–213, 2014
12. BONY, J.-M.: Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires. *Ann. Sci. École Norm. Sup. (4)* **14**(2), 209–246, 1981
13. CAMERON, S.: Global well-posedness for the two-dimensional Muskat problem with slope less than 1. *Anal. PDE* **12**(4), 997–1022, 2019
14. CASTRO, A., CÓRDOBA, D., FEFFERMAN, C., GANCEDO, F.: Breakdown of smoothness for the Muskat problem. *Arch. Ration. Mech. Anal.* **208**(3), 805–909, 2013
15. CASTRO, A., CÓRDOBA, D., FEFFERMAN, C.L., GANCEDO, F., LÓPEZ-FERNÁNDEZ, María: Rayleigh Taylor breakdown for the Muskat problem with applications to water waves. *Ann. Math.* **175**(2), 909–948, 2012
16. CHANG-LARA, H.A., GUILLEN, N., SCHWAB, R.W.: Some free boundary problems recast as nonlocal parabolic equations, preprint, [arXiv:1807.02714](https://arxiv.org/abs/1807.02714), 2018
17. CHEN, X.: The Hele-Shaw problem and area-preserving curve-shortening motions. *Arch. Ration. Mech. Anal.* **123**(2), 117–151, 1993
18. CHENG, C.H., GRANERO-BELINCHÓN, R., SHKOLLER, S.: Well-posedness of the Muskat problem with H^2 initial data. *Adv. Math.* **286**, 32–104, 2016
19. CONSTANTIN, P., CÓRDOBA, D., GANCEDO, F., RODRIGUEZ-PIAZZA, L., STRAIN, R.M.: On the Muskat problem: global in time results in 2D and 3D. *Am. J. Math.* **138**(6), 1455–1494, 2016
20. CONSTANTIN, P., CÓRDOBA, D., GANCEDO, F., STRAIN, R.M.: On the global existence for the Muskat problem. *J. Eur. Math. Soc.* **15**, 201–227, 2013
21. CONSTANTIN, P., GANCEDO, F., SHVYDKOY, R., VICOL, V.: Global regularity for 2D Muskat equations with finite slope. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **34**(4), 1041–1074, 2017
22. CONSTANTIN, P., PUGH, M.: Global solutions for small data to the Hele-Shaw problem. *Nonlinearity* **6**(3), 393–415, 1993
23. CÓRDOBA, A., CÓRDOBA, D., GANCEDO, F.: Interface evolution: the Hele-Shaw and Muskat problems. *Ann. Math.* **173**(1), 477–542, 2011
24. CÓRDOBA, A., CÓRDOBA, D., GANCEDO, F.: Porous media: the Muskat problem in three dimensions. *Anal. PDE* **6**(2), 447–497, 2013

25. CÓRDOBA, D., GANCEDO, F.: Contour dynamics of incompressible 3-D fluids in a porous medium with different densities. *Commun. Math. Phys.* **273**(2), 445–471, 2007
26. CÓRDOBA, D., GANCEDO, F.: A maximum principle for the Muskat problem for fluids with different densities. *Commun. Math. Phys.* **286**(2), 681–696, 2009
27. CÓRDOBA, D., GRANERO-BELINCHÓN, R., ORIVE, R.: The confined Muskat problem: differences with the deep water regime. *Commun. Math. Sci.* **12**(3), 423–455, 2014
28. CÓRDOBA, D., LAZAR, O.: Global well-posedness for the 2d stable Muskat problem in $H^3/2$, preprint, [arXiv:1803.07528](https://arxiv.org/abs/1803.07528), 2018
29. DENG, F., LEI, Z., LIN, F.: On the two-dimensional Muskat problem with monotone large initial data. *Commun. Pure Appl. Math.* **70**(6), 1115–1145, 2017
30. de POYFERRÉ, T.: A priori estimates for water waves with emerging bottom. *Arch. Ration. Mech. Anal.* **232**(2), 763–812, 2019
31. de POYFERRÉ, T., NGUYEN, H.Q.: A paradifferential reduction for the gravity–capillary waves system at low regularity and applications. *Bull. Soc. Math. Fr.* **145**(4), 643–710, 2017
32. de POYFERRE, T., NGUYEN, H.Q.: Strichartz estimates and local existence for the gravity–capillary water waves with non-Lipschitz initial velocity. *J. Differ. Equ.* **261**(1), 396–438, 2016
33. ESCHER, J., SIMONETT, G.: Classical solutions for Hele-Shaw models with surface tension. *Adv. Differ. Equ.* **2**, 619–642, 1997
34. ESCHER, J., MATIOC, B.V.: On the parabolicity of the Muskat problem: well-posedness, fingering, and stability results. *Z. Anal. Anwend.* **30**(2), 193–218, 2011
35. FLYNN, P., NGUYEN, H.Q.: The vanishing surface tension limit of the Muskat problem, preprint [arXiv:2001.10473](https://arxiv.org/abs/2001.10473), 2020
36. GANCEDO, F.: A survey for the Muskat problem and a new estimate. *SeMA* **74**(1), 21–35, 2017
37. GÓMEZ-SERRANO, J., GRANERO-BELINCHÓN, R.: On turning waves for the inhomogeneous Muskat problem: a computer-assisted proof. *Nonlinearity* **27**(6), 1471–1498, 2014
38. GRANERO-BELINCHÓN, R.: Global existence for the confined Muskat problem. *SIAM J. Math. Anal.* **46**(2), 1651–1680, 2014
39. GANCEDO, F., GARCÍA-JUÁREZ, E., PATEL, N., STRAIN, R.M.: On the Muskat problem with viscosity jump: global in time results. *Adv. Math.* **345**, 552–597, 2019
40. GRANERO-BELINCHÓN, R., LAZAR, O.: Growth in the Muskat problem, preprint, [arXiv:1904.00294](https://arxiv.org/abs/1904.00294), 2019
41. GRANERO-BELINCHÓN, R., SHKOLLER, S.: Well-posedness and decay to equilibrium for the Muskat problem with discontinuous permeability. *Trans. Amer. Math. Soc.* **372**(4), 2255–2286, 2019
42. GUO, Y., HALLSTROM, C., SPIRN, D.: Dynamics near unstable, interfacial fluids. *Commun. Math. Phys.* **270**(3), 635–689, 2007
43. HELE-SHAW, H.S.: The flow of water. *Nature* **58**, 34–36, 1898
44. HELE-SHAW, H.S.: On the motion of a viscous fluid between two parallel plates. *Trans. R. Inst. Nav. Arch.* **40**, 218, 1898
45. HÖRMANDER, L.: *Lectures on Nonlinear Hyperbolic Differential Equations*, volume 26 of *Mathématiques & Applications (Berlin) [Mathematics & Applications]*. Springer, Berlin, 1997
46. HUNTER, J.K., IFRIM, M., TATARU, D.: Two dimensional water waves in holomorphic coordinates. *Commun. Math. Phys.* **346**, 483–552, 2016
47. LANNES, D.: Well-posedness of the water waves equations. *J. Am. Math. Soc.* **18**(3), 605–654, 2005
48. LIONS, J.L., MAGENES, E.: *Problèmes aux limites non homogènes et applications*, vol. 1. Dunod, Paris 1968
49. LEONI, G., TICE, I.: Traces for homogeneous Sobolev spaces in infinite strip-like domains. *J. Funct. Anal.* **277**(7), 2288–2380, 2019

50. MATIOC, B.-V.: The muskat problem in 2D: equivalence of formulations, well-posedness, and regularity results. *Anal. PDE* **12**(2), 281–332, 2018
51. MATIOC, B.-V.: Viscous displacement in porous media: the Muskat problem in 2D. *Trans. Am. Math. Soc.* **370**(10), 7511–7556, 2018
52. MÉTIVIER, G.: *Para-Differential Calculus and Applications to the Cauchy Problem for Nonlinear Systems*, volume 5 of *Centro di Ricerca Matematica Ennio De Giorgi (CRM) Series*. Edizioni della Normale, Pisa, 2008
53. MUSKAT, M.: Two fluid systems in porous media. The encroachment of water into an oil sand. *Physics* **5**, 250–264, 1934
54. NGUYEN, H.Q.: A sharp Cauchy theory for 2D gravity–capillary water waves. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **34**(7), 1793–1836, 2017
55. NGUYEN, H.Q.: On well-posedness of the Muskat problem with surface tension. [arXiv:1907.11552](https://arxiv.org/abs/1907.11552) [math.AP], 2019
56. PERNAS-CASTAÑO, T.: Local-existence for the inhomogeneous Muskat problem. *Non-linearity* **30**(5), 2063, 2017
57. SAFFMAN, P.G., TAYLOR, G.: The penetration of a fluid into a porous medium or Hele-Shaw cell containing a more viscous liquid. *Proc. R. Soc. Lond. Ser. A* **245**, 312–329, 1958. (2 plates)
58. SAFONOV, M.V.: Boundary estimates for positive solutions to second order elliptic equations. [arXiv:0810.0522](https://arxiv.org/abs/0810.0522), 2008
59. SIEGEL, M., CAFLISCH, R., HOWISON, S.: Global existence, singular solutions, and Ill-posedness for the Muskat problem. *Commun. Pure Appl. Math.* **57**, 1374–1411, 2004
60. STRICHARTZ, R.S.: “Graph paper” trace characterizations of functions of finite energy. *J. Anal. Math.* **128**, 239–260, 2016
61. WU, S.: Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent. Math.* **130**, 39–72, 1997
62. WU, S.: Well-posedness in Sobolev spaces of the full water wave problem in 3-D. *J. Am. Math. Soc.* **12**(2), 445–495, 1999
63. YI, F.: Local classical solution of Muskat free boundary problem. *J. Partial Differ. Equ.* **9**, 84–96, 1996

HUY Q. NGUYEN & BENOÎT PAUSADER

Department of Mathematics,

Brown University,

Providence

RI

02912 USA.

e-mail: hnguyen@math.brown.edu

and

BENOÎT PAUSADER

e-mail: benoit.pausader@math.brown.edu

(Received July 8, 2019 / Accepted February 3, 2020)

© Springer-Verlag GmbH Germany, part of Springer Nature (2020)