ELSEVIER

Contents lists available at ScienceDirect

Carbohydrate Research

journal homepage: www.elsevier.com/locate/carres



Synthesis of 2-azido-2-deoxy- and 2-acetamido-2-deoxy-D-manno derivatives as versatile building blocks



Catherine Alex^a, Satsawat Visansirikul^{a,b}, Yongzhen Zhang^c, Jagodige P. Yasomanee^a, Jeroen Codee^c, Alexei V. Demchenko^{a,*}

- ^a Department of Chemistry and Biochemistry, University of Missouri St. Louis, One University Boulevard, St. Louis, Missouri, 63121, USA
- b Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Mahidol University, 447 Sri-Ayuddhaya Road, Rajathevee, Bangkok, 10400, Thailand
- ^c Bio-organic Synthesis, Leiden Institute of Chemistry, Leiden University, Einsteinweg 55, 2333 CC, Leiden, the Netherlands

ABSTRACT

Reported herein is the synthesis of a number of building blocks of 2-amino-2-deoxy-p-mannose from common p-glucose precursors.

1. Introduction

All oligosaccharides comprise an oligomeric sequence wherein monomers are linked via O-glycosidic linkages. This type of linkage is obtained by a glycosylation reaction, which, in spite of significant progress, remains challenging due to the requirement to achieve high stereocontrol [1,2] and suppress side reactions [3,4]. Due to significant advances, many glycans can now be obtained by using chemical methods and automated platforms in solution and on solid phases [5]. However, poor accessibility to sugar building blocks hampers development of all synthetic approaches. Things are further complicated because "unlike the synthesis of peptides and oligonucleotides, there are no universal building blocks or methods for the synthesis of all glycans." [6] In spite of general improvements in application of protecting groups in the mainstream carbohydrate research [7-12], building blocks for the introduction of uncommon (rare) sugars remain largely underdeveloped or not available at all [13]. Reported herein are reliable and scalable procedures for the synthesis of mannosamine building blocks.

N-Acetylmannosamine (ManNAc) is the biosynthetic precursor of N-acetylneuraminic acid [14,15] and a constituent of many bacterial glycans [16–18]. Hence, a considerable interest has been devoted to the synthesis of ManNAc and its derivatives. Nevertheless, mannosamine remains prohibitively expensive as the starting material both for the laboratory and industrial application. The total synthesis from p-arabinose reported by O'Neill [19] and alkaline-mediated epimerization of C-2 in N-acetylglucosamine by Spivak and Roseman [20] are the classics of the synthesis of ManNAc. More recently, building blocks of ManNAc have been prepared via the inversion of configuration at C-2 of p-glucose with nitrogen nucleophiles [21–23].

Reported herein are two modes to obtain ManNAc derivatives from methyl 4,6-O-benzylidene- α and β -D-glucopyranosides 1 and 6 [24,25]. As previously shown by Knapp et al., [23] high regioselectivity of sulfonation of diol 1 with triflic anhydride in pyridine toward the C-2 position could be achieved at low temperature. The resulting 2-O-triflate was then subjected to the nucleophilic displacement with sodium azide in DMF to afford 2-azido-2-deoxy-p-mannopyranoside intermediate 2 (Scheme 1). This two-step protocol represent a common approach to obtain rare or inaccessible aminosugars [26]. The latter can be used for a number of pathways that is showcased herein by two synthetic transformations. First, treatment of compound 2 with zinc in the presence of acetic anhydride provided direct transformation of the 2-azido into 2-acetamido group. As a result, ManNAc derivative 3 was obtained in 78% yield. It should be noted that derivatives 2 and 3 can be used as glycosyl acceptors or as intermediates to access other functionalized derivatives. In addition, while we were specifically targeting 2-azido and 2-acetamido derivatives, many other N-protecting groups can be introduced after the azide reduction step, if so desired.

We also performed acetolysis of intermediate 2 with acetic anhydride in the presence of H_2SO_4 . This reaction led to concomitant replacement of methyl glycoside and benzylidene acetal with acetyl groups. As a result, tetraacetate 4 was smoothly produced in 80% yield. The latter derivative can be used as a versatile intermediate for further functionalization. Herein, tetraacetate 4 was reacted with ethane thiol in the presence of BF_3 etherate. As a result, ethylthio glycoside 5 was produced in 58% yield. While thioglycosides can be used as versatile building blocks and as glycosyl donors, the synthesis of other types of glycosyl donors or building blocks can be readily achieved using similar

E-mail address: demchenkoa@umsl.edu (A.V. Demchenko).

^{2.} Results and Discussion

^{*} Corresponding author.

Scheme 1. The synthesis of $\mbox{\tiny D-mannosamine}$ building blocks 2–5 from D-gluco derivative 1.

synthetic routes.

The second approach to produce mannosamine derivatives described herein involves adaptation of the O-trichloroacetimidoyl protecting group migration approach developed by van der Marel and coworkers for the synthesis of other aminosugar targets [27,28]. This was affected by regioselective O-imidoylation at the C-3 position of diol 6 (Scheme 2). Preferential substitution at the C-3 position was achieved with trichloroacetonitrile in the presence of DBU at room temperature. Subsequent in-situ sulfonation with triflic anhydride in pyridine at C-2 followed by the treatment with DIPEA led to the formation of 2,3-N,Ooxazoline 7, all in one pot. The intermediate 7 was isolated in 44% yield over three steps. The fair yield is due to marginal regioselectivity obtained during the first step and the necessity to separate product 7 from the resulting side products. The oxazoline ring in derivative 7 was hydrolyzed by reaction with sodium hydroxide. The amine group in the resulting partially protected derivative was then chemoselectively Nacetylated to afford ManNAc derivative 8. While we were specifically interested in obtaining 2-acetamido derivatives, many other N-

protecting groups can be introduced after the oxazoline ring removal, if so desired. The resulting 3-OH derivative 8 can be used as a glycosyl acceptor directly, or employed in further functionalization per specific synthetic targets.

In summary, presented herein is an efficient synthesis of a variety of 2-amino-2-deoxy derivatives of p-mannose. We investigated gram-scale syntheses, and the developed protocols appear to be sufficiently straightforward and reliable to be attempted at a greater scale if needed. The utility of these building blocks has been demonstrated by the synthesis of glycosyl acceptors 2, 3 and 8 as well as thioglycoside donor 5. While we were specifically targeting 2-azido and 2-acetamido derivatives, many other *N*-protecting groups can be introduced, if necessary. Further application of these versatile building blocks, both as glycosyl donors and as glycosyl acceptors, are currently underway in our laboratories.

3. Experimental

General methods. Solvents were purified according to standard procedures [29]. Reactions were performed using commercial reagents (Millipore Sigma, Carbosynth, TCI, or Acros), and monitored by TLC on Kieselgel 60 F₂₅₄ (EM Science). Compounds were detected by UV light and by charring with 10% sulfuric acid in methanol. Solvents were removed under reduced pressure at < 40 °C. Column chromatography was performed on silica gel 60 (70–230 mesh). Optical rotations were measured at 'Jasco P-2000' polarimeter at 22 °C. $^{1}{\rm H}$ and $^{13}{\rm C}$ NMR spectra were recorded at 300 and 75 MHz, respectively, with a Bruker Avance spectrometer. The chemical shifts are referenced to the signal of the residual CHCl₃ ($\delta_{\rm H}=7.27$ ppm, $\delta_{\rm C}=77.23$ ppm) for solutions in CDCl₃. COSY and HMQC experiments were done to aid nuclei assignments. HRMS were recorded with a JEOL MStation (JMS-700) Mass Spectrometer. Melting points were measured at Digimelt MPA 160 melting point apparatus.

Methyl 2-azido-4,6-*O*-benzylidene-2-deoxy-α-p-mannopyranoside (2). Pyridine (14 mL, 173.8 mmol) and trifluoromethanesulfonic anhydride (Tf₂O, 0.65 mL, 3.87 mmol) were added at -30 °C under argon to a stirred solution of methyl 4,6-*O*-benzylidene-α-p-glucopyranoside (1, 0.99 g, 3.52 mmol) in CH₂Cl₂ (14 mL) and the resulting mixture was stirred at -30 °C for 3 h. The volatiles were removed under reduced pressure, and the residue was dried in *vacuo* for 2 h. Sodium azide (NaN₃, 1.14 g, 17.6 mmol) was added to a solution of the residue in DMF (15 mL) and the resulting mixture was stirred at 75 °C for 12 h. After that, the reaction mixture was allowed to cool to rt, volatiles were removed under reduced pressure, and the residue was

Scheme 2. The synthesis of ${\tt D\text{-}mannosamine}$ building block 8.

purified by column chromatography on silica gel (EtOAc-hexane gradient elution) to afford the title compound (0.874 g, 2.84 mmol) in 80% as a pale-yellow syrup. Analytical data for **2**: R_f 0.7 (EtOAc-hexanes, 2/3, v/v). [α]_D +70.8 (c 1.0, CHCl₃); ¹H NMR: δ , 2.60 (s, 1H, 3-OH), 3.39 (s, 3H, OCH₃), 3.72–3.87 (m, 2H, H-5, 6a), 3.91 (dd, 1H, $J_{4,5}$ = 7.6 Hz, H-4), 3.99 (dd, 1H, $J_{2,3}$ = 3.9 Hz, H-2), 4.21–4.32 (m, 2H, $J_{3,4}$ = 7.6 Hz, H-3, 6b), 4.70 (d, 1H, $J_{1,2}$ = 1.1 Hz, H-1), 5.58 (s, 1H, CHPh), 7.34–7.44 (m, 3H, aromatic), 7.45–7.54 (m, 2H, aromatic) ppm; ¹³C NMR: δ , 55.3 (OCH₃), 63.4 (C-5), 63.8 (C-2), 68.8 (C-6), 69.0 (C-3), 79.1 (C-4), 100.2 (C-1), 102.4 (CHPh), 126.4, 128.5, 129.4, 137.2 (6C, aromatic) ppm; HR-FAB MS: [M+Na] + calcd for C₁₄H₁₇N₃O₅Na, 330.1066; found, 330.1069. Partial characterization data [α]_D²⁵ +69.5 (c 1.1, CHCl₃) [30] and ¹³C NMR for **2** were reported previously [22].

2-acetamido-4,6-O-benzylidene-2-deoxy-α-D-mannopyranoside (3). Acetic acid (2.0 mL, 30.7 mmol), acetic anhydride (Ac₂O, 2.0 mL, 18.6 mmol), and zinc dust (2.0 g, 26.9 mmol) were added to a solution of methyl 2-azido-4,6-O-benzylidene-2-deoxy-α-Dmannopyranoside (2, 0.83 g, 2.72 mmol) in THF (10 mL) and the resulting mixture was stirred under argon for 1 h at rt. The solids were filtered off through a pad of Celite, volatiles were removed under reduced pressure, and the residue was purified by column chromatography on silica gel (MeOH-CH2Cl2 gradient elution) to afford the title compound (0.68 g, 2.10 mmol) as white crystals in 78% yield. Analytical data for 3: m.p. 227–228 °C (CH_2Cl_2 – hexanes); $[\alpha]_D$ + 40.5 (c 1.0, CHCl₃); R_f 0.6 (MeOH/CH₂Cl₂, 1/9, v/v); ¹H NMR: δ , 2.07 (s, 3H, COCH₃), 3.38 (s, 3H, OCH₃), 3.66 (dd, 1H, $J_{4.5} = 9.4$ Hz, H-4), 3.75-3.92 (m, 2H, H-5, 6a), 4.25-4.34 (m, 2H, H-3, 6b), 4.45 (br ddd, 1H, H-2), 4.77 (d, 1H, $J_{1,2} = 0.8$ Hz, H-1), 5.59 (s, 1H, CHPh), 5.85 (br d, 1H, $J_{2,NH} = 7.0$ Hz, NH), 7.33–7.39 (m, 3H, aromatic), 7.45–7.51 (m, 2H, aromatic) ppm; 13 C NMR: δ , 23.5 (COCH₃), 53.6 (C-2), 55.4 (OCH₃), 62.8 (C-5), 67.2 (C-3), 68.9 (C-6), 79.7 (C-4), 100.8 (C-1), 102.5 (CHPh), 126.4, 128.5, 129.4, 137.1 (6C, aromatic), 171.8 (C=O) ppm; HR-FAB MS: $[M + Na]^+$ calcd for $C_{16}H_{21}NO_6Na$, 346.1267; found, 346.1261.

1,3,4,6-Tetra-O-acetyl-2-azido-2-deoxy-p-mannopyranose (4). Sulfuric acid (0.2 mL, 3.73 mmol) was added dropwise to a solution of 2 (0.76 g, 2.47 mmol) in acetic anhydride (10.0 mL, 106.0 mmol) at 0 °C. The external cooling was removed, and the resulting mixture was stirred under argon for 4 h at rt. Sat. aq. NaHCO3 was added carefully to the reaction mixture until gas evolution has ceased. EtOAc (30 mL) was added, and the resulting mixture was washed with sat. aq. NaHCO3. The organic layer was separated, dried over Na2SO4, and filtered. Volatiles were removed under reduced pressure and the residue was co-evaporated with toluene. The resulting residue was purified by column chromatography on silica gel (ethyl acetate-hexanes gradient elution) to afford the title compound (0.74 g, 1.97 mmol) as a pale-yellow amorphous solid in 80% yield ($\alpha/\beta = 7/1$). Analytical data for 4: R_f 0.5 (EtOAc/hexanes, 1/1, v/v); ¹H NMR: δ , 2.11, 2.15, 2.17, 2.22 (4 s, 12H, $4 \times COCH_3$, 4.04-4.20 (m, 3H, H-2, 5, 6a), 4.30 (dd, 1H, J = 4.5, 12.4 Hz, H-6a), 5.39–5.50 (m, 2H, H-3, 4), 6.17 (d, 1H, J = 1.9 Hz, H-1) ppm; 13 C NMR: δ , 20.6, 20.7, 20.8, 20.9, 60.5, 61.8, 65.3, 70.6, 70.7, 91.3, 168.3, 169.4, 170.1, 170.8 ppm; HR-FAB MS: [M+Na] + calcd for C₁₄H₁₉N₃O₉Na, 396.1019; found, 396.1024.

Ethyl 3,4,6-tri-O-acetyl-2-azido-2-deoxy-1-thio-α-p-mannopyranoside (5). Boron trifluoride diethyl etherate (3.05 mL, 24.05 mmol) was added dropwise to a solution of 4 (2.24 g, 6.01 mmol) and ethane thiol (0.89 mL, 12.03 mmol) in CH₂Cl₂ (50 mL) at 0 °C. The external cooling was then removed and the resulting mixture was stirred under argon for 16 h at rt. After that, the reaction mixture was diluted with CH₂Cl₂ (~150 mL) and was washed with H₂O (50 mL), sat. aq. NaHCO₃ (50 mL), and H₂O (2 x 50 mL). The organic layer was separated, dried, and volatiles were removed under reduced pressure. The residue was purified by column chromatography on silica gel (ethyl acetate-hexanes gradient elution) to afford the title compound (1.31 g, 3.49 mmol) as a pale-yellow syrup in 58% yield. Analytical data for 5: R_f 0.7 (EtOAc/hexanes, 1/1, v/v); $[\alpha]_D$ +26.6 (c 1.0, CHCl₃); ¹H NMR: δ , 1.35 (t, 3H,

J=7.4 Hz, SCH₂CH₃), 2.09, 2.13, 2.13 (3 s, 9H, COCH₃), 2.59–2.78 (m, 2H, SCH₂CH₃), 4.09–4.17 (m, 2H, H-2, 6a), 4.29–4.42 (m, 2H, H-5, 6b), 5.30–5.42 (m, 3H, H-1, 3, 4) ppm; ¹³C NMR: δ, 14.7, 20.6, 20.7, 20.8, 25.5, 62.1, 62.8, 66.1, 68.8, 71.3, 82.3, 169.6, 170.0, 170.8 ppm; HR-FAB MS: [M+Na] + calcd for C₁₄H₂₁N₃O₇SNa, 398.0998; found, 398.1003.

4,6-O-benzylidene-2-deoxy-2N,3O-(2,2,2-trichloro-Methyl ethylidene)-β-D-mannopyranoside (7). Trichloroacetonitrile (CCl₃CN, 535 µL, 5.33 mmol) and activated molecular sieves (3 Å, 1.20 g) were added to a solution of 6 (1.25 g, 4.44 mmol) in CH₂Cl₂ (50 mL) and the resulting mixture was stirred under argon for 15 min at rt. After that, DBU (67 µL, 0.44 mmol) was added, and the reaction mixture was stirred for 5 h at rt. The mixture was then cooled to -30 °C, pyridine (1.79 mL, 22.2 mmol) and Tf₂O (897 μ L, 5.33 mmol) were added, and the resulting mixture was stirred under argon for 2 h at - 30 °C. The cooling was removed, N,N-diisopropylethylamine (DIPEA, 7.7 mL, 44.4 mmol) was added, and the bright-red colored mixture was stirred under argon for 12 h at rt. The volatiles were removed under reduced pressure and the residue was purified by column chromatography on silica gel (EtOAc-hexanes gradient elution) to afford the title compound (805 mg, 1.97 mmol) as colorless crystals in 44% yield. Analytical data for 7: R_f 0.6 (EtOAc/hexanes, 2/3, v/v); m.p. 164–165 °C (from aq. MeOH); $[α]_D$ –76.5 (*c* 1.0, CHCl₃); ¹H NMR: δ, 3.46 (s, 3H, OCH₃), 3.64 (m, 1H, $J_{5,6a} = 10.1$ Hz, $J_{5,6b} = 4.4$ Hz, H-5), $3.72 \text{ (dd, 1H, } J_{6a,6b} = 10.0 \text{ Hz, H-6a), } 4.38 \text{ (dd, 1H, H-6b), } 4.56-5.65$ (m, 2H, H-2, 4), 4.92 (d, 1H, $J_{1,2} = 3.5$ Hz, H-1), 5.02 (dd, 1H, $J_{2.3} = 7.4, J_{3.4} = 9.8 \text{ Hz}, \text{ H-3}, 5.61 \text{ (s, 1H, CHPh)}, 7.34-7.42 \text{ (m, 3H, CHPh)}$ aromatic), 7.46–7.55 (m, 2H, aromatic) ppm; 13 C NMR: δ , 56.1 (OCH₃), 64.5 (C-5), 67.4 (C-2), 70.3 (C-6), 76.9 (C-4), 77.4 (CCl₃), 82.6 (C-3), 97.5 (C-1), 101.8 (CHPh), 126.4, 128.5, 129.5, 136.9 (6C, aromatic), 164.7 (C=N) ppm; HR-FAB MS: $[M+Na]^+$ calcd for $C_{16}H_{16}Cl_3NO_5Na$, 429.9992; found, 429.9980.

2-acetamido-4,6-O-benzylidene-2-deoxy-β-D-mannopyranoside (8). 1 M aq. NaOH (10 mL) was added dropwise to a solution of 7 (559 mg, 1.37 mmol) in MeOH (30 mL) and the resulting mixture was kept for 12 h at reflux (~65 °C). The reaction mixture was allowed to cool to rt, the volatiles were removed under reduced pressure, and the residue was dried in vacuo for 1 h. The crude residue was dissolved in MeOH (15 mL), Ac2O (0.65 mL, 6.91 mmol) was added, and the resulting mixture was stirred under argon for 2 h at rt. After that, volatiles were removed, and the residue was purified by column chromatography on silica gel (MeOH-CH₂Cl₂ gradient elution) to afford the title compound (252 mg, 0.78 mmol) as a white amorphous solid in 58% yield. Analytical data for 8: R_f 0.5 (MeOH/CH₂Cl₂, 1/9, v/v); m.p. 217–219 °C (from CH_2Cl_2 -hexane); $[\alpha]_D$ – 42.0 (c 1.0, $CHCl_3$); ¹H NMR: δ , 2.11 (s, 3H, COCH₃), 3.46 (m, 1H, $J_{5,6a} = 10.2$, $J_{5,6b} = 4.9$ Hz, H-5), 3.52 (s, 3H, OCH₃), 3.70 (dd, 1H, $J_{4,5} = 9.4$ Hz, H-4), 3.80 (dd, 1H, J_{6a} $_{6b}$ = 10.2 Hz, H-6a), 4.00 (dd, 1H, $J_{3,4}$ = 9.5 Hz, H-3), 4.34 (dd, 1H, H-6b), 4.52 (br dd, 1H, $J_{2,3} = 4.0$ Hz, H-2), 4.64 (d, 1H, $J_{1,2} = 1.7$ Hz, H-1), 5.57 (s, 1H, CHPh), 5.98 (br d, 1H, $J_{2,NH} = 6.0$ Hz, NH), 7.30–7.40 (m, 3H, aromatic), 7.44–7.53 (m, 2H, aromatic) ppm; 13 C NMR: δ , 23.3 (COCH₃), 54.9 (C-2), 57.3 (OCH₃), 67.4 (C-5), 68.6 (C-6), 71.6 (C-3), 79.7 (C-4), 100.6 (C-1), 102.3 (CHPh), 126.4, 128.4, 129.3, 137.1 (6C, aromatic), 173.8 (C=O) ppm; HR-FAB MS: [M+Na]⁺ calcd for C₁₆H₂₁NO₆Na, 346.1267; found, 346.1254.

Author contributions

The manuscript was written through contributions of all authors./ All authors have given approval to the final version of the manuscript.

Declaration of competing interest

The authors have no conflicts to declare.

Acknowledgement

This work was supported by the National Science Foundation (CHE-1800350) and National Institute of General Medical Sciences (GM120673).

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.carres.2019.107900.

References

- L.K. Mydock, A.V. Demchenko, Mechanism of chemical O-glycosylation: from early studies to recent discoveries, Org. Biomol. Chem. 8 (2010) 497–510.
- [2] D. Crich, Mechanism of a chemical glycosylation reaction, Acc. Chem. Res. 43 (2010) 1144–1153.
- [3] A.V. Demchenko (Ed.), Handbook of Chemical Glycosylation: Advances in Stereoselectivity and Therapeutic Relevance, Wiley-VCH, Weinheim, Germany, 2008
- [4] H.M. Christensen, S. Oscarson, H.H. Jensen, Common side reactions of the glycosyl donor in chemical glycosylation, Carbohydr. Res. 408 (2015) 51–95.
- [5] M. Panza, S.G. Pistorio, K.J. Stine, A.V. Demchenko, Automated chemical oligosaccharide synthesis: novel approach to traditional challenges, Chem. Rev. 118 (2018) 8105–8150.
- [6] L.C. Hsieh-Wilson, M.E. Griffin, Improving biologic drugs via total chemical synthesis. Science 342 (2013) 1332–1333.
- [7] S.M. Polyakova, A.V. Nizovtsev, R.A. Kuznetskiy, N.V. Bovin, New protecting groups in oligosaccharide synthesis, Russ. Chem. Bull. (2015) 974–989.
- [8] K. Ágoston, H. Streicher, P. Fügedi, Orthogonal protecting group strategies in carbohydrate chemistry, Tetrahedron: Asymmetry 27 (2016) 707–728.
- [9] M. Jager, A.J. Minnaard, Regioselective modification of unprotected glycosides, Chem. Commun. 52 (2016) 656–664.
- [10] S.S. Kulkarni, C.C. Wang, N.M. Sabbavarapu, A.R. Podilapu, P.H. Liao, S.C. Hung, One-pot" protection, glycosylation, and protection-glycosylation strategies of carbohydrates, Chem. Rev. 118 (2018) 8025–8104.
- [11] A.G. Volbeda, G.A. van der Marel, J.D.C. Codée, Protecting group strategies in carbohydrate chemistry, in: S. Vidal (Ed.), Protecting Groups – Strategies and Applications in Carbohydrate Chemistry, Wiley-VCH, Weinheim, 2019, pp. 1–28.
- [12] T. Wang, A.V. Demchenko, Synthesis of carbohydrate building blocks via regioselective uniform protection/deprotection strategies, Org. Biomol. Chem. 17 (2019) 4934–4950.
- [13] M. Emmadi, S.S. Kulkarni, Recent advances in synthesis of bacterial rare sugar building blocks and their applications, Nat. Prod. Rep. 31 (2014) 870–879.
- [14] A. Varki, Biological roles of oligosaccharides: all of the theories are correct, Glycobiology 3 (1993) 97–130.

- [15] T. Angata, A. Varki, Chemical diversity in the sialic acids and related a-keto acids: an evolutionary perspective, Chem. Rev. 102 (2002) 439–469.
- [16] E. Danieli, D. Proietti, G. Brogioni, M.R. Romano, E. Cappelletti, M. Tontini, F. Berti, L. Lay, P. Costantino, R. Adamo, Synthesis of Staphylococcus aureus type 5 capsular polysaccharide repeating unit using novel L-FucNAc and D-FucNAc synthons and immunochemical evaluation, Bioorg, Med. Chem. 20 (2012) 6403–6415.
- [17] I.A. Gagarinov, T. Fang, L. Liu, A.D. Srivastava, G.J. Boons, Synthesis of Staphylococcus aureus type 5 trisaccharide repeating unit: solving the problem of lactamization, Org. Lett. 17 (2015) 928–931.
- [18] S. Visansirikul, J.P. Yasomanee, P. Pornsuriyasak, M.N. Kamat, N.M. Podvalnyy, C.P. Gobble, M. Thompson, S.A. Kolodziej, A.V. Demchenko, A concise synthesis of the repeating unit of capsular polysaccharide Staphylococcus aureus type 8, Org. Lett. 17 (2015) 2382–2384.
- [19] A.N. O'Neill, Asymmetric synthesis of D- and L-mannosamine, Can. J. Chem. 37 (1959) 1747–1753.
- [20] C.T. Spivak, S. Roseman, Preparation of N-Acetyl-D-mannosamine (2-acetamido-2-deoxy-D-mannose) and D-mannosamine hydrochloride (2-amino-2-deoxy-D-mannose), J. Am. Chem. Soc. 81 (1959) 2403–2404.
- [21] V. Draghetti, L. Poletti, D. Prosperi, L. Lay, A convenient multigram preparation of functionalized 2-azido-2-deoxy-D-mannose as a useful orthogonally protected building block for oligosaccharide synthesis, J. Carbohydr. Chem. 20 (2001) 813–819.
- [22] A. Popelova, K. Kefurt, M. Hlavackova, J. Moravcova, A concise synthesis of 4nitrophenyl 2-azido-2-deoxy- and 2-acetamido-2-deoxy-D-mannopyranosides, Carbohydr. Res. 340 (2005) 161–166.
- [23] S. Knapp, P.J. Kukkola, S. Sharma, T.G.M. Dhar, A.B.J. Naughton, Amino alcohol and amino sugar synthesis by benzoylcarbamate cyclization, J. Org. Chem. 55 (1990) 5700–5710.
- [24] M.E. Evans, Methyl 4,6-O-Benzylidene- α and - β -D-Glucosides, Carbohydr. Res. 21 (1972) 473–475.
- [25] P. Pornsuriyasak, J.P. Yasomanee, E.I. Balmond, C.D. Meo, A.V. Demchenko, Synthesis of 4,6-O-benzylidene acetals of methyl α-D-glucopyranoside, ethyl 1-thio-β-D-gluco- and galactopyranoside, in: G. van der Marel, J. Codee (Eds.), Carbohydrate Chemistry: Proven Synthetic Methods, vol. 2, CRC Press, 2014, pp. 189–196.
- [26] S.R. Sanapala, S.S. Kulkarni, Expedient route to access rare deoxy amino L-sugar building blocks for the assembly of bacterial glycoconjugates, J. Am. Chem. Soc. 138 (2016) 4938–4947.
- [27] L.J. van den Bos, J.D.C. Codée, J.H. van Boom, H.S. Overkleeft, G.A. van der Marel, A novel strategy towards the synthesis of orthogonally functionalised 4-aminoglycosides, Org. Biomol. Chem. 1 (2003) 4160–4165.
- [28] A.E. Christina, V.M. Blas Ferrando, F. de Bordes, W.A. Spruit, H.S. Overkleeft, J.D. Codee, G.A. van der Marel, Multigram-scale synthesis of an orthogonally protected 2-Acetamido-4-amino-2,4,6-trideoxy-D-galactose (AAT) building block, Carbohydr. Res. 356 (2012) 282–287.
- [29] W.L.F. Armarego, C.L.L. Chai, Purification of Laboratory Chemicals, fifth ed., Butterworth-Heinemann, Amsterdam - Boston - London, 2003.
- [30] T. Sugawara, K. Igarashi, Synthesis of a trisaccharide component of the capsular polysaccharide of Streptococcus pneumoniae type 19F, Carbohydr. Res. 172 (1988) 195–207.