



Parameters in Categorified Quantum Groups

Aaron D. Lauda¹

Received: 21 December 2018 / Accepted: 3 April 2019 / Published online: 12 April 2019
© Springer Nature B.V. 2019

Abstract

In this note we give explicit isomorphisms of 2-categories between various versions of the categorified quantum group associated to a simply-laced Kac-Moody algebra. These isomorphisms are convenient when working with the categorified quantum group. They make it possible to translate results from the \mathfrak{gl}_n variant of the 2-category to the \mathfrak{sl}_n variant and transfer results between various conventions in the literature. We also extend isomorphisms of finite type KLR algebras for different choices of parameters to the level of 2-categories.

Keywords Categorification · Quantum group · KLR-algebra

1 Introduction

Brundan has shown that the categorified quantum group associated to \mathfrak{g} is essentially unique [2]. There remain several choices to be made:

- a choice of scalars Q determining the KLR algebra R_Q categorifying the positive half of $\dot{U}(\mathfrak{g})$,
- overall scalings of generating 2-morphisms that affect the precise form of the relations. These parameters include the values of degree zero bubbles and the dependence of these choices on the weight $\lambda \in X$.

Different choices for the parameters can result in different behaviors of the resulting 2-categories.

Skew Howe duality has proven to be a powerful tool in higher representation theory and its applications to link homology [3, 4, 13, 15, 16]. However, this technique is fundamentally a \mathfrak{gl}_n phenomenon, though many of the results in the literature express the results in terms of \mathfrak{sl}_n . Hence, one has to move between the categorified quantum groups for \mathfrak{gl}_n and \mathfrak{sl}_n . This creates complications because in the current literature the categorifications of \mathfrak{sl}_n and \mathfrak{gl}_n are not connected in a straightforward manner. The parameters in each case seem oddly related. Further, many results are formulated for some fixed choice of parameters in the categorified

Presented by: Alistair Savage

✉ Aaron D. Lauda
lauda@usc.edu

¹ Department of Mathematics, University of Southern California, Los Angeles, CA, USA

quantum group. For example, in type A the connection with Soergel bimodules relies on one choice of coefficients [14], while the connection to cohomology rings of Grassmannians utilizes another [8, 10, 11]. Translating results from one choice to another can be rather complicated since one must paste together and extend the needed rescaling 2-functors from partial results scattered throughout the literature, and some of these isomorphisms require weight dependent rescalings that depend on values of weights modulo 4.

In this note we define a general version of the categorified quantum group and give explicit 2-functors for translating results from one formulation to the other. This allows for an immediate translation from \mathfrak{gl}_n coefficients to \mathfrak{sl}_n coefficients with an explicit rescaling 2-functor collecting various results from the literature. We also make explicit certain isomorphisms that can be defined between KLR-algebras R_Q and $R_{Q'}$ for different choices of scalars when the underlying graph of the simply-laced Kac-Moody algebra is a tree, in particular, a Dynkin diagram. Here we show how to extend these isomorphisms to isomorphisms of 2-categories for the corresponding categorified quantum groups. This map hasn't appeared in the literature before. The author found himself often reconstructing these results and decided to produce this note.

1.1 Versions of Categorified Quantum Groups

In [8] a 2-category $\mathcal{U}(\mathfrak{g})$ was defined for any symmetrizable Kac-Moody algebra \mathfrak{g} . They conjectured that this 2-category categorifies the integral form of Lusztig's idempotent version $\dot{\mathcal{U}}_q(\mathfrak{g})$ of the quantum group and proved the conjecture for $\mathfrak{g} = \mathfrak{sl}_n$, following earlier work for $\mathfrak{g} = \mathfrak{sl}_2$ from [10]. Later Webster proved the conjecture in general [18]. An important feature of the 2-category $\mathcal{U}(\mathfrak{g})$ is its diagrammatic nature, making it possible to represent 2-morphisms using a planar diagrammatic calculus. However, this 2-category is only cyclic up to a sign, meaning that planar deformations of a diagram representing a 2-morphism only gives rise to scalar multiples of the same 2-morphism.

The KLR algebra governs the upward oriented strands in $\mathcal{U}(\mathfrak{g})$. In [9, 17] more general coefficients Q for KLR algebras were introduced and a more general non-cyclic 2-category $\mathcal{U}_Q(\mathfrak{g})$ was defined in [5] for an arbitrary KLR algebra R_Q . Closely related categories were also studied in [17]. Composing biadjunction units and counits with other generators produces certain endomorphisms of the identity map in each weight λ called *dotted bubbles*. With the appropriate number of dots, these dotted bubbles have degree zero and take some value called *bubble parameters* in the ground field \mathbb{k}^\times :

$$\begin{array}{ccc} \lambda & & \lambda \\ \text{---} & = c_{i,\lambda}^+, & \text{---} \\ \lambda_i - 1 & \text{---} & \lambda_i - 1 \end{array} \quad (1.1)$$

In the 2-category $\mathcal{U}_Q(\mathfrak{g})$ all these bubble parameters are normalized to 1.

A categorification $\mathcal{U}(\mathfrak{gl}_n)$ of quantum \mathfrak{gl}_n and its Schur quotients were defined in [14]. Unlike the 2-category $\mathcal{U}_Q(\mathfrak{g})$, this 2-category is cyclic. However, the bubble parameters in $\mathcal{U}(\mathfrak{gl}_n)$ take values of ± 1 depending explicitly on the $\mathfrak{gl}(n)$ weights. Further, the relations that give isomorphisms lifting the \mathfrak{sl}_2 relations have a significant sign difference from $\mathcal{U}_Q(\mathfrak{g})$ that force the values of degree zero bubbles to alternate sign as the weight changes along \mathfrak{sl}_2 strings. Additionally, the 2-category $\mathcal{U}(\mathfrak{gl}_n)$ has a specific choice of scalars Q for the KLR algebra that depends on an orientation of the underlying quiver.

In [12, Section 3.9] a version of the \mathfrak{sl}_2 2-category $\mathcal{U}_\beta(\mathfrak{sl}_2)$ was defined with arbitrary values of bubbles parameters and a more general \mathfrak{sl}_2 relation

that generalizes the corresponding relation from the $\mathcal{U}(\mathfrak{gl}_n)$ when $\beta_{i,\lambda} = +1$ and $\mathcal{U}_Q(\mathfrak{g})$ when $\beta_{i,\lambda} = -1$ for all $i \in I$ and $\lambda \in X$. For any choice of parameters $\beta_{i,\lambda}$, the 2-category $\mathcal{U}_\beta(\mathfrak{sl}_2)$ is isomorphic to the original 2-category $\mathcal{U}(\mathfrak{sl}_2)$ from [10], although this isomorphism requires rescalings of caps and cups that depend on the weight modulo 4. These parameters $\beta_{i,\lambda}$ control the values of higher degree bubbles and the definition of the ‘fake bubbles’ as defined by the infinite Grassmannian equation

The parameters $\beta_{i,\lambda}$ also impact how bubble parameters change as weights change along sl_2 strings.

In [1] a cyclic variant $\mathcal{U}_Q^{cyc}(\mathfrak{g})$ of the categorified quantum group was defined utilizing more general bubble coefficients. This 2-category has \mathfrak{sl}_2 relations with all $\beta_{i,\lambda} = -1$ so that bubble coefficients $c_{i,\lambda}^+$ are constant along \mathfrak{sl}_2 strings and $c_{i,\lambda}^- = (c_{i,\lambda}^+)^{-1}$. In [1, Theorem 2.1] an explicit isomorphism of 2-categories is defined from $\mathcal{U}_Q^{cyc}(\mathfrak{g})$ to $\mathcal{U}_Q(\mathfrak{g})$. A minimal presentation of this category can be determined from [2].

Here we introduce a variant $\mathcal{U}_{Q,\beta}(\mathfrak{g}) = \mathcal{U}_{Q,\beta}^{cyc}(\mathfrak{g})$ of the categorified quantum group (Definition 2.3) that generalizes all of the variants discussed above. We then give a 2-isomorphism (Theorem 3.1) showing this general 2-category is isomorphic to the cyclic 2-category $\mathcal{U}_Q^{cyc}(\mathfrak{g})$ and hence $\mathcal{U}_Q(\mathfrak{g})$. Again, this is entirely expected by Brundan's uniqueness of categorification result. This 2-isomorphism makes it possible to directly translate results from the \mathfrak{gl}_n version of the categorified quantum group to the \mathfrak{sl}_n variant.

In Sections 3.2 and 3.3 we provide new results showing how to extend various isomorphisms of KLR-algebras to the level of 2-categories.

2 The Categorified Quantum Group $\mathcal{U}_{0,\beta}(g)$

Below we indicate in red the changes in the definition of the categorified quantum group, so that experts can easily see the differences.

2.0.1 Cartan Data

For this article we restrict our attention to simply-laced Kac-Moody algebras. These algebras are associated to a symmetric Cartan data consisting of

- a free \mathbb{Z} -module X (the weight lattice),
- for $i \in I$ (I is an indexing set) there are elements $\alpha_i \in X$ (simple roots) and $\Lambda_i \in X$ (fundamental weights),
- for $i \in I$ an element $h_i \in X^\vee = \text{Hom}_{\mathbb{Z}}(X, \mathbb{Z})$ (simple coroots),
- a bilinear form $\langle \cdot, \cdot \rangle$ on X .

Write $\langle \cdot, \cdot \rangle: X^\vee \times X \rightarrow \mathbb{Z}$ for the canonical pairing. This data should satisfy:

- $\langle \alpha_i, \alpha_i \rangle = 2$ for any $i \in I$,
- $\langle \alpha_i, \alpha_j \rangle \in \{0, -1\}$ for $i, j \in I$ with $i \neq j$,
- $\langle i, \lambda \rangle := \langle h_i, \lambda \rangle = \langle \alpha_i, \lambda \rangle$ for $i \in I$ and $\lambda \in X$,
- $\langle h_j, \Lambda_i \rangle = \delta_{ij}$ for all $i, j \in I$.

Hence $(a_{ij})_{i,j \in I}$ is a symmetrizable generalized Cartan matrix, where $a_{ij} = \langle h_i, \alpha_j \rangle = \langle \alpha_i, \alpha_j \rangle$. We will sometimes denote the bilinear pairing $\langle \alpha_i, \alpha_j \rangle$ by $i \cdot j$ and abbreviate $\langle i, \lambda \rangle$ to λ_i . We denote by $X^+ \subset X$ the dominant weights which are of the form $\sum_i \lambda_i \Lambda_i$ where $\lambda_i \geq 0$.

2.0.2 Parameters

We introduce new parameters following [12] that extend the bubble parameters introduced for the cyclic version of the quantum group in [1].

Definition 2.1 Associated to a Cartan datum we define *bubble parameters* β to be a set consisting of

- $\beta_i = \beta_{i,\lambda} \in \mathbb{k}^\times$ for $i \in I$ and $\lambda \in X$,
- $c_{i,\lambda}^+ \in \mathbb{k}^\times$ for $i \in I$ and $\lambda \in X$,
- $c_{i,\lambda}^- \in \mathbb{k}^\times$ for $i \in I$ and $\lambda \in X$.

Note that we do not require that $c_{i,\lambda}^- = (c_{i,\lambda}^+)^{-1}$ as was done in [1].

Definition 2.2 Associated to a symmetric Cartan datum define a *choice of scalars* Q consisting of:

- $\{t_{ij} \mid \text{for all } i, j \in I \text{ with } i \neq j\}$,

such that

- $t_{ij} \in \mathbb{k}^\times$,
- $t_{ij} = t_{ji}$ when $a_{ij} = 0$.

It is convenient to define $t_{ii} := -\beta_i = -\beta_{i,\lambda}$ to express the compatibility condition (2.2) in a uniform manner. A choice of bubble parameters is said to be *compatible with the scalars* Q if

$$c_{i,\lambda}^+ c_{i,\lambda}^- = -\frac{1}{\beta_i} = \frac{1}{t_{ii}}, \quad (2.1)$$

$$c_{i,\lambda \pm \alpha_j}^{\pm} = t_{ij} c_{i,\lambda}^{\pm}. \quad (2.2)$$

Such a compatible choice of scalars can be chosen for any t_{ij} by fixing an arbitrary choice of $c_{i,\lambda}^+$ for a fixed coset representative in every coset of the root lattice in the weight lattice, and then extending to the rest of the coset using the compatibility conditions.

For any choice of bubble parameters compatible with the choice of scalars Q the values along an \mathfrak{sl}_2 -string depend on $\beta_{i,\lambda}$ since

$$c_{i,\lambda + \alpha_i} = t_{ii} c_{i,\lambda} = (-\beta_{i,\lambda}) c_{i,\lambda},$$

so that for all $k \in \mathbb{Z}$ we have $c_{i,\lambda + k\alpha_i} = (-\beta_{i,\lambda})^k c_{i,\lambda}$.

2.1 The General Cyclic Form of the Categorified Quantum Group

Definition 2.3 Let β be a choice of bubble parameters that is compatible with a choice of scalars Q . The 2-category $\mathcal{U}_{Q,\beta}(\mathfrak{g}) := \mathcal{U}_{Q,\beta}^{\text{cyc}}(\mathfrak{g})$ is the graded linear 2-category consisting of:

- **Objects** λ for $\lambda \in X$.
- **1-morphisms** are formal direct sums of (shifts of) compositions of

$$\mathbb{1}_\lambda, \quad \mathbb{1}_{\lambda + \alpha_i} \mathcal{E}_i = \mathbb{1}_{\lambda + \alpha_i} \mathcal{E}_i \mathbb{1}_\lambda, \quad \text{and} \quad \mathbb{1}_{\lambda - \alpha_i} \mathcal{F}_i = \mathbb{1}_{\lambda - \alpha_i} \mathcal{F}_i \mathbb{1}_\lambda$$

for $i \in I$ and $\lambda \in X$. We denote the grading shift by $\langle 1 \rangle$, so that for each 1-morphism x in \mathcal{U} and $t \in \mathbb{Z}$ we a 1-morphism $x\langle t \rangle$.

- **2-morphisms** are \mathbb{k} -vector spaces spanned by compositions of coloured, decorated tangle-like diagrams illustrated below.

In this 2-category (and those throughout the paper) we read diagrams from right to left and bottom to top. The identity 2-morphism of the 1-morphism $\mathcal{E}_i \mathbb{1}_\lambda$ is represented by an upward oriented line labelled by i and the identity 2-morphism of $\mathcal{F}_i \mathbb{1}_\lambda$ is represented by a downward such line.

The 2-morphisms satisfy the following relations:

- (1) The 1-morphisms $\mathcal{E}_i \mathbb{1}_\lambda$ and $\mathcal{F}_i \mathbb{1}_\lambda$ are biadjoint (up to a specified degree shift). These conditions are expressed diagrammatically as

$$\begin{array}{ccc} \text{Diagram with strands } \lambda + \alpha_i \text{ and } i & = & \text{Diagram with strands } \lambda + \alpha_i \text{ and } i \\ \text{Diagram with strands } \lambda \text{ and } i & & \text{Diagram with strands } \lambda \text{ and } i \end{array} \quad (2.4)$$

(2) The 2-morphisms are cyclic with respect to this biadjoint structure.

$$\begin{array}{ccc} \text{Diagram with strands } i \text{ and } \lambda + \alpha_i & := & \text{Diagram with strands } \lambda \text{ and } i \\ \text{Diagram with strands } \lambda \text{ and } i & & \text{Diagram with strands } i \text{ and } \lambda + \alpha_i \end{array} \quad (2.5)$$

The cyclic relations for crossings are given by

$$\begin{array}{ccc} \text{Diagram with strands } i \text{ and } j & := & \text{Diagram with strands } i \text{ and } j \\ \text{Diagram with strands } i \text{ and } j & & \text{Diagram with strands } i \text{ and } j \end{array} \quad (2.6)$$

Sideways crossings are equivalently defined by the following identities:

$$\begin{array}{ccc} \text{Diagram with strands } i \text{ and } j & := & \text{Diagram with strands } i \text{ and } j \\ \text{Diagram with strands } i \text{ and } j & & \text{Diagram with strands } i \text{ and } j \end{array} \quad (2.7)$$

(3) The \mathcal{E} 's (respectively \mathcal{F} 's) carry an action of the KLR algebra for a fixed choice of parameters Q . The KLR algebra R_Q associated to a fixed set of parameters Q is defined by finite \mathbb{k} -linear combinations of braid-like diagrams in the plane, where each strand is labelled by a vertex $i \in I$. Strands can intersect and can carry dots, but triple intersections are not allowed. Diagrams are considered up to planar isotopy that do not change the combinatorial type of the diagram. We recall the local relations:

i) The quadratic KLR relations are

$$\begin{array}{ccc} \text{Diagram with strands } i \text{ and } j & = & \begin{cases} 0 & \text{if } i \cdot j = 2, \\ t_{ij} \begin{array}{c} \uparrow \\ i \end{array} \begin{array}{c} \uparrow \\ j \end{array} & \text{if } i \cdot j = 0, \\ t_{ij} \begin{array}{c} \uparrow \\ i \end{array} \begin{array}{c} \bullet \\ j \end{array} + t_{ji} \begin{array}{c} \uparrow \\ i \end{array} \begin{array}{c} \bullet \\ j \end{array} & \text{if } i \cdot j = -1, \end{cases} \\ \text{Diagram with strands } i \text{ and } j & & \text{Diagram with strands } i \text{ and } j \end{array} \quad (2.8)$$

ii) The nilHecke dot sliding relations

$$\begin{array}{ccccccc} \text{Diagram with strands } i \text{ and } j & - & \text{Diagram with strands } i \text{ and } j & = & \text{Diagram with strands } i \text{ and } j & - & \text{Diagram with strands } i \text{ and } j \\ \text{Diagram with strands } i \text{ and } j & & \text{Diagram with strands } i \text{ and } j & & \text{Diagram with strands } i \text{ and } j & & \text{Diagram with strands } i \text{ and } j \end{array} \quad (2.9)$$

iii) For $i \neq j$ the dot sliding relations

$$\begin{array}{ccc} \text{Diagram: two strands } i \text{ and } j \text{ crossing, } i \text{ over } j, \text{ with a dot on } i. & = & \text{Diagram: two strands } i \text{ and } j \text{ crossing, } i \text{ over } j, \text{ with a dot on } j. \end{array} \quad (2.10)$$

hold.

iv) Unless $i = k$ and $i \cdot j = -1$ the relation

$$\begin{array}{ccc} \text{Diagram: three strands } i, j, k \text{ crossing in a triangle, } i \text{ over } j \text{ over } k. & = & \text{Diagram: three strands } i, j, k \text{ crossing in a triangle, } i \text{ over } k \text{ over } j. \end{array} \quad (2.11)$$

holds. Otherwise, $i \cdot j = -1$ and

$$\begin{array}{ccc} \text{Diagram: three strands } i, j, i \text{ crossing in a triangle, } i \text{ over } j \text{ over } i. & - & \text{Diagram: three strands } i, j, i \text{ crossing in a triangle, } i \text{ over } i \text{ over } j. \end{array} = t_{ij} \begin{array}{c} \uparrow \\ i \\ \uparrow \\ j \\ \uparrow \\ i \end{array} \quad (2.12)$$

(4) When $i \neq j$ one has the mixed relations relating $\mathcal{E}_i \mathcal{F}_j$ and $\mathcal{F}_j \mathcal{E}_i$:

$$\begin{array}{ccc} \text{Diagram: two strands } j \text{ and } i \text{ crossing, } j \text{ over } i, \text{ with a dot on } j. & = & \text{Diagram: two strands } i \text{ and } j \text{ crossing, } i \text{ over } j, \text{ with a dot on } i. \end{array} \quad (2.13)$$

(5) Negative degree bubbles are zero. That is for all $m \in \mathbb{Z}_{>0}$ one has

$$\begin{array}{ccc} \text{Diagram: a circle with a dot at } i, \text{ degree } \lambda. & = 0 & \text{if } m < \lambda_i - 1, \\ \text{Diagram: a circle with a dot at } m, \text{ degree } \lambda. & = 0 & \text{if } m < -\lambda_i - 1 \end{array} \quad (2.14)$$

Furthermore, dotted bubbles of degree zero are scalar multiples of the identity 2-morphism determined by the bubble parameters

$$\begin{array}{ccc} \text{Diagram: a circle with a dot at } i, \text{ degree } \lambda. & = c_{i,\lambda}^+ \cdot \text{Id}_{\mathbb{1}_\lambda} & \text{for } \lambda_i \geq 1, \\ \text{Diagram: a circle with a dot at } -\lambda_i - 1, \text{ degree } \lambda. & = c_{i,\lambda}^- \cdot \text{Id}_{\mathbb{1}_\lambda} & \text{if } \lambda_i \leq -1. \end{array} \quad (2.15)$$

We introduce formal symbols called *fake bubbles*. These are positive degree endomorphisms of $\mathbb{1}_\lambda$ that carry a formal label by a negative number of dots.

- Degree zero fake bubbles are equal to

$$\begin{array}{ccc} \text{Diagram: a circle with a dot at } i, \text{ degree } \lambda. & = c_{i,\lambda}^+ \cdot \text{Id}_{\mathbb{1}_\lambda} & \text{for } \lambda_i \leq 0, \\ \text{Diagram: a circle with a dot at } -\lambda_i - 1, \text{ degree } \lambda. & = c_{i,\lambda}^- \cdot \text{Id}_{\mathbb{1}_\lambda} & \text{if } \lambda_i \geq 0 \end{array} \quad (2.16)$$

(compare with (2.15)).

- Higher degree fake bubbles for $\lambda_i < 0$ are defined inductively as

$$\lambda_i - 1 + j \circlearrowleft^{\lambda} = \begin{cases} -\frac{1}{c_{i,\lambda}} \sum_{\substack{x+y=j \\ y \geq 1}} \begin{array}{c} i \\ \lambda_i - 1 \\ +x \end{array} \circlearrowleft^{\lambda} & \text{if } 0 < j < -\lambda_i + 1; \\ 0, & \text{if } j < 0. \end{cases} \quad (2.17)$$

- Higher degree fake bubbles for $\lambda_i > 0$ are defined inductively as

$$\text{Diagram 18: A circular diagram with a point } \lambda_i - 1 + j \text{ on the left boundary. A curved arrow labeled } i \text{ enters from the left, loops around the point, and exits to the right. A curved arrow labeled } \lambda \text{ enters from the right, loops around the point, and exits to the left. The point is labeled } \lambda_i - 1 + j. \text{ The diagram is followed by an equals sign and a piecewise function.}$$

$$= \begin{cases} -\frac{1}{c_{i,\lambda}^+} \sum_{\substack{x+y=j \\ x \geq 1}} \begin{matrix} i \\ \lambda_i - 1 + j \\ +x \end{matrix} & \text{if } 0 < j < \lambda_i + 1; \\ 0, & \text{if } j < 0. \end{cases} \quad (2.18)$$

The above relations are sometimes referred to as the *infinite Grassmannian relations* which are given by comparing coefficients of t in the expression¹:

$$\begin{aligned} & \left(\begin{array}{c} i \\ \text{---} \\ \lambda \\ \text{---} \\ \lambda_{i-1} \end{array} \right) t^0 + \left(\begin{array}{c} i \\ \text{---} \\ \lambda \\ \text{---} \\ \lambda_{i-1}+1 \end{array} \right) t^1 + \dots + \left(\begin{array}{c} i \\ \text{---} \\ \lambda \\ \text{---} \\ \lambda_{i-1}+r \end{array} \right) t^r + \dots \right) \\ & \times \left(\begin{array}{c} i \\ \text{---} \\ \lambda \\ \text{---} \\ -\lambda_{i-1} \end{array} \right) t^0 + \left(\begin{array}{c} i \\ \text{---} \\ \lambda \\ \text{---} \\ -\lambda_{i-1}+1 \end{array} \right) t + \dots + \left(\begin{array}{c} i \\ \text{---} \\ \lambda \\ \text{---} \\ -\lambda_{i-1}+s \end{array} \right) t^s + \dots \right) = -\frac{1}{\beta_{i,\lambda}}. \quad (2.19) \end{aligned}$$

(6) The \mathfrak{sl}_2 relations (which we also refer to as the \mathcal{EF} and \mathcal{FE} decompositions) are:

¹Here the product formula for the infinite Grassmannian relation is equal to $-\frac{1}{\beta_{i,\lambda}}$ rather than 1 as is usually the case. This is one motivation for considering all $\beta_{i,\lambda} = -1$ as a natural choice.

It is sometimes convenient to use a shorthand notation for the bubbles that emphasizes their degrees.

$$\begin{array}{ccc} \text{Diagram 1} & := & \text{Diagram 2} \\ \text{Diagram 1} & := & \text{Diagram 2} \end{array} \quad (2.21)$$

Diagram 1: A circle with a dot at the bottom-left. The top arc is labeled i , the right arc is labeled λ , and the bottom arc is labeled $*+r$. Diagram 2: A circle with a dot at the bottom-left. The top arc is labeled i , the right arc is labeled λ , and the bottom arc is labeled $\lambda_i - 1 + r$. Diagram 1: A circle with a dot at the bottom-left. The top arc is labeled i , the right arc is labeled λ , and the bottom arc is labeled $*+r$. Diagram 2: A circle with a dot at the bottom-left. The top arc is labeled i , the right arc is labeled λ , and the bottom arc is labeled $-\lambda_i - 1 + r$.

2.2 The Usual Cyclic Form of the 2-Category

Definition 2.4 The cyclic 2-category $\mathcal{U}_Q^{cyc}(\mathfrak{g})$ defined in [1] is a specialization of $\mathcal{U}_{Q,\beta}(\mathfrak{g})$ corresponding to taking $\beta_{i,\lambda} = -1$ for all $i \in I$ and $\lambda \in X$. In that case, $c_{i,\lambda}^- = (c_{i,\lambda}^+)^{-1}$ and we will use the notation $c_{i,\lambda} := c_{i,\lambda}^+$, so that $c_{i,\lambda}^{-1} = c_{i,\lambda}^-$. Since $\beta_{i,\lambda} = -1$, it follows that the coefficients $c_{i,\lambda}$ are constant along \mathfrak{sl}_2 strings $c_{i,\lambda} = c_{i,\lambda \pm k\alpha_i}$ for all k .

Remark 2.5 The original version of the 2-category $\mathcal{U}(\mathfrak{g})$ from [5, 8] is not a specialization of $\mathcal{U}_{Q,\beta}(\mathfrak{g})$ (outside of \mathfrak{sl}_2) since these 2-categories are all cyclic. However, [1, Theorem 2.1] gives an explicit isomorphism of categories $\mathcal{M}: \mathcal{U}_Q^{cyc}(\mathfrak{g}) \rightarrow \mathcal{U}(\mathfrak{g})$ given by

$$\mathcal{M}\left(\begin{array}{c} \lambda \\ i \curvearrowright \end{array}\right) = c_{i,\lambda} \begin{array}{c} \lambda \\ i \curvearrowright \end{array} \quad \mathcal{M}\left(\begin{array}{c} i \\ \curvearrowleft \lambda \end{array}\right) = c_{i,\lambda}^{-1} \begin{array}{c} i \\ \curvearrowleft \lambda \end{array} \quad (2.22)$$

so that a sideways crossing is rescaled by

$$\mathcal{M}\left(\begin{array}{c} \lambda \\ j \times \curvearrowleft_i \end{array}\right) = t_{ij}^{-1} \begin{array}{c} \lambda \\ j \times \curvearrowleft_i \end{array} \quad (2.23)$$

2.3 The $\mathfrak{gl}(n)$ -Version of the 2-Category

The weight lattice of $U_q(\mathfrak{gl}_n)$ is given by tuples $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}^n$. Such a weight determines an \mathfrak{sl}_n weight $\bar{\lambda} = (\bar{\lambda}_1, \dots, \bar{\lambda}_{n-1}) \in \mathbb{Z}^{n-1}$ such that $\bar{\lambda}_k := \lambda_k - \lambda_{k+1}$. Let $\varepsilon_i = (0, \dots, 0, 1, 0, \dots, 0) \in \mathbb{Z}^n$ with a single 1 in the i th position. Let $\alpha_i = \varepsilon_i - \varepsilon_{i+1} \in \mathbb{Z}^n$ for $1 \leq i \leq n-1$.

Here we will be interested in starting from a \mathfrak{sl}_n weight $\mu = (\mu_1, \dots, \mu_{n-1})$ and producing a \mathfrak{gl}_n weight, though such an assignment is not unique. For $d \in \mathbb{Z}$ define a map

$$\phi_{n,d}: \mathbb{Z}^{n-1} \longrightarrow \mathbb{Z}^n \cup \{*\} \quad (2.24)$$

by sending $\phi_{n,d}(\mu) = \lambda$, where λ uniquely satisfies

$$\lambda_i - \lambda_{i+1} = \mu_i \quad (2.25)$$

$$\sum_{i=1}^n \lambda_i = d \quad (2.26)$$

if such a solution exists and $\phi_{n,d}(\mu) = *$ otherwise. Write $(\mathbb{Z}^n)_d$ for the set of weights $\lambda \in \mathbb{Z}^n$ such that $\sum_i \lambda_i = d$. Let $\dot{U}_q(\mathfrak{gl}_n)$ denote Lusztig's idempotent form of the quantum group $U_q(\mathfrak{gl}_n)$ and set $\dot{U}_q(\mathfrak{gl}_n)^d$ to be the subalgebra of $\dot{U}_q(\mathfrak{gl}_n)$ spanned by weights $\lambda \in (\mathbb{Z}^n)_d$.

For a given μ , its image $\phi_{n,d}(\mu)$ has a solution if and only if

$$d \equiv \sum_{i=1}^{n-1} i \mu_i \pmod{n}.$$

In particular, for a fixed $d \in \mathbb{Z}$, then exactly one of the maps $\phi_{n,d+k}$ will have a solution for $k = 0, 1, \dots, n-1$. Hence, for all $d \in \mathbb{Z}$ there is an isomorphism²

$$\dot{U}_q(\mathfrak{sl}_n) \cong \bigoplus_{k=d}^{d+n-1} \dot{U}_q(\mathfrak{gl}_n)^k \quad (2.27)$$

determined on weights by sending μ to $\phi_{n,d+k}(\mu)$ where $d+k \equiv \sum_{i=1}^{n-1} i \mu_i \pmod{n}$, see for example [14] or [6, Remark 2.2] and the references therein.

Definition 2.6 Fix a compatible choice of scalars and bubble coefficients Q and β . The 2-category $\mathcal{U}_{Q,\beta}(\mathfrak{gl}_n)$ is the graded additive 2-category with

- **Objects** objects $\lambda \in \mathbb{Z}^n$,
- **1-morphisms** are formal direct sums of (shifts of) compositions of

$$\mathbb{1}_\lambda, \quad \mathbb{1}_{\lambda+\alpha_i} \mathcal{E}_i = \mathbb{1}_{\lambda+\alpha_i} \mathcal{E}_i \mathbb{1}_\lambda, \quad \text{and} \quad \mathbb{1}_{\lambda-\alpha_i} \mathcal{F}_i = \mathbb{1}_{\lambda-\alpha_i} \mathcal{F}_i \mathbb{1}_\lambda$$

for $1 \leq i \leq n-1$ and $\lambda \in \mathbb{Z}^n$. We denote the grading shift by $\langle 1 \rangle$, so that for each 1-morphism x in \mathcal{U} and $t \in \mathbb{Z}$ we a 1-morphism $x \langle t \rangle$.

- **2-morphisms** the same as those in the 2-category $\mathcal{U}_{Q,\beta}(\mathfrak{sl}_n)$

and relations identical to those in $\mathcal{U}_{Q,\beta}(\mathfrak{sl}_n)$ with each $\lambda_i := \langle i, \lambda \rangle$ for $\lambda \in \mathbb{Z}^{n-1}$ replaced by $\bar{\lambda}_i$.

Then the isomorphism (2.27) motivates the following 2-functors defined for each $d \in \mathbb{Z}$

$$\begin{aligned} \phi_{n,d}: \mathcal{U}_{Q,\beta}(\mathfrak{sl}_n) &\longrightarrow \mathcal{U}_{Q,\beta}(\mathfrak{gl}_n) \\ \mu &\mapsto \phi_{n,d}(\mu) \\ \mathcal{E}_i \mathbb{1}_\mu &\mapsto \mathcal{E}_i \mathbb{1}_{\phi_{n,d}(\mu)} \end{aligned} \quad (2.28)$$

which send the generating 1-morphisms and 2-morphisms of $\mathcal{U}_{Q,\beta}(\mathfrak{sl}_n)$ to the corresponding ones in $\mathcal{U}_{Q,\beta}(\mathfrak{gl}_n)$. Using \mathfrak{gl}_n weights it is possible to specify an explicit choice of compatible scalars and bubble coefficients.

Definition 2.7 The 2-category $\mathcal{U}(\mathfrak{gl}_n)$ defined in [14] corresponds to the choice of scalars

$$t_{ij} = \begin{cases} -1 & \text{if } i = j \\ -1 & \text{if } i \cdot j = -1 \text{ and } i \longrightarrow j \\ 1 & \text{otherwise,} \end{cases}$$

(so that $\beta_{i,\lambda} = -t_{ii} = 1$) and bubble parameters

$$c_{i,\lambda}^+ = (-1)^{\lambda_{i+1}} \quad \text{for } \bar{\lambda}_i \geq 0, \quad c_{i,\lambda}^- = (-1)^{\lambda_{i+1}-1} \quad \text{for } \bar{\lambda}_i \leq 0.$$

Hence, the [14] 2-category $\mathcal{U}(\mathfrak{gl}_n)$ is a specialization of $\mathcal{U}_{Q,\beta}(\mathfrak{gl}_n)$.

3 Isomorphism of 2-Categories

To aid the reader, our convention throughout this section is to indicate diagrams in the image of a 2-isomorphism in blue.

²Thanks to Hoel Queffelec for pointing out this fact.

3.1 Cyclic to General 2-Isomorphisms

Recall that the cyclic 2-category $\mathcal{U}_Q^{cyc}(\mathfrak{g})$ is a specialization of $\mathcal{U}_{Q,\beta}(\mathfrak{g})$ corresponding to taking $\beta_{i,\lambda} = -1$ for all $i \in I$ and $\lambda \in X$. We denote the bubble parameters in $\mathcal{U}_Q^{cyc}(\mathfrak{g})$ using the notation $c_{i,\lambda} := \mathbf{c}_{i,\lambda}^+$ and $c_{i,\lambda}^{-1} = \mathbf{c}_{i,\lambda}^-$, since for this choice of $\beta_{i,\lambda}$ we have $\mathbf{c}_{i,\lambda}^- = (\mathbf{c}_{i,\lambda}^+)^{-1}$.

Theorem 3.1 *There is a 2-isomorphism of graded additive \mathbb{k} -linear categories*

$$\mathcal{U}_Q^{cyc}(\mathfrak{g}) \longrightarrow \mathcal{U}_{Q,\beta}(\mathfrak{g}) \quad (3.1)$$

which acts as the identity on objects and 1-morphisms and rescales generating 2-morphisms as follows

$$F \left(\begin{array}{c} \uparrow \\ | \\ i \end{array} \right) = \begin{array}{c} \uparrow \\ | \\ i \end{array} \quad F \left(\begin{array}{c} \uparrow \uparrow \\ i \times j \end{array} \right) = \begin{array}{c} \uparrow \uparrow \\ i \times j \end{array}$$

$$F \left(\begin{array}{c} \lambda \\ i \curvearrowright \end{array} \right) = \begin{cases} \frac{c_{i,\lambda}}{c_{i,\lambda}^+} i \xrightarrow{\lambda} & \text{if } \lambda_i \equiv_4 0 \text{ or } \lambda_i \equiv_4 1 \\ c_{i,\lambda} i \xrightarrow{\lambda} & \text{otherwise.} \end{cases} \quad (3.2)$$

$$F \left(\begin{array}{c} \lambda \\ \curvearrowright i \end{array} \right) = \begin{cases} \frac{1}{c_{i,\lambda}^-} \xleftarrow{\lambda} i & \text{if } \lambda_i \equiv_4 0 \text{ or } \lambda_i \equiv_4 1 \\ \xleftarrow{\lambda} i & \text{otherwise.} \end{cases} \quad (3.3)$$

$$F \left(\begin{array}{c} \uparrow \\ \lambda \end{array} \right) = \begin{cases} \frac{1}{c_{i,\lambda}^+} \xleftarrow{\lambda} i & \text{if } \lambda_i \equiv_4 2 \text{ or } \lambda_i \equiv_4 3 \\ \xleftarrow{\lambda} i & \text{otherwise} \end{cases} \quad (3.4)$$

$$F \left(\begin{array}{c} i \\ \lambda \end{array} \right) = \begin{cases} \frac{c_{i,\lambda}^{-1}}{c_{i,\lambda}^+} i \xleftarrow{\lambda} & \text{if } \lambda_i \equiv_4 2 \text{ or } \lambda_i \equiv_4 3 \\ c_{i,\lambda}^{-1} i \xleftarrow{\lambda} & \text{otherwise} \end{cases}$$

Proof This assignment preserves the scalars Q , however by definition we have $t_{ii}^{cyc} = -\beta_i^{cyc} = +1$ in $\mathcal{U}_Q^{cyc}(\mathfrak{g})$ and $t_{ii} = -\beta_i$ in $\mathcal{U}_{Q,\beta}(\mathfrak{g})$. Since F does not effect upward oriented dots and crossings this rescaling has no effect on the KLR-relations.

To see that these definitions preserve the adjunction axioms (2.3) and (2.4) for caps and cups it is helpful to note that by the properties of compatible bubble coefficients we have

$$\mathbf{c}_{i,\lambda-\alpha_i}^- \stackrel{(2.2)}{=} t_{ii} \mathbf{c}_{i,\lambda}^- = -\beta_{i,\lambda} \mathbf{c}_{i,\lambda}^- \stackrel{(2.1)}{=} -\beta_{i,\lambda} \left(\frac{1}{-\beta_{i,\lambda} \mathbf{c}_{i,\lambda}^+} \right) = \frac{1}{\mathbf{c}_{i,\lambda}^+}. \quad (3.5)$$

The preservation of the dot cyclicity relation (2.5) follows.

A careful case by case analysis shows that crossing cyclicity (2.6) is preserved. This verification produces rescalings that depend of the values of λ_i and λ_j . For example, if $\lambda_i \equiv 4 0$ and $\lambda_j \equiv 4 0$ there is no rescaling factor on either side of Eq. 2.6. However, for $\lambda_i \equiv 4 2$ and $\lambda_j \equiv 4 1$ both sides are rescaled by a factor of $t_{ji} t_{ij}^{-1} / c_{j,\lambda}^+$.

To verify the remaining relations it is helpful to compute the image of some important composite morphisms. By definition, the images of real bubbles are given by

$$F \left(\begin{array}{c} i \\ \text{---} \\ \bullet \\ \text{---} \\ *+r \end{array} \lambda \right) = \frac{c_{i,\lambda}}{c_{i,\lambda}^+} \begin{array}{c} i \\ \text{---} \\ \bullet \\ \text{---} \\ *+r \end{array} \lambda \quad F \left(\begin{array}{c} i \\ \text{---} \\ \bullet \\ \text{---} \\ *+r \end{array} \lambda \right) = \frac{c_{i,\lambda}^{-1}}{c_{i,\lambda}^+} \begin{array}{c} i \\ \text{---} \\ \bullet \\ \text{---} \\ *+r \end{array} \lambda \quad (3.6)$$

so it is clear that the degree zero bubble relations are preserved (a clockwise bubble is equal to $c_{i,\lambda}$ in $\mathcal{U}_Q^{cyc}(\mathfrak{g})$ and $c_{i,\lambda}^+$ in $\mathcal{U}_{Q,\beta}(\mathfrak{g})$). For fake bubbles it follows by induction using Eqs. 2.17 and 2.18 that

$$F \left(\begin{array}{c} i \\ \text{---} \\ \bullet \\ \text{---} \\ *+r \end{array} \lambda \right) = -\beta_{i,\lambda} \frac{c_{i,\lambda}^-}{c_{i,\lambda}^{-1}} \begin{array}{c} i \\ \text{---} \\ \bullet \\ \text{---} \\ *+r \end{array} \lambda \quad F \left(\begin{array}{c} i \\ \text{---} \\ \bullet \\ \text{---} \\ *+r \end{array} \lambda \right) = -\beta_{i,\lambda} \frac{c_{i,\lambda}^+}{c_{i,\lambda}} \begin{array}{c} i \\ \text{---} \\ \bullet \\ \text{---} \\ *+r \end{array} \lambda \quad (3.7)$$

Hence, the right most terms in the EF -relations (2.20) rescale as the product of the rescalings for the real bubble together with the oppositely oriented fake bubble. In both cases this product is $-\beta_{i,\lambda}$.

One can also show that

$$F \left(\begin{array}{c} i \\ \text{---} \\ \text{---} \\ \text{---} \\ j \end{array} \lambda \right) = F \left(\begin{array}{c} i \\ \text{---} \\ \text{---} \\ \text{---} \\ j \end{array} \lambda \right) = c_{i,\lambda} \cdot c_{i,\lambda+\alpha_j-\alpha_i}^{-1} \frac{1}{c_{i,\lambda+\alpha_j-\alpha_i}^-} \frac{1}{c_{i,\lambda}^+} \begin{array}{c} i \\ \text{---} \\ \text{---} \\ \text{---} \\ j \end{array} \lambda \quad (3.8)$$

$$F \left(\begin{array}{c} i \\ \text{---} \\ \text{---} \\ \text{---} \\ j \end{array} \lambda \right) = F \left(\begin{array}{c} i \\ \text{---} \\ \text{---} \\ \text{---} \\ j \end{array} \lambda \right) = c_{i,\lambda} \cdot c_{i,\lambda+\alpha_j-\alpha_i}^{-1} \frac{1}{c_{i,\lambda+\alpha_j-\alpha_i}^-} \frac{1}{c_{i,\lambda}^+} \begin{array}{c} i \\ \text{---} \\ \text{---} \\ \text{---} \\ j \end{array} \lambda \quad (3.9)$$

where the contribution of the bubble parameters from $\mathcal{U}_{Q,\beta}(\mathfrak{g})$ is $\frac{1}{c_{i,\lambda+\alpha_j-\alpha_i}^-} \frac{1}{c_{i,\lambda}^+} \stackrel{(3.5)}{=} \frac{c_{i,\lambda+\alpha_j}^+}{c_{i,\lambda}^+} = t_{ij}$ while the contribution from $\mathcal{U}_Q^{cyc}(\mathfrak{g})$ is

$$c_{i,\lambda} \cdot c_{i,\lambda+\alpha_j-\alpha_i}^{-1} = (t_{ij}^{cyc})^{-1}.$$

In particular, when $i \neq j$ the contributions cancel to give a rescaling coefficient of 1 and when $i = j$ only the $t_{ii} = -\beta_{i,\lambda}$ contributes. Hence, the map F preserves the mixed relation (2.13) and sends the usual \mathfrak{sl}_2 relation to the $\beta_{i,\lambda}$ modified form in Eq. 2.20. \square

3.2 Rescaling NilHecke Subalgebras

The nilHecke algebra remains invariant if we rescale a dot by an arbitrary invertible parameter along with the crossing by the inverse of this parameter. Here we record how the 2-category $\mathcal{U}_{Q,\beta}(\mathfrak{g})$ is effected by this transformation.

Proposition 3.2 *Given bubble parameters β with compatible choice of scalars Q , and $D_i \in \mathbb{k}^\times$ for $i \in I$, define bubble coefficients $\widehat{\beta}$ with*

$$\widehat{c}_{i,\lambda}^+ := D_i^{-\lambda_i+1} c_{i,\lambda}^+, \quad \widehat{c}_{i,\lambda}^- := D_i^{\lambda_i+1} c_{i,\lambda}^-, \quad \widehat{\beta}_{i,\lambda} := D_i^{-2} \beta_{i,\lambda}. \quad (3.10)$$

and scalars \widehat{Q} with

$$\begin{aligned} \widehat{t}_{ii} &= -\widehat{\beta}_{i,\lambda} = -D_i^{-2} \beta_{i,\lambda} = D_i^{-2} t_{ii} \\ \widehat{t}_{ij} &= D_i t_{ij} \quad \text{if } (\alpha_i, \alpha_j) = -1, \\ \widehat{t}_{ik} &= t_{ik} \quad \text{if } (\alpha_i, \alpha_k) = 0. \end{aligned} \quad (3.11)$$

Then \widehat{Q} and $\widehat{\beta}$ define bubble parameters with a compatible choice of scalars.

Proof One can check that with these assignments we have

$$\widehat{c}_{i,\lambda}^+ \widehat{c}_{i,\lambda}^- = -\frac{1}{\widehat{\beta}_i} \quad (3.12)$$

so that Eq. 2.1 is satisfied with the new bubble coefficients. We also have

$$\begin{aligned} \widehat{c}_{i,\lambda+\alpha_i}^+ &= D_i^{-\lambda_i-1} c_{i,\lambda+\alpha_i}^+ = t_{ii} D_i^{-\lambda_i-1} c_{i,\lambda}^+ = t_{ii} D_i^{-2} \widehat{c}_{i,\lambda}^+ = \widehat{t}_{ii} \widehat{c}_{i,\lambda}^+ \\ \widehat{c}_{i,\lambda+\alpha_j}^+ &= D_i^{-\lambda_i+2} c_{i,\lambda+\alpha_j}^+ = D_i^{-\lambda_i+2} t_{ij} c_{i,\lambda}^+ = D_i t_{ij} \widehat{c}_{i,\lambda}^+ \quad \text{if } (\alpha_i, \alpha_j) = -1 \\ \widehat{c}_{i,\lambda+\alpha_k}^+ &= D_i^{-\lambda_i+1} c_{i,\lambda+\alpha_k}^+ = D_i^{-\lambda_i+1} t_{ik} c_{i,\lambda}^+ = t_{ik} \widehat{c}_{i,\lambda}^+ \quad \text{if } (\alpha_i, \alpha_k) = 0, \end{aligned}$$

so that Eq. 2.2 is also preserved. Equation 2.2 for c^- follows similarly. \square

Proposition 3.3 *For each $i \in I$ let $D_i \in \mathbb{k}$. Then there is an isomorphism of 2-categories*

$$\beth: \mathcal{U}_{Q,\beta}(\mathfrak{g}) \longrightarrow \mathcal{U}_{\widehat{Q},\widehat{\beta}}(\mathfrak{g}) \quad (3.13)$$

defined by

$$\beth \left(\begin{array}{c} \uparrow \\ \bullet \\ i \end{array} \middle| \lambda \right) = D_i \begin{array}{c} \uparrow \\ \bullet \\ i \end{array} \quad \beth \left(\begin{array}{c} \nearrow \nwarrow \\ i \end{array} \middle| \lambda \right) = D_i^{-1} \begin{array}{c} \nearrow \nwarrow \\ i \end{array} \quad (3.14)$$

where the coefficients \widehat{Q} and $\widehat{\beta}$ are defined in Eq. 3.2.

Proof The rescaling of the i -labelled dot and the ii -crossing are necessarily inverse because of the nilHecke dot sliding relation (2.10). Since caps and cups are untouched, it is immediate that adjunction axioms and cyclicity are preserved. For this rescaling of the ii -crossing it is clear by Eq. 2.8 that the coefficients t_{ij} must map to $D_i^{-1} \widehat{t}_{ij}$ and the other KLR relations follow similarly. Likewise, the definition of bubble parameters (2.15) determines $\widehat{c}_{i,\lambda}^+$ and

$\widehat{c}_{i,\lambda}$. The only relation that requires care is the \mathfrak{sl}_2 -relations (2.20). For example, the first \mathfrak{sl}_2 relation follows since for $\lambda_i \geq 0$ the degree $2j$ fake bubble is given by

$$\square \left(\begin{array}{c} i \\ -\lambda_i - 1 + j \end{array} \right) := D_i^{-\lambda_i - 1 + j} \left(\begin{array}{c} i \\ -\lambda_i - 1 + j \end{array} \right) \quad (3.15)$$

which follows by induction since

$$\begin{aligned} \square \left(\begin{array}{c} i \\ -\lambda_i - 1 + j \end{array} \right) &:= -\frac{1}{\widehat{c}_{i,\lambda}^+} \square \left(\sum_{\substack{x+y=j \\ x \geq 1}} \begin{array}{c} i \\ \lambda \\ \lambda_i - 1 + x \end{array} \right) \left(\begin{array}{c} i \\ -\lambda_i - 1 + y \end{array} \right) \\ &= -\frac{D_i^{-\lambda_i + 1}}{\widehat{c}_{i,\lambda}^+} \square \left(\sum_{\substack{x+y=j \\ x \geq 1}} \begin{array}{c} i \\ \lambda \\ \lambda_i - 1 + x \end{array} \right) \left(\begin{array}{c} i \\ -\lambda_i - 1 + y \end{array} \right) \\ &= -\frac{D_i^{-\lambda_i + 1}}{\widehat{c}_{i,\lambda}^+} D_i^{\lambda_i - 1 + x} D_i^{-\lambda_i - 1 + y} \sum_{\substack{x+y=j \\ x \geq 1}} \begin{array}{c} i \\ \lambda \\ \lambda_i - 1 + x \end{array} \left(\begin{array}{c} i \\ -\lambda_i - 1 + y \end{array} \right) \\ &= -\frac{D_i^{-\lambda_i + 1}}{\widehat{c}_{i,\lambda}^+} \sum_{\substack{x+y=j \\ x \geq 1}} \begin{array}{c} i \\ \lambda \\ \lambda_i - 1 + x \end{array} \left(\begin{array}{c} i \\ -\lambda_i - 1 + y \end{array} \right) = D_i^{-\lambda_i - 1 + j} \left(\begin{array}{c} i \\ \lambda \\ \lambda_i - 1 + j \end{array} \right) \end{aligned}$$

Hence, the \mathfrak{sl}_2 relation follows since

$$\begin{aligned} \square \left(\beta_{i,\lambda} \sum_{\substack{f_1 + f_2 + f_3 \\ = \lambda_i - 1}} \begin{array}{c} i \\ + f_2 \\ - f_1 \end{array} \right) &= \beta_{i,\lambda} \sum_{\substack{f_1 + f_2 + f_3 \\ = \lambda_i - 1}} D_i^{f_1 + f_3} D_i^{-\lambda_i - 1 + f_2} \begin{array}{c} i \\ + f_2 \\ - f_1 \end{array} \\ &= \widehat{\beta}_{i,\lambda} \sum_{\substack{f_1 + f_2 + f_3 \\ = \lambda_i - 1}} \begin{array}{c} i \\ + f_2 \\ - f_1 \end{array} \end{aligned}$$

and

$$\square \left(\beta_{i,\lambda} \begin{array}{c} i \\ \diagup \quad \diagdown \\ i \end{array} \right) = D_i^{-2} \beta_{i,\lambda} \begin{array}{c} i \\ \diagup \quad \diagdown \\ i \end{array} = \widehat{\beta}_{i,\lambda} \begin{array}{c} i \\ \diagup \quad \diagdown \\ i \end{array}$$

The other \mathfrak{sl}_2 relation follows similarly. \square

3.3 Rescalings for a Preferred Choice of Scalars Q

The choice of scalars Q controls the form of the KLR algebra R_Q that governs the upward oriented strands. The algebras R_Q are governed by the products $v_{ij} = t_{ij}^{-1} t_{ji}$ taken over all pairs $i, j \in I$. The products v_{ij} can be thought of as a \mathbb{k}^\times -valued 1-cocycle on the graph Γ associated to the symmetric Cartan data; we call two choices *cohomologous* if these 1-cocycles are in the same cohomology class. For cohomologous choices of scalars Q and Q' the algebras R_Q and $R'_{Q'}$ are isomorphic, though not canonically. If Γ is a tree, in particular, a Dynkin diagram, then all choices of scalars are cohomologous. However, an explicit isomorphism of algebras has not been presented in the literature. Here we give this isomorphism $R_Q \rightarrow R_{Q'}$ for any choices of scalars Q and Q' and discuss how to extend to the level of 2-categories $\mathcal{U}_{Q,\beta}(\mathfrak{g})$ and $\mathcal{U}_{Q',\beta}(\mathfrak{g})$. In [7] Kashiwara studies the effect of these parameters on simple modules for the KLR algebra and show that these generically correspond to upper global canonical bases.

3.3.1 KLR Isomorphisms for Trees

Let Γ be the underlying graph associated to a simply-laced Kac-Moody algebra. If Γ is a tree, then any two vertices $i, j \in I$ of the graph are connected by a unique path. To specify the isomorphism we designate a distinguished vertex $r \in I$ as the root of the tree. This defines an orientation on the graph in which all edges are oriented away from r . Define the *level* of a vertex $i \in I$ to be its distance from the root r . We write $i < j$ if vertex i occurs in a lower level than vertex j .

Denote by P_i the unique directed path of edges from the root vertex r to vertex i . For any edge $e \in \Gamma$ we write $e \in P_i$ if e appears in the path from r to i . Let $s(e) \in I$ denote the source vertex of the edge e and $t(e) \in I$ denote the target of the directed edge e .

Define parameters

$$D_i := \prod_{e \in P_i} v'_{s(e)t(e)} v_{t(e)s(e)}, \quad D_i^{-1} := \prod_{e \in P_i} v'_{t(e)s(e)} v_{s(e)t(e)} \quad (3.16)$$

so that D_i^{-1} is the inverse of D_i , since $v_{ij} = t_{ij}^{-1} t_{ji}$ so that $v_{ji} = v_{ij}^{-1}$. For the root of the tree $r \in I$ we have $D_r = 1$.

Proposition 3.4 *Let Q and Q' be choices of scalars for a KLR algebra associated to a simply-laced Kac-Moody algebra whose underlying graph is a tree. Then associated to a fixed choice of root for the tree, there is an algebra isomorphism*

$$\natural: R_Q \longrightarrow R_{Q'} \quad (3.17)$$

defined by

$$\natural \left(\begin{array}{c} \uparrow \\ \bullet \quad \lambda \\ i \end{array} \right) = D_i \begin{array}{c} \uparrow \\ \bullet \quad \lambda \\ i \end{array} \quad (3.18)$$

$$\mathfrak{J} \left(\begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} \right) = \begin{cases} D_i^{-1} \begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} & \text{if } i = j, \\ t_{ji}(t'_{ji})^{-1} D_j \begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} & \text{if } i < j \text{ and } i \cdot j = -1, \\ t_{ij}(t'_{ij})^{-1} \begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} & \text{if } i < j \text{ and } i \cdot j = 0, \\ \begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} & \text{otherwise,} \end{cases}$$

with parameters D_i for $i \in I$ defined in Eq. 3.16.

Proof It is immediate that the nilHecke relations are preserved because the i colored dot is rescaled inversely to the ii colored crossing. If $i < j$ and $i \cdot j = -1$, then the path P_i is identical to the path P_j with the addition of the edge from $i \rightarrow j$. Hence,

$$D_j = \prod_{e \in P_j} v'_{s(e)t(e)} v_{t(e)s(e)} = v'_{ij} v_{ji} \prod_{e \in P_i} v'_{s(e)t(e)} v_{t(e)s(e)} = v'_{ij} v_{ji} D_i \quad (3.19)$$

and in particular,

$$t_{ji}(t'_{ji})^{-1} D_j = t_{ji}(t'_{ji})^{-1} v'_{ij} v_{ji} D_i = t_{ij}(t'_{ij})^{-1} D_i. \quad (3.20)$$

Hence, if $i < j$ and $i \cdot j = -1$ then

$$\mathfrak{J} \left(\begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} \right) = t_{ji}(t'_{ji})^{-1} D_j \begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} = t_{ij}(t'_{ij})^{-1} D_i \begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} \quad (3.21)$$

which, together with the dot rescaling, transforms the quadratic KLR relation in parameters Q to the corresponding relation with parameters Q' . A similar argument establishes the quadratic KLR relation for $j < i$ and $i \cdot j = -1$. The quadratic relation for $i \cdot j = 0$ is immediate.

All of the cubic KLR relations follow immediately except for the case (2.12). If $i < j$ and $i \cdot j = -1$ then

$$\mathfrak{J} \left(\begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} - \begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} \right) \stackrel{(3.20)}{=} t_{ij}(t'_{ij})^{-1} \left(\begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} - \begin{array}{c} \text{Diagram: two strands } i \text{ and } j \text{ crossing, with a dot at the crossing labeled } \lambda. \end{array} \right)$$

so that the nontrivial cubic KLR relation also follows. \square

3.3.2 KLR rescaling for 2-categories

The isomorphism $\mathfrak{J}: R_Q \rightarrow R_{Q'}$ of KLR algebras can be extended to an isomorphism of 2-categories. Without rescaling the caps and cups, the assignment (3.18) will rescale the bubble parameters as in Eq. 3.10, but these new parameters need not be compatible with the coefficients Q' . If we assume that a set of bubble parameters β' compatible with Q' have been specified, then it is possible to rescale appropriately to extend \mathfrak{J} to an isomorphism of 2-categories.

Theorem 3.5 Let β , respectively β' , be bubble parameters with compatible choice of scalars Q , respectively Q' . Then the algebra isomorphism $\natural: R_Q \rightarrow R_{Q'}$ from Eq. 3.18 extends to an isomorphism of 2-categories

$$\natural: \mathcal{U}_{Q,\beta} \longrightarrow \mathcal{U}_{Q',\beta'} \quad (3.22)$$

given by

$$\natural \left(\begin{array}{c} \lambda \\ i \curvearrowleft \curvearrowright \\ i \end{array} \right) = D_i^{-\lambda_i+1} \frac{c_{i,\lambda}^+}{c_{i,\lambda}^+} \begin{array}{c} \lambda \\ i \curvearrowleft \curvearrowright \\ i \end{array} \quad \natural \left(\begin{array}{c} i \\ \cup \\ \lambda \end{array} \right) = D_i^{\lambda_i+1} \frac{c_{i,\lambda+\alpha_i}^{'+}}{c_{i,\lambda+\alpha_i}^+} \begin{array}{c} i \\ \cup \\ \lambda \end{array} \quad (3.23)$$

and sending the other cap and cup to themselves.

Proof Note that the coefficient for the rightward pointing cup can be rewritten

$$D_i^{\lambda_i+1} \frac{c_{i,\lambda+\alpha_i}^{'+}}{c_{i,\lambda+\alpha_i}^+} \stackrel{(2.2)}{=} D_i^{\lambda_i+1} \frac{t'_{ii}}{t_{ii}} \frac{c_{i,\lambda}^+}{c_{i,\lambda}^+} \stackrel{(2.1)}{=} D_i^{\lambda_i+1} \frac{c_{i,\lambda}^-}{c_{i,\lambda}^-}.$$

Using this it is not difficult to show that the adjunction axioms and dot cyclicity are preserved. It is also straightforward to show that bubble coefficients are preserved. For example,

$$c_{i,\lambda}^+ = \natural \left(\begin{array}{c} i \\ \circlearrowleft \\ \lambda \\ \lambda_{i-1} \end{array} \right) = D_i^{\lambda_i-1} \cdot D_i^{-\lambda_i+1} \frac{c_{i,\lambda}^+}{c_{i,\lambda}^+} \begin{array}{c} i \\ \circlearrowleft \\ \lambda \\ \lambda_{i-1} \end{array} \quad (3.24)$$

For crossing cyclicity let γ be the rescaling of the ij crossing (for arbitrary relationship between i and j) so that

$$\begin{aligned} \natural \left(\begin{array}{c} j \\ \cup \\ i \\ \lambda \\ i \downarrow \\ j \downarrow \end{array} \right) &= \gamma D_j^{-\lambda_j+1} \frac{c_{j,\lambda}^+}{c_{j,\lambda}^+} \cdot D_i^{-\lambda_i} \frac{c_{i,\lambda-\alpha_j}^+}{c_{i,\lambda-\alpha_j}^+} \cdot D_i^{\lambda_i-1} \frac{c_{i,\lambda}^+}{c_{i,\lambda}^+} \\ &\cdot D_j^{\lambda_j} \frac{c_{j,\lambda-\alpha_i}^+}{c_{j,\lambda-\alpha_i}^+} \begin{array}{c} j \\ \cup \\ i \\ \lambda \\ i \downarrow \\ j \downarrow \end{array} \\ &= \gamma D_j D_i^{-1} \frac{c_{j,\lambda-\alpha_i}^+}{c_{j,\lambda}^+} \frac{c_{i,\lambda}^+}{c_{i,\lambda-\alpha_j}^+} \frac{c_{i,\lambda-\alpha_j}^+}{c_{i,\lambda}^+} \frac{c_{j,\lambda}^+}{c_{j,\lambda-\alpha_i}^+} \begin{array}{c} j \\ \cup \\ i \\ \lambda \\ i \downarrow \\ j \downarrow \end{array} \end{aligned}$$

$$\begin{aligned}
 &= \gamma D_j D_i^{-1} t_{ji}^{-1} t'_{ij} t_{ij}^{-1} t_{ji} \\
 &\quad \text{Diagram: } \begin{array}{c} j \quad i \\ \text{---} \\ \text{---} \end{array} \text{ with a blue } \lambda \text{-labeled crossing } i \downarrow j \\
 &= \gamma \quad \text{Diagram: } \begin{array}{c} j \quad i \\ \text{---} \\ \text{---} \end{array} \text{ with a blue } \lambda \text{-labeled crossing } i \downarrow j \quad (3.25) \\
 &= \gamma \quad \text{Diagram: } \begin{array}{c} j \quad i \\ \text{---} \\ \text{---} \end{array} \text{ with a blue } \lambda \text{-labeled crossing } i \downarrow j \\
 &= \mathbb{J} \left(\begin{array}{c} j \quad i \\ \text{---} \\ \text{---} \end{array} \text{ with a blue } \lambda \text{-labeled crossing } i \downarrow j \right)
 \end{aligned}$$

Where in Eq. 3.25 we used properties of the coefficients Q and Q' from Definition 2.2 and Eq. 3.19 when $(\alpha_i, \alpha_j) = -1$.

Similarly, if γ represents the rescaling of an ij -labelled crossing then

$$\begin{aligned}
 \mathbb{J} \left(\begin{array}{c} i \\ \text{---} \\ i \downarrow j \end{array} \text{ with a blue } \lambda \text{-labeled crossing } i \downarrow j \right) &= \mathbb{J} \left(\begin{array}{c} i \uparrow j \\ \text{---} \\ i \downarrow j \end{array} \text{ with a blue } \lambda \text{-labeled crossing } i \downarrow j \right) = \gamma \cdot D_i^{-\lambda_i+1} \frac{c_{i,\lambda}^+}{c_{i,\lambda}^{i+}} \cdot D_i^{\lambda_i-1+(\alpha_i, \alpha_j)} \frac{c_{i,\lambda+\alpha_j}^{i+}}{c_{i,\lambda+\alpha_j}^+} \\
 &= \gamma \cdot D_i^{(\alpha_i, \alpha_j)} \frac{c_{i,\lambda+\alpha_j}^{i+}}{c_{i,\lambda}^{i+}} \cdot \frac{c_{i,\lambda}^+}{c_{i,\lambda+\alpha_j}^+} \stackrel{(2.2)}{=} \gamma \cdot D_i^{(\alpha_i, \alpha_j)} t'_{ij} t_{ij}^{-1} \quad (3.26)
 \end{aligned}$$

Using the specific value of γ from Eq. 3.18 that depends on the relationship of i and j in the ij -crossing, it follows that

$$\gamma \cdot D_i^{(\alpha_i, \alpha_j)} t'_{ij} t_{ij}^{-1} = \begin{cases} t'_{ii} / t_{ii} = \beta'_i / \beta_i & \text{if } i = j, \\ 1 & \text{otherwise,} \end{cases} \quad (3.27)$$

where we have made use of Eq. 3.20 when $i < j$ and $i \cdot j = -1$. This, and a similar computation with the other orientation, shows that the mixed relations are preserved.

For the \mathfrak{sl}_2 relation we note that bubbles (real or fake) rescale as

$$\mathbb{J} \left(\begin{array}{c} i \\ \text{---} \\ *+j \end{array} \text{ with a blue } \lambda \text{-labeled bubble } *+j \right) = D_i^j \frac{c_{i,\lambda}^+}{c_{i,\lambda}^{i+}} \quad \mathbb{J} \left(\begin{array}{c} i \\ \text{---} \\ *+j \end{array} \text{ with a blue } \lambda \text{-labeled bubble } *+j \right) = D_i^j \frac{c_{i,\lambda+\alpha_i}^{i+}}{c_{i,\lambda+\alpha_i}^+} \quad (3.28)$$

Hence,

$$\begin{aligned}
 & \left(\beta_{i,\lambda} \sum_{\substack{f_1+f_2+f_3 \\ =\lambda_i-1}} D_i^{f_1+f_3} \cdot D_i^{f_2} \frac{c_{i,\lambda+a_i}^+}{c_{i,\lambda}^+} \cdot D_i^{-\lambda_i+1} \frac{c_{i,\lambda}^+}{c_{i,\lambda+a_i}^+} \right) = \beta_{i,\lambda} \sum_{\substack{f_1+f_2+f_3 \\ =\lambda_i-1}} D_i^{f_1+f_3} \cdot D_i^{f_2} \frac{c_{i,\lambda+a_i}^+}{c_{i,\lambda}^+} \cdot D_i^{-\lambda_i+1} \frac{c_{i,\lambda}^+}{c_{i,\lambda+a_i}^+} \\
 & = \beta_{i,\lambda} \sum_{\substack{f_1+f_2+f_3 \\ =\lambda_i-1}} \frac{c_{i,\lambda}^+}{c_{i,\lambda+a_i}^+} \cdot \frac{c_{i,\lambda+a_i}^+}{c_{i,\lambda}^+} \stackrel{(2.2)}{=} \beta_{i,\lambda} \sum_{\substack{f_1+f_2+f_3 \\ =\lambda_i-1}} t_{ii}^{-1} t_{ii}'^{-\lambda_i-1} \frac{c_{i,\lambda}^+}{c_{i,\lambda+a_i}^+} \\
 & = \beta'_{i,\lambda} \sum_{\substack{f_1+f_2+f_3 \\ =\lambda_i-1}} \frac{c_{i,\lambda}^+}{c_{i,\lambda+a_i}^+} \cdot D_i^{-\lambda_i-1} \frac{c_{i,\lambda}^+}{c_{i,\lambda+a_i}^+} \quad (3.29)
 \end{aligned}$$

This computation together with Eqs. 3.26 and 3.27 prove this \mathfrak{sl}_2 -relation. The other is proven similarly. \square

Acknowledgments The author is partially supported by the NSF grants DMS-1255334 and DMS-1664240. He would also like to thank Joshua Sussan and Hoel Queffelec for comments on an early version of this note.

References

1. Beliakova, A., Habiro, K., Lauda, A.D., Webster, B.: Cyclicity for categorified quantum groups. *J. Algebra* **452**, 118–132 (2016). arXiv:1506.04671
2. Brundan, J.: On the definition of Kac-Moody 2-category. *Math. Ann.* **364**(1-2), 353–372 (2016). arXiv:1501.00350
3. Cautis, S.: Clasp technology to knot homology via the affine Grassmannian. *Math. Ann.* **363**(3-4), 1053–1115 (2015). arXiv:1207.2074
4. Cautis, S., Kamnitzer, J., Licata, A.: Categorical geometric skew Howe duality. *Inventiones Math.* **180**(1), 111–159 (2010)
5. Cautis, S., Lauda, A.D.: Implicit structure in 2-representations of quantum groups. *Selecta Mathematica*, pp. 1–44. arXiv:1111.1431 (2014)
6. Rose, D.E.V., Queffelec, H., Sartori, A.: Annular evaluation and link homology. arXiv:1802.04131 (2018)
7. Kashiwara, M.: Notes on parameters of quiver Hecke algebras. *Proc. Japan Acad. Ser. A Math. Sci.* **88**(7), 97–102 (2012)
8. Khovanov, M., Lauda, A.D.: A diagrammatic approach to categorification of quantum groups III. *Quantum Topology* **1**, 1–92 (2010). arXiv:0807.3250
9. Khovanov, M., Lauda, A.D.: A diagrammatic approach to categorification of quantum groups II. *Trans. Amer. Math. Soc.* **363**, 2685–2700 (2011). arXiv:0804.2080
10. Lauda, A.D.: A categorification of quantum $\mathfrak{sl}(2)$. *Adv. Math.* **225**, 3327–3424 (2008). arXiv:0803.3652
11. Lauda, A.D.: Categorified quantum $\mathfrak{sl}(2)$ and equivariant cohomology of iterated flag varieties. *Algebras and Representation Theory*, pp. 1–30. arXiv:0803.3848 (2009)
12. Lauda, A.D.: An introduction to diagrammatic algebra and categorified quantum \mathfrak{sl}_2 . *Bullet. Inst. Math. Acad. Sin.* **7**, 165–270 (2012). arXiv:1106.2128
13. Lauda, A.D., Queffelec, H., Rose, D.E.V.: Khovanov homology is a skew Howe 2-representation of categorified quantum \mathfrak{sl}_m . *Algebr. Geom. Topol.* **15**(5), 2517–2608 (2015). arXiv:1212.6076
14. Mackaay, M., Stošić, M., Vaz, P.: A diagrammatic categorification of the q -Schur algebra. *Quantum Topol.* **4**(1), 1–75 (2013)
15. Mackaay, M., Webster, B.: Categorified skew Howe duality and comparison of knot homologies. *Adv. Math.* **330**, 876–945 (2018). arXiv:1502.06011

16. Queffelec, H., Rose, D.E.V.: The \mathfrak{sl}_n foam 2-category: a combinatorial formulation of Khovanov-Rozansky homology via categorical skew Howe duality. *Adv. Math.* **302**, 1251–1339 (2016). arXiv:[1405.5920](https://arxiv.org/abs/1405.5920)
17. Rouquier, R.: 2-Kac-Moody algebras. arXiv:[0812.5023](https://arxiv.org/abs/0812.5023) (2008)
18. Webster, B.: Knot invariants and higher representation theory. *Mem. Amer. Math. Soc.* **250**(1191), v+141 (2017). arXiv:[1001.2020](https://arxiv.org/abs/1001.2020)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.