

OPERATIONS IN ÉTALE AND MOTIVIC COHOMOLOGY

BERT GUILLOU AND CHUCK WEIBEL

ABSTRACT. We classify all étale cohomology operations on $H_{\text{ét}}^n(-, \mu_{\ell}^{\otimes i})$, showing that they were all constructed by Epstein. We also construct operations P^a on the mod- ℓ motivic cohomology groups $H^{p,q}$, differing from Voevodsky's operations. We use them to classify all motivic cohomology operations on $H^{p,1}$ and $H^{1,q}$ and suggest a general classification.

In the last decade, several papers have given constructions of cohomology operations on motivic and étale cohomology, following the earlier work of Jardine [J], Kriz-May [KM], and Voevodsky [V2, V1]; see [BJ, BJ1, Jo, M1, V3, V4]. The goal of this paper is to provide, for each n and i , a classification of all such operations on the étale groups $H_{\text{ét}}^n(-, \mu_{\ell}^{\otimes i})$ and the motivic groups $H^{n,i}(-, \mathbb{F}_{\ell})$, similar to Cartan's classification of operations on singular cohomology $H_{\text{top}}^n(-, \mathbb{F}_{\ell})$ in [C]. We succeed for étale operations and partially succeed for motivic operations.

We work over a fixed field k and fix a prime ℓ with $1/\ell \in k$. By definition, an (unstable) étale cohomology operation on $H_{\text{ét}}^n(-, \mu_{\ell}^{\otimes i})$ over k is a natural transformation $H_{\text{ét}}^n(-, \mu_{\ell}^{\otimes i}) \rightarrow H_{\text{ét}}^p(-, \mu_{\ell}^{\otimes q})$ of set-valued functors from the category of (smooth) simplicial schemes over k (for some p and q). Similarly, an (unstable) motivic cohomology operation on $H^{n,i}$ over k is a natural transformation $H^{n,i} \rightarrow H^{p,q}$ of functors defined on this category, where $H^{p,q}(X)$ denotes the Nisnevich cohomology $H_{\text{nis}}^p(X, \mathbb{F}_{\ell}(q))$, and the cochain complex $\mathbb{F}_{\ell}(q)$ is defined in [V2] or [MVW]. Fixing k , n , and i , the set of all unstable cohomology operations forms a ring. The product of θ_1 and θ_2 is the operation $x \mapsto \theta_1(x) \cdot \theta_2(x)$.

Our classification theorems describe the ring of all operations in terms of certain specific operations. Thus we begin with their construction, in Sections 1, 4, and 6.

In étale cohomology with constant coefficients, operations P^a were constructed by Epstein [E] and used by Raynaud in [R]. A second construction is given by Peter May in [M]. As both are based on Steenrod's original construction in [SE], they agree. The upshot is that Cartan's ring $H_{\text{top}}^*(K_n)$ of operations on $H_{\text{top}}^n(-, \mathbb{F}_{\ell})$ embeds into the ring of all étale operations on $H_{\text{ét}}^n(-, \mu_{\ell}^{\otimes i})$; we refer the reader to Definition 0.1 below for a precise description of Cartan's ring.

For étale cohomology with twisted coefficients $\mu_{\ell}^{\otimes i}$, Epstein's approach [E] and May's approach [M] give apparently different constructions of étale operations P^a . We will show in Corollary 4.7 that the two constructions give the same operations.

Epstein's construction is more easily accessible to algebraic geometers, because it uses equivariant sheaf cohomology and is an application of the method described in his 1966 paper [E]. After stating Epstein's result in Theorem 1.3, we indicate the key points in his construction that we will need to compare with May's construction.

Received by the editors January 26, 2017, and, in revised form, April 18, 2018.

2010 *Mathematics Subject Classification*. Primary 14F42, 55S05; Secondary 14F20.

The classification of étale cohomology operations is given in Sections 2 and 3. Theorem 3.5 gives the general result: the ring of all (unstable) étale operations on $H_{\text{ét}}^n(-, \mu_{\ell}^{\otimes i})$ over k is the tensor product $H_{\text{ét}}^*(k(\zeta), \mathbb{F}_{\ell}) \otimes H_{\text{top}}^*(K_n)$, where ζ is a primitive ℓ th root of unity. Thus all (unstable) étale operations on $H_{\text{ét}}^n(-, \mu_{\ell}^{\otimes i})$ over k are $H_{\text{ét}}^*(k(\zeta), \mathbb{F}_{\ell})$ -linear combinations of monomials in the operations P^I . Our proof starts with the special case in which $\zeta \in k$. In this case, it is a result of Breen and Jardine that the graded ring of all étale operations on $H_{\text{ét}}^n(-, \mu_{\ell}^{\otimes i})$ is $H_{\text{ét}}^*(k, \mu_{\ell}^{\otimes *}) \otimes H_{\text{top}}^*(K_n)$.

In Section 4, we present May's construction, using the notion of a suitable pair (\mathcal{K}, θ) (see 4.2) and show in Corollary 4.7 that the étale operations P^a coincide. Our construction of motivic cohomology operations will use May's construction. The brief Section 5 relates the discussion of Section 4 to the operad-based approach of Hinich and Schechtman [HS].

In Section 6, we use the Norm Residue Theorem to construct motivic operations P^a (see Definition 6.5). We show they are compatible with the étale operations and stable under simplicial suspension, and we verify the usual properties in Section 7. The operation P^0 is the Frobenius $H^{n,i} \rightarrow H^{n,i\ell}$ on motivic cohomology, induced by the ℓ th power map $\mathbb{F}_{\ell}(i) \rightarrow \mathbb{F}_{\ell}(i\ell)$; see Proposition 8.4. One new result concerning Voevodsky's operations is that for $n > i$ and $x \in H^{2n,i}$ we have $P_V^n(x) = [\zeta^{\otimes(\ell-1)}]^{(n-i)} \cup x^{\ell}$, where $[\zeta^{\otimes(\ell-1)}]$ is the canonical element of $H^{0,\ell-1}(k)$ (see Corollary 8.10). This extends Lemma 9.8 of [V1], which states that $P^n(x) = x^{\ell}$ for $x \in H^{2n,n}(X)$.

The classification of motivic cohomology operations is complicated by the presence of more operations than those constructed by Voevodsky or via Steenrod-Epstein methods. One example is that an ℓ -torsion element t in the Brauer group of k gives an operation $H^{1,2} \rightarrow H^{3,3}$ by

$$H^{1,2}(X) \cong H_{\text{ét}}^1(X, \mu_{\ell}^{\otimes 2}) \xrightarrow{\cup t} H_{\text{ét}}^3(X, \mu_{\ell}^{\otimes 3}) \cong H^{3,3}(X).$$

Also unexpectedly, we may also use t and the Bockstein β to get an operation $H^{1,2}(X) \rightarrow H^{4,3}(X)$ (see Example 11.5 below). When k contains a primitive ℓ th root of unity ζ , we also have an interesting operation $H^{1,2}(X) \rightarrow H^{2,1}(X) = \text{Pic}(X)/\ell$: divide by the Bott element $[\zeta] \in H^{0,1}(k)$ and then apply the Bockstein; see Proposition 11.2.

In Section 10, we determine the ring of all motivic cohomology operations on $H^{n,1}$. If $\ell \neq 2$, it is the algebra $H^{*,*}(k) \otimes H_{\text{top}}^*(K_n)$, where $H^{*,*}(k)$ is the motivic cohomology of k and $H_{\text{top}}^*(K_n)$ is Cartan's ring, described in Definition 0.1 below. Many of these operations fail to be stable operations because they do not fit into a sequence of operations compatible with the motivic t -suspension $X \mapsto S_t^1 X$; for this reason, we call them *unstable*.

In Section 11, we determine the ring of (unstable) cohomology operations on $H^{1,i}$. When k contains the ℓ th roots of unity, this is the graded polynomial ring over $H^{*,*}(k)$ on operations $\gamma : H^{1,i}(X) \cong H^{1,1}(X)$ and its Bockstein, where γ is given by the Norm Residue Theorem 6.2. For general fields, it is the Galois-invariant subring. The operations on $H^{1,2}$ referred to above arise in this way.

Finally Section 12 contains a conjecture about what the general classification might be for $H^{n,i}$ when $n, i > 1$.

Since it is the topological prototype of our classification theorem, we conclude this introduction with a description of the ring of all singular cohomology operations

on $H_{\text{top}}^n(-, \mathbb{F}_\ell)$. Serre observed in [S50, 28.1] (cf. [EM, p. 513]) that the ring of operations from $H_{\text{top}}^n(-, \mathbb{F}_\ell)$ to $H_{\text{top}}^*(-, \mathbb{F}_\ell)$ is isomorphic to the cohomology $H_{\text{top}}^*(K_n)$ of the Eilenberg-Mac Lane space $K_n = K(\mathbb{F}_\ell, n)$; the structure of this ring was determined by Serre and Cartan in [S50], [C], [C1]. The following description is taken from [McC, 6.19].

Definition 0.1. For $\ell > 2$, let Λ_n denote the free graded-commutative \mathbb{F}_ℓ -algebra generated by the elements $P^I(\iota_n)$, where $I = (\epsilon_0, s_1, \epsilon_1, \dots, s_k, \epsilon_k)$ is an admissible sequence satisfying either $e(I) < n$ or else $e(I) = n$ and $\epsilon_0 = 1$.

Here the *excess* of I is defined to be $e(I) = 2 \sum (s_i - \ell s_{i+1} - \epsilon_i) + \sum_{i=0}^k \epsilon_i$, where $s_i = 0$ for $i > k$, and I is *admissible* if $s_i \geq \ell s_{i+1} + \epsilon_i$ for all $i < k$.

When $\ell = 2$, Λ_n denotes the free graded-commutative \mathbb{F}_2 -algebra generated by the elements $Sq^I(\iota_n)$, with $I = (s_1, \dots, s_k)$ admissible ($s_i \geq 2s_{i+1}$) and $e(I) < n$, where the excess is $e(I) = \sum (s_i - 2s_{i+1}) = s_1 - \sum_{i>1} s_i$.

We will write $H_{\text{top}}^*(K_n)$ for Λ_n because of the following result.

Theorem 0.2 (Cartan–Serre). *The ring $H_{\text{top}}^*(K_n, \mathbb{F}_\ell)$ of cohomology operations from $H_{\text{top}}^n(-, \mathbb{F}_\ell)$ to $H_{\text{top}}^*(-, \mathbb{F}_\ell)$ is isomorphic to Λ_n .*

For example, every operation on $H_{\text{top}}^2(-, \mathbb{F}_\ell)$ is a polynomial in id , β , the $P^I\beta$, and (if $\ell \neq 2$) the $\beta P^I\beta$ (where $P^I = P^{\ell^k} \dots P^\ell P^1$). This is because the only admissible sequences with excess < 2 are 0, (1), and $(0, \ell^k, 0, \dots, \ell, 0, 1, 1)$.

1. EPSTEIN'S ÉTALE CONSTRUCTION

Cohomology operations in étale cohomology were constructed by David Epstein long ago in the 1966 paper [E] as a special case of operations constructed in an axiomatic framework; see Sections 10 and 11.1 of [E]. Epstein's construction was made explicit by Michèle Raynaud [R, 4.4] for étale cohomology with constant coefficients. Alternative constructions were later given by L. Breen [Br, III.4] and J.F. Jardine [J, 1.4], [J1, §2], and [M1].

In Epstein's approach, one starts with an \mathbb{F}_ℓ -linear tensor abelian category \mathcal{Sh} (such as sheaves of \mathbb{F}_ℓ -modules on a site), a left exact functor $H^0(X, -)$ (global sections over X), and a commutative associative ring object \mathcal{A} of \mathcal{Sh} .

In this section, we consider the situation in which \mathcal{Sh} is the category of étale sheaves of \mathbb{F}_ℓ -modules on the big étale site of simplicial schemes X_\bullet over a base S , which we may assume is $\text{Spec}(k)$. The ring object \mathcal{A} will be the graded étale sheaf $\bigoplus_{i=0}^\infty \mu_\ell^{\otimes i}$.

Definition 1.1. If X_\bullet is a simplicial scheme over S , the étale site $\text{Et}(X_\bullet)$ is the category whose objects are pairs $(n, U \rightarrow X_n)$ with $U \rightarrow X_n$ étale. A morphism to (m, U') is an ordinal map $[m] \rightarrow [n]$, together with a map $U \rightarrow U'$ forming a commutative square with $X_n \rightarrow X_m$. A *covering* of (n, U) is a family of maps $U_i \rightarrow U$ over X_n so that the $U_i \rightarrow U$ are an étale cover of U .

The étale site of a simplicial scheme X_\bullet first arose in [D, 5.1.8]; our description is based upon the definition of the étale site $\text{Et}(X_\bullet)$ in [F, 1.4].

Recall that if $X = X_\bullet$ is a simplicial scheme and \mathcal{F} is a sheaf of \mathbb{F}_ℓ -modules, then the functor $\Gamma(\mathcal{F}) = \mathcal{F}(X)$ is defined as the equalizer of $\mathcal{F}(X_0) \rightrightarrows \mathcal{F}(X_1)$, and the cohomology functors $H_{\text{ét}}^i(X, \mathcal{F})$ are defined as its derived functors; see [D, 5.2.2]

or [F, 2.3]. If X_\bullet is a constant simplicial scheme, $H_{\text{et}}^i(X, \mathcal{F})$ is the usual étale cohomology of X .

The derived functors of $H^0(X, -) : \mathcal{S}h \rightarrow \mathbb{F}_\ell\text{-mod}$ are just the usual étale cohomology groups, because the usual Godement resolution of a sheaf \mathcal{F} [Milne, p. 90] is a flasque resolution by sheaves of \mathbb{F}_ℓ -modules which are injective objects of $\mathcal{S}h$.

The étale Bockstein $\beta : H_{\text{et}}^n(X, \mu_\ell^{\otimes i}) \rightarrow H_{\text{et}}^{n+1}(X, \mu_\ell^{\otimes i})$ is defined as the connecting map in the cohomology sequence for $0 \rightarrow \mu_\ell^{\otimes i} \rightarrow \mu_{\ell^2}^{\otimes i} \rightarrow \mu_\ell^{\otimes i} \rightarrow 0$. (This sequence is the tensor product of the sequence $0 \rightarrow \mathbb{Z}/\ell \rightarrow \mathbb{Z}/\ell^2 \rightarrow \mathbb{Z}/\ell \rightarrow 0$ with $\mu_{\ell^2}^{\otimes i}$.) By definition, β is natural in X but can depend on the choice of i ; see Remark 3.3.1 below. If $\mathcal{B} = \bigoplus_{i=0}^{\infty} \mu_{\ell^2}^{\otimes i}$, we may also regard β as the connecting map in the cohomology sequence for $0 \rightarrow \mathcal{A} \xrightarrow{i} \mathcal{B} \rightarrow \mathcal{A} \rightarrow 0$.

Lemma 1.2. *The étale Bockstein is a derivation. That is, if $u \in H_{\text{et}}^n(X, \mu_\ell^{\otimes i})$ and $v \in H_{\text{et}}^m(X, \mu_\ell^{\otimes j})$, then $\beta(u \cup v) = \beta(u) \cup v + (-1)^n u \cup \beta(v)$.*

Proof. (Folklore) Choose a flasque Godement-style resolution $\mathcal{B} \rightarrow \mathcal{I}$ whose stalks are free (=injective) \mathbb{Z}/ℓ^2 -modules, and write $\bar{\mathcal{I}}$ for $\mathcal{I}/\ell\mathcal{I}$, so that $\mathcal{A} \rightarrow \bar{\mathcal{I}}$ is also a flasque resolution. Lifting cycles \bar{u} and \bar{v} representing u and v to chains $u' \in \mathcal{I}^n(X)$ and $v' \in \mathcal{I}^m(X)$, $\beta(u)$ and $\beta(v)$ are represented by u'' and v'' , defined by $\delta(u') = i(u'')$ and $\delta(v') = i(v'')$.

The cup product $u \cup v$ is represented by the image of $\bar{u} \otimes \bar{v}$ under the map $m : \bar{\mathcal{I}} \otimes \bar{\mathcal{I}} \rightarrow \bar{\mathcal{I}}$ resolving $\mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$; see [D-4.5, 1.2.2]. Since the coboundary on $\mathcal{I} \otimes \mathcal{I}$ satisfies

$$\delta(u' \otimes v') = i(u'') \otimes v' + (-1)^n u' \otimes i(v'') = i(u'' \otimes \bar{v}) + (-1)^n i(\bar{u} \otimes v'')$$

it follows that $\beta(u \cup v)$ is represented by $m(u'' \otimes \bar{v}) + (-1)^n m(\bar{u} \otimes v'')$, i.e., by $\beta(u) \cup v + (-1)^n u \cup \beta(v)$. \square

Remark 1.2.1. The same proof works in the motivic setting to show that the motivic Bockstein is also a derivation, a fact stated in [V1, (8.1)].

Epstein defines an operation $P^0 : H_{\text{et}}^n(X, \mu_\ell^{\otimes i}) \rightarrow H_{\text{et}}^n(X, \mu_\ell^{\otimes i\ell})$ in [E, 7.1] and shows in [E, 7.3] that P^0 is the canonical Frobenius isomorphism induced by $\mu_\ell^{\otimes i} \cong \mu_\ell^{\otimes i\ell}$. (Our pairing $H_{\text{et}}^0(X, \mu_\ell^{\otimes i}) \otimes H_{\text{et}}^0(X, \mu_\ell^{\otimes j}) \rightarrow H_{\text{et}}^0(X, \mu_\ell^{\otimes i+j})$ is the pairing $SA \otimes TB \rightarrow U(A \otimes B)$ in [E, (3.2.1)].) Epstein also defines an operation

$$Q^0 : H_{\text{et}}^n(X, \mu_\ell^{\otimes i}) \rightarrow H_{\text{et}}^{n+1}(X, \mu_\ell^{\otimes i\ell});$$

we'll see in Proposition 3.3 below that $Q^0 = \beta P^0$ and $Q^a = \beta P^a$, where β is the Bockstein on $H_{\text{et}}^*(X, \mu_\ell^{\otimes i\ell})$. With this dictionary, Epstein's theorem specializes to yield:

Theorem 1.3. *For each odd prime ℓ , there are additive cohomology operations*

$$P^a : H_{\text{et}}^n(X, \mu_\ell^{\otimes i}) \rightarrow H_{\text{et}}^{n+2a(\ell-1)}(X, \mu_\ell^{\otimes i\ell}), \quad a \geq 0,$$

natural in X , satisfying the usual relations: $P^a x = x^\ell$ if $n = 2a$, $P^a x = 0$ if $n < 2a$, the Cartan relation $P^a(xy) = \sum P^i(x)P^j(y)$, and Adem relations for both $P^a P^b$ ($a < b\ell$) and $P^a \beta P^b$ ($a \leq b\ell$).

When $\ell = 2$, there are Steenrod operations $Sq^a : H_{\text{et}}^n(X, \mu_2^{\otimes i}) \rightarrow H_{\text{et}}^{n+a}(X, \mu_2^{\otimes 2i})$ or $H_{\text{et}}^n(X, \mathbb{Z}/2) \rightarrow H_{\text{et}}^{n+a}(X, \mathbb{Z}/2)$, satisfying the usual relations.

Proof. The existence and basic properties are given in Chapter 7 of [E]; additivity is 6.7. The Adem relations are established in [E, 9.7–8], using the dictionary that $Q^b = \beta P^b$ and $P^a \beta P^b = P^a Q^b$. Naturality follows from [E, 11.1(8)]. \square

Remark 1.3.1.

- (i) There are canonical isomorphisms $\mu_\ell \cong \mu_\ell^{\otimes \ell}$ and $\mu_\ell^{\otimes i} \cong \mu_\ell^{\otimes i \ell}$. We have emphasized the difference in weight because of our application to motivic operations.
- (ii) Epstein also constructs operations Q^a ; they equal βP^a if we use the sheaf $\mathcal{A} = \bigoplus_{i=0}^{\infty} \mu_\ell^{\otimes i}$ but can differ if we use the smaller sheaf $A = \bigoplus_{i=0}^d \mu_\ell^{\otimes i}$, where $d = [k(\zeta_\ell) : k]$; see Propositions 2.4 and 3.3.
- (iii) If Z is a closed simplicial subscheme of X (see [D] or [F]), we get cohomology operations P^a on the relative groups $H_{\text{ét}}^a(X, Z; \mu_\ell^{\otimes i})$, natural in the pair (X, Z) , by replacing $H^0(X, -)$ by the left exact functor $H_{\text{ét}}^0(X, Z; -)$.

In order to compare to May's construction and to classify operations, we will need a rephrasing of one of the key results from [E], using the language of equivariant sheaf cohomology.

Definition 1.4. If G is a finite group, we write \mathcal{Sh}_G for the category of G -equivariant objects of \mathcal{Sh} , i.e., objects B equipped with a homomorphism $G \rightarrow \text{End}(B)$. If B is in \mathcal{Sh}_G , then $H^0(X, B)$ is a G -module, and we define the left exact functor $H_G^0(X, -)$ on the category \mathcal{Sh}_G by the formula $H_G^0(X, B) = H^0(X, B)^G$. We write $H_G^*(X, -)$ for the derived functors of $H_G^0(X, -)$.

We will use the following result in Section 6.

Theorem 1.5. *Let A be a bounded below cochain complex of objects of \mathcal{Sh} on which a finite group G acts trivially. Then there is a natural isomorphism*

$$H^*(G, \mathbb{F}_\ell) \otimes H^*(X, A) \xrightarrow{\sim} H_G^*(X, A).$$

If A is a sheaf of dg commutative algebras, this is an algebra isomorphism.

Proof (See [E, 4.4.4]). Fix an injective resolution $A \xrightarrow{\sim} I^*$ in \mathcal{Sh} . Choosing a resolution $F_* \rightarrow \mathbb{F}_\ell$ by finitely generated free $\mathbb{F}_\ell[G]$ -modules, set $F^* = \text{Hom}(F_*, \mathbb{F}_\ell)$, so $H^*(G, \mathbb{F}_\ell)$ is the cohomology of $(F^*)^G$. Since G acts trivially on A , we have quasi-isomorphisms of complexes in \mathcal{Sh}_G : $A \xrightarrow{\sim} I^* = \mathbb{F}_\ell \otimes I^* \xrightarrow{\sim} \text{Tot}(F^* \otimes I^*)$. Since each F^n is a free $\mathbb{F}_\ell[G]$ -module of finite rank, $F^n \otimes I^q \cong \text{Hom}(F_n, I^q)$ is injective in \mathcal{Sh}_G , and $A \rightarrow \text{Tot}(F^* \otimes I^*)$ is an injective resolution in \mathcal{Sh}_G . Hence $H_G^*(X, A)$ is the cohomology of the total complex of

$$H_G^0(X, F^* \otimes I^*) = (F^*)^G \otimes I^*(X).$$

The Künneth formula tells us that $H^*((F^*)^G \otimes I^*(X))$ is the tensor product of the cohomology of $(F^*)^G$ and $I^*(X)$, i.e., of $H^*(G, \mathbb{F}_\ell)$ and $H^*(X, A)$.

We omit the standard proof that a commutative associative product on A induces an algebra structure on $H^*(X, A)$ and $H_G^*(X, A)$ and that the isomorphism of Theorem 1.5 commutes with products. \square

Recall that for any sheaf (or complex) A , the symmetric group S_n acts on $A^{\otimes n}$ by permuting factors (with the usual sign change for tensor products of complexes). If $A \xrightarrow{\sim} B$ is a quasi-isomorphism, then so is $A^{\otimes n} \xrightarrow{\sim} B^{\otimes n}$ by the following useful lemma.

Lemma 1.6. *If C , C' , and D are bounded below cochain complexes of sheaves of \mathbb{F}_ℓ -modules and $f : C \xrightarrow{\sim} C'$ is a quasi-isomorphism, then so is $f \otimes 1 : C \otimes D \rightarrow C' \otimes D$.*

Proof. Let K denote the cone of f ; it is acyclic. Since \otimes is an exact functor, every sheaf of \mathbb{F}_ℓ -modules is flat, and $K \otimes D$ is acyclic by the Künneth formula [WH, 3.6.1]. Since $C \otimes D \rightarrow C' \otimes D \rightarrow K \otimes D$ is a distinguished triangle, the result follows. \square

Remark 1.6.1. If A is a sheaf of commutative \mathbb{F}_ℓ -algebras and $A \xrightarrow{\sim} I$ is an injective resolution, the choice of a lift $I^{\otimes 2} \rightarrow I$ of $A^{\otimes 2} \rightarrow A$ makes I a sheaf of homotopy commutative, homotopy unital, and homotopy associative dg algebras, as $A^{\otimes 2} \xrightarrow{\sim} I^{\otimes 2}$. Such a lift exists and is unique up to chain homotopy, by the comparison theorem [WH, 2.3.7, 10.4.7].

Similar remarks hold when A is a sheaf of bounded below, homotopy associative and commutative dg algebras, using the total complex I of a Cartan-Eilenberg resolution; see [WH, 5.7.9 and Ex. 5.7.2].

Let π be a Sylow ℓ -subgroup of S_ℓ . Choosing an injective resolution $A^{\otimes \ell} \rightarrow J^*$ in Sh_π , the comparison theorem lifts the equivariant quasi-isomorphism $A^{\otimes \ell} \rightarrow I^{\otimes \ell}$ to an equivariant map $I^{\otimes \ell} \rightarrow J^*$, unique up to chain homotopy.

Since $H^*(X, A)$ is the cohomology of $I(X)$, we can represent any element of $H^n(X, A)$ by an n -cocycle $u \in I^n(X)$. The $n\ell$ -cocycle $u \otimes \cdots \otimes u$ of $I(X)^{\otimes \ell}$ is π -invariant, because the generator of π acts as multiplication by $(-1)^{n(\ell-1)}$, which is the identity on any \mathbb{F}_ℓ -module. Its image Pu in $J^{n\ell}(X)$ is also π -invariant. Epstein shows in [E, 5.1.3] that $P(u + dv) = Pu + dw$ for $v \in I^{n-1}(X)$ and $w \in J^{n\ell-1}(X)$, so the cohomology class of Pu is independent of the choice of cocycle u .

Definition 1.7. The *reduced power map* is defined to be the map on cohomology associated to $u \mapsto u \otimes \cdots \otimes u$:

$$P : H^n(X, A) \rightarrow H_\pi^{n\ell}(X, A^{\otimes \ell}).$$

Now let π denote the cyclic group of order ℓ . We will write $W_* \rightarrow \mathbb{F}_\ell$ for the standard periodic $\mathbb{F}_\ell[\pi]$ -resolution [WH, 6.2.1], with generator e_k of $W_k \cong \mathbb{F}_\ell[\pi]$, and set $W^* = \text{Hom}(W_*, \mathbb{F}_\ell)$; thus $H^*(\pi, \mathbb{F}_\ell)$ is the cohomology of $(W^*)^\pi$.

Now suppose that there is a π -equivariant map $A^{\otimes \ell} \xrightarrow{m} B$ and that π acts trivially on B . (When A is a commutative ring, multiplication $A^{\otimes \ell} \rightarrow A$ is a π -equivariant map.) We write m_* for the induced map $H_\pi^*(X, A^{\otimes \ell}) \rightarrow H_\pi^*(X, B)$. By Theorem 1.5, $m_* P(u) \in H_\pi^*(X, B)$ has an expansion $\sum w_k \otimes D_k(u)$, where $w_k \in H^k(\pi, \mathbb{F}_\ell)$ are the (dual) basis elements of [SE, V.5.2]: if $\ell > 2$, then $w_0 = 1$, $w_2 = \beta w_1$, $w_{2i} = w_2^i$, and $w_{2i+1} = w_1 w_2^i$. If $\ell > 2$ and $n \geq 2a$, Epstein defines

$$(1.8) \quad P^a : H^n(X, A) \rightarrow H^{n+2a(\ell-1)}(X, B), \quad P^a u = (-1)^a \nu_n D_{(n-2a)(\ell-1)}(u),$$

where

$$\nu_n = (-1)^r \left(\frac{\ell-1}{2} \right)!^{-n} \quad \text{and} \quad r = \frac{(\ell-1)(n^2+n)}{4}.$$

(See [E, 7.1], [SE, VII.6.1], and [SE-err].) If $n < 2a$, then Epstein defines $P^a = 0$.

When $\ell = 2$, Epstein defines operations Sq^i by $Sq^i(u) = D_{n-i}(u)$ for $n \geq i$ and $Sq^i(u) = 0$ for $n < i$.

Remark 1.8.1. Epstein also defines operations $Q^a = (-1)^{a+1} \nu_n D_{(n-2a)(\ell-1)-1}(u)$ in this setting, with $Q^a = 0$ when $n \leq 2a$, and establishes Adem relations for them as well.

Of course, Epstein's construction mimicks Steenrod's construction of D_k , P^a , and Q^a (see [SE], VII.3.2 and VII.6.1). In Steenrod's setting one can lift to integral cochains; with this assumption, Steenrod proves that $\beta D_{2k} = -D_{2k+1}$ and hence that $\beta P^a = Q^a$; see [SE, VII.4.6] and [SE-err]. As we mentioned in Remark 1.3.1, the formula $Q^a = \beta P^a$ may not hold when the sheaf A fails to distinguish between $\mu_\ell^{\otimes i}$ and $\mu_\ell^{\otimes i\ell}$. (See Propositions 2.4 and 3.3 and Theorem 8.11.)

Recall that the simplicial suspension SX of a simplicial scheme X is again a simplicial scheme. There is a canonical isomorphism $\tilde{H}_{\text{ét}}^n(X, \mu_\ell^{\otimes i}) \xrightarrow{\cong} \tilde{H}_{\text{ét}}^{n+1}(SX, \mu_\ell^{\otimes i})$.

Proposition 1.9. *The operations P^a are simplicially stable in the sense that they commute with simplicial suspension: there are commutative diagrams for all X and all n and i , with $N = n + 2a(\ell - 1)$:*

$$\begin{array}{ccc} \tilde{H}_{\text{ét}}^n(X, \mu_\ell^{\otimes i}) & \xrightarrow{P^a} & \tilde{H}_{\text{ét}}^N(X, \mu_\ell^{\otimes i\ell}) \\ \cong \downarrow & & \cong \downarrow \\ \tilde{H}_{\text{ét}}^{n+1}(SX, \mu_\ell^{\otimes i}) & \xrightarrow{P^a} & \tilde{H}_{\text{ét}}^{N+1}(SX, \mu_\ell^{\otimes i\ell}). \end{array}$$

Proof. The proofs of Lemmas 1.2 and 2.1 of [SE] go through, using simplicial homotopy invariance ($H_{\text{ét}}^*(X) \cong H_{\text{ét}}^*(X \times \Delta^1)$) of étale cohomology and excision. \square

If $f : X \rightarrow Y$ is a finite map, then $f_* : \mathcal{Sh}(X) \rightarrow \mathcal{Sh}(Y)$ is an exact functor. It follows that a π -equivariant map $A^{\otimes \ell} \xrightarrow{m} B$ of étale sheaves on X induces a π -equivariant map $f_* m : (f_* A)^{\otimes \ell} \cong f_*(A^{\otimes \ell}) \rightarrow f_* B$ and hence operations P^a , Q^a from $H^*(Y, f_* A)$ to $H^*(Y, f_* B)$.

Proposition 1.10. *Suppose that $f : X \rightarrow Y$ is a finite map and $A^{\otimes \ell} \xrightarrow{m} B$ is a π -equivariant map of sheaves on X . Then Epstein's P^a and Q^a commute with the isomorphism $f_* : H^*(Y, f_* A) \xrightarrow{\cong} H^n(X, A)$.*

Proof. Let I and J be the injective resolutions of A and $A^{\otimes \ell}$ in the construction 1.7 of the power map P for A . Then $f_* I$ and $f_* J$ are injective resolutions of $f_* A$ and $f_* A^{\otimes \ell}$, and if u is a cocycle in $I^n(X) = (f_* I^n)(Y)$, then $u \otimes \cdots \otimes u$ is a cocycle in $I(X)^{\otimes \ell} \cong (f_* I)(Y)^{\otimes \ell}$, and $I^*(X) \rightarrow J^*(X)$ is identified with $f_* I^*(Y) \rightarrow f_* J^*(Y)$. It follows that the left square commutes in the diagram

$$\begin{array}{ccccc} H^n(Y, f_* A) & \xrightarrow{P} & H_{\pi}^{n\ell}(Y, f_* A^{\otimes \ell}) & \xrightarrow{m_*} & H^{n\ell}(Y, f_* B) \\ \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\ H^n(X, A) & \xrightarrow{P} & H_{\pi}^{n\ell}(X, A^{\otimes \ell}) & \xrightarrow{m_*} & H^{n\ell}(X, B). \end{array}$$

The right square commutes by the functoriality of f_* . Expanding $m_* P(f_* u)$ as in Theorem 1.5, the result follows from (1.8) and Remark 1.8.1. \square

2. THE ÉTALE STEENROD ALGEBRA WHEN $\zeta_\ell \in k$

In this section and the next we determine the algebra of all étale cohomology operations $H_{\text{ét}}^n(-, \mu_\ell^{\otimes i}) \rightarrow H_{\text{ét}}^*(-, \mu_\ell^{\otimes *})$ for each n and i , over a field k containing

$1/\ell$. We work in the big étale site of smooth simplicial schemes over k (see Definition 1.1).

Recall from SGA 4 (V.2.1.2 in [Ver]) that if M is a (simplicial) étale sheaf of \mathbb{F}_ℓ -modules, then the sheaf cohomology groups $H_{\text{ét}}^*(X, M)$ are isomorphic to the (hyper) Ext-groups $\text{Ext}^*(\mathbb{F}_\ell[X], M)$ in the category of étale sheaves of \mathbb{F}_ℓ -modules. (Here we regard M as a cochain complex using Dold-Kan.) If K is a second simplicial étale sheaf of \mathbb{F}_ℓ -modules, one writes $H_{\text{ét}}^*(K, M)$ for $\text{Ext}^*(K, M)$.

It is well known that cohomology operations $H_{\text{ét}}^n(-, L) \rightarrow H_{\text{ét}}^*(-, M)$ are in 1–1 correspondence with elements of $H_{\text{ét}}^*(K, M)$, where K denotes the standard simplicial Eilenberg-Mac Lane sheaf $K(L, n)$ associated to L . If M is a ring, these operations form a ring; the product of θ_1 and θ_2 is $x \mapsto \theta_1(x) \cdot \theta_2(x)$.

Now fix i and consider cohomology operations $H_{\text{ét}}^n(-, \mu_\ell^{\otimes i}) \rightarrow H_{\text{ét}}^*(-, \mu_\ell^{\otimes j})$. As observed above, they are in 1–1 correspondence with elements of $H_{\text{ét}}^*(K, \mu_\ell^{\otimes j})$, where K denotes the simplicial Eilenberg-Mac Lane scheme $K(\mu_\ell^{\otimes i}, n)$. For example, the identity operation on $H_{\text{ét}}^n(-, \mu_\ell^{\otimes i})$ corresponds to $\iota_n \in H_{\text{ét}}^n(K, \mu_\ell^{\otimes i})$, and the étale Bockstein $\beta : H_{\text{ét}}^n(X, \mu_\ell^{\otimes i}) \rightarrow H_{\text{ét}}^{n+1}(X, \mu_\ell^{\otimes i})$ corresponds to $\beta(\iota_n) \in H_{\text{ét}}^{n+1}(K, \mu_\ell^{\otimes i})$.

We first discuss the case of constant coefficients ($M = \mathbb{F}_\ell$), which is known and due to Breen [Br, 4.3–4] and Jardine [J]. The graded ring of all unstable étale cohomology operations from $H_{\text{ét}}^n(-, \mathbb{F}_\ell)$ to $H_{\text{ét}}^*(-, \mathbb{F}_\ell)$ is isomorphic to the cohomology ring $H_{\text{ét}}^*(K_n, \mathbb{F}_\ell)$, where $K_n = K(\mathbb{F}_\ell, n)$ is the constant simplicial sheaf classifying elements of $H_{\text{ét}}^n(-, \mathbb{F}_\ell)$. By Theorem 1.3, there is a ring homomorphism from the classical unstable Steenrod algebra $H_{\text{top}}^*(K_n)$ of Definition 0.1 to $H_{\text{ét}}^*(K_n, \mathbb{F}_\ell)$.

There is also a ring homomorphism from $H_{\text{ét}}^*(k, \mathbb{F}_\ell)$ to $H_{\text{ét}}^*(K_n, \mathbb{F}_\ell)$, sending a to the constant operation $\theta(x) = a$. It is injective and is induced by $K_n \rightarrow \text{Spec } k$. These induce a graded algebra homomorphism from $H_{\text{ét}}^*(k, \mathbb{F}_\ell) \otimes_{\mathbb{F}_\ell} H_{\text{top}}^*(K_n)$ to $H_{\text{ét}}^*(K_n, \mathbb{F}_\ell)$. Note that $H_{\text{ét}}^*(k, \mathbb{F}_\ell) \otimes_{\mathbb{F}_\ell} H_{\text{top}}^*(K_n)$ is free as a left $H_{\text{ét}}^*(k, \mathbb{F}_\ell)$ -module; a basis is given by those monomials P^I in the Steenrod operations P^a and βP^a for which I has excess $< n$, exactly as in the topological case.

We summarize the above discussion.

Theorem 2.1 (Breen-Jardine). *The ring of all étale cohomology operations on $H_{\text{ét}}^n(-, \mathbb{F}_\ell)$ is the graded tensor product $H_{\text{ét}}^*(k, \mathbb{F}_\ell) \otimes H_{\text{top}}^*(K_n)$: every operation is a polynomial in the operations P^I with coefficients in $H_{\text{ét}}^*(k, \mathbb{F}_\ell)$.*

Examples 2.2. When $k = \mathbb{C}$, $H_{\text{ét}}^*(\mathbb{C}, \mathbb{F}_\ell) \cong \mathbb{F}_\ell$ and étale operations are classified by $H_{\text{top}}^*(K_n)$. The action of the P^I is compatible with the canonical comparison isomorphism $H_{\text{ét}}^*(X, \mathbb{F}_\ell) \cong H_{\text{top}}^*(X(\mathbb{C}), \mathbb{F}_\ell)$. This is clear from the constructions in [E] and [J].

When $k = \mathbb{R}$ and $\ell = 2$, the ring of étale cohomology operations over \mathbb{R} is the graded polynomial ring $H_{\text{top}}^*(K_n)[\sigma]$, generated over \mathbb{F}_2 by σ in degree 1 and the $Sq^I(\iota_n)$ with I admissible and $e(I) < n$. This is because $H_{\text{ét}}^*(\mathbb{R}, \mathbb{F}_2) = \mathbb{F}_2[\sigma]$.

Remark 2.2.1. Let $H_{\text{top}}^*(K_n)^{\text{ind}}$ denote the indecomposable subspace of $H_{\text{top}}^*(K_n)$; by Definition 0.1 it has the $P^I(\iota_n)$ as a basis, and $H_{\text{top}}^p(K_n)^{\text{ind}}$ injects into the degree $p - n$ part of the topological Steenrod algebra. It follows that the vector space $H_{\text{ét}}^*(k, \mathbb{F}_\ell) \otimes_{\mathbb{F}_\ell} H_{\text{top}}^*(K_n)^{\text{ind}}$ embeds into the algebra of *stable* étale operations with a degree shift; cf. [J]. Note that the multiplication \circ in the stable algebra is different: $P^a \circ \lambda = \sum P^i(\lambda)P^j$, $\lambda \in H_{\text{ét}}^*(k, \mathbb{F}_\ell)$, and $a > 0$, arising from the Cartan formula.

Example 2.3. When k^\times contains μ_ℓ but not μ_{ℓ^2} , the étale sheaves $\mu_{\ell^2}^{\otimes i}$ are distinct nontrivial extensions of \mathbb{F}_ℓ by \mathbb{F}_ℓ ($i = 0, \dots, \ell - 1$), and the associated boundary maps ∂_i in the exact sequence

$$(2.3.1) \quad \cdots H^n(X, \mathbb{F}_\ell) \rightarrow H^n(X, \mu_{\ell^2}^{\otimes i}) \rightarrow H^n(X, \mathbb{F}_\ell) \xrightarrow{\partial_i} H^{n+1}(X, \mathbb{F}_\ell) \cdots$$

are cohomology operations on $H_{\text{ét}}^n(-, \mathbb{F}_\ell)$; ∂_0 is the Bockstein β . By Theorem 2.1, the ∂_i are linear combinations of the generators β of $H_{\text{top}}^1(K_n)$ and elements of $H_{\text{ét}}^1(k, \mathbb{F}_\ell)$, such as $z = \zeta^{-1}\beta(\zeta)$, $\zeta \in \mu_\ell$. Note that z is nontrivial as $\mu_{\ell^2} \not\subset k^\times$. It is an exercise to check that z is independent of the choice of generator ζ of μ_ℓ .

Lemma 2.3.2. *If $\mu_\ell \subset k^\times$, the cohomology operation ∂_i of (2.3.1) is $\beta - iz$.*

Proof. The sheaf ring $\bigoplus \mu_{\ell^2}^{\otimes j}$ acts on the extensions $0 \rightarrow \mathbb{F}_\ell \rightarrow \mu_{\ell^2}^{\otimes i} \rightarrow \mathbb{F}_\ell \rightarrow 0$, so there is a pairing between $H_{\text{ét}}^*(k, R)$ and the direct sum of the cohomology sequences (2.3.1). Set $H^{0j} = H_{\text{ét}}^0(k, \mu_{\ell^2}^{\otimes j})$; when $j \neq 0 \pmod{\ell}$, this group is \mathbb{Z}/ℓ . This yields a commutative diagram for each i and j :

$$\begin{array}{ccccc} H^{0j} \otimes H_{\text{ét}}^n(X, \mu_{\ell^2}^{\otimes i}) & \longrightarrow & H^{0j} \otimes H_{\text{ét}}^n(X, \mathbb{F}_\ell) & \xrightarrow{1 \otimes \partial_i} & H^{0j} \otimes H_{\text{ét}}^{n+1}(X, \mathbb{F}_\ell) \\ \cup \downarrow & & \cup \downarrow \simeq & & \cup \downarrow \simeq \\ H_{\text{ét}}^n(X, \mu_{\ell^2}^{\otimes i+j}) & \longrightarrow & H_{\text{ét}}^n(X, \mathbb{F}_\ell) & \xrightarrow{1 \otimes \partial_{i+j}} & H_{\text{ét}}^{n+1}(X, \mathbb{F}_\ell). \end{array}$$

Now set $j = \ell - i$, so $\mu_{\ell^2}^{\otimes i+j} = \mu_{\ell^2}^0 = \mathbb{Z}/\ell^2$. Then the bottom right map ∂_{i+j} is the Bockstein β . For $u \in H_{\text{ét}}^n(X, \mathbb{F}_\ell)$ and $[\zeta^{\otimes j}] \in H^{0j}$ we have

$$\beta([\zeta^{\otimes j}] \cup u) = [\zeta^{\otimes j}] \cup \beta(u) + jz \cup [\zeta^{\otimes j}] \cup u = [\zeta^{\otimes j}] \cup (\beta(u) + jz).$$

As this equals $[\zeta^{\otimes j}] \cup \partial_i(u)$, we are done. \square

Proposition 2.4. *If $\mu_\ell \subset k$, Epstein's cohomology operation Q^0 agrees with the Bockstein β on $H_{\text{ét}}^n(X, \mathbb{F}_\ell)$, and $Q^a = \beta P^a$.*

Proof. Jardine's argument in [J, pp. 108–114] that Epstein's Sq^1 is the Bockstein when $\ell = 2$ applies when $\ell > 2$ as well and proves that Epstein's Q^0 is the Bockstein operation. For Q^a , we invoke the Adem relation $Q^a = Q^a P^0 = Q^0 P^a$ [E, 9.8(4)]. \square

3. THE ÉTALE STEENROD ALGEBRA WHEN $\zeta_\ell \notin k$

We now discuss the twisted coefficient case ($M = \mu_\ell^{\otimes i}$). This reduces to Theorem 2.1 when k contains a primitive ℓ th root of unity: since the sheaves $\mu_\ell^{\otimes i}$ are all isomorphic to \mathbb{F}_ℓ , the ring of étale operations on $H_{\text{ét}}^n(-, \mu_\ell^{\otimes i})$ is just $H_{\text{ét}}^*(k, \mathbb{F}_\ell) \otimes H_{\text{top}}^*(K_n)$. Since this is always the case when $\ell = 2$, we shall restrict to the case of an odd prime ℓ .

Fix a field k with $1/\ell \in k$, and let G be the Galois group of the extension $k(\zeta)/k$, where ζ denotes a primitive ℓ th root of unity. Then G is cyclic of order $d = [k(\zeta) : k]$, $d \mid \ell - 1$, and $1 \mapsto \zeta^{\otimes d}$ defines an isomorphism $\mathbb{F}_\ell \xrightarrow{\cong} \mu_\ell^{\otimes d}$. Since $i \equiv j \pmod{d}$ implies that $\mu_\ell^{\otimes i} \cong \mu_\ell^{\otimes j}$, we are led to consider the \mathbb{Z}/d -graded étale sheaf of Kummer algebras

$$A = \bigoplus_{i=0}^{d-1} \mu_\ell^{\otimes i}.$$

Thus our problem is to determine the ring $H_{\text{et}}^*(K_n, A) = \bigoplus_{j=0}^{d-1} H_{\text{et}}^*(K_n, \mu_\ell^{\otimes i})$.

Epstein defines cohomology operations P_A^a and Q_A^a on each $H_{\text{et}}^n(-, A)$; setting $N = n + 2a(\ell - 1)$, their components are operations

$$P_A^a : H_{\text{et}}^n(X, \mu_\ell^{\otimes i}) \rightarrow H_{\text{et}}^N(X, \mu_\ell^{\otimes i}) \quad \text{and} \quad Q_A^a : H_{\text{et}}^n(X, \mu_\ell^{\otimes i}) \rightarrow H_{\text{et}}^{N+1}(X, \mu_\ell^{\otimes i}).$$

By naturality with respect to the homomorphism $\mathcal{A} = \bigoplus_{i=0}^{\infty} \mu_\ell^{\otimes i} \rightarrow A$ [E, 6.2], the operations P^a and Q^a defined using \mathcal{A} are compatible with the operations P_A^a and Q_A^a . Since the cup product with $\zeta^{\otimes i(\ell-1)} \in H_{\text{et}}^0(k, \mu_\ell^{\otimes i(\ell-1)})$ induces an isomorphism $H_{\text{et}}^N(X, \mu_\ell^{\otimes i}) \cong H_{\text{et}}^N(X, \mu_\ell^{\otimes i\ell})$, this means that

$$(3.1) \quad P^a(u) = \zeta^{\otimes i(\ell-1)} \cup P_A^a(u) \quad \text{and} \quad Q^a(u) = \zeta^{\otimes i(\ell-1)} \cup Q_A^a(u).$$

Since ℓ does not divide $|G|$, Maschke's Theorem gives an identification of the étale sheaf $\mathbb{F}_\ell[G]$ with the direct sum of the sheaves of irreducible $\mathbb{F}_\ell[G]$ -modules $\mu_\ell^{\otimes i}$, i.e., with A . For any X , Shapiro's Lemma provides an isomorphism

$$(3.2) \quad H_{\text{et}}^*(X, A) \xrightarrow{\cong} H_{\text{et}}^*(X(\zeta), \mathbb{F}_\ell),$$

where $X(\zeta)$ denotes $X \times_k \text{Spec}(k(\zeta))$. In fact, $A = \pi_* \mathbb{F}_\ell \simeq R\pi_* \mathbb{F}_\ell$, where $\pi : \text{Spec}(k(\zeta))_{\text{et}} \rightarrow \text{Spec}(k)_{\text{et}}$. Taking $X = \text{Spec}(k)$ yields $H_{\text{et}}^*(k, A) \cong H_{\text{et}}^*(k(\zeta), \mathbb{F}_\ell)$.

Any $\mathbb{F}_\ell[G]$ -module M is the sum of its isotypical summands, the isotypical summand for $\mu_\ell^{\otimes i}$ being $\text{Hom}_G(\mu_\ell^{\otimes i}, M) \cong \text{Hom}_G(\mathbb{F}_\ell, \mu_\ell^{\otimes -i} \otimes M)$. In particular, the action of G on $X(\zeta)$ decomposes $H_{\text{et}}^*(X(\zeta), \mathbb{F}_\ell)$ into its isotypical pieces. Because the $\mu_\ell^{\otimes i}$ are the isotypical summands of $A = \pi_* \mathbb{F}_\ell$, the summand $H_{\text{et}}^*(X, \mu_\ell^{\otimes i})$ in (3.2) is the isotypical summand of $H_{\text{et}}^*(X(\zeta), \mathbb{F}_\ell)$ for $\mu_\ell^{\otimes i}$. Thus the injection

$$H_{\text{et}}^n(X, \mu_\ell^{\otimes i}) \rightarrow H_{\text{et}}^n(X(\zeta), \mu_\ell^{\otimes i}) \cong H_{\text{et}}^n(X(\zeta), \mathbb{F}_\ell)$$

sends u to $\zeta^{-i} \otimes \pi^*(u)$, where $\pi^*(u)$ is the image of u in $H_{\text{et}}^n(X(\zeta), \mu_\ell^{\otimes i})$.

Example 3.2.1. The product of $\zeta^{-1} \in H_{\text{et}}^0(k(\zeta), \mu_\ell^{\otimes -1})$ and $\beta(\zeta) \in H_{\text{et}}^1(k(\zeta), \mu_\ell)$ is an element $\zeta^{-1} \cup \beta(\zeta)$ of $H_{\text{et}}^1(k(\zeta), \mathbb{F}_\ell)$. As it is fixed by G , it descends to an element of $H_{\text{et}}^1(k, \mathbb{F}_\ell)$, which we will call z . Thus $\pi^*(z) = \zeta^{-1} \cup \beta(\zeta)$. As observed after (2.3.1), $z = 0$ iff $k(\zeta)$ contains primitive ℓ^2 -roots of unity.

Because the Frobenius is the identity on A , P_A^0 is the identity operation by [E, 8.3.4], and $P^0(u) = \zeta^{\otimes i(\ell-1)} \cup u$. We can now relate Epstein's operation Q^0 to the étale Bockstein β and relate his Q^a to βP^a .

Proposition 3.3. *Suppose that $\zeta_\ell \notin k$, and let $z \in H_{\text{et}}^1(k, \mathbb{F}_\ell)$ be the element of Example 3.2.1. Then, as cohomology operations on $H_{\text{et}}^n(X, \mu_\ell^{\otimes i})$ for $(0 \leq i < d)$:*

$$\begin{aligned} Q_A^0 &= \beta - i z \quad \text{and} \quad Q^0 = \beta P^0 = \zeta^{\otimes i(\ell-1)} \cup (\beta - i z); \\ Q_A^a &= \beta P_A^a - i z \cup P_A^a \quad \text{and} \quad Q^a = \beta P^a \quad \text{for } a > 0. \end{aligned}$$

The case when $\zeta_\ell \in k$ ($d = 1$) is handled by Proposition 2.4. Note that by our convention, the β in $Q^a = \beta P^a$ is the Bockstein on $H^*(X, \mu_\ell^{\otimes i\ell})$.

Proof. Given the isomorphism (3.2), Proposition 1.10 for Q_A^0 implies that the following diagram commutes:

$$\begin{array}{ccc} \bigoplus_{i=0}^{d-1} H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i}) & \xrightarrow{Q_A^0} & \bigoplus_{i=0}^{d-1} H_{\text{ét}}^{n+1}(X, \mu_{\ell}^{\otimes i\ell}) \\ \zeta^{-i} \cup \pi^* \downarrow \cong & & \zeta^{-i\ell} \cup \pi^* \downarrow \cong \\ H_{\text{ét}}^n(X(\zeta), \mathbb{F}_{\ell}) & \xrightarrow{Q^0 = \beta} & H_{\text{ét}}^{n+1}(X(\zeta), \mathbb{F}_{\ell}). \end{array}$$

For all $u \in H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i})$, $\beta(\pi^* u) = \pi^* \beta(u)$ by naturality. Since $\pi^*(z) = \zeta^{-1} \beta(\zeta)$, the diagram implies that $(\zeta^{-i\ell} \cup \pi^*)(Q_A^0 u)$ equals

$$\begin{aligned} \beta(\zeta^{-i} \cup \pi^* u) &= \zeta^{-i} \cup \beta(\pi^* u) + \beta(\zeta^{-i}) \cup \pi^*(u) \\ &= \zeta^{-i} \cup \pi^* \beta(u) - i \zeta^{-i-1} \beta(\zeta) \cup \pi^*(u) \\ &= (\zeta^{-i} \cup \pi^*)(\beta(u) - i z \cup u). \end{aligned}$$

Since $\zeta^{-i} \cup \pi^*$ is the isomorphism (3.2), the identity $Q_A^0(u) = \beta(u) - i z \cup u$ follows. By (3.1), we have $Q^0 u = \zeta^{i(\ell-1)} \cup Q_A^0 u$, which agrees with $\beta P^0 u = \beta(\zeta^{i(\ell-1)} \cup u)$.

Finally, the identity for Q_A^a and Q^a follows by invoking the Adem relations $Q_A^a P_A^0 = Q_A^0 P_A^a$, $Q^a P^0 = Q^0 P^a$, and $P^0 P^a = P^a P^0$ [E, 9.8(1,4)]:

$$Q_A^a = Q_A^a P_A^0 = Q_A^0 P_A^a = (\beta - i z) P_A^a \quad \text{and} \quad Q^a P^0 = Q^0 P^a = \beta P^0 P^a = \beta P^a P^0.$$

Now use the fact that P^0 is invertible. \square

Remark 3.3.1. If $i \equiv j \pmod{d}$, the étale sheaves $\mu_{\ell}^{\otimes i}$ and $\mu_{\ell}^{\otimes j}$ are isomorphic. This does not affect the operation Q_A^0 , but it changes the Bockstein because the isomorphism $H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i}) \cong H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes j})$ sends u to $v = \zeta^{j-i} \cup u$. Setting $j = dq + i$, Lemma 1.2 yields

$$\beta(v) = \zeta^{\otimes dq} \cup \beta(u) + dq \zeta^{\otimes dq-1} \beta(\zeta) \cup u = \zeta^{\otimes dq} \cup \{\beta(u) + (dq) z \cup u\}.$$

Using the Bockstein and Epstein's operations P^a , we have operations P^I defined on $H_{\text{ét}}^n(-, \mu_{\ell}^{\otimes i})$ for every admissible sequence I in the sense of Definition 0.1.

In order to classify all operations on $H_{\text{ét}}^n$, we first consider the case $n = 1$. In topology, the ring of operations on $H^1(-, \mathbb{F}_{\ell})$ is $H_{\text{top}}^*(K_1) \cong \mathbb{F}_{\ell}[u, v]/(u^2)$, where $u = P^0$ is in degree 1, corresponding to the identity operation, and v is in degree 2, corresponding to the Bockstein operation. By Theorem 1.3, there is a canonical map from $\mathbb{F}_{\ell}[u, v]/(u^2)$ to étale cohomology operations from $H_{\text{ét}}^1(-, \mu_{\ell}^{\otimes i})$ to $H_{\text{ét}}^*(-, \mu_{\ell}^{\otimes i})$, sending u to the identity, v to the Bockstein $\beta : H_{\text{ét}}^1(-, \mu_{\ell}^{\otimes i}) \rightarrow H_{\text{ét}}^2(-, \mu_{\ell}^{\otimes i})$, and v^m to $x \mapsto \beta(x)^m$.

For any i , the basechange $\mu_{\ell}^{\otimes i}(\zeta)$ of the algebraic group $\mu_{\ell}^{\otimes i}$ is isomorphic to $\mathbb{F}_{\ell}(\zeta)$, the constant sheaf \mathbb{F}_{ℓ} on the big étale site of $k(\zeta)$. The induced isomorphism $(B\mu_{\ell}^{\otimes i})(\zeta) \cong (B\mathbb{F}_{\ell})(\zeta)$ induces an isomorphism of cohomology groups, which immediately yields the following calculation.

Recall that $A = \bigoplus_{i=0}^{d-1} \mu_{\ell}^{\otimes i}$.

Proposition 3.4. *The graded algebra of cohomology operations from $H_{\text{ét}}^1(-, \mu_{\ell}^{\otimes i})$ to $\bigoplus_{j=0}^{d-1} H_{\text{ét}}^*(-, \mu_{\ell}^{\otimes j})$ is isomorphic to the $H_{\text{ét}}^*(k(\zeta), \mathbb{F}_{\ell})$ -module*

$$H^*(B\mu_{\ell}^{\otimes i}, A) \cong H^*(B\mu_{\ell}^{\otimes i}(\zeta), \mathbb{F}_{\ell}) \cong H^*(k(\zeta), \mathbb{F}_{\ell}) \otimes \mathbb{F}_{\ell}[u, v]/(u^2), \quad \beta(u) = v.$$

Every operation on $H_{\text{ét}}^1(-, \mu_{\ell}^{\otimes i})$ is uniquely a sum of operations $\phi(x) = cx^{\varepsilon} \beta(x)^m$, where $\varepsilon \in \{0, 1\}$, $m \geq 0$, and $c \in H_{\text{ét}}^(k, \mu_{\ell}^{\otimes j})$ for some j .*

The operations u and v on H_{et}^1 are of course $u(x) = x$ and $v(x) = \beta(x)$. Proposition 3.4 is the case $n = 1$ of the following result.

Theorem 3.5. *For each i and $n \geq 1$, the ring of all étale cohomology operations from $H_{\text{et}}^n(-, \mu_{\ell}^{\otimes i})$ to $H_{\text{et}}^*(-, A)$ is the free left $H_{\text{et}}^*(k(\zeta), \mathbb{F}_{\ell})$ -module $H_{\text{et}}^*(k(\zeta), \mathbb{F}_{\ell}) \otimes H_{\text{top}}^*(K_n)$.*

If $c \in H_{\text{et}}^(k, \mu_{\ell}^{\otimes j})$, the operation corresponding to the monomial $c P^{I_1} \cdots P^{I_r}$ sends $H_{\text{et}}^n(-, \mu_{\ell}^{\otimes i})$ to $H_{\text{et}}^*(-, \mu_{\ell}^{\otimes ri+j})$.*

Proof. We first show that the basechange $K(\mu_{\ell}^{\otimes i}, n) \times_k \text{Spec}(k(\zeta))$ is the space $K(\mu_{\ell}^{\otimes i}, n)$ over $k(\zeta)$. This is clear for $n = 0$ and follows inductively from the construction of $K(A, n+1)$ via the bar construction on $K(A, n)$, together with the observation that $(X \times_k Y) \times_k \text{Spec}(k(\zeta))$ is $X(\zeta) \times_{k(\zeta)} Y(\zeta)$.

By (3.2), the cohomology of $K(\mu_{\ell}^{\otimes i}, n)$ with coefficients in A is the same as the cohomology of $K(\mu_{\ell}^{\otimes i}, n) \times_k \text{Spec}(k(\zeta))$ with coefficients in \mathbb{F}_{ℓ} . The Breen-Jardine result, Theorem 2.1, shows that this is $H_{\text{et}}^*(k(\zeta), \mathbb{F}_{\ell}) \otimes H_{\text{top}}^*(K_n)$. \square

4. MAY'S ADJOINT CONSTRUCTION

A somewhat different approach to constructing cohomology operations was given by Peter May in [M]. Because we will need May's version of Kudo's Theorem (in Theorem 9.5 below), we need to know how the two constructions compare.

First, we need a chain level version of the Steenrod-Epstein function

$$m_* P : H^n(X, A) \rightarrow H_{\pi}^{n\ell}(X, A)$$

used in (1.8) to define P^a when A is a sheaf of commutative algebras. Fix an injective resolution $A \xrightarrow{\sim} I^*$ (in \mathcal{Sh}), an injective resolution $A^{\otimes \ell} \xrightarrow{\sim} I^{\otimes \ell} \xrightarrow{\sim} J^*$ in the category \mathcal{Sh}_{π} of π -equivariant sheaves, and an injective resolution $I^{\otimes \ell} \xrightarrow{\sim} J_S^*$ in the category $\mathcal{Sh}_{S_{\ell}}$ of S_{ℓ} -equivariant sheaves, as in Section 1.

The multiplication map $m : A^{\otimes \ell} \rightarrow A$ is equivariant for both the action of S_{ℓ} and its subgroup π on $A^{\otimes \ell}$. As we observed in the proof of Theorem 1.5, $A \rightarrow \text{Tot}(W^* \otimes I^*)$ is an injective resolution in \mathcal{Sh}_{π} . The comparison theorem lifts the resolution $A^{\otimes \ell} \rightarrow J^*$ to an equivariant map $J^* \rightarrow \text{Tot}(W^* \otimes I^*)$ over $A^{\otimes \ell} \rightarrow A$; taking sections over X yields a map $J^*(X) \rightarrow \text{Tot}(W^* \otimes I^*(X))$, natural in X . This induces an equivariant map of complexes of π -sheaves

$$m_{\pi} : I^{\otimes \ell} \xrightarrow{\sim} J^* \rightarrow \text{Tot}(W^* \otimes I^*).$$

Consider the isomorphism $\eta : W^* \otimes I^* \rightarrow \text{Hom}(W_*, I^*)$, defined on sections by

$$\eta_U(f \otimes x)(w) = (-1)^{|x||w|} f(w)x, \quad f \in W^*, w \in W_*, x \in I^*(U).$$

If $\{w_k \in W^k\}$ is the dual basis for $\{e_k \in W_k\}$ we have $\eta(w_j \otimes x)(e_k) = (-1)^{k|x|} \delta_{jk} x$. The composition ηm_{π} sends $I^{\otimes \ell}$ to $\text{Hom}(W_*, I^*)$. It is the (signed) adjoint

$$(4.1) \quad \theta : W_* \otimes I^{\otimes \ell} \rightarrow I^*$$

of the map $\eta m_{\pi}(X)$ which forms the basis for May's approach; see [M, 2.1]. In this approach, we fix a projective resolution of \mathbb{F}_{ℓ} as an S_{ℓ} -module, $V_* \rightarrow \mathbb{F}_{\ell}$, and a π -equivariant map $j : W_* \rightarrow V_*$ over \mathbb{F}_{ℓ} .

Similarly, suppose that \mathcal{K} is a sheaf of bounded below, homotopy associative dg algebras, and $\mathcal{K} \rightarrow I$ is an injective replacement (so that a lift $I \otimes I \rightarrow I$ gives I

the structure of a homotopy associative dg algebra). Given a π -equivariant map $m : \mathcal{K}^{\otimes \ell} \rightarrow \text{Hom}(W_*, \mathcal{K})$, the comparison theorem lifts

$$\mathcal{K}^{\otimes \ell} \xrightarrow{m} \text{Hom}(W_*, \mathcal{K}) \xrightarrow{\simeq} \text{Hom}(W_*, I)$$

to an equivariant map $m_\pi : I^{\otimes \ell} \rightarrow \text{Hom}(W_*, I)$, whose adjoint is again a map θ of the form given in (4.1).

Definition 4.2. Suppose that \mathcal{K} is a sheaf of homotopy associative dg \mathbb{F}_ℓ -algebras on some site and $\theta : W_* \otimes \mathcal{K}^{\otimes \ell} \rightarrow \mathcal{K}$ is a morphism of complexes in \mathcal{Sh}_π . We say that (\mathcal{K}, θ) is *suitable* if (i) the restriction of θ to $\mathcal{K}^{\otimes \ell} = \mathbb{F}_\ell\{e_0\} \otimes \mathcal{K}^{\otimes \ell}$ is chain homotopic to the iterated product $\mathcal{K}^{\otimes \ell} \rightarrow \mathcal{K}$ (in some order) and (ii) θ is chain homotopic to a composite

$$W_* \otimes \mathcal{K}^{\otimes \ell} \xrightarrow{j} V_* \otimes \mathcal{K}^{\otimes \ell} \xrightarrow{\phi} \mathcal{K},$$

where ϕ is the restriction to \mathcal{Sh}_π of a morphism of complexes in \mathcal{Sh}_{S_ℓ} .

A *morphism* of suitable pairs $(\mathcal{K}, \theta) \rightarrow (\mathcal{K}', \theta')$ is a morphism $f : \mathcal{K} \rightarrow \mathcal{K}'$ for which $f\theta$ is chain homotopic (over $\mathbb{F}_\ell[\pi]$) to $\theta'(1 \otimes f^{\otimes \ell})$. It is a *perfect* morphism if $f\theta = \theta'(1 \otimes f^{\otimes \ell})$. May writes $\mathcal{C}(\pi, \infty, \mathbb{F}_\ell)$ for the category of suitable pairs.

Taking sections over X , the pair $(\mathcal{K}(X), \theta_X)$ satisfies May's axioms in [M, 2.1], where θ_X is the induced map $W_* \otimes \mathcal{K}(X)^{\otimes \ell} \rightarrow W_* \otimes \mathcal{K}^{\otimes \ell}(X) \rightarrow \mathcal{K}(X)$. If f is a (perfect) morphism, then for any X , $f_X : \mathcal{K}(X) \rightarrow \mathcal{K}'(X)$ is a (perfect) morphism in the sense of [M, 2.1].

Remark 4.2.1. If Λ is any commutative ring, Definition 4.2 makes sense for any sheaf of homotopy associative dg Λ -algebras; we say that (K, θ) is *suitable for Λ* . Following [M, p. 161], we say that a suitable (K, θ) is *reduced* if it is obtained by reduction mod ℓ from a pair $(\tilde{K}, \tilde{\theta})$ which is suitable for \mathbb{Z}/ℓ^2 , such that \tilde{K} is a flat \mathbb{Z}/ℓ^2 -module. Since $0 \rightarrow K \rightarrow \tilde{K} \rightarrow K \rightarrow 0$ is an exact sequence of chain complexes, this data suffices to yield a Bockstein $\beta : H^N(K) \rightarrow H^{n+1}(K)$.

Example 4.3. Let \mathcal{C} be an acyclic operad of dg vector spaces over \mathbb{F}_ℓ . Then we may take $V_* = \mathcal{C}(\ell)$, since it is a resolution of \mathbb{F}_ℓ . If \mathcal{C} acts on \mathcal{K} , \mathcal{K} is homotopy associative, and θ is the composition of $j \otimes 1 : W_* \otimes \mathcal{K}^{\otimes \ell} \rightarrow \mathcal{C}(\ell) \otimes \mathcal{K}^{\otimes \ell}$ with the structure map $\mathcal{C}(\ell) \otimes \mathcal{K}^{\otimes \ell} \rightarrow \mathcal{K}$, then (\mathcal{K}, θ) is a suitable pair.

Definition 4.4 (May). Suppose that (\mathcal{K}, θ) is suitable, and set $K = \mathcal{K}(X)$. Define the function $D_k^M : K^n \rightarrow K^{n\ell-k}$ by the formula

$$D_k^M(u) = \theta(e_k \otimes u^{\otimes \ell}),$$

and define the Steenrod operations $P_M^a : H^n(K) \rightarrow H^{n+2a(\ell-1)}(K)$ (and Q_M^a) by

$$P_M^a(u) = (-1)^a \nu_n D_{(n-2a)(\ell-1)}^M(u) \quad \text{and} \quad Q_M^a(u) = (-1)^a \nu_n D_{(n-2a)(\ell-1)-1}^M(u)$$

(see [M, pp. 162, 182]; May's $\nu(-n)$ is our ν_n). As with Epstein's construction, $P_M^a = 0$ when $n < 2a$, and $Q_M^a = 0$ when $n \leq 2a$. May notes in [M, 2.3–2.5] that the D_k^M and hence the P_M^a, Q_M^a are additive and functorial for morphisms of suitable pairs.

These operations are natural in X , because for every morphism $f : Y \rightarrow X$ in the site, the restriction $f^* : \mathcal{K}(X) \rightarrow \mathcal{K}(Y)$ gives a perfect morphism $(\mathcal{K}(X), \theta_X) \rightarrow (\mathcal{K}(Y), \theta_Y)$ of objects in the sense of [M, 2.1].

A morphism of suitable pairs $(\mathcal{K}, \theta) \rightarrow (\mathcal{K}', \theta')$ induces a map $H^n \mathcal{K}(X) \rightarrow H^n \mathcal{K}'(X)$ compatible with the operations P_M^a on $H^n \mathcal{K}(X)$ and the corresponding operations P_M^a on $H^n \mathcal{K}'(X)$ in the sense that $P_M^a f^* = f^* P_M^a$. This follows from [M, 3.1(iii)].

Lemma 4.4.1. *If (K, θ) is reduced, then $Q_M^a = \beta P_M^a$, where β is the Bockstein.*

Proof. The proof of [M, 2.3(v)] applies; it suffices to show that $\beta D_{2i} = D_{2i-1}$. Given u in K^n with $d(u) = 0$, lift u to $\tilde{u} \in \tilde{K}^n$ and let b be such that $d(\tilde{u}) = \ell b$, so $\beta(u) = b$. Since $d(\tilde{u}^{\otimes \ell}) = N(\ell b \otimes \tilde{u}^{\otimes \ell-1})$, where $N = \sum \{\sigma \in \pi\}$ and $\theta\beta = \beta\theta$, May's calculation in $W_* \otimes \tilde{K}^{\otimes \ell}$ [M, p. 163] goes through to show that

$$\beta D_{2i}(u) = \theta\beta(e_{2i} \otimes u^{\otimes \ell}) \equiv \theta(e_{2i-1} \otimes u^{\otimes \ell}) = D_{2i-1}(u) \text{ modulo boundaries. } \square$$

Lemma 4.5. *If A is a sheaf of commutative dg \mathbb{F}_ℓ -algebras, $A \xrightarrow{\sim} I$ is an injective resolution, and θ is as in (4.1), the pair (I, θ) is suitable. Hence this data yields cohomology operations $P_M^a : H^n(X, A) \rightarrow H^{n+2a(\ell-1)}(X, A)$, natural in X .*

Proof. By Remark 1.6.1, I is a sheaf of homotopy associative dg algebras. By construction, the restriction of θ to $I^{\otimes \ell} = W_0 \otimes I^{\otimes \ell}$ is chain homotopic to the given map $I^{\otimes \ell} \rightarrow I$. To see that axiom 4.2(ii) is satisfied, set $V^* = \text{Hom}(V_*, \mathbb{F}_\ell)$ and note that, by the proof of Theorem 1.5, $A \rightarrow \text{Tot}(V^* \otimes I)$ is an injective resolution in $\mathcal{S}h_{S_\ell}$. The construction before Definition 1.7 yields a map $m_S : I^{\otimes \ell} \xrightarrow{\sim} J_S^* \rightarrow V^* \otimes I$ whose adjoint $V_* \otimes I^{\otimes \ell} \rightarrow I$ is ϕ . The comparison theorem for $\mathcal{S}h_\pi$ provides a lift $J_S \rightarrow J$ over $I^{\otimes \ell}$ such that the map

$$I^{\otimes \ell} \xrightarrow{m_S} \text{Hom}(V_*, I) \xrightarrow{j^*} \text{Hom}(W_*, I)$$

is chain homotopic over $\mathbb{F}_\ell[\pi]$ to the map m_π , as required. The final assertion follows because $H^*(X, A) = H^* I(X)$. \square

In fact, the operations P_M^a are independent of the choice of resolution $A \xrightarrow{\sim} I$; this follows from the following lemma, whose proof is the same as the proof of Lemma 4.5, with A replaced by \mathcal{K} . (See the discussion before Definition 4.2.)

Lemma 4.5.1. *Let (\mathcal{K}, θ_0) be a suitable pair, where \mathcal{K} is a sheaf of bounded below, homotopy associative dg \mathbb{F}_ℓ -algebras. If $\eta : \mathcal{K} \xrightarrow{\sim} I$ is an injective replacement and θ is the map of (4.1), then the pair (I, θ) is suitable, and η is a perfect morphism $(\mathcal{K}, \theta_0) \rightarrow (I, \theta)$ of suitable pairs.*

The sign differences in the formulas for P^a and P_M^a (and for Q^a and Q_M^a) are explained by the following calculation.

Proposition 4.6. *For u in $H^n(X, A)$, $D_k^M = (-1)^k D_k$.*

Proof. Consider the isomorphism $\phi : W^* \otimes I^*(X) \rightarrow \text{Hom}(W_*, I^*(X))$, defined above. The adjoint θ of ϕm_π is the composite

$$W_* \otimes I(X)^{\otimes \ell} \cong I(X)^{\otimes \ell} \otimes W_* \xrightarrow{\phi m_\pi \otimes 1} \text{Hom}(W_*, I(X)) \otimes W_* \xrightarrow{\eta} I(X),$$

where the first map is the signed symmetry isomorphism and η is evaluation. We now compute that

$$\begin{aligned}
D_k^M(u) &= \theta(e_k \otimes u^{\otimes \ell}) = (-1)^{kn\ell} \eta(\phi[m_\pi(u^{\otimes \ell})] \otimes e_k) \\
&= (-1)^{kn\ell} \eta\left(\phi\left[\sum w_j \otimes D_j(u)\right] \otimes e_k\right) \\
&= (-1)^{kn\ell} \sum_j \phi[w_j \otimes D_j(u)](e_k) \\
&= (-1)^{kn\ell} (-1)^{k(n\ell-k)} D_k(u) = (-1)^k D_k(u). \quad \square
\end{aligned}$$

Recall that Epstein's operations P^a and Q^a are defined in (1.8) and Remark 1.8.1.

Corollary 4.7. *May's operations P_M^a and Q_M^a coincide with Epstein's P^a and Q^a .*

Lemma 4.8. *Set $m = (\ell - 1)/2$. Then for each $u \in I^n(X)$:*

- (i) $dP^a(u) = P^a(du)$ and $dQ^a(u) = -Q^a(du)$; and
- (ii) if u is a cocycle representing $x \in H^n(X, A)$, then $P^a(u)$ and $Q^a(u)$ are cocycles representing $P^a(x)$ and $Q^a(x)$, respectively.

Proof. In Theorem 3.1 of [M], May shows that (i) $dP_M^a(u) = P_M^a(du)$ and $dQ_M^a(u) = -Q_M^a(du)$, and (ii) if u is a cocycle representing $x \in H^n(X, A)$, then $P_M^a(u)$ and $Q_M^a(u)$ are cocycles representing $P_M^a(x)$ and $Q_M^a(x)$. The result is immediate from Corollary 4.7. \square

Let $G \cong \pi \wr \pi$ denote a Sylow ℓ -subgroup of S_{ℓ^2} . Then there is a free $\mathbb{F}_\ell[G]$ -module resolution $W_* \otimes W_*^{\otimes \ell} \rightarrow \mathbb{F}_\ell$. If $'V_* \rightarrow \mathbb{F}_\ell$ is a projective resolution of \mathbb{F}_ℓ as an S_{ℓ^2} -module, then there is a G -module morphism $\sigma : W_* \otimes W_*^{\otimes \ell} \rightarrow 'V_*$ over \mathbb{F}_ℓ .

Definition 4.9. Following [M, 4.1], we say that a suitable (\mathcal{K}, θ) is an *Adem object* if there is an S_{ℓ^2} -equivariant morphism $'V_* \otimes \mathcal{K}^{\otimes \ell^2} \rightarrow \mathcal{K}$ whose composition with $\sigma \otimes 1$ is G -homotopic to

$$(W_* \otimes W_*^{\otimes \ell}) \otimes \mathcal{K}^{\otimes \ell^2} \cong W_* \otimes (W_* \otimes \mathcal{K}^{\otimes \ell})^{\otimes \ell} \xrightarrow{\theta^{\otimes \ell}} W_* \otimes \mathcal{K}^{\otimes \ell} \xrightarrow{\theta} \mathcal{K}.$$

May proves in [M, 4.7] that the P^a satisfy the Adem relations whenever (\mathcal{K}, θ) is an Adem object.

Example 4.9.1. Given $A \xrightarrow{\sim} I$, $A^{\otimes \ell} \rightarrow \text{Tot}('V^* \otimes I)$ is an injective resolution in $\mathcal{Sh}_{S_{\ell^2}}$, and we get an S_{ℓ^2} -equivariant map $I^{\otimes \ell^2} \rightarrow \text{Hom}('V_*, I)$ whose adjoint $'V_* \otimes I^{\otimes \ell^2} \rightarrow I$ makes (I, θ) into an Adem object. We omit the routine details.

Similarly, if (\mathcal{K}, θ_0) is an Adem object, $\mathcal{K} \rightarrow I$ an injective replacement, and (I, θ) is as in Lemma 4.5.1, then the map $'V_* \otimes \mathcal{K}^{\otimes \ell^2} \rightarrow \mathcal{K} \rightarrow I$ lifts to an S_{ℓ^2} -equivariant map $'V_* \otimes I^{\otimes \ell^2} \rightarrow I$; this map makes (I, θ) into an Adem object. Again, we omit the routine details.

Definition 4.10. If $(\mathcal{K}_1, \theta_1)$ and $(\mathcal{K}_2, \theta_2)$ are suitable pairs, the tensor product $(\mathcal{K}_1 \otimes \mathcal{K}_2, \theta_1 \otimes \theta_2)$ is also a suitable pair, where $\theta_1 \otimes \theta_2$ is described in [M, 2.1].

A suitable pair (\mathcal{K}, θ) is a *Cartan object* if the product $(\mathcal{K} \otimes \mathcal{K}, \theta \otimes \theta) \rightarrow (\mathcal{K}, \theta)$ is a morphism of suitable pairs.

Example 4.10.1. If A is a sheaf of commutative algebras, then so is $A \otimes A$, and the product $A \otimes A \rightarrow A$ makes A into a Cartan object. If $A \xrightarrow{\sim} I$ is an injective

resolution, then the pair (I, θ) of Lemma 4.5 is also a Cartan object, because the two morphisms $W_* \otimes (I \otimes I)^{\otimes \ell} \rightarrow I$ are isomorphic to the two morphisms $W_* \otimes (A \otimes A)^{\otimes \ell} \rightarrow A$ in the derived category and (because I is a bounded below complex of injectives) therefore chain homotopic.

For exactly the same reasons, if (\mathcal{K}, θ_0) is a Cartan object and $\mathcal{K} \rightarrow I$ is an injective replacement, then so is (I, θ) .

5. OPERADS AND OPERATIONS

In [M1], May gave a different approach to power operations in sheaf cohomology. In this section, we give a short discussion of this approach.

Let \mathcal{Z} denote the *Eilenberg-Zilber operad* in the category of \mathbb{F}_ℓ -modules, defined by Hinich and Schechtman in [HS]. By an action of \mathcal{Z} on a cochain complex of sheaves C we mean a collection of sheaf morphisms

$$\mathcal{Z}(n) \otimes C^{\otimes n} \longrightarrow C$$

satisfying appropriate equivariance, associativity, and unit axioms. The choice of an element $m \in \mathcal{Z}(2)$ determines a product $C^{\otimes 2} \rightarrow C$ and makes C into a homotopy associative dg algebra. Because each $\mathcal{Z}(n)$ is an acyclic complex of $\mathbb{F}_\ell[S_n]$ -modules, we have quasi-isomorphisms $W_* \xrightarrow{\sim} \mathcal{Z}(\ell)$ and $'V_* \xrightarrow{\sim} \mathcal{Z}(\ell^2)$. We thus have a natural map

$$\theta : W_* \otimes C^{\otimes \ell} \xrightarrow{\sim} \mathcal{Z}(\ell) \otimes C^{\otimes n} \longrightarrow C.$$

Lemma 5.1. *If \mathcal{Z} acts on a cochain complex C , then (C, θ) is a suitable pair in the sense of Definition 4.2. It is also an Adem and a Cartan object (Definitions 4.9 and 4.10).*

Proof. This is an exercise in the axioms of operads, left to the reader. The axiom about $\mathcal{Z}(\ell) \otimes \mathcal{Z}(\ell)^{\otimes \ell} \rightarrow \mathcal{Z}(\ell^2)$ is used to show (C, θ) is an Adem object, and the axiom about $\mathcal{Z}(2) \otimes \mathcal{Z}(\ell)^{\otimes 2} \rightarrow \mathcal{Z}(2\ell)$ is used to show it is a Cartan object. \square

Example 5.2. Let $F^\bullet \mathcal{F}$ denote the Godement resolution of a sheaf \mathcal{F} . Since every skyscraper sheaf of \mathbb{F}_ℓ -modules is an injective sheaf, this is an injective resolution of \mathcal{F} . If A is any sheaf of commutative \mathbb{F}_ℓ -algebras, the results of Section 4 apply. Alternatively, Hinich and Schechtman showed in [HS] that \mathcal{Z} acts on $F^\bullet A$, so $(F^\bullet A, \theta)$ is not only suitable but is both an Adem and a Cartan object by the lemma above.

By Lemma 4.5, this data provides cohomology operations P^a on $H^*(X, A)$, natural in X . Of course, this construction is little more than a reinterpretation of May's procedure (in Theorem 4.8 of [M1]), where he shows that the Eilenberg-Zilber operad acts on the sections $F^\bullet A(X)$, giving cohomology operations.

Remark 5.3. If A is a sheaf of étale algebras, the direct image $\alpha_*(F^\bullet A)$ is also a complex of injective Nisnevich sheaves. However, although the operad \mathcal{Z} acts on the good truncations $\tau^{\leq i} \alpha_*(F^\bullet A)$, it only acts up to homotopy on an injective replacement, such as the complex I_{nis} of Corollary 6.4 below. Thus one needs to take care when using this approach to construct motivic cohomology operations in the next section.

6. MOTIVIC STEENROD OPERATIONS

In this section we construct operations P^a on the motivic cohomology groups $H^{n,i}(X) = H^{n,i}(X, \mathbb{F}_\ell)$, $n \geq 2a$, compatible with the operations P^a in étale cohomology (defined in Theorem 1.3) in the sense that there are commutative diagrams

$$(6.1) \quad \begin{array}{ccc} H^{n,i}(X, \mathbb{F}_\ell) & \xrightarrow{P^a} & H^{n+2a(\ell-1), i\ell}(X, \mathbb{F}_\ell) \\ \downarrow & & \downarrow \\ H_{\text{et}}^n(X, \mu_\ell^{\otimes i}) & \xrightarrow{P^a} & H_{\text{et}}^{n+2a(\ell-1)}(X, \mu_\ell^{\otimes i\ell}). \end{array}$$

Let α_* denote the direct image functor from the étale site to the Nisnevich site. If A is any étale sheaf, then we may regard $R\alpha_* A$ as a complex of Nisnevich sheaves such that $H_{\text{nis}}^*(X, R\alpha_* A) \cong H_{\text{et}}^*(X, A)$. If $A \rightarrow I$ is an injective resolution, then $\alpha_* I$ is a complex of injective Nisnevich sheaves representing $R\alpha_* A$.

Let $\tau^{\leq i} A$ denote the good truncation of A in cohomological degrees at most i ; $H^n(\tau^{\leq i} A)$ is $H^n(A)$ for $i \leq n$ and zero for $n > i$ (cf. [WH, 1.2.7]). If $B^n = 0$ for $n < 0$, there is a natural transformation $(\tau^{\leq i} A) \otimes B \rightarrow \tau^{\leq i}(A \otimes B)$.

The following theorem, due to Voevodsky and Rost, is sometimes known as the Beilinson Conjecture; it is equivalent to the Norm Residue Theorem; see [SV], [W], [V4, 6.17], or [HW, Thm. C].

Norm Residue Theorem 6.2. *For any field of characteristic $\neq \ell$, the canonical map $\mathbb{F}_\ell(i) \rightarrow \tau^{\leq i} R\alpha_* \mu_\ell^{\otimes i}$ is a quasi-isomorphism of complexes of Nisnevich sheaves on the category of smooth simplicial schemes. Hence for any X we have*

$$H^{n,i}(X, \mathbb{F}_\ell) \cong H_{\text{nis}}^n(X, \tau^{\leq i} R\alpha_* \mu_\ell^{\otimes i}).$$

In particular, if $n \leq i$, then $H^{n,i}(X, \mathbb{F}_\ell) \cong H_{\text{et}}^n(X, \mu_\ell^{\otimes i})$.

We apply the above constructions to the étale sheaf of dg algebras $\mathcal{A} = \bigoplus_{i=0}^\infty \mu_\ell^{\otimes i}$. Choose injective resolutions $\mu_\ell^{\otimes i} \rightarrow I(i)$ and write I for the complex of étale sheaves $\bigoplus_{i=0}^\infty I(i)$. Then $\mathcal{A} \xrightarrow{\sim} I$ is an injective resolution; by Remark 1.6.1, I is an étale sheaf of homotopy-associative graded dg algebras, and (I, θ) is suitable by Lemma 4.5.

Now consider the Nisnevich sheaf of dg algebras $\alpha_* I$. Using the natural transformation $(\alpha_* I)^{\otimes \ell} \rightarrow \alpha_*(I^{\otimes \ell})$, we obtain a map $\alpha_* \theta : W_* \otimes (\alpha_* I)^{\otimes \ell} \rightarrow \alpha_* I$ such that $(\alpha_* I, \alpha_* \theta)$ is suitable in the sense of Definition 4.2. It is also clear that $(\alpha_* I(X), \alpha_* \theta_X)$ is both an Adem object and a Cartan object, natural in X (see Examples 4.9.1 and 4.10.1).

Of course, nothing new has happened; for each X we have $\alpha_* I(X) = I(X)$ and $(\alpha_* \theta)_X = \theta_X$ by the definition of direct image, so the resulting cohomology operations on $H^* \alpha_* I(X) = H_{\text{et}}^*(X, \mu_\ell^{\otimes *})$ are the same as those constructed in Definition 4.4 and hence (by Corollary 4.7) are the same as those in Theorem 1.3.

We now consider the effect of truncation. If $\mathcal{K} = \bigoplus_{i=0}^\infty \mathcal{K}_i$ is a sheaf of (homotopy associative) graded dg algebras, set $\tau \mathcal{K} = \bigoplus \tau^{\leq i} \mathcal{K}_i$. The products

$$\tau^{\leq i_1} \mathcal{K}_{i_1} \otimes \tau^{\leq i_2} \mathcal{K}_{i_2} \rightarrow \tau^{\leq i_1 + i_2} (\mathcal{K}_{i_1} \otimes \mathcal{K}_{i_2}) \rightarrow \tau^{\leq i_1 + i_2} \mathcal{K}_{i_1 + i_2}$$

make $\tau \mathcal{K}$ a sheaf of (homotopy associative) graded dg algebras. If (\mathcal{K}, θ) is suitable (Definition 4.2) and $\sum i_r = n$, the component maps

$$W_* \otimes \bigotimes_{r=1}^{\ell} (\tau^{\leq i_r} \mathcal{K}_{i_r}) \rightarrow \tau^{\leq n} (W_* \otimes \bigotimes \mathcal{K}_{i_r}) \xrightarrow{\theta} \tau^{\leq n} \mathcal{K}_n$$

assemble to define a map $W_* \otimes (\tau\mathcal{K})^{\otimes \ell} \rightarrow \tau\mathcal{K}$ we shall call $\tau\theta$.

Lemma 6.3. *If (\mathcal{K}, θ) is suitable, then so is $(\tau\mathcal{K}, \tau\theta)$, and $\tau\mathcal{K} \rightarrow \mathcal{K}$ defines a perfect morphism of suitable pairs.*

If (\mathcal{K}, θ) is an Adem object (resp., a Cartan object), so is $(\tau\mathcal{K}, \tau\theta)$.

Proof. Axiom (i) is trivial, and axiom (ii) follows from the commutative diagram

$$\begin{array}{ccc} W_* \otimes \bigotimes_{r=1}^{\ell} (\tau^{\leq i_r} \mathcal{K}_{i_r}) & \rightarrow & \tau^{\leq j} (W_* \otimes \bigotimes \mathcal{K}_{i_r}) \\ \downarrow j & & \downarrow j \\ V_* \otimes \bigotimes_{r=1}^{\ell} (\tau^{\leq i_r} \mathcal{K}_{i_r}) & \rightarrow & \tau^{\leq j} (V_* \otimes \bigotimes \mathcal{K}_{i_r}) \xrightarrow{\phi} \tau^{\leq j} \mathcal{K}_j. \end{array}$$

The final two assertions are easily verified using similar diagrams. \square

Example 6.3.1. When $\mathcal{K} = \alpha_* I = \bigoplus \alpha_* I(i)$, we see that $(\tau \alpha_* I, \tau\theta)$ is suitable, where $\tau \alpha_* I = \bigoplus \tau^{\leq i} I(i)$. Since $(\alpha_* I, \theta)$ is both an Adem object and a Cartan object, so is $(\tau \alpha_* I, \tau\theta)$.

Because each $\alpha_* I(i)^n$ is an injective Nisnevich sheaf and the sheaf $Z^i(\alpha_* I(i))$ of i -cycles has an injective resolution starting with $\alpha_* I(i)^i$, $\tau^{\leq i} \alpha_* I(i)$ is quasi-isomorphic to a chain complex $I(i)_{\text{nis}}^*$ of injective Nisnevich sheaves on X with $I(i)_{\text{nis}}^n = \alpha_* I(i)^n$ for $n \leq i$, and $I(i)_{\text{nis}}$ represents $\tau^{\leq i} R\alpha_* \mu_{\ell}^{\otimes i}$. By Theorem 6.2, $H^{n,i}(X, \mathbb{F}_{\ell})$ is the n th cohomology of the cochain complex $I(i)_{\text{nis}}^*(X)$.

By Remark 1.6.1, $I_{\text{nis}} = \bigoplus I(i)_{\text{nis}}$ is a Nisnevich sheaf of homotopy-associative dg algebras, and the products $I(i)_{\text{nis}} \otimes I(j)_{\text{nis}} \rightarrow I(i+j)_{\text{nis}}$, representing the pairings

$$\tau^{\leq i} R\alpha_* \mu_{\ell}^{\otimes i} \otimes \tau^{\leq j} R\alpha_* \mu_{\ell}^{\otimes j} \rightarrow \tau^{\leq i+j} R\alpha_* \mu_{\ell}^{\otimes i+j},$$

induce the product in motivic cohomology by [SV, 7.1].

Corollary 6.4. *The pair $(I_{\text{nis}}, \theta')$ is suitable and is both an Adem and a Cartan object. In addition, $\tau \alpha_* I \rightarrow I_{\text{nis}}$ defines a perfect morphism $(\tau \alpha_* I, \tau\theta) \rightarrow (I_{\text{nis}}, \theta')$.*

Proof. By Example 6.3.1, $(\tau \alpha_* I, \tau\theta)$ is suitable and both an Adem and a Cartan object. By Lemma 4.5.1, $(I_{\text{nis}}, \theta')$ is suitable, and η is a perfect morphism. As observed in Examples 4.9.1 and 4.10.1, $(I_{\text{nis}}, \theta')$ is both an Adem and a Cartan object. \square

Remark 6.4.1. By the comparison theorem, $\tau \alpha_* I \rightarrow \alpha_* I$ lifts to a morphism $f : I_{\text{nis}} \rightarrow \alpha_* I$. We will see in Lemma 6.6 that f defines a morphism of suitable pairs $(I_{\text{nis}}, \theta') \rightarrow (\alpha_* I, \alpha_* \theta)$.

Recall that $H^{n,i}(X, \mathbb{F}_{\ell}) = H_{\text{nis}}^n(X, \mathbb{F}_{\ell}(i))$. By the Norm Residue Theorem 6.2,

$$\bigoplus_i H^{n,i}(X, \mathbb{F}_{\ell}) = \bigoplus_i H_{\text{nis}}^n(X, I(i)_{\text{nis}}) = H^n I_{\text{nis}}(X).$$

Definition 4.4, applied to $(I_{\text{nis}}, \theta')$, yields operations P^a and Q^a on the cohomology of $I_{\text{nis}}(X)$, sending $\bigoplus_i H^{n,i}(X)$ to $\bigoplus_i H^{N,i\ell}(X)$ and $\bigoplus_i H^{N+1,i\ell}(X)$. By construction, these operations multiply the weight i by ℓ . We shall now formalize this observation.

Definition 6.5 (P^a). The motivic cohomology operations are the operations

$$P^a : H^{n,i}(X) \rightarrow H^{N,i\ell}(X) \quad \text{and} \quad Q^a : H^{n,i}(X) \rightarrow H^{N+1,i\ell}(X),$$

$N = n + 2a(\ell - 1)$, defined by the suitable pair $(I_{\text{nis}}, \theta')$ of Corollary 6.4, using the identification of $H^{n,i}(X)$ with $H^n I_{\text{nis}}(X)$. If $n < 2a$, we define $P^a = 0$.

If $\ell = 2$, we write Sq^{2a} for P^a and Sq^{2a+1} for βP^a , so that Sq^a takes $H^{n,i}(X)$ to $H^{n+a,2i}(X)$. If $n < a$, then Sq^a is zero on $H^{n,i}(X)$.

By Lemma 4.5, these operations are natural in X .

Remark 6.5.1. These motivic cohomology operations are almost surely the operations defined by Kriz and May in [KM, I.7.2] and by Joshua in [Jo, §8]; compare with [BJ].

Lemma 6.6. *The motivic cohomology operations P^a and Q^a are compatible with the étale cohomology operations P^a and Q^a in the sense that the diagram (6.1) commutes.*

Proof. We need to show that $f\theta'$ is chain homotopic to $(\alpha_*\theta)(1 \otimes f^{\otimes \ell})$. Consider the following diagram of complexes in $\mathcal{S}h_\pi$.

$$\begin{array}{ccccc} W_* \otimes (\tau\alpha_* I)^{\otimes \ell} & \xrightarrow{1 \otimes \epsilon^{\otimes \ell}} & W_* \otimes I_{\text{nis}}^{\otimes \ell} & \xrightarrow{1 \otimes f^{\otimes \ell}} & W_* \otimes (\alpha_* I)^{\otimes \ell} \\ \downarrow \tau\alpha_* \theta & & \downarrow \theta' & & \downarrow \alpha_* \theta \\ \tau\alpha_* I & \xrightarrow{\epsilon} & I_{\text{nis}} & \xrightarrow{f} & \alpha_* I \end{array}$$

By Lemma 6.3, the bottom composite $\tau\alpha_* I \rightarrow \alpha_* I$ defines a perfect morphism of suitable pairs, meaning that the outer square commutes (see Definition 4.2). By Corollary 6.4, $\epsilon : \tau\alpha_* I \xrightarrow{\sim} I_{\text{nis}}$ defines a perfect morphism of suitable pairs, meaning that the left square commutes. Because ϵ is a quasi-isomorphism, so is $1 \otimes \epsilon^{\otimes \ell}$, by Lemma 1.6. It follows that the right square commutes in the derived category. Because $\alpha_* I$ is a bounded below complex of injectives, this implies that the right square commutes up to chain homotopy equivalence; see [WH, 10.4.7]. \square

Remark 6.6.1. It is an easy exercise to show that the motivic and étale Bockstein operations are compatible, using $\tau^{\leq i} \mu_{\ell^2}^{\otimes i}$. We omit the details.

7. MOTIVIC FORMULAS

We now show that the motivic cohomology operations P^a of Definition 6.5 enjoy familiar properties.

Proposition 7.1. *If $u \in H^{2n,i}(X)$, then $P^n(u) = u^\ell$.*

Proof. This is [M, 2.4]. \square

We now turn to the Adem relations. Recall that by convention $\binom{n}{k}$ is zero if $k < 0$. Thus the sums below run over $t \leq a/\ell$.

Theorem 7.2 (Adem relations). *If $\ell > 2$ and $a < b\ell$, then*

$$\begin{aligned} P^a P^b &= \sum_{s+t=b} (-1)^{a+t} \binom{(\ell-1)s-1}{a-t\ell} P^{a+s} P^t, \\ P^a \beta P^b &= \sum_{s+t=b} (-1)^{a+t} \binom{(\ell-1)s}{a-t\ell} \beta P^{a+s} P^t + (-1)^{a+t} \binom{(\ell-1)s-1}{a-t\ell-1} \beta P^{a+s} \beta P^t. \end{aligned}$$

If $\ell = 2$ and $a < 2b$, then

$$Sq^a Sq^b = \sum_{s+t=b} \binom{s-1}{a-2t} Sq^{a+s} Sq^t.$$

Proof. By Corollary 6.4, $(I_{\text{nis}}, \theta')$ is an Adem object. As noted in [M, 4.7] (see Definition 4.9), this implies that the P^a and Q^a satisfy the Adem relations. We have replaced Q^a by βP^a (using Theorem 8.11) for the sake of familiarity. \square

Bistable Operations 7.3. In [V1], Voevodsky defines (bistable) cohomology operations P_V^a on $H^{n,i}(-, \mathbb{F}_\ell)$ of bidegree $(2a(\ell-1), a(\ell-1))$. These satisfy $P_V^0 x = x$ for all x , $P_V^a x = x^\ell$ for $x \in H^{2a,a}(X, \mathbb{F}_\ell)$, and $P_V^a = 0$ on $H^{n,i}$ if $i \leq a$ and $n < i+a$; the usual Adem relations hold when $\ell > 2$. The cohomological degrees of P^a and P_V^a are the same, namely $2a(\ell-1)$, but the weights differ if $a \neq i$: if $a < i$, then P_V^a has lower weight, but if $a > i$, then P^a has lower weight.

When $\ell = 2$, Voevodsky's operations Sq_V^{2i} have bidegree $(2i, i)$ and $Sq_V^{2i+1} = \beta Sq_V^{2i}$. They satisfy a modified Cartan formula [V1, 9.7] which differs from our Cartan formula (Theorem 7.4) by the presence of a factor of $[\zeta]$ in some terms.

Remark 7.3.1. Brosnan and Joshua have observed in [BJ, 2.1] and [BJ1, 1.1(iii)] that the motivic-to-étale map sends P_V^a to P^a and Sq_V^a to Sq^a . The key is to observe that Voevodsky's total power operation [V1, 5.3] is compatible with Epstein's reduced power map (Definition 1.7 above).

Theorem 7.4 (Cartan formula). *Let $u \in H^{n,i}(X)$ and $v \in H^{m,j}(Y)$. Then in $H^{*,(i+j)\ell}(X \times Y)$ we have*

$$P^a(u \cup v) = \sum_{s+t=a} P^s(u) \cup P^t(v), \quad \ell > 2,$$

and $Sq^a(u \cup v) = \sum_{s+t=a} Sq^s(u) \cup Sq^t(v)$ when $\ell = 2$. There is a similar formula for $Q^a(u \cup v)$.

Proof. By Corollary 6.4, $(I_{\text{nis}}, \theta')$ is a Cartan object (Definition 4.10). The formula now follows from [M, 2.7] (for both P^a and Q^a). \square

Cohomology operations on $H^{n,0}$ are easy to describe because of the following characterization. Recall that $\pi_0 X$ denotes the set of connected components of a scheme X ; if X_\bullet is a simplicial scheme, $\pi_0 X_\bullet$ denotes the simplicial set $n \mapsto \pi_0(X_n)$.

Lemma 7.5. *Let A be any abelian group. If X_\bullet is a smooth simplicial scheme, the motivic cohomology ring $H^{*,0}(X_\bullet, A)$ is isomorphic to the topological cohomology ring $H_{\text{top}}^*(\pi_0 X_\bullet, A)$ of the simplicial set $\pi_0 X_\bullet$.*

Proof. For smooth connected X we have $H^{n,0}(X, A) = H_{\text{nis}}^n(X, A) = 0$ for $n > 0$ and $H^{0,0}(X, A) = A$, almost by definition; see [MVW, 3.4]. Hence the spectral sequence $E_1^{p,q} = H^q(X_p, A) \Rightarrow H^{p+q,0}(X)$ degenerates to the cohomology of the chain complex $\text{Hom}(\pi_0 X_\bullet, A)$, which is $H_{\text{top}}^*(\pi_0 X_\bullet, A)$. For a simplicial set K such as $\pi_0 X_\bullet$, the construction of the product in motivic cohomology [MVW, 3.11] shows that $H_{\text{top}}^*(K) \cong H^{*,0}(K)$ is an isomorphism of rings. \square

Any simplicial set $X = X_\bullet$ may be regarded as a discrete simplicial scheme; in degree n it is the disjoint union of copies of $\text{Spec}(k)$, indexed by X_n .

Theorem 7.6. *For any simplicial set X , there is a natural isomorphism*

$$H^{*,*}(k, \mathbb{F}_\ell) \otimes H_{\text{top}}^*(X, \mathbb{F}_\ell) \cong H^{*,*}(k, \mathbb{F}_\ell) \otimes H^{*,0}(X, \mathbb{F}_\ell) \xrightarrow{\cup} H^{*,*}(X, \mathbb{F}_\ell).$$

Proof. By Lemma 7.5, $H_{\text{top}}^*(X, \mathbb{F}_\ell) \cong H^{*,0}(X, \mathbb{F}_\ell)$. Thus the map exists and is natural in X by the above remarks. It is an isomorphism for spheres by [V1, (2.7)]. If $X^{(i)}$ denotes the i -skeleton of X , the cone C_i of $X^{(i-1)} \rightarrow X^{(i)}$ is a bouquet of i -spheres, and the map is compatible with the exact sequence

$$\cdots \tilde{H}^n(C_i, \mathbb{F}_\ell) \rightarrow H^n(X^{(i)}, \mathbb{F}_\ell) \rightarrow H^n(X^{(i-1)}, \mathbb{F}_\ell) \cdots$$

of [V2, Lemma 8.2]. The result now follows by induction on i . \square

Applying this to the classifying space K_n for simplicial cohomology, we obtain:

Corollary 7.7. *The ring of motivic cohomology operations on $H^{n,0}(-, \mathbb{F}_\ell)$ is isomorphic to $H^{*,*}(k, \mathbb{F}_\ell) \otimes H_{\text{top}}^*(K_n) \cong H^{*,*}(K_n, \mathbb{F}_\ell)$ as a free left $H^{*,*}(k, \mathbb{F}_\ell)$ -module.*

If K is a simplicial set, the isomorphism $H^{*,0}(K, \mathbb{F}_\ell) \cong H_{\text{top}}^*(K, \mathbb{F}_\ell)$ is compatible with the action of the P^I . This is clear from Lemma 6.6 and Example 2.2.

Example 7.7.1. Let Δ^1 denote the simplicial 1-simplex and let $s \in H^{1,0}(\Delta^1, \partial\Delta^1)$ be the generator. By the above comparison with topology, $P^0(s) = s$. By definition, $P^a(s) = 0$ for $a > 0$.

Recall that the simplicial suspension SX of a pointed simplicial scheme X is again a simplicial scheme. Multiplication by the element s of Example 7.7.1 induces a canonical isomorphism $H^{n,i}(X, \mathbb{F}_\ell) \xrightarrow{\cong} H^{n+1,i}(SX, \mathbb{F}_\ell)$. (Compare to Lemmas 1.2 and 2.1 of [SE].)

Proposition 7.8. *The motivic operations P^a and Q^a are simplicially stable in the sense that they commute with simplicial suspension: there are commutative diagrams for all X , n , and i , with $N = n + 2a(\ell - 1)$, the diagram for P^a being*

$$\begin{array}{ccc} H^{n,i}(X) & \xrightarrow{P^a} & H^{N,i\ell}(X) \\ \cong \downarrow & & \cong \downarrow \\ H^{n+1,i}(SX) & \xrightarrow{P^a} & H^{N+1,i\ell}(SX). \end{array}$$

Proof. By the Cartan formula Theorem 7.4, $P^a(sx) = P^0(s)P^a(x) = s \cdot P^a(x)$. \square

Recall from [V1] that although each $H^{n,i}(-, \mathbb{F}_\ell)$ is defined as a contravariant functor on the category of smooth simplicial schemes, it is homotopy invariant and factors through the pointed motivic homotopy category Ho_\bullet . It is an elementary observation that any natural transformation between homotopy invariant functors, defined on the category of smooth simplicial schemes, must factor through Ho_\bullet . In particular, cohomology operations $H^{n,i}(-, \mathbb{F}_\ell) \rightarrow H^{p,q}(-, \mathbb{F}_\ell)$ may be regarded as natural transformations between functors defined on Ho_\bullet .

Example 7.9. The classifying space $K = K(\mathbb{F}_\ell(i), n)$ for $H^{n,i}(-, \mathbb{F}_\ell)$ is an object of Ho_\bullet ; as observed in [V1, p. 3] we have $H^{n,i}(X, \mathbb{F}_\ell) \cong \text{Hom}_{\text{Ho}_\bullet}(X_+, K)$. By the Yoneda Lemma, cohomology operations $H^{n,i}(-, \mathbb{F}_\ell) \rightarrow H^{p,q}(-, \mathbb{F}_\ell)$ correspond to elements of $H^{p,q}(K, \mathbb{F}_\ell)$. For example, under the map from $H^{p,q} = H^{p,q}(k, \mathbb{F}_\ell)$ to $H^{p,q}(K, \mathbb{F}_\ell)$ induced by the structure map $K \rightarrow \text{Spec}(k)$, $c \in H^{p,q}$ corresponds to the constant operation sending every element of $H^{n,i}(X, \mathbb{F}_\ell)$ to the image of c in $H^{p,q}(X, \mathbb{F}_\ell)$. Similarly, the canonical class $\iota \in H^{n,i}(K, \mathbb{F}_\ell)$ corresponds to the identity operation.

If $n \geq i$, we see from [V3, 3.27] that the summands of the motive $\mathbb{F}_{\ell\text{tr}}(K)$ having smallest weight or degree are $\mathbb{F}_{\ell}(i)[n]$ and $\mathbb{F}_{\ell}(i)[n+1]$. It follows that every cohomology operation $H^{n,i} \rightarrow H^{p,q}$ with $p \leq n+1$ is an \mathbb{F}_{ℓ} -linear combination of the Bockstein, the identity, and constant operations. More precisely:

- (a) if $p < n$, then $H^{p,*}(K) = H^{p,*}$, corresponding to constant operations;
- (b) $H^{n,*}(K) \cong H^{0,*} \cdot \iota \oplus H^{n,*}$, with $(b\iota, c)$ corresponding to the operation $\phi(x) = bx + c$;
- (c) $H^{n+1,*}(K) \cong H^{0,*} \cdot \beta(\iota) \oplus H^{1,*} \cdot \iota \oplus H^{n+1,*}$, with (a, b, c) corresponding to the operation $\phi(x) = a\beta(x) + bx + c$.

If $n < i$, this is no longer the case. In Example 11.5 below, we show that there is a weight-reducing operation $H^{1,2}(-, \mathbb{F}_{\ell}) \rightarrow H^{2,1}(-, \mathbb{F}_{\ell})$ for all k and a weight-preserving operation $H^{1,2}(-, \mathbb{F}_{\ell}) \rightarrow H^{3,2}(-, \mathbb{F}_{\ell})$ for most k . For another example, suppose that $\zeta \in k$ and $n \leq i$. Then cupping with $[\zeta] \in H^{0,1}(k, \mathbb{F}_{\ell})$ is an isomorphism by Theorem 6.2; its inverse (defined when $n < i$) is an operation $H^{n,i}(-, \mathbb{F}_{\ell}) \rightarrow H^{n,i-1}(-, \mathbb{F}_{\ell})$.

8. RELATION TO PERIODICITY

Sometimes we can deduce motivic operations from étale operations. For example, if $n \leq i$ (and hence $n \leq i\ell$), then the diagram (6.1) allows us to identify the motivic operation $P^0 : H^{n,i}(X) \rightarrow H^{n,i\ell}(X)$ with the étale operation $P^0 : H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i}) \cong H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i\ell})$ and thus conclude that P^0 is an isomorphism in this range. The same reasoning, using the Norm Residue Theorem 6.2, shows that if $n \leq i$ and $n + 2a(\ell - 1) \leq i\ell$, the motivic and étale operations P^a agree on $H^{n,i}(X) \cong H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i})$ and also agree with $b^{(i-a)(\ell-1)/d} P_V^a$, where $b \in H^{0,d}(k)$ is defined as follows.

Fix a primitive ℓ th root of unity, ζ , in an extension field of k ; this choice determines a generator $[\zeta]$ of $H^0(k(\zeta), \mu_{\ell})$. If $[k(\zeta) : k] = d$, then $H^{0,d}(k) \cong H_{\text{ét}}^0(k, \mu_{\ell}^{\otimes d}) \cong H_{\text{ét}}^0(k, \mathbb{F}_{\ell})$, and the element $[\zeta]^d = [\zeta^{\otimes d}]$ descends to a “periodicity” element b in $H^{0,d}(k)$. By abuse of notation, if $d|m$ we write $[\zeta]^m$ for the element $b^{m/d}$ of $H^{0,m}(k)$. (If $d = 1$, then $b = [\zeta]$.)

Note that multiplication by b is a map from $H^{n,i}(X)$ to $H^{n,i+d}(X)$; by the Norm Residue Theorem 6.2, it is an isomorphism when $i \geq n$. By construction, this is the map in cohomology induced by the change-of-truncation map

$$(8.1) \quad \mathbb{F}_{\ell}(i) \cong \tau^{\leq i} R\alpha_* \mu_{\ell}^{\otimes i} \rightarrow \tau^{\leq i+d} R\alpha_* \mu_{\ell}^{\otimes i} \cong \mathbb{F}_{\ell}(i+d)$$

associated to the isomorphism of étale sheaves $\mu_{\ell}^{\otimes i} \rightarrow \mu_{\ell}^{\otimes i+d}$ sending the generator $\zeta^{\otimes i}$ to the generator $\zeta^{\otimes i+d}$.

Write $H^{n,i}(X)[1/b]$ for the colimit of

$$H^{n,i}(X) \xrightarrow{b} H^{n,i+d}(X) \xrightarrow{b} \dots \xrightarrow{b} H^{n,i+jd}(X) \xrightarrow{b} \dots$$

From the diagram

$$\begin{array}{ccccccc} H^{n,i}(X) & \xrightarrow{b} & H^{n,i+d}(X) & \xrightarrow{b} & \dots H^{n,i+jd}(X) & \xrightarrow{b} & \dots \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i}) & \xrightarrow{\cong} & H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i+d}) & \xrightarrow{\cong} & \dots H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i+jd}) & \xrightarrow{\cong} & \dots \end{array}$$

we obtain a natural transformation from $H^{n,i}(X)[1/b]$ to $H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i})$.

We can formulate this in the motivic derived category DM , using the étale-to-Nisnevich change of topology map α . Recall from [MVW, 10.2] that $H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i})$ is isomorphic to

$$\text{Hom}_{DM_{\text{ét}}}(\alpha^* \mathbb{F}_{\ell, \text{tr}} X, \mathbb{F}_{\ell}(i)[n]) \cong \text{Hom}_{DM}(\mathbb{F}_{\ell, \text{tr}} X, R\alpha_* \mu_{\ell}^{\otimes i}).$$

The map $\mathbb{F}_{\ell}(i) = \tau^{\leq i} R\alpha_* \mu_{\ell}^{\otimes i} \rightarrow R\alpha_* \mu_{\ell}^{\otimes i}$ is compatible with the map $\mathbb{F}_{\ell}(i) \rightarrow \mathbb{F}_{\ell}(i+d) \rightarrow R\alpha_* \mu_{\ell}^{\otimes i+d}$, so it factors through a map $\mathbb{F}_{\ell}(i)[1/b] \rightarrow R\alpha_* \mu_{\ell}^{\otimes i}$, where $\mathbb{F}_{\ell}(i)[1/b]$ denotes the (homotopy) colimit in DM of

$$\mathbb{F}_{\ell}(i) \xrightarrow{b} \mathbb{F}_{\ell}(i+d) \xrightarrow{b} \mathbb{F}_{\ell}(i+2d) \xrightarrow{b} \cdots \xrightarrow{b} \mathbb{F}_{\ell}(i+jd) \xrightarrow{b} \cdots.$$

The following calculation is originally due to Levine [L].

Theorem 8.2. *For each i , $\mathbb{F}_{\ell}(i)[1/b] \rightarrow R\alpha_* \mu_{\ell}^{\otimes i}$ is an isomorphism in DM .*

For X smooth and all n , $H^{n,i}(X)[1/b] \rightarrow H_{\text{ét}}^n(X, \mu_{\ell}^{\otimes i})$ is an isomorphism.

Proof. Any complex C is the homotopy colimit of the change-of-truncation maps $\tau^{\leq m} C \rightarrow \tau^{\leq m+1} C$. For $C = R\alpha_* \mu_{\ell}^{\otimes i}$, this yields the first assertion. The second assertion is an immediate consequence of this and the fact that $\mathbb{F}_{\ell, \text{tr}} X$ is a compact object in DM , so $\text{Hom}_{DM}(\mathbb{F}_{\ell, \text{tr}} X, -)$ commutes with homotopy colimits. \square

Our next goal is to compare P^0 to the cohomology of the change-of-truncation map $\tau^{\leq i} R\alpha_* \mu_{\ell}^{\otimes i} \rightarrow \tau^{\leq i\ell} R\alpha_* \mu_{\ell}^{\otimes i}$ of (8.1).

Lemma 8.3. *The Frobenius map $\mathbb{F}_{\ell}(i) \xrightarrow{\Phi} \mathbb{F}_{\ell}(i\ell)$ in motivic cohomology is chain homotopic to the change-of-truncation map*

$$\mathbb{F}_{\ell}(i) \cong \tau^{\leq i} R\alpha_* \mu_{\ell}^{\otimes i} \rightarrow \tau^{\leq i\ell} R\alpha_* \mu_{\ell}^{\otimes i} \cong \mathbb{F}_{\ell}(i\ell).$$

The Frobenius $H^{n,i}(X) \xrightarrow{\Phi} H^{n,i\ell}(X)$ is multiplication by $b^{i(\ell-1)/d} = [\zeta^{\otimes i(\ell-1)}]$.

Proof. The Frobenius endomorphism is the identity on the étale sheaf of rings $A = \bigoplus_{i=0}^d \mu_{\ell}^{\otimes i}$, so if we fix i and an injective replacement $\mu_{\ell}^{\otimes i} \rightarrow I$, the Frobenius on $\mu_{\ell}^{\otimes i}$ lifts to a map $f_i : I \rightarrow I$ which is chain homotopic to the identity. Since the product in motivic cohomology is induced from the product on $R\alpha_* \mu_{\ell}^{\otimes i} = \alpha_* I$, the Frobenius in motivic cohomology is represented by the good truncation in degrees at most $i\ell$ of the composite $\tau^{\leq i} \alpha_* I \subset \alpha_* I \xrightarrow{f_i} \alpha_* I$. Since good truncation preserves chain homotopy, it is chain homotopic to the canonical map $\tau^{\leq i} \alpha_* I \subset \tau^{\leq i\ell} \alpha_* I$.

The final assertion follows from (8.1). \square

Proposition 8.4. *The map $P^0 : H^{n,i}(X) \rightarrow H^{n,i\ell}(X)$ is multiplication by $b^{i(\ell-1)/d}$. Equivalently, P^0 is the cohomology of the change-of-truncation map*

$$\tau^{\leq i} R\alpha_* \mu_{\ell}^{\otimes i} \rightarrow \tau^{\leq i\ell} R\alpha_* \mu_{\ell}^{\otimes i}.$$

Proof. Recall that the Godement resolution $\mathcal{F} \rightarrow S^{\bullet}(\mathcal{F})$ is a functorial simplicial resolution of any sheaf \mathcal{F} by flasque sheaves. Letting S_i^{\bullet} denote the total complex of the Godement resolution of $\tau^{\leq i} R\alpha_* \mu_{\ell}^{\otimes i}$, it follows that the product on $\bigoplus \tau^{\leq i} R\alpha_* \mu_{\ell}^{\otimes i}$ induces a product pairing $S_i^n \otimes S_j^n \rightarrow S_{i+j}^n$ for all n . In particular, the Frobenius on $R\alpha_* \mu_{\ell}^{\otimes i}$ induces a map $S_i^n \rightarrow S_{i\ell}^n$.

In [E, 11.1], Epstein shows that the Godement resolution satisfies the conditions of his Section 8. By functoriality, the equivariant map $(R\alpha_* \mu_{\ell}^{\otimes i})^{\otimes \ell} \rightarrow R\alpha_* \mu_{\ell}^{\otimes i\ell}$ constructed after Theorem 6.2 lifts to an equivariant map $(S_i^{\bullet})^{\otimes \ell} \rightarrow S_{i\ell}^{\bullet}$. This is the analogue of [E, 8.3.2] and is exactly what we need in order for the proof of

[E, 8.3.4] to work. Thus if we represent $v \in H^{n,i}(X)$ by a cocycle u in the algebra $H^0(X, S^n)$, then $P^0 v$ is represented by the element u^ℓ of $H^0(X, S^n)$. Therefore P^0 is represented by the Frobenius. \square

Example 8.4.1. Recall that $b \in H^{0,d}(k)$. Since $P^0(b) = b^\ell$ (by Proposition 8.4), the Cartan formula Theorem 7.4 yields $P^a(bx) = b^\ell P^a(x)$.

Recall from [V3, 2.60] that a *split proper Tate motive* is a direct sum of Tate motives $\mathbb{F}_\ell(i)[2i+j]$ with $j \geq 0$. If the weights i are at least n , then we say the motive has weight $\geq n$. Note that the cohomology of $\mathbb{F}_\ell(i)[2i+j]$ is a free bigraded $H^{*,*}$ -module of rank 1 with a generator in bidegree $(2i+j, i)$.

It follows that we have a Künneth formula (see [W, 4.1]): if $\mathbb{F}_{\ell,\text{tr}}(Y)$ is a split proper Tate motive, then $H^{*,*}(Y)$ is a free bigraded $H^{*,*}$ -module, and

$$(8.5) \quad H^{*,*}(X \times Y) \cong H^{*,*}(X) \otimes_{H^{*,*}} H^{*,*}(Y).$$

Example 8.6. Let $K = K(\mathbb{F}_\ell(i), n)$ be the Eilenberg-Mac Lane space classifying $H^{n,i}(-, \mathbb{F}_\ell)$; if $n \geq 2i \geq 0$, then $M = \mathbb{F}_{\ell,\text{tr}}(K)$ is a split proper Tate motive of weight $\geq i$, by [V3, 3.28]. It follows that $H^{*,*}(K^{\otimes p})$ is the p -fold tensor product of $H^{*,*}(K)$ with itself over $H^{*,*}$.

Recall from [MVW, 3.1] that $\mathbb{F}_\ell(i)[i]$ is represented by the abelian presheaf $\mathbb{F}_{\ell,\text{tr}}(\mathbb{G}_m^{\wedge i})$, so $\mathbb{F}_\ell(i)[n]$ is represented by the simplicial abelian presheaf associated to $\mathbb{F}_{\ell,\text{tr}}(\mathbb{G}_m^{\wedge i})[n-i]$ when $n \geq i$. From the adjunction

$$\text{Hom}_{\text{Ho}_\bullet}(X, u\mathbb{F}_\ell(i)[n]) \cong \text{Hom}_{DM}(\mathbb{F}_{\ell,\text{tr}}(X), \mathbb{F}_\ell(i)[n]) = H^{n,i}(X)$$

(see Example 7.9) we see that the classifying space $K(\mathbb{F}_\ell(i), n)$ of $H^{n,i}(-, \mathbb{F}_\ell)$ is the simplicial abelian presheaf $G = u\mathbb{F}_\ell(i)[n]$ underlying $\mathbb{F}_\ell(i)[n]$; see [V3, p. 5].

Lemma 8.7. *If $\mathbb{F}_{\ell,\text{tr}}(Y)$ is a split proper Tate motive, then multiplication by b^e is an injection from $H^{p,q}(Y, \mathbb{F}_\ell)$ into $H^{p,q+de}(Y, \mathbb{F}_\ell)$ and hence into $H^{p,q}(Y, \mathbb{F}_\ell)[1/b]$.*

Proof. It suffices to consider $\mathbb{F}_{\ell,\text{tr}}(Y) = \mathbb{F}_\ell(i)[2i+j]$. There is no harm in increasing e so that $(\ell-1)|de$. Set $p' = p - 2i - j$ and $q' = q - i$. Since $H^{*,*}(Y)$ is a free $H^{*,*}(k)$ -module of rank 1, the assertion for $H^{p,q}(Y)$ amounts to the assertion that either $H^{p',q'}(k) = 0$ (and injectivity is obvious) or else $0 \leq p' \leq q'$ and $H^{p',q'}(k) \cong H_{\text{et}}^{p'}(k, \mu_\ell^{\otimes q'})$. In the latter case, we also have $H^{p',q'+de}(k) \cong H_{\text{et}}^{p'}(k, \mu_\ell^{\otimes q'+de})$, and the isomorphism is induced from the isomorphism $\mu_\ell^{\otimes q'} \cong \mu_\ell^{\otimes q'+de}$. \square

In the next proposition, we write K for $K(\mathbb{F}_\ell(i), n)$. For each p and q , there is a canonical map $H^{p,q}(K, \mathbb{F}_\ell) \rightarrow H_{\text{et}}^p(K, \mu_\ell^{\otimes q})$. It sends the motivic operations P^a of Definition 6.5 to the étale operations P^a of Theorem 1.3.

Proposition 8.8. *If $n \geq 2i$, the canonical map is an injection, from the set $H^{p,q}(K, \mathbb{F}_\ell)$ of motivic cohomology operations $H^{n,i} \rightarrow H^{p,q}$ to the set $H_{\text{et}}^p(K, \mu_\ell^{\otimes q})$ of étale cohomology operations $H_{\text{et}}^n(-, \mu_\ell^{\otimes i}) \rightarrow H_{\text{et}}^p(-, \mu_\ell^{\otimes q})$.*

Proof. By the usual transfer argument, we may assume that $\zeta \in k$. Let K denote the Eilenberg-Mac Lane space classifying $H^{n,i}(-, \mathbb{F}_\ell)$. By Example 8.6, $\mathbb{F}_{\ell,\text{tr}}(K)$ is a split proper Tate motive. By Lemma 8.7 and Levine's Theorem 8.2, $H^{p,q}(K, \mathbb{F}_\ell)$ injects into $H^{p,q}(K, \mathbb{F}_\ell)[1/b] \cong H_{\text{et}}^p(K, \mu_\ell^{\otimes q})$. Thus the group $H^{p,q}(K, \mathbb{F}_\ell)$ of motivic cohomology operations injects into the group $H_{\text{et}}^p(K, \mu_\ell^{\otimes q})$ of étale cohomology operations. \square

Corollary 8.9. *Suppose that $n \geq 2i$ and $n \geq 2a$. Then for $x \in H^{n,i}(X)$:*

- (1) *If $a \leq i$, $P^a(x) = [\zeta]^{(i-a)(\ell-1)} \cup P_V^a(x)$.*
- (2) *If $a \geq i$, $P_V^a(x) = [\zeta]^{(a-i)(\ell-1)} \cup P^a(x)$.*

Proof (Cf. [BJ, Thm. 1.1]). The two sides have the same bidegree and agree with $P^a(x)$ in étale cohomology by Lemma 6.6 and Remark 7.3.1. \square

Corollary 8.10. *If $n \geq i$ and $x \in H^{2n,i}$, then $P_V^n(x) = [\zeta]^{(n-i)(\ell-1)} \cup x^\ell$.*

Proof. This is the case $a = n$ of Corollary 8.9, as $P^n(x) = x^\ell$ (Proposition 7.1). \square

We can now show that the motivic Q^a equals βP^a .

Theorem 8.11. *The motivic operations Q^a on $H^{n,i}$ satisfy $Q^a = \beta P^a$.*

Proof. Set $K = K(\mathbb{F}_\ell(i), n)$, so that motivic cohomology operations $H^{n,i} \rightarrow H^{N,i\ell}$ correspond to elements of $H^{N,i\ell}(K)$ (see Example 8.6). Now the identity on $H^{n,i}$ is represented by the canonical element ι of $H^{n,i}(K)$, and the motivic cohomology operations Q^a and βP^a are represented by the elements $Q^a(\iota)$ and $\beta P^a(\iota)$ of $H^{N,i\ell}(K)$.

We first consider Q^0 . The map $H^{n+1,i\ell}(K) \rightarrow H_{\text{et}}^{n+1}(K, \mu_\ell^{\otimes i\ell})$ is an isomorphism if $n \leq i\ell$ by Theorem 6.2 and is an injection if $n \geq 2i$ by Proposition 8.8. By Lemma 6.6 and Remark 6.6.1, we have a commutative diagram:

$$\begin{array}{ccc} H^{n,i}(K) & \xrightarrow{Q^0 - \beta P^0} & H^{n+1,i\ell}(K) \\ \downarrow & & \text{into} \downarrow \\ H_{\text{et}}^n(K, \mu_\ell^{\otimes i}) & \xrightarrow{Q^0 - \beta P^0} & H_{\text{et}}^{n+1}(K, \mu_\ell^{\otimes i\ell}). \end{array}$$

The bottom map is zero by Proposition 3.3. It follows that $Q^0(\iota) = \beta P^0(\iota)$ in $H^{n+1,i\ell}(K)$ and hence that $Q^0 = \beta P^0$ as motivic cohomology operations.

Now suppose that $a > 0$, and set $N = n + a(\ell - 1) + 1$. If $n \geq 2i$, we consider the commutative diagram:

$$\begin{array}{ccc} H^{n,i}(K) & \xrightarrow{Q^a - \beta P^a} & H^{N,i\ell}(K) \\ \downarrow & & \text{into} \downarrow \\ H_{\text{et}}^{n,i}(K, \mu_\ell^{\otimes i}) & \xrightarrow{Q^a - \beta P^a} & H_{\text{et}}^N(K, \mu_\ell^{\otimes i\ell}). \end{array}$$

The lower horizontal map is zero by Proposition 3.3. The right vertical map is an injection by Proposition 8.8, because $\mathbb{F}_{\ell,\text{tr}}(K)$ is a split proper Tate motive (by Example 8.6). It follows that $Q^a(\iota) = \beta P^a(\iota)$ in $H^{N,i\ell}(K)$ and hence that $Q^a = \beta P^a$ as cohomology operations on $H^{n,i}$.

If $n < 2i$, we consider suspension $S^d K$, where $d = 2i - n$. By Proposition 7.8, we have a commutative diagram with vertical isomorphisms:

$$\begin{array}{ccc} H^{n,i}(K) & \xrightarrow{Q^a - \beta P^a} & H^{N,i\ell}(K) \\ \cong \downarrow & & \cong \downarrow \\ H^{2i,i}(S^d K) & \xrightarrow{Q^a - \beta P^a} & H^{N+d,i\ell}(SK). \end{array}$$

By the above argument, $Q^a = \beta P^a$ on $H^{2i,i}(S^d K)$. It follows that $Q^a = \beta P^a$ on $H^{n,i}(K)$ and hence as cohomology operations on $H^{n,i}$. \square

9. BOREL'S THEOREM

In order to go from $H^{1,*}$ to $H^{n,*}$, we need a slight generalization of Borel's theorem [McC, 6.21], one which accounts for the coefficient ring $H^{*,*} = H^{*,*}(\mathrm{Spec} k)$.

Definition 9.1. Let H^* be a graded-commutative \mathbb{F}_ℓ -algebra. If W^* is a graded H^* -algebra, an ℓ -simple system of generators of W^* over H^* is a totally ordered set of elements x_i such that W^* is a free left H^* -module on the monomials $x_{i_1}^{m_1} \cdots x_{i_k}^{m_k}$, where the i 's are in order and $0 \leq m_j < \ell$ (with $m_j \leq 1$ if $\deg(x_j)$ is odd).

Theorem 9.2. Let H^* be a graded-commutative \mathbb{F}_ℓ -algebra with $H^0 = \mathbb{F}_\ell$, and suppose that $\{E_r^{*,*}, d_r\}$ is a 1st-quadrant spectral sequence of graded-commutative H^* -algebras converging to H^* . Set $V^* = E_2^{*,0}$ and $W^* = E_2^{0,*}$, and suppose that

(i) $E_2^{*,*} \cong W^* \otimes_{H^*} V^*$ as algebras, and that

(ii) the H^* -algebra W^* has an ℓ -simple system of generators $\{x_i\}$, each of which is transgressive.

Then V^* is the tensor product of H^* and a free graded-commutative \mathbb{F}_ℓ -algebra on generators $y_i = \tau(x_i)$ and (when $\ell \neq 2$ and $\deg(x_j)$ is even) $z_j = \tau(y_j \otimes x_j^{\ell-1})$. (Here τ is the transgression.)

Proof. The proof of Borel's Theorem in [McC, 6.21] goes through. \square

Let G be a simplicial sheaf of groups, such as $B\mu_\ell$. We use the bar construction to form the bisimplicial classifying spaces $B_\bullet G$ (with G^p in simplicial degree p) of a simplicial sheaf of groups G and $E_\bullet G$ (with G^{p+1} in simplicial degree p). We write the canonical projection as $E_\bullet G \xrightarrow{\pi} B_\bullet G$. The Leray spectral sequence [Milne, III.1.18] becomes

$$(9.3) \quad E_2^{p,q} = H^p(B_\bullet G, R^q\pi_* A) \Rightarrow H^{p+q}(E_\bullet G, A).$$

Proposition 9.4. Suppose that G is a simplicial sheaf of groups on a site over k and A is a sheaf of homotopy-associative dg \mathbb{F}_ℓ -algebras satisfying $H^0(G, A) = H^0(k, A)$ as well as the Künneth condition that

$$H^*(U, A) \otimes_{H^*(k, A)} H^*(G, A) \xrightarrow{\cong} H^*(U \times G, A)$$

is an isomorphism for all U in the site of $B_\bullet G$. Then the Leray spectral sequence (9.3) satisfies condition (i) of Borel's Theorem 9.2 with

$$E_2^{p,q} = H^p(B_\bullet G, A) \otimes_{H^*(k, A)} H^q(G, A).$$

Proof. For simplicity of notation, let us write \otimes_H for $\otimes_{H^*(k, A)}$. We first claim that the higher direct images $R^q\pi_*(A)$ are $A \otimes_H H^*(G, A)$. To see this, recall that $R^q\pi_*(A)$ is the sheafification of the presheaf that to a map $U \rightarrow B_p G$ associates $H^q(\pi^{-1}U, A)$, where $\pi^{-1}U = E_\bullet G \times_{B_p G} U$ is $U \times G$. By hypothesis, $H^*(\pi^{-1}U, A)$ is $H^*(U, A) \otimes_H H^*(G, A)$. The claim follows, since sheafification commutes with $\otimes_H H^*(G, A)$, and the sheaf associated to $H^q(-, A)$ is A if $q = 0$ and zero for $q > 0$. Thus we have $E_2^{p,q} = H^p(B_\bullet G, A) \otimes_H H^q(G, A)$.

It remains to check the Borel condition. By hypothesis, $H^0(G, A) = H^0(k, A)$, so $E_2^{p,0} = H^p(B_\bullet G, A)$, and $H^0(B_\bullet G, A) = H^0(k, A)$. Thus we have

$$E_2^{0,q} = H^0(B_\bullet G, A) \otimes_H H^q(G, A) = H^q(G, A).$$

The fact that the spectral sequence is multiplicative follows from the fact that A is a sheaf of algebras and the work of Massey [Mass]. \square

Kudo's Theorem 9.5. *Suppose G and A satisfy the hypotheses of Proposition 9.4. If $x \in H^n(G, A)$ transgresses to $y \in H^{n+1}(B_\bullet G, A)$, then*

- (1) $\beta(x)$ transgresses to $-\beta(y)$,
- (2) $P^a(x)$ transgresses to $P^a(y)$, and
- (3) if $n = 2a$, then $x^{\ell-1} \otimes y$ transgresses to $-Q^a(y)$.

Any simplicially stable operation commutes with the transgression; see [McC, 6.5]. Hence part (2) of Theorem 9.5 is immediate whenever we know that P^a is simplicially stable. This is so for the operations P^a in étale and motivic cohomology (by Propositions 1.9 and 7.8).

Proof (Cf. [M, 3.4]). As in the proof of Theorem 1.5, we fix a quasi-isomorphism $A \xrightarrow{\sim} I^*$. Let $f = \pi^*$ and $g = i^*$ be the canonical maps $I(G) \xleftarrow{g} I(E_\bullet G) \xleftarrow{f} I(B_\bullet G)$ coming from $G \xrightarrow{i} E_\bullet G \xrightarrow{\pi} B_\bullet G$. The assertion that x transgresses to y means that there is a cocycle b in $I^{n+1}(B_\bullet G)$ representing y and an element u in $I^n(E_\bullet G)$, such that $f(b) = du$ and $g(u)$ is a cocycle representing x .

Since the Bockstein satisfies $g(\beta u) = \beta g(u)$ and $f(\beta b) = \beta(f(b)) = \beta(du) = -d(\beta u)$, we see that $\beta(x)$, which is represented by $g(\beta u)$, transgresses to $-\beta(y)$.

Recall from Section 4 that b and u determine a cocycle $P^a(b)$ in $I^*(B_\bullet G)$ representing $P^a(y)$ and an element $P^a(u)$ in $I^*(E_\bullet G)$ so that $P^a(x)$ is represented by $P^a g(u) = g P^a(u)$. By Lemma 4.8, we have

$$f P^a(b) = P^a f(b) = P^a(du) = d P^a(u).$$

It follows that $P^a(x)$ transgresses to $P^a(y)$.

Since b is a cocycle, $Q^a(b)$ represents $Q^a(y)$, and by Lemma 4.8 we have

$$f Q^a(b) = Q^a f(b) = Q^a(du) = -d(Q^a u).$$

Thus the class of $Q^a(u)$ transgresses to $-Q^a(y)$, and it suffices to show that $Q^a(u)$ represents $x^{\ell-1} \otimes y$ under the isomorphism $E_2^{p,q} \cong H^p(B_\bullet G) \otimes H^q(G)$ of Proposition 9.4.

Recall from (1.8) that $\nu_n = (-1)^r m!^{-n}$, where $m = (\ell - 1)/2$. We have $\nu_n = (-1)^a$, because $n = 2a$, $(m!)^2 = (-1)^{m+1}$, and $r \equiv am \pmod{2}$. We now follow p. 167 of [M] up to (9). Starting from $u \in I^n(X)$, May produces elements t_i in $I^{\otimes \ell}(X)$ and a family of elements $\{c_a\}$, $\{c'_a\}$ in $C_* \otimes I^{\otimes \ell}(X)$, depending naturally on u , such that

$$Q_M^a(u) = (-1)^a \nu(1 - n) \theta(c'_a) = m! \theta(c'_a).$$

The analysis of the terms in c'_a at the top of p. 171 of [M] shows that there is a term c'' such that $c' - d(c'')$ is $(-1)^m m! z$ plus terms mapped by θ into lower parts of the filtration, where $z = e_0 \otimes u \otimes \cdots \otimes u \otimes du$, and that $\theta(z)$ represents $x^{\ell-1} \otimes y$. Therefore, up to terms in lower parts of the filtration we have $Q_M^a(u) = m! \theta(c'_a) = (-1)^m (m!)^2 \theta(z) = -\theta(z)$. Since we saw in Corollary 4.7 that $Q^a(u) = Q_M^a(u)$, the result follows. \square

We illustrate the use of Proposition 9.4 with the étale topology. First, consider the étale sheaf $G = \mu_\ell$. If μ_ℓ is connected, then it does not satisfy the Künneth condition of Proposition 9.4 for $U = \text{Spec}(\bar{k})$. Indeed, $H^0(\mu_\ell, \mathbb{F}_\ell) = \mathbb{F}_\ell$, yet $H^0(G \times \text{Spec}(\bar{k}), \mathbb{F}_\ell) = \prod_1^\ell \mathbb{F}_\ell$. However, things change if we consider the étale sheaf $\mathcal{A} = \bigoplus_{i=0}^{d-1} \mu_\ell^{\otimes i}$ of Section 2.

Lemma 9.6. $H_{\text{ét}}^*(X \times \mu_\ell^{\otimes i}, \mathcal{A}) \cong H_{\text{ét}}^*(X, \mathcal{A}) \otimes_{H^*(k, \mathcal{A})} H_{\text{ét}}^*(\mu_\ell^{\otimes i}, \mathcal{A})$.

Proof. As an étale sheaf of \mathbb{F}_ℓ -modules, constant over $k(\zeta)$, $\mathbb{F}_\ell[\mu_\ell^{\otimes i}]$ is a direct sum of the locally constant sheaves $\mu_\ell^{\otimes \alpha}$, each of which is an invertible object. Because $\mathbb{F}_\ell[X \times \mu_\ell^{\otimes i}] \cong \mathbb{F}_\ell[X] \otimes \mathbb{F}_\ell[\mu_\ell^{\otimes i}]$, $H_{\text{et}}^n(X \times \mu_\ell^{\otimes i}, \mu_\ell^{\otimes q})$ equals

$$\begin{aligned} \text{Ext}^n(\mathbb{F}_\ell[X] \otimes \mathbb{F}_\ell[\mu_\ell^{\otimes i}], \mu_\ell^{\otimes q}) &\cong \text{Ext}^n(\mathbb{F}_\ell[X], \mathcal{R}\mathcal{H}\text{om}(\mathbb{F}_\ell[\mu_\ell^{\otimes i}], \mu_\ell^{\otimes q})) \\ &\cong \text{Ext}^n(\mathbb{F}_\ell[X], \mathcal{R}\mathcal{H}\text{om}(\mathbb{F}_\ell, \oplus \mu_\ell^{\otimes q-\alpha})) \\ &\cong \bigoplus_{\alpha} \text{Ext}^n(\mathbb{F}_\ell[X], \mu_\ell^{\otimes q-\alpha}) \cong \bigoplus_{\alpha} H_{\text{et}}^n(X, \mu_\ell^{\otimes q-\alpha}). \end{aligned}$$

The pairing $H_{\text{et}}^*(X, \mathcal{A}) \otimes_{\mathbb{F}_\ell} H_{\text{et}}^*(\mu_\ell^{\otimes i}, \mathcal{A}) \rightarrow H_{\text{et}}^*(X \times \mu_\ell^{\otimes i}, \mathcal{A})$ is the direct sum over α , s , and t of the top row in the commutative diagram

$$\begin{array}{ccc} \text{Ext}^*(\mathbb{F}_\ell[X], \mu_\ell^{\otimes s}) \otimes_{\mathbb{F}_\ell} \text{Ext}^*(\mu_\ell^{\otimes \alpha}, \mu_\ell^{\otimes t}) & \longrightarrow & \text{Ext}^*(\mathbb{F}_\ell[X] \otimes \mu_\ell^{\otimes \alpha}, \mu_\ell^{\otimes s+t}) \\ \cong \downarrow & & \downarrow \cong \\ \text{Ext}^*(\mathbb{F}_\ell[X], \mu_\ell^{\otimes s}) \otimes_{\mathbb{F}_\ell} \text{Ext}^*(\mathbb{F}_\ell, \mu_\ell^{\otimes t-\alpha}) & \longrightarrow & \text{Ext}^*(\mathbb{F}_\ell[X], \mu_\ell^{\otimes s+t-\alpha}). \end{array}$$

Since $H_{\text{et}}^*(k, \mathcal{A}) = \text{Ext}^*(\mathbb{F}_\ell, \mathcal{A}) \cong \bigoplus_t \text{Ext}(\mathbb{F}_\ell, \mu_\ell^{\otimes t-\alpha})$ for each α , setting $s = q - \alpha$ and summing over s and t yields the result. \square

Corollary 9.7. *If Y is a coproduct of schemes which are finite products of $\mu_\ell^{\otimes i}$, then*

$$H_{\text{et}}^*(X \times Y, \mathcal{A}) \cong H_{\text{et}}^*(X, \mathcal{A}) \otimes_{H_{\text{et}}^*(k, \mathcal{A})} H_{\text{et}}^*(Y, \mathcal{A}).$$

Example 9.8. The ring of all étale cohomology operations from $H_{\text{et}}^2(-, \mu_\ell^{\otimes i})$ to $H_{\text{et}}^*(-, \mu_\ell^{\otimes *})$ is the free left $H_{\text{et}}^*(k, \mathcal{A})$ -module on generators in $H^{*,i}(K_2)$: monomials in the identity ($\text{id} \in H_{\text{et}}^2(K_2, \mu_\ell^{\otimes i})$), β , the $P^I\beta$, and the $\beta P^I\beta$ ($P^I = P^{\ell^\nu} \cdots P^\ell P^1$). This result, proven in Theorem 3.5 above, can also be obtained from the Leray spectral sequence (9.3).

Each term in the simplicial sheaf $B_{\bullet} \mu_\ell^{\otimes i}$ is a coproduct of products of $\mu_\ell^{\otimes i}$, so Corollary 9.7 and Proposition 9.4 imply that the Leray spectral sequence satisfies condition (i) of Borel's Theorem 9.2. The explicit description of $H_{\text{et}}^*(B\mu_\ell^{\otimes i}, \mathcal{A})$ in Proposition 3.4 as $H_{\text{et}}^*(k, \mathcal{A}) \otimes \mathbb{F}_\ell[u, v]/(u^2)$ shows that it has an ℓ -simple system of generators: u and the $x_\nu = v^{\ell^\nu}$ for $\nu \geq 0$. The transgression τ sends u to ι , so $v = \beta u$ transgresses to $-\beta$, by Kudo's Theorem 9.5(1). Thus condition (ii) is also satisfied, and Borel's Theorem states that $H_{\text{et}}^*(K_2, \mathcal{A})$ is the free graded-commutative $H^{*,*}$ -algebra on generators $\iota \in H^{2,i}(K_2)$,

$$y_\nu = \tau(x_\nu) \in H^{2\ell^\nu+1}(K_2, \mu_\ell^{\otimes i\ell^\nu}) \quad \text{and} \quad z_\nu = \tau(x_\nu^{\ell-1} \otimes y_\nu) \in H^{2\ell^\nu+1+2}(K_2, \mu_\ell^{\otimes i\ell^\nu+1}).$$

Note that $y_0 = \beta(\iota)$. Since $x_{\nu+1} = x_\nu^{\ell^\nu} = P^{\ell^\nu}x_\nu$, Kudo's Theorem 9.5(2) and an inductive argument show that $y_{\nu+1}$ is $P^{\ell^\nu}y_\nu$ and also $P^{\ell^\nu} \cdots P^\ell P^1 \beta$. This completes the proof for $\ell = 2$.

For $\ell > 2$, it remains to show that $-z_\nu$ is $\beta P^{\ell^\nu}(y_\nu) = \beta P^{\ell^\nu} \cdots P^\ell P^1 \beta$. This follows from Kudo's Theorem 9.5(3), using Proposition 3.3 to write βP^a for Q^a .

10. MOTIVIC OPERATIONS ON WEIGHT 1 COHOMOLOGY

We now turn to natural operations defined on the motivic cohomology groups with weight 1, i.e., $H^{n,1}(X) = H^{n,1}(X, \mathbb{F}_\ell)$. We begin with the case $n = 1$.

Let μ_ℓ be the group scheme of ℓ th roots of unity. On pp. 130–131 of [MV], Morel and Voevodsky define a simplicial Nisnevich sheaf $B_{\text{et}}\mu_\ell$ and observe that

it classifies the étale cohomology group $H_{\text{ét}}^1(-, \mu_\ell)$, and hence the motivic group $H^{1,1}$ by Theorem 6.2, in the sense that $[X_+, B_{\text{ét}}\mu_\ell] \cong H^{1,1}(X)$ for every smooth simplicial scheme X over k .

Following [V1, p. 17], we write $B\mu_\ell$ for the geometric classifying space of μ_ℓ , constructed in [MV, p. 133] (where the notation $B_{gm}\mu_\ell$ was used). By [MV, 4.2.7], $B\mu_\ell$ is \mathbb{A}^1 -equivalent to $B_{\text{ét}}\mu_\ell$, so it also classifies $H^{1,1}$.

When $\ell = 2$, the generator $[\zeta]$ of $H^{0,1}(k) = \mu_2(k)$ and its Bockstein, the element $[-1] \in H^{1,1}(k) = k^\times/k^{\times\ell}$, play an important role.

Proposition 10.1. *There are elements $u \in H^{1,1}(B\mu_\ell)$, $v \in H^{2,1}(B\mu_\ell)$ such that*

$$H^{*,*}(B\mu_\ell) \cong \begin{cases} H^{*,*}(k) \otimes \mathbb{F}_\ell[u, v]/(u^2), & \ell \neq 2, \\ H^{*,*}(k) \otimes \mathbb{F}_\ell[u, v]/(u^2 + [-1]u + [\zeta]v), & \ell = 2. \end{cases}$$

Thus every cohomology operation on $H^{1,1}(X)$ is uniquely a sum of the operations $x \mapsto cx^\varepsilon \beta(x)^m$, where $c \in H^{s,j}(k)$, $m \geq 0$, and $0 \leq \varepsilon \leq 1$.

Proof. This is the special case $F_\bullet = S^0$ in Proposition 6.10 of [V1]. Note that the operation $cx^\varepsilon(\beta x)^m$ has bidegree $(s + 2m + \varepsilon - 1, j + m + \varepsilon - 1)$. \square

As in Example 9.8, we can use this as the starting point to describe all motivic operations on $H^{*,1}$. For example, $G = B\mu_\ell$ is a simplicial group-scheme whose class in DM is a split proper Tate motive, so the Künneth formula (8.5) holds. Since $H^{0,q}(B\mu_\ell) \cong H^{0,q}(k)$ by Proposition 10.1, Proposition 9.4 applies to show that the Leray spectral sequence has the form

$$(10.2) \quad E_2^{p,q} = H^{p,*}(B_\bullet B\mu_\ell) \otimes_{H^{*,*}(k)} H^{q,*}(B\mu_\ell) \Rightarrow H^{p+q,*}(E_\bullet B\mu_\ell) \cong H^{p+q,*}(k).$$

Corollary 10.3. *If $\ell \neq 2$, the ring of cohomology operations on $H^{2,1}$ is the tensor product of $H^{*,*}(k, \mathbb{F}_\ell)$ and the free graded-commutative algebra generated by the identity of $H^{2,1}$, the Bockstein β , the $P^I \beta$, and the $\beta P^I \beta$ where $P^I = P^{\ell^\nu} \cdots P^\ell P^1$.*

For $\ell = 2$, the ring of cohomology operations on $H^{2,1}$ is the tensor product of $H^{,*}(k, \mathbb{F}_2)$ and the free graded-commutative algebra generated by the identity of $H^{2,1}$, $Sq^1, \dots, Sq^I = Sq_V^I$, where $Sq^I = Sq^{2^\nu} \cdots Sq^2 Sq^1$.*

Proof. By [MV, 4.1.16], $B\mu_\ell$ is $K(\mathbb{F}_\ell(1), 1)$. From the sequence

$$\begin{aligned} \text{Hom}_{\text{Ho}_\bullet}(X_+, E_\bullet B\mu_\ell) &\rightarrow \text{Hom}_{\text{Ho}_\bullet}(X_+, B_\bullet B\mu_\ell) \rightarrow \text{Hom}_{\text{Ho}_\bullet}(X_+, B\mu_\ell[1]) \\ &\rightarrow \text{Hom}_{\text{Ho}_\bullet}(X_+, E_\bullet B\mu_\ell[1]) \end{aligned}$$

it follows that the simplicial scheme $B_\bullet B\mu_\ell$ is $K(\mathbb{F}_\ell(1), 2)$. Thus we merely need to compute the motivic cohomology of $K(\mathbb{F}_\ell(1), 2)$ using (10.2).

By Proposition 10.1, $H^{*,*}(B\mu_\ell)$ has an ℓ -simple system of generators over $H^{*,*}$ consisting of u and the $x_\nu = v^{\ell^\nu}$ for $\nu \geq 0$. Since the Künneth formula (8.5) holds, Proposition 9.4 implies that the hypotheses of Borel's Theorem 9.2 and Kudo's Theorem 9.5 hold for (10.2). Therefore $H^{*,*}(B_\bullet B\mu_\ell)$ is the tensor product of $H^{*,*}$ and the free graded-commutative \mathbb{F}_ℓ -algebra on generators ι , $y_\nu = \tau(x_\nu)$ — and z_ν if $\ell > 2$. Since u transgresses to ι , $x_0 = v = \beta(u)$ and $x_{\nu+1} = P^{\ell^\nu} x_\nu$, Kudo's Theorem 9.5 implies (by induction) that $y_0 = \beta(\iota)$, $y_{\nu+1} = P^{\ell^\nu} y_\nu = P^I \beta(\iota)$, and (using Theorem 8.11) that z_ν is $-\beta P^I \beta(\iota)$. \square

To describe cohomology operations on $H^{n,1}$, we use the algebra $H_{\text{top}}^*(K_n)$, defined in Definition 0.1. We bigrade it by giving it the weight grading that P^I has weight $\ell^k - 1$, where $I = (\epsilon_0, s_1, \epsilon_1, \dots, s_k, \epsilon_k)$.

Theorem 10.4. *For each $n \geq 1$, the ring of all motivic cohomology operations on $H^{n,1}$ is isomorphic to the free left $H^{*,*}$ -module $H^{*,*}(k) \otimes H_{\text{top}}^*(K_n)$ in which the P^I are bigraded according to Definition 6.5.*

Thus every cohomology operation on $H^{n,1}(X)$ is a sum of the operations $x \mapsto c(P^{I_1}x)(P^{I_2}x) \cdots (P^{I_s}x)$, where $c \in H^{,*}(k)$ and each I_j satisfies the excess condition of Definition 0.1.*

Proof. We proceed by induction on n , the cases $n = 1, 2$ being given above. Set $K_n = K(\mathbb{F}_\ell(1), n)$, so $K_{n+1} = B_\bullet(K_n)$, and suppose inductively that the algebra $H^{*,*}(K_n)$ is given as described in the theorem, so that it has an ℓ -simple system of generators consisting of the $P^I(\iota_n)$ with I admissible and $e(I) < n$ (or $e(I) = n$ and $\epsilon_1 = 1$), and ℓ^ν powers of the $P^I(\iota_n)$ of even degree.

Since $\mathbb{F}_{\ell,\text{tr}}(K_n)$ is a split proper Tate motive by [V3, 3.28], the Künneth condition (8.5) of Proposition 9.4 holds. Hence the hypotheses of Borel's Theorem 9.2 are satisfied, and the Leray spectral sequence (9.3) has the form

$$E_2^{p,q} = H^{p,*}(K_{n+1}) \otimes_{H^{*,*}(k)} H^{q,*}(K_n) \Rightarrow H^{p+q,*}(E) \cong H^{p+q,*}(k).$$

Therefore $H^{*,*}(K_{n+1})$ is the tensor product of $H^{*,*}$ and a free graded-commutative \mathbb{F}_ℓ -algebra on certain generators; it remains to establish that they are the ones described in the theorem. But, except for weight considerations, this is exactly the same as in the topological case, as presented on p. 200 of [McC]. Of course, the weight of the $x_I = P^I(\iota_n)$ is the same as the weight of $y_I = P^I(\iota_{n+1})$. Inspection of the weights of the new generators $P^{\ell^s} \cdots P^s y_I$ (when x_I has degree $2s$) shows that each additional P^{ℓ^s} multiplies the weight by ℓ , as required. \square

11. MOTIVIC OPERATIONS ON DEGREE 1 COHOMOLOGY

We now turn to operations defined on $H^{1,*}$. Here we encounter new cohomology operations arising from the Norm Residue Theorem 6.2, representing a negative twist. Here are a couple of examples.

Example 11.1. There are operations $H^{1,r(\ell-1)}(X) \xrightarrow{\sim} H^{1,1}(X)$, since both groups are naturally isomorphic to $H_{\text{et}}^1(X, \mu_\ell)$. An element $\eta \in H_{\text{et}}^1(k, \mu_\ell^{\otimes 2-i})$ determines a natural transformation $H^{1,i}(X) \rightarrow H^{2,2}(X)$.

The case $k = k(\zeta)$. If k contains a primitive ℓ th root of unity ζ , the classification is immediate from Proposition 10.1. Let $[\zeta]$ be the class of ζ in $H^{0,1}(k) \cong \mu_\ell$.

Proposition 11.2. *Suppose that $\zeta \in k$ and $i > 1$. Then there is a natural isomorphism $\gamma : H^{1,i}(X) \xrightarrow{\sim} H^{1,1}(X)$, and $[\zeta]^{i-1} \cup \gamma(x) = x$.*

Every motivic cohomology operation on $H^{1,i}$ is uniquely a sum of the operations $x \mapsto c(\gamma x)^\varepsilon \beta(\gamma x)^m$, where $c \in H^{,*}(k)$, $0 \leq \varepsilon \leq 1$, and $m \geq 0$.*

Proof. By Theorem 6.2, $H^{1,i}(X) \cong H_{\text{et}}^1(X, \mu_\ell^{\otimes i})$ for all $i > 0$. Since multiplication by $[\zeta^{i-1}]$ is an isomorphism between $H_{\text{et}}^n(X, \mu_\ell)$ and $H_{\text{et}}^n(X, \mu_\ell^{\otimes i})$, its inverse isomorphism γ is natural. Via γ , operations on $H^{1,i}$ correspond to operations on $H^{1,1}$, which are described in Proposition 10.1. \square

For example if $i \geq 2$ and $\eta \in H_{\text{et}}^1(k, \mu_\ell^{\otimes 2-i})$, then the operation $H^{1,i} \rightarrow H^{2,2}$ of Example 11.1 is the operation $x \mapsto c(\gamma x)$ of Proposition 11.2, where $c = \eta \cup [\zeta]^{i-1}$.

Remark 11.2.1. If $c \in H^{s,j}(k)$, then $\phi(x) = c(\gamma x)^\varepsilon (\beta \gamma x)^m$ is a cohomology operation of bidegree $(s + \varepsilon + 2m - 1, j + \varepsilon + m - i)$. In particular γ is a cohomology operation of bidegree $(0, 1 - i)$.

Galois descent. We now consider the situation in which $\mu_\ell \not\subset k$. Clearly, not all cohomology operations defined over $k(\zeta)$ are defined over k . However, some of these operations do descend, such as those in Example 11.1.

It is convenient to consider the étale cohomology of k as being bigraded, by integers $n \geq 0$ and $i \in \mathbb{Z}$, with $H_{\text{ét}}^n(k, \mu_\ell^{\otimes i})$ in bidegree (n, i) . Thus the motivic cohomology ring $H^{*,*}(k)$ is a bigraded subring of $H_{\text{ét}}^*(k, \mu_\ell^{\otimes *})$.

Definition 11.3. For each integer b , let $\zeta^{-b} H^{*,*}(k)$ denote the direct sum of all $H_{\text{ét}}^s(k, \mu_\ell^{\otimes t})$ with $0 \leq s \leq t+b$. This is a bigraded $H^{*,*}(k)$ -submodule of $H_{\text{ét}}^*(k, \mu_\ell^{\otimes *})$. It is a cyclic module if and only if $\zeta^b \in k$, when it is the $H^{*,*}(k)$ -submodule generated by $[\zeta^b] \in H_{\text{ét}}^0(k, \mu_\ell^{\otimes -b})$.

Theorem 11.4. Fix an integer $i \geq 2$. Then the ring of cohomology operations on $H^{1,i}$ is the direct sum of copies of $\zeta^{-b} H^{*,*}(k)$, $b = (i-1)(\varepsilon+m)$, over integers $m \geq 0$, and $\varepsilon \in \{0, 1\}$. If $0 \leq s \leq t+b$, the operation corresponding to $c \in H_{\text{ét}}^s(k, \mu_\ell^{\otimes t})$, m , and ε sends $H^{1,i}(X)$ to $H^{s+\varepsilon+2m, t+b+\varepsilon+m}(X)$:

$$\phi(\zeta^{i-1} \cup y) = (\zeta^b \cup c)y^\varepsilon \beta(y)^m.$$

Proof. Let G denote the Galois group of $k(\zeta)/k$. Since $H^{*,*}(X)$ is the G -invariant summand of $H^{*,*}(X(\zeta))$, a motivic operation $H^{1,i}(X) \rightarrow H^{*,*}(X)$ is the same thing as a G -invariant operation $H^{1,i}(X) \rightarrow H^{*,*}(X(\zeta))$. Given $x \in H^{1,i}(X)$, there is a unique $y \in H^{1,1}(X(\zeta))$ so that $x = [\zeta]^{i-1} \cup y$, where $[\zeta] \in H^{0,1}(k(\zeta))$. By Proposition 11.2, we are reduced to determining when G acts trivially on $c'y^\varepsilon(\beta y)^m$. Since $y^\varepsilon(\beta y)^m$ is in the summand of $H^{*,*}(k(\zeta))$ which is isotypical for $\mu_\ell^{\otimes -b}$, this holds if and only if c' is in the summand of $H^{s,j}(k(\zeta))$ which is isotypical for $\mu_\ell^{\otimes b}$. By (3.2), there is a unique $c \in H^{s,j-b}(k)$ so that $c' = [\zeta]^b \cup c$. \square

Example 11.5 ($b = 1$). An element c in $H_{\text{ét}}^1(k, \mathbb{F}_\ell) = \text{Hom}(\text{Gal}(\bar{k}/k), \mathbb{F}_\ell)$ determines operations $C : H^{1,2}(X) \rightarrow H^{2,1}(X)$ and $\phi : H^{1,2}(X) \rightarrow H^{3,2}(X)$. If $y \in H^{1,1}(X(\zeta))$ is such that $x = [\zeta] \cup y$, then, regarding ζc as an element of $H^{1,1}(k(\zeta))$, we have $C(x) = (\zeta c)y$ and $\phi(x) = (\zeta c)\beta(y)$. Of course, we can identify C with the map $H_{\text{ét}}^1(X, \mu_\ell) \xrightarrow{\cup c} H_{\text{ét}}^2(X, \mu_\ell)$.

An element t in $H_{\text{ét}}^2(k, \mu_\ell)$ (the ℓ -torsion subgroup of the Brauer group of k) determines operations $H^{1,2}(X) \rightarrow H^{3,3}(X)$ and $H^{1,2}(X) \rightarrow H^{4,3}(X)$. Writing $x = [\zeta] \cup y$ in $H^{1,2}(X(\zeta))$, the operations followed by the inclusion $H^{*,*}(X) \subset H^{*,*}(X(\zeta))$ send x to $([\zeta] \cup t)y$ and $([\zeta] \cup t)\beta(y)$, respectively. As mentioned in the introduction, we can identify the first operation with $H_{\text{ét}}^1(X, \mu_\ell^{\otimes 2}) \xrightarrow{\cup t} H_{\text{ét}}^3(X, \mu_\ell^{\otimes 3})$.

12. CONJECTURAL MATTER

In the preceding two sections we have classified motivic cohomology operations on $H^{n,i}$ when $n = 1$ or $i = 1$. We have also classified operations whose targets lie inside the “étale zone” where $n \leq i$. We know little about the intermediate zone where $i < n < 2i$. In this section we make some guesses about operations in the “topological zone” where $n \geq 2i$.

Example 12.1. There are many operations defined on $H^{n,2}$, $n \geq 2$. Let us compare Voevodsky’s operation P_V^1 (landing in $H^{n+2\ell-2, \ell+1}$) with our operation P^1 (landing in $H^{n+2\ell-2, 2\ell}$). Thus P^1 has the same bidegree as $[\zeta]^{\ell-1} P_V^1$, where $[\zeta] \in H^{0,1}(k)$. If $n \geq 4$, we have $P^1 = [\zeta]^{\ell-1} P_V^1$ by Corollary 8.9. If $n = 2$ we also have $P^1 = [\zeta]^{\ell-1} P_V^1$ because they induce the same étale operation (P^1) from

$H^{2,2}(X) \cong H^2_{\text{et}}(X, \mu_{\ell}^{\otimes 2})$ to $H^{2\ell, 2\ell}(X) \cong H^{2\ell}_{\text{et}}(X, \mu_{\ell}^{\otimes 2\ell})$. We do not know if P^1 and $[\zeta]^{\ell-1} P_V^1$ agree on $H^{3,2}$.

Suppose that ϕ is a motivic cohomology operation on $H^{n,i}$ where $n \geq 2i$. Passing to étale cohomology sends ϕ to an étale operation, which by Theorem 3.5 is a polynomial in the étale operations P^I . By Proposition 8.8, some multiple of the Bott element b sends ϕ to operations $b^N \phi$ which are in the subalgebra generated by the motivic operations P^I defined in Definition 6.5. It remains to determine what those powers are.

The following result of Voevodsky [V1, 3.6–7] shows that all nontrivial operations in the topological zone increase n .

Lemma 12.2 (Voevodsky). *There are no motivic cohomology operations from $H^{2i,i}$ to $H^{n,j}$ when $j < i$ or when $i = j$ and $(n,j) \neq (2i,i)$. The module of motivic cohomology operations from $H^{2i,i}$ to $H^{*,i}$ is isomorphic to \mathbb{F}_{ℓ} , on the identity.*

Conjecture 12.3. *Assume that k contains all primitive ℓ th roots of unity and that $n \geq 2i$. Then the module of all motivic cohomology operations on $H^{n,i}(-, \mathbb{F}_{\ell})$ is the tensor product of $H^{*,*}$ and a free graded polynomial algebra over \mathbb{F}_{ℓ} with generators all $P^I P_V^J$, where $I = (\epsilon_0, s_1, \epsilon_1, \dots, s_k, \epsilon_k)$, $J = (s_{k+1}, \epsilon_{k+1}, \dots, s_m, \epsilon_m)$ subject to the conditions that (a) the concatenation IJ is admissible with excess $e(IJ)$ either < 4 or else $\epsilon_0 = 1$ and $e(IJ) = 4$ and (b) for all $j > k$, $s_j < i + (\ell - 1) \sum_{j+1}^m s_i$.*

For $(n, i) = (4, 2)$ this conjecture implies that among the polynomial generators for the motivic operations on $H^{4,2}$ we find $P^{\ell^2+\ell+1} \beta P_V^{\ell+1} \beta P_V^1 \beta$. If $\ell = 2$, we may rewrite these operations as $Sq^{14} Sq_V^7 Sq_V^3 Sq_V^1$; compare with [V3, 3.57].

Lemma 12.4. *If Conjecture 12.3 holds for $H^{2i,i}$, then it holds for all $H^{n,i}$ with $n \geq 2i$.*

Proof. We consider the Leray spectral sequence (9.3) for $G = K(\mathbb{F}_{\ell}(i), n)$ and $K = B_{\bullet} G = K(\mathbb{F}_{\ell}(i), n+1)$ when $n \geq 2i$. By induction, $H^{*,*}(G)$ is a polynomial algebra over $H^{*,*}$ with an ℓ -simple system $\{x_i\}$ of generators. By [V3, 3.28], $\mathbb{F}_{\ell, \text{tr}}(G)$ is a split proper Tate motive, so the Künneth condition of Proposition 9.4 holds, and Borel's Theorem 9.2 implies that $H^{*,*}(K)$ is the tensor product of $H^{*,*}$ and a free graded-commutative \mathbb{F}_{ℓ} -algebra on generators $y_i = \tau(x_i)$ and, when $\deg(x_j)$ is even and $\ell > 2$, $z_j = \tau(x_j^{\ell-1} \otimes y_j)$.

We now use the fact that the transgression commutes with any (S^1) -stable cohomology operation, such as P_V^J ; see [McC, 6.5]. Since the tautological element ι_n of $H^{n,i}(G)$ transgresses to the tautological element ι_{n+1} of $H^{n+1,i}(K)$, the generator $x_j = P^I P_V^J(\iota_n)$ transgresses to $y_j = P^I P_V^J(\iota_{n+1})$ by Kudo's Theorem 9.5. This finishes the proof for $\ell = 2$.

If ℓ is odd and $x_j = P^I P_V^J(\iota_n)$ has degree $2a$, the transgression z_j of $x_j^{\ell-1} \otimes y_j$ is $-\beta P^a P^I P_V^J(\iota_{n+1})$ by Kudo's Theorem 9.5(3). This finishes the proof for ℓ odd. \square

ACKNOWLEDGMENTS

The authors would like to thank Peter May for his encouragement and for providing us with a copy of [SE-err]. They would also like to thank a referee for pointing out the need for a dg approach to motivic operations, which we carry out in Section 6, and for insisting on a clear presentation of Q^0 and a careful treatment of the Bockstein.

REFERENCES

- [Br] Lawrence Breen, *Extensions du groupe additif* (French), Inst. Hautes Études Sci. Publ. Math. **48** (1978), 39–125. MR516914
- [BJ] P. Brosnan and R. Joshua, *Cohomology operations in motivic, étale and de Rham-Witt cohomology*, preprint, 2007.
- [BJ1] Patrick Brosnan and Roy Joshua, *Comparison of motivic and simplicial operations in mod-l-motivic and étale cohomology*, Feynman amplitudes, periods and motives, Contemp. Math., vol. 648, Amer. Math. Soc., Providence, RI, 2015, pp. 29–56, DOI 10.1090/conm/648/12997. MR3415409
- [C] Séminaire Henri Cartan de l’Ecole Normale Supérieure, 1954/1955. *Algèbres d’Eilenberg-MacLane et homotopie* (French), Secrétariat mathématique, 11 rue Pierre Curie, Paris, 1955. MR0087934
- [C1] Henri Cartan, *Sur les groupes d’Eilenberg-MacLane. II* (French), Proc. Nat. Acad. Sci. U. S. A. **40** (1954), 704–707. MR0065161
- [D] Pierre Deligne, *Théorie de Hodge. III* (French), Inst. Hautes Études Sci. Publ. Math. **44** (1974), 5–77. MR0498552
- [D-4.5] A. Grothendieck and P. Deligne, *La classe de cohomologie associée à un cycle* (French), Cohomologie étale, Lecture Notes in Math., vol. 569, Springer, Berlin, 1977, pp. 129–153. MR3727436
- [EM] Samuel Eilenberg and Saunders MacLane, *On the groups $H(\Pi, n)$. III*, Ann. of Math. (2) **60** (1954), 513–557, DOI 10.2307/1969849. MR0065163
- [E] D. B. A. Epstein, *Steenrod operations in homological algebra*, Invent. Math. **1** (1966), 152–208, DOI 10.1007/BF01389726. MR0199240
- [F] Eric M. Friedlander, *Étale homotopy of simplicial schemes*, Annals of Mathematics Studies, vol. 104, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1982. MR676809
- [HRD] Robin Hartshorne, *Residues and duality*, Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20, Springer-Verlag, Berlin-New York, 1966. MR0222093
- [HS] V. A. Hinich and V. V. Schechtman, *On homotopy limit of homotopy algebras*, K-theory, arithmetic and geometry (Moscow, 1984), Lecture Notes in Math., vol. 1289, Springer, Berlin, 1987, pp. 240–264, DOI 10.1007/BFb0078370. MR923138
- [HW] C. Haesemeyer and C. Weibel, *The norm residue theorem in motivic cohomology*, Princeton Univ. Press, 2019. Available at math.rutgers.edu/~weibel.
- [J] J. F. Jardine, *Steenrod operations in the cohomology of simplicial presheaves*, Algebraic K-theory: connections with geometry and topology (Lake Louise, AB, 1987), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. 279, Kluwer Acad. Publ., Dordrecht, 1989, pp. 103–116, DOI 10.1007/978-94-009-2399-7_5. MR1045847
- [J1] J. F. Jardine, *Higher spinor classes*, Mem. Amer. Math. Soc. **110** (1994), no. 528, vi+88, DOI 10.1090/memo/0528. MR1211372
- [Jo] R. Joshua, *Motivic E_∞ algebras and the motivic DGA*, preprint, 2012. Available at <http://www.math.ohio-state.edu/~joshua/pub.html>.
- [KM] Igor Kríž and J. P. May, *Operads, algebras, modules and motives* (English, with English and French summaries), Astérisque **233** (1995), iv+145pp. MR1361938
- [L] Marc Levine, *Inverting the motivic Bott element*, K-Theory **19** (2000), no. 1, 1–28, DOI 10.1023/A:1007874218371. MR1740880
- [Mass] W. S. Massey, *Products in exact couples*, Ann. of Math. (2) **59** (1954), 558–569, DOI 10.2307/1969719. MR0060829
- [M] J. Peter May, *A general algebraic approach to Steenrod operations*, The Steenrod Algebra and its Applications (Proc. Conf. to Celebrate N. E. Steenrod’s Sixtieth Birthday, Battelle Memorial Inst., Columbus, Ohio, 1970), Lecture Notes in Mathematics, Vol. 168, Springer, Berlin, 1970, pp. 153–231. MR0281196
- [M1] J. P. May, *Operads and sheaf cohomology*, preprint, 2004. Available at <http://www.math.uchicago.edu/~may/PAPERS/Esheaf.pdf>.
- [McC] John McCleary, *A user’s guide to spectral sequences*, 2nd ed., Cambridge Studies in Advanced Mathematics, vol. 58, Cambridge University Press, Cambridge, 2001. MR1793722

- [Milne] James S. Milne, *Étale cohomology*, Princeton Mathematical Series, vol. 33, Princeton University Press, Princeton, N.J., 1980. MR559531
- [MV] Fabien Morel and Vladimir Voevodsky, A^1 -homotopy theory of schemes, Inst. Hautes Études Sci. Publ. Math. **90** (1999), 45–143 (2001). MR1813224
- [MVW] Carlo Mazza, Vladimir Voevodsky, and Charles Weibel, *Lecture notes on motivic cohomology*, Clay Mathematics Monographs, vol. 2, American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2006. MR2242284
- [R] Michèle Raynaud, *Modules projectifs universels* (French), Invent. Math. **6** (1968), 1–26, DOI 10.1007/BF01389829. MR0236164
- [SE] N. E. Steenrod, *Cohomology operations*, Lectures by N. E. Steenrod written and revised by D. B. A. Epstein, Annals of Mathematics Studies, No. 50, Princeton University Press, Princeton, N.J., 1962. MR0145525
- [SE-err] N. Steenrod and D. Epstein, Errata for *Cohomology Operations*, 5 pp., circa 1965.
- [S50] Jean-Pierre Serre, *Cohomologie modulo 2 des complexes d'Eilenberg-MacLane* (French), Comment. Math. Helv. **27** (1953), 198–232, DOI 10.1007/BF02564562. MR0060234
- [SV] Andrei Suslin and Vladimir Voevodsky, *Bloch-Kato conjecture and motivic cohomology with finite coefficients*, The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), NATO Sci. Ser. C Math. Phys. Sci., vol. 548, Kluwer Acad. Publ., Dordrecht, 2000, pp. 117–189. MR1744945
- [Ver] J.-L. Verdier, *Cohomologie dans les topos*, SGA 4, Lecture Notes in Math. 270, Springer, Berlin, 1972, pp. 1–82.
- [V1] Vladimir Voevodsky, *Reduced power operations in motivic cohomology*, Publ. Math. Inst. Hautes Études Sci. **98** (2003), 1–57, DOI 10.1007/s10240-003-0009-z. MR2031198
- [V2] Vladimir Voevodsky, *Motivic cohomology with $\mathbb{Z}/2$ -coefficients*, Publ. Math. Inst. Hautes Études Sci. **98** (2003), 59–104, DOI 10.1007/s10240-003-0010-6. MR2031199
- [V3] Vladimir Voevodsky, *Motivic Eilenberg-MacLane spaces*, Publ. Math. Inst. Hautes Études Sci. **112** (2010), 1–99, DOI 10.1007/s10240-010-0024-9. MR2737977
- [V4] Vladimir Voevodsky, *On motivic cohomology with \mathbb{Z}/l -coefficients*, Ann. of Math. (2) **174** (2011), no. 1, 401–438, DOI 10.4007/annals.2011.174.1.11. MR2811603
- [WH] Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994. MR1269324
- [W] C. Weibel, *The norm residue isomorphism theorem*, J. Topol. **2** (2009), no. 2, 346–372, DOI 10.1112/jtopol/jtp013. MR2529300

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, LEXINGTON, KENTUCKY 40506
Email address: bertguillou@uky.edu

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY 08901
Email address: weibel@math.rutgers.edu