

NONCONFORMING VIRTUAL ELEMENT METHOD FOR $2m$ TH ORDER PARTIAL DIFFERENTIAL EQUATIONS IN $\mathbb{R}^n$

LONG CHEN AND XUEHAI HUANG

ABSTRACT. A unified construction of the H^m -nonconforming virtual elements of any order k is developed on any shape of polytope in $\mathbb{R}^n$ with constraints $m \leq n$ and $k \geq m$. As a vital tool in the construction, a generalized Green's identity for H^m inner product is derived. The H^m -nonconforming virtual element methods are then used to approximate solutions of the m -harmonic equation. After establishing a bound on the jump related to the weak continuity, the optimal error estimate of the canonical interpolation, and the norm equivalence of the stabilization term, the optimal error estimates are derived for the H^m -nonconforming virtual element methods.

1. INTRODUCTION

We intend to construct H^m -nonconforming virtual elements of order $k \in \mathbb{N}$ on a very general polytope $K \subset \mathbb{R}^n$ in any dimension and any order with constraints $m \leq n$ and $k \geq m$. Since an m th order derivative of polynomial degree $m - 1$ or less would be zero, the constraint $k \geq m$ is required to ensure that the virtual element spaces possess meaningful approximation in the H^m -seminorm. Due to a technical reason, our attention is restricted to the case $m \leq n$ in this paper, and the case $m > n$ is postponed for future works. The virtual element was described as a generalization of the finite element on a general polytope in [12, 13], thus it is helpful to recall the definition of the finite element first.

A finite element on K was defined as a triple $(K, V_K, \mathcal{N}_K)$ in [23], where V_K is the finite-dimensional space of shape functions and $\mathcal{N}_K$ is the set of degrees of freedom (d.o.f.). The set $\mathcal{N}_K$ forms a basis of $(V_K)'$ the dual space of the space of shape functions. The shape functions of the finite element are usually polynomials, and their basis functions being dual to the degrees of freedom $\mathcal{N}_K$ have to be explicitly constructed for the implementation, which is painful for high order cases (either k, m , or n is large).

We can also represent the virtual element as a triple $(K, \mathcal{N}_K, V_K)$. Here we reorder V_K and $\mathcal{N}_K$ to emphasize that the set of the degrees of freedom $\mathcal{N}_K$ is crucial in the construction of the virtual element and that the space of shape functions V_K is virtual. Indeed, after having the degrees of freedom $\mathcal{N}_K$, we may attach different spaces. The space of shape functions V_K is only required to include all polynomials

Received by the editor November 7, 2018, and, in revised form, November 9, 2018, July 7, 2019, and September 9, 2019.

2010 *Mathematics Subject Classification.* Primary 65N30, 65N12, 65N22.

The first author was supported by NSF DMS-1913080.

The second author was supported by the National Natural Science Foundation of China Project 11771338, the Fundamental Research Funds for the Central Universities 2019110066, and Zhejiang Provincial Natural Science Foundation of China Project LY17A010010.

©2019 American Mathematical Society

of the total degree up to k for the approximation property. Different from the finite element, one advantage of the virtual element is that the basis functions of V_K are not explicitly required in the implementation. When forming the linear system of the virtual element method, the computation of all the to-be-required quantities can be transferred to the computation using the degrees of freedom.

Construction of H^m -conforming or nonconforming elements has been an active topic in the field of the finite element methods in recent years. Some H^m -conforming finite elements with polynomial shape functions were designed on the simplices in [4, 7, 15, 35, 41] and on the hyperrectangles in [27, 29, 42]. Recently an H^m -conforming virtual element for polyharmonic problems with arbitrary m in two dimensions was introduced and studied in [6]. For general m , nonconforming elements on the simplices are easier to construct than conforming ones. In [36, 37], Wang and Xu constructed the minimal H^m -nonconforming elements in any dimension with constraint $m \leq n$. Recently Wu and Xu extended these minimal H^m -nonconforming elements to $m = n + 1$ by enriching the space of shape functions with bubble functions in [40], and to arbitrary m and n by using the interior penalty technique in [39]. In two dimensions, Hu and Zhang designed the H^m -nonconforming elements on the triangle for any m in [30]. On the other hand, the H^2 -conforming virtual element, the C^0 -type H^2 -nonconforming virtual element, and the fully H^2 -nonconforming virtual element on the polygon with any shape in two dimensions were developed in [19], [43], and [5, 44], respectively. In [38], a nonconforming Crouzeix–Raviart type, i.e., H^1 -nonconforming finite element, was advanced on the polygon.

Although the H^m -conforming virtual element has been devised for $n = 2$ in [6] for arbitrary m , generalization to dimension $n > 2$ seems nontrivial. As we show in this paper, the H^m -nonconforming virtual element can be constructed in a universal way for all $n \geq m$ and allows unified error analysis.

We shall construct the H^m -nonconforming virtual element in any order on the polytope with any shape in any dimension (with constraints $k \geq m$ and $m \leq n$). The vital tool is the following generalized Green's identity for the H^m space:

$$(1.1) \quad (\nabla^m u, \nabla^m v)_K = ((-\Delta)^m u, v)_K + \sum_{j=1}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} \left(D_{F,\alpha}^{2m-j-|\alpha|}(u), \frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} \right)_F,$$

which is proved by the mathematical induction and integration by parts. Here $\mathcal{F}^j(K)$ is the set of all $(n-j)$ -dimensional faces of the polytope K , A_j the set consisting of all n -dimensional multi-indexes $\alpha = (\alpha_1, \dots, \alpha_n)$ with $\alpha_{j+1} = \dots = \alpha_n = 0$, $D_{F,\alpha}^{2m-j-|\alpha|}(u)$ some $(2m-j-|\alpha|)$ th order derivatives of u on F , and $\frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha}$ the multi-indexed normal derivatives on F .

Imagining u in the Green's identity (1.1) as a polynomial of degree k temporarily, we acquire the degrees of freedom $\mathcal{N}_k(K)$ from the right-hand side of the Green's identity (1.1), and the space $V_k(K)$ of shape functions is defined inherently by requiring the first terms in the inner product to be in polynomial spaces. Namely, the right-hand side of (1.1) provides a natural duality of $V_k(K)$ and $\mathcal{N}_k(K)$. As a result we construct the fully H^m -nonconforming virtual element $(K, \mathcal{N}_k(K), V_k(K))$ completely based on the Green's identity (1.1). If K is a simplex and $k = m$, the virtual element $(K, \mathcal{N}_k(K), V_k(K))$ is reduced to the nonconforming finite element

in [37], and hence we generalize the nonconforming finite element in [37] to high order $k > m$ and arbitrary polytopes. In two dimensions, we also recover the fully H^2 -nonconforming virtual element in [5, 44].

After introducing the local H^m projection Π^K and a stabilization term using d.o.f., we propose H^m -nonconforming virtual element methods for solving the m -harmonic equation. We assume the mesh $\mathcal{T}_h$ admits a virtual quasi-uniform triangulation, and each element in $\mathcal{T}_h$ is star-shaped. A bound on the jump $[\![\nabla_h^s v_h]\!]$ is derived using the weak continuity and the trace inequality, with which we show the discrete Poincaré inequality for the global virtual element space. The optimal error estimate of the canonical interpolation $I_h u$ is achieved after establishing a Galerkin orthogonality of $u - I_h u$. By employing the bubble function technique which was frequently used in proving the efficiency of the a posteriori error estimators, the inverse inequality for polynomials, the generalized Green's identity, and the trace inequality, we acquire the norm equivalence of the standard stabilization using l^2 inner products of degree of freedoms on $\ker(\Pi^K)$. The optimal error estimates are derived for the H^m -nonconforming virtual element methods by further estimating the consistency error.

The shape functions of the virtual element spaces are not explicitly known; in particular, the output of the method is a vector of degrees of freedom and not an explicit function. In order to explicitly represent the solution, one employs some suitable polynomial projector, which is typically piecewise defined and discontinuous over the polytopal decomposition. However, since the degrees of freedom in the interior of each element for the virtual elements can be eliminated by the static condensation, similarly as the hybridizable discontinuous Galerkin methods [31], the virtual element methods possess fewer globally decoupled degrees of freedom than the usual discontinuous Galerkin methods. Furthermore, the nonconforming virtual element can be constructed in a universal way which allows unified error analysis and is employed for theoretical purposes, independently of the way one wants to represent the solution.

The rest of this paper is organized as follows. In Section 2, we present some notation and the construction of the fully H^1 - and H^2 -nonconforming virtual elements. The general fully H^m -nonconforming virtual element is designed in Section 3. The corresponding H^m -nonconforming virtual element method and its error estimate are shown in Sections 4 and 5, respectively. A conclusion is given in Section 6. Finally, we prove the norm equivalence in Appendix A and give a remark on the implementation in Appendix B.

2. PRELIMINARIES

2.1. Notation. Assume that $\Omega \subset \mathbb{R}^n$ ($n \geq 2$) is a bounded polytope. For any nonnegative integer r and $1 \leq \ell \leq n$, denote the set of r -tensor spaces over $\mathbb{R}^\ell$ by $\mathbb{T}_\ell(r) := (\mathbb{R}^\ell)^r = \prod_{j=1}^r \mathbb{R}^\ell$. Given a bounded domain $K \subset \mathbb{R}^n$ and a nonnegative integer k , let $H^k(K; \mathbb{T}_\ell(r))$ be the usual Sobolev space of functions over K taking values in the tensor space $\mathbb{T}_\ell(r)$. The corresponding norm and seminorm are denoted, respectively, by $\|\cdot\|_{k,K}$ and $|\cdot|_{k,K}$. It is customary to rewrite $H^k(K; \mathbb{T}_\ell(0))$ as $H^k(K)$. For any $F \subset \partial K$, denote by $\nu_{K,F}$ the unit outward normal to ∂K . Without causing any confusion, for simplicity we will abbreviate $\nu_{K,F}$ as ν . Define $H_0^k(K)$ as the closure of $C_0^\infty(K)$ with respect to the norm $\|\cdot\|_{k,K}$, i.e.

(cf. [1, Theorem 5.37]),

$$H_0^k(K) := \left\{ v \in H^k(K) : v = \frac{\partial v}{\partial \nu} = \cdots = \frac{\partial^{k-1} v}{\partial \nu^{k-1}} = 0 \text{ on } \partial K \right\},$$

and define $H_0^1(K; \mathbb{T}_\ell(r))$ in a similar way. Let $(\cdot, \cdot)_K$ be the standard inner product on $L^2(K; \mathbb{T}_\ell(r))$. If K is Ω , we abbreviate $\|\cdot\|_{k,K}$, $|\cdot|_{k,K}$ and $(\cdot, \cdot)_K$ by $\|\cdot\|_k$, $|\cdot|_k$ and $(\cdot, \cdot)$, respectively. The notation $\mathbb{P}_k(K)$ stands for the set of all polynomials over K with the total degree no more than k , and denote by $\mathbb{P}_k(K; \mathbb{T}_\ell(r))$ the tensorial version space of $\mathbb{P}_k(K)$. Let Q_k^K be the L^2 -orthogonal projection onto $\mathbb{P}_k(K; \mathbb{T}_\ell(r))$.

For an n -dimensional multi-index $\alpha = (\alpha_1, \dots, \alpha_n)$ with $\alpha_i \in \mathbb{Z}^+ \cup \{0\}$, define $|\alpha| := \sum_{i=1}^n \alpha_i$. For $0 \leq \ell \leq n$, let A_ℓ be the set consisting of all multi-indexes α with $\sum_{i=\ell+1}^n \alpha_i = 0$, i.e., the nonzero index only exists for $1 \leq i \leq \ell$. For any nonnegative integer k , define the scaled monomial $\mathbb{M}_k(K)$ on an ℓ -dimensional domain K ,

$$\mathbb{M}_k(K) := \left\{ \left(\frac{\mathbf{x} - \mathbf{x}_K}{h_K} \right)^\alpha, \alpha \in A_\ell, |\alpha| \leq k \right\},$$

where h_K is the diameter of K and $\mathbf{x}_K$ is the centroid of K . Also, $\mathbb{M}_k(K) := \emptyset$ whenever $k < 0$.

Let $\{\mathcal{T}_h\}$ be a regular family of partitions of Ω into nonoverlapping simple polytopal elements with $h := \max_{K \in \mathcal{T}_h} h_K$. Let $\mathcal{F}_h^r$ be the set of all $(n-r)$ -dimensional faces of the partition $\mathcal{T}_h$ for $r = 1, 2, \dots, n$, and its boundary part

$$\mathcal{F}_h^{r,\partial} := \{F \in \mathcal{F}_h^r : F \subset \partial\Omega\},$$

and interior part $\mathcal{F}_h^{r,i} := \mathcal{F}_h^r \setminus \mathcal{F}_h^{r,\partial}$. Moreover, we set for each $K \in \mathcal{T}_h$

$$\mathcal{F}^r(K) := \{F \in \mathcal{F}_h^r : F \subset \partial K\}.$$

The superscript r in $\mathcal{F}_h^r$ represents the co-dimension of an $(n-r)$ -dimensional face F . As we shall show later the degree of freedom will be associated to the r normal vectors of F . Similarly, for $F \in \mathcal{F}_h^r$ and $j = 0, 1, \dots, n-r$ with $r = 1, 2, \dots, n$, we define

$$\mathcal{F}^j(F) := \{e \in \mathcal{F}_h^{r+j} : e \subset \overline{F}\}.$$

Here j is the co-dimension relative to the face F . Apparently $\mathcal{F}^0(F) = F$.

For any $F \in \mathcal{F}_h^r$, let $\nu_{F,1}, \dots, \nu_{F,r}$ be its mutually perpendicular unit normal vectors, and define the surface gradient on F as

$$(2.1) \quad \nabla_F v := \nabla v - \sum_{i=1}^r \frac{\partial v}{\partial \nu_{F,i}} \nu_{F,i},$$

namely the projection of ∇v to the face F , which is independent of the choice of the normal vectors. When v is a tensor, the surface gradient $\nabla_F v$ is defined elementwisely in convention, which is a one-order higher tensor. And denote by div_F the corresponding surface divergence. For any $F \in \mathcal{F}_h^r$ and $\alpha \in A_r$ for $r = 1, \dots, m$, set

$$\frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} := \frac{\partial^{|\alpha|} v}{\partial \nu_{F,1}^{\alpha_1} \cdots \partial \nu_{F,r}^{\alpha_r}}.$$

For any $(n-2)$ -dimensional face $e \in \mathcal{F}_h^2$, denote

$$\partial^{-1} e := \{F \in \mathcal{F}_h^1 : e \subset \partial F\}.$$

Similarly, for any $(n-1)$ -dimensional face $F \in \mathcal{F}_h^1$, let

$$\partial^{-1}F := \{K \in \mathcal{T}_h : F \in \mathcal{F}^1(K)\}.$$

For nonnegative integers m and k , let

$$H^m(\mathcal{T}_h) := \{v \in L^2(\Omega) : v|_K \in H^m(K) \text{ for each } K \in \mathcal{T}_h\},$$

$$\mathbb{P}_k(\mathcal{T}_h) := \{v \in L^2(\Omega) : v|_K \in \mathbb{P}_k(K) \text{ for each } K \in \mathcal{T}_h\}.$$

For a function $v \in H^m(\mathcal{T}_h)$, equip the usual broken H^m -type norm and seminorm

$$\|v\|_{m,h} := \left(\sum_{K \in \mathcal{T}_h} \|v\|_{m,K}^2 \right)^{1/2}, \quad |v|_{m,h} := \left(\sum_{K \in \mathcal{T}_h} |v|_{m,K}^2 \right)^{1/2}.$$

We introduce jumps on $(n-1)$ -dimensional faces. Consider two adjacent elements K^+ and K^- sharing an interior $(n-1)$ -dimensional face F . Denote by ν^+ and ν^- the unit outward normals to the common face F of the elements K^+ and K^- , respectively. For a scalar-valued or tensor-valued function v , write $v^+ := v|_{K^+}$ and $v^- := v|_{K^-}$. Then define the jump on F as follows:

$$[[v]] := v^+ \nu_{F,1} \cdot \nu^+ + v^- \nu_{F,1} \cdot \nu^-.$$

On a face F lying on the boundary $\partial\Omega$, the above term is defined by $[[v]] := v \nu_{F,1} \cdot \nu$.

Throughout this paper, we also use “ $\lesssim \dots$ ” to mean that “ $\leq C \dots$ ”, where C is a generic positive constant independent of mesh size h , but may depend on the chunkiness parameter of the polytope, the degree of polynomials k , the order of differentiation m , and the dimension of space n , which may take different values at different appearances. Also, $A \asymp B$ means $A \lesssim B$ and $B \lesssim A$. Hereafter, we always assume $k \geq m$.

We summarize important notation in Tables 1 and 2 below.

TABLE 1. Notation of the mesh, elements, and faces.

m	order of differentiation H^m	n	dimension of space $\mathbb{R}^n$	$m \leq n, k \geq m$
k	degree of polynomial $\mathbb{P}_k$	r	co-dimension of a face	$0 \leq r \leq n$
$\mathcal{T}_h$	a mesh of Ω	K	a polytope element	$K \in \mathcal{T}_h$
$\mathcal{F}_h^r$	$(n-r)$ -dimensional face	F	a typical face	$F \in \mathcal{F}_h^r$
$\partial^{-1}e$	all faces surrounding e	$\partial^{-1}F$	elements containing F	$e \in \mathcal{F}_h^2, F \in \mathcal{F}_h^1$

TABLE 2. Notation for differentiation

$\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$	an n -dimensional multi-index
A_r set of multi-index	$\alpha = (\alpha_1, \dots, \alpha_r, 0, \dots, 0)$ for $\alpha \in A_r$
$\nu_{F,1}, \dots, \nu_{F,r}$	r linearly independent unit normal vectors for $F \in \mathcal{F}_h^r$
$\nabla_F v := \nabla v - \sum_{i=1}^r \frac{\partial v}{\partial \nu_{F,i}} \nu_{F,i}$	surface gradient on F
$D_{F,\alpha}^j(v)$	a j th order derivative of v on F
$\frac{\partial^{ \alpha } v}{\partial \nu_F^\alpha} := \frac{\partial^{ \alpha } v}{\partial \nu_{F,1}^{\alpha_1} \cdots \partial \nu_{F,r}^{\alpha_r}}$	a multi-indexed normal derivative on F

2.2. H^1 -nonconforming virtual element. To drive the H^m -nonconforming virtual element in a unified framework, we first revisit the simplest case for the purpose of discovering the underlying mechanism.

Taking any $K \in \mathcal{T}_h$, let $u \in H^2(K)$, and let $v \in H^1(K)$. Applying the integration by parts, it holds that

$$(2.2) \quad (\nabla u, \nabla v)_K = -(\Delta u, v)_K + \sum_{F \in \mathcal{F}^1(K)} \left(\frac{\partial u}{\partial \nu_{K,F}}, v \right)_F.$$

Imaging $u \in \mathbb{P}_k(K)$, we are inspired by the Green's identity (2.2) to advance the following local degrees of freedom (dofs) $\mathcal{N}_k(K)$ of the H^1 nonconforming virtual element:

$$(2.3) \quad \frac{1}{|F|} (v, q)_F \quad \forall q \in \mathbb{M}_{k-1}(F) \text{ on each } F \in \mathcal{F}^1(K),$$

$$(2.4) \quad \frac{1}{|K|} (v, q)_K \quad \forall q \in \mathbb{M}_{k-2}(K).$$

The local space of the H^1 -nonconforming virtual element is

$$V_k(K) := \left\{ u \in H^1(K) : \Delta u \in \mathbb{P}_{k-2}(K), \frac{\partial u}{\partial \nu_{K,F}}|_F \in \mathbb{P}_{k-1}(F) \quad \forall F \in \mathcal{F}^1(K) \right\}$$

for $k \geq 1$. This is the H^1 -nonconforming virtual element constructed in [8]; see also [32].

2.3. H^2 -nonconforming virtual element. Then we consider the case $m = 2$. For each $F \in \mathcal{F}^1(K)$ and any function $v \in H^4(K)$, set

$$\begin{aligned} M_{\nu\nu}(v) &:= \nu_{F,1}^\top (\nabla^2 v) \nu_{K,F}, \\ M_{\nu t}(v) &:= (\nabla^2 v) \nu_{K,F} - M_{\nu\nu}(v) \nu_{F,1}, \\ Q_\nu(v) &:= \nu_{K,F}^\top \operatorname{div}(\nabla^2 v) + \operatorname{div}_F M_{\nu t}(v). \end{aligned}$$

In two dimensions, $M_{\nu\nu}(v)$, $M_{\nu t}(v)$, and $Q_\nu(v)$ are called the normal bending moment, twisting moment, and effective transverse shear force, respectively, when v is the deflection of a thin plate in the context of elastic mechanics [25, 34].

Lemma 2.1. *For any $u \in H^4(K)$ and $v \in H^2(K)$, it holds that*

$$\begin{aligned} (\nabla^2 u, \nabla^2 v)_K &= (\Delta^2 u, v)_K + \sum_{F \in \mathcal{F}^1(K)} \left[(M_{\nu\nu}(u), \frac{\partial v}{\partial \nu_{F,1}})_F - (Q_\nu(u), v)_F \right] \\ (2.5) \quad &+ \sum_{e \in \mathcal{F}^2(K)} \sum_{F \in \mathcal{F}^1(K) \cap \partial^{-1}e} (\nu_{F,e}^\top M_{\nu t}(u), v)_e. \end{aligned}$$

Proof. Using integration by parts, we get

$$(\operatorname{div}(\nabla^2 u), \nabla v)_K = -(\Delta^2 u, v)_K + \sum_{F \in \mathcal{F}^1(K)} (\nu_{K,F}^\top \operatorname{div}(\nabla^2 u), v)_F,$$

and for each $F \in \mathcal{F}^1(K)$,

$$(M_{\nu t}(u), \nabla_F v)_F = -(\operatorname{div}_F M_{\nu t}(u), v)_F + \sum_{e \in \mathcal{F}^1(F)} (\nu_{F,e}^\top M_{\nu t}(u), v)_e.$$

Then we acquire from the last two identities and splitting the gradient into the tangential and normal components that

$$\begin{aligned}
 (\nabla^2 u, \nabla^2 v)_K &= -(\operatorname{div}(\nabla^2 u), \nabla v)_K + \sum_{F \in \mathcal{F}^1(K)} ((\nabla^2 u) \nu_{K,F}, \nabla v)_F \\
 &= -(\operatorname{div}(\nabla^2 u), \nabla v)_K + \sum_{F \in \mathcal{F}^1(K)} (M_{\nu\nu}(u), \frac{\partial v}{\partial \nu_{F,1}})_F \\
 &\quad + \sum_{F \in \mathcal{F}^1(K)} (M_{\nu t}(u), \nabla_F v)_F \\
 &= (\Delta^2 u, v)_K + \sum_{F \in \mathcal{F}^1(K)} \left[(M_{\nu\nu}(u), \frac{\partial v}{\partial \nu_{F,1}})_F - (Q_\nu(u), v)_F \right] \\
 &\quad + \sum_{F \in \mathcal{F}^1(K)} \sum_{e \in \mathcal{F}^1(F)} (\nu_{F,e}^\top M_{\nu t}(u), v)_e,
 \end{aligned}$$

which ends the proof. $\square$

Inspired by the Green's identity (2.5), for any element $K \in \mathcal{T}_h$ and integer $k \geq 2$, the local degrees of freedom $\mathcal{N}_k(K)$ of the H^2 nonconforming virtual element are given as follows:

$$(2.6) \quad \frac{1}{|K|} (v, q)_K \quad \forall q \in \mathbb{M}_{k-4}(K),$$

$$(2.7) \quad \frac{1}{|F|} (v, q)_F \quad \forall q \in \mathbb{M}_{k-3}(F) \text{ on each } F \in \mathcal{F}^1(K),$$

$$(2.8) \quad \frac{1}{|F|^{(n-2)/(n-1)}} \left(\frac{\partial v}{\partial \nu_{F,1}}, q \right)_F \quad \forall q \in \mathbb{M}_{k-2}(F) \text{ on each } F \in \mathcal{F}^1(K),$$

$$(2.9) \quad \frac{1}{|e|} (v, q)_e \quad \forall q \in \mathbb{M}_{k-2}(e) \text{ on each } e \in \mathcal{F}^2(K).$$

The local space of the H^2 nonconforming virtual element is

$$\begin{aligned}
 V_k(K) &:= \{u \in H^2(K) : \Delta^2 u \in \mathbb{P}_{k-4}(K), M_{\nu\nu}(u)|_F \in \mathbb{P}_{k-2}(F), Q_\nu(u)|_F \in \mathbb{P}_{k-3}(F), \\
 &\quad \sum_{F \in \mathcal{F}^1(K) \cap \partial^{-1}e} \nu_{F,e}^\top M_{\nu t}(u)|_e \in \mathbb{P}_{k-2}(e) \quad \forall F \in \mathcal{F}^1(K), e \in \mathcal{F}^2(K)\}.
 \end{aligned}$$

Remark 2.2. In two dimensions, the degrees of freedom (2.9) will be reduced to the function values on the vertices of K . Then the virtual element $(K, \mathcal{N}_k(K), V_k(K))$ is the same as that in [5, 44].

Remark 2.3. If the element $K \in \mathcal{T}_h$ is a simplex and $k = 2$, the degrees of freedom (2.6)–(2.7) disappear, and the degrees of freedom (2.8)–(2.9) are the same as the Morley–Wang–Xu element's degrees of freedom in [36]. Indeed, the virtual element $(K, \mathcal{N}_k(K), V_k(K))$ coincides with the Morley–Wang–Xu element in [36] when $k = 2$ and K is a simplex.

3. H^m -NONCONFORMING VIRTUAL ELEMENT WITH $1 \leq m \leq n$

In this section, we will construct the H^m -nonconforming virtual element. It has been illustrated in Sections 2.2–2.3 that the Green's identity plays a vital role in deriving the H^1 and H^2 nonconforming virtual elements. To this end, we shall first derive a generalized Green's identity for the H^m space.

3.1. Generalized Green's identity. For any scalar or tensor-valued smooth function v , nonnegative integer j , $F \in \mathcal{F}_h^r$ with $1 \leq r \leq n$, and $\alpha \in A_r$, we use $D_{F,\alpha}^j(v)$ to denote some j th order derivatives of v restrict on F , which may take different expressions at different appearances.

Lemma 3.1. *Let $K \in \mathcal{T}_h$, let $F \in \mathcal{F}^r(K)$ with $1 \leq r \leq n-1$, and let s be a positive integer satisfying $s \leq n-r$. There exist differential operators $D_{e,\alpha}^{s-j-|\alpha|}$ for $j = 0, \dots, s$, $e \in \mathcal{F}^j(F)$, and $\alpha \in A_{r+j}$ with $|\alpha| \leq s-j$ such that for any $\tau \in H^s(F; \mathbb{T}_n(s))$ and $(\nabla^s v)|_F \in L^2(F; \mathbb{T}_n(s))$, it holds that*

$$(3.1) \quad (\tau, \nabla^s v)_F = \sum_{j=0}^s \sum_{e \in \mathcal{F}^j(F)} \sum_{\substack{\alpha \in A_{r+j} \\ |\alpha| \leq s-j}} \left(D_{e,\alpha}^{s-j-|\alpha|}(\tau), \frac{\partial^{|\alpha|} v}{\partial \nu_e^\alpha} \right)_e.$$

Proof. We adopt the mathematical induction to prove the identity (3.1). When $s = 1$, we get from (2.1) and integration by parts

$$\begin{aligned} (\tau, \nabla v)_F &= \sum_{i=1}^r (\tau, \frac{\partial v}{\partial \nu_{F,i}} \nu_{F,i})_F + (\tau, \nabla_F v)_F \\ &= \sum_{i=1}^r (\nu_{F,i}^\top \tau, \frac{\partial v}{\partial \nu_{F,i}})_F - (\operatorname{div}_F \tau, v)_F + \sum_{e \in \mathcal{F}^1(F)} (\nu_{F,e}^\top \tau, v)_e. \end{aligned}$$

Thus the identity (3.1) holds for $s = 1$.

Next assume that the identity (3.1) is true for $s = \ell - 1$ with $2 \leq \ell \leq n - r$, and then let us prove it is also true for $s = \ell$. We get from (2.1) and integration by parts

$$\begin{aligned} (\tau, \nabla^\ell v)_F &= \sum_{i=1}^r (\tau \nu_{F,i}, \nabla^{\ell-1} \frac{\partial v}{\partial \nu_{F,i}})_F + (\tau, \nabla_F \nabla^{\ell-1} v)_F \\ &= \sum_{i=1}^r (\tau \nu_{F,i}, \nabla^{\ell-1} \frac{\partial v}{\partial \nu_{F,i}})_F - (\operatorname{div}_F \tau, \nabla^{\ell-1} v)_F + \sum_{e \in \mathcal{F}^1(F)} (\tau \nu_{F,e}, \nabla^{\ell-1} v)_e. \end{aligned}$$

Applying the assumption with $s = \ell - 1$ to the right-hand side of the last equation term by term, we have

$$\begin{aligned} (\tau \nu_{F,i}, \nabla^{\ell-1} \frac{\partial v}{\partial \nu_{F,i}})_F &= \sum_{j=0}^{\ell-1} \sum_{e \in \mathcal{F}^j(F)} \sum_{\substack{\alpha \in A_{r+j} \\ |\alpha| \leq \ell-1-j}} \left(D_{e,\alpha}^{\ell-1-j-|\alpha|}(\tau \nu_{F,i}), \frac{\partial^{|\alpha|}}{\partial \nu_e^\alpha} \left(\frac{\partial v}{\partial \nu_{F,i}} \right) \right)_e, \\ (\operatorname{div}_F \tau, \nabla^{\ell-1} v)_F &= \sum_{j=0}^{\ell-1} \sum_{e \in \mathcal{F}^j(F)} \sum_{\substack{\alpha \in A_{r+j} \\ |\alpha| \leq \ell-1-j}} \left(D_{e,\alpha}^{\ell-1-j-|\alpha|}(\operatorname{div}_F \tau), \frac{\partial^{|\alpha|} v}{\partial \nu_e^\alpha} \right)_e, \\ (\tau \nu_{F,e}, \nabla^{\ell-1} v)_e &= \sum_{j=0}^{\ell-1} \sum_{\tilde{e} \in \mathcal{F}^j(e)} \sum_{\substack{\alpha \in A_{r+1+j} \\ |\alpha| \leq \ell-1-j}} \left(D_{\tilde{e},\alpha}^{\ell-1-j-|\alpha|}(\tau \nu_{F,e}), \frac{\partial^{|\alpha|} v}{\partial \nu_{\tilde{e}}^\alpha} \right)_{\tilde{e}}. \end{aligned}$$

Finally we conclude (3.1) for $s = \ell$ by combining the last four equations and the fact that $\nu_{F,i}$ is a linear combination of $\nu_{e,1}, \dots, \nu_{e,r+j}$ if $e \in \mathcal{F}^j(F)$. $\square$

For each term on the right-hand side of (3.1), the total number of differentiation of the integrand is $s - j$. In view of Stokes's theorem, $e \in \mathcal{F}^j(F)$ can be thought of as $e \in \partial^j F$ so that the total number of differentiations is s which matches that of the left-hand side. The bounds on s in Lemma 3.1 imply $r + s \leq n$; then $\mathcal{F}^s(F) \subset \mathcal{F}^{r+s}(K)$ is well-defined for $F \in \mathcal{F}^r(K)$. Hence we can recursively apply the Stokes theorem until there is no derivative of v on the lowest-dimensional faces.

We give two examples of identity (3.1). When $n \geq 2$, $s = 1$, and $1 \leq r \leq n - 1$, the explicit expression of (3.1) is that for any $F \in \mathcal{F}^r(K)$, $\tau \in H^1(F; \mathbb{R}^n)$, and $(\nabla v)|_F \in L^2(F; \mathbb{R}^n)$,

$$(\tau, \nabla v)_F = -(\operatorname{div}_F \tau, v)_F + \sum_{i=1}^r (\nu_{F,i}^\top \tau, \frac{\partial v}{\partial \nu_{F,i}})_F + \sum_{e \in \mathcal{F}^1(F)} (\nu_{F,e}^\top \tau, v)_e.$$

If $n = 3$ and $s = 2$, then $r = 1$. Also, the explicit expression of (3.1) is that for any $F \in \mathcal{F}^1(K)$, $\tau \in H^2(F; \mathbb{T}_3(2))$, and $(\nabla^2 v)|_F \in L^2(F; \mathbb{T}_3(2))$,

$$\begin{aligned} (\tau, \nabla^2 v)_F &= (\operatorname{div}_F \operatorname{div}_F \tau, v)_F - (\nu_{F,1}^\top (\operatorname{div}_F \tau) + \operatorname{div}_F (\tau \nu_{F,1}), \frac{\partial v}{\partial \nu_{F,1}})_F \\ &\quad + (\nu_{F,1}^\top \tau \nu_{F,1}, \frac{\partial^2 v}{\partial \nu_{F,1}^2})_F - \sum_{e \in \mathcal{F}^1(F)} (\nu_{F,e}^\top (\operatorname{div}_F \tau) + \operatorname{div}_e (\tau \nu_{F,e}), v)_e \\ &\quad + \sum_{e \in \mathcal{F}^1(F)} \sum_{i=1}^2 (\nu_{e,i}^\top \tau \nu_{F,e} + (\nu_{e,i}^\top \nu_{F,1}) \nu_{F,1}^\top \tau \nu_{F,e}, \frac{\partial v}{\partial \nu_{e,i}})_e \\ &\quad + \sum_{e \in \mathcal{F}^1(F)} \sum_{\delta \in \mathcal{F}^1(e)} (\nu_{e,\delta}^\top \tau \nu_{F,e}) (\delta) v(\delta). \end{aligned}$$

Theorem 3.2. *Let $1 \leq m \leq n$. There exist differential operators $D_{F,\alpha}^{2m-j-|\alpha|}$ for $j = 1, \dots, m$, $F \in \mathcal{F}^j(K)$, and $\alpha \in A_j$ with $|\alpha| \leq m - j$ such that it holds for any $u \in H^m(K)$ satisfying $(-\Delta)^m u \in L^2(K)$ and $D_{F,\alpha}^{2m-j-|\alpha|} u \in L^2(F)$, and any $v \in H^m(K)$*

$$\begin{aligned} (\nabla^m u, \nabla^m v)_K &= ((-\Delta)^m u, v)_K \\ (3.2) \quad &+ \sum_{j=1}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} \left(D_{F,\alpha}^{2m-j-|\alpha|} u, \frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} \right)_F. \end{aligned}$$

Proof. By the density argument, we can assume $u \in H^{2m}(K)$. We still use the mathematical induction to prove the identity (3.2). The identity (3.2) for $m = 1$ and $m = 2$ is just the identities (2.2) and (2.5), respectively. Assume the identity (3.2) is true for $m = \ell - 1$ with $3 \leq \ell \leq n$; then let us prove that it is also true for $m = \ell$.

Applying the integration by parts,

$$\begin{aligned} (\nabla^\ell u, \nabla^\ell v)_K &= -(\operatorname{div} \nabla^\ell u, \nabla^{\ell-1} v)_K + \sum_{F \in \mathcal{F}^1(K)} ((\nabla^\ell u) \nu_{K,F}, \nabla^{\ell-1} v)_F \\ &= (\nabla^{\ell-1} (-\Delta u), \nabla^{\ell-1} v)_K + \sum_{F \in \mathcal{F}^1(K)} ((\nabla^\ell u) \nu_{K,F}, \nabla^{\ell-1} v)_F. \end{aligned}$$

Since the identity (3.2) holds for $m = \ell - 1$, we have

$$\begin{aligned} & (\nabla^{\ell-1}(-\Delta u), \nabla^{\ell-1}v)_K \\ &= ((-\Delta)^\ell u, v)_K + \sum_{j=1}^{\ell-1} \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq \ell-1-j}} \left(D_{F,\alpha}^{2(\ell-1)-j-|\alpha|}(-\Delta u), \frac{\partial^{|\alpha|}v}{\partial \nu_F^\alpha} \right)_F. \end{aligned}$$

Taking $\tau = (\nabla^\ell u)\nu_{K,F}$, $s = \ell - 1$, and $r = 1$ in (3.1), we get

$$\begin{aligned} ((\nabla^\ell u)\nu_{K,F}, \nabla^{\ell-1}v)_F &= \sum_{j=0}^{\ell-1} \sum_{e \in \mathcal{F}^j(F)} \sum_{\substack{\alpha \in A_{1+j} \\ |\alpha| \leq \ell-1-j}} \left(D_{e,\alpha}^{\ell-1-j-|\alpha|}((\nabla^\ell u)\nu_{K,F}), \frac{\partial^{|\alpha|}v}{\partial \nu_e^\alpha} \right)_e \\ &= \sum_{j=1}^{\ell} \sum_{e \in \mathcal{F}^{j-1}(F)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq \ell-j}} \left(D_{e,\alpha}^{\ell-j-|\alpha|}((\nabla^\ell u)\nu_{K,F}), \frac{\partial^{|\alpha|}v}{\partial \nu_e^\alpha} \right)_e. \end{aligned}$$

Therefore we finish the proof by combining the last three equations. $\square$

Examples for $m = 1, 2, n \geq m$, and $m = n = 3$ can be found in Appendix B.

3.2. Virtual element space. Inspired by identity (3.2), for any element $K \in \mathcal{T}_h$ and integer $k \geq m$, the local degrees of freedom $\mathcal{N}_k(K)$ are given as follows:

$$(3.3) \quad \frac{1}{|K|}(v, q)_K \quad \forall q \in \mathbb{M}_{k-2m}(K),$$

$$(3.4) \quad \frac{1}{|F|^{(n-j-|\alpha|)/(n-j)}} \left(\frac{\partial^{|\alpha|}v}{\partial \nu_F^\alpha}, q \right)_F \quad \forall q \in \mathbb{M}_{k-(2m-j-|\alpha|)}(F)$$

on each $F \in \mathcal{F}^j(K)$, where $j = 1, \dots, m$, $\alpha \in A_j$, and $|\alpha| \leq m - j$. Denote by N_K the number of local degrees of freedom (3.3)–(3.4).

We present a heuristic explanation of the scaling factor in (3.4). Let $\hat{K} = \{\hat{\mathbf{x}} \in \mathbb{R}^n : \hat{\mathbf{x}} = \frac{1}{h_K}(\mathbf{x} - \mathbf{x}_K) \quad \forall \mathbf{x} \in K\}$, and let an affine mapping $\Psi : \hat{\mathbf{x}} \in \mathbb{R}^n \rightarrow \Psi(\hat{\mathbf{x}}) = h_K \hat{\mathbf{x}} + \mathbf{x}_K \in \mathbb{R}^n$. Then $h_{\hat{K}} \approx 1$ and $\Psi(\hat{K}) = K$. For any function $v(\mathbf{x})$ defined on K , let $\hat{v}(\hat{\mathbf{x}}) := v(\Psi(\hat{\mathbf{x}}))$, which is defined on $\hat{K}$. By the scaling argument, we have

$$\left(\frac{\partial^{|\alpha|}v}{\partial \nu_F^\alpha}, q \right)_F = h_K^{n-j-|\alpha|} \left(\frac{\partial^{|\alpha|}\hat{v}}{\partial \nu_{\hat{F}}^\alpha}, \hat{q} \right)_{\hat{F}}.$$

By the mesh conditions (A1)–(A2) in Section 4.2, it holds that $|F| \approx h_K^{n-j}$. Thus there exists a constant $C > 0$ being independent of h_K such that $|F|^{(n-j-|\alpha|)/(n-j)} = Ch_K^{n-j-|\alpha|}$. Then

$$\frac{1}{|F|^{(n-j-|\alpha|)/(n-j)}} \left(\frac{\partial^{|\alpha|}v}{\partial \nu_F^\alpha}, q \right)_F = \frac{1}{C} \left(\frac{\partial^{|\alpha|}\hat{v}}{\partial \nu_{\hat{F}}^\alpha}, \hat{q} \right)_{\hat{F}} = \frac{1}{C_1} \frac{1}{|\hat{F}|^{(n-j-|\alpha|)/(n-j)}} \left(\frac{\partial^{|\alpha|}\hat{v}}{\partial \nu_{\hat{F}}^\alpha}, \hat{q} \right)_{\hat{F}},$$

where $C_1 = C/|\hat{F}|^{(n-j-|\alpha|)/(n-j)}$ is independent of h_K . Hence all the degrees of freedom in (3.3)–(3.4) share the same order of magnitude.

Again due to the first terms in the inner products of the right-hand side of (3.2), and the degrees of freedom (3.3)–(3.4), it is inherent to define the local space of the

H^m -nonconforming virtual element as

$$\begin{aligned} V_k(K) &:= \{u \in H^m(K) : (-\Delta)^m u \in \mathbb{P}_{k-2m}(K), \\ &\quad D_{F,\alpha}^{2m-j-|\alpha|}(u)|_F \in \mathbb{P}_{k-(2m-j-|\alpha|)}(F) \quad \forall F \in \mathcal{F}^j(K), \\ &\quad j = 1, \dots, m, \alpha \in A_j, \text{ and } |\alpha| \leq m-j\}, \end{aligned}$$

where the differential operators $D_{F,\alpha}^{2m-j-|\alpha|}$ are introduced in Theorem 3.2.

In the following we shall prove that $(K, \mathcal{N}_k(K), V_k(K))$ forms a finite element triple in the sense of Ciarlet [23]. Unlike the traditional finite element, in the virtual element only the set of the degrees of freedom $\mathcal{N}_k(K)$ needs to be explicitly known. The “virtual” space $V_k(K)$ is only needed for the purpose of analysis and the specific formulation for $D_{F,\alpha}^{2m-j-|\alpha|}$ is not needed in the definition of $V_k(K)$.

The following property is the direct result of (3.1) and the definition of the degrees of freedom (3.4).

Lemma 3.3. *Let $K \in \mathcal{T}_h$, let $F \in \mathcal{F}^r(K)$ with $1 \leq r \leq m$, and let $s \leq m-r$ be a nonnegative integer satisfying $k \geq 2m-(r+s)$. For any $\tau \in \mathbb{P}_{k-(2m-r-s)}(F; \mathbb{T}_n(s))$ and $(\nabla^s v)|_F \in L^2(F; \mathbb{T}_n(s))$, the term*

$$(\tau, \nabla^s v)_F$$

is uniquely determined by the degrees of freedom $\left(\frac{\partial^{|\alpha|} v}{\partial \nu_e^\alpha}, q\right)_e$ for all $0 \leq j \leq s$, $e \in \mathcal{F}^j(F)$, $\alpha \in A_{r+j}$ with $|\alpha| \leq s-j$, and $q \in \mathbb{M}_{k-(2m-r-j-|\alpha|)}(e)$.

Lemma 3.4. *We have $\mathbb{P}_k(K) \subseteq V_k(K)$ and*

$$(3.5) \quad \dim V_k(K) = \dim \mathcal{N}_k(K).$$

Proof. For any $q \in \mathbb{P}_k(K)$, it is obvious that

$$(-\Delta)^m q \in \mathbb{P}_{k-2m}(K), \quad D_{F,\alpha}^{2m-j-|\alpha|} q|_F \in \mathbb{P}_{k-(2m-j-|\alpha|)}(F).$$

Hence it holds that $\mathbb{P}_k(K) \subseteq V_k(K)$. Since all the differential operators in the definition of $V_k(K)$ are linear, $V_k(K)$ is a vector space.

Next we count the dimension of $V_k(K)$. Consider the local polyharmonic equation with the Neumann boundary condition

$$(3.6) \quad \begin{cases} (-\Delta)^m u = f_1 & \text{in } K, \\ D_{F,\alpha}^{2m-j-|\alpha|}(u) = g_j^{F,\alpha} & \text{on each } F \in \mathcal{F}^j(K) \\ & \text{with } j = 1, \dots, m, \alpha \in A_j, \text{ and } |\alpha| \leq m-j, \end{cases}$$

where $f_1 \in \mathbb{P}_{k-2m}(K)$, $g_j^{F,\alpha} \in \mathbb{P}_{k-(2m-j-|\alpha|)}(F)$. Applying the generalized Green’s identity (3.2), the weak formulation of (3.6) is

$$(3.7) \quad (\nabla^m u, \nabla^m v)_K = (f_1, v)_K + \sum_{j=1}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} \left(g_j^{F,\alpha}, \frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} \right)_F$$

for any $v \in H^m(K)$. If we take $v = q \in \mathbb{P}_{m-1}(K)$ in (3.7), we have the compatibility condition of the data

$$(3.8) \quad (f_1, q)_K + \sum_{j=1}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} \left(g_j^{F,\alpha}, \frac{\partial^{|\alpha|} q}{\partial \nu_F^\alpha} \right)_F = 0 \quad \forall q \in \mathbb{P}_{m-1}(K).$$

On the other hand, given $f_1 \in \mathbb{P}_{k-2m}(K)$, $g_j^{F,\alpha} \in \mathbb{P}_{k-(2m-j-|\alpha|)}(F)$ satisfying the compatibility condition (3.8), the weak formulation of the Neumann problem of the local polyharmonic equation (3.6) is: find $u \in H^m(K)/\mathbb{P}_{m-1}(K)$ such that

$$(\nabla^m u, \nabla^m v)_K = (f_1, v)_K + \sum_{j=1}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} \left(g_j^{F,\alpha}, \frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} \right)_F$$

for all $v \in H^m(K)/\mathbb{P}_{m-1}(K)$. The well-posedness of this variational formulation is guaranteed by the Lax–Milgram lemma [9, 23], and specifically the well-posedness of polyharmonic equations with various boundary conditions can be found in [2, 26].

Therefore $\dim(V_k(K)/\mathbb{P}_{m-1}(K))$ equals

$$\dim \mathbb{P}_{k-2m}(K) + \sum_{j=1}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} \dim \mathbb{P}_{k-(2m-j-|\alpha|)}(F) - \dim \mathbb{P}_{m-1}(K),$$

where the dimension of the constraint for the data is subtracted. When counting $\dim V_k(K)$, we should add back the dimension of the kernel space, i.e., solution spaces of $(\nabla^m u, \nabla^m v)_K = 0$, which implies $\dim V_k(K) = \dim \mathbb{P}_{k-2m}(K) + \sum_{j=1}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} \dim \mathbb{P}_{k-(2m-j-|\alpha|)}(F) = \dim \mathcal{N}_k(K)$. $\square$

Lemma 3.5. *The degrees of freedom (3.3)–(3.4) are unisolvent for the local virtual element space $V_k(K)$.*

Proof. Let $v \in V_k(K)$ and suppose all the degrees of freedom (3.3)–(3.4) vanish. We get from (3.7) that

$$\|\nabla^m v\|_{0,K}^2 = 0.$$

Thus $v \in \mathbb{P}_{m-1}(K)$. By Lemma 3.3 with $s = m - r$, we have for any $F \in \mathcal{F}^r(K)$ with $1 \leq r \leq m$,

$$(3.9) \quad (\tau, \nabla^s v)_F = 0 \quad \forall \tau \in \mathbb{P}_0(F; \mathbb{T}_n(s)).$$

Due to (3.9) with $r = 1$ and the fact that $v \in \mathbb{P}_{m-1}(K)$, it follows that $v \in \mathbb{P}_{m-2}(K)$. Recursively applying (3.9) with $r = 2, \dots, m$ gives $v = 0$. This ends the proof. $\square$

Remark 3.6. If the element $K \in \mathcal{T}_h$ is a simplex and $k = m$, the degrees of freedom (3.3) disappear, and the degrees of freedom (3.4) are the same as those of the nonconforming finite element in [37]. Since $\mathbb{P}_m(K) \subseteq V_k(K)$, the virtual element $(K, \mathcal{N}_k(K), V_k(K))$ coincides with the nonconforming finite element in [37] when K is a simplex and $k = m$, which is the minimal finite element for the $2m$ th order partial differential equations in $\mathbb{R}^n$. In other words, we generalize the nonconforming finite element in [37] to high order $k > m$ and arbitrary shape of polytopes.

3.3. Local projections. For each $K \in \mathcal{T}_h$, define a local H^m projection $\Pi^K : H^m(K) \rightarrow \mathbb{P}_k(K)$ as follows: given $v \in H^m(K)$, let $\Pi_k^K v \in \mathbb{P}_k(K)$ be the solution of the problem

$$(3.10) \quad (\nabla^m \Pi_k^K v, \nabla^m q)_K = (\nabla^m v, \nabla^m q)_K \quad \forall q \in \mathbb{P}_k(K),$$

$$(3.11) \quad \sum_{F \in \mathcal{F}^r(K)} Q_0^F(\nabla^{m-r} \Pi_k^K v) = \sum_{F \in \mathcal{F}^r(K)} Q_0^F(\nabla^{m-r} v), \quad r = 1, \dots, m.$$

The number of equations in (3.11) is

$$\sum_{r=1}^m C_{n+m-1-r}^{n-1} = C_{n+m-1}^n = \dim(\mathbb{P}_{m-1}(K)).$$

Then the well-posedness of (3.10)–(3.11) can be shown by the similar argument as in the proof of Lemma 3.5. To simplify the notation, we will write it as Π^K .

Obviously we have

$$(3.12) \quad |\Pi^K v|_{m,K} \leq |v|_{m,K} \quad \forall v \in H^m(K).$$

We show that the projection $\Pi^K u$ is computable using the degrees of freedom (3.3)–(3.4).

Lemma 3.7. *The operator $\Pi^K : H^m(K) \rightarrow \mathbb{P}_k(K)$ is a projector, i.e.,*

$$(3.13) \quad \Pi^K v = v \quad \forall v \in \mathbb{P}_k(K),$$

and the projector Π^K can be computed using only the degrees of freedom (3.3)–(3.4).

Proof. We first show that Π^K is a projector. Let $p = \Pi^K v - v \in \mathbb{P}_k(K)$. Taking $q = p$ in (3.10), we get $\nabla^m p = 0$, i.e., $p \in \mathbb{P}_{m-1}(K)$. By (3.11),

$$\sum_{F \in \mathcal{F}^r(K)} Q_0^F(\nabla^{m-r} p) = 0, \quad r = 1, \dots, m.$$

Therefore $p = 0$, which means Π^K is a projector.

Next, by applying the identity (3.2) and the right-hand side of (3.10)

$$(\nabla^m v, \nabla^m q)_K = (v, (-\Delta)^m q)_K + \sum_{j=1}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} \left(\frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha}, D_{F,\alpha}^{2m-j-|\alpha|}(q) \right)_F.$$

Hence we conclude from the degrees of freedom (3.3)–(3.4) and Lemma 3.3 with $s = m - r$ that the right-hand sides of (3.10)–(3.11) are computable. $\square$

Remark 3.8. $D_{F,\alpha}^{2m-j-|\alpha|}$ is needed in the computation of Π^K . But since $q \in \mathbb{P}_k(K)$ and $\nabla^m q \in \mathbb{P}_{k-m}(K)$, few terms are left for moderate k .

Let $W_k(K) := V_k(K)$ for $k \geq 3m - 1$ or $m \leq k \leq 2m - 1$. To compute the L^2 projection onto $\mathbb{P}_{m-1}(K)$ for $2m \leq k < 3m - 1$, following [3], define

$$\begin{aligned} \tilde{V}_k(K) &:= \{v \in H^m(K) : (-\Delta)^m v \in \mathbb{P}_{m-1}(K), D_{F,\alpha}^{2m-j-|\alpha|}(v)|_F \in \mathbb{P}_{k-(2m-j-|\alpha|)}(F), \\ &\quad \forall F \in \mathcal{F}^j(K), j = 1, \dots, m, \alpha \in A_j, \text{ and } |\alpha| \leq m - j\}, \end{aligned}$$

$$W_k(K) := \{v \in \tilde{V}_k(K) : (v - \Pi^K v, q)_K = 0 \quad \forall q \in \mathbb{P}_{k-2m}^\perp(K)\},$$

where $\mathbb{P}_{k-2m}^\perp(K) \subset \mathbb{P}_{m-1}(K)$ is the orthogonal complement space of $\mathbb{P}_{k-2m}(K)$ in $\mathbb{P}_{m-1}(K)$ with respect to the inner product $(\cdot, \cdot)_K$. It is apparent that $\mathbb{P}_k(K) \subset W_k(K)$ and that the local space $W_k(K)$ shares the same degrees of freedom as $V_k(K)$. That is, for the same $\mathcal{N}_k(K)$, we can associate different “virtual” spaces and thus have a different interpretation.

Lemma 3.9. *The degrees of freedom (3.3)–(3.4) are unisolvent for the local virtual element space $W_k(K)$.*

Proof. It is enough to only consider the case $2m \leq k < 3m-1$. Take any $v \in W_k(K)$ all of whose degrees of freedom (3.3)–(3.4) disappear. Then $\Pi^K v = 0$. By the definition of $W_k(K)$, we have

$$(v, q)_K = 0 \quad \forall q \in \mathbb{P}_{k-2m}^\perp(K),$$

which together with (3.3) implies

$$(v, q)_K = 0 \quad \forall q \in \mathbb{P}_{m-1}(K).$$

Applying the argument in Lemma 3.5 to the space $\tilde{V}_k(K)$ with vanishing degrees of freedom (3.4) and the last equation, we know that $v = 0$. $\square$

In the original space $V_k(K)$, the volume moment (cf. (3.3)), is only defined up to degree $k-2m$ which cannot compute the L^2 -projection to $\mathbb{P}_{m-1}$ when k is small. For $2m \leq k < 3m-1$, a desirable property of the local virtual element space $W_k(K)$ is that the L^2 -projection Q_{m-1}^K is computable if all the degrees of freedom (3.3)–(3.4) are known. Indeed, it follows from the definition of $W_k(K)$ that

$$(Q_{m-1}^K - Q_{k-2m}^K)(v - \Pi^K v) = Q_{m-1}^K(I - Q_{k-2m}^K)(v - \Pi^K v) = 0 \quad \forall v \in W_k(K),$$

which provides a way to compute the L^2 -projection

$$(3.14) \quad Q_{m-1}^K v = Q_{k-2m}^K v + Q_{m-1}^K \Pi^K v - Q_{k-2m}^K \Pi^K v \quad \forall v \in W_k(K).$$

Denote by $I_K : H^m(K) \rightarrow W_k(K)$ the canonical interpolation operator based on the degrees of freedom in (3.3)–(3.4). Namely, given a $u \in H^m(K)$, $I_K u \in W_k(K)$ so that $\chi(u) = \chi(I_K u)$ for all $\chi \in \mathcal{N}_k(K)$. As a direct corollary of Lemma 3.7, we have the following identity.

Corollary 3.10. *For any $v \in H^m(K)$, it holds that*

$$(3.15) \quad \Pi^K(v) = \Pi^K(I_K v).$$

4. DISCRETE METHOD

We will present the virtual element method for the polyharmonic equation based on the virtual element $(K, \mathcal{N}_k(K), V_k(K))$ or $(K, \mathcal{N}_k(K), W_k(K))$ when L^2 -projection is needed.

4.1. Discretization. Consider the polyharmonic equation with a homogeneous Dirichlet boundary condition

$$(4.1) \quad \begin{cases} (-\Delta)^m u = f & \text{in } \Omega, \\ u = \frac{\partial u}{\partial \nu} = \dots = \frac{\partial^{m-1} u}{\partial \nu^{m-1}} = 0 & \text{on } \partial\Omega, \end{cases}$$

where $f \in L^2(\Omega)$ and $\Omega \subset \mathbb{R}^n$ with $1 \leq m \leq n$. The weak formulation of the polyharmonic equation (4.1) is to find $u \in H_0^m(\Omega)$ such that

$$(4.2) \quad (\nabla^m u, \nabla^m v) = (f, v) \quad \forall v \in H_0^m(\Omega).$$

Since

$$\|\nabla^m v\|_0 \approx \|v\|_m \quad \text{and} \quad (f, v) \lesssim \|f\|_0 \|v\|_m \quad \forall v \in H_0^m(\Omega),$$

it follows from the Lax–Milgram lemma that the variational formulation (4.2) is well-posed.

Define the global virtual element space as

$V_h := \{v_h \in L^2(\Omega) : v_h|_K \in W_k(K) \text{ for each } K \in \mathcal{T}_h; \text{ the degrees of freedom}$

$$\begin{aligned} & \left(\frac{\partial^{|\alpha|} v_h}{\partial \nu_F^\alpha}, q \right)_F \text{ are continuous through } F \text{ for all} \\ & q \in \mathbb{P}_{k-(2m-j-|\alpha|)}(F), \alpha \in A_j \text{ with } |\alpha| \leq m-j, F \in \mathcal{F}^j(K), \\ & \text{and } j = 1, \dots, m; \left(\frac{\partial^{|\alpha|} v_h}{\partial \nu_F^\alpha}, q \right)_F = 0 \text{ if } F \subset \partial\Omega \}. \end{aligned}$$

Define the local bilinear form $a_{h,K}(\cdot, \cdot) : W_k(K) \times W_k(K) \rightarrow \mathbb{R}$ as

$$a_{h,K}(w, v) := (\nabla^m \Pi^K w, \nabla^m \Pi^K v)_K + S_K(w - \Pi^K w, v - \Pi^K v),$$

where the stabilization term

$$(4.3) \quad S_K(w, v) := h_K^{n-2m} \sum_{i=1}^{N_K} \chi_i(w) \chi_i(v) \quad \forall w, v \in W_k(K),$$

where χ_i is the i th local degree of freedom in (3.3)–(3.4) for $i = 1, \dots, N_K$. The global bilinear form $a_h(\cdot, \cdot) : V_h \times V_h \rightarrow \mathbb{R}$ is

$$a_h(w_h, v_h) := \sum_{K \in \mathcal{T}_h} a_{h,K}(w_h, v_h).$$

Remark 4.1. The stabilization (4.3) resembles the original recipe in [12, 13]. This classical stabilization is easy to implement and was used to develop the discrete Galerkin orthogonality (5.1) and the stability bounds in Appendix A. However, the stabilization (4.3) usually suffers from conditioning and stability issues, especially for high-order k and differential problems with large m . Instead, several other stabilizations have been devised and investigated in [10, 11, 14, 24, 33] to cure these issues. The numerical results in [33] show that these stabilizations have almost the same effect on the condition number of the stiffness matrix for $n = 2$. The classical stabilization, the D-recipe stabilization in [11], and the D-recipe stabilization with only boundary dofs were compared in three dimensions in [24]; as a result the D-recipe stabilization outperforms the other two.

Define $\Pi_h : V_h \rightarrow \mathbb{P}_k(\mathcal{T}_h)$ as $(\Pi_h v)|_K := \Pi^K(v|_K)$ for each $K \in \mathcal{T}_h$, and let $Q_h^l : L^2(\Omega) \rightarrow \mathbb{P}_l(\mathcal{T}_h)$ be the L^2 -orthogonal projection onto $\mathbb{P}_l(\mathcal{T}_h)$. Then for any $v \in L^2(\Omega)$,

$$(Q_h^l v)|_K := Q_l^K(v|_K) \quad \forall K \in \mathcal{T}_h.$$

We compute the right-hand side according to the following cases:

$$(4.4) \quad \langle f, v_h \rangle := \begin{cases} (f, \Pi_h v_h), & m \leq k \leq 2m-1, \\ (f, Q_h^{m-1} v_h), & 2m \leq k \leq 3m-2, \\ (f, Q_h^{k-2m} v_h), & 3m-1 \leq k. \end{cases}$$

We will need this definition of the right-hand side in order to get an optimal order of convergence; see Lemma 5.7.

Remark 4.2. When $m \leq k \leq 2m-1$, which is an important range for large m as high-order methods are harder to implement, there is no need to compute a new projection and no need to modify the local virtual element space. For $k \geq 2m$, however, an L^2 -projection to a higher degree polynomial space is needed to control the consistency error; see Section 5.2.

With the previous preparations, we propose the nonconforming virtual element method for the polyharmonic equation (4.1) in any dimension: find $u_h \in V_h$ such that

$$(4.5) \quad a_h(u_h, v_h) = \langle f, v_h \rangle \quad \forall v_h \in V_h.$$

In the rest of this section, we shall prove the well-posedness of the discretization (4.5) by establishing the coercivity and continuity of the bilinear form $a_h(\cdot, \cdot)$.

4.2. Mesh conditions. We impose the following conditions on the mesh $\mathcal{T}_h$:

- (A1) Each element $K \in \mathcal{T}_h$ and each face $F \in \mathcal{F}_h^r$ for $1 \leq r \leq n-1$ is star-shaped with a uniformly bounded star-shaped constant.
- (A2) There exists a quasiuniform simplicial mesh $\mathcal{T}_h^*$ such that each $K \in \mathcal{T}_h$ is a union of some simplexes in $\mathcal{T}_h^*$.

Notice that (A1) and (A2) imply $\text{diam}(F) \approx \text{diam}(K)$ for all $F \in \mathcal{F}^r(K)$, $1 \leq r \leq n-1$.

For a star-shaped domain D , there exists a ball $B_D \subset D$ with radius $\rho_D h_D$ and a Lipschitz isomorphism $\Phi : B_D \rightarrow D$ such that $|\Phi|_{1,\infty,B_D}$ and $|\Phi^{-1}|_{1,\infty,D}$ are bounded by a constant depending only on the chunkiness parameter ρ_D . Then several trace inequalities of $H^1(D)$ can be established with a constant depending only on ρ_D [17, (2.18)]. In particular, we shall use

$$(4.6) \quad \|v\|_{0,\partial D}^2 \lesssim h_D^{-1} \|v\|_{0,D}^2 + h_D |v|_{1,D}^2 \quad \forall v \in H^1(D).$$

The condition (A2) is inspired by the virtual triangulation condition used in [18, 22]. The simplicial mesh $\mathcal{T}_h^*$ will serve as a bridge to transfer the results from finite element methods to virtual element methods.

Very recently, some geometric assumptions being the relaxation of conditions (A1)–(A2) were suggested in [20, 21] under which a refined error analysis was developed for the linear conforming and nonconforming virtual element methods of the Poisson equation, i.e., the H^1 case. For high order H^m , $m > 1$, elements, we will investigate such a relaxation in future works.

4.3. Weak continuity. Based on Lemma 3.3, the space V_h has the weak continuity; that is, for any $F \in \mathcal{F}_h^1$, $v_h \in V_h$, and nonnegative integer $s \leq m-1$,

$$(4.7) \quad (\llbracket \nabla_h^s v_h \rrbracket, \tau)_F = 0 \quad \forall \tau \in \mathbb{P}_{k-(2m-1-s)}(F; \mathbb{T}_n(s)),$$

$$(4.8) \quad Q_0^e(\llbracket \nabla_h^s v_h \rrbracket|_F) = 0 \quad \forall e \in \mathcal{F}^{m-s-1}(F),$$

where ∇_h is the elementwise gradient with respect to the partition $\mathcal{T}_h$. We shall derive some bound on the jump $\llbracket \nabla_h^s v_h \rrbracket$ using the weak continuity and the trace inequality.

By the weak continuity (4.7), the mean value of $\nabla_h^s v_h$ over F is continuous only when $s \geq 2m-1-k$. For $s < 2m-1-k$, the mean value of $\nabla_h^s v_h$ is merely continuous over some low-dimensional face of F ; cf. (4.8). As a concrete example, consider the Morley element in three dimensions. The mean value of $\nabla_h v_h$ over faces is continuous, but the mean value of v_h is only continuous on edges rather than over faces.

Recall the following error estimates of the L^2 -projection.

Lemma 4.3. *Let $\ell \in \mathbb{N}$. For each $K \in \mathcal{T}_h$ and $\mathcal{F}^1(K)$, we have for any $v \in H^{\ell+1}(K)$*

$$(4.9) \quad \|v - Q_\ell^K v\|_{0,K} \lesssim h_K^{\ell+1} |v|_{\ell+1,K},$$

$$(4.10) \quad \|v - Q_\ell^F v\|_{0,F} \lesssim h_K^{\ell+1/2} |v|_{\ell+1,K}.$$

Then recall the Bramble–Hilbert Lemma (cf. [16, Lemma 4.3.8]).

Lemma 4.4. *Let $\ell \in \mathbb{N}$, and let $K \in \mathcal{T}_h \cup \mathcal{T}_h^*$. There exists a linear operator $T_\ell^K : L^1(K) \rightarrow \mathbb{P}_\ell(K)$ such that for any $v \in H^{\ell+1}(K)$,*

$$(4.11) \quad \|v - T_\ell^K v\|_{j,K} \lesssim h_K^{\ell+1-j} |v|_{\ell+1,K} \quad \text{for } 0 \leq j \leq \ell + 1.$$

Notice that the constants in (4.9)–(4.11) depend on the star-shaped constant, i.e., the chunkiness parameter ρ_K , and also depend on the degree ℓ .

Similarly, define $T_h : L^2(\Omega) \rightarrow \mathbb{P}_k(\mathcal{T}_h)$ as

$$(T_h v)|_K := T_k^K(v|_K) \quad \forall K \in \mathcal{T}_h.$$

Lemma 4.5. *Given $F \in \mathcal{F}_h^1$ and a positive integer $s < m$, assume for any $e \in \mathcal{F}^r(F)$ with $r = 0, 1, \dots, m-1-s$ that*

$$(4.12) \quad \|\llbracket \nabla_h^s v_h \rrbracket\|_{0,e} \lesssim \sum_{K \in \partial^{-1}F} h_K^{m-s-(r+1)/2} |v_h|_{m,K} \quad \forall v_h \in V_h.$$

Then we have for any $e \in \mathcal{F}^r(F)$ with $r = 0, 1, \dots, m-s$,

$$(4.13) \quad \|\llbracket \nabla_h^{s-1} v_h \rrbracket\|_{0,e} \lesssim \sum_{K \in \partial^{-1}F} h_K^{m-s-(r-1)/2} |v_h|_{m,K} \quad \forall v_h \in V_h.$$

Proof. We use the mathematical induction on r to prove (4.13). First consider $r = 0$. Take some $e_i \in \mathcal{F}^i(F)$ for $i = 1, \dots, m-s$ such that $e_i \in \mathcal{F}^1(e_{i-1})$ with $e_0 = F$. In the following we shall use Q_0^e as the L^2 -orthogonal projection onto the constant tensor space on e which can be understood as a tensor defined on the whole space. Employing the trace inequality (4.6), we get from (4.12) with $r = i-1$

$$\begin{aligned} & h_F^{i/2} \|\llbracket \nabla_h^{s-1} v_h \rrbracket - Q_0^F(\llbracket \nabla_h^{s-1} v_h \rrbracket)\|_{0,e_i} \\ & \lesssim h_F^{(i-1)/2} \|\llbracket \nabla_h^{s-1} v_h \rrbracket - Q_0^F(\llbracket \nabla_h^{s-1} v_h \rrbracket)\|_{0,e_{i-1}} + h_F^{(i+1)/2} \|\llbracket \nabla_h^s v_h \rrbracket\|_{0,e_{i-1}} \\ & \lesssim h_F^{(i-1)/2} \|\llbracket \nabla_h^{s-1} v_h \rrbracket - Q_0^F(\llbracket \nabla_h^{s-1} v_h \rrbracket)\|_{0,e_{i-1}} + \sum_{K \in \partial^{-1}F} h_K^{m-s+1/2} |v_h|_{m,K}. \end{aligned}$$

By this recursive inequality and the approximation properties of the L^2 -projection, it holds that

$$\begin{aligned} & h_F^{(m-s)/2} \|\llbracket \nabla_h^{s-1} v_h \rrbracket - Q_0^F(\llbracket \nabla_h^{s-1} v_h \rrbracket)\|_{0,e_{m-s}} \\ & \lesssim \|\llbracket \nabla_h^{s-1} v_h \rrbracket - Q_0^F(\llbracket \nabla_h^{s-1} v_h \rrbracket)\|_{0,F} + \sum_{K \in \partial^{-1}F} h_K^{m-s+1/2} |v_h|_{m,K} \\ & \lesssim h_F \|\llbracket \nabla_h^s v_h \rrbracket\|_{0,F} + \sum_{K \in \partial^{-1}F} h_K^{m-s+1/2} |v_h|_{m,K}. \end{aligned}$$

On the other side, we have from (4.8) that

$$\begin{aligned}
& \|[\nabla_h^{s-1} v_h]\|_{0,F} = \|[\nabla_h^{s-1} v_h] - Q_0^{e_{m-s}}([\nabla_h^{s-1} v_h])\|_{0,F} \\
& = \|[\nabla_h^{s-1} v_h] - Q_0^F([\nabla_h^{s-1} v_h]) - Q_0^{e_{m-s}}([\nabla_h^{s-1} v_h] - Q_0^F([\nabla_h^{s-1} v_h]))\|_{0,F} \\
& \leq \|[\nabla_h^{s-1} v_h] - Q_0^F([\nabla_h^{s-1} v_h])\|_{0,F} + \|Q_0^{e_{m-s}}([\nabla_h^{s-1} v_h] - Q_0^F([\nabla_h^{s-1} v_h]))\|_{0,F} \\
& \lesssim h_F \|[\nabla_h^s v_h]\|_{0,F} + h_F^{(m-s)/2} \|Q_0^{e_{m-s}}([\nabla_h^{s-1} v_h] - Q_0^F([\nabla_h^{s-1} v_h]))\|_{0,e_{m-s}} \\
& \leq h_F \|[\nabla_h^s v_h]\|_{0,F} + h_F^{(m-s)/2} \|[\nabla_h^{s-1} v_h] - Q_0^F([\nabla_h^{s-1} v_h])\|_{0,e_{m-s}}.
\end{aligned}$$

Hence (4.13) with $r = 0$ follows from the last two inequalities and (4.12) with $r = 0$.

Assume (4.13) holds with $r = j < m - s$. Let $e \in \mathcal{F}^{j+1}(F)$. Take some $e_j \in \mathcal{F}^j(F)$ satisfying $e \in \mathcal{F}^1(e_j)$. Using the trace inequality (4.6) again, we know that

$$\|[\nabla_h^{s-1} v_h]\|_{0,e} \lesssim h_e^{-1/2} \|[\nabla_h^{s-1} v_h]\|_{0,e_j} + h_e^{1/2} \|[\nabla_h^s v_h]\|_{0,e_j},$$

which combined with the assumptions means (4.13) is true with $r = j + 1$. $\square$

Again consider the Morley–Wang–Xu element in three dimensions [36], i.e., $m = 2$ and $s = 1$. The inequality (4.12) is just

$$\|[\nabla_h v_h]\|_{0,F} \lesssim \sum_{K \in \partial^{-1}F} h_K^{1/2} |v_h|_{2,K} \quad \forall v_h \in V_h,$$

for each face $F \in \mathcal{F}_h^1$. Then by Lemma 4.5 we will get from (4.8) that

$$\|v_h\|_{0,F} + h_K^{1/2} \|v_h\|_{0,e} \lesssim \sum_{K \in \partial^{-1}F} h_K^{3/2} |v_h|_{2,K} \quad \forall v_h \in V_h,$$

for any face $F \in \mathcal{F}_h^1$ and any edge $e \in \mathcal{F}_h^2$. We refer to [36, Lemma 5] for these estimates on tetrahedra.

Lemma 4.6. *For each $F \in \mathcal{F}_h^1$ and nonnegative integer $s < m$, it holds that*

$$(4.14) \quad \|[\nabla_h^s v_h]\|_{0,F} \lesssim \sum_{K \in \partial^{-1}F} h_K^{m-s-1/2} |v_h|_{m,K} \quad \forall v_h \in V_h.$$

Proof. It is sufficient to prove that for $s = m - 1, m - 2, \dots, 0$ and any $e \in \mathcal{F}^r(F)$ with $r = 0, 1, \dots, m - 1 - s$, it holds that

$$\|[\nabla_h^s v_h]\|_{0,e} \lesssim \sum_{K \in \partial^{-1}F} h_K^{m-s-(r+1)/2} |v_h|_{m,K} \quad \forall v_h \in V_h.$$

According to Lemma 4.5 and the mathematical induction, we only need to show

$$\|[\nabla_h^{m-1} v_h]\|_{0,F} \lesssim \sum_{K \in \partial^{-1}F} h_K^{1/2} |v_h|_{m,K} \quad \forall v_h \in V_h.$$

In fact, due to (4.8) and (4.10), we get

$$\|[\nabla_h^{m-1} v_h]\|_{0,F} = \|[\nabla_h^{m-1} v_h] - Q_0^F([\nabla_h^{m-1} v_h])\|_{0,F} \lesssim \sum_{K \in \partial^{-1}F} h_K^{1/2} |v_h|_{m,K}.$$

This ends the proof. $\square$

Given the virtual triangulation $\mathcal{T}_h^*$, for each nonnegative integer $r < m$, define the tensorial $(m - r)$ th order Lagrange element space associated with $\mathcal{T}_h^*$:

$$S_h^{m,r} := \{\tau_h \in H_0^1(\Omega; \mathbb{T}_n(r)) : \tau_h|_K \in \mathbb{P}_{m-r}(K; \mathbb{T}_n(r)) \quad \forall K \in \mathcal{T}_h^*\}.$$

Lemma 4.7. *Let $r = 0, 1, \dots, m-1$. For any $v_h \in V_h$, there exists $\tau_r = \tau_r(v_h) \in S_h^{m,r}$ such that*

$$(4.15) \quad |\nabla_h^r v_h - \tau_r|_{j,h} \lesssim h^{m-r-j} |v_h|_{m,h} \quad \text{for } j = 0, 1, \dots, m-r.$$

Proof. Let $w_h \in L^2(\Omega; \mathbb{T}_n(r))$ be defined as

$$w_h|_K = T_{m-r-1}^K(\nabla^r(v_h|_K)) \quad \forall K \in \mathcal{T}_h^*.$$

Since w_h is a piecewise tensorial polynomial, by Lemma 3.1 in [37], there exists $\tau_r \in S_h^{m,r}$ such that

$$|w_h - \tau_r|_{j,h}^2 \lesssim \sum_{F \in \mathcal{F}_h^1(\mathcal{T}_h^*)} h_F^{1-2j} \|[[w_h]]\|_{0,F}^2,$$

where $\mathcal{F}_h^1(\mathcal{T}_h^*)$ is the set of all $(n-1)$ -dimensional faces of the partition $\mathcal{T}_h^*$. Then it follows from (4.11) and (4.14) that

$$\begin{aligned} |w_h - \tau_r|_{j,h}^2 &\lesssim \sum_{F \in \mathcal{F}_h^1(\mathcal{T}_h^*)} h_F^{1-2j} \|[[w_h - \nabla_h^r v_h]]\|_{0,F}^2 + \sum_{F \in \mathcal{F}_h^1} h_F^{1-2j} \|[[\nabla_h^r v_h]]\|_{0,F}^2 \\ &\lesssim h^{2(m-r-j)} |v_h|_{m,h}^2. \end{aligned}$$

Here we have used the fact that the jump $[[\nabla_h^r v_h]]$ is zero on $F \in \mathcal{F}_h^1(\mathcal{T}_h^*) \setminus \mathcal{F}_h^1(\mathcal{T}_h)$. Applying (4.11) again gives

$$|\nabla_h^r v_h - w_h|_{j,h} \lesssim h^{m-r-j} |v_h|_{m,h}.$$

Finally, combining the last two inequalities indicates (4.15). $\square$

Lemma 4.8. *We have the discrete Poincaré inequality*

$$(4.16) \quad \|v_h\|_{m,h} \lesssim |v_h|_{m,h} \quad \forall v_h \in V_h.$$

Proof. By picking $\tau_r \in H_0^1(\Omega; \mathbb{T}_n(r))$ as in Lemma 4.7, due to (4.15) and the Poincaré inequality, we have for $r = 0, 1, \dots, m-1$,

$$\begin{aligned} \|\nabla_h^r v_h\|_0 &\leq \|\nabla_h^r v_h - \tau_r\|_0 + \|\tau_r\|_0 \lesssim |v_h|_{m,h} + |\tau_r|_1 \\ &\leq |v_h|_{m,h} + |\nabla_h^r v_h - \tau_r|_{1,h} + |\nabla_h^r v_h|_{1,h} \\ &\lesssim |v_h|_{m,h} + \|\nabla_h^{r+1} v_h\|_0, \end{aligned}$$

which leads to (4.16). $\square$

The discrete Poincaré inequality (4.16) means

$$\|v_h\|_{m,h} \approx |v_h|_{m,h} \quad \forall v_h \in V_h,$$

i.e., $|\cdot|_{m,h}$ is a norm on the space V_h .

4.4. Norm equivalence and well-posedness of the discretization. Denote by $\ker(\Pi^K) \subset W_k(K)$ the kernel space of the operator Π^K . By (3.13) and Lemma 3.9, both $|\cdot|_{m,K}$ and $S_K^{1/2}(\cdot, \cdot)$ are norms on the finite-dimensional space $\ker(\Pi^K)$. Then we have the following norm equivalence.

Theorem 4.9. *Assume the mesh $\mathcal{T}_h$ satisfies conditions (A1) and (A2). For any $K \in \mathcal{T}_h$, the following norm equivalence holds:*

$$(4.17) \quad S_K(v, v) \approx |v|_{m,K}^2 \quad \forall v \in \ker(\Pi^K),$$

where the constant is independent of h_K , but may depend on the chunkiness parameter ρ_K , the degree of polynomials k , the order of differentiation m , the dimension

of space n , and the shape regularity and quasiuniform constants of the virtual triangulation $\mathcal{T}_h^*$.

Using the generalized scaling argument, i.e., applying an affine map $\hat{\mathbf{x}} = (\mathbf{x} - \mathbf{x}_K)/h_K$, it is easy to show that the norm equivalence constant is independent of the diameter of K .

The constant in (4.17), however, could still depend on the geometry of K , and a clear dependence is not easy to characterize. For finite element space defined on simplexes, the shape functions are usually polynomials and there exists an affine map to the reference element $\hat{K}$. The norm equivalence on the reference element can be used. Since the Jacobi matrix is constant, the norms $H^m(K)$ and $H^m(\hat{K})$ can be clearly characterized using the geometry of the simplex, e.g., the angles of a triangle in 2D.

Now for a general polytope K , there does not exist an affine-equivalent reference polytope $\hat{K}$. For a star-shaped and Lipschitz continuous domain, one can use the isomorphism $\Phi : K \rightarrow B_K$, but $\Phi \in W^{1,\infty}(B_K)$ only. One can apply the norm equivalence on B_K , but how the norm $H^m(K)$ is related to $H^m(B_K)$, for $m > 1$, is not clear.

We shall prove the norm equivalence (4.17) with mesh conditions (A1)–(A2) in Appendix A.

By the Cauchy–Schwarz inequality and the norm equivalence (4.17), we have

$$(4.18) \quad S_K(w, v) \lesssim |w|_{m,K} |v|_{m,K} \quad \forall w, v \in \ker(\Pi^K),$$

which implies the continuity of $a_h(\cdot, \cdot)$:

$$(4.19) \quad a_h(w_h, v_h) \lesssim |w_h|_{m,h} |v_h|_{m,h} \quad \forall w_h, v_h \in V_h + \mathbb{P}_k(\mathcal{T}_h).$$

Next we verify the coercivity of $a_h(\cdot, \cdot)$.

Lemma 4.10. *For any $v_h \in V_h + \mathbb{P}_k(\mathcal{T}_h)$, it holds that*

$$(4.20) \quad |v_h|_{m,h}^2 \lesssim a_h(v_h, v_h).$$

Proof. Since Π^K is the H^m -orthogonal projection,

$$|v_h|_{m,K}^2 = |\Pi^K(v_h|_K)|_{m,K}^2 + |v_h - \Pi^K(v_h|_K)|_{m,K}^2.$$

Applying (4.17), we have

$$(4.21) \quad \begin{aligned} |v_h|_{m,K}^2 &\lesssim |\Pi^K(v_h|_K)|_{m,K}^2 + S_K(v_h - \Pi^K(v_h|_K), v_h - \Pi^K(v_h|_K)) \\ &= a_{h,K}(v_h, v_h), \end{aligned}$$

which implies (4.20). $\square$

Therefore the nonconforming virtual element method (4.5) is uniquely solvable by the Lax–Milgram lemma.

5. ERROR ANALYSIS

In this section, we will develop the error analysis of the nonconforming virtual element method (4.5) for the H^m -problem.

5.1. Interpolation error estimate. We first explore a discrete Galerkin orthogonality of $u - I_h u$ for the nonconforming element, where I_h defined on $H^m(\Omega)$ is the global canonical interpolation operator based on the degrees of freedom in (3.3)–(3.4), i.e., $(I_h v)|_K := I_K(v|_K)$ for any $v \in H^m(\Omega)$ and $K \in \mathcal{T}_h$. A similar result was given in [28, (3.3)].

Lemma 5.1. *For each $K \in \mathcal{T}_h$, any $v \in H^m(\Omega)$, and $w \in H^m(K)$, it holds that*

$$(5.1) \quad a_{h,K}(v - I_h v, w) = 0.$$

Proof. It follows from (3.15) and the definition of $S_K(\cdot, \cdot)$ that

$$\begin{aligned} a_{h,K}(v - I_h v, w) &= (\nabla^m \Pi^K(v - I_h v), \nabla^m \Pi^K w)_K \\ &\quad + S_K(v - I_h v - \Pi^K(v - I_h v), w - \Pi^K w) \\ &= S_K(v - I_h v, w - \Pi^K w) = 0. \end{aligned}$$

In the last step we use the fact that v and $I_h v$ share the same degrees of freedom, and thus the stabilization $S_K(v - I_h v, w - \Pi^K w)$ using d.o.f. vanishes; cf. (4.3). $\square$

Remark 5.2. Lemma 5.1 holds true by virtue of the choice (4.3); indeed, any stabilization equivalent to (4.3) which annihilates if all the degrees of freedom are zero would be fine.

With the help of the discrete Galerkin orthogonality, we present the following interpolation error estimate.

Lemma 5.3. *For each $K \in \mathcal{T}_h$ and any $v \in H^{k+1}(K)$, we have*

$$(5.2) \quad |v - I_h v|_{m,K} \lesssim h_K^{k+1-m} |v|_{k+1,K}.$$

Proof. Applying (4.21) and (5.1) with $w = (T_h v - I_h v)|_K$, we have

$$\begin{aligned} |T_h v - I_h v|_{m,K}^2 &\lesssim a_{h,K}(T_h v - I_h v, T_h v - I_h v) = a_{h,K}(T_h v - v, T_h v - I_h v) \\ &\lesssim |v - T_h v|_{m,K} |T_h v - I_h v|_{m,K}, \end{aligned}$$

which indicates

$$|T_h v - I_h v|_{m,K} \lesssim |v - T_h v|_{m,K}.$$

Hence

$$|v - I_h v|_{m,K} \leq |v - T_h v|_{m,K} + |T_h v - I_h v|_{m,K} \lesssim |v - T_h v|_{m,K}.$$

Therefore (5.2) follows from (4.11). $\square$

5.2. Consistency error estimate. Due to (3.13) and (3.10), we have the following k -consistency.

Lemma 5.4. *For any $p \in \mathbb{P}_k(K)$ and any $v \in W_k(K)$, it holds that*

$$(5.3) \quad a_{h,K}(p, v) = (\nabla^m p, \nabla^m v)_K.$$

To estimate the consistency error of the discretization, we split it into two cases, i.e., $k \geq 2m - 1$ and $m \leq k < 2m - 1$. For the first case, the weak continuity (4.7), that is the projection $Q_{k-(m+i)}^F(\nabla_h^{m-(i+1)} v_h)$ is continuous across $F \in \mathcal{F}_h^1$ for $i = 0, \dots, m - 1$, is sufficient to derive the optimal consistency error estimate.

Lemma 5.5. *Let $u \in H_0^m(\Omega) \cap H^{k+1}(\Omega)$ be the solution of the polyharmonic equation (4.1). Assume $k \geq 2m - 1$. Then it holds that*

$$(5.4) \quad (\nabla^m u, \nabla_h^m v_h) - (f, v_h) \lesssim h^{k+1-m} |u|_{k+1} |v_h|_{m,h} \quad \forall v_h \in V_h.$$

Proof. First we notice that

$$(5.5) \quad (\nabla^m u, \nabla_h^m v_h) - (f, v_h) = \sum_{i=0}^{m-1} (-1)^i \left((\operatorname{div}^i \nabla^m u, \nabla_h^{m-i} v_h) + (\operatorname{div}^{i+1} \nabla^m u, \nabla_h^{m-(i+1)} v_h) \right).$$

For each term on the right-hand side of (5.5), applying integration by parts, (4.7) with $s = m - (i + 1)$, (4.10), and (4.14), we get

$$\begin{aligned} & (-1)^i \left((\operatorname{div}^i \nabla^m u, \nabla_h^{m-i} v_h) + (\operatorname{div}^{i+1} \nabla^m u, \nabla_h^{m-(i+1)} v_h) \right) \\ &= (-1)^i \sum_{K \in \mathcal{T}_h} ((\operatorname{div}^i \nabla^m u) \nu, \nabla_h^{m-(i+1)} v_h)_{\partial K} \\ &= (-1)^i \sum_{F \in \mathcal{F}_h^1} ((\operatorname{div}^i \nabla^m u) \nu_{F,1}, [\nabla_h^{m-(i+1)} v_h])_F \\ &= (-1)^i \sum_{F \in \mathcal{F}_h^1} ((\operatorname{div}^i \nabla^m u) \nu_{F,1} - Q_{k-(m+i)}^F((\operatorname{div}^i \nabla^m u) \nu_{F,1}), [\nabla_h^{m-(i+1)} v_h])_F \\ &\lesssim h^{k+1-m} |u|_{k+1} |v_h|_{m,h}, \end{aligned}$$

as required. $\square$

When the order k is not high enough, the mean value of $\nabla_h^s v_h$ is only continuous over some low-dimensional face of F for $s < 2m - 1 - k$. In this case, we divide the consistency error into two parts. The first part is estimated by using the weak continuity (4.7) as in the proof of (5.4), while the second part is estimated by using the weak continuity (4.8) through employing the Lagrange element space as a bridge.

Lemma 5.6. *Let $u \in H_0^m(\Omega) \cap H^{2m-1}(\Omega)$ be the solution of the polyharmonic equation (4.1). Assume $m \leq k < 2m - 1$. Then it holds that*

$$(5.6) \quad (\nabla^m u, \nabla_h^m v_h) - (f, v_h) \lesssim \left(\sum_{i=k+1-m}^{m-1} h^i |u|_{m+i} + h^m \|f\|_0 \right) |v_h|_{m,h} \quad \forall v_h \in V_h.$$

Proof. Similarly as in (5.5), we have

$$(5.7) \quad (\nabla^m u, \nabla_h^m v_h) - (f, v_h) = E_1 + E_2 + E_3,$$

where

$$\begin{aligned} E_1 &:= \sum_{i=0}^{k-m} (-1)^i \left((\operatorname{div}^i \nabla^m u, \nabla_h^{m-i} v_h) + (\operatorname{div}^{i+1} \nabla^m u, \nabla_h^{m-(i+1)} v_h) \right), \\ E_2 &:= \sum_{i=k-m+1}^{m-2} (-1)^i \left((\operatorname{div}^i \nabla^m u, \nabla_h^{m-i} v_h) + (\operatorname{div}^{i+1} \nabla^m u, \nabla_h^{m-(i+1)} v_h) \right), \\ E_3 &:= ((-\operatorname{div})^{m-1} \nabla^m u, \nabla_h v_h) - (f, v_h). \end{aligned}$$

By the same argument as in the proof of Lemma 5.5, we have

$$(5.8) \quad E_1 \lesssim h^{k+1-m} |u|_{k+1} |v_h|_{m,h}.$$

Next let us estimate E_2 and E_3 . By (4.15), for each $k - m + 1 \leq i \leq m - 1$, there exists $\tau_{m-(i+1)} \in S_h^{m, m-(i+1)}$ such that

$$(5.9) \quad |\nabla_h^{m-(i+1)} v_h - \tau_{m-(i+1)}|_{j,h} \lesssim h^{i+1-j} |v_h|_{m,h} \quad \text{for } j = 0, 1.$$

Since $\tau_{m-(i+1)} \in H_0^1(\Omega; \mathbb{T}_n(m - (i + 1)))$, we get for $i = k - m + 1, \dots, m - 2$

$$(5.10) \quad (\operatorname{div}^i \nabla^m u, \nabla \tau_{m-(i+1)}) + (\operatorname{div}^{i+1} \nabla^m u, \tau_{m-(i+1)}) = 0,$$

$$(5.11) \quad ((-\operatorname{div})^{m-1} \nabla^m u, \nabla \tau_0) - (f, \tau_0) = 0.$$

For $i = k - m + 1, \dots, m - 2$, it follows from (5.9)–(5.10) that

$$\begin{aligned} & (-1)^i \left((\operatorname{div}^i \nabla^m u, \nabla_h^{m-i} v_h) + (\operatorname{div}^{i+1} \nabla^m u, \nabla_h^{m-(i+1)} v_h) \right) \\ &= (-1)^i (\operatorname{div}^i \nabla^m u, \nabla_h (\nabla_h^{m-(i+1)} v_h - \tau_{m-(i+1)})) \\ & \quad + (-1)^i (\operatorname{div}^{i+1} \nabla^m u, \nabla_h^{m-(i+1)} v_h - \tau_{m-(i+1)}) \\ & \lesssim h^i |u|_{m+i} |v_h|_{m,h} + h^{i+1} |u|_{m+i+1} |v_h|_{m,h}. \end{aligned}$$

Thus we obtain

$$(5.12) \quad E_2 \lesssim \sum_{i=k+1-m}^{m-1} h^i |u|_{m+i} |v_h|_{m,h}.$$

Similarly, employing (5.11) and (5.9), we get

$$\begin{aligned} E_3 &= ((-\operatorname{div})^{m-1} \nabla^m u, \nabla_h v_h) - (f, v_h) \\ &= ((-\operatorname{div})^{m-1} \nabla^m u, \nabla_h (v_h - \tau_0)) - (f, v_h - \tau_0) \\ &\lesssim h^{m-1} |u|_{2m-1} |v_h|_{m,h} + h^m \|f\|_0 |v_h|_{m,h}, \end{aligned}$$

which together with (5.7)–(5.8) and (5.12) ends the proof. $\square$

We then consider the perturbation of the right-hand side. Namely, replace the L^2 -inner product (f, v_h) by an approximated one $\langle f, v_h \rangle$ defined in (4.4).

Lemma 5.7. *Let $u \in H_0^m(\Omega) \cap H^r(\Omega)$ with $r = \max\{k+1, 2m-1\}$ be the solution of the polyharmonic equation (4.1). Assume $f \in H^\ell(\mathcal{T}_h)$ with $\ell = \max\{0, k+1-2m\}$. Then it holds for any $v_h \in V_h$ that*

$$(5.13) \quad \begin{aligned} & (\nabla^m u, \nabla_h^m v_h) - \langle f, v_h \rangle \\ & \lesssim h^{k+1-m} (\|u\|_r + h\|f\|_0 + h^{\max\{0, 2m-k-1\}} |f|_{\ell,h}) |v_h|_{m,h}. \end{aligned}$$

Proof. It follows from (5.4) and (5.6) that

$$(\nabla^m u, \nabla_h^m v_h) - (f, v_h) \lesssim h^{k+1-m} (\|u\|_r + h\|f\|_0) |v_h|_{m,h}.$$

For $m \leq k \leq 2m - 1$, we get from the local Poincaré inequality (A.11)

$$(f, v_h) - \langle f, v_h \rangle = (f, v_h - \Pi_h v_h) \lesssim h^m \|f\|_0 |v_h|_{m,h}.$$

For $k \geq 2m$, it holds from (4.9) that

$$\begin{aligned}
 (f, v_h) - \langle f, v_h \rangle &= \left(f, v_h - Q_h^{\max\{m-1, k-2m\}} v_h \right) \\
 &= \left(f - Q_h^{k-2m} f, v_h - Q_h^{\max\{m-1, k-2m\}} v_h \right) \\
 &\leq \|f - Q_h^{k-2m} f\|_0 \|v_h - Q_h^{m-1} v_h\|_0 \\
 &\lesssim h^{k+1-m} |f|_{k+1-2m, h} |v_h|_{m, h}.
 \end{aligned}$$

Thus we conclude (5.13) from the last three inequalities. $\square$

5.3. Error estimate. Now we are in a position to present the optimal order convergence of our nonconforming virtual element method.

Theorem 5.8. *Let $u \in H_0^m(\Omega) \cap H^r(\Omega)$ with $r = \max\{k+1, 2m-1\}$ be the solution of the polyharmonic equation (4.1), and let $u_h \in V_h$ be the nonconforming virtual element method (4.5). Assume the mesh $\mathcal{T}_h$ satisfies conditions (A1) and (A2). Assume $f \in H^\ell(\mathcal{T}_h)$ with $\ell = \max\{0, k+1-2m\}$. Then it holds that*

$$(5.14) \quad |u - u_h|_{m, h} \lesssim h^{k+1-m} (\|u\|_r + h\|f\|_0 + h^{\max\{0, 2m-k-1\}} |f|_{\ell, h}).$$

Proof. Let $v_h = I_h u - u_h$. From (4.19), (5.2), and (4.11), it holds that

$$\begin{aligned}
 &a_h(I_h u - T_h u, v_h) + (\nabla_h^m(T_h u - u), \nabla_h^m v_h) \\
 &\lesssim |I_h u - T_h u|_{m, h} |v_h|_{m, h} + |u - T_h u|_{m, h} |v_h|_{m, h} \\
 (5.15) \quad &\lesssim (|u - I_h u|_{m, h} + |u - T_h u|_{m, h}) |v_h|_{m, h} \lesssim h^{k+1-m} |u|_{k+1} |v_h|_{m, h}.
 \end{aligned}$$

Employing (4.20), (4.5), and (5.3), we have

$$\begin{aligned}
 |I_h u - u_h|_{m, h}^2 &\lesssim a_h(I_h u - u_h, v_h) = a_h(I_h u, v_h) - \langle f, v_h \rangle \\
 &= a_h(I_h u - T_h u, v_h) + a_h(T_h u, v_h) - \langle f, v_h \rangle \\
 &= a_h(I_h u - T_h u, v_h) + (\nabla_h^m T_h u, \nabla_h^m v_h) - \langle f, v_h \rangle \\
 &= a_h(I_h u - T_h u, v_h) + (\nabla_h^m(T_h u - u), \nabla_h^m v_h) \\
 &\quad + (\nabla_h^m u, \nabla_h^m v_h) - \langle f, v_h \rangle.
 \end{aligned}$$

Then we get from (5.15) and (5.13) that

$$|I_h u - u_h|_{m, h} \lesssim h^{k+1-m} (\|u\|_r + h\|f\|_0 + h^{\max\{0, 2m-k-1\}} |f|_{\ell, h}).$$

Finally we derive (5.14) by combining the last inequality and (5.2). $\square$

6. CONCLUSION

In view of a generalized Green's identity for the H^m inner product, we have constructed the H^m -nonconforming virtual elements of any order k on any shape of polytope in $\mathbb{R}^n$ with constraints $m \leq n$ and $k \geq m$ in a unified way. A rigorous and detailed convergence analysis is developed for the H^m -nonconforming virtual element methods, and the optimal error estimates are achieved. When $m > n$, the generalized Green's identity for the H^m inner product, the key tool in this paper, will involve the derivative terms on zero-dimensional subsurfaces, i.e., nodes of the mesh. We will postpone the case $m > n$ to future works.

This paper was motivated by the theoretical purposes. The numerical investigation of the virtual element method proposed in this paper is also postponed to future works.

APPENDIX A. NORM EQUIVALENCE

As mentioned before, it is difficult to derive the norm equivalence (4.17) directly from the norm equivalence of the finite-dimensional space due to the absence of an affine-equivalent reference polytope. We shall prove the norm equivalence (4.17) in this appendix by assuming that the mesh $\mathcal{T}_h$ satisfies conditions (A1) and (A2). For $m = 1$, proofs on the norm equivalence for H^1 conforming VEM can be found in [14, 17, 22].

With the help of the virtual triangulation $\mathcal{T}_h^*$, we can prove the inverse inequality of polynomial spaces on K following the proof in [22, Lemma 3.1]:

$$(A.1) \quad \|g\|_{0,K} \lesssim h_K^{-i} \|g\|_{-i,K} \quad \forall g \in \mathbb{P}_k(K), \quad i = 1, 2, \dots, m,$$

where the constant depends only on the degree of polynomials k , the order of differentiation m , the dimension of space n , and the shape regularity and quasi-uniformity of the virtual triangulation $\mathcal{T}_h^*(K)$.

On the polynomial space, we have the normal equivalence of the L^2 -norm of g and l^2 -norm of its d.o.f. Let $g = \sum_i g_i m_i$ be a polynomial on F , where $F \in \mathcal{F}^j(K)$ with $j \geq 1$. Denote by $\mathbf{g} = (g_i)$ the coefficient vector. Then the following norm equivalence holds (cf. [22, Lemma 4.1]):

$$(A.2) \quad h_F^{(n-j)/2} \|\mathbf{g}\|_{l^2} \lesssim \|g\|_{0,F} \lesssim h_F^{(n-j)/2} \|\mathbf{g}\|_{l^2}.$$

Take an element $K \in \mathcal{T}_h$. For any $F \in \mathcal{F}^j(K)$ with $j \geq 1$, let $\mathbb{R}_F^{n-j}$ be the $(n-j)$ -dimensional affine space passing through F , $\mathcal{F}_F^j(K) := \{e \in \mathcal{F}^j(K) : e \subset \mathbb{R}_F^{n-j}\}$, and let

$$\lambda_{F,i} := \nu_{F,i}^\top \frac{\mathbf{x} - \mathbf{x}_F}{h_K}, \quad i = 1, \dots, j.$$

Apparently $\lambda_{F,i}|_F = 0$, i.e., the $(n-1)$ -dimensional face $\lambda_{F,i} = 0$ passes through F . If K is a simplex and $F \in \mathcal{F}^1(K)$, $-(\nu_{F,1}^\top \nu_{K,F}) \lambda_{F,1}$ is just the barycenter coordinate when h_K represents the height of K corresponding to the base F , and $\mathcal{F}_F^j(K) = \{F\}$ if K is strictly convex. For any $F \in \mathcal{F}^j(K)$ with $j \geq 1$, and $F' \in \mathcal{F}^j(K) \setminus \mathcal{F}_F^j(K)$, let $\nu_{F,F'}$ be some unit normal vector of F' such that the hyperplane $\nu_{F,F'}^\top (\mathbf{x} - \mathbf{x}_{F'}) = 0$ does not pass through F . Define bubble functions

$$b_K := \prod_{F \in \mathcal{F}^1(K)} \lambda_{F,1},$$

$$b_F := \left(\prod_{F' \in \mathcal{F}^j(K) \setminus \mathcal{F}_F^j(K)} \nu_{F,F'}^\top \frac{\mathbf{x} - \mathbf{x}_{F'}}{h_K} \right) \left(\prod_{F' \in \mathcal{F}_F^j(K)} \prod_{e \in \mathcal{F}^1(F')} \nu_{F',e}^\top \frac{\mathbf{x} - \mathbf{x}_e}{h_K} \right),$$

for each $F \in \mathcal{F}^j(K)$ with $1 \leq j \leq n$. Notice that both b_K and b_F are polynomials.

Lemma A.1. *Let $K \in \mathcal{T}_h$. It holds that*

$$(A.3) \quad h_K^m \|(-\Delta)^m v\|_{0,K} \lesssim \|\nabla^m v\|_{0,K} \quad \forall v \in V_k(K) \cup W_k(K).$$

Proof. Let $\phi_K := b_K^{2m} (-\Delta)^m v \in H_0^m(K)$; then $\|\phi_K\|_{0,K} \approx \|(-\Delta)^m v\|_{0,K}$. Using the scaling argument, integration by parts, and the inverse inequality for polynomials (A.1), we get

$$\begin{aligned} \|(-\Delta)^m v\|_{0,K}^2 &\lesssim ((-\Delta)^m v, \phi_K)_K = (\nabla^m v, \nabla^m \phi_K)_K \\ &\lesssim \|\nabla^m v\|_{0,K} \|\nabla^m \phi_K\|_{0,K} \lesssim h_K^{-m} \|\nabla^m v\|_{0,K} \|\phi_K\|_{0,K} \\ &\lesssim h_K^{-m} \|\nabla^m v\|_{0,K} \|(-\Delta)^m v\|_{0,K}, \end{aligned}$$

which induces the required inequality. $\square$

Lemma A.2. *Let $K \in \mathcal{T}_h$. For any positive integer $j \leq m$, $F \in \mathcal{F}^j(K)$, and $\alpha \in A_j$ with $|\alpha| \leq m - j$, we have*

$$\begin{aligned}
 & \sum_{F' \in \mathcal{F}_F^j(K)} h_K^{m-|\alpha|-j/2} \|D_{F',\alpha}^{2m-j-|\alpha|}(v)\|_{0,F'} \\
 & \lesssim \|\nabla^m v\|_{0,K} + h_K^m \|(-\Delta)^m v\|_{0,K} \\
 & + \sum_{\ell=1}^{j-1} \sum_{e \in \mathcal{F}^\ell(K)} \sum_{\substack{\beta \in A_\ell \\ |\beta| \leq m-\ell}} h_K^{m-|\beta|-\ell/2} \|D_{e,\beta}^{2m-\ell-|\beta|}(v)\|_{0,e} \\
 (A.4) \quad & + \sum_{F' \in \mathcal{F}_F^j(K)} \sum_{\substack{\beta \in A_j \\ |\alpha| < |\beta| \leq m-j}} h_K^{m-|\beta|-\ell/2} \|D_{F',\beta}^{2m-j-|\beta|}(v)\|_{0,F'}
 \end{aligned}$$

for all $v \in V_k(K) \cup W_k(K)$.

Proof. Since $D_{F',\alpha}^{2m-j-|\alpha|}(v)|_{F'}$ is a polynomial for each $F' \in \mathcal{F}_F^j(K)$, we can regard $D_{F',\alpha}^{2m-j-|\alpha|}(v)|_{F'}$ as the function on the $(n-j)$ -dimensional affine space $\mathbb{R}_F^{n-j}$. Then we extend the polynomial $D_{F',\alpha}^{2m-j-|\alpha|}(v)|_{F'}$ to $\mathbb{R}^n$. For any $\mathbf{x} \in \mathbb{R}^n$, let $\mathbf{x}_F^P$ be the projection of $\mathbf{x}$ on $\mathbb{R}_F^{n-j}$. Define

$$E_K(D_{F',\alpha}^{2m-j-|\alpha|}(v))(\mathbf{x}) := D_{F',\alpha}^{2m-j-|\alpha|}(v)(\mathbf{x}_F^P).$$

Let $\mathbb{R}_{F'}^n := \{\mathbf{x} \in \mathbb{R}^n : \mathbf{x}_F^P \in F'\}$, and let ϕ_F be a piecewise polynomial defined as

$$\phi_F(\mathbf{x}) = \begin{cases} \frac{1}{\alpha!} h_K^{|\alpha|} b_{F'}^{2m} E_K(D_{F',\alpha}^{2m-j-|\alpha|}(v)) \prod_{i=1}^j \lambda_{F',i}^{\alpha_i}, & \mathbf{x} \in \mathbb{R}_{F'}^n, F' \in \mathcal{F}_F^j(K), \\ 0, & \mathbf{x} \in \mathbb{R}^n \setminus \bigcup_{F' \in \mathcal{F}_F^j(K)} \mathbb{R}_{F'}^n, \end{cases}$$

where $\alpha! = \alpha_1! \cdots \alpha_j!$. Then we have

$$(A.5) \quad \|\phi_F\|_{0,K} \lesssim \sum_{F' \in \mathcal{F}_F^j(K)} h_K^{|\alpha|+j/2} \|D_{F',\alpha}^{2m-j-|\alpha|}(v)\|_{0,F'},$$

$$\begin{aligned}
 \left. \frac{\partial^{|\alpha|} \phi_F}{\partial \nu_{F'}^\alpha} \right|_{F'} &= \frac{1}{\alpha!} h_K^{|\alpha|} b_{F'}^{2m} E_K(D_{F',\alpha}^{2m-j-|\alpha|}(v)) \prod_{i=1}^j \frac{\partial^{|\alpha|} (\lambda_{F',i}^{\alpha_i})}{\partial \nu_{F'}^\alpha} \\
 &= b_{F'}^{2m} E_K(D_{F',\alpha}^{2m-j-|\alpha|}(v)).
 \end{aligned}$$

Hence

$$(A.6) \quad \|D_{F',\alpha}^{2m-j-|\alpha|}(v)\|_{0,F'}^2 \approx \left(D_{F',\alpha}^{2m-j-|\alpha|}(v), \frac{\partial^{|\alpha|} \phi_F}{\partial \nu_{F'}^\alpha} \right)_{F'}.$$

For each $e \in \mathcal{F}^\ell(K)$ with $\ell = j+1, \dots, m$, it follows from the fact $b_{F'}|_e = 0$ that

$$\left. \frac{\partial^{|\beta|} \phi_F}{\partial \nu_e^\beta} \right|_e = 0 \quad \forall \beta \in A_\ell \text{ with } |\beta| \leq m - \ell.$$

Similarly we have for each $e \in \mathcal{F}^j(K) \setminus \mathcal{F}_F^j(K)$ that

$$\left. \frac{\partial^{|\beta|} \phi_F}{\partial \nu_e^\beta} \right|_e = 0 \quad \forall \beta \in A_j \text{ with } |\beta| \leq m - j.$$

For any $\beta \in A_j$, $|\beta| < |\alpha|$, since $\frac{\partial^{|\beta|}}{\partial \nu_{F'}^\beta} \left(\prod_{i=1}^j \lambda_{F',i}^{\alpha_i} \right) \Big|_{F'} = 0$, it yields $\frac{\partial^{|\beta|} \phi_F}{\partial \nu_{F'}^\beta} \Big|_{F'} = 0$. For any $\beta \in A_j$, $|\beta| = |\alpha|$, but $\beta \neq \alpha$, noting that $\frac{\partial \lambda_{F',i}}{\partial \nu_{F',\ell}} = 0$ for $i \neq \ell$, we also have $\frac{\partial^{|\beta|} \phi_F}{\partial \nu_{F'}^\beta} \Big|_{F'} = 0$. Based on the previous discussion, we obtain from (A.6), the generalized Green's identity (3.2), and the density argument that

$$\begin{aligned} \sum_{F' \in \mathcal{F}_F^j(K)} \|D_{F',\alpha}^{2m-j-|\alpha|}(v)\|_{0,F'}^2 &\simeq (\nabla^m v, \nabla^m \phi_F)_K - ((-\Delta)^m v, \phi_F)_K \\ &\quad - \sum_{\ell=1}^{j-1} \sum_{e \in \mathcal{F}^\ell(K)} \sum_{\substack{\beta \in A_\ell \\ |\beta| \leq m-\ell}} \left(D_{e,\beta}^{2m-\ell-|\beta|}(v), \frac{\partial^{|\beta|} \phi_F}{\partial \nu_e^\beta} \right)_e \\ &\quad - \sum_{F' \in \mathcal{F}_F^j(K)} \sum_{\substack{\beta \in A_j \\ |\alpha| < |\beta| \leq m-j}} \left(D_{F',\beta}^{2m-j-|\beta|}(v), \frac{\partial^{|\beta|} \phi_F}{\partial \nu_{F'}^\beta} \right)_{F'}. \end{aligned}$$

Employing the Cauchy-Schwarz inequality and the inverse inequality for polynomials, it follows that

$$\begin{aligned} &\sum_{F' \in \mathcal{F}_F^j(K)} \|D_{F',\alpha}^{2m-j-|\alpha|}(v)\|_{0,F'}^2 \\ &\lesssim h_K^{-m} \|\nabla^m v\|_{0,K} \|\phi_F\|_{0,K} + \|(-\Delta)^m v\|_{0,K} \|\phi_F\|_{0,K} \\ &\quad + \sum_{\ell=1}^{j-1} \sum_{e \in \mathcal{F}^\ell(K)} \sum_{\substack{\beta \in A_\ell \\ |\beta| \leq m-\ell}} h_K^{-|\beta|-\ell/2} \|D_{e,\beta}^{2m-\ell-|\beta|}(v)\|_{0,e} \|\phi_F\|_{0,K} \\ &\quad + \sum_{F' \in \mathcal{F}_F^j(K)} \sum_{\substack{\beta \in A_j \\ |\alpha| < |\beta| \leq m-j}} h_K^{-|\beta|-j/2} \|D_{F',\beta}^{2m-j-|\beta|}(v)\|_{0,F'} \|\phi_F\|_{0,K}, \end{aligned}$$

which combined with (A.5) implies (A.4). $\square$

Lemma A.3. *For any $K \in \mathcal{T}_h$, it holds that*

$$(A.7) \quad ((-\Delta)^m v, v)_K \lesssim h_K^m \|(-\Delta)^m v\|_{0,K} S_K^{1/2}(v, v) \quad \forall v \in \ker(\Pi^K).$$

Proof. If $m \leq k \leq 2m-1$, by the definition of $W_k(K) = V_k(K)$, $(-\Delta)^m v = 0$, thus (A.7) is obvious. Now let us prove (A.7) for $k \geq 2m$. When $2m \leq k < 3m-1$, since $v \in \ker(\Pi^K)$, it follows from (3.14) that

$$\begin{aligned} ((-\Delta)^m v, v)_K &= ((-\Delta)^m v, Q_{m-1}^K v)_K \\ &= ((-\Delta)^m v, Q_{k-2m}^K v)_K = (Q_{k-2m}^K ((-\Delta)^m v), v)_K. \end{aligned}$$

If $k \geq 3m-1$, then $W_k(K) = V_k(K)$, and we also have

$$((-\Delta)^m v, v)_K = (Q_{k-2m}^K ((-\Delta)^m v), v)_K.$$

Therefore, to derive (A.7) for $k \geq 2m$, it is sufficient to prove that

$$(A.8) \quad (Q_{k-2m}^K ((-\Delta)^m v), v)_K \lesssim h_K^m \|(-\Delta)^m v\|_{0,K} S_K^{1/2}(v, v) \quad \forall v \in \ker(\Pi^K).$$

Let N be the dimension of the space $\mathbb{P}_{k-2m}(K)$. Then there exist constants c_i , $i = 1, \dots, N$, such that

$$Q_{k-2m}^K((-\Delta)^m v) = \sum_{i=1}^N c_i q_i,$$

where $\mathbb{M}_{k-2m}(K) := \{q_1, \dots, q_N\}$; thus

$$(Q_{k-2m}^K((-\Delta)^m v), v)_K = |K| \sum_{i=1}^N c_i \chi_i(v).$$

Applying the norm equivalence on the polynomial space $\mathbb{P}_{k-2m}(K)$ (cf. (A.2)) we get

$$\|Q_{k-2m}^K((-\Delta)^m v)\|_{0,K}^2 \approx h_K^n \sum_{i=1}^N c_i^2.$$

Hence

$$\begin{aligned} (Q_{k-2m}^K((-\Delta)^m v), v)_K &\lesssim h_K^{n/2} \|Q_{k-2m}^K((-\Delta)^m v)\|_{0,K} \left(\sum_{i=1}^N \chi_i^2(v) \right)^{1/2} \\ &\lesssim h_K^m \|Q_{k-2m}^K((-\Delta)^m v)\|_{0,K} S_K^{1/2}(v, v), \end{aligned}$$

which implies (A.8). $\square$

Lemma A.4. *For any $K \in \mathcal{T}_h$, it holds that*

$$(A.9) \quad \|\nabla^m v\|_{0,K}^2 \lesssim S_K(v, v) \quad \forall v \in \ker(\Pi^K).$$

Proof. By the generalized Green's identity (3.2),

$$\begin{aligned} (A.10) \quad \|\nabla^m v\|_{0,K}^2 &= ((-\Delta)^m v, v)_K \\ &+ \sum_{j=1}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} \left(D_{F,\alpha}^{2m-j-|\alpha|}(v), \frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} \right)_F. \end{aligned}$$

Since $v \in W_k(K)$, we have $D_{F,\alpha}^{2m-j-|\alpha|}(v)|_F \in \mathbb{P}_{k-(2m-j-|\alpha|)}(F)$ for any $F \in \mathcal{F}^j(K)$. Let N_F be the dimension of the space $\mathbb{P}_{k-(2m-j-|\alpha|)}(F)$. Then there exist constants c_i , $i = 1, \dots, N_F$, such that

$$\left(D_{F,\alpha}^{2m-j-|\alpha|}(v), \frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} \right)_F = h_K^{n-j-|\alpha|} \sum_{i=1}^{N_F} c_i \chi_i(v).$$

Applying the norm equivalence on the polynomial space $\mathbb{P}_{k-(2m-j-|\alpha|)}(F)$ (cf. (A.2)) we get

$$\|D_{F,\alpha}^{2m-j-|\alpha|}(v)\|_{0,F}^2 \approx h_K^{n-j} \sum_{i=1}^{N_F} c_i^2.$$

Hence

$$\begin{aligned} \left(D_{F,\alpha}^{2m-j-|\alpha|}(v), \frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} \right)_F &\lesssim h_K^{(n-j)/2-|\alpha|} \|D_{F,\alpha}^{2m-j-|\alpha|}(v)\|_{0,F} \left(\sum_{i=1}^{N_F} \chi_i^2(v) \right)^{1/2} \\ &\lesssim h_K^{m-|\alpha|-j/2} \|D_{F,\alpha}^{2m-j-|\alpha|}(v)\|_{0,F} S_K^{1/2}(v, v). \end{aligned}$$

Applying (A.4) recursively, it follows that

$$\left(D_{F,\alpha}^{2m-j-|\alpha|}(v), \frac{\partial^{|\alpha|}v}{\partial\nu_F^\alpha}\right)_F \lesssim (\|\nabla^m v\|_{0,K} + h_K^m \|(-\Delta)^m v\|_{0,K}) S_K^{1/2}(v, v).$$

Then we derive from (A.10), (A.7), and (A.3) that

$$\|\nabla^m v\|_{0,K}^2 \lesssim (\|\nabla^m v\|_{0,K} + h_K^m \|(-\Delta)^m v\|_{0,K}) S_K^{1/2}(v, v) \lesssim \|\nabla^m v\|_{0,K} S_K^{1/2}(v, v),$$

which induces (A.9). $\square$

We then prove another side of the norm equivalence (4.17).

Lemma A.5. *For any $K \in \mathcal{T}_h$ and nonnegative integer $s \leq m$, we have the local Poincaré inequality*

$$(A.11) \quad \sum_{j=0}^{m-s} \sum_{F \in \mathcal{F}^j(K)} h_K^{s+j/2} \|\nabla^s v\|_{0,F} \lesssim h_K^m \|\nabla^m v\|_{0,K} \quad \forall v \in \ker(\Pi^K).$$

Proof. It is sufficient to prove that

$$(A.12) \quad \sum_{j=0}^{m-s} \sum_{F \in \mathcal{F}^j(K)} h_K^{s+j/2} \|\nabla^s v\|_{0,F} \lesssim \sum_{\ell=0}^{m-s-1} \sum_{e \in \mathcal{F}^\ell(K)} h_K^{s+1+\ell/2} \|\nabla^{s+1} v\|_{0,e},$$

for $s = 0, 1, \dots, m-1$. Thanks to (3.11), it follows that

$$\begin{aligned} h_K^{j/2} \|\nabla^s v\|_{0,F} &= h_K^{j/2} \left\| \nabla^s v - \frac{1}{\#\mathcal{F}^{m-s}(K)} \sum_{e \in \mathcal{F}^{m-s}(K)} Q_0^e(\nabla^s v) \right\|_{0,F} \\ &\lesssim h_K^{j/2} \sum_{e \in \mathcal{F}^{m-s}(K)} \|\nabla^s v - Q_0^e(\nabla^s v)\|_{0,F} \\ &= h_K^{j/2} \sum_{e \in \mathcal{F}^{m-s}(K)} \|\nabla^s v - Q_0^K(\nabla^s v) - Q_0^e(\nabla^s v - Q_0^K(\nabla^s v))\|_{0,F} \\ &\lesssim h_K^{j/2} \|\nabla^s v - Q_0^K(\nabla^s v)\|_{0,F} + \sum_{e \in \mathcal{F}^{m-s}(K)} h_K^{(m-s)/2} \|Q_0^e(\nabla^s v - Q_0^K(\nabla^s v))\|_{0,e} \\ &\leq h_K^{j/2} \|\nabla^s v - Q_0^K(\nabla^s v)\|_{0,F} + \sum_{e \in \mathcal{F}^{m-s}(K)} h_K^{(m-s)/2} \|\nabla^s v - Q_0^K(\nabla^s v)\|_{0,e}. \end{aligned}$$

On the other hand, applying the trace inequality (4.6) recursively, we get from (4.9) that

$$\begin{aligned} &h_K^{j/2} \|\nabla^s v - Q_0^K(\nabla^s v)\|_{0,F} + \sum_{e \in \mathcal{F}^{m-s}(K)} h_K^{(m-s)/2} \|\nabla^s v - Q_0^K(\nabla^s v)\|_{0,e} \\ &\lesssim \|\nabla^s v - Q_0^K(\nabla^s v)\|_{0,K} + \sum_{\ell=0}^{m-s-1} \sum_{e \in \mathcal{F}^\ell(K)} h_K^{1+\ell/2} \|\nabla^{s+1} v\|_{0,e} \\ &\lesssim \sum_{\ell=0}^{m-s-1} \sum_{e \in \mathcal{F}^\ell(K)} h_K^{1+\ell/2} \|\nabla^{s+1} v\|_{0,e}. \end{aligned}$$

Combining the last two inequalities yields

$$h_K^{j/2} \|\nabla^s v\|_{0,F} \lesssim \sum_{\ell=0}^{m-s-1} \sum_{e \in \mathcal{F}^\ell(K)} h_K^{1+\ell/2} \|\nabla^{s+1} v\|_{0,e},$$

which indicates (A.12). Thus the Poincaré inequality (A.11) holds. $\square$

Lemma A.6. *For any $K \in \mathcal{T}_h$, it holds that*

$$(A.13) \quad S_K(v, v) \lesssim \|\nabla^m v\|_{0,K}^2 \quad \forall v \in \ker(\Pi^K).$$

Proof. Due to the definition of the degrees of freedom, we have

$$\begin{aligned} S_K(v, v) &= h_K^{n-2m} \sum_{i=1}^{N_K} \chi_i^2(v) \\ &\lesssim \sum_{j=0}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} h_K^{2|\alpha|-2m+j} \left\| Q_{k-(2m-j-|\alpha|)}^F \left(\frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} \right) \right\|_{0,F}^2 \\ &\leq \sum_{j=0}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} h_K^{2|\alpha|-2m+j} \left\| \frac{\partial^{|\alpha|} v}{\partial \nu_F^\alpha} \right\|_{0,F}^2 \\ &\leq \sum_{j=0}^m \sum_{F \in \mathcal{F}^j(K)} \sum_{\substack{\alpha \in A_j \\ |\alpha| \leq m-j}} h_K^{2|\alpha|-2m+j} \|\nabla^{|\alpha|} v\|_{0,F}^2, \end{aligned}$$

which together with the Poincaré inequality (A.11) implies (A.13). $\square$

At last, combining (A.9) and (A.13) gives the norm equivalence (4.17); cf. Theorem 4.9.

APPENDIX B. EXAMPLES OF GREEN'S FORMULA

Take $K \in \mathcal{T}_h$. The explicit expression of (3.2) for $m = 1$ with $n \geq m$ is no more than (2.2), i.e.,

$$(\nabla u, \nabla v)_K = -(\Delta u, v)_K + \sum_{F \in \mathcal{F}^1(K)} \left(\frac{\partial u}{\partial \nu_{K,F}}, v \right)_F \quad \forall u \in H^2(K), v \in H^1(K).$$

Also, the explicit expression of (3.2) for $m = 2$ with $n \geq m$ is exactly (2.5), i.e., for any $u \in H^4(K)$ and $v \in H^2(K)$, it holds that

$$\begin{aligned} (\nabla^2 u, \nabla^2 v)_K &= (\Delta^2 u, v)_K + \sum_{F \in \mathcal{F}^1(K)} \left[\left(M_{\nu\nu}(u), \frac{\partial v}{\partial \nu_{F,1}} \right)_F - (Q_\nu(u), v)_F \right] \\ &\quad + \sum_{e \in \mathcal{F}^2(K)} \sum_{F \in \mathcal{F}^1(K) \cap \partial^{-1}e} (\nu_{F,e}^\top M_{\nu t}(u), v)_e. \end{aligned}$$

When $m = n = 3$, the explicit expression of (3.2) is that for any $u \in H^6(K)$ and $v \in H^3(K)$,

$$\begin{aligned}
& (\nabla^3 u, \nabla^3 v)_K + (\Delta^3 u, v)_K \\
&= \sum_{F \in \mathcal{F}^1(K)} \left(\nu_{K,F}^\top \operatorname{div}^2(\nabla^3 u) + \operatorname{div}_F(\operatorname{div}(\nabla^3 u) \nu_{K,F}) + \operatorname{div}_F \operatorname{div}_F((\nabla^3 u) \nu_{K,F}), v \right)_F \\
&\quad - \sum_{F \in \mathcal{F}^1(K)} \left(\nu_{F,1}^\top (\operatorname{div}_F((\nabla^3 u) \nu_{K,F})) + \operatorname{div}_F(((\nabla^3 u) \nu_{K,F}) \nu_{F,1}), \frac{\partial v}{\partial \nu_{F,1}} \right)_F \\
&\quad - \sum_{F \in \mathcal{F}^1(K)} \left(\nu_{F,1}^\top \operatorname{div}(\nabla^3 u) \nu_{K,F}, \frac{\partial v}{\partial \nu_{F,1}} \right)_F \\
&\quad + \sum_{F \in \mathcal{F}^1(K)} \left(\nu_{F,1}^\top ((\nabla^3 u) \nu_{K,F}) \nu_{F,1}, \frac{\partial^2 v}{\partial \nu_{F,1}^2} \right)_F \\
&\quad - \sum_{F \in \mathcal{F}^1(K)} \sum_{e \in \mathcal{F}^1(F)} \left(\nu_{F,e}^\top (\operatorname{div}_F((\nabla^3 u) \nu_{K,F})) + \operatorname{div}_e(((\nabla^3 u) \nu_{K,F}) \nu_{F,e}), v \right)_e \\
&\quad - \sum_{F \in \mathcal{F}^1(K)} \sum_{e \in \mathcal{F}^1(F)} (\nu_{F,e}^\top \operatorname{div}(\nabla^3 u) \nu_{K,F}, v)_e \\
&\quad + \sum_{F \in \mathcal{F}^1(K)} \sum_{e \in \mathcal{F}^1(F)} \sum_{i=1}^2 \left(\left(\nu_{e,i} + (\nu_{e,i}^\top \nu_{F,1}) \nu_{F,1}^\top \right) ((\nabla^3 u) \nu_{K,F}) \nu_{F,e}, \frac{\partial v}{\partial \nu_{e,i}} \right)_e \\
&\quad + \sum_{F \in \mathcal{F}^1(K)} \sum_{e \in \mathcal{F}^1(F)} \sum_{\delta \in \mathcal{F}^1(e)} \left(\nu_{e,\delta}^\top ((\nabla^3 u) \nu_{K,F}) \nu_{F,e} \right) (\delta) v(\delta).
\end{aligned}$$

Consider the lowest-order case $k = m = 3$. The last identity will be reduced to

$$\begin{aligned}
& (\nabla^3 v, \nabla^3 q)_K \\
&= \sum_{F \in \mathcal{F}^1(K)} \left(\frac{\partial^2 v}{\partial \nu_{F,1}^2}, \nu_{F,1}^\top ((\nabla^3 q) \nu_{K,F}) \nu_{F,1} \right)_F \\
&\quad + \sum_{F \in \mathcal{F}^1(K)} \sum_{e \in \mathcal{F}^1(F)} \sum_{i=1}^2 \left(\frac{\partial v}{\partial \nu_{e,i}}, \left(\nu_{e,i} + (\nu_{e,i}^\top \nu_{F,1}) \nu_{F,1}^\top \right) ((\nabla^3 q) \nu_{K,F}) \nu_{F,e} \right)_e \\
&\quad + \sum_{F \in \mathcal{F}^1(K)} \sum_{e \in \mathcal{F}^1(F)} \sum_{\delta \in \mathcal{F}^1(e)} v(\delta) \left(\nu_{e,\delta}^\top ((\nabla^3 q) \nu_{K,F}) \nu_{F,e} \right) (\delta)
\end{aligned}$$

for any $v \in H^3(K)$ and $q \in \mathbb{P}_3(K)$, which will be used to compute the projector $\Pi^K : H^3(K) \rightarrow \mathbb{P}_3(K)$. The degrees of freedom are

$$\left(\frac{\partial^2 v}{\partial \nu_{F,1}^2}, 1 \right)_F, \quad \left(\frac{\partial v}{\partial \nu_{e,1}}, 1 \right)_e, \quad \left(\frac{\partial v}{\partial \nu_{e,2}}, 1 \right)_e, \quad v(\delta)$$

on each $F \in \mathcal{F}^1(K)$, $e \in \mathcal{F}^2(K)$, and $\delta \in \mathcal{F}^3(K)$.

ACKNOWLEDGMENTS

The authors would like to thank Dr. Shuhao Cao for the insightful discussion and the anonymous referees for their incisive comments.

REFERENCES

- [1] R. A. Adams and J. J. F. Fournier, *Sobolev Spaces*, 2nd ed., Pure and Applied Mathematics (Amsterdam), vol. 140, Elsevier/Academic Press, Amsterdam, 2003. MR2424078
- [2] S. Agmon, A. Douglis, and L. Nirenberg, *Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I*, Comm. Pure Appl. Math. **12** (1959), 623–727, DOI 10.1002/cpa.3160120405. MR125307
- [3] B. Ahmad, A. Alsaedi, F. Brezzi, L. D. Marini, and A. Russo, *Equivalent projectors for virtual element methods*, Comput. Math. Appl. **66** (2013), no. 3, 376–391, DOI 10.1016/j.camwa.2013.05.015. MR3073346
- [4] P. Alfeld and M. Sirvent, *The structure of multivariate superspline spaces of high degree*, Math. Comp. **57** (1991), no. 195, 299–308, DOI 10.2307/2938675. MR1079007
- [5] P. F. Antonietti, G. Manzini, and M. Verani, *The fully nonconforming virtual element method for biharmonic problems*, Math. Models Methods Appl. Sci. **28** (2018), no. 2, 387–407, DOI 10.1142/S0218202518500100. MR3741104
- [6] P. F. Antonietti, G. Manzini, and M. Verani, *The conforming virtual element method for polyharmonic problems*, Comput. Math. Appl., 2019. <https://doi.org/10.1016/j.camwa.2019.09.022>.
- [7] J. Argyris, I. Fried, and D. Scharpf, *The TUBA family of plate elements for the matrix displacement method*, The Aeronautical Journal of the Royal Aeronautical Society **72** (1968), 701–709.
- [8] B. Ayuso de Dios, K. Lipnikov, and G. Manzini, *The nonconforming virtual element method*, ESAIM Math. Model. Numer. Anal. **50** (2016), no. 3, 879–904, DOI 10.1051/m2an/2015090. MR3507277
- [9] I. Babuška, *Error-bounds for finite element method*, Numer. Math. **16** (1971), 322–333.
- [10] L. Beirão da Veiga, A. Chernov, L. Mascotto, and A. Russo, *Exponential convergence of the hp virtual element method in presence of corner singularities*, Numer. Math. **138** (2018), no. 3, 581–613, DOI 10.1007/s00211-017-0921-7. MR3767695
- [11] L. Beirão da Veiga, F. Dassi, and A. Russo, *High-order virtual element method on polyhedral meshes*, Comput. Math. Appl. **74** (2017), no. 5, 1110–1122, DOI 10.1016/j.camwa.2017.03.021. MR3689939
- [12] L. Beirão da Veiga, F. Brezzi, A. Cangiani, G. Manzini, L. D. Marini, and A. Russo, *Basic principles of virtual element methods*, Math. Models Methods Appl. Sci. **23** (2013), no. 1, 199–214, DOI 10.1142/S0218202512500492. MR2997471
- [13] L. Beirão da Veiga, F. Brezzi, L. D. Marini, and A. Russo, *The hitchhiker’s guide to the virtual element method*, Math. Models Methods Appl. Sci. **24** (2014), no. 8, 1541–1573, DOI 10.1142/S021820251440003X. MR3200242
- [14] L. Beirão da Veiga, C. Lovadina, and A. Russo, *Stability analysis for the virtual element method*, Math. Models Methods Appl. Sci. **27** (2017), no. 13, 2557–2594, DOI 10.1142/S021820251750052X. MR3714637
- [15] J. H. Bramble and M. Zlámal, *Triangular elements in the finite element method*, Math. Comp. **24** (1970), 809–820, DOI 10.2307/2004615. MR282540
- [16] S. C. Brenner and L. R. Scott, *The Mathematical Theory of Finite Element Methods*, 3rd ed., Texts in Applied Mathematics, vol. 15, Springer, New York, 2008. MR2373954
- [17] S. C. Brenner and L.-Y. Sung, *Virtual element methods on meshes with small edges or faces*, Math. Models Methods Appl. Sci. **28** (2018), no. 7, 1291–1336, DOI 10.1142/S0218202518500355. MR3815658
- [18] F. Brezzi, A. Buffa, and K. Lipnikov, *Mimetic finite differences for elliptic problems*, M2AN Math. Model. Numer. Anal. **43** (2009), no. 2, 277–295, DOI 10.1051/m2an:2008046. MR2512497
- [19] F. Brezzi and L. D. Marini, *Virtual element methods for plate bending problems*, Comput. Methods Appl. Mech. Engrg. **253** (2013), 455–462, DOI 10.1016/j.cma.2012.09.012. MR3002804
- [20] S. Cao and L. Chen, *Anisotropic error estimates of the linear virtual element method on polygonal meshes*, SIAM J. Numer. Anal. **56** (2018), no. 5, 2913–2939, DOI 10.1137/17M1154369. MR3857893

- [21] S. Cao and L. Chen, *Anisotropic error estimates of the linear nonconforming virtual element methods*, SIAM J. Numer. Anal. **57** (2019), no. 3, 1058–1081, DOI 10.1137/18M1196455. MR3948241
- [22] L. Chen and J. Huang, *Some error analysis on virtual element methods*, Calcolo **55** (2018), no. 1, Art. 5, 23, DOI 10.1007/s10092-018-0249-4. MR3760899
- [23] P. G. Ciarlet, *The Finite Element Method for Elliptic Problems*, North-Holland Publishing Co., Amsterdam-New York-Oxford, 1978. Studies in Mathematics and its Applications, Vol. 4. MR0520174
- [24] F. Dassi and L. Mascotto, *Exploring high-order three dimensional virtual elements: bases and stabilizations*, Comput. Math. Appl. **75** (2018), no. 9, 3379–3401, DOI 10.1016/j.camwa.2018.02.005. MR3785566
- [25] K. Feng and Z.-C. Shi, *Mathematical Theory of Elastic Structures*, Springer-Verlag, Berlin; Science Press Beijing, Beijing, 1996. Translated from the 1981 Chinese original; Revised by the authors. MR1392473
- [26] F. Gazzola, H.-C. Grunau, and G. Sweers, *Polyharmonic Boundary Value Problems: Positivity Preserving and Nonlinear Higher Order Elliptic Equations in Bounded Domains*, Lecture Notes in Mathematics, vol. 1991, Springer-Verlag, Berlin, 2010. MR2667016
- [27] J. Hu, Y. Huang, and S. Zhang, *The lowest order differentiable finite element on rectangular grids*, SIAM J. Numer. Anal. **49** (2011), no. 4, 1350–1368, DOI 10.1137/100806497. MR2817542
- [28] J. Hu, Z. Shi, and J. Xu, *Convergence and optimality of the adaptive Morley element method*, Numer. Math. **121** (2012), no. 4, 731–752, DOI 10.1007/s00211-012-0445-0. MR2945614
- [29] J. Hu and S. Zhang, *The minimal conforming H^k finite element spaces on R^n rectangular grids*, Math. Comp. **84** (2015), no. 292, 563–579, DOI 10.1090/S0025-5718-2014-02871-8. MR3290955
- [30] J. Hu and S. Zhang, *A canonical construction of H^m -nonconforming triangular finite elements*, Ann. Appl. Math. **33** (2017), no. 3, 266–288. MR3821070
- [31] J. Huang and X. Huang, *A hybridizable discontinuous Galerkin method for Kirchhoff plates*, J. Sci. Comput. **78** (2019), no. 1, 290–320, DOI 10.1007/s10915-018-0780-0. MR3902886
- [32] X. Liu, J. Li, and Z. Chen, *A nonconforming virtual element method for the Stokes problem on general meshes*, Comput. Methods Appl. Mech. Engrg. **320** (2017), 694–711, DOI 10.1016/j.cma.2017.03.027. MR3646370
- [33] L. Mascotto, *Ill-conditioning in the virtual element method: stabilizations and bases*, Numer. Methods Partial Differential Equations **34** (2018), no. 4, 1258–1281, DOI 10.1002/num.22257. MR3798428
- [34] J. N. Reddy, *Theory and Analysis of Elastic Plates and Shells*, CRC Press, New York, second edition, 2006.
- [35] A. Ženíšek, *Interpolation polynomials on the triangle*, Numer. Math. **15** (1970), 283–296, DOI 10.1007/BF02165119. MR275014
- [36] M. Wang and J. Xu, *The Morley element for fourth order elliptic equations in any dimensions*, Numer. Math. **103** (2006), no. 1, 155–169, DOI 10.1007/s00211-005-0662-x. MR2207619
- [37] M. Wang and J. Xu, *Minimal finite element spaces for $2m$ -th-order partial differential equations in R^n* , Math. Comp. **82** (2013), no. 281, 25–43, DOI 10.1090/S0025-5718-2012-02611-1. MR2983014
- [38] Y. Wang, *A nonconforming Crouzeix-Raviart type finite element on polygonal meshes*, Math. Comp. **88** (2019), no. 315, 237–271, DOI 10.1090/mcom/3334. MR3854057
- [39] S. Wu and J. Xu, *$\mathcal{P}_m$ interior penalty nonconforming finite element methods for $2m$ -th order PDEs in R^n* , arXiv:1710.07678, 2017.
- [40] S. Wu and J. Xu, *Nonconforming finite element spaces for $2m$ th order partial differential equations on R^n simplicial grids when $m = n + 1$* , Math. Comp. **88** (2019), no. 316, 531–551, DOI 10.1090/mcom/3361. MR3882275
- [41] S. Zhang, *A family of 3D continuously differentiable finite elements on tetrahedral grids*, Appl. Numer. Math. **59** (2009), no. 1, 219–233, DOI 10.1016/j.apnum.2008.02.002. MR2474112
- [42] S. Zhang, *On the full C_1 - Q_k finite element spaces on rectangles and cuboids*, Adv. Appl. Math. Mech. **2** (2010), no. 6, 701–721. MR2719052

- [43] J. Zhao, S. Chen, and B. Zhang, *The nonconforming virtual element method for plate bending problems*, Math. Models Methods Appl. Sci. **26** (2016), no. 9, 1671–1687, DOI 10.1142/S021820251650041X. MR3529253
- [44] J. Zhao, B. Zhang, S. Chen, and S. Mao, *The Morley-type virtual element for plate bending problems*, J. Sci. Comput. **76** (2018), no. 1, 610–629, DOI 10.1007/s10915-017-0632-3. MR3812981

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT IRVINE, IRVINE, CALIFORNIA 92697

Email address: `chenlong@math.uci.edu`

SCHOOL OF MATHEMATICS, SHANGHAI UNIVERSITY OF FINANCE AND ECONOMICS, SHANGHAI 200433, PEOPLE'S REPUBLIC OF CHINA

Email address: `huang.xuehai@sufe.edu.cn`