

Measurements of the transverse-momentum-dependent cross sections of J/ψ production at mid-rapidity in proton+proton collisions at $\sqrt{s} = 510$ and 500 GeV with the STAR detector

J. Adam,¹² L. Adamczyk,² J. R. Adams,³⁵ J. K. Adkins,²⁶ G. Agakishiev,²⁴ M. M. Aggarwal,³⁷ Z. Ahammed,⁵⁷ I. Alekseev,^{3,31} D. M. Anderson,⁵¹ R. Aoyama,⁵⁴ A. Aparin,²⁴ D. Arkhipkin,⁵ E. C. Aschenauer,⁵ M. U. Ashraf,⁵³ F. Atetalla,²⁵ A. Attri,³⁷ G. S. Averichev,²⁴ V. Bairathi,³² K. Barish,⁹ A. J. Bassill,⁹ A. Behera,⁴⁹ R. Bellwied,¹⁹ A. Bhasin,²³ A. K. Bhati,³⁷ J. Bielcik,¹³ J. Bielcikova,³⁴ L. C. Bland,⁵ I. G. Bordyuzhin,³ J. D. Brandenburg,^{5,46} A. V. Brandin,³¹ J. Bryslawskij,⁹ I. Bunzarov,²⁴ J. Butterworth,⁴² H. Caines,⁶⁰ M. Calderón de la Barca Sánchez,⁷ D. Cebra,⁷ I. Chakaberia,^{25,46} P. Chaloupka,¹³ B. K. Chan,⁸ F.-H. Chang,³³ Z. Chang,⁵ N. Chankova-Bunzarova,²⁴ A. Chatterjee,⁵⁷ S. Chattopadhyay,⁵⁷ J. H. Chen,⁴⁷ X. Chen,⁴⁵ J. Cheng,⁵³ M. Cherney,¹² W. Christie,⁵ H. J. Crawford,⁶ M. Csanád,¹⁵ S. Das,¹⁰ T. G. Dedovich,²⁴ I. M. Deppner,¹⁸ A. A. Derevschikov,³⁹ L. Didenko,⁵ C. Dilks,³⁸ X. Dong,²⁷ J. L. Drachenberg,¹ J. C. Dunlop,⁵ T. Edmonds,⁴⁰ N. Elsey,⁵⁹ J. Engelage,⁶ G. Eppley,⁴² R. Esha,⁸ S. Esumi,⁵⁴ O. Evdokimov,¹¹ J. Ewigleben,²⁸ O. Eyser,⁵ R. Fatemi,²⁶ S. Fazio,⁵ P. Federic,³⁴ J. Fedorisin,²⁴ Y. Feng,⁴⁰ P. Filip,²⁴ E. Finch,⁴⁸ Y. Fisysak,⁵ L. Fulek,² C. A. Gagliardi,⁵¹ T. Galatyuk,¹⁴ F. Geurts,⁴² A. Gibson,⁵⁶ D. Grosnick,⁵⁶ A. Gupta,²³ W. Guryn,⁵ A. I. Hamad,²⁵ A. Hamed,⁵¹ J. W. Harris,⁶⁰ L. He,⁴⁰ S. Heppelmann,⁷ S. Heppelmann,³⁸ N. Herrmann,¹⁸ L. Holub,¹³ Y. Hong,²⁷ S. Horvat,⁶⁰ B. Huang,¹¹ H. Z. Huang,⁸ S. L. Huang,⁴⁹ T. Huang,³³ X. Huang,⁵³ T. J. Humanic,³⁵ P. Huo,⁴⁹ G. Igo,⁸ W. W. Jacobs,²¹ A. Jentsch,⁵² J. Jia,^{5,49} K. Jiang,⁴⁵ S. Jowzaee,⁵⁹ X. Ju,⁴⁵ E. G. Judd,⁶ S. Kabana,²⁵ S. Kagamaster,²⁸ D. Kalinkin,²¹ K. Kang,⁵³ D. Kapukchyan,⁹ K. Kauder,⁵ H. W. Ke,⁵ D. Keane,²⁵ A. Kechechyan,²⁴ M. Kelsey,²⁷ Y. V. Khyzhniak,³¹ D. P. Kikoła,⁵⁸ C. Kim,⁹ T. A. Kinghorn,⁷ I. Kisel,¹⁶ A. Kisiel,⁵⁸ M. Kocan,¹³ L. Kochenda,³¹ L. K. Kosarzewski,¹³ L. Kramarik,¹³ P. Kravtsov,³¹ K. Krueger,⁴ N. Kulathunga Mudiyansele,¹⁹ L. Kumar,³⁷ R. Kunnawalkam Elayavalli,⁵⁹ J. H. Kwasizur,²¹ R. Lacey,⁴⁹ J. M. Landgraf,⁵ J. Lauret,⁵ A. Lebedev,⁵ R. Lednicky,²⁴ J. H. Lee,⁵ C. Li,⁴⁵ W. Li,⁴⁷ W. Li,⁴² X. Li,⁴⁵ Y. Li,⁵³ Y. Liang,²⁵ R. Licenik,¹³ T. Lin,⁵¹ A. Lipiec,⁵⁸ M. A. Lisa,³⁵ F. Liu,¹⁰ H. Liu,²¹ P. Liu,⁴⁹ P. Liu,⁴⁷ T. Liu,⁶⁰ X. Liu,³⁵ Y. Liu,⁵¹ Z. Liu,⁴⁵ T. Ljubicic,⁵ W. J. Llope,⁵⁹ M. Lomnitz,²⁷ R. S. Longacre,⁵ S. Luo,¹¹ X. Luo,¹⁰ G. L. Ma,⁴⁷ L. Ma,¹⁷ R. Ma,⁵ Y. G. Ma,⁴⁷ N. Magdy,¹¹ R. Majka,⁶⁰ D. Mallick,³² S. Margetis,²⁵ C. Markert,⁵² H. S. Matis,²⁷ O. Matonoha,¹³ J. A. Mazer,⁴³ K. Meehan,⁷ J. C. Mei,⁴⁶ N. G. Minaev,³⁹ S. Mioduszewski,⁵¹ D. Mishra,³² B. Mohanty,³² M. M. Mondal,²² I. Mooney,⁵⁹ Z. Moravcova,¹³ D. A. Morozov,³⁹ Md. Nasim,⁸ K. Nayak,¹⁰ J. M. Nelson,⁶ D. B. Nemes,⁶⁰ M. Nie,⁴⁶ G. Nigmatkulov,³¹ T. Niida,⁵⁹ L. V. Nogach,³⁹ T. Nonaka,¹⁰ G. Odyniec,²⁷ A. Ogawa,⁵ K. Oh,⁴¹ S. Oh,⁶⁰ V. A. Okorokov,³¹ B. S. Page,⁵ R. Pak,⁵ Y. Panebratsev,²⁴ B. Pawlik,³⁶ D. Pawlowska,⁵⁸ H. Pei,¹⁰ C. Perkins,⁶ R. L. Pinter,¹⁵ J. Pluta,⁵⁸ J. Porter,²⁷ M. Posik,⁵⁰ N. K. Pruthi,³⁷ M. Przybycien,² J. Putschke,⁵⁹ A. Quintero,⁵⁰ S. K. Radhakrishnan,²⁷ S. Ramachandran,²⁶ R. L. Ray,⁵² R. Reed,²⁸ H. G. Ritter,²⁷ J. B. Roberts,⁴² O. V. Rogachevskiy,²⁴ J. L. Romero,⁷ L. Ruan,⁵ J. Rusnak,³⁴ O. Rusnakova,¹³ N. R. Sahoo,⁵¹ P. K. Sahu,²² S. Salur,⁴³ J. Sandweiss,⁶⁰ J. Schambach,⁵² W. B. Schmidke,⁵ N. Schmitz,²⁹ B. R. Schweid,⁴⁹ F. Seck,¹⁴ J. Seger,¹² M. Sergeeva,⁸ R. Seto,⁹ P. Seyboth,²⁹ N. Shah,⁴⁷ E. Shahaliev,²⁴ P. V. Shanmuganathan,²⁸ M. Shao,⁴⁵ F. Shen,⁴⁶ W. Q. Shen,⁴⁷ S. S. Shi,¹⁰ Q. Y. Shou,⁴⁷ E. P. Sichtermann,²⁷ S. Siejka,⁵⁸ R. Sikora,² M. Simko,³⁴ JSingh,³⁷ S. Singha,²⁵ D. Smirnov,⁵ N. Smirnov,⁶⁰ W. Solyst,²¹ P. Sorensen,⁵ H. M. Spinka,⁴ B. Srivastava,⁴⁰ T. D. S. Stanislaus,⁵⁶ M. Stefaniak,⁵⁸ D. J. Stewart,⁶⁰ M. Strikhanov,³¹ B. Stringfellow,⁴⁰ A. A. P. Suaide,⁴⁴ T. Sugiura,⁵⁴ M. Sumner,³⁴ B. Summa,³⁸ X. M. Sun,¹⁰ Y. Sun,⁴⁵ Y. Sun,²⁰ B. Surrow,⁵⁰ D. N. Svirida,³ P. Szymanski,⁵⁸ A. H. Tang,⁵ Z. Tang,⁴⁵ A. Taranenko,³¹ T. Tarnowsky,³⁰ J. H. Thomas,²⁷ A. R. Timmins,¹⁹ D. Tlustý,¹² T. Todoroki,⁵ M. Tokarev,²⁴ C. A. Tomkiel,²⁸ S. Trentalange,⁸ R. E. Tribble,⁵¹ P. Tribedy,⁵ S. K. Tripathy,²² O. D. Tsai,⁸ B. Tu,¹⁰ T. Ullrich,⁵ D. G. Underwood,⁴ I. Upsal,^{46,5} G. Van Buren,⁵ J. Vanek,³⁴ A. N. Vasiliev,³⁹ I. Vassiliev,¹⁶ F. Videbæk,⁵ S. Vokal,²⁴ S. A. Voloshin,⁵⁹ F. Wang,⁴⁰ G. Wang,⁸ P. Wang,⁴⁵ Y. Wang,¹⁰ Y. Wang,⁵³ J. C. Webb,⁵ L. Wen,⁸ G. D. Westfall,³⁰ H. Wieman,²⁷ S. W. Wissink,²¹ R. Witt,⁵⁵ Y. Wu,²⁵ Z. G. Xiao,⁵³ G. Xie,¹¹ W. Xie,⁴⁰ H. Xu,²⁰ N. Xu,²⁷ Q. H. Xu,⁴⁶ Y. F. Xu,⁴⁷ Z. Xu,⁵ C. Yang,⁴⁶ Q. Yang,⁴⁶ S. Yang,⁵ Y. Yang,³³ Z. Ye,⁴² Z. Ye,¹¹ L. Yi,⁴⁶ K. Yip,⁵ I. -K. Yoo,⁴¹ H. Zbroszczyk,⁵⁸ W. Zha,⁴⁵ D. Zhang,¹⁰ L. Zhang,¹⁰ S. Zhang,⁴⁵ S. Zhang,⁴⁷ X. P. Zhang,⁵³ Y. Zhang,⁴⁵ Z. Zhang,⁴⁷ J. Zhao,⁴⁰ C. Zhong,⁴⁷ C. Zhou,⁴⁷ X. Zhu,⁵³ Z. Zhu,⁴⁶ M. Zurek,²⁷ and M. Zyzak¹⁶

(STAR Collaboration)

¹Abilene Christian University, Abilene, Texas 79699

²AGH University of Science and Technology, FPACS, Cracow 30-059, Poland

³Alikhanov Institute for Theoretical and Experimental Physics, Moscow 117218, Russia

⁴Argonne National Laboratory, Argonne, Illinois 60439

- ⁵Brookhaven National Laboratory, Upton, New York 11973
- ⁶University of California, Berkeley, California 94720
- ⁷University of California, Davis, California 95616
- ⁸University of California, Los Angeles, California 90095
- ⁹University of California, Riverside, California 92521
- ¹⁰Central China Normal University, Wuhan, Hubei 430079
- ¹¹University of Illinois at Chicago, Chicago, Illinois 60607
- ¹²Creighton University, Omaha, Nebraska 68178
- ¹³Czech Technical University in Prague, FNSPE, Prague 115 19, Czech Republic
- ¹⁴Technische Universität Darmstadt, Darmstadt 64289, Germany
- ¹⁵Eötvös Loránd University, Budapest, Hungary H-1117
- ¹⁶Frankfurt Institute for Advanced Studies FIAS, Frankfurt 60438, Germany
- ¹⁷Fudan University, Shanghai, 200433
- ¹⁸University of Heidelberg, Heidelberg 69120, Germany
- ¹⁹University of Houston, Houston, Texas 77204
- ²⁰Huzhou University, Huzhou, Zhejiang 313000
- ²¹Indiana University, Bloomington, Indiana 47408
- ²²Institute of Physics, Bhubaneswar 751005, India
- ²³University of Jammu, Jammu 180001, India
- ²⁴Joint Institute for Nuclear Research, Dubna 141 980, Russia
- ²⁵Kent State University, Kent, Ohio 44242
- ²⁶University of Kentucky, Lexington, Kentucky 40506-0055
- ²⁷Lawrence Berkeley National Laboratory, Berkeley, California 94720
- ²⁸Lehigh University, Bethlehem, Pennsylvania 18015
- ²⁹Max-Planck-Institut für Physik, Munich 80805, Germany
- ³⁰Michigan State University, East Lansing, Michigan 48824
- ³¹National Research Nuclear University MEPhI, Moscow 115409, Russia
- ³²National Institute of Science Education and Research, HBNI, Jatni 752050, India
- ³³National Cheng Kung University, Tainan 70101
- ³⁴Nuclear Physics Institute of the CAS, Rez 250 68, Czech Republic
- ³⁵Ohio State University, Columbus, Ohio 43210
- ³⁶Institute of Nuclear Physics PAN, Cracow 31-342, Poland
- ³⁷Panjab University, Chandigarh 160014, India
- ³⁸Pennsylvania State University, University Park, Pennsylvania 16802
- ³⁹NRC "Kurchatov Institute", Institute of High Energy Physics, Protvino 142281, Russia
- ⁴⁰Purdue University, West Lafayette, Indiana 47907
- ⁴¹Pusan National University, Pusan 46241, Korea
- ⁴²Rice University, Houston, Texas 77251
- ⁴³Rutgers University, Piscataway, New Jersey 08854
- ⁴⁴Universidade de São Paulo, São Paulo, Brazil 05314-970
- ⁴⁵University of Science and Technology of China, Hefei, Anhui 230026
- ⁴⁶Shandong University, Qingdao, Shandong 266237
- ⁴⁷Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800
- ⁴⁸Southern Connecticut State University, New Haven, Connecticut 06515
- ⁴⁹State University of New York, Stony Brook, New York 11794
- ⁵⁰Temple University, Philadelphia, Pennsylvania 19122
- ⁵¹Texas A&M University, College Station, Texas 77843
- ⁵²University of Texas, Austin, Texas 78712
- ⁵³Tsinghua University, Beijing 100084
- ⁵⁴University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan
- ⁵⁵United States Naval Academy, Annapolis, Maryland 21402
- ⁵⁶Valparaiso University, Valparaiso, Indiana 46383
- ⁵⁷Variable Energy Cyclotron Centre, Kolkata 700064, India
- ⁵⁸Warsaw University of Technology, Warsaw 00-661, Poland
- ⁵⁹Wayne State University, Detroit, Michigan 48201
- ⁶⁰Yale University, New Haven, Connecticut 06520

(Dated: August 27, 2019)

We present measurements of the differential cross sections of inclusive J/ψ meson production as a function of transverse momentum ($p_T^{J/\psi}$) using the $\mu^+\mu^-$ and e^+e^- decay channels in proton+proton collisions at center-of-mass energies of 510 and 500 GeV, respectively, recorded by the STAR detector at the Relativistic Heavy Ion Collider. The measurement from the $\mu^+\mu^-$ channel is for $0 < p_T^{J/\psi} < 9$ GeV/c and rapidity range $|y^{J/\psi}| < 0.4$, and that from the e^+e^- channel is for $4 < p_T^{J/\psi} < 20$ GeV/c

and $|y^{J/\psi}| < 1.0$. The $\psi(2S)$ to J/ψ ratio is also measured for $4 < p_T^{\text{meson}} < 12$ GeV/ c through the e^+e^- decay channel. Model calculations, which incorporate different approaches toward the J/ψ production mechanism, are compared with experimental results and show reasonable agreement within uncertainties.

Keywords: STAR, J/ψ , $\psi(2S)$ cross section, $\mu^+\mu^-$, e^+e^-

I. INTRODUCTION

The J/ψ meson is a bound state of charm and anti-charm quarks ($c\bar{c}$) which was discovered several decades ago [1]. In hadronic collisions at energies reached at the Relativistic Heavy Ion Collider (RHIC), J/ψ are primarily produced via inelastic scattering by two gluons into charm and anti-charm quarks, followed by hadronization of the $c\bar{c}$ pair [2, 3]. Studying the J/ψ production provides valuable knowledge for the understanding of Quantum Chromodynamics (QCD) in both perturbative and non-perturbative regimes. The production of the $c\bar{c}$ pair can be calculated using the perturbative approach, however, the evolution of a $c\bar{c}$ pair into a J/ψ meson is non-perturbative, and the theoretical description remains a challenge. Different theoretical approaches have been proposed to describe the J/ψ production mechanism [4–8]. However, these descriptions have difficulties in explaining the experimental results of production cross section and polarization simultaneously. Therefore, precise measurements of the J/ψ cross section in elementary collisions at different collision energies are essential for investigating the J/ψ production mechanism. Moreover, as an important probe of the hot and dense medium, known as quark-gluon plasma (QGP), it is necessary to have a good understanding of the J/ψ production mechanism in elementary collisions in order to help understand the modification to its production in heavy-ion collisions, which has been proposed and widely pursued to study the properties of QGP [9].

There are three notable models for J/ψ production which differ mainly in the description of the non-perturbative process. These are the Color Singlet Model (CSM) [4], Non-Relativistic QCD formalism (NRQCD) [5] and the Color Evaporation Model (CEM) [6]. In the CSM, it is assumed that the hadronization process does not change the quantum numbers of the $c\bar{c}$ pair. The initially produced $c\bar{c}$ can then bind to a given charmonium state only if it is created in a color-singlet state with matching angular-momentum quantum numbers. The Next-to-Next-to-Leading-Order CSM (NNLO* CSM) has been tested for the S -wave quarkonium states in the Tevatron and LHC data. However, this model is not able to calculate the full NNLO contribution, or provide the predictions for the P -wave states, due to limitations of accuracy in the NLO calculation [10]. Therefore, it is expected to underestimate the production for the quarkonium states which have significant contributions from the decays of excited states, known as feed-down contributions [11, 12]. In the NRQCD approach, the charmonium can be produced from both the CS state

and a color-octet (CO) state. The color neutralization of the CO state is achieved by radiating soft gluons during the hadronization process. In the CEM, the produced $c\bar{c}$ pair is assumed to evolve into a J/ψ with a certain probability if its invariant mass is below the threshold for producing a $D\bar{D}$ pair. In this model, spin is always summed over which prevents it from predicting the J/ψ polarization. A recent improvement to the CEM (Improved CEM (ICEM) [7]) overcomes this issue by sorting out different spin states and is able to predict the polarization of the quarkonium states. In the low transverse momentum (p_T) range of the charmonium, the $c\bar{c}$ cross section becomes difficult to calculate at collider center-of-mass energies since the dynamics are sensitive to the large logarithms of small Bjorken x . A newly developed Color Glass Condensate (CGC) effective theory of small- x QCD provides a viable path towards calculating the J/ψ cross section at low p_T ($p_T < \sim M$, where M is the quarkonium mass) by combining the CGC effective theory with the NRQCD formalism [8].

This paper presents the measurements of the J/ψ production cross sections covering a wide p_T range from 0 to 20 GeV/ c in proton+proton collisions at center-of-mass energies of 510 and 500 GeV at RHIC. These cross sections are measured in two decay channels, which include the $\mu^+\mu^-$ channel, for $0 < p_T^{J/\psi} < 9$ GeV/ c and J/ψ rapidity ($|y^{J/\psi}| < 0.4$), and the e^+e^- channel, for $4 < p_T^{J/\psi} < 20$ GeV/ c and $|y^{J/\psi}| < 1.0$, respectively. The measured cross sections contain the direct production of J/ψ , contributions from excited charmonium states, and from decays of bottom-flavored hadrons. The first two are often categorized as prompt J/ψ as they are produced at the collision vertex and cannot be experimentally separated. The last one is often called non-prompt J/ψ , while the detector setup used in this analysis cannot experimentally distinguish it from prompt J/ψ . The feed-down contribution to the J/ψ production is an additional complication in understanding the J/ψ production mechanism as nearly 30 – 40% of the inclusive J/ψ yields come from the decay of excited charmonium states [13, 14]. Many experiments have already presented the results of heavy quarkonium production in electron+positron, hadron+hadron, and heavy-ion collisions [15, 16]. The latest measurements from the LHC [17–19] probe the high p_T production cross sections in proton+proton collisions with center-of-mass energies of 7, 8, and 13 TeV. The large kinematic range of the J/ψ measurement at the highest beam energies at RHIC (510 and 500 GeV) provides valuable insights to the J/ψ production mechanism. Additionally, the $\psi(2S)$ to J/ψ ratio is measured in the e^+e^- decay channel in the p_T range of $4 < p_T^{\text{meson}} <$

12 GeV/c. This measurement could help constrain the feed-down contribution to the J/ψ from the excited charmonium states.

The paper is organized as follows: the STAR detector will be discussed in section II, and the analyses of the $\mu^+\mu^-$ and e^+e^- decay channels will be described in detail in section III and IV, respectively. The results from these two different channels will be presented and compared to different theoretical models in section V. Finally, conclusions will be given in section VI.

II. THE STAR DETECTOR

The STAR detector is optimized for high energy nuclear physics. It has excellent particle identification capability and a large acceptance at mid-rapidity. The heart of the STAR detector is the Time Projection Chamber (TPC). The TPC is the primary tracking detector for charged particles and provides particle identification via measurements of their ionization energy losses (dE/dx) [20]. It covers the full azimuthal range ($0 \leq \phi < 2\pi$) and a large pseudorapidity range ($|\eta| < 1$). The p_T of charged particles are measured from the curvature of their trajectories in the 0.5 Tesla solenoidal field [21]. There are 30 iron bars, known as “backlegs” outside the coil to provide the return flux path for the magnetic field. These are 61 cm thick at a radius of 363 cm corresponding to about 5 interaction lengths. These backlegs play an essential role in enhancing the muon purity by absorbing the background hadrons from collisions. The hadron rejection rate is about 99% as shown in the simulation study [22]. The Muon Telescope Detector (MTD) is a fast detector which uses Multi-gap Resistive Plate Chamber technology to record signals, also referred to as “hits”, generated by charged particles traversing it. It provides single-muon and dimuon triggers depending on the number of hits within a predefined online timing window. The MTD modules are installed at a radius of about 403 cm, and the full MTD detector covers about 45% in azimuth within $|\eta| < 0.5$ [22]. The timing resolution of the MTD is ~ 100 ps and the spatial resolutions are ~ 1 -2 cm in both $r\phi$ and z directions as demonstrated in the cosmic-ray data [23]. The data used in this analysis were taken during the run in which the MTD detector was 63% completed. The Barrel Electromagnetic Calorimeter (BEMC) is a lead-scintillator sampling calorimeter with 23 radiation lengths [24]. The BEMC, being a thick absorber, is dedicated to measuring energies of particles with electromagnetic interactions, such as electrons and positrons. The BEMC is physically segmented into a total of 4800 towers with a granularity of 0.05×0.05 in $\Delta\phi \times \Delta\eta$. The energy deposited in the towers is used as a trigger to record rare events. The Vertex Position Detectors (VPD) [25] and the Beam Beam Counters (BBC) [26] are scintillator-based detectors located on both sides of the main detector, and they cover pseudorapidity ranges from 4.4 to 4.9 and 2 to 5, respectively.

III. MEASUREMENT OF $J/\psi \rightarrow \mu^+\mu^-$ SIGNAL

A. Data and Monte Carlo

Data for the $\mu^+\mu^-$ channel in this analysis were collected by the STAR detector during the 2013 RHIC proton+proton run at a collision energy of 510 GeV. The corresponding integrated luminosity sampled by the MTD dimuon trigger, which requires at least two coincidence hits on the MTD, as well as signals in the VPD and the two BBCs, within the bunch crossing is 22.0 pb^{-1} . Events used in the analysis are required to have a valid reconstructed vertex with at least two tracks that are associated with corresponding MTD hits.

A Monte Carlo (MC) simulation sample was generated by a single-particle generator with flat distributions in p_T , ϕ and y for the J/ψ signal. These simulated signals were passed through a full GEANT3 [27] STAR detector simulation and then “embedded” into real data events. These embedded events were reconstructed using the same reconstruction procedure used for real data. The kinematic distributions of the embedded J/ψ were weighted by the p_T spectrum of J/ψ in proton+proton collisions at a collision energy of 510 GeV, determined via interpolation through a global fit of world-wide measurements of J/ψ cross sections [28]. Due to the systematic uncertainties on various distortion corrections for the TPC, the p_T resolution of the reconstructed muon in MC was retuned to match the reconstructed J/ψ signal mass shape in data.

B. Muon candidate selection

Muon candidates for reconstructing the J/ψ signal must satisfy the following selection criteria: p_T^μ is greater than 1.3 GeV/c to ensure the track can reach the MTD detector; the pseudorapidity of the track is within the MTD acceptance, $|\eta^\mu| < 0.5$; the distance of closest approach (DCA) to the collision vertex must be less than 3 cm to suppress background tracks from pile-up events and secondary decay vertices; the number of TPC clusters used in track reconstruction is more than 15 (the maximum possible is 45) to ensure good momentum resolution; the number of TPC clusters used for the dE/dx measurement should be more than 10 to have good dE/dx resolution; the ratio of the number of TPC clusters used over the number of possible clusters is at least 0.52 to avoid double counting for the same tracks from track splitting. Tracks are propagated from the interaction vertex to the MTD and required to match the MTD hits geometrically which fired the trigger. In addition, the muon candidates were selected by an advanced muon identification method called the Likelihood Ratio method which is described in Ref. [29]. The rapidity of the $\mu^+\mu^-$ pairs should be smaller than 0.4 to reduce the edge effect from the J/ψ kinematic acceptance which will be described in the next section. Figure 1 shows

the invariant mass spectrum of the $\mu^+\mu^-$ pairs with the selection criteria described above applied to both candidate daughters. This can be well described by a single Gaussian as signal plus second-order polynomial function as background. A total of 1154 ± 54 final J/ψ candidates are observed within the kinematic phase space of $0 < p_T^{J/\psi} < 9$ GeV/c and $|y^{J/\psi}| < 0.4$.




FIG. 1. The $\mu^+\mu^-$ invariant mass spectrum in proton+proton collisions at $\sqrt{s} = 510$ GeV. The red solid line depicts a fit using a Gaussian function (blue line) for J/ψ signal and a second-order polynomial function (red dashed line) for background.

C. Acceptance and efficiency

The differential production cross section multiplied by the $J/\psi \rightarrow \mu^+\mu^-$ branching ratio (BR), $(5.961 \pm 0.033)\%$ [30], is given by

$$\text{BR} \times \frac{d^2\sigma}{2\pi p_T dp_T dy} = \frac{N_{J/\psi \rightarrow \mu^+\mu^-}^{\text{corr.}}}{(2\pi p_T) \cdot \int \mathcal{L} dt \cdot \Delta p_T \cdot \Delta y}, \quad (1)$$

where $N_{J/\psi \rightarrow \mu^+\mu^-}^{\text{corr.}}$ is the efficiency-corrected number of J/ψ candidates. $\int \mathcal{L} dt$ is the corresponding integrated luminosity. Δp_T and Δy are the corresponding bin widths in p_T and y of the $\mu^+\mu^-$ pairs, respectively. For each $\mu^+\mu^-$ candidate, a weighting factor (w_i) is multiplied to correct for the detector acceptance (\mathcal{A}) and the total reconstruction efficiency ($\varepsilon_{\text{reco.}}$), to obtain $N_{J/\psi \rightarrow \mu^+\mu^-}^{\text{corr.}} = \sum_{i=1}^{N_{J/\psi}} w_i$, where $w_i^{-1} = \mathcal{A} \times \varepsilon_{\text{reco.}}$ and i indicates the i th candidate. This minimizes the potential bias from the efficiency correction due to the gaps between the MTD modules and restricted pseudorapidity coverage.

The detector acceptance, \mathcal{A} , is the probability of detecting muons having certain kinematics, namely $p_T^\mu >$

1.3 GeV/c and $|\eta| < 0.5$, from the J/ψ decay within the detector fiducial volume. The \mathcal{A} can be factorized into the J/ψ decay kinematic acceptance and the MTD geometric acceptance, $\mathcal{A} = \mathcal{A}_{J/\psi} \times \mathcal{A}_{\text{MTD}}$. The acceptance can be determined with the muon angular distribution calculated in the J/ψ rest frame by the following formula [31]:

$$\frac{d^2 N}{d \cos \theta^* d \phi^*} \propto 1 + \lambda_\theta \cos^2 \theta^* + \lambda_\phi \sin^2 \theta^* \cos 2\phi^* + \lambda_{\theta\phi} \sin 2\theta^* \cos \phi^*, \quad (2)$$

where θ^* is the polar angle between the μ^+ momentum in the J/ψ rest frame and the direction of the J/ψ momentum in the laboratory frame; ϕ^* is the azimuthal angle between the J/ψ production plane (defined in the J/ψ rest frame by the momenta of the incoming protons) and the J/ψ decay plane in the lab frame; and λ_i are the parameters for different polarization configurations. Similar to the analyses carried out by other experiments [11, 17], five extreme configurations are considered to cover the polarization phase space: unpolarized, $\lambda_\theta = \lambda_\phi = \lambda_{\theta\phi} = 0$; longitudinally polarized, $\lambda_\theta = -1, \lambda_\phi = \lambda_{\theta\phi} = 0$; zero transversely polarized, $\lambda_\theta = +1, \lambda_\phi = \lambda_{\theta\phi} = 0$; positively transversely polarized, $\lambda_\theta = +1, \lambda_\phi = +1, \lambda_{\theta\phi} = 0$; and negatively transversely polarized, $\lambda_\theta = +1, \lambda_\phi = -1, \lambda_{\theta\phi} = 0$. Figure 2 shows the J/ψ decay kinematics acceptance and the MTD geometric acceptance as a function of J/ψ p_T with different polarization assumptions, respectively. There is a significant difference in the J/ψ decay kinematic acceptance for different polarization assumptions at p_T around 2 GeV/c, and the fractional difference becomes smaller at higher p_T . On the other hand, the MTD geometric acceptance is almost independent of the J/ψ polarization configuration.

The total J/ψ reconstruction efficiency, $\varepsilon_{\text{reco.}}$, includes the VPD requirement, the TPC tracking efficiency, the vertex-finding efficiency, the dimuon triggering, the MTD in-situ response and matching, and muon-identification efficiencies, as shown in the following:

$$\varepsilon_{\text{reco.}} = \varepsilon_{\text{VPD}} \times \varepsilon_{\text{vtx}} \times \varepsilon_{\text{trig.}} \times (\varepsilon_{\text{trk}}^1 \cdot \varepsilon_{\text{MTD}}^1 \cdot \varepsilon_{\text{ID}}^1) \times (\varepsilon_{\text{trk}}^2 \cdot \varepsilon_{\text{MTD}}^2 \cdot \varepsilon_{\text{ID}}^2), \quad (3)$$

where the superscripts, 1 and 2, indicate the first and second muon from a J/ψ candidate. The VPD efficiency (ε_{VPD}) is obtained from the zero-bias MC sample. The TPC tracking efficiency (ε_{trk}) is calculated from the $J/\psi \rightarrow \mu^+\mu^-$ MC sample. The vertex finding efficiency (ε_{vtx}) is obtained from data directly and is about 95% across the entire J/ψ p_T region. The MTD trigger efficiency ($\varepsilon_{\text{trig.}}$) includes the trigger electronics efficiency which varies from 95% at low p_T to more than 99% at high p_T , and the online timing window cut efficiency which reaches a plateau of 99.9%. The MTD efficiency (ε_{MTD}) is determined from the cosmic ray data for the in-situ response efficiency and from the MC sample for the matching efficiency. It is evaluated as a function of




FIG. 2. (a) The J/ψ decay kinematic acceptance and (b) the MTD geometric acceptance as a function of J/ψ p_T with different polarization assumptions in the $J/\psi \rightarrow \mu^+\mu^-$ analysis. The black solid line is the unpolarized, the red dashed line is longitudinally polarized, the pink dotted line is zero transversely polarized, the blue dashed-dotted line is positively transversely polarized, and the green dashed-dotted-dotted line is negatively transversely polarized.

muon p_T for each MTD backleg and module separately. Finally, the muon identification efficiency is calculated from the MC events and the plateau efficiency is above 95% [29]. Figure 3 shows the individual and total efficiencies used for the J/ψ cross section measurement as a function of $p_T^{J/\psi}$.

The $N_{J/\psi \rightarrow \mu^+\mu^-}^{\text{corr.}}$ in different p_T regions are extracted using the χ^2 fit with several combinations of signal and background models of the efficiency-corrected $\mu^+\mu^-$ mass distributions. The signal shape is modeled by a single




FIG. 3. The total J/ψ efficiency as a function of J/ψ p_T in the $J/\psi \rightarrow \mu^+\mu^-$ analysis. Individual contributions are also shown. The green solid circles are the TPC tracking efficiency; the magenta open triangles are the trigger efficiency; the blue open circles are the MTD efficiency including response and matching; the red open squares are the muon identification efficiency; and the blue solid triangles are the total reconstruction efficiency.

Gaussian, double Gaussian or Crystal-Ball function, and the background shape can be well described by the same-sign muon track pairs or a polynomial function at different orders. The averaged result from the various fits with different shapes for signal and background is used as the mean of $N_{J/\psi}^{\text{corr.}}$, and the maximum deviation from the mean is assigned as the signal extraction systematic uncertainty. Figure 4 shows an example of the fit results using a single Gaussian as the signal function and same-sign muon track pairs as the background template for different p_T bins.

D. Uncertainties

The statistical uncertainty is about 10.6–18.8% for different J/ψ p_T regions. There are several systematic uncertainties considered in this analysis. The maximum deviation in the results for variations in cuts/methods is taken as the systematic uncertainty for each source listed below.

- The luminosity estimate and the in-bunch pile-up effect contribute global uncertainties of 8.1% [32] and 7.7%, respectively. The latter is estimated by comparing the results with different selections of the difference of the z position measured by the VPD and TPC detectors.




FIG. 4. Fits to the efficiency-corrected $M_{\mu\mu}$ spectra of the J/ψ candidates in different p_T regions. The solid red line is a combined fit to the signal and background with a single Gaussian plus a background template from the pairs of same-sign charged tracks in the TPC.

- The uncertainty in the number of J/ψ candidates extraction is evaluated using different signal and background models as described in the previous section. It contributes about 0.3–4.8% uncertainty depending on J/ψ p_T .
 - The uncertainty in the TPC tracking efficiency is estimated by comparing the results using different TPC track quality selection criteria. Since it is difficult to obtain a p_T dependent uncertainty due to the low statistics, the p_T -integrated uncertainty of 10.6% is applied to the entire p_T range.
 - The uncertainty in the MTD trigger includes two components: (i) the trigger electronics which is dominated by the statistical uncertainty in calculating the MTD trigger efficiency; (ii) the online timing window cut which is evaluated by the difference between the 2013 and 2015 data-taking. A total 3.6% uncertainty is assigned.
 - The MTD efficiency includes three sources: (i) statistical precision of the cosmic ray data; (ii) different fit templates used for determining the response efficiency which is the main contributor; (iii) difference in the matching efficiency between cosmic ray data and simulation. The resulting uncertainty is between 1.9–7.6%.
 - The uncertainty in the muon identification efficiency is determined by comparing the efficiencies from the data-driven method and the MC sample. It contributes a 5.2–8.7% uncertainty depending on J/ψ p_T [29].
 - The uncertainty in vertex finding efficiency is estimated by comparing the efficiencies from data-driven and zero-bias MC sample. A 4.1% uncertainty is assigned.
- The systematic uncertainties from the MTD geometric acceptance is negligible. Systematic uncertainties from different sources are added in quadrature. Figure 5 shows all the uncertainties as a function of J/ψ p_T .

E. Cross-section for $J/\psi \rightarrow \mu^+ \mu^-$

We measure the invariant differential cross section multiplied by the $\mu^+ \mu^-$ branching ratio of the J/ψ meson as a function of J/ψ p_T in a fiducial volume defined by $p_T^\mu > 1.3$ GeV/c and $|\eta^\mu| < 0.5$ (fiducial cross section) and in a full muon decay phase space with $|y^{J/\psi}| < 0.4$ (full cross section). Figure 6 shows the fiducial and full cross sections of the J/ψ production. The fiducial cross section is calculated using the fiducial weight, $w_i^{\text{fid}} = \mathcal{A}_{\text{MTD}} \times \varepsilon_{\text{reco.}}$, in which the kinematic acceptance is not included. This eliminates the large unknown effect




FIG. 5. The statistical and individual systematic uncertainties as a function of J/ψ p_T . The black dashed-dotted line is the statistical uncertainty; the orange dashed-dotted-dotted line is the signal extraction uncertainty; the red dashed line is the TPC tracking efficiency uncertainty; the blue shaded line is the MTD response and matching efficiency uncertainty; the blue dotted line is MTD trigger efficiency uncertainty; the green shaded line is the muon identification uncertainty; the magenta dashed line is the vertex finding efficiency uncertainty; the black dashed line is the total systematic uncertainty; A common luminosity uncertainty of 11.2% is not included.

from the polarization assumption. The full cross section uses the full weight, $w_i^{\text{full}} = \mathcal{A}_{J/\psi} \times \mathcal{A}_{\text{MTD}} \times \varepsilon_{\text{reco.}}$, to correct for the efficiency effect on each J/ψ candidate. The central values of the full cross section in this analysis are derived under the unpolarized assumption. The gray shaded band indicates the maximum span of the cross-sections with different polarization assumptions, denoted as the “polarization envelope”.

Table I summarizes the results on fiducial and full cross sections of the J/ψ production in different p_T bins. The integrated fiducial and full cross sections up to 9 GeV/c of J/ψ p_T within $|y^{J/\psi}| < 0.4$ are 10.3 ± 0.9 (stat.) ± 1.6 (sys.) ± 1.1 (lumi.) nb and 67 ± 6 (stat.) ± 10 (sys.) $^{+200}_{-18}$ (pol.) ± 7 (lumi.) nb, respectively.

IV. MEASUREMENT OF $J/\psi \rightarrow e^+e^-$ SIGNAL

A. Data set and analysis

The proton+proton collision data at $\sqrt{s} = 500$ GeV, used in the e^+e^- analysis, were recorded by the STAR detector in 2011. The integrated luminosity of the data set is 22.1 pb^{-1} sampled by the BEMC trigger which requires a BEMC tower with a transverse energy deposit




FIG. 6. The fiducial (open black stars) and full (solid blue stars) cross sections multiplied by the branching ratio as a function of J/ψ p_T . The bars and boxes are the statistical and systematic uncertainties, respectively. The gray shaded band is the polarization envelope. A common luminosity uncertainty of 11.2% is not included.

larger than 4.3 GeV [15]. The e^\pm candidates are reconstructed and identified using information from the TPC and BEMC detectors. The track quality requirements are that each track has at least 25 out of 45 possible hits in the TPC, the number of hits for the dE/dx measurement must be larger than 15 to ensure a good dE/dx resolution, and tracks are reconstructed within the TPC acceptance of $|\eta| < 1$. The electron and positron candidates are then identified by their ionization energy loss ($\langle dE/dx \rangle$) in the TPC. The normalized $\langle dE/dx \rangle$ is defined as follows:

$$n\sigma_e = \frac{1}{\sigma_{dE/dx}} \ln\left(\frac{\langle dE/dx \rangle_e^{\text{me.}}}{\langle dE/dx \rangle_e^{\text{exp.}}}\right), \quad (4)$$

where $\langle dE/dx \rangle_e^{\text{me.}}$ and $\langle dE/dx \rangle_e^{\text{exp.}}$ are the measured $\langle dE/dx \rangle$ and the expected $\langle dE/dx \rangle$ value for electron, and the $\sigma_{dE/dx}$ is the experimental $\ln(dE/dx)$ resolution. The $n\sigma_e$ requirement for the triggered e^\pm candidates is set to be $|n\sigma_e| < 2$. The triggered e^\pm candidate is also required to have $p_T > 3.5$ GeV/c, its track must have DCA from the primary vertex less than 1 cm to reduce contamination from pile-up tracks, and it is required to match to a BEMC trigger tower in which the ADC value is larger than 290, corresponding to a deposited energy of 4.3 GeV. A cut on the ratio of the momentum measured by the TPC to the energy deposited in the BEMC towers, $0.3 < pc/E < 1.5$, is used to further suppress contribution of hadrons in triggered electron selection. However, the non-triggered e^\pm candidate is only required to be a TPC track with $p_T > 1$ GeV/c and DCA < 3 cm. The looser DCA requirement is applied to increase the statis-

$p_T^{J/\psi}$ range (GeV/c)	$p_T^{J/\psi}$ position (GeV/c)	$\text{BR} \times \frac{d\sigma_{\text{fid}}^2}{2\pi p_T dp_T dy} \pm \delta_{\text{stat.}} \pm \delta_{\text{sys.}}$ (pb/(GeV/c) ²)	$\text{BR} \times \frac{d\sigma_{\text{full}}^2}{2\pi p_T dp_T dy} \pm \delta_{\text{stat.}} \pm \delta_{\text{sys.}}^{+\delta_{\text{pol. upper}} - \delta_{\text{pol. lower}}}$ (pb/(GeV/c) ²)
0.0 - 1.5	0.67	$(9.3 \pm 1.6 \pm 1.5) \times 10^2$	$(5.4 \pm 0.9 \pm 0.9 {}^{+11.5}_{-1.3}) \times 10^3$
1.5 - 3.0	2.07	$(2.2 \pm 0.2 \pm 0.3) \times 10^2$	$(1.8 \pm 0.2 \pm 0.3 {}^{+7.3}_{-0.5}) \times 10^3$
3.0 - 5.0	3.65	$(4.4 \pm 0.4 \pm 0.6) \times 10^1$	$(2.6 \pm 0.3 \pm 0.4 {}^{+7.2}_{-0.7}) \times 10^2$
5.0 - 7.0	5.68	$(6.7 \pm 0.9 \pm 0.9) \times 10^0$	$(2.6 \pm 0.3 \pm 0.3 {}^{+4.3}_{-0.6}) \times 10^1$
7.0 - 9.0	7.73	$(1.7 \pm 0.3 \pm 0.2) \times 10^0$	$(4.6 \pm 0.9 \pm 0.6 {}^{+3.7}_{-0.9}) \times 10^0$

TABLE I. A summary of the fiducial and full differential cross sections of the J/ψ production via the $\mu^+\mu^-$ decay channel in proton+proton collisions at $\sqrt{s} = 510$ GeV. The fiducial volume is defined as $p_T^\mu > 1.3$ GeV/c and $|\eta^\mu| < 0.5$. A common luminosity uncertainty of 11.2% is not included.

tics, since lower p_T tracks are more affected by multiple scattering.




FIG. 7. The invariant mass distributions of e^+e^- pairs before and after the like-sign background (the green histogram) subtraction, as shown in black histogram and red solid circles, respectively. The blue curve is a fit to the mass spectrum. The green-dashed line indicates the residual background and a Crystal-Ball function is used to describe the J/ψ and $\psi(2S)$ signals, where $\psi(2S)$ is shown in the insert. The error bars depict the statistical uncertainties.

Figure 7 shows both J/ψ and $\psi(2S)$ signals which are reconstructed from e^+e^- pairs, which include the triggered electron pairing with either another triggered electron or with a non-triggered electron. The signal shape for J/ψ candidates is obtained from the MC simulation, which includes track momentum resolution and electron bremsstrahlung radiation in the detector. On the other hand, the background includes the combinatorial background evaluated using the like-sign e^+e^+ and e^-e^- pairs

within the same event (green histogram) and the residual background, which mainly comes from the Drell-Yan process, $c\bar{c}$ and $b\bar{b}$ decays. The invariant mass distribution after the like-sign background subtraction is fitted with a J/ψ signal shape combined with an exponential function. The raw J/ψ yield is obtained by counting the bin contents after subtracting the residual background in the mass range of $2.7 < M_{ee} < 3.3$ GeV/c². There are 9581 ± 207 J/ψ signals in $4 < p_T^{J/\psi} < 20$ GeV/c. About $\sim 10\%$ of J/ψ candidates are reconstructed outside this mass window based on MC simulations, and the J/ψ raw yield as a function of p_T is corrected for this effect. A total of 350 ± 89 $\psi(2S)$ signals are obtained in the mass counting range of $3.5 < M_{ee} < 3.8$ GeV/c². The $\psi(2S)$ signals are extracted by using the same method as for J/ψ , but with a linear function to describe the residual background. Figure 8 shows invariant mass distributions of e^+e^- pairs (black histogram) and the like-sign (filled histogram) e^+e^+ and e^-e^- pairs in three representative p_T bins.

The measurement of the J/ψ differential production cross section multiplied by BR for the e^+e^- decay channel, $(5.971 \pm 0.032)\%$ [30], is defined as:

$$\text{BR} \times \frac{d^2\sigma}{2\pi p_T dp_T dy} = \frac{N_{J/\psi \rightarrow e^+e^-}^{\text{raw}}}{(2\pi p_T) \cdot \int \mathcal{L} dt \cdot \mathcal{A}\varepsilon \cdot \Delta p_T \cdot \Delta y}, \quad (5)$$

where BR is the branching ratio for the $J/\psi \rightarrow e^+e^-$ decay channel; $N_{J/\psi \rightarrow e^+e^-}^{\text{raw}}$ is the raw number of reconstructed J/ψ via the e^+e^- pairs; $\mathcal{A}\varepsilon$ is the detector's geometric acceptance times the detection efficiency of the J/ψ candidates; $\int \mathcal{L} dt$, Δp_T , and Δy have the same meanings as of that in the $\mu^+\mu^-$ decay channel.

The total J/ψ detection efficiency, $\mathcal{A}\varepsilon$, includes the detector acceptance, the mass bin counting efficiency, and individual efficiency of the electron candidates including the TPC tracking efficiency, the electron identification, the selection on the number of hits for the dE/dx measurement, the additional efficiency for the trigger, and the cut on pc/E for the triggered electrons. The de-




FIG. 8. Dielectron mass distributions after the like-sign background subtraction (red solid circles) in different p_T ranges. The solid blue line is a combined fit to the signal and residual background with a Crystal-Ball function plus an exponential function. The error bars depict the statistical error.

cay electron's momentum resolution and additional p_T smearing are also included in the calculation of the J/ψ detection efficiency. The efficiencies for the number of dE/dx hits and $n\sigma_e$ cuts are assessed using a pure electron sample from photon conversion data, while all the other acceptance and efficiencies are obtained from MC simulation with the STAR detector geometry. Figure 9 shows the individual efficiencies for the triggered electron candidates as a function of p_T^e .




FIG. 9. The individual electron efficiencies for the triggered electrons as a function of p_T^e including pc/E cut (magenta solid circles), the number of hits for the dE/dx measurement (blue up-triangles), identification (gray down-triangles), tracking (black open circles), and trigger (green solid boxes).

The detection efficiency for $\psi(2S)$ candidates is obtained in the same way as for J/ψ . The relatively larger invariant mass of $\psi(2S)$ enhances the trigger efficiency

in the low p_T range while the slightly larger opening angle between the electron and positron daughters decaying from the $\psi(2S)$ will result in a smaller acceptance and thus a lower detection efficiency at high p_T . Figure 10 shows the detection efficiency for J/ψ and $\psi(2S)$, as well as the $\psi(2S)$ to J/ψ efficiency ratio as a function of $p_T^{e^+e^-}$.




FIG. 10. (a) The J/ψ and $\psi(2S)$ detection efficiencies as a function of p_T as shown in the black solid circles and the blue open circles, respectively. (b) The detection efficiency ratio of $\psi(2S)$ to J/ψ as a function of p_T and the red dashed-line is a line at unity.

B. Uncertainties

The systematic uncertainties for the final J/ψ cross section are estimated by varying analysis selections in

both data and MC simulation and comparing the corresponding J/ψ cross section to the nominal value. The systematic uncertainties considered in this analysis are the following:

- The uncertainty in the luminosity contributes an overall 8.1% uncertainty [32]. The uncertainty in the in-bunch pile-up effect is negligible due to the low instantaneous luminosity in the 2011 data-taking and the high vertex finding efficiency for the BEMC triggered events [33].
- The J/ψ extraction uncertainty is estimated by using different fitting ranges and different residual background shapes. It contributes a 0.2–12.7% uncertainty depending on J/ψ p_T .
- The uncertainty in the TPC tracking efficiency is estimated in the same way as in the $\mu^+\mu^-$ decay channel, and it contributes a 4–14%.
- The uncertainty in the trigger efficiency is evaluated by comparing BEMC response in data and MC simulation, and the contribution is a 0.3–11.8% in various J/ψ p_T range.
- The uncertainty in the electron identification is estimated by comparing the difference between photonic electron's $n\sigma_e$ distribution at different p_T ranges, and it contributes an overall 1% uncertainty.
- The J/ψ internal conversion is estimated to be 4% within the mass counting range of 2.7 to 3.3 GeV/c^2 .

The systematic uncertainty from the vertex finding is negligible. Systematic uncertainties from different sources are added in quadrature. Figure 11 shows all the uncertainties as a function of J/ψ p_T . For the $\psi(2S)$ to J/ψ ratio, most of the systematic uncertainties cancel in the ratio, except for the signal extraction and trigger efficiency. They are evaluated the same way as for the J/ψ . They contribute 5.5% and 5.3%, respectively, to the uncertainty of the ratio measurement.

C. Cross-section for $J/\psi \rightarrow e^+e^-$

Figure 12 shows the fiducial and full production cross sections measured in the e^+e^- decay channel for $4 < p_T^{J/\psi} < 20 \text{ GeV}/c$ and $|y| < 1.0$. Table II summarizes the cross sections of the J/ψ production as a function of p_T . Similarly as for the $\mu^+\mu^-$ channel, the full cross section is calculated under the unpolarized assumption and the polarization envelope is obtained from the same five extreme cases. This provides us the information of the J/ψ production cross section within the full J/ψ decay phase space. On the other hand, the fiducial cross section only accesses the restricted phase space, but it is independent




FIG. 11. The statistical and individual systematic uncertainties as a function of J/ψ p_T . The black dashed-dotted line is the statistical uncertainty. The violet dashed-dotted-dotted line is the signal extraction uncertainty; the red dashed line is the TPC tracking uncertainty; the yellow dashed line is the electron identification ($n\sigma_e$) uncertainty; the green shaded line is the trigger uncertainty; the blue dashed line is the J/ψ internal conversion uncertainty; the black dashed line is the total systematic uncertainty; A common luminosity uncertainty of 8.1% is not included.

from any polarization assumptions. The integrated fiducial and full cross sections from 4 to 20 GeV/c of J/ψ p_T are 2.90 ± 0.08 (stat.) ± 0.22 (sys.) ± 0.24 (lumi.) nb and 10.7 ± 0.5 (stat.) ± 0.8 (sys.) $^{+5.9}_{-2.2}$ (pol.) ± 0.9 (lumi.) nb, respectively.

The cross section ratio of $\psi(2S)$ over J/ψ is 0.038 ± 0.010 (stat.) ± 0.003 (sys.) measured in the p_T range of $4 < p_T^{\text{meson}} < 12 \text{ GeV}/c$, which is shown in Fig. 13. The result is consistent with other experimental measurements [34–38] and there is no obvious dependence on the collision energy observed. The ICeM model [7] prediction is consistent with our data within uncertainties.

V. COMBINED RESULTS

Figure 14 shows the differential cross section of inclusive J/ψ production in proton+proton collisions at $\sqrt{s} = 510$ and 500 GeV measured by the STAR experiment combining the $\mu^+\mu^-$ and e^+e^- decay channels. Please note that there is a $\sim 3\%$ difference between the cross sections at 510 and 500 GeV collision energies and the difference between the different rapidity coverages is negligible [39]. The unpolarized-assumption result is compared to the NRQCD [5] and ICeM [7] calculations of J/ψ production, which includes feed-down contributions from excited charmonium states. The prediction

$p_T^{J/\psi}$ range (GeV/c)	$p_T^{J/\psi}$ position (GeV/c)	$\text{BR} \times \frac{d\sigma_{\text{fid}}^2}{2\pi p_T dp_T dy} \pm \delta_{\text{stat.}} \pm \delta_{\text{sys.}}$ (pb/(GeV/c) ²)	$\text{BR} \times \frac{d\sigma_{\text{full}}^2}{2\pi p_T dp_T dy} \pm \delta_{\text{stat.}} \pm \delta_{\text{sys.}}^{+\delta_{\text{pol. upper}} - \delta_{\text{pol. lower}}}$ (pb/(GeV/c) ²)
4.0 - 4.5	4.23	$(1.64 \pm 0.20 \pm 0.12) \times 10^1$	$(1.32 \pm 0.16 \pm 0.10^{+0.72}_{-0.35}) \times 10^2$
4.5 - 5.0	4.73	$(1.88 \pm 0.16 \pm 0.15) \times 10^1$	$(9.0 \pm 0.8 \pm 0.7^{+5.2}_{-2.0}) \times 10^1$
5.0 - 5.5	5.23	$(1.36 \pm 0.07 \pm 0.11) \times 10^1$	$(4.6 \pm 0.2 \pm 0.4^{+2.6}_{-0.8}) \times 10^1$
5.5 - 6.0	5.73	$(10.3 \pm 0.7 \pm 0.8) \times 10^0$	$(2.69 \pm 0.17 \pm 0.20^{+1.59}_{-0.39}) \times 10^1$
6.0 - 6.5	6.23	$(7.6 \pm 0.4 \pm 0.6) \times 10^0$	$(1.64 \pm 0.09 \pm 0.13^{+0.97}_{-0.24}) \times 10^1$
6.5 - 7.0	6.73	$(5.40 \pm 0.29 \pm 0.31) \times 10^0$	$(10.6 \pm 0.6 \pm 0.6^{+5.4}_{-1.6}) \times 10^0$
7.0 - 7.5	7.23	$(3.15 \pm 0.20 \pm 0.20) \times 10^0$	$(5.8 \pm 0.4 \pm 0.4^{+2.9}_{-0.9}) \times 10^0$
7.5 - 8.0	7.73	$(2.41 \pm 0.16 \pm 0.16) \times 10^0$	$(4.26 \pm 0.28 \pm 0.28^{+1.76}_{-0.61}) \times 10^0$
8.0 - 8.5	8.23	$(1.40 \pm 0.11 \pm 0.13) \times 10^0$	$(2.40 \pm 0.19 \pm 0.23^{+0.97}_{-0.33}) \times 10^0$
8.5 - 9.0	8.73	$(10.6 \pm 0.8 \pm 0.6) \times 10^{-1}$	$(1.75 \pm 0.14 \pm 0.11^{+0.65}_{-0.23}) \times 10^0$
9.0 - 9.5	9.24	$(7.5 \pm 0.7 \pm 0.4) \times 10^{-1}$	$(12.0 \pm 1.1 \pm 0.7^{+4.0}_{-1.6}) \times 10^{-1}$
9.5 - 10.0	9.73	$(5.26 \pm 0.48 \pm 0.33) \times 10^{-1}$	$(8.3 \pm 0.7 \pm 0.5^{+2.4}_{-1.1}) \times 10^{-1}$
10.0 - 10.5	10.23	$(3.26 \pm 0.43 \pm 0.21) \times 10^{-1}$	$(5.06 \pm 0.67 \pm 0.32^{+1.57}_{-0.62}) \times 10^{-1}$
10.5 - 11.0	10.74	$(2.31 \pm 0.35 \pm 0.14) \times 10^{-1}$	$(3.51 \pm 0.52 \pm 0.22^{+1.02}_{-0.42}) \times 10^{-1}$
11.0 - 12.0	11.44	$(1.58 \pm 0.20 \pm 0.13) \times 10^{-1}$	$(2.33 \pm 0.30 \pm 0.19^{+0.62}_{-0.27}) \times 10^{-1}$
12.0 - 13.0	12.45	$(7.5 \pm 1.5 \pm 0.6) \times 10^{-2}$	$(10.7 \pm 2.1 \pm 0.9^{+2.3}_{-1.2}) \times 10^{-2}$
13.0 - 15.0	13.83	$(3.5 \pm 0.7 \pm 0.6) \times 10^{-2}$	$(4.9 \pm 1.0 \pm 0.8^{+1.0}_{-0.5}) \times 10^{-2}$
15.0 - 17.0	15.85	$(2.02 \pm 0.39 \pm 0.31) \times 10^{-2}$	$(2.7 \pm 0.5 \pm 0.4^{+0.4}_{-0.3}) \times 10^{-2}$
17.0 - 20.0	18.20	$(0.84 \pm 0.18 \pm 0.17) \times 10^{-2}$	$(1.07 \pm 0.23 \pm 0.22^{+0.13}_{-0.09}) \times 10^{-2}$

TABLE II. A summary of the fiducial and full differential cross sections for the inclusive J/ψ production via the e^+e^- decay channel in proton+proton collisions at $\sqrt{s} = 500$ GeV. A common luminosity uncertainty of 8.1% is not included.

from the CGC effective theory coupled with NRQCD (CGC+NRQCD) [8] lies systematically above the data at low p_T , however, it is consistent with the data within the polarization envelope. The NLO NRQCD [5] result does a reasonably good job in describing the data above 6 GeV/c. The ICEM calculation [7] can cover the entire p_T range and is also consistent with the data within the polarization envelope. The feed-down contributions from B -hadrons are about 10–20% in the $p_T < 10$ GeV/c region and nearly 40% in our maximum p_T bin (20 GeV/c) as measured by other experiments [17, 18]. Therefore, to present a fair comparison, all of the predictions are adjusted to include this contribution using the FONLL calculation [40], shown as a green band in Fig. 14 (a).

The scaling behavior of particle production with $x_T = 2p_T/\sqrt{s}$ is characteristic for production through fragmentation due to hard scatterings. The x_T scaling ($E \frac{d^3\sigma}{dp^3} =$

$g(x_T)/s^{n/2}$) has been tested for pions, protons, and the J/ψ for various collision energies [41], where n is a free parameter which can be interpreted as the number of active partons involved in hadron production. Figure 15 shows the x_T dependence of protons, pions, and J/ψ . The J/ψ measured in 510 and 500 GeV proton+proton collisions has been fit to extract the parameter n , with $n = 5.6 \pm 0.1$. This value is consistent with $n = 5.6 \pm 0.2$ as found in a previous STAR measurement [42], as well as other previous measurements [34, 43–51] at high- p_T , and this value is close to the CO and CEM predictions, which are $n \sim 6$ [52, 53] and smaller than that from NNLO* CSM prediction which is $n \sim 8$ [54]. The broken scaling at low- p_T is due to the onset of soft processes [42].




FIG. 12. The fiducial (open black stars) and full (solid red stars) cross sections multiplied by the branching ratio as a function of J/ψ p_T . The boxes are the total systematic uncertainty. The bars are the statistical uncertainty and are too small to be visible in the figure. The gray shaded band is the polarization envelope. A common luminosity uncertainty of 8.1% is not included.




FIG. 13. The cross section ratio of $\psi(2S)$ over J/ψ as a function of their p_T measured by STAR (red solid star) is compared to results from CDF (open circles), PHENIX (open boxes), HERA-B (solid-down-triangles and open-up-triangles), ATLAS (gray crosses), CMS (gray solid and open triangle-crosses) experiments, and the prediction from ICeM (purple band). The bar and box indicates the statistical and total systematic uncertainty, respectively.




FIG. 14. (a) The J/ψ differential full production cross sections as a function of $p_T^{J/\psi}$ in proton+proton collisions at $\sqrt{s} = 510$ and 500 GeV measured through the $\mu^+\mu^-$ (blue stars) and e^+e^- decay channels (red circles). The shaded region around the data points denotes the polarization envelope and the green curve is the estimation of the B -hadrons feed-down from FONLL [40]. (b, c, d) Ratios of data and different model calculations to the Levy fit function. The bars and boxes indicate the statistical and total systematic uncertainty, respectively. A common luminosity uncertainties of 11.2% and 8.1% for the $\mu^+\mu^-$ and e^+e^- decay channels are not included.

VI. CONCLUSIONS

Differential cross sections for the J/ψ meson in proton+proton collisions at $\sqrt{s} = 510$ and 500 GeV at RHIC are measured using the $\mu^+\mu^-$ and e^+e^- decay channels. The results cover a wide $p_T^{J/\psi}$ range from 0 to 20 GeV/ c within $|y^{J/\psi}| < 0.4$ and 1.0 for the $\mu^+\mu^-$ and e^+e^- channel, respectively. Two different measurements of the inclusive J/ψ production cross section have been presented. The first is a fiducial cross section measurement utilizing only a restricted phase space as defined by detector acceptance. It is independent of the assumptions regarding polarization which results in a large systematic uncer-




FIG. 15. The x_T dependence of pion, proton, and J/ψ from different experiments.

tainty at low p_T . This allows direct comparisons between measurements and theoretical calculations in the future, for more discriminating tests of the models. The second is a full cross section measurement, accessing the full J/ψ decay phase space, depending highly on assumptions regarding polarization. The integrated fiducial and full production cross sections measured for inclusive J/ψ mesons within $0 < p_T^{J/\psi} < 9$ GeV/c are 10.3 ± 0.9 (stat.) ± 1.6 (sys.) ± 1.1 (lumi.) nb and 67 ± 6 (stat.) ± 10 (sys.) $^{+200}_{-18}$ (pol.) ± 7 (lumi.) nb, respectively, via the $\mu^+\mu^-$ channel. For $4 < p_T^{J/\psi} < 20$ GeV/c, they are 2.90 ± 0.08 (stat.) ± 0.22 (sys.) ± 0.24 (lumi.) nb and 10.7 ± 0.5 (stat.) ± 0.8 (sys.) $^{+5.9}_{-2.2}$ (pol.) ± 0.9 (lumi.) nb, respectively, via the e^+e^- channel. The calculations from CGC+NRQCD, NLO NRQCD and ICM [5, 8], which cover low, high, and both p_T regions respectively, give a reasonable description for the data within the polarization envelope. The x_T dependence for J/ψ production is also presented for 510 and 500 GeV proton+proton collisions. The result is consistent with measurements at

other collision energies from other collaborations. The ratio of $\psi(2S)$ to J/ψ for p_T from 4–12 GeV/c is measured to be 0.038 ± 0.010 (stat.) ± 0.003 (sys.). It is consistent with results from other experiments and there is no obvious collision energy dependence. Since the J/ψ production mechanism is not yet fully understood, it is important to continue confronting the models that incorporate the most current understanding with new data. A more discriminating comparison to theoretical models at low p_T can be performed in the future, if the calculations are carried out within the fiducial volume of the STAR detector, eliminating the uncertainty due to the J/ψ polarization. The results presented in this paper, together with cross section measurements at other energies, and measurements of the polarization, contribute to the goal of better understanding the production of heavy quarkonium in hadronic collisions.

VII. ACKNOWLEDGMENTS

We thank the RHIC Operations Group and RCF at BNL, the NERSC Center at LBNL, and the Open Science Grid consortium for providing resources and support. This work was supported in part by the Office of Nuclear Physics within the U.S. DOE Office of Science, the U.S. National Science Foundation, the Ministry of Education and Science of the Russian Federation, National Natural Science Foundation of China, Chinese Academy of Science, the Ministry of Science and Technology of China and the Chinese Ministry of Education, the National Research Foundation of Korea, Czech Science Foundation and Ministry of Education, Youth and Sports of the Czech Republic, Hungarian National Research, Development and Innovation Office, New National Excellency Programme of the Hungarian Ministry of Human Capacities, Department of Atomic Energy and Department of Science and Technology of the Government of India, the National Science Centre of Poland, the Ministry of Science, Education and Sports of the Republic of Croatia, RosAtom of Russia and German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF) and the Helmholtz Association.

-
- [1] J.J. Aubert *et al.*, Phys. Rev. Lett. **33**, 1404 (1974); J.-E. Augustin *et al.*, Phys. Rev. Lett. **33**, 1406 (1974).
 - [2] G.C. Nayak *et al.*, Phys. Lett. B **613**, 45 (2005); G.C. Nayak *et al.*, Phys. Rev. D **72**, 114012 (2005); G.C. Nayak *et al.*, Phys. Rev. D **74**, 074007 (2006).
 - [3] A. Andronic *et al.*, Eur. Phys. J. C **76**, 107 (2016).
 - [4] C.-H. Chang, Nucl. Phys. B **172**, 425 (1980); R. Baier and R. Rückl, Phys. Lett. B **102**, 364 (1981); Z. Phys. C **19**, 251 (1983); E.L. Berger, D.L. Jones, Phys. Rev. D **23**, 1521 (1981).
 - [5] Y.-Q. Ma *et al.*, Phys. Rev. Lett. **106**, 042002 (2011).
 - [6] H. Fritzsch, Phys. Lett. B **67**, 217 (1977); F. Halzen, Phys. Lett. B **69**, 105 (1977); M. Gluck, J.F. Owens, and E. Reya, Phys. Rev. D **17**, 2324 (1978); V.D. Barger, W.Y. Keung, and R.J.N. Phillips, Phys. Lett. B **91**, 253 (1980); J.F. Amundson, O.J.P. Eboli, E.M. Gregores and F. Halzen, Phys. Lett. B **372**, 127 (1996); J.F. Amundson, O.J.P. Eboli, E.M. Gregores and F. Halzen, Phys. Lett. B **390**, 323 (1997).
 - [7] Y.-Q. Ma and R. Vogt, Phys. Rev. D **94**, 114029 (2016).
 - [8] Y.-Q. Ma and R. Venugopalan, Phys. Rev. Lett. **113**, 192301 (2014).

- [9] L. Adamczyk *et al.* (STAR Collaboration), Phys. Rev. C **90**, 024906 (2014); L. Adamczyk *et al.* (STAR Collaboration), Phys. Lett. B **722**, 55 (2013); J. Adam *et al.* (ALICE Collaboration), JHEP **07**, 051 (2015); V. Khachatryan *et al.* (CMS Collaboration), Eur. Phys. J. C **77**, 252 (2017); A.M. Sirunyan *et al.* (CMS Collaboration), Eur. Phys. J. C **78**, 509 (2018).
- [10] J.P. Lansberg, Eur. Phys. J. C **61**, 693 (2009).
- [11] G. Aad *et al.* (ATLAS Collaboration), Phys. Rev. D **87**, 052004 (2013).
- [12] S. Chatrchyan *et al.* (CMS Collaboration), Phys. Lett. B **727**, 101 (2013).
- [13] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **79**, 578 (1997).
- [14] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **98**, 232001 (2007).
- [15] J. Adam *et al.* (STAR Collaboration), Phys. Lett. B **786**, 87 (2018).
- [16] M. Kramer, Prog. Part. Nucl. Phys. **47**, 141 (2001); J.P. Lansberg, Int. J. Mod. Phys. A **21**, 3857 (2006); J.P. Lansberg *et al.*, AIP Conf. Proc. **1038**, 15 (2008).
- [17] G. Aad *et al.* (ATLAS Collaboration), Nucl. Phys. B **850**, 387 (2011); G. Aad *et al.* (ATLAS Collaboration), Eur. Phys. J. C **76**, 283 (2016).
- [18] V. Khachatryan *et al.* (CMS Collaboration), Eur. Phys. J. C **71**, 1575 (2011); V. Khachatryan *et al.* (CMS Collaboration), Phys. Rev. Lett. **114**, 191802 (2015); A. M. Sirunyan *et al.* (CMS Collaboration), Phys. Lett. B **780**, 251 (2018).
- [19] R. Aaij *et al.* (LHCb Collaboration), Eur. Phys. J. C **71**, 1645 (2011).
- [20] M. Anderson *et al.*, Nucl. Instr. and Meth. A **499**, 659 (2003).
- [21] F. Bergsma *et al.*, Nucl. Instr. and Meth. A **499**, 633 (2003).
- [22] L. Ruan *et al.*, J. Phys. G **36**, 095001 (2009).
- [23] C. Yang *et al.*, Nucl. Instr. and Meth. A **762**, 1 (2014).
- [24] M. Beddo *et al.*, Nucl. Instr. and Meth. A **499**, 725 (2003).
- [25] W. Llope, *et al.*, Nucl. Instr. and Meth. A **522**, 252 (2004).
- [26] J. Kiryluk, AIP Conf. Proc. **675**, 424 (2003).
- [27] CERN Program Library Long Writeup W5013 (1994).
- [28] W. Zha *et al.*, Phys. Rev. C **93**, 024919 (2016).
- [29] T.C. Huang *et al.*, Nucl. Instr. and Meth. A **833**, 88 (2016).
- [30] M. Tanabashi *et al.* (Particle Data Group), Phys. Rev. D **98**, 030001 (2018).
- [31] P. Faccioli *et al.*, Eur. Phys. J. C **69**, 657 (2010).
- [32] H. Agakishiev *et al.* (STAR Collaboration), Phys. Rev. D **83**, 052006 (2011).
- [33] R. Reed *et al.*, J. Phys.: Conf. Ser. **219**, 032020 (2010).
- [34] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **79**, 572 (1997).
- [35] A. Adare *et al.* (PHENIX Collaboration), Phys. Rev. D **85**, 092004 (2012).
- [36] I. Abt *et al.* (HERA-B Collaboration), Eur. Phys. J. C **49**, 545 (2007).
- [37] G. Aad *et al.* (ATLAS Collaboration), Eur. Phys. J. C, **76**, 283 (2016).
- [38] S. Chatrchyan *et al.* (CMS Collaboration), JHEP **02**, 011 (2012).
- [39] T. Sjöstrand *et al.*, Comput. Phys. Commun. **191**, 159 (2015).
- [40] <http://www.lpthe.jussieu.fr/~cacciari/fonll/fonllform.html>
- [41] A.G. Clark *et al.*, Phys. Lett. B **74**, 267 (1978); A.L.S. Angelis *et al.* (CCOR Collaboration), Phys. Lett. B **79**, 505 (1978); S.S. Adler *et al.* (PHENIX Collaboration), Phys. Rev. C **69**, 034910 (2004).
- [42] B.I. Abelev *et al.* (STAR Collaboration), Phys. Rev. C **80**, 041902 (2009).
- [43] D.E. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
- [44] C. Albajar *et al.* (UA1 Collaboration), Phys. Lett. B **256**, 112 (1991).
- [45] A. Adare *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **98**, 232002 (2007).
- [46] C. Kourkoumelis *et al.*, Phys. Lett. B **91**, 481 (1980).
- [47] J. Adams *et al.* (STAR Collaboration), Phys. Lett. B **637**, 161 (2006).
- [48] J. Adams *et al.* (STAR Collaboration), Phys. Lett. B **616**, 8 (2005).
- [49] M. Banner *et al.*, Phys. Lett. B **115**, 59 (1982).
- [50] B. Alper *et al.*, Nucl. Phys. B **100**, 237 (1975).
- [51] D. Antreasyan *et al.*, Phys. Rev. D **19**, 764 (1979).
- [52] G.C. Nayak, M.X. Liu, and F. Cooper, Phys. Rev. D **68**, 034003 (2003).
- [53] M. Bedjidian *et al.* (2004), arXiv:hep-ph/0311048.
- [54] P. Artoisenet *et al.*, Phys. Rev. Lett. **101**, 152001 (2008).