


One-pot Aminoalkylation of Aldehydes: Diastereoselective Synthesis of Vicinal Diamines with Azaaryl methylamines

Haixing Guan,[†] Xianzhong Cao,[†] Patrick J. Walsh,^{*,‡} and Jianyou Mao,^{*,†}

[†]Institute of Advanced Synthesis, School of Chemistry and Molecular Engineering, Nanjing Tech University, 30 South Puzhu Road, Nanjing 211816, P. R. China

[‡]Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, Pennsylvania 19104-6323, USA

Supporting Information Placeholder



ABSTRACT: A one-pot synthesis of vicinal diamines with azaaryl methylamines and aldehydes is reported. A diverse array of vicinal diamines could be achieved in up to 92% yield with good to excellent diastereoselectivities (up to 20:1). The tandem reaction takes place under mild conditions and provides an alternative strategy for the synthesis of vicinal diamines.

Vicinal diamines constitute an important structural class with broad applications in natural products, chemotherapy, medicinal chemistry, and organic synthesis.¹ Due to their wide applications, their synthesis continues to attract significant attention, where the demand for more diastereoselective and straightforward syntheses still remains high. Furthermore, the rapid introduction of nitrogens into organic structures is a long-standing challenge in medicinal chemistry.²

Traditional routes for diamine syntheses generally involve: (1) direct inter- or intra-molecular damination of unfunctionalized alkenes;³ (2) addition of organometallic reagents to bisimine derivatives;⁴ (3) reductive homocoupling of imines;⁵ (4) ring opening of aziridines with nitrogen nucleophiles;⁶ and (5) Mannich type reactions employing nitroalkanes followed by reduction of the corresponding nitro groups.⁷ These routes have a number of drawbacks. For example, the first three routes are limited to the synthesis of symmetric vicinal diamines whereas the latter two require additional steps to prepare the starting aziridines or use an external reductant to reduce the undesired nitro groups.

The direct addition of α -amino anions to imines is a very important tool to construct a diverse array of vicinal diamines.⁸ To the best of our knowledge, however, a one-pot synthesis of vicinal diamines from aldehydes is unknown. Recently, our team developed a one-pot chemoselective C–N and C–C bond forming reaction for the aminobenzylation of aldehydes with toluene derived pronucleophiles (Scheme 1a).⁹ More recently, this method has been made catalytic in base.¹⁰


Further adaptation of the chemistry led to a novel one-pot route to 2-arylinoles from readily available 2-fluoro toluenes and benzonitriles (Scheme 1b).¹¹ We put forth evidence that these reactions are facilitated by formation of a cation- π interactions between the main group metal (Cs^+) and the toluene derivative, facilitating the deprotonation of the benzylic methyl group^{9,11–12}. Herein, we advance a straightforward and practical one-pot synthesis of vicinal diamines employing azaaryl methylamines and aldehydes. The tandem reaction provides an alternative strategy for the synthesis of biologically active compounds.¹³

Scheme 1. Tandem synthesis of bioactive fine chemicals.


a) One-pot aminobenzylation of aldehydes



b) One-pot synthesis of indoles




c) One-pot synthesis of vicinal diamines (*this work*)



Based on our previous studies, we began to investigate employing benzaldehyde (**1a**) and 4-(pyridin-2-ylmethyl)morpholine (**2a**) as model substrates (Table 1). We first screened three different bases [LiN(SiMe₃)₂, NaN(SiMe₃)₂, and KN(SiMe₃)₂] using THF as solvent at 10 °C for 12 h (Table 1, entries 1–3). According to the screening results, NaN(SiMe₃)₂ was identified as the best candidate among these three bases, giving **3aa** in 84% AY with 4.5 : 1 dr (entry 2, AY = assay yield determined by ¹H NMR integration of the unpurified reaction mixture against an internal standard). It is important to emphasize that MN(SiMe₃)₂ (M = Li, Na, K) plays two vital roles: (1) condensation with benzaldehyde and aza-Peterson elimination to in situ generate *N*-(trimethylsilyl)imines; and (2) direct deprotonation of benzylic C–H bonds of 4-(pyridin-2-ylmethyl)morpholine (**2a**) (Scheme 1c). Based on the conditions of entry 2, we next tested five ethereal solvents (ⁱPr₂O, TBME, CPME, DME and Et₂O). Based on consideration of diastereomeric ratios and yield, ⁱPr₂O outperformed other solvents (5.2 : 1 dr, 86% AY, entry 4) and was used going forward. It is noteworthy that TBME also gave good results (90% yield, 4.6 : 1 dr, entry 5). Since amine additives impact the aggregation of silyl amide bases,¹⁴ and therefore, their reactivity and selectivity, we examined two commonly used polyamine ligands: *N,N,N',N'*-pentamethyldiethylenetriamine (PMDTA, 1.0 equiv) and TMEDA (1.0 equiv) under the conditions of entry 2. PMDTA improved both the yield and diastereoselectivity, affording the desired product in 95% yield with 6.0 : 1 dr (entry 10). Ultimately, the optimized conditions for this tandem reaction are 1.0 equiv of benzaldehyde, 1.5 equiv of **2a**, 2.0 equiv of NaN(SiMe₃)₂, and 1.0 equiv of PMDTA in ⁱPr₂O at 10 °C for 12 h. After derivatization of **3aa** by 3,5-dinitrobenzoyl chloride, the corresponding *anti*- and *syn*-diastereomers of **4aa** could be separated. According to ¹H NMR of **4aa** (see Supporting Information), the coupling constant between the two protons alpha to the nitrogens of the major isomer of **4aa** is 10.5 Hz whereas the minor component of **4aa** 3.7 Hz, leading to the assignment of the major diastereomer as *anti*-**4aa**.^{8h} This spectroscopic difference between the diastereoisomers was used to judge *syn/anti* selectivity in the following studies, which show *anti* selectivity in all cases.

Table 1. Optimization of Reaction Conditions^a.




| entry | base | additive | solvent | AY(%) ^b | dr ^c |
|-------|--------------------------------------|----------|--------------------------------|--------------------|-----------------|
| 1 | LiN(SiMe ₃) ₂ | -- | THF | 90 | 1.6:1 |
| 2 | NaN(SiMe ₃) ₂ | -- | THF | 84 | 4.5:1 |
| 3 | KN(SiMe ₃) ₂ | -- | THF | Trace | -- |
| 4 | NaN(SiMe ₃) ₂ | -- | ⁱ Pr ₂ O | 86 | 5.2:1 |
| 5 | NaN(SiMe ₃) ₂ | -- | TBME | 90 | 4.6:1 |
| 6 | NaN(SiMe ₃) ₂ | -- | CPME | 76 | 5.8:1 |
| 7 | NaN(SiMe ₃) ₂ | -- | DME | 52 | 4.3:1 |

| | | | | | |
|----|--------------------------------------|-------|--------------------------------|----|-------|
| 8 | NaN(SiMe ₃) ₂ | -- | Et ₂ O | 87 | 4.4:1 |
| 9 | NaN(SiMe ₃) ₂ | TMEDA | ⁱ Pr ₂ O | 81 | 5.5:1 |
| 10 | NaN(SiMe ₃) ₂ | PMDTA | ⁱ Pr ₂ O | 95 | 6.0:1 |

^aReactions conducted under argon on 0.1 mmol scale. ^bYields were determined by ¹H NMR analysis of unpurified reaction mixtures with internal standard dimethyl terephthalate. ^cThe dr was determined by LC-MS analysis of unpurified reaction mixtures.

With the optimized conditions in hand, the scope of aryl aldehydes was examined (Scheme 2). In general, a diverse array of substituents on the aldehydes was tolerated under the mild conditions. The parent benzaldehyde (**1a**) was successfully transformed to **3aa** in 95% yield with 6.0 : 1 dr. Benzaldehydes substituted by electron donating groups, such as 4-methyl (**1b**), 4-methoxy (**1c**) and 4-phenoxy (**1d**) exhibited good reactivity, giving **3ba**, **3ca** and **3da** in 73%, 70% and 79% yields with similar diastereoselectivities of 4.2 : 1, 6.7 : 1 and 4.5 : 1, respectively. It is interesting that 4-methylbenzaldehyde is a good substrate, given that deprotonation of the relatively acidic methyl group of the aldehyde or imine could be envisioned to be problematic. Substrates containing extended π -systems, such as 2-naphthyl aldehyde (**1e**), furnished the corresponding product **3ea** in 77% yield with 5.6 : 1 dr. Benzaldehydes bearing halogens, such as 4-chlorobenzaldehyde, were also good substrates, as exemplified by the generation of **3fa** (80% yield and 6.0 : 1 dr). This product is primed for further functionalization by cross-coupling methods. Since fluorinated compounds play an extremely important role in medicinal chemistry,¹⁵ we next examined fluorinated benzaldehydes. Both 4-fluoro- (**1g**) and 4-trifluoromethoxy-benzaldehyde (**1h**) were excellent coupling partners, affording **3ga** and **3ha** in 91% (5.8 : 1 dr) and 87% yields (dr = 5.5 : 1) respectively. Benzaldehydes containing aromatic substituents such as 4-(2-thienyl) (**1i**) and 4-(*N*-pyrrolyl) (**1j**) provided products **3ia** and **3ja** in 85% (5.8 : 1 dr) and 82% (5.1 : 1 dr) yields, respectively. To our surprise, 4-phenyl benzaldehyde and 4-(2-pyridyl) benzaldehyde reacted to give the expected products **3ka** and **3la** in very good yields (83% and 86%) with excellent diastereoselectivities (16:1 and >20:1), respectively. In addition to benzaldehydes, heterocyclic aldehydes such as nicotinaldehyde (**1m**), quinoline-6-carbaldehyde (**1n**), benzofuran-5-carbaldehyde (**1o**) and thiophene-2-carbaldehyde (**1p**) were transformed to their corresponding products in good yields (76–86%) and diastereoselectivities (4.6 : 1 to 6.3 : 1).


Scheme 2. Substrate Scope of Aldehydes^a



^aReactions conducted under argon on 0.2 mmol scale. Yields are isolated yield. The dr was determined by LC-MS analysis of unpurified reaction mixtures.

Subsequently, the scope of azaaryl methylamines was explored (Scheme 3), beginning with the cyclic amine moiety. When the morpholino group of **2a** was replaced by thiomorpholino (**2b**), piperidine (**2c**) and 1-methylpiperazine (**2d**), the corresponding products **4ab**, **4ac** and **4ad** were obtained in 89%, 83% and 81% yields with dr values of 7 : 1, 12 : 1 and > 20 : 1, respectively. Acyclic amines *N,N*-dimethyl- (**2e**), *N*-benzyl-*N*-methyl- (**2f**) and *N,N*-diethyl- (**2g**) derivatives were also good coupling partners and their respective products **4ae**, **4af** and **4ag** were recovered in 70–86% yield and with 4.2 : 1 to >20 : 1 dr. To test the azaaryl group, 3-aminomethyl isoquinoline (**2h**) and 4-aminomethyl pyridine (**2i**) derivatives were examined and the corresponding products (**4ah** and **4ai**) were formed in 70% and 80% yield with 7 : 1 and 2 : 1 dr, respectively. Unfortunately, 3-aminomethyl pyridine derivatives and diphenylmethanes, which are significantly less acidic, did not react under our conditions. To obtain acceptable diastereoselectivity, 2-aminomethyl substituted pyridine derivatives, which can chelate to the main group metal of the base, are required.


Scheme 3. Substrate Scope of Azaaryl methylamines^a



^aReactions conducted under argon on 0.2 mmol scale. Yields are isolated. The dr was determined by LC-MS analysis of unpurified reaction mixtures.

To investigate the potential scalability (Scheme 4), a 5 mmol scale reaction with **1I** and **2a** was conducted. The desired product **3la** was obtained in 88% yield with greater than 20 : 1 dr. Condensation of **3la** with 3,5-dinitrobenzoyl chloride (1.2 equiv) and triethylamine (1.2 equiv) formed the expected amide, the structure of which was determined to be the *anti*-**4la'** product by X-ray crystallographic analysis (see Supporting Information for details).

Scheme 4. Scale-up to 5 mmol.



In conclusion, we have reported a straightforward and practical one-pot synthesis of vicinal diamines between azaaryl methylamines and aldehydes. The tandem reaction takes place without transition metal catalysts and provides an effective strategy to synthesize biologically active vicinal diamines. A wide range of electron-donating and withdrawing aryl substrates, including heterocycles, are viable substrates in our reaction. The synthetic value and practicality of this method were demonstrated with a highly diastereoselective gram scale reaction. We anticipate that this straightforward avenue to stitch together molecules with 3–4 nitrogen centers from readily available precursors will render it immediately useful to medicinal chemists. Further efforts will focus on the identification of a catalytic asymmetric route to these diamines.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website
Experimental procedures, characterization data, and NMR spectra (PDF)
CIF for compound **4la'** (CIF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: pwalsh@sas.upenn.edu

*E-mail: ias_jymao@njtech.edu.cn.

ORCID

Jianyou Mao: 0000-0003-0581-3978

Patrick J. Walsh: 0000-0001-8392-4150

ACKNOWLEDGMENT

The authors acknowledge National Natural Science Foundation of China (21801128 to J.M.), Natural Science Foundation of Jiangsu Province, China (BK20170965 to J.M.) and the Nanjing Tech University (3980001601 and 39837112) for financial support. PJW thanks the US National Science Foundation (CHE-1902509).

REFERENCES

(1) (a) Lucet, D.; Le Gall, T.; Mioskowski, C. The chemistry of vicinal diamines. *Angew. Chem., Int. Ed.* **1998**, *37*, 2580; (b) Viso, A.; Fernandez de la Pradilla, R.; Garcia, A.; Flores, A. α,β -Diamino acids. Biological significance and synthetic approaches. *Chem. Rev.* **2005**, *105*, 3167; (c) Kotti, S. R. S. S.; Timmons, C.; Li, G. Vicinal diamino functionalities as privileged structural elements in biologically active compounds and exploitation of their synthetic chemistry. *Chem. Biol. Drug Des.* **2006**, *67*, 101.

(2) Blakemore, D. C.; Castro, L.; Churcher, I.; Rees, D. C.; Thomas, A. W.; Wilson, D. M.; Wood, A. Organic synthesis provides opportunities to transform drug discovery. *Nat. Chem.* **2018**, *10*, 383.

(3) (a) Shen, S.-J.; Zhu, C.-L.; Lu, D.-F.; Xu, H. Iron-Catalyzed Direct Olefin Diazidation via Peroxyester Activation Promoted by Nitrogen-Based Ligands. *ACS Catal.* **2018**, *8*, 4473; (b) Yuan, Y.-A.; Lu, D.-F.; Chen, Y.-R.; Xu, H. Iron-Catalyzed Direct Diazidation for a Broad Range of Olefins. *Angew. Chem., Int. Ed.* **2016**, *55*, 534; (c) Zhao, B.; Peng, X.; Cui, S.; Shi, Y. Cu(I)-Catalyzed Regioselective Diamination of Conjugated Dienes via Dual Mechanistic Pathways. *J. Am. Chem. Soc.* **2010**, *132*, 11009.

(4) (a) Neumann, W. L.; Rogic, M. M.; Dunn, T. J. The stereoselective synthesis of functionalized vicinal diamine systems by the double allylation reactions of protected 1,2-bisimine precursors. *Tetrahedron Lett.* **1991**, *32*, 5865; (b) Alexakis, A.; Tomassini, A.; Chouillet, C.; Roland, S.; Mangeney, P.; Bernardinelli, G. A new efficient synthesis of (R,R)-2,2'-bipyrrolidine: an interesting chiral 1,2-diamine with C2 symmetry. *Angew. Chem., Int. Ed.* **2000**, *39*, 4093; (c) Alvaro, G.; Savoia, D. Addition of organometallic reagents to imines bearing stereogenic N-substituents. Stereochemical models explaining the 1,3-asymmetric induction. *Synlett* **2002**, 651; (d) Sun, X.; Wang, S.; Sun, S.; Zhu, J.; Deng, J. Highly diastereoselective and enantioselective addition of organometallic reagents to a chiral C2-symmetrical bisimine. *Synlett* **2005**, 2776; (e) Vemula, R.; Wilde, N. C.; Goreti, R.; Corey, E. J. A New, Short, and Stereocontrolled Synthesis of C2-Symmetric 1,2-Diamines. *Org. Lett.* **2017**, *19*, 3883.

(5) (a) Zhong, Y.-W.; Izumi, K.; Xu, M.-H.; Lin, G.-Q. Highly Diastereoselective and Enantioselective Synthesis of Enantiopure C2-Symmetrical Vicinal Diamines by Reductive Homocoupling of Chiral N-tert-Butanesulfinyl Imines. *Org. Lett.* **2004**, *6*, 4747; (b) Faugeroux, V.; Genisson, Y. The imino-pinacol coupling reaction. *Curr. Org. Chem.* **2008**, *12*, 751; (c) Rieke, R. D.; Kim, S.-H. New reagent for reductive coupling of carbonyl and imine compounds: highly reactive manganese-mediated pinacol coupling of aryl aldehydes, aryl ketones, and aldimines. *J. Org. Chem.* **1998**, *63*, 5235; (d) Smith, J. G.; Ho, I. Reductive dimerization of Schiff bases by alkali metals. Isomerization of the dimeric dianions. *J. Org. Chem.* **1972**, *37*, 653; (e) Campos, P. J.; Arranz, J.; Rodriguez, M. A. Photoreductive Coupling of Aldimines. Synthesis of C2 Symmetrical Diamines. *Tetrahedron* **2000**, *56*, 7285; (f) Kitajima, K.; Nagatomo, R.; Fujimoto, T. Diastereoselective imino-pinacol coupling of chiral imines promoted by Mg(0) and a Grignard reagent. *Tetrahedron* **2015**, *71*, 977.

(6) (a) Huang, Y.-Y.; Lv, Z.-C.; Yang, X.; Wang, Z.-L.; Zou, X.-X.; Zhao, Z.-N.; Chen, F. Nucleophilic ring opening of aziridines with amines under catalyst- and solvent-free conditions. *Green Chem.* **2017**, *19*, 924; (b) Hajra, S.; Singha Roy, S.; Biswas, A.; Saleh, S. A. Catalyst-Free Ring Opening of Spiroaziridine Oxindoles by Heteronucleophiles: An Approach to the Synthesis of Enantiopure 3-Substituted Oxindoles. *J. Org. Chem.* **2018**, *83*, 3633; (c) Samanta, S.; Chatterjee, R.; Mahato, S.; Hajra, A.; Santra, S.; Zyryanov, G. V.; Majee, A. Synthesis of diverse β -(nitrooxy)-substituted amines by regioselective ring-opening of aziridines under neat conditions. *Synth. Commun.* **2018**, *48*, 1857.

(7) (a) Noble, A.; Anderson, J. C. Nitro-Mannich Reaction. *Chem. Rev.* **2013**, *113*, 2887; (b) Marcantonio, E.; Petrini, M. Recent Developments in the Stereoselective Synthesis of Nitrogen-Containing Heterocycles using N-Acylimines as Reactive Substrates. *Adv. Synth. Catal.* **2016**, *358*, 3657; (c) Dudek, A.; Mlynarski, J. Iron-Catalyzed Asymmetric Nitro-Mannich Reaction. *J. Org. Chem.* **2017**, *82*, 11218; (d) Hajra, S.; Jana, B. Quinine-Based Trifunctional Organocatalyst for Tandem Aza-Henry Reaction-Cyclization: Asymmetric Synthesis of Spiroxindole-Pyrrolidine/Piperidines. *Org. Lett.* **2017**, *19*, 4778; (e) Lu, N.; Li, R.; Wei, Z.; Cao, J.; Liang, D.; Lin, Y.; Duan, H. Enantio- and Diastereoselective Nitro-Mannich Reaction of α -Aryl Nitromethanes with Amidosulfones Catalyzed by Phase-Transfer Catalysts. *J. Org. Chem.* **2017**, *82*, 4668; (f) Prasad, S. S.; Senthilkumar, S.; Srivastava, A.; Baskaran, S. Iminosugar C-Nitromethyl Glycosides and Divergent Synthesis of Bicyclic Iminosugars. *Org. Lett.* **2017**, *19*, 4403; (g) Ouyang, L.; Zhan, L.; Li, J.; Zhang, Q.; Qi, C.; Wu, W.; Jiang, H. Pd-Catalyzed Three-Component Reaction of Anilines, Ethyl Vinyl Ether, and Nitro-Paraffin: Assembly of β -Nitroamines. *Org. Lett.* **2018**, *20*, 550; (h) Sprague, D. J.; Singh, A.; Johnston, J. N. Diastereo- and enantioselective additions of α -nitro esters to imines for anti- α,β -diamino acid synthesis with α -alkyl-substitution. *Chem. Sci.* **2018**, *9*, 2336.

(8) (a) Kise, N.; Kashiwagi, K.; Watanabe, M.; Yoshida, J.-i. Stereoselective Synthesis of trans-4,5-Disubstituted 1,3-Imidazolidin-2-ones by Reaction of an α -Nitrogen Carbanion with Imines. *J. Org. Chem.* **1996**, *61*, 428; (b) Park, Y. S.; Boys, M. L.; Beak, P. (-)-Sparteine-Mediated α -Lithiation of N-Boc-N-(p-methoxyphenyl)benzylamine: Enantioselective Syntheses of (S) and (R) Mono- and Disubstituted N-Boc-benzylamines. *J. Am. Chem. Soc.* **1996**, *118*, 3757; (c) Ooi, T.; Kameda, M.; Fujii, J.; Maruoka, K. Catalytic Asymmetric Synthesis of a Nitrogen Analogue of Dialkyl Tartrate by Direct Mannich Reaction under Phase-Transfer Conditions. *Org. Lett.* **2004**, *6*, 2397; (d) Chen, Y.-J.; Seki, K.; Yamashita, Y.; Kobayashi, S. Catalytic Carbon-Carbon Bond-Forming Reactions of Aminoalkane Derivatives with Imines. *J. Am. Chem. Soc.* **2010**, *132*, 3244; (e) Kano, T.; Sakamoto, R.; Akakura, M.; Maruoka, K. Stereocontrolled Synthesis of Vicinal Diamines by Organocatalytic Asymmetric Mannich Reaction of N-Protected Aminocetaldehydes: Formal Synthesis of (-)-Aglastatin A. *J. Am. Chem. Soc.* **2012**, *134*, 7516; (f) Mitchell, E. A.; Peschiulli, A.; Lefevre, N.; Meerpoel, L.; Maes, B. U. W. Direct α -Functionalization of Saturated Cyclic Amines. *Chem. - Eur. J.* **2012**, *18*, 10092; (g) Ding, L.; Chen, J.; Hu, Y.; Xu, J.; Gong, X.; Xu, D.; Zhao, B.; Li, H. Aminative Umpolung of Aldehydes to α -Amino Anion Equivalents for Pd-Catalyzed Allylation: An Efficient Synthesis of Homoallylic Amines. *Org. Lett.* **2014**, *16*, 720; (h) Liu, X.; Gao, A.; Ding, L.; Xu, J.; Zhao, B. Aminative Umpolung Synthesis of Aryl Vicinal Diamines from Aromatic Aldehydes. *Org. Lett.* **2014**, *16*, 2118; (i) Lin, C.-Y.; Sun, Z.; Xu, Y.-J.; Lu, C.-D. Synthesis of Aryl anti-Vicinal Diamines via Aza-Brook Rearrangement-Initiated Nucleophilic Addition of α -Silylamines to Imines. *J. Org. Chem.* **2015**, *80*, 3714; (j) Lin, S.; Kawato, Y.; Kumagai, N.; Shibusaki, M. Catalytic Asymmetric Mannich-Type Reaction of N-Alkylidene- α -Aminoacetonitrile with Ketimines. *Angew. Chem., Int. Ed.* **2015**, *54*, 5183; (k) Li, K.; Weber, A. E.; Tseng, L.; Malcolmson, S. J. Diastereoselective and Enantiospecific Synthesis of 1,3-Diamines via 2-Azaallyl Anion Benzylidene Ring-Opening of Aziridines. *Org. Lett.* **2017**, *19*, 4239; (l) Liu, F.; Zhao, G.; Cai, W.; Xu, D.; Zhao, B. Aminative Umpolung cyclization for synthesis of chiral exocyclic vicinal diamines. *Org. Biomol. Chem.* **2018**, *16*, 7498; (m) Shao, X.; Li, K.; Malcolmson, S. J. Enantioselective Synthesis of anti-1,2-Diamines by Cu-Catalyzed Reductive Couplings of Azadienes with Aldimines and Ketimines. *J. Am. Chem. Soc.* **2018**, *140*, 7083; (n) Tang, S.; Zhang, X.; Sun, J.; Niu, D.; Chrunga, J. J. 2-Azaallyl Anions, 2-Azaallyl Cations, 2-Azaallyl Radicals, and Azomethine Ylides. *Chem. Rev.* **2018**, *118*, 10393; (o) Xia, Q.; Tian, H.; Dong, J.; Qu, Y.; Li, L.; Song, H.; Liu, Y.; Wang, Q. N-Arylamines Coupled with Aldehydes, Ketones, and Imines by Means of Photocatalytic Proton-Coupled Electron Transfer. *Chem. - Eur. J.* **2018**, *24*, 9269; (p) Wang, M.; Fan, Q.; Jiang, X. Nitrogen-4

Iodine Exchange of Diaryliodonium Salts: Access to Acridine and Carbazole. *Org. Lett.* **2018**, *20*, 216.

(9) Wang, Z.; Zheng, Z.; Xu, X.; Mao, J.; Walsh, P. J. One-pot aminobenzylation of aldehydes with toluenes. *Nat. Commun.* **2018**, *9*, 1.

(10) Wang, R.; Ma, M.; Gong, X.; Fan, X.; Walsh, P. J. Reductive Cross-Coupling of Aldehydes and Imines Mediated by Visible Light Photoredox Catalysis. *Org. Lett.* **2019**, *21*, 23.

(11) Mao, J.; Wang, Z.; Xu, X.; Liu, G.; Jiang, R.; Guan, H.; Zheng, Z.; Walsh, P. J. Synthesis of Indoles through Domino Reactions of 2-Fluorotoluenes and Nitriles. *Angew. Chem., Int. Ed.* **2019**, *58*, 11033.

(12) (a) Pardue, D. B.; Gustafson, S. J.; Periana, R. A.; Ess, D. H.; Cundari, T. R. Computational study of carbon-hydrogen bond deprotonation by alkali metal superbases. *Comput. Theor. Chem.* **2013**, *1019*, 85; (b) Sha, S.-C.; Tcyruulinikov, S.; Li, M.; Hu, B.; Fu, Y.; Kozlowski, M. C.; Walsh, P. J. Cation- π Interactions in the Benzylic Arylation of Toluenes with Bimetallic Catalysts. *J. Am. Chem. Soc.* **2018**, *140*, 12415.

(13) Arndts, D.; Loesel, W.; Roos, O.; (Boehringer Ingelheim KG, Germany). Application: DE, 1993, p 11 pp.

(14) (a) Lappert, M. F.; Slade, M. J.; Singh, A.; Atwood, J. L.; Rogers, R. D.; Shakir, R. Structure and reactivity of sterically hindered lithium amides and their diethyl etherates: crystal and molecular structures of $[\text{Li}\{\text{N}(\text{SiMe}_3)_2\}\{\text{OEt}_2\}]_2$ and tetrakis(2,2,6,6-tetramethylpiperidinatolithium). *J. Am. Chem. Soc.* **1983**, *105*, 302; (b) Henderson, K. W.; Walther, D. S.; Williard, P. G. Identification of a Unimetal Complex of Bases by ^6Li NMR Spectroscopy and Single-Crystal Analysis. *J. Am. Chem. Soc.* **1995**, *117*, 8680; (c) Armstrong, D. R.; Henderson, K. W.; Kennedy, A. R.; Kerr, W. J.; Mair, F. S.; Moir, J. H.; Moran, P. H.; Snaith, R. Structural studies of the chiral lithium amides $[\{\text{PhC}(\text{H})\text{Me}\}_2\text{NLi}]$ and $[\text{PhCH}_2\{\text{PhC}(\text{H})\text{Me}\}\text{NLi}\cdot\text{THF}]$ derived from α -methylbenzylamine. *J. Chem. Soc., Dalton Trans.* **1999**, 4063.

(15) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. Fluorine in medicinal chemistry. *Chem. Soc. Rev.* **2008**, *37*, 320.