

# Copper-Catalyzed Intermolecular Difunctionalization of Styrenes with Thiosulfonates and Arylboronic Acids via a Radical Relay Pathway

Qingjin Liang,<sup>†,‡</sup> Patrick J. Walsh<sup>§</sup> and Tiezheng Jia<sup>\*,‡</sup>

<sup>†</sup>School of Chemistry and Chemical Engineering, Harbin Institute of Technology, Harbin 150001, P. R. China

<sup>‡</sup>Department of Chemistry, Southern University of Science and Technology, Shenzhen, Guangdong, 518055, P. R. China

<sup>§</sup>Roy and Diana Vagelos Laboratories, Penn/Merck Laboratory for High-Throughput Experimentation, Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, Pennsylvania 19104-6323, United States

**KEYWORDS:** *copper catalysis • radical relay • sulfones • alkenes • difunctionalization*

**ABSTRACT:** A general and practical copper-catalyzed intermolecular difunctionalization strategy of styrenes with methyl thiosulfonates and arylboronic acids has been developed. This method provides an efficient and straightforward avenue to a broad range of 2,2-diarylethylsulfone derivatives from readily available methyl thiosulfonates and commercially available styrene and arylboronic acid derivatives. The diverse substrate scope attests to the high functional group tolerance of this reaction. The mild nature of this protocol make it suitable for late-stage functionalization of bioactive natural products. Mechanistic investigations support the role of sulfonyl radicals and corroborate a copper-catalyzed radical relay pathway.


Sulfones are prevalent scaffolds in natural products,<sup>1</sup> synthetic bioactive molecules,<sup>2</sup> and marketed therapeutics (such as SB-3CT for the treatment of migraine headaches<sup>3</sup> and anti-androgen receptor Casodex<sup>34</sup>). In fact, a study of drug likeness analysis found that alkyl aryl sulfones are among the top five drug skeletal fragments.<sup>5</sup> Sulfones also find widespread applications in material science<sup>6</sup> and agricultural industries.<sup>7</sup> They serve as important intermediates in organic synthesis,<sup>8</sup> as exemplified by the well-known Julia–Lythgoe olefination.<sup>9</sup> Therefore, methods for the construction of sulfones have attracted significant attention.

Despite the importance of sulfones in the applications outlined above, 2,2-diarylethylsulfones have received little attention, probably due to a lack of efficient and straightforward synthetic approaches.


Aliphatic sulfones can be prepared from  $\alpha,\beta$ -unsaturated sulfone precursors<sup>10</sup> through hydrogenation, including asymmetric hydrogenations.<sup>11</sup> In 1998, the Ranu group reported the Pd/C-catalyzed hydrogenation of  $\alpha,\beta$ -unsaturated sulfones using ammonium formate.<sup>12</sup> Inspired by this study, an elegant enantioselective rhodium-catalyzed hydrogenation of  $\alpha,\beta$ -unsaturated sulfones was developed by the Hou group in 2019 (Scheme 1a).<sup>13</sup> In this procedure, both the (*E*)- and (*Z*)-isomers gave excellent enantioselectivities leading to opposite configurations of the products. The synthesis of the (*E*)- and (*Z*)- $\alpha,\beta$ -unsaturated sulfone precursors required 4 steps. In 2004, Lautens and coworkers described a rhodium-catalyzed Heck-type coupling of arylboronic acids with  $\alpha,\beta$ -unsaturated sulfones.<sup>14</sup> By employing a rhodium catalyst bearing an enantioenriched tetrafluorobenzobarrelene-based ligand, an enantioselective 1,4-addition of boronic acids to  $\alpha,\beta$ -unsaturated sulfones was accomplished by Hayashi and his group (Scheme 1b).<sup>15</sup> Here also, the requisite  $\alpha,\beta$ -unsaturated sulfones required several steps to prepare,<sup>15</sup> reducing the overall synthetic efficiency.

Scheme 1. Previous Synthetic Approaches to 2,2-Diarylethylsulfones.

a) Hydrogenation: Ranu (1998), Hou (2019)



b) 1,4-Addition: Lautens (2004), Hayashi (2012)




In recent years, copper-catalyzed coupling reactions between *in situ* formed radicals and arylboronic acids have drawn much attention due to copper's low cost, bio-compatibility, and complimentary reactivity to heavier d-block metals.<sup>16</sup> Copper-catalyzed difunctionalization of alkenes represents one of the most powerful and straightforward approaches to rapidly increase molecular complexity, as two chemical bonds are generated in one step.<sup>17</sup> We envisioned that the 2,2-diarylethylsulfones outlined above could be accessed via a three-component process involving the arylsulfonylation of styrenes (Figure 1).

To develop the arylsulfonylation of styrenes, three crucial requirements would need to be satisfied (Figure 1). First, a sulfonyl radical source that is compatible with the catalyst and intermediates in the arylsulfonylation would need to be identified.<sup>18</sup> Second, the competitive two-component coupling between sulfonyl radicals and boronic acids would need to be minimized (see byproduct 1, Figure 1).<sup>19</sup> Finally, an appropriate oxidant for the selective oxidation of Cu(I) to Cu(II) must be chosen that would not oxidize other reactive intermediates (such as oxidize a punitive benzyl

radical to the corresponding cation, leading to byproducts **2**,<sup>10a</sup> **3**,<sup>20</sup> or **4**,<sup>21</sup> (Figure 1).

The choice of sulfonyl radical source is important to develop user-friendly general methods to make C–S bonds. Several commonly used sulfonyl radical sources suffer from drawbacks that decrease their attractiveness. For instance, sulfonyl chlorides require expensive silver salts to trap chloride and prevent its undesired radical background additions.<sup>22</sup> The activation of sulfonyl hydrazides usually employ high temperatures, and the release of nitrogen gas raises safety concerns.<sup>23</sup> For our difunctionalization of styrenes to afford 2,2-diarylethylsulfone scaffolds, methyl thiosulfonates were chosen as the sulfonyl radical source.<sup>24</sup> Methyl thiosulfonates are readily available, odorless, and safe. Herein, we report an unprecedented copper-catalyzed intermolecular difunctionalization of styrenes with methyl thiosulfonates and arylboronic acids to afford 2,2-diarylethylsulfones (Figure 1). Our copper-catalyzed difunctionalization reaction features: (a) commercially available styrenes and arylboronic acids and readily available methyl thiosulfonates as starting materials; (b) cheap and abundant copper catalysts; and (c) mild reaction conditions with broad functional group compatibility. It represents an efficient and streamlined method to synthesize 2,2-diarylethylsulfone scaffolds.

Figure 1. Synthesis of 2,2-Diarylsulfones via a Difunctionalization of Styrenes Strategy and the Potential Challenges.



For the reaction optimization, we chose styrene (**1a**), *S*-methyl 4-methylbenzenesulfonothioate (**2a**) and phenylboronic acid (**3a**) as the model substrates. When 2.0 equiv  $K_2CO_3$  was used as base, 2.0 equiv  $BzOO^tBu$  as oxidant, 10 mol %  $Cu(OTf)_2$  and 15 mol % 4,4'-di-*tert*-butyl-2,2'-bipyridine (dtbbpy) as the catalyst at 50 °C in acetone, the desired product **4aa** was isolated in 34% AY (AY = assay yield, determined by  $^1H$  NMR integration of the crude product against an internal standard, Table 1, entry 1). Encouraged by this result, three additional bases ( $NaHCO_3$ ,  $CS_2CO_3$ , and  $KHCO_3$ ) were examined. These experiments revealed that  $KHCO_3$  was the top choice, generating **4aa** in 48% AY (Table 1, entry 4 vs. entries 1–3). Thus,  $KHCO_3$  was used for further optimization. Three additional copper salts [ $CuBr$ ,  $Cu(NCCH_3)_4PF_6$ , and  $CuI$ ] were surveyed (Table 1, entries 5–7). The yield of **4aa** increased to 60% when  $CuI$  was utilized (Table 1, entry 7). In subsequent solvent screens, *N,N*-dimethylformamide (DMF) led to formation of **4aa** in 63% yield. This reaction also resulted in the formation of 20% AY of 4-tolyl phenyl sulfone (**4aa'**) from the two-component coupling side reaction between **2a** and **3a** (Table 1, entry 8). Formation of 4-tolyl phenyl sulfone (**4aa'**) was not observed in the prior experiments in acetone (entries 1–7). This observation inspired us to turn to mix solvents with DMF and acetone (Table 1, entries 9–11). When the volume ratio of acetone to DMF was 6 : 1, 74% AY of **4aa** was obtained (Table 1, entry 10). In contrast, an acetone to DMF ratio of 8 : 1, furnished the product in 66% AY (Table 1, entry 11). Upon decreasing the temperature from 50 to 40 °C, **4aa** was formed in 82% AY and 80% isolated yield (Table 1, entry 12). When the reaction was carried out at room temperature,

the AY dropped to 48% (Table 1, entry 13). We next examine the catalyst loading. When reducing the catalyst loading to 5 mol % (Table 1, entry 14) the AY dropped to 72%. Thus, subsequent reactions were conducted with 10 mol % copper.

In a series of control experiments,  $Cu(I)$ , dtbbpy and  $KHCO_3$  were demonstrated to be essential for the catalytic reaction. No desired product was observed in the absence of any of these components (Table 1, entries 15–17). Therefore, the optimized conditions were: 10 mol %  $CuI$  and 15 mol% dtbbpy as the catalyst, styrene (**1a**) as the limiting reagent, 2.0 equiv of thiosulfonate (**2a**), 2.5 equiv of **3a**, 2.5 equiv of  $KHCO_3$ , and 3.0 equiv  $BzOO^tBu$  as both oxidant and radical initiator in a mix solvent of acetone and DMF (6 : 1 volume ratio) at 40 °C for 12 h (for additional screening results, see SI).

Table 1. Optimization of the Copper-Catalyzed Difunctionalization of Styrene with **2a** and **3a**.<sup>a</sup>


| $Ph-CH=CH_2 + TsSMe + PhB(OH)_2 \xrightarrow[\text{base, solvent}]{[Cu]/\text{ligand}} Ph-CH(Ph)-CH_2-Ts$ |                    |            |                                 |      |                                    |
|-----------------------------------------------------------------------------------------------------------|--------------------|------------|---------------------------------|------|------------------------------------|
| entry                                                                                                     | catalyst           | base       | solvent                         | T/°C | <b>4aa</b><br>(AY <sup>b</sup> /%) |
| 1 <sup>c</sup>                                                                                            | $Cu(OTf)_2$        | $K_2CO_3$  | acetone                         | 50   | 34                                 |
| 2 <sup>c</sup>                                                                                            | $Cu(OTf)_2$        | $NaHCO_3$  | acetone                         | 50   | 17                                 |
| 3 <sup>c</sup>                                                                                            | $Cu(OTf)_2$        | $CS_2CO_3$ | acetone                         | 50   | N.P                                |
| 4 <sup>c</sup>                                                                                            | $Cu(OTf)_2$        | $KHCO_3$   | acetone                         | 50   | 48                                 |
| 5 <sup>c</sup>                                                                                            | $CuBr$             | $KHCO_3$   | acetone                         | 50   | trace                              |
| 6 <sup>c</sup>                                                                                            | $Cu(CH_3CN)_4PF_6$ | $KHCO_3$   | acetone                         | 50   | 40                                 |
| 7 <sup>c</sup>                                                                                            | $CuI$              | $KHCO_3$   | acetone                         | 50   | 60                                 |
| 8                                                                                                         | $CuI$              | $KHCO_3$   | DMF                             | 50   | 63                                 |
| 9                                                                                                         | $CuI$              | $KHCO_3$   | acetone: DMF = 4:1 <sup>d</sup> | 50   | 68                                 |
| 10                                                                                                        | $CuI$              | $KHCO_3$   | acetone: DMF = 6:1 <sup>d</sup> | 50   | 74                                 |
| 11                                                                                                        | $CuI$              | $KHCO_3$   | acetone: DMF = 8:1 <sup>d</sup> | 50   | 66                                 |
| 12                                                                                                        | $CuI$              | $KHCO_3$   | acetone: DMF = 6:1 <sup>d</sup> | 40   | 82(80 <sup>e</sup> )               |
| 13                                                                                                        | $CuI$              | $KHCO_3$   | acetone: DMF = 6:1 <sup>d</sup> | R.T  | 48                                 |
| 14 <sup>f</sup>                                                                                           | $CuI$              | $KHCO_3$   | acetone: DMF = 6:1 <sup>d</sup> | 40   | 72                                 |
| 15 <sup>g</sup>                                                                                           | $CuI$              | $KHCO_3$   | Acetone: DMF = 6:1 <sup>d</sup> | 40   | trace                              |
| 16                                                                                                        | -                  | $KHCO_3$   | acetone: DMF = 6:1 <sup>d</sup> | 40   | N.P                                |

17      CuI      -      acetone: 40      N.P  
 DMF =  
 6:1<sup>d</sup>

<sup>a</sup>Reaction conditions: **1a** (0.1 mmol), **2a** (2.0 equiv), **3a** (2.5 equiv), base (2.5 equiv), BzOO<sup>t</sup>Bu (3.0 equiv), copper catalyst (10 mol %), dtbbpy (15 mol %), acetone (0.6 mL) and DMF (0.1 mL), 40 °C for 12 h. <sup>b</sup>AY = Assay yield, determined by <sup>1</sup>H NMR analysis of the crude reaction mixtures using 0.1 mmol CH<sub>2</sub>Br<sub>2</sub> (7.0 μL) as internal standard. <sup>c</sup>**3a** (2.0 equiv), KHCO<sub>3</sub> (2.0 equiv), BzOO<sup>t</sup>Bu (2.0 equiv) and solvent (1.0 mL). <sup>d</sup>Volume ratio. <sup>e</sup>Isolated yield. <sup>f</sup>CuI (5 mol %), dtbbpy (7.5 mol %). <sup>g</sup>Without dtbbpy.

With the optimized conditions in hand, we evaluated the substrate scope of arylboronic acids in the three-component coupling reaction to generate 2,2-diarylethylsulfones. As shown in Table 2, the parent phenylboronic acid gave the product **4aa** in 80% yield. Arylboronic acids with electronegative or electron-withdrawing groups reacted smoothly with **1a** and **2a** to give the target products **4ab**, **4ac**, **4ad** and **4ae** in 56–83% yields. Likewise, *ortho*- and *meta*-substituted phenylboronic acids furnished the products **4af–h** in 57–71% yield. Of note, arylboronic acids possessing aldehyde (**3i**), ketone (**3j**) and ester (**3k**), provided **4ai–k** in 52–88% yields. Heterocycles are important in medicinal chemistry, but often challenging to install.<sup>25</sup> When 2-thiophenyl and 3-thiophenyl boronic acids were employed, **4al** and **4am** were generated from **1a** and **2a** under our standard reaction conditions in 32 and 44% yields, respectively.

Table 2. Substrate Scope of Arylboronic Acids in the Copper-Catalyzed Difunctionalization with **1a** and **2a**.<sup>a</sup>




<sup>a</sup>Reaction conditions: **1a** (0.1 mmol), **2a** (2.0 equiv), **3** (2.5 equiv), KHCO<sub>3</sub> (2.5 equiv), BzOO<sup>t</sup>Bu (3.0 equiv), CuI (10 mol %), dtbbpy (15 mol %), acetone (0.6 mL) and DMF (0.1 mL), 40 °C for 12 h.

We next turned our attention to the substrate scope of styrenes in the three-component difunctionalization with **2a** and **3a**. Overall, this method proved general with respect to the styrenes, providing access to a wide range of 2,2-diarylethylsulfones (Table 3). Styrenes with electron-donating groups, such as *para*-<sup>t</sup>Bu, -OMe or -Ph were well tolerated, providing **5aa–c** in 70–87% yields. Styrenes bearing electron-withdrawing 4-F, 4-Cl, 4-Br and 4-NO<sub>2</sub> (**1e–g**) were good coupling partners, delivering the desired products **4ab**,

**4ae** and **5ad** in 62–66% yields. When sterically hindered 2-methyl styrene (**1h**) was utilized as the coupling partner, the corresponding product **5ae** was afforded in 65% yield. In addition, *meta*-bromostyrene (**1i**) exhibited good reactivity generating the expected product **5af** in 79% yield. 1-(Chloromethyl)-4-vinylbenzene is a substrate that would react rapidly with any anionic sulfur-based nucleophiles. This substrate was suitable for the multicomponent reaction, affording **5ag** in 64% yield, leaving the *para*-chloromethyl group for further modification. As noted earlier, heterocycles are important in medicinal chemistry.<sup>25</sup> Our catalytic protocol proved applicable to a variety of heterocyclic styrene derivatives.<sup>2</sup> Pyridyl (**1k**, **1l**), thiophenyl (**1m**, **1n**), furanyl (**1p**), quinolonyl (**1q**), isoquinolonyl (**1r**), and indolyl (**1s**) alkenes furnished the desired 2,2-diarylethylsulfones (**5ah–l**, **4al–m**, and **5aj–n**) in 30–64% yields. The internal styrene derivative, indene (**1t**), was successfully applied to this transformation, delivering the corresponding *trans*-sulfone product **5ao** as a single diastereomer in 42% yield.

Table 3. Substrate Scope of Styrenes in Copper-Catalyzed Difunctionalization with **2a** and **3a**.<sup>a</sup>




<sup>a</sup>Reaction conditions: **1** (0.1 mmol), **2a** (2.0 equiv), **3a** (2.5 equiv), KHCO<sub>3</sub> (2.5 equiv), BzOO<sup>t</sup>Bu (3.0 equiv), CuI (10 mol %), dtbbpy (15 mol %), acetone (0.6 mL) and DMF (0.1 mL), 40 °C for 12 h.

The generality of methyl thiosulfonates in copper-catalyzed difunctionalization with **1a** and **3a** was next investigated. The parent 2,2-diphenylethyl phenyl sulfone (**6aa**) was generated in 70% yield. The electron-donating 4-methoxy substituent exerts a negligible effect on the outcome of the transformation, affording **6ab** in 76% yield. Aryl thiosulfonates with electronegative or electron-withdrawing groups, such as 4-F, 4-Cl, 4-Br and 4-NO<sub>2</sub>, coupled under the standard conditions, providing **6ac–6af** in 51–73% yields. Using sterically demanding 2-methyl thiosulfonate (**2h**) as coupling partner in the difunctionalization led to formation of **6ag** in 75% yield. An acetyl group on the phenyl sulfone (**2i**) was tolerated owing to the mild reaction conditions, furnishing the product **6ah** in 60% yield. 2,2-Diphenylethyl 2-thiophenyl sulfone (**6ai**), a heteroaryl sulfone, was produced in 62% yield. It is noteworthy that methyl thiosulfonate **2j**, an example of an alkyl thiosulfonate, also exhibited good reactivity in the copper-

catalyzed difunctionalization with **1a** and **3a**, providing **6aj** in 50% yield under the optimized conditions.


Table 4. Substrate Scope of Thiosulfonates in Copper-Catalyzed Difunctionalization with **1a** and **3a**.<sup>a</sup>



<sup>a</sup>Reaction conditions: **1a** (0.1 mmol), **2** (2.0 equiv), **3a** (2.5 equiv),  $\text{KHCO}_3$  (2.5 equiv),  $\text{BzOO}^t\text{Bu}$  (3.0 equiv),  $\text{CuI}$  (10 mol %), dtbbpy (15 mol %), acetone (0.6 mL) and DMF (0.1 mL), 40 °C for 12 h.

To explore the potential synthetic utility of our method, late-stage functionalization of bioactive natural products was conducted. Alkenes bearing cromaril, estrone and tocol, three well known bioactive natural products, proceeded smoothly under the optimized reaction conditions to provide **5ap**, **5aq**, and **5ar** in synthetically useful yields (40–52%) (Scheme 2).

Scheme 2. Late-Stage Functionalization of Natural Products Based on Copper-Catalyzed Difunctionalization of Styrenes with **2a** and **3a**.




Several experiments to probe the mechanism of this catalytic transformation were performed (Scheme 3). First, when 2 equiv of radical scavenger 2,2,6,6-tetramethylpiperidinyl-1-oxide (TEMPO) was added to the reaction under otherwise standard conditions, the formation of product **4aa** was inhibited and **1a** was recovered in near quantitative assay yield, as determined by GC (Scheme 3.1). Both *p*-methyl benzene sulfonate anion (**7a**) and 2,2,6,6-tetramethylpiperidine (**8a**) were detected in the reaction mixture via HRMS analysis (see Table S8). Based on these observations, we speculate that TEMPO captured the sulfonyl radical, which was followed by reduction to **7a** and **8a**. These results are consistent with a radical pathway. When 2.0 equiv of the spin trap 5,5-


dimethyl-1-pyrroline *N*-oxide (DMPO) was added to the model reaction, the formation of **9a** was confirmed by HRMS (Scheme 3b) and electron paramagnetic resonance spectroscopy (EPR) (Scheme 3c). This result supports the intermediacy of the sulfonyl radical in this process (Scheme 3b). In addition, a series of UV–vis experiments was performed to probe the oxidation states of the copper catalyst (Scheme 3d). When copper(I) iodide was mixed together with dtbbpy ligand, the UV absorption of  $\text{Cu(I)}$  complex is ca. 431 nm (Scheme 3d). However, the UV band shifted to 703 nm when copper(I) iodide and dtbbpy were mixed together with  $\text{BzOO}^t\text{Bu}$ , which is consistent with the d-d transition of  $\text{Cu(II)}$  species.<sup>26</sup> To verify the existence of  $\text{Cu(II)}$  species, the solution of  $\text{CuI}$ , dtbbpy and  $\text{BzOO}^t\text{Bu}$  was monitored by EPR, which clearly exhibited signals for a  $\text{Cu(II)}$  species (see Figure S2 in the SI). When the reaction mixture was monitored by UV, the copper absorption is approximately 684 nm, which was attributed to a  $\text{Cu(II)}$  species as the catalyst resting state.

Scheme 3. Control Experiments to Probe the Reaction Mechanism.


### 1) Radical trapping experiments with TEMPO




### 2) EPR studies with DMPO



### 3) EPR spectra of 9a (the fitting data and experimental data)



### 4) UV-Vis spectra studies




Based on the experimental results above, and related literature reports,<sup>16c</sup> a plausible reaction mechanism is illustrated in Figure 3.  $\text{CuI}$  binds the ligand dtbbpy to give  $(\text{dtbbpy})\text{CuI}$  (**A**), which reacts



with  $\text{BzOO}^t\text{Bu}$  to generate the copper(II) complex **B**, and  $\bullet\text{OCMe}_3$  (**E**).<sup>27</sup> Then  $\bullet\text{OCMe}_3$  (**E**) abstracts hydrogen from DMF to generate the formyl radical **F**.<sup>28</sup> A radical relay process between **2a** and the formyl radical **F** affords sulfonyl radical (**G**), and generates the thiocarbamate, which was formed in 72% AY by  $^1\text{H}$  NMR spectroscopy (and fully characterized, see SI for details).  $\text{Ts}\bullet$  (**G**) undergoes addition to styrene **1a**, generating the benzyl radical **H**. Meanwhile, the  $\text{Cu(II)}$  complex **B** reacts with arylboronic acid **3a** via transmetalation to form the aryl copper(II) species **C**, which then undergoes oxidative trapping with the benzylic radical **H** to afford a reactive  $\text{Cu(III)}$  intermediate **D**.<sup>29</sup> Complex **D** undergoes reductive elimination to furnish **4aa** and regenerate the copper(I) species **A** to close the catalytic cycle. As mentioned earlier, the reaction forming **5ag**, containing a benzylic chloromethyl group, suggests that anionic sulfur species are unlikely to be intermediates in this process. Anionic sulfur compounds undergo rapid  $\text{S}_{\text{N}}2$  reactions with benzyl chloride derivatives.

Figure 2. Proposed Mechanism of Copper-Catalyzed Difunctionalization of Styrenes.



In summary, a novel copper-catalyzed intermolecular difunctionalization of styrenes with arylboronic acids and methyl thiosulfonates has been developed. A variety of styrenes, methyl thiosulfonates and arylboronic acids were compatible with the catalytic conditions, resulting in rapid access to 2,2-diarylethylsulfone derivatives in a one-pot procedure. Key to success of this three component coupling hinges on the use of methyl thiosulfonates as sulfonyl radical source, which suppress the direct arylation of sulfonyl radical with arylboronic acids when reactions are conducted in acetone/DMF solvent systems. Importantly, owing to mild reaction conditions, our method is suited for the late-stage functionalization of complex natural products. The mechanistic studies point toward a sulfonyl radical initialized radical relay pathway. The development of enantioselective version of the strategy described herein is currently ongoing in our laboratory.

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

## AUTHOR INFORMATION

### Corresponding Author

\*E-mail: jiatz@sustech.edu.cn

## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENT

We thank the Science and Technology Innovation Commission of Shenzhen Municipality (JCYJ20180302180256215) to provide financial support. SUSTech is greatly appreciated for providing startup fund to T.J.. We are also very grateful to Dr. Yang Yu (SUSTech) for HRMS and Dr. Zhou Tang (SUSTech) for EPR fitting data. PJW thanks the US NSF (CHE-1902509).

## REFERENCES

- (1) (a) Trost, B. M. Chemical Chameleons. *Organosulfones as Synthetic Building Blocks*. *Bull. Chem. Soc. Jpn.* **1988**, *61*, 107-124. (b) Oida, S.; Tajima, Y.; Konosu, Y.; Nakamura, Y.; Somada, A.; Tanaka, T.; Habuki, S.; Harasaki, T.; Kamai, Y.; Fukuoka, T.; Ohya, S.; Tasuda, H. Synthesis and Antifungal Activities of R-102557 and Related Dioxane-Triazole Derivatives. *Chem. Pharm. Bull.* **2000**, *48*, 694-707. (c) Back, T. G. Design and synthesis of some biologically interesting natural and unnatural products based on organosulfur and selenium chemistry. *Can. J. Chem.* **2009**, *87*, 1657-1674.
- (2) (a) Eto, H.; Kaneko, Y.; Takeda, S.; Tokizawa, M.; Sato, S.; Yoshida, K.; Namiki, S.; Ogawa, M.; Maebashi, K.; Ishida, K.; Matsumoto, M.; Asaoka, T. New Antifungal 1,2,4-Triazoles with Difluoro(substituted sulfonyl)methyl Moiety. *Chem. Pharm. Bull.* **2001**, *49*, 173-182. (b) Fortugno, C.; Varchi, G.; Guerrini, A.; Carrupt, P.-A.; Bertucci, C. Optical biosensor analysis in studying new synthesized bicalutamide analogs binding to androgen receptor. *J. Pharm. Biomed. Anal.* **2014**, *95*, 151-157. (c) Shen, Y.; Zifcsak, C. A.; Shea, J. E.; Lao, X.; Bollt, O.; Li, X.; Lisko, J. G.; Therooff, J. P.; Scaife, C. L.; Ator, M. A.; Ruggeri, B. A.; Dorsey, B. D.; Kuwada, S. K. Design, Synthesis, and Biological Evaluation of Sulfonyl Acrylonitriles as Novel Inhibitors of Cancer Metastasis and Spread. *J. Med. Chem.* **2015**, *58*, 1140-1158.
- (3) Lee, M.; Ikejiri, M.; Klimpel, D.; Toth, M.; Espahbodi, M.; Heseck, D.; Forbes, C.; Kumarasiri, M.; Noll, B. C.; Chang, M.; Mobashery, S. Structure-Activity Relationship for Thiirane-Based Gelatinase Inhibitors. *ACS Med. Chem. Lett.* **2012**, *3*, 490-495.
- (4) Lopez de Compadre, R. L.; Pearlstein, R. A.; Hopfinger, A. J.; Seydel, J. K. A quantitative structure-activity relationship analysis of some 4-aminodiphenyl sulfone antibacterial agents using linear free energy and molecular modeling methods. *J. Med. Chem.* **1987**, *30*, 900-906.
- (5) Wang, J.; Hou, T. Drug and Drug Candidate Building Block Analysis. *J. Chem. Inf. Model.* **2010**, *50*, 55-67.
- (6) Ashcroft, C. P.; Hellier, P.; Pettman, A.; Watkinson, S. Second-Generation Process Research Towards Eletriptan: A Fischer Indole Approach. *Org. Process Res. Dev.* **2011**, *15*, 98-103.
- (7) Madasu, S. B.; Vekariya, N. A.; Kiran, M. N.; Gupta, B.; Islam, A.; Douglas, P. S.; Babu, K. R. Synthesis of compounds related to the anti-migraine drug eletriptan hydrobromide. *Beilstein J. Org. Chem.* **2012**, *8*, 1400-1405.
- (8) (a) Patai, S.; Rappoport, Z.; Stirling, C. *The Chemistry of Sulphones and Sulphoxides*. Wiley: Chichester, 1988. (b) Simpkins, N. S. *Sulphones in Organic Synthesis*. Pergamon Press: Oxford, 1993. (c) Magnus, P. D. Recent Developments in Sulfone Chemistry. *Tetrahedron* **1977**, *33*, 2019-2045.
- (9) (a) Julia, M.; Paris, J. M. Syntheses using Sulfones. 5. Method for General Synthesis of Doubles. *Tetrahedron Lett.* **1973**, *14*, 4833-4836. (b) Blakemore, P. R. The modified Julia olefination: alkene synthesis via the condensation of metallated heteroarylalkylsulfones with carbonyl compounds. *J. Chem. Soc., Perkin Trans. 1* **2002**, 2563-2585.
- (10) (a) Fang, Y.; Luo, Z.; Xu, X. M. Recent advances in the synthesis of vinyl sulfones. *RSC Adv.* **2016**, *6*, 59661-59676. (b) Liwosz, T. W.; Chemler, S. R. Copper-Catalyzed Oxidative Amination and Allylic Amination of Alkenes. *Chem. Eur. J.* **2013**, *19*, 12771-12777.
- (11) (a) Zhou, T.; Peters, B.; Maldonado, M. F.; Govender, T.; Andersson, P. G. Enantioselective Synthesis of Chiral Sulfones by Ir-Catalyzed Asymmetric Hydrogenation: A Facile Approach to the Preparation of Chiral Allylic and Homoallylic Compounds. *J. Am. Chem. Soc.* **2012**, *134*, 13592-13595. (b) Jiang, J.; Wang, Y.; Zhang, X. Rhodium-Catalyzed Asymmetric Hydrogenation of  $\beta$ -Acetylamino Acrylosulfones: A Practical Approach to Chiral  $\beta$ -Amido Sulfones. *ACS Catal.*

**2014**, 4, 1570-1573. (c) Peters, B. K.; Zhou, T.; Rujirawanich, J.; Cadu, A.; Singh, T.; Rabten, W.; Kerdphon, S.; Andersson, P. G. An Enantioselective Approach to the Preparation of Chiral Sulfones by Ir-Catalyzed Asymmetric Hydrogenation. *J. Am. Chem. Soc.* **2014**, 136, 16557-16562. (d) Shi, L.; Wei, B.; Yin, X.; Xue, P.; Lv, H.; Zhang, X. Rh-Catalyzed Asymmetric Hydrogenation of  $\alpha$ -Substituted Vinyl Sulfones: An Efficient Approach to Chiral Sulfones. *Org. Lett.* **2017**, 19, 1024-1027.

(12) Ranu, B. C.; Guchhait, S. K.; Ghosh, K. Chemoselective Hydrogenation of  $\alpha,\beta$ -Unsaturated Sulfones and Phosphonates via Palladium-Assisted Hydrogen Transfer by Ammonium Formate. *J. Org. Chem.* **1998**, 63, 5250-5251.

(13) Yan, Q.; Xiao, G.; Wang, Y.; Zi, G.; Zhang, Z.; Hou, G. Highly Efficient Enantioselective Synthesis of Chiral Sulfones by Rh-Catalyzed Asymmetric Hydrogenation. *J. Am. Chem. Soc.* **2019**, 141, 1749-1756.

(14) Lautens, M.; Mancuso, J.; Grover, H. Rhodium-Catalyzed Heck-Type Coupling of Boronic Acids with Activated Alkenes in an Aqueous Emulsion. *Synthesis* **2004**, 2006-2014.

(15) Nishimura, T.; Takiguchi, Y.; Hayashi, T. Effect of Chiral Diene Ligands in Rhodium-Catalyzed Asymmetric Addition of Arylboronic Acids to  $\alpha,\beta$ -Unsaturated Sulfonyl Compounds. *J. Am. Chem. Soc.* **2012**, 134, 9086-9089.

(16) (a) Evano, G.; Blanchard, N. *Copper-Mediated Cross-Coupling Reactions*. Wiley: New Jersey, 2014. (b) Shaughnessy, K. H.; Ciganek, E.; DeVasher, R. B. *Copper-Catalyzed Amination of Aryl and Alkenyl Electrophiles* Wiley: New Jersey, 2017. (c) Wang, F.; Chen, P.; Liu, G. Copper-Catalyzed Radical Relay for Asymmetric Radical Transformations. *Acc. Chem. Res.* **2018**, 51, 2036-2046.

(17) (a) Lan, X.-W.; Wang, N.-X.; Xing, Y. Recent Advances in Radical Difunctionalization of Simple Alkenes. *Eur. J. Org. Chem.* **2017**, 5821-5851. (b) Zhang, J.-S.; Liu, L.; Chen, T.; Chen, L.-B. Transition-Metal-Catalyzed Three-Component Difunctionalizations of Alkenes. *Chem. Asian J.* **2018**, 13, 2277-2291. (c) Lin, J.; Song, R.-J.; Hu, M.; Li, J.-H. Recent Advances in the Intermolecular Oxidative Difunctionalization of Alkenes. *Chem. Rec.* **2019**, 19, 440-451. (d) Fu, X.; Zhao, W. Progress in Difunctionalization of Alkenes. *Chin. J. Org. Chem.* **2019**, 39, 625-647.

(18) (a) Bertrand, M. P.; Ferreri, C. *Sulfur-Centered Radicals*. Wiley-VCH: Weinheim: 2001; Vol. 2. (b) Pan, X.-Q.; Zou, J.-P.; Yi, W.-B.; Zhang, W. Recent advances in sulfur- and phosphorous-centered radical reactions for the formation of S-C and P-C bonds. *Tetrahedron* **2015**, 71, 7481-7529. (c) Zheng, D.; Yu, J.; Wu, J. Generation of Sulfonyl Radicals from Aryldiazonium Tetrafluoroborates and Sulfur Dioxide: The Synthesis of 3-Sulfonated Coumarins. *Angew. Chem., Int. Ed.* **2016**, 55, 11925-11929.

(19) (a) Beaulieu, C.; Guay, D.; Wang, Z.; Evans, D. A. A mild and efficient new synthesis of aryl sulfones from boronic acids and sulfinic acid salts. *Tetrahedron Lett.* **2004**, 45, 3233-3236. (b) Huang, F.; Batey, R. A. Cross-coupling of organoboronic acids and sulfinate salts using catalytic copper(II) acetate and 1,10-phenanthroline: synthesis of aryl and alkenylsulfones. *Tetrahedron* **2007**, 63, 7667-7672. (c) Kar, A.; Sayyed, I. A.; Lo, W. F.; Kaiser, H. M.; Beller, M.; Tse, M. K. A General Copper-Catalyzed Sulfonylation of Arylboronic Acids. *Org. Lett.* **2007**, 9, 3405-3408.

(20) (a) Lu, Q. Q.; Zhang, J.; Wei, F. L.; Qi, Y.; Wang, H. M.; Liu, Z. L.; Lei, A. W. Aerobic Oxysulfonylation of Alkenes Leading to

Secondary and Tertiary beta-Hydroxysulfones. *Angew. Chem., Int. Ed.* **2013**, 52, 7156-7159. (b) Terent'ev, A. O.; Mulina, O. M.; Pargach, D. A.; Demchuk, D. V.; Syroeshkin, M. A.; Nikishin, G. I. Copper(I)-mediated synthesis of  $\beta$ -hydroxysulfones from styrenes and sulfonylhydrazides: an electrochemical mechanistic study. *RSC Adv.* **2016**, 6, 93476-93485. (c) Taniguchi, T.; Idota, A.; Ishibashi, H. Iron-catalyzed sulfonyl radical formations from sulfonylhydrazides and oxidative addition to alkenes. *Org. Biomol. Chem.* **2011**, 9, 3151-3153.

(21) (a) Wei, W.; Liu, C.; Yang, D.; Wen, J.; You, J.; Suo, Y.; Wang, H. Copper-catalyzed direct oxysulfonylation of alkenes with dioxygen and sulfonylhydrazides leading to beta-ketosulfones. *Chem. Commun.* **2013**, 49, 10239-10241. (b) Huang, S.; Thirupathi, N.; Tung, C. H.; Xu, Z. Copper-Catalyzed Oxidative Trifunctionalization of Olefins: An Access to Functionalized beta-Keto Thiosulfones. *J. Org. Chem.* **2018**, 83, 9449-9455.

(22) Li, X.-F.; Lin, J.-S.; Liu, X.-Y. Catalytic Radical Intramolecular Aminoperfluoroalkylation and Aminodifluoromethylation of Unactivated Alkenes with Fluoro-alkylsulfonyl Chlorides. *Synthesis* **2017**, 49, 4213-4220.

(23) Kumar, S.; Singh, R.; Singh, K. N. AIBN-Initiated Denitrative Cross-Coupling Reactions of  $\beta$ -Nitrostyrenes with Sulfonyl Hydrazides/Disulfides: A Metal-free Approach towards Vinyl Sulfones. *Asian J. Org. Chem.* **2018**, 7, 359-362.

(24) (a) Shyam, P. K.; Son, S.; Jang, H.-Y. Copper-Catalyzed Sulfonylation of Alkenes and Amines by Using Thiosulfonates as a Sulfonylating Agent. *Eur. J. Org. Chem.* **2017**, 5025-5031. (b) Son, S.; Shyam, P. K.; Park, H.; Jeong, I.; Jang, H.-Y. Complementary Strategy for Regioselective Synthesis of Diverse  $\beta$ -Hydroxysulfones from Thiosulfonates. *Eur. J. Org. Chem.* **2018**, 3365-3371.

(25) (a) Taylor, R. D.; MacCoss, M.; Lawson, A. D. G. Rings in Drugs. *J. Med. Chem.* **2014**, 57, 5845-5859. (b) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals. *J. Med. Chem.* **2014**, 57, 10257-10274.

(26) Tromp, M.; van Strijdonck, G. P. F.; van Berkel, S. S.; van den Hoogenband, A.; Feiters, M. C.; de Bruin, B.; Fiddy, S. G.; van der Eerden, A. M. J.; van Bokhoven, J. A.; van Leeuwen, P. W. N. M.; Koningsberger, D. C. Multitechnique Approach to Reveal the Mechanism of Copper(II)-Catalyzed Arylation Reactions. *Organometallics* **2010**, 29, 3085-3097.

(27) (a) Kharasch, M. S.; Sosnovsky, G.; Yang, N. C. Reactions of t-Butyl Peresters. 1. The Reaction of Peresters with Olefins. *J. Am. Chem. Soc.* **1959**, 81, 5819-5824. (b) Xiong, Y.; Sun, Y.; Zhang, G. Copper-Catalyzed Synthesis of  $\beta$ -Azido Sulfonates or Fluorinated Alkanes: Divergent Reactivity of Sodium Sulfonates. *Org. Lett.* **2018**, 20, 6250-6254.

(28) Xiang, J.-n.; Yuan, Y.-q.; Guo, S.-r. Cu(OAc)<sub>2</sub>-Catalyzed Thiolation of Acyl C-H Bonds with Thiols Using TBHP as an Oxidant. *Synlett* **2013**, 24, 443-448.

(29) Wang, F.; Wang, D.; Mu, X.; Chen, P.; Liu, G. Copper-Catalyzed Intermolecular Trifluoromethylarylation of Alkenes: Mutual Activation of Arylboronic Acid and CF<sub>3</sub><sup>+</sup> Reagent. *J. Am. Chem. Soc.* **2014**, 136, 10202-10205.

