



# Variation of key elements in soils and plant tissues in subalpine forests of the northern Rocky Mountains, U.S.A.

David P. Pompeani<sup>1</sup>, Kendra K. McLauchlan<sup>1</sup>, Barrie V. Chileen<sup>1</sup>, Kyra D. Wolf<sup>2</sup>, Philip E. Higuera<sup>2,3</sup>

<sup>1</sup>Department of Geography, Kansas State University, Manhattan, KS 66506

<sup>2</sup>Systems Ecology Intercollegiate Graduate Program, University of Montana, Missoula, MT 59812

<sup>3</sup>Department of Ecosystem and Conservation Sciences, University of Montana, Missoula, MT 59812

## Abstract

The essential elements for the structure and function of forest ecosystems are found in relatively predictable proportions in living tissues and soils; however, both the degree of spatial variability in elemental concentrations and their relationship with wildfire history are unclear. Quantifying the association between nutrient concentrations in living plant tissue and surface soils within fire-affected forests can help determine how these elements contribute to biogeochemical resilience. Here, we present elemental concentration data (C, N, P, K, Ca, Mg, S, Fe, Mn, Zn) from 72 foliar and 44 soil samples from a network of 15 sites located in the fire-prone subalpine forests of the northern Rocky Mountains, U.S.A. Plant functional type is strongly correlated with carbon (C) and nitrogen (N)—C concentrations are highest in coniferous needles, and N concentrations are highest in broadleaved plant species. The average N/P ratio of foliage among samples is  $9.8 \pm 0.6$  ( $\mu \pm 95\%$  confidence). This suggests that N is the limiting nutrient for these plants, however several factors can complicate the use of N/P ratios to evaluate nutrient status. Average C concentrations in organic soil horizons that were burned in regionally extensive fires in 1910 or 1918 CE are lower than those from sites that burned prior to 1901 CE ( $p < 0.05$ ). This difference suggests that wildfires reduced the pool of soil C and that the legacy of these fires can be measured a century later. Our results help aid in modeling how changing wildfire regimes will influence biogeochemical cycling in subalpine forests.

## 1. Introduction

Although living plant tissue is primarily composed of carbon (C), there are approximately 20 other elements that are necessary for biochemical reactions and growth. These other key elements—including nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), and sulfur (in the form of  $\text{SO}_4$ )—play important roles in



41 regulating terrestrial ecosystem processes. The quantities of these elements are controlled  
42 by a number of factors including bedrock parent material, soil composition (Kramer et  
43 al., 2017), vegetation type (Hu et al., 2001), and climate (Campbell et al., 2009). It is  
44 generally thought that plant tissue and soil nutrient concentrations can provide  
45 information about element limitation to growth (Wardle et al., 2004; Boerner,  
46 1984; Schreeg et al., 2014). However, relatively few studies have analyzed the  
47 concentrations of a large suite of elements in both plant and soil samples from one region.

48 Plant traits, such as biochemical, physiological, and anatomical features measured  
49 at the individual level (Violle et al., 2007), reflect the outcome of evolutionary processes  
50 responding to environmental constraints (Valladares et al., 2007). Traits determine how  
51 primary producers respond to abiotic and biotic environmental factors and influence a  
52 host of other ecosystem services (Kattge et al., 2011). Plant functional type (PFT) models  
53 capture a substantial fraction of the observed trait variation across plant species. A  
54 relatively small number of PFTs (5-20) have been used to represent the functional  
55 diversity of >300,000 documented plant species on Earth in global vegetation models  
56 (Kattge et al., 2011). Growth-limiting elements in foliar material derived from soils, such  
57 as phosphorus, make nutrient concentrations among the most plastic of plant traits  
58 (Wright et al., 2004) because the availability of these nutrients varies spatially as a result  
59 of parent bedrock composition. Nutrient concentrations in plants are influenced by a  
60 number of other factors, including the age of the tissue and overall age of the plant  
61 (Turner et al., 1977; Reich et al., 1992; Schreeg et al., 2014; Luo et al., 2017). Additionally,  
62 sunlight and location on a tree branch can also affect nutrient concentration in leaves  
63 (Schreeg et al., 2014), and weather-related factors such as precipitation and temperatures  
64 can affect nutrient levels on a seasonal basis (Turner et al., 1977). Therefore, samples of  
65 plant tissues taken over a short time period for several species across one region are  
66 particularly useful for differentiating factors that influence foliar nutrient concentrations  
67 and subsequent bioavailability of nutrients in leaf litter and upper soil horizons (Qualls  
68 and Haines, 1991; Taylor et al., 1989).

69 High severity fires can profoundly impact the cycling of C, N, and other nutrients  
70 in plants and soils (Chen et al., 2017; Fletcher et al., 2014; Certini, 2005; Dunnette et al.,  
71 2014), and could therefore affect the ability of the forest to regenerate to pre-fire  
72 conditions (Smithwick, 2011; McLauchlan et al., 2014). Several mechanisms determine  
73 how forests respond to fire, and how cycling of C, N, and other elements subsequently  
74 change (Pellegrini et al., 2018; Certini, 2005). For example, combustion of vegetation and  
75 soils from frequent, low-intensity burning can lead to the loss of C and N to the  
76 atmosphere (Pellegrini et al., 2014; Reich et al., 2001; Deluca and Sala, 2006; Yelenik et  
77 al., 2013). Fires also release nutrients from plant tissues as ash, which can potentially  
78 increase post-fire vegetation growth through soil nutrient enrichment (Boerner et al.,  
79 2009; Hurteau and Brooks, 2011). Over longer time scales of centuries to millennia,  
80 ecosystem modeling informed with paleofire history records suggest that changing fire



regimes can have substantial and long-lasting impacts on C and nutrient cycling (Hudiburg et al., 2017; Kelly et al., 2016).

Here we provide data from 72 leaf and 44 soil samples from a network of 15 sites in the coniferous subalpine forests of the northern Rocky Mountains, U.S.A. (Fig. 1, Table 1). The samples were collected during the middle of the growing season from sites with similar elevation, climate, and bedrock geology, but with varying fire histories over the 20<sup>th</sup> century. Our approach isolates local plant and soil forming processes that affect the distribution of key elements, and compares sites that are reported to have burned in extensive fires in 1910 and 1918 CE to those that have not burned since before 1901 CE (Morgan et al., 2014). We evaluate nutrient status and characterize patterns of element distribution within and among the leaves of several tree and shrub species in the context of associated bioavailable nutrient data from upper soil horizons. We use statistical methods to infer the transfer and distribution of key elements in soil and vegetation.

## 2. Methods

### 2.1 Study area

The study sites are located between 1623–2011 m above sea level in the northern Bitterroot Mountains along the border of Idaho and Montana, U.S.A. (Fig. 1). The vegetation at all sites is subalpine forest dominated by subalpine fir (*Abies lasiocarpa*), Engelmann spruce (*Picea engelmannii*), lodgepole pine (*Pinus contorta*), and mountain hemlock (*Tsuga mertensiana*). Of the 14 sites, seven experienced wildfire in 1910 or 1918 CE, while the other seven have no indication of fire activity back to 1901 CE (Morgan et al., 2014) (Table 1). The regional climate is classified as modified maritime with warm, dry summers and cool, wet winters. The study region is underlain with partially metamorphosed argillites of the Belt Basin Supergroup, known as Belt meta-sedimentary rocks. Glacial till and shallow entisols overlay the bedrock (Sasich and Lamotte-Hagen, 1989).

### 2.2 Fieldwork

Soil cores, approximately 2 cm in diameter and 25 cm deep, and foliage samples were collected at 15 sites along a 100-km northwest to southeast transect (46.78°–47.36° N; 114.76°–115.57° W) from 12 July to 25 August, 2017 CE (Fig. 1). At each soil core site, except for Hoodoo, associated leaf samples were collected from nearby trees and shrubs, avoiding new growth and juvenile trees. The majority of the foliar samples consist of coniferous needles from eight species (n=68): *Abies lasiocarpa*, *Larix occidentalis*, *Pinus albicaulis*, *Pinus contorta*, *Picea engelmannii*, *Taxus brevifolia*, *Thuja plicata*, and *Tsuga mertensiana*. We also collected two samples each of the



broadleaf species green alder (*Alnus viridis*; n=2) and false huckleberry (*Menziesia ferruginea*; n=2). Foliar samples were taken directly from living vegetation located approximately 2 m above the soil surface. Soil cores were sampled by horizons: ‘litter’ was unconsolidated plant material from the surface; ‘organic’ soil was identified visually and sampled at a depth between approximately 0–10 cm; the ‘mineral’ soil layer was sampled between 10–25 cm below the surface depending on overall soil depth. Samples were stored in airtight plastic containers for transport back to the laboratory.

### 2.3 Laboratory analyses

Samples were dried, homogenized, and sent to the Department of Agronomy Soil Testing Lab at Kansas State University. Foliar samples were weighed into 50 ml Kimax digestion tubes for elemental analysis. Boiling chips and 5 ml of nitric acid at 50% strength were added to the tubes, which were covered with plastic wrap to react overnight. The following day, 5 ml of perchloric acid was added to predigested plant material and tubes were then placed on a cold techator digestion block. Temperatures were set to 200°C and digests were heated for approximately three hours or until white fumes appeared and the acid was clear (and colorless when cooled to room temperature) (Gieseking et al., 1935). Tubes were diluted to 25 or 50 ml with deionized water and mixed by inverting twice. Each digested sub-sample was then analyzed for P, Mg, K, Ca, iron (Fe), manganese (Mn), zinc (Zn), and SO<sub>4</sub> using a Varian Model 720-ES ICP Optical Emission Spectrometer. Analytical precision ( $\sigma$ ) for element analysis was <0.0001%.

Base cations (Ca, Mg, K, and Na) were extracted from soil sub-samples using 1 M ammonium acetate (pH 7.0). The resulting supernatant was then analyzed using a Varian Model 720-ES ICP Optical Emission Spectrometer. Trace elements (e.g. Mn, Zn, etc.) were not measured on soil sub-samples. Soils were not analyzed for P.

Total carbon and nitrogen concentrations (weight %) in both plant and soil sub-samples were measured using a LECO CN 2000 combustion analyzer. Analytical precision ( $\sigma$ ) was approximately 0.06% for carbon and 0.006% for nitrogen.

### 2.4 Statistics

To compare elemental concentrations among different sample sites, we used box plots, unpaired t-tests (assuming unequal variance), principal components analysis (PCA), and Pearson correlation coefficients. We calculated principal components using the ade4 package in R (Dray and Dufour, 2007). Correlation tables were calculated using the R base stats package (R Core Team).

## 3. Results and discussion



### 3.1 Patterns in soil and foliar elements

Soil C and N concentrations were highest on average (34.94% and 1.32%, respectively) within litter material and declined with depth through organic (18.30% and 0.84%) and mineral (6.05% and 0.29%) soil horizons (Table S1, Fig. 2). The concentration of other nutrients such as K, Ca, and Mg were highest on average within the litter (0.016%, 0.115%, and 0.012%, respectively) and organic soil horizons (0.019%, 0.104%, and 0.013%), and decreased in the mineral soil horizon (0.009%, 0.046%, 0.007%). Na concentrations displayed the opposite pattern and increased from litter (<0.001%) to mineral (0.001%) soil horizon.

The pattern of five elements in the soil—C, N, Ca, K, and Mg—correspond with the dominant role of decomposing plant material controlling the concentration of nutrients in forested soils (Aerts and Chapin, 1999). These results are consistent with a global compilation of 10,000 soil profiles that indicate ranking of nutrient concentrations from shallow to deep (Jobbagy and Jackson, 2001) (Fig. 2, Fig. S1). The decline in the bioavailable K, Ca, and Mg in the mineral horizon suggests removal by mineralization or microbial immobilization (Qualls et al., 1991; Qualls, 2000).

Foliage contained C concentrations between 47–52%, with the exception of *Larix occidentalis*, *Alnus viridis*, and *Menziesia ferruginea*, which had slightly lower concentrations of C between 43–47% (Fig. 3, Table S2). N concentrations were generally low (0.75–1.89%) in coniferous needles, and high in leaves of the broadleaved species *Alnus viridis* and *Menziesia ferruginea* (2.5–3.5%). P, K, and Ca concentrations were variable within and between tree and shrub species, with no clear pattern differentiating needle and broadleaf species (Fig. 3).

There was considerable variability in trace element concentrations among samples of foliar material (Fig. 3). Mg was below 0.20% in all leaf material, except for *Alnus viridis* (0.35%) and *Menziesia ferruginea* (0.50%). Fe, Mn, and Zn concentrations were highly variable within and between species. *Menziesia ferruginea* contained high concentrations of Mn (0.48%) and Zn (0.01%). SO<sub>4</sub> concentrations ranged between 0.01–0.15% for all foliar material, except for higher values found in *Alnus viridis* (0.18%).

Element concentrations did not vary with site latitude or elevation likely as a result of the constricted latitude and elevation range of the study sites. Soil and foliar N, K, and Mg were all positively correlated ( $p < 0.001$ ) with each other ( $r = 0.42$ – $0.75$ ) (Figs. S1, S2, S3, and S4). In soils, Ca was positively correlated ( $p < 0.01$ ) with Mg (0.74), K (0.54), N (0.47), and C (0.39), whereas in plant tissues Ca was poorly correlated ( $p > 0.05$ ) with P (−0.21), K (−0.21), and Mg (0.15), C (0.05), and N (0.02). Foliar C was negatively correlated ( $p < 0.05$ ) with Mg (−0.53), N (−0.41), P (−0.38), and K (−0.30). In soils, Na was inversely related ( $p < 0.05$ ) to both C (−0.52) and N (−0.42). In vegetation, Zn was correlated ( $p < 0.05$ ) with Mn (0.54), K (0.33), and C (−0.26). Trace elements Fe and Mn were weakly correlated ( $p > 0.05$ ) with C, N, P, K, and Mg in plant tissues.



A multivariate analysis of elemental concentrations clearly separated foliar ( $n=72$ ) from litter/soil samples ( $n=44$ ) (principal component analysis [PCA]; Fig. 4). Axis 1 of the PCA explained 58.7% of total variation in the soil and foliar elements. The concentrations of several elements were negatively loaded on Axis 1: C (-0.905), K (-0.809), Ca (-0.791), Mg (-0.799), and N (-0.706). C/N was correlated to Axis 1 (-0.535) but more weakly than C, N, and the base nutrients. Axis 2 explained 22.4% of the variation and was primarily driven by N and Mg (0.577, 0.381, respectively), and was strongly influenced by the high foliar N concentrations of *Alnus viridis* ( $n=2$ ). Foliage samples from different plant species were dispersed along Axis 2.

Leaf carbon, nitrogen, and phosphorus content are considered important biochemical traits in PFT modeling, as seen in the TRY plant database (Kattge et al., 2011). We measured additional elemental concentrations to understand inter- and intra-species variation in biogeochemical traits. In live vegetation, N, P, K, Mg, and  $\text{SO}_4$  are all positively correlated (Fig. S2), indicating that leaf tissues are built in predictable stoichiometric ratios, but the variation in the concentration of those nutrients among sampling sites suggests that the availability of these elements in soils and bedrock likewise varied. Ca was weakly correlated with N, P, K, Mg, and  $\text{SO}_4$  in leaves. Ca concentration is strongly affected by foliage age (Turner et al., 1977), which represents another source of variation in the sampled foliage not controlled for in this study.

We found substantial differences in the foliar nutrient concentrations between evergreen and deciduous plant types (Fig. 3). N, Mg, and  $\text{SO}_4$  were found at higher concentrations in broadleaf material relative to evergreen needles, probably due to higher rates of metabolic activity and photosynthesis in broadleaf material during the summer (Linzon et al., 1979). Leaves from *Alnus*, a nitrogen-fixing plant, contained greater N (2.55–3.56%) relative to non-fixing species (Taylor et al., 1989). Leaf N concentrations ranged from 0.75 to 1.89% in the evergreen conifer species (i.e., excluding *Larix occidentalis*) ( $n=63$ ) —which is slightly lower but consistent with ( $1.15 \pm 0.24\%$ ) a previously reported global average of  $1.21 \pm 0.01\%$  N for needleleaf evergreens ( $n=5558$ ) (Kattge et al., 2011). These data suggest that nutrient variation between plant functional types (i.e. evergreen needleleaf versus deciduous) is much greater than inter- or intra-specific variation within PFTs. This supports the use of broadly-defined PFTs in biogeochemical models of ecosystem processes.

### 3.2 Fire legacies on nutrient availability

We compared average elemental concentrations (i.e. C, N, Ca, K, and Mg) in organic soil horizons taken from sample sites burned in 1910 or 1918 CE with sites that were burned prior to 1901 CE (i.e., the start of the Fire Atlas published by Morgan et al. (2014)) (Fig. 1, Table 1) to investigate the long-term effects of wildfires on soil nutrient availability. Soil samples from the organic horizon that were burned in the early 20<sup>th</sup>



241 century contained significantly less C on average than samples from sites that burned  
242 prior to 1901 CE ( $t = -2.26$ ,  $df = 9.60$ ,  $p = 0.0484$ ). In contrast, N ( $t = -1.46$ ,  $df = 10.54$ ,  $p =$   
243  $0.1731$ ) and other soil nutrient concentrations (Fig. 5; Ca, K, Mg not shown) did not  
244 significantly differ between these two populations.

245 The significant difference in soil C suggests that a single fire event in the early  
246 20<sup>th</sup> century at several sites reduced the pool of soil C, and that the legacies of those fires  
247 are still detectable today (Fig. 5). These results are consistent with ecosystem modeling  
248 experiments, which find that both fire frequency and severity are the dominant drivers of  
249 C dynamics in sub-alpine coniferous forests over centennial to millennial timescales  
250 (Hudiburg et al., 2017). Furthermore, the greater variance in C concentrations observed in  
251 soils burned in the early 20<sup>th</sup> century ( $\sigma^2 = 37.31$ ) versus soils not burned ( $\sigma^2 = 12.44$ ) (Fig.  
252 5) may also reflect aspects of fire history, and specifically high spatial variability in fire  
253 severity and post-fire recovery.

254 The results from this study are broadly consistent with ecosystem model  
255 simulations suggesting that changing fire regimes under climate change scenarios have  
256 the potential to alter C stored in forested ecosystems by changing the frequency and  
257 intensity of wildfire events (Hudiburg et al., 2017; Kelly et al., 2016). We show that even  
258 in a relatively small region of subalpine forests in the Rocky Mountains, an assumption  
259 of a single value for C stocks would not hold, and are instead highly dependent on a  
260 spatially heterogeneous fire history extending back for at least a century. The importance  
261 of this variability in determining post-fire C dynamics implies that equilibrium scenarios  
262 extrapolated from a single fire event in one location are a poor assumption when  
263 simulating fire regimes in Earth System models at spatial scales larger than an individual  
264 site (Hudiburg et al., 2017).

### 266 3.3 Evaluating nutrient limitations

268 The process whereby forest vegetation typically returns to pre-fire conditions after  
269 several years or decades following wildfire events is referred to as biogeochemical  
270 resilience (McLauchlan et al., 2014; Smithwick, 2011). This resilience is determined, in  
271 part, by the availability of growth limiting nutrients, such as N and P (Güsewell,  
272 2004; Schreeg et al., 2014). In non-N<sub>2</sub>-fixing plant species, foliar N and P concentrations  
273 reflect soil N and P availability (Reich and Oleksyn, 2004). Previous studies have shown  
274 that a N/P mass ratio in foliar material of about 10–20 is optimal for plant growth (Aerts  
275 and Chapin, 1999; Ingestad and Lund, 1979). N/P ratios >16 can indicate P-limited  
276 biomass productions, whereas N/P ratios <14 are suggestive of N-limited plant growth  
277 (Aerts and Chapin, 1999; Koerselman and Meuleman, 1996).

278 The average ( $\pm$  95% confidence intervals) N/P values for sampled foliage ( $n = 72$ )  
279 was  $9.8 \pm 0.6$ . This suggests a N-limited growing environment. Although N/P ratios of  
280 leaves are often used to evaluate nutrient status in plants, several factors could complicate



281 this interpretation: (1) N/P ratios of old leaves may be different than new foliage growth  
282 (Schreeg et al., 2014). When plants are older, nutrients are reallocated to active  
283 meristems (e.g., young leaves, shoot tips) (Güsewell, 2004). We did not account for  
284 foliage or plant age during sampling. (2) All foliage was recovered in the summer,  
285 therefore N/P ratios could be low because samples were taken during the period of active  
286 growth (Méndez and Karlsson, 2005; Rivas-Ubach et al., 2012). (3) As a result of N  
287 limitation, biomass allocation to roots can increase at the expense of foliage (Andrews et  
288 al., 1999; De Groot et al., 2003). (4) P uptake can be enhanced in response to deficiencies  
289 through several mechanisms, such as exudation of enzymes or acids (Dakora and Phillips,  
290 2002) and association with mycorrhizal fungi (Colpaert et al., 1999). (5) Productivity  
291 might also be limited by other elements (van Duren and Peggel, 2000), solar irradiance,  
292 and/or climatic factors (Spink et al., 1998).

293 To assess potential nutrient limitations resulting from wildfire, we compared  
294 average N and P concentrations in foliar material from plants grown in soils that were  
295 burned in 1910 and 1918 CE ( $n=29$ ) with sites not burned ( $n=43$ ) since the start of the  
296 Fire Atlas (Morgan et al., 2014). We found no significant difference in average N ( $t=-$   
297  $0.13$ ,  $df=47.76$ ,  $p=0.8948$ ) or P ( $t=0.99$ ,  $df=53.48$ ,  $p=0.3258$ ) concentrations between  
298 the two sample groups. Likewise, average C ( $t=-0.40$ ,  $df=65.96$ ,  $p=0.6901$ )  
299 concentrations did not significantly differ between foliar samples taken from burned  
300 versus unburned locations. This indicates that, although past wildfires reduced the pool of  
301 soil C, they did not affect the concentration of growth limiting nutrients measured in  
302 vegetation a century later.

#### 305 4. Conclusions

307 Analyses of foliar and soil samples from a network of sites located in the northern  
308 U.S. Rocky Mountains indicate the spatial distribution of key elements in subalpine  
309 forested ecosystems. The concentration of C and nutrients (N, K, Mg, Ca) in soils is  
310 highest in the upper litter and organic horizons and decreases at depth in the mineral soil,  
311 consistent with previous studies. Comparing the two plant functional types, needle leaf  
312 plants contain higher concentrations of C, while broadleaf material is enriched in N and  
313 other trace elements (Mg,  $SO_4$ ). Sites that were burned in a regionally extensive wildfire  
314 in the early 20<sup>th</sup> century contained significantly lower C on average in the organic soil  
315 horizon, compared with sites that burned prior to the 20<sup>th</sup> century. This highlights the  
316 important role of wildfires as a dominant driver of soil C dynamics in sub-alpine forests,  
317 with legacies that can last for more than a century (Hudiburg et al., 2017). Furthermore,  
318 the high degree of variance in soil C concentrations among burned soils is consistent with  
319 the inherent spatial heterogeneity in fire severity seen in contemporary fires. This spatial  
320 heterogeneity adds to additional complexity Earth Systems Modeling efforts to represent  
321 fire across space and time. The low average values of foliar N/P ratios ( $9.8 \pm 0.6$ ) suggest



that N may be in low supply to plants. Therefore N availability in soils may play an important role in understanding the biogeochemical resilience of coniferous forests to wildfires. Finally, these data contribute empirical data to efforts to model the biogeochemical consequences of wildfires in the subalpine forests of western North America.

#### *Data Availability*

Foliar data are available in the TRY Plant Trait Database.

#### *Supplemental information*

Supplementary information is available online.

#### *Author contribution*

DPP and KKM designed the study, analyzed the data, and prepared the paper with contributions from BC, KW, and PEH. KW and PEH selected study sites. KW, PEH, and KKM collected the samples. BC conducted laboratory analyses to acquire data and assisted with data analysis.

#### *Competing interests*

The authors declare that they have no conflict of interest.

#### *Acknowledgments*

This work was supported by NSF award DEB-1655179 to KKM, and DEB-1655121 to PEH. DPP was supported by NSF DEB-1145815. K. Lowe, J. Roozeboom and A. Marcotte provided analytical assistance. We thank other Big Burns team members T. Hudiburg, K. Bartowitz, and B. Shuman for field assistance and valuable discussions.




365 Table 1. Study sites

| Site name      | Elevation m asl | Latitude | Longitude | Year of last major fire (CE) <sup>1</sup> |
|----------------|-----------------|----------|-----------|-------------------------------------------|
| Cliff          | 1810            | 47.1390  | -115.1919 | 1910                                      |
| Heart          | 1682            | 47.2856  | -115.4529 | unknown                                   |
| Hoodoo         | 1817            | 46.9789  | -115.0012 | unknown                                   |
| Hub            | 1819            | 47.2756  | -115.3757 | unknown                                   |
| Kid            | 1909            | 46.7782  | -114.8146 | 1910                                      |
| Little Montana | 2011            | 46.8151  | -114.7652 | unknown                                   |
| Lost Lake      | 1840            | 47.1047  | -115.1274 | 1910                                      |
| Lower Bonanza  | 1922            | 47.0916  | -115.1330 | 1910                                      |
| Lower Oregon   | 1811            | 47.0554  | -115.0913 | unknown                                   |
| Missoula       | 1807            | 47.0671  | -115.1162 | 1910                                      |
| Moore          | 1629            | 47.1818  | -115.2521 | unknown                                   |
| Silver         | 1623            | 47.3596  | -115.5659 | 1918                                      |
| Surveyor       | 1831            | 46.8241  | -114.7585 | unknown                                   |
| Upper Bonanza  | 1922            | 47.0916  | -115.1330 | 1910                                      |
| Upper Oregon   | 1811            | 47.0554  | -115.0913 | unknown                                   |

<sup>1</sup>This coverage includes fire perimeters recorded from 1901-2008 CE (Morgan et al., 2014).



370



371

372

373

374

Fig. 1) (Bottom) Map of the study area with 1910 CE fire extent polygons (Morgan et al., 2014). (Upper right) Regional map of the study region.




Fig. 2) Boxplots displaying element concentrations in soil samples.





Fig. 3) Boxplots displaying element concentrations in foliar samples. Subtypes are *Abies lasiocarpa* (ABLA), *Alnus viridis* (ALVI), *Larix occidentalis* (LAOC), *Menziesia*



381 *ferruginea* (MEFE), *Pinus albicaulis* (PIAL), *Pinus contorta* (PICO), *Picea engelmannii*  
 382 (PIEN), *Taxus brevifolia* (TABR), *Thuja plicata* (THPL), and *Tsuga mertensiana*  
 383 (TSME).  
 384



385  
 386 Fig. 4) Principal component analysis of soil and vegetation element concentrations.  
 387




Fig. 5) Boxplots of C and N from the organic soil horizon burned ( $n=7$ ) in 1910 CE and 1918 CE versus sites not burned ( $n=7$ ) during the historical record (Morgan et al., 2014). An unpaired t-test indicates that soil C concentrations are significantly ( $p=0.0484$ ) lower in burned sites.

## References

- Aerts, R., and Chapin, F. S.: The Mineral Nutrition of Wild Plants Revisited: A Re-evaluation of Processes and Patterns, *Adv. Ecol. Res.*, 30, 1-67, [https://doi.org/10.1016/s0065-2504\(08\)60016-1](https://doi.org/10.1016/s0065-2504(08)60016-1), 1999.
- Andrews, M., Sprent, J. I., Raven, J. A., and Eady, P. E.: Relationships between shoot to root ratio, growth and leaf soluble protein concentration of *Pisum sativum*, *Phaseolus vulgaris*, and *Triticum aestivum* under different nutrient deficiencies, *Plant, Cell & Environment*, 22, 949-958, <https://doi.org/10.1046/j.1365-3040.1999.00452.x>, 1999.
- Boerner, R. E., Huang, J., and Hart, S. C.: Impacts of fire and fire surrogate treatments on forest soil properties: a meta-analytical approach, *Ecol. Appl.*, 19, 338-358, <https://doi.org/10.1890/07-1767.1>, 2009.
- Boerner, R. E. J.: Foliar nutrient dynamics and nutrient use efficiency of four deciduous tree species in relation to site fertility, *J. Appl. Ecol.*, 21, 1029-1040, <https://doi.org/10.2307/2405065>, 1984.
- Campbell, J. L., Rustad, L. E., Boyer, E. W., Christopher, S. F., Driscoll, C. T., Fernandez, I. J., Groffman, P. M., Houle, D., Kiebusch, J., Magill, A. H., Mitchell, M. J., and Ollinger, S. V.: Consequences of climate change for biogeochemical cycling in forests of northeastern North America. This article is one of a selection of papers from NE Forests 2100: A Synthesis of Climate Change Impacts on Forests of the Northeastern US and Eastern Canada, *Can. J. Forest Res.*, 39, 264-284, <https://doi.org/10.1139/x08-104>, 2009.
- Certini, G.: Effects of fire on properties of forest soils: a review, *Oecologia*, 143, 1-10, <https://doi.org/10.1007/s00442-004-1788-8>, 2005.



- 420 Chen, G., Hayes, D. J., and David McGuire, A.: Contributions of wildland fire to  
421 terrestrial ecosystem carbon dynamics in North America from 1990 to 2012, *Global*  
422 *Biogeochem. Cy.*, 31, 878-900, <https://doi.org/10.1002/2016gb005548>, 2017.
- 423 Colpaert, J. V., Van Tichelen, K. K., Van Assche, J. A., and Van Laere, A.: Short-term  
424 phosphorus uptake rates in mycorrhizal and non-mycorrhizal roots of intact *Pinus*  
425 *sylvestris* seedlings, *New Phytol.*, 143, 589-597, <https://doi.org/10.1046/j.1469-8137.1999.00471.x>, 1999.
- 426 Dakora, F. D., and Phillips, D. A.: Root exudates as mediators of mineral acquisition in  
427 low-nutrient environments, *Plant Soil*, 245, 35-47,  
428 <https://doi.org/10.1023/A:1020809400075>, 2002.
- 429 De Groot, C., Marcelis, L. F. M., van den Boogard, R., Kaiser, W. M., and Lambers, H.:  
430 Interactions of nitrogen and phosphorus nutrition in determining growth, *Plant Soil*, 248,  
431 257-268, <https://doi.org/10.1023/A:1022323215010>, 2003.
- 432 Deluca, T. H., and Sala, A.: Frequent fire alters nitrogen transformations in ponderosa  
433 pine stands of the inland northwest, *Ecology*, 87, 2511-2522,  
434 [https://doi.org/10.1890/0012-9658\(2006\)87\[2511:FFANTI\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[2511:FFANTI]2.0.CO;2), 2006.
- 435 Dray, S., and Dufour, A. B.: The ade4 package: implementing the duality diagram for  
436 ecologists, *J. Stat. Softw.*, 22, <https://doi.org/10.18637/jss.v022.i04>, 2007.
- 437 Dunnette, P. V., Higuera, P. E., McLauchlan, K. K., Derr, K. M., Briles, C. E., and  
438 Keefe, M. H.: Biogeochemical impacts of wildfires over four millennia in a Rocky  
439 Mountain subalpine watershed, *New Phytol.*, 203, <https://doi.org/900-912>,  
440 10.1111/nph.12828, 2014.
- 441 Fletcher, M.-S., Wolfe, B. B., Whitlock, C., Pompeani, D. P., Heijnis, H., Haberle, S. G.,  
442 Gadd, P. S., Bowman, D. M. J. S., and Linder, P.: The legacy of mid-Holocene fire on a  
443 Tasmanian montane landscape, *J. Biogeogr.*, 41, 476-488,  
444 <https://doi.org/10.1111/jbi.12229>, 2014.
- 445 Gieseking, J. E., Snider, H. J., and Getz, C. A.: Destruction of organic matter in plant  
446 material by the use of nitric and perchloric acids, *Industrial and Engineering Chemistry*  
447 *Analytical Edition*, 7, 185-186, <https://doi.org/10.1021/ac50095a021>, 1935.
- 448 Güsewell, S.: N : P ratios in terrestrial plants: variation and functional significance, *New*  
449 *Phytol.*, 164, 243-266, <https://doi.org/10.1111/j.1469-8137.2004.01192.x>, 2004.
- 450 Hu, F. S., Finney, B. P., and Brubaker, L. B.: Effects of Holocene *Alnus* Expansion on  
451 Aquatic Productivity, Nitrogen Cycling, and Soil Development in Southwestern Alaska,  
452 *Ecosystems*, 4, 358-368, <https://doi.org/10.1007/s10021-001-0017-0>, 2001.
- 453 Hudiburg, T. W., Higuera, P. E., and Hicke, J. A.: Fire-regime variability impacts forest  
454 carbon dynamics for centuries to millennia, *Biogeosciences*, 14, 3873-3882,  
455 <https://doi.org/10.5194/bg-14-3873-2017>, 2017.
- 456 Hurteau, M. D., and Brooks, M. L.: Short- and long-term effects of fire on carbon in US  
457 dry temperate forest systems, *Biosciences*, 6, 139-146,  
458 <https://doi.org/10.1525/bio.2011.61.2.9>, 2011.
- 459 Ingestad, T., Lund, A.B.: Nitrogen stress in birch seedlings. II. N, K, P, Ca, and Mg  
460 nutrition, *Physiol. Plantarum*, 45, 149-157, <https://doi.org/10.1111/j.1399-3054.1979.tb01678.x>, 1979.
- 461 Jobbagy, E. G., and Jackson, R. B.: The distribution of soil nutrients with depth: Global  
462 patterns and the imprint of plants, *Biogeochemistry*, 53, 51-77,  
463 <https://doi.org/10.1023/A:1010760720215>, 2001.



- 466 Kattge, J., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., BöNisch, G., Garnier, E.,  
467 Westoby, M., Reich, P. B., Wright, I. J., Cornelissen, J. H. C., Violle, C., Harrison, S. P.,  
468 Van Bodegom, P. M., Reichstein, M., Enquist, B. J., Soudzilovskaia, N. A., Ackerly, D.  
469 D., Anand, M., Atkin, O., Bahn, M., Baker, T. R., Baldocchi, D., Bekker, R., Blanco, C.  
470 C., Blonder, B., Bond, W. J., Bradstock, R., Bunker, D. E., Casanoves, F., Cavender-  
471 Bares, J., Chambers, J. Q., Chapin Iii, F. S., Chave, J., Coomes, D., Cornwell, W. K.,  
472 Craine, J. M., Dobrin, B. H., Duarte, L., Durka, W., Elser, J., Esser, G., Estiarte, M.,  
473 Fagan, W. F., Fang, J., Fernández-MÉNdez, F., Fidelis, A., Finegan, B., Flores, O.,  
474 Ford, H., Frank, D., Freschet, G. T., Fyllas, N. M., Gallagher, R. V., Green, W. A.,  
475 Gutierrez, A. G., Hickler, T., Higgins, S. I., Hodgson, J. G., Jalili, A., Jansen, S., Joly, C.  
476 A., Kerkhoff, A. J., Kirkup, D., Kitajima, K., Kleyer, M., Klotz, S., Knops, J. M. H.,  
477 Kramer, K., KÜHn, I., Kurokawa, H., Laughlin, D., Lee, T. D., Leishman, M., Lens, F.,  
478 Lenz, T., Lewis, S. L., Lloyd, J., LlusiÀ, J., Louault, F., Ma, S., Mahecha, M. D.,  
479 Manning, P., Massad, T., Medlyn, B. E., Messier, J., Moles, A. T., MÜLLer, S. C.,  
480 Nadrowski, K., Naeem, S., Niinemets, Ü., NÖLlert, S., NÜSke, A., Ogaya, R., Oleksyn,  
481 J., Onipchenko, V. G., Onoda, Y., OrdoÑEz, J., Overbeck, G., Ozinga, W. A., PatiÑO,  
482 S., Paula, S., Pausas, J. G., PeÑUelas, J., Phillips, O. L., Pillar, V., Poorter, H., Poorter,  
483 L., Poschlod, P., Prinzing, A., Proulx, R., Rammig, A., Reinsch, S., Reu, B., Sack, L.,  
484 Salgado-Negret, B., Sardans, J., Shiodera, S., Shipley, B., Siefert, A., Sosinski, E.,  
485 Soussana, J. F., Swaine, E., Swenson, N., Thompson, K., Thornton, P., Waldram, M.,  
486 Weiher, E., White, M., White, S., Wright, S. J., Yguel, B., Zaehle, S., Zanne, A. E., and  
487 Wirth, C.: TRY - a global database of plant traits, *Glob. Change Biol.*, 17, 2905-2935,  
488 <https://doi.org/10.1111/j.1365-2486.2011.02451.x>, 2011.  
489 Kelly, R., Genet, H., McGuire, A. D., and Hu, F. S.: Palaeodata-informed modelling of  
490 large carbon losses from recent burning of boreal forests, *Nat. Clim. Change*, 6, 79-84,  
491 <https://doi.org/10.1038/nclimate2832>, 2016.  
492 Koerselman, W., and Meuleman, A. F. M.: The vegetation N:P ratio: a new tool to detect  
493 the nature of nutrient limitation, *J. Appl. Ecol.*, 33, 1441-1450,  
494 <https://doi.org/10.2307/2404783>, 1996.  
495 Kramer, M. G., Lajtha, K., and Aufdenkampe, A. K.: Depth trends of soil organic matter  
496 C:N and  $^{15}\text{N}$  natural abundance controlled by association with minerals,  
497 *Biogeochemistry*, 136, 237-248, <https://doi.org/10.1007/s10533-017-0378-x>, 2017.  
498 Linzon, S. N., Temple, P. J., and Pearson, R. G.: Sulfur Concentrations in Plant Foliage  
499 and Related Effects, *J. Air Waste Ma. Journal of the Air Pollution Control Association*,  
500 29, 520-525, <https://doi.org/10.1080/00022470.1979.10470822>, 1979.  
501 Luo, Z., Guan, H., Zhang, X., and Liu, N.: Photosynthetic capacity of senescent leaves  
502 for a subtropical broadleaf deciduous tree species *Liquidambar formosana* Hance, *Sci.*  
503 *Rep.*, 7, 6323, <https://doi.org/10.1038/s41598-017-06629-7>, 2017.  
504 McLauchlan, K. K., Higuera, P. E., Gavin, D. G., Perakis, S. S., Mack, M. C., Alexander,  
505 H., Battles, J., Biondi, F., Buma, B., Colombaroli, D., Enders, S. K., Engstrom, D. R.,  
506 Hu, F. S., Marlon, J. R., Marshall, J., McGlone, M., Morris, J. L., Nave, L. E., Shuman,  
507 B., Smithwick, E. A. H., Urrego, D. H., Wardle, D. A., Williams, C. J., and Williams, J.  
508 J.: Reconstructing Disturbances and Their Biogeochemical Consequences over Multiple  
509 Timescales, *BioScience*, 64, 105-116, <https://doi.org/10.1093/biosci/bit017>, 2014.



- 510 Méndez, M., and Karlsson, P. S.: Nutrient stoichiometry in *Pinguicula vulgaris*: Nutrient  
511 availability, plant size, and reproductive status, *Ecology*, 86, 982-991,  
512 <https://doi.org/10.1890/04-0354>, 2005.
- 513 Morgan, P., Heyerdahl, E. K., Miller, C., Wilson, A. M., and Gibson, C. E.: Northern  
514 Rockies pyrogeography: An example of fire atlas utility, *Fire Ecol.*, 10, 1-13,  
515 <https://doi.org/10.4996/fireecology.1001001>, 2014.
- 516 Pellegrini, A. F. A., Hoffmann, W. A., and Franco, A. C.: Carbon accumulation and  
517 nitrogen pool recovery during transitions from savanna to forest in central Brazil,  
518 *Ecology*, 95, 342-352, <https://doi.org/10.1890/13-0290.1>, 2014.
- 519 Pellegrini, A. F. A., Ahlstrom, A., Hobbie, S. E., Reich, P. B., Nieradzik, L. P., Staver, A.  
520 C., Scharenbroch, B. C., Jumpponen, A., Anderegg, W. R. L., Randerson, J. T., and  
521 Jackson, R. B.: Fire frequency drives decadal changes in soil carbon and nitrogen and  
522 ecosystem productivity, *Nature*, 553, 194-198, <https://doi.org/10.1038/nature24668>,  
523 2018.
- 524 Qualls, R. G., and Haines, B. L.: Geochemistry of Dissolved Organic Nutrients in Water  
525 Percolating through a Forest Ecosystem, *Soil Sci. Soc. Am. J.*, 55, 1112-1123,  
526 <https://doi.org/10.2136/sssaj1991.03615995005500040036x>, 1991.
- 527 Qualls, R. G., Haines, B. L., and Swank, W. T.: Fluxes of dissolved organic nutrients and  
528 humic substances in a deciduous forest, *Ecology*, 72, 254-266,  
529 <https://doi.org/10.2307/1938919>, 1991.
- 530 Qualls, R. G.: Comparison of the behavior of soluble organic and inorganic nutrients in  
531 forest soils, *Forest Ecol. Manag.*, 138, 29-50, [https://doi.org/10.1016/S0378-1127\(00\)00410-2](https://doi.org/10.1016/S0378-1127(00)00410-2), 2000.
- 532 Reich, P. B., Walters, M. B., and Ellsworth, D. S.: Leaf life-span in relation to leaf, plant,  
533 and stand characteristics among diverse ecosystems, *Ecol. Monogr.*, 62, 365-392,  
534 <https://doi.org/10.2307/2937116>, 1992.
- 535 Reich, P. B., Peterson, D. W., Wedin, D. A., and Wrage, K.: Fire and vegetation effects  
536 on productivity and nitrogen cycling across a forest-grassland continuum, *Ecology*, 82,  
537 1703-1719, [https://doi.org/10.1890/0012-9658\(2001\)082\[1703:FAVEOP\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[1703:FAVEOP]2.0.CO;2),  
538 2001.
- 539 Reich, P. B., and Oleksyn, J.: Global patterns of plant leaf N and P in relation to  
540 temperature and latitude, *P. Natl. Acad. Sci. U.S.A.*, 101, 11001-11006,  
541 <https://doi.org/10.1073/pnas.0403588101>, 2004.
- 542 Rivas-Ubach, A., Sardans, J., Perez-Trujillo, M., Estiarte, M., and Penuelas, J.: Strong  
543 relationship between elemental stoichiometry and metabolome in plants, *P. Natl. Acad. Sci. U.S.A.*, 109, 4181-4186, <https://doi.org/10.1073/pnas.1116092109>, 2012.
- 544 Schreeg, L. A., Santiago, L. S., Wright, S. J., and Turner, B. L.: Stem, root, and older leaf  
545 N:P ratios are more responsive indicators of soil nutrient availability than new foliage,  
546 *Ecology*, 95, 2062-2068, <https://doi.org/10.1890/13-1671.1>, 2014.
- 547 Smithwick, E. A. H.: Pyrogeography and Biogeochemical Resilience, in: *The Landscape Ecology of Fire*, *Ecol. Stud.*, Springer, Dordrecht, 143-163, [https://doi.org/10.1007/978-94-007-0301-8\\_6](https://doi.org/10.1007/978-94-007-0301-8_6), 2011.
- 548 Spink, A., Sparks, R. E., van Oorschot, M., and Verhoeven, J. T. A.: Nutrient dynamics  
549 of large river floodplains, *Regulated Rivers: Research and Management*, 14, 203-216,  
550 [https://doi.org/10.1002/\(SICI\)1099-1646\(199803/04\)14:2<203::AID-RRR498>3.0.CO;2-7](https://doi.org/10.1002/(SICI)1099-1646(199803/04)14:2<203::AID-RRR498>3.0.CO;2-7),  
551 1998.



556 Taylor, B. R., Parsons, W. F. J., and Parkinson, D.: Decomposition of *Populus*  
557 *tremuloides* leaf litter accelerated by addition of *Alnus crispa* litter, *Can. J. Forest Res.*,  
558 19, 674-679, <https://doi.org/10.1139/x89-104>, 1989.  
559 Turner, J., Dice, S. F., Cole, D. W., and Gessel, S. P.: Variation of nutrients in forest tree  
560 foliage - a review, College of Forest Resources, University of Washington, Seattle,  
561 Washington, 1977.  
562 Valladares, F., Gianoli, E., and Gomez, J. M.: Ecological limits to plant phenotypic  
563 plasticity, *New Phytol.*, 176, 749-763, <https://doi.org/10.1111/j.1469-8137.2007.02275.x>,  
564 2007.  
565 van Duren, I. C., and Peggel, D. M.: Nutrient limitations in wet, drained and rewetted fen  
566 meadows: evaluation of methods and results, *Plant Soil*, 220, 35-47,  
567 <https://doi.org/10.1023/A:1004735618905>, 2000.  
568 Violle, C., Navas, M. L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., and Garnier,  
569 E.: Let the concept of trait be functional!, *Oikos*, 116, 882-892,  
570 <https://doi.org/10.1111/j.2007.0030-1299.15559.x>, 2007.  
571 Wardle, D. A., Walker, L. R., and Bardgett, R. D.: Ecosystem properties and forest  
572 decline in contrasting long-term chronosequences, *Science*, 305, 509-513,  
573 <https://doi.org/10.1126/science.1098778>, 2004.  
574 Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F.,  
575 Cavender-Bares, J., Chapin, T., Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E.,  
576 Groom, P. K., Gullas, J., Hikosaka, K., Lamont, B. B., Lee, T., Lee, W., Lusk, C.,  
577 Midgley, J. J., Navas, M. L., Niinemets, Ü., Oleksyn, J., Osada, N., Poorter, H., Poot, P.,  
578 Prior, L., Pyankov, V. L., Roumet, C., Thomas, S. C., Tjoelker, M. G., Veneklaas, E. J.,  
579 and Villar, R.: The worldwide leaf economics spectrum, *Nature*, 428, 821-827, 2004.  
580 Yelenik, S., Perakis, S., and Hibbs, D.: Regional constraints to biological nitrogen  
581 fixation in post-fire forest communities, *Ecology*, 94, 739-750,  
582 <https://doi.org/10.1890/12-0278.1>, 2013.  
583  
584  
585