

Collision and Breakup of Fractal Particle Agglomerates in a Shear Flow

Farzad F. Dizaji, Jeffrey S. Marshall and John R. Grant
Department of Mechanical Engineering
The University of Vermont

Corresponding Author: Jeffrey S. Marshall, Department of Mechanical Engineering, The University of Vermont, Burlington, VT 05405, U.S.A. PHONE: 1 (802) 656-3826, EMAIL: jmarshal@uvm.edu.

Keywords: particle agglomeration; agglomerate collisions; turbulent agglomeration

Abstract

A computational study was performed both of a single agglomerate and of the collision of two agglomerates in a shear flow. The agglomerates were extracted from a direct numerical simulation of a turbulent agglomeration process, and had the loosely-packed fractal structure typical of agglomerate structures formed in turbulent agglomeration processes. The computation was performed using a discrete-element method for adhesive particles with four-way coupling, accounting both for forces between the fluid and the particles (and vice versa) as well as force transmission directly between particles via particle collisions. In addition to understanding and characterizing the particle dynamics, the study focused on illuminating the fluid flow field induced by the agglomerate in the presence of a background shear and the effect of collisions on this particle-induced flow. Perhaps the most interesting result of the current work was the observation that the flow field induced by a particle agglomerate rotating in a shear flow has the form of two tilted vortex rings with opposite sign circulation. These rings are surrounded by a sea of stretched vorticity from the background shear flow. The agglomerate rotates in the shear flow, but at a slower rate than the ambient fluid elements. In the computations with two colliding agglomerates, we observed cases resulting in agglomerate merger, bouncing and fragmentation. However, the bouncing cases were all observed to also result in an exchange of particles between the two colliding agglomerates, so that they were influenced both by elastic rebound of the agglomerate structures as well as by tearing away of particulate matter between the agglomerates. Overall, the problems of agglomerate-flow interaction and of the collision of two agglomerates in a shear flow are considerably richer in physical phenomena and more complex than can be described by the common approximation that represents each agglomerate by an 'equivalent sphere'.

1. Introduction

Collision of particle agglomerates with each other and with container walls or other obstacles in turbulent flow fields is important during both the agglomerate formation and breakup processes. The significance of agglomerate collisions has been studied for important industrial processes such as drug particle dispersion in dry powder inhalers (Tong et al., 2013, 2016; Yang et al., 2014), cyclone operation (Tong et al., 2010), and particle filtration (Iimura et al., 2009a,b). Similar agglomerate-agglomerate collision processes occur in astrophysics during formation of protoplanetary disks (Ormel et al, 2007, 2009) and in the dynamics of planetary rings (Schäfer et al., 2007).

The development of particle agglomerates in turbulent flows occurs through a series of processes in which individual particles collide and adhere to form small agglomerates, and these small agglomerates then collide and adhere to each other to form larger agglomerates, and so forth (Dizaji and Marshall, 2016, 2017). As they increase in size, the agglomerates begin to lose particles by processes such as erosion of small groups of particles from an agglomerate surface or rupture of the agglomerate into smaller pieces in response to the fluctuating turbulent shear flow (Serra et al., 1997; Higashitani et al., 2001), eventually balancing the agglomerate formation processes to achieve a quasi-equilibrium state (provided that the turbulence itself is in an equilibrium state). As discussed by Sayvet and Navard (2000), a dominant agglomerate breakup process for turbulent flows at lower shear stress values is fragmentation of agglomerates during collisions with other agglomerates. The question of whether two colliding agglomerates will merge together, bounce off each other, or split apart into a larger number of fragments is thus one of central importance for a wide range of processes. All three of these outcomes were observed under different conditions in a microgravity experiment of particle agglomerates in a

vibrating box by Brisset et al. (2016) for different values of the collision velocity and in a normal gravity experiment by Ihalainen et al. (2012) in which agglomerates were impacted onto a flat surface.

An important simplification that is often made in modeling turbulent agglomeration is replacement of a particle agglomerate by single 'effective particle', often selected as a sphere with the same mass as the agglomerate. This assumption is integral to the traditional population balance model for agglomerate formation (Smoluchowski, 1917; Lu and Wang, 2006; Reinhold and Briesen, 2012), and it plays an important part in many analytical statistical models for the early stages of agglomerate formation in turbulence (Brunk et al., 1998; Chun and Koch, 2005; Koch and Pope, 2002; Wang et al., 1998). The equivalent sphere assumption is also used in the 'extended hard-sphere' discrete-element method (DEM), which seeks to use the hard-sphere approach for binary collisions to study formation of particle agglomerates (Kosinski and Hoffmann, 2010; Balakin et al., 2011). All such applications of this equivalent sphere approximation must impose some external criterion for whether or not an agglomerate will stick or bounce upon collision. Although the equivalent sphere approximation is commonly made for simulation of turbulent flows of adhesive particles, the accuracy of this approximation has not been addressed in detail. Can mechanical properties be assigned to the effective particle such that its collision with another effective particle accurately approximates the collision of two agglomerates? Are agglomerate collisions more complex than can be represented by a simple stick or bounce decision? Addressing these questions is one of the primary objectives of the current paper.

There is a fairly large literature on use of the discrete element method for examining collision of tightly-packed agglomerates with a wall (Kafui and Thornton, 2000; Lian et al.,

1998; Moreno et al., 2003; Moreno-Atanasio et al., 2006; Thornton et al., 1999; Ning et al., 1997; Thornton and Liu, 2004), with each other (Kun and Herrmann, 1999; Schäfer et al., 2007; Seizinger and Kley, 2013; Tong et al., 2009), or with some other obstacle, such as a cylinder or sphere in the flow field (Iimura et al., 2009a,b; Yang et al., 2014). Experimental studies of compressed particle aggregates with each other (Beitz et al., 2011) and with a wall (Samimi et al., 2004) have also been reported. Much of this work is motivated by the problem of deagglomeration of particles in dry powder inhalers (Tong et al., 2013, 2016; Yang et al., 2014), used to break up agglomerates and deliver small drug particles to the lungs, where they are absorbed. In this application, the particles are initially compressed into tightly-packed aggregates at the time of manufacture, which then need to be broken up to release the small drug particles at time of use. Alternatively, ice particles can form tightly-packed aggregates in planetary rings (Schäfer et al., 2007), and the dynamics of their collision plays a central role in understanding the ring dynamics.

A useful definition of agglomerate strength was given by Moreno-Atanasio and Ghadiri (2006), based on the work of Rumpf (1962), as “the force that is required to break all contacts simultaneously on a prescribed failure plane”. This force depends both on the strength of the individual contacts and the number of contacts in the failure plane. The number of contacts in any given cross-sectional plane increases with the agglomerate fractal dimension, with higher values for tightly-packed agglomerates with fractal dimension close to $d_f \cong 3$ and lower values for the loosely-structured agglomerates more typically formed in turbulent flocculation processes, with fractal dimension closer to $d_f \sim 2$. For instance, in experiments with turbulent agglomeration of latex particles in stirred tanks, Selomulya et al. (2001) reported d_f between 1.7 - 2.1 and Waldner et al. (2005) reported d_f between 1.8 - 2.6. The above definition of

agglomerate strength is based on the idea of pulling an agglomerate apart in tension, whereas the agglomerate response to collision is also dependent on its behavior under compression. In compressive deformation, agglomerates with lower values of particle concentration are more susceptible to buckling of force chains due to having fewer surrounding particles (Marangoni and Narine, 2001). The sensitivity of agglomerate collisions to particle concentration c (or void fraction $\varepsilon = 1 - c$) was noted in DEM simulations by Gunkelmann et al. (2016), who in a study of head-on collision of two agglomerates in a vacuum found that agglomerates with higher porosities are more fragile during collision and have higher tendency to fragment. These conclusions are also supported by the simulations of Nguyen et al. (2014) of the collision of a loose-structured agglomerate of fine particles with a larger spherical particle, who found a higher tendency of the loose-structured agglomerate to fragment compared to simulations with highly packed agglomerates.

The current paper examines the collision of two particle agglomerates in a shear flow under conditions typical of agglomerate collision in turbulent flows. The primary objective of the paper is to address two issues: (1) to understand the flow field induced by a particle agglomerate in a shear flow and (2) to evaluate the accuracy of the equivalent sphere approximation by examining the physics of actual agglomerate-agglomerate collisions with loosely-structured agglomerates. The agglomerate collision is computed using a CFD-DEM approach based on the soft-sphere method with four-way coupling. The CFD-DEM approach does not resolve flow around individual particles, but instead it introduces a distributed body force that accounts for the influence of particles on the bulk fluid flow. The bulk flow within the agglomerates can be resolved by the fluid flow computation since the agglomerates selected consist of several hundred particles, and so are much larger than the individual particle size. Loosely-structured

agglomerates are first generated from a direct numerical simulation of turbulent agglomeration (Dijazi and Marshall, 2016), and from which agglomerates are extracted and placed in a shear flow. We first examine agglomerate evolution and breakup in shear with no collision, and then examine the effect of two-agglomerate collision on agglomerate merger, bouncing, and fragmentation. The paper differs from previous work in its focus on agglomerate collision in shear flows, in its use of loosely-structured agglomerates typical of turbulent flocculation processes, and in its focus on fluid flow effects on the agglomerate collision.

2. Computational Method

The agglomerate breakup and collision is computed using the four-way coupled CFD-DEM approach, using an adhesive soft-sphere discrete-element method for the particles and a high-order finite-difference method for the fluid. The DEM approach is based on evolving the motion of each individual particle by solution of the particle momentum and angular momentum equations, while accounting for the many different forces and torques acting between the particles due to collision and van der Waals adhesion effects as well as between the particles and the surrounding fluid via a set of model equations. The flow around each individual particle is not resolved, but rather the computational method introduces the force imposed by the particles on the fluid as a smooth body force field, which is generated by the combined forces acting on many particles in a local region. A conservative particle blob method (Marshall and Sala, 2013) is used in the current paper to translate between forces on individual particles and the body force acting on the fluid grid in a manner that is well suited for cases where the ratio of particle diameter to the grid cell spacing is of order unity.

The soft-sphere DEM formulation can be used both to simulate isolated particles and to simulate particles contained in agglomerates. An agglomerate is defined in the current paper as an assemblage of particles in which each particle in the agglomerate is in contact with at least one other particle in the agglomerate, in such a manner that one can continuously travel between any two particles in the agglomerate by following a chain of contacts. With use of the soft-sphere DEM approach, the motion and deformation of the agglomerate is simulated by evolving the motion and rotation of its constituent parts.

As is standard in DEM, the drag on each particle is given by the Stokes drag expression multiplied by a particle crowding factor that accounts for the effect of surrounding particles on the drag force and an inertia factor that accounts for finite particle inertia. The particle crowding factor was determined empirically as a function of the local particle concentration and the particle Reynolds number based on experiments with a fluidized particle bed (Di Felice, 1994). While this approach is commonly employed, we note that it does not account for the effects of strongly heterogeneous concentration along the sides of the agglomerate. The pairwise-interaction extended point-particle (PIEPP) method recently proposed by Akiki et al. (2017a,b) might be one approach that could be used to account for the effect of heterogeneity in future studies of agglomerate flows. However, even without such corrections previous studies with the CFD-DEM method, such as that of Bosse et al. (2005) for a particle suspension droplet falling under gravity, have produced predictions for agglomerate formation and dynamics in excellent agreement with experimental observations. A review of the CFD-DEM approach by Zhu et al. (2007) provides a detailed discussion of the modeling approximations used in this method.

The computations proceed in two parts. The first part is concerned with the initial formation of agglomerates in a turbulent flow, and the approach used for these computations

have been described in detail in a previous paper (Dizaji and Marshall, 2017). The second part conducts a detailed examination of the collision process that occurs when either one or two of the agglomerates are extracted and placed in a plane shear flow, which is intended to represent a very small section of the overall turbulent flow. A summary of the DEM and CFD methods used to simulate the particles and the fluid flow, and of the turbulent flow computations used to initialize the agglomerate structure, is given below.

2.1. Discrete element method

The computations of particle agglomerate breakup and collision are performed using a soft-sphere adhesive discrete element method (Marshall, 2009). Because particle collision and adhesion processes involve a wide range of time scales, a multiple time step algorithm is used in the current paper in which the fluid time step $\Delta t = O(\ell / u_0)$, the particle time step $\Delta t_p = O(d / u_0)$, and the collision time step $\Delta t_c = O(d(\rho_p^2 / E_p^2 u_0)^{1/5})$ satisfy $\Delta t > \Delta t_p > \Delta t_c$. Here d is the particle diameter, ρ_p is the particle density, and E_p is the particle elastic modulus. The method follows the motion of individual particles in the three-dimensional fluid flow by solution of the particle momentum and angular momentum equations

$$m \frac{d\mathbf{v}}{dt} = \mathbf{F}_F + \mathbf{F}_A, \quad I \frac{d\mathbf{\Omega}}{dt} = \mathbf{M}_F + \mathbf{M}_A, \quad (1)$$

subject to forces and torques induced by the fluid flow ($\mathbf{F}_F$, $\mathbf{M}_F$) and by particle collision and van der Waals adhesion ($\mathbf{F}_A$, $\mathbf{M}_A$). In this equation, m is the particle mass, I is the moment of inertia, and $\mathbf{v}$ and $\mathbf{\Omega}$ are the particle velocity and rotation rate, respectively. The dominant fluid

force is the drag force, given by the Stokes drag law modified to account for the effects of particle inertia and local particle crowding as

$$\mathbf{F}_d = 3\pi\mu d(\mathbf{u} - \mathbf{v})f, \quad (2)$$

where $\mathbf{u}$ is the fluid velocity evaluated at the particle centroid. The friction factor $f = C_I C_C$ is written as the product of an inertial correction term C_I and a particle crowding correction term C_C . An expression for the inertial correction was given by Schiller and Naumann (1933) as

$$C_I = 1 + 0.15 \text{Re}_p^{0.687}, \quad (3)$$

where $\text{Re}_p = \rho_f d v_s / \mu$ is the particle Reynolds number and $v_s = |\mathbf{v} - \mathbf{u}|$ is the magnitude of the particle slip velocity relative to the fluid. This expression is valid to within 5% of comparison experimental data for particle Reynolds number up to about 800. An expression for the crowding correction factor was determined empirically by Di Felice (1994) for particle Reynolds numbers in the range 0.01 to 10^4 as a function of the void fraction ε as

$$C_C = \varepsilon^{1-\zeta}, \quad \zeta = 3.7 - 0.65 \exp\left(-\frac{1}{2}[1.5 - \ln(\text{Re}_p)]^2\right). \quad (4)$$

This expression approaches the Wen and Yu (1966) expression for low particle Reynolds number. A viscous fluid torque arises from a difference in rotation rate of the particle and the local fluid element (Crowe et al., 2012), and is given by

$$\mathbf{M}_F = -\pi\mu d^3(\boldsymbol{\Omega} - \frac{1}{2}\boldsymbol{\omega}), \quad (5)$$

where $\boldsymbol{\omega}$ is the fluid vorticity vector. While the drag is the primary fluid force acting on the particles, we also include in the computations several secondary forces such as the added mass force and the Saffman and Magnus lift forces (Saffman, 1965; Rubinow and Keller, 1961).

Particle collision and van der Waals adhesion forces are simulated using the classical Johnson-Kendall-Roberts (JKR) theory (Johnson et al., 1971). This theory assumes that particle diameter is much larger than the effective length scale over which the van der Waals adhesion force acts (approximately 10nm), so that the adhesion force can be assumed to only act within the flattened contact region between two colliding particles. No adhesion force acts between the particles prior to or following collision, when the particles are not in contact. Since the size of the contact region depends on both the elastic and adhesive forces, these forces are nonlinearly combined to yield the total contact/adhesion force on any pair of particles in contact with each other.

The collision and adhesion force and torque fields acting on particle i , with radius r_i , are given by

$$\mathbf{F}_A = F_n \mathbf{n} + F_s \mathbf{t}_S, \quad \mathbf{M}_A = r F_s (\mathbf{n} \times \mathbf{t}_S) + M_r (\mathbf{t}_R \times \mathbf{n}), \quad (6)$$

where $\mathbf{n} = (\mathbf{x}_j - \mathbf{x}_i) / |\mathbf{x}_j - \mathbf{x}_i|$ is the unit normal vector oriented along the line connecting the centers of the two colliding particles, i and j . The normal component of the collision and adhesion force F_n is further divided into an elastic-adhesion part F_{ne} and a dissipative part F_{nd} . The sliding resistance is composed of a force with magnitude F_s acting in a direction $\mathbf{t}_s$, corresponding to the direction of relative motion of the particle surfaces at the contact point projected onto the contact plane (the plane orthogonal to $\mathbf{n}$), as well as a related torque in the $\mathbf{n} \times \mathbf{t}_s$ direction. The rolling resistance, which arises due to the effects of particle adhesion, exerts a torque of magnitude M_r on the particle in the $\mathbf{t}_R \times \mathbf{n}$ direction, where $\mathbf{t}_R$ is the direction of the “rolling” velocity. While all of these various collision-adhesion forces and torques are included in the current computations, the dynamics of small adhesive particles are dominated by the normal elastic-adhesive force and the rolling resistance torque.

The effective elastic modulus E and the effective radius R are defined by

$$\frac{1}{E} \equiv \frac{1 - \sigma_i^2}{E_i} + \frac{1 - \sigma_j^2}{E_j}, \quad \frac{1}{R} \equiv \frac{1}{r_i} + \frac{1}{r_j}, \quad (7)$$

where E_i , σ_i , and r_i are the elastic modulus, Poisson ratio, and radius of particle i , respectively.

The adhesive force between the two particles depends on the surface energy potential γ , where the work required to separate two spheres colliding over a contact region of radius $a(t)$ is given by $2\pi\gamma a^2$ in the absence of further elastic deformation. Particle normal elastic rebound force and van der Waals adhesion force are simulated using the JKR theory, which can be written in terms

of the contact region radius $a(t)$ and the normal particle overlap $\delta_N = r_i + r_j - |\mathbf{x}_i - \mathbf{x}_j|$ as (Chokshi et al. 1993)

$$\frac{\delta_N}{\delta_c} = 6^{1/3} \left[2 \left(\frac{a}{a_o} \right)^2 - \frac{4}{3} \left(\frac{a}{a_o} \right)^{1/2} \right], \quad \frac{F_{ne}}{F_c} = 4 \left(\frac{a}{a_o} \right)^3 - 4 \left(\frac{a}{a_o} \right)^{3/2}, \quad (8)$$

The critical overlap δ_c , the critical normal force F_c , and the equilibrium contact region radius a_o are given by (Johnson et al. 1971)

$$F_c = 3\pi\gamma R, \quad \delta_c = \frac{a_o^2}{2(6)^{1/3} R}, \quad a_o = \left(\frac{9\pi\gamma R^2}{E} \right)^{1/3}. \quad (9)$$

As two particles move away from each other following collision, they remain in contact until the point where $F_n = -F_c$ and $\delta_N = -\delta_c$ due to the necking of the material in the contact region. Beyond this state any further separation leads the two particles to break apart.

The effect of lubrication forces within the fluid squeeze-film within the contact region is to limit the minimum approach distance between the particles (i.e., the contact region gap size) and to reduce the particle restitution coefficient. Experimental studies of particle collisions at different Stokes numbers (e.g., Joseph et al., 2001) indicate that the coefficient of restitution is essentially zero when the Stokes number is less than about 10 due to dissipation in the squeeze-film. We use the model of Tsuji et al. (1992) for the dissipative part of the normal collision force F_{nd} and set the damping parameter such that the restitution coefficient vanishes.

The second major effect of particle adhesion is to introduce a torque that resists particle rolling. For uniform-size spherical particles, the “rolling velocity” $\mathbf{v}_L$ of particle i is given by (Bagi and Kuhn 2004)

$$\mathbf{v}_L = -R(\boldsymbol{\Omega}_i - \boldsymbol{\Omega}_j) \times \mathbf{n} . \quad (10)$$

A linear expression for the rolling resistance torque M_r is postulated as

$$M_r = -k_R \xi , \quad (11)$$

where $\xi = (\int_{t_0}^t \mathbf{v}_L(\tau) d\tau) \cdot \mathbf{t}_R$ is the rolling displacement in the direction $\mathbf{t}_R = \mathbf{v}_L / |\mathbf{v}_L|$. Rolling involves an upward motion of the particle surfaces within one part of the contact region and a downward motion in the other part of the contact region. The presence of an adhesion force between the two contacting surfaces introduces a torque resisting rolling of the particles. An expression for the rolling resistance due to van der Waals adhesion was derived by Dominik and Tielens (1995), which yields the coefficient k_R as

$$k_R = 4F_c(a/a_0)^{3/2} . \quad (12)$$

Dominik and Tielens (1995) further argue that the critical resistance occurs when the rolling displacement ξ achieves a critical value, corresponding to a critical rolling angle $\theta_{crit} = \xi_{crit} / R$.

For $\theta > \theta_{crit}$, the rolling displacement ξ in (11) is replaced by ξ_{crit} . Data for critical rolling angle for particles having diameter of approximately 10 μm were reported by Ding et al. (2008), who found critical rolling angles θ_{crit} of between 0.02 and 0.06 radians.

A simplified expression for the effect of van der Waals adhesion on tangential sliding resistance was proposed by Thornton (1991). In this model, the sliding resistance force F_s is given by a spring-like expression of the form (Cleary et al., 1998)

$$F_s = -k_T \left(\int_{t_0}^t \mathbf{v}_s(\xi) d\xi \right) \cdot \mathbf{t}_s \quad (13)$$

when F_s is less than a critical value F_{crit} . In (13), the sliding velocity $\mathbf{v}_s(t)$ is the relative tangential surface velocity of the particles at the contact point projection. The tangential stiffness coefficient k_T is derived by Mindlin (1949) and can be written in terms of the contact region radius $a(t)$ as

$$k_T = 8G a(t) . \quad (14)$$

The critical sliding force is approximated using the expression

$$F_{crit} = \mu_f |F_{ne} + 2F_c|, \quad (15)$$

where F_c is the critical force for pull-off given in (9) and μ_f is the friction coefficient. The expression (15) was shown by Thornton (1991) to provide results in reasonable agreement to experiments. For $|F_s| \geq F_{crit}$, the sliding resistance is given by the Amonton expression

$$F_s = -F_{crit}.$$

2.2. Agglomerate formation

The agglomerates are formed using a turbulent agglomeration process similar to that described by Dizaji and Marshall (2017). The computations employed a pseudo-spectral method for forced turbulence in a triply-periodic domain with side length 2π and with 128 grid points in each direction. The turbulence is initiated with random perturbations and allowed to develop with no particles until it approached a quasi-steady state corresponding to microscale Reynolds number $Re_\lambda = u_0 \lambda / \nu = 99$. Particles are then added to the computation, with 46,656 particles spread randomly over the flow field with diameter $d = 0.04$ and particle-to-fluid density ratio $\rho_p / \rho_f = 10$. Over time as the particles are advected by the flow, small agglomerates first form and then collide with each other to form progressively larger agglomerates.

The computation was stopped once the agglomerates achieved a broad range of sizes. One common way to measure the size of an agglomerate is the radius of gyration R_g , which for an agglomerate with N particles is defined by

$$R_g = \left[\sum_{i=1}^N |\mathbf{x}_i - \bar{\mathbf{x}}|^2 \right]^{1/2}. \quad (16)$$

In this equation, $\bar{\mathbf{x}}$ denotes the position vector of the agglomerate centroid and $\mathbf{x}_i$ is the centroid of the i^{th} particle within the agglomerate. Particle agglomerates admit a power law relating N and R_g given by (Adachi and Ooi, 1990)

$$N = K \left(R_g / r_p \right)^{d_f}, \quad (17)$$

where K is a coefficient (called the fractal pre-factor), r_p is the individual particle radius, and the exponent d_f is the fractal dimension of the agglomerate. The value of d_f for particle agglomerates varies over the interval $1 \leq d_f \leq 3$ depending on the agglomeration formation mechanism (Brasil et al., 2001); however, typical values for turbulent particle agglomeration processes are between about 1.7 - 2.8 (Selomulya et al., 2001; Waldner et al.). A log-log plot of N versus R_g / r_p for the current turbulent flow simulation is given in Figure 1. The best-fit line to the computational predictions has slope $d_f = 2.12$, which is consistent with the range of fractal dimension observed in the experimental turbulent particle agglomeration studies listed above.

2.3. Shear flow simulation

The agglomerates extracted from the turbulent agglomeration calculation described in Section 2.2 are immersed in a linear shear flow, where the initial configuration appears as shown in Figure 2a for cases with a single agglomerate in the shear flow and as shown in Figure 2b for cases with agglomerate collision. Over time, the shear flow is modified by the presence of the particles, as described below. The fluid flow is assumed to be incompressible and is governed by the continuity and momentum equations of the form

$$\nabla \cdot \mathbf{u} = 0, \quad (18a)$$

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho_f} \nabla p + \nu \nabla^2 \mathbf{u} + \mathbf{F}_p. \quad (18b)$$

In this equation, $\mathbf{u}$, p and $\mathbf{F}_p$ are the fluid velocity, the pressure and the particle-induced body force per unit mass, respectively. The void fraction $\varepsilon = 1 - c$ was not included in (18) since our computations indicate that local void fraction remains above 90% even within the agglomerates for current computations due to the loose structuring of agglomerates typical of turbulent flow.

Since the grid cell size is on the same order as the particle size in these computations, we have not used the standard point-force approach in which the force imposed on the fluid by a particle is assigned to the grid cell containing the particle center. Instead, in computing both the particle-induced body force and the particle concentration field, we have smoothed the particle field with use of the conservative particle blob method proposed by Marshall and Sala (2013). In this method, the particle body force field $\mathbf{F}_p(\mathbf{x}, t)$ is written as the sum of some number N particle ‘blobs’, centered at positions $\mathbf{x}_n$, as

$$\mathbf{F}_p(\mathbf{x}, t) = \sum_{n=1}^N \mathbf{A}_n f_w(\mathbf{x} - \mathbf{x}_n, R_n). \quad (19)$$

The Gaussian weight f_w is a function of position and of the characteristic blob ‘radius’ R_n , and can be written as

$$f_w(\mathbf{x} - \mathbf{x}_n, R_n) = \frac{2}{3\pi R_n^3} \exp[-|\mathbf{x} - \mathbf{x}_n|^2 / R_n^2]. \quad (20)$$

The blob amplitude, $\mathbf{A}_n$, is given by

$$\mathbf{A}_n = \frac{(-\mathbf{F}_{f,n})}{G_{cell} \sum_{j=1}^Q f_w(\mathbf{g}_j - \mathbf{x}_n, R_n)}, \quad (21)$$

where $\mathbf{g}_j$ is the location of the centroid of grid cell j , $\mathbf{x}_n$ is the centroid of particle n , G_{cell} is the grid cell volume, and $\mathbf{F}_{f,n}$ is the fluid-induced force acting on particle n (which imposes an equal and opposite force $-\mathbf{F}_{f,n}$ back on the fluid). The force $\mathbf{F}_{f,n}$ is given by the sum of the drag force in (2) plus minor forces such as lift, added mass force, and pressure gradient force. Each particle distributes part of its force to a set Q of surrounding grid cells, and the sum in the denominator of (21) is evaluated over all grid cells in this set Q . With the choice (21) for blob amplitude, the discrete-to-continuum conversion operation is discretely conservative. This method can be applied to the particle concentration simply by replacing the particle force with the particle volume.

The fluid flow computations were performed using a fractional-step method (Rai and Moin, 1991; Verzicco and Orlandi, 1996; Uhlmann, 2005), with time advancement performed using a third-order Runge-Kutta method for convective terms and the 2nd order Crank-Nicholson method for viscous terms. Algorithms for all spatial derivatives except the convective terms are approximated using second-order centered finite differences (three-point stencil) on a non-staggered grid. The discretized equations for the k th Runge-Kutta step are given by

$$\begin{aligned}\tilde{\mathbf{u}} = & \mathbf{u}^{k-1} + \Delta t \left(2\alpha_k \nu \nabla^2 \mathbf{u}^{k-1} - 2\alpha_k \nabla p^{k-1} \right) \\ & - \Delta t \left\{ \gamma_k [(\mathbf{u} \cdot \nabla) \mathbf{u} - \mathbf{F}_p]^{k-1} + \zeta_k [(\mathbf{u} \cdot \nabla) \mathbf{u} - \mathbf{F}_p]^{k-2} \right\},\end{aligned}\quad (22a)$$

$$\nabla^2 \mathbf{u}^* - \frac{\mathbf{u}^*}{\alpha_k \nu \Delta t} = -\frac{\tilde{\mathbf{u}}}{\alpha_k \nu \Delta t} + \nabla^2 \mathbf{u}^{k-1}, \quad (22b)$$

$$\nabla \cdot (\nabla \phi^k) = \frac{\nabla \cdot \mathbf{u}^*}{2\alpha_k \Delta t}, \quad (22c)$$

$$\mathbf{u}^k = \mathbf{u}^* - 2\alpha_k \Delta t \nabla \phi^k, \quad (22d)$$

$$p^k = p^{k-1} + \phi^k - \alpha_k \nu \Delta t \nabla^2 \phi^k, \quad (22e)$$

where α_k , γ_k , and ζ_k are coefficients given by Rai and Moin (1991). Continuity is enforced by a projection method leading to equation (22c) for the pseudo-pressure, denoted by ϕ . In the multigrid solution of this equation, the five-point stencil produced by successive application of the gradient operation followed by the divergence operation was employed, rather than the finite-difference approximation to the Laplacian. The Crank-Nicholson method was used to solve the Helmholtz problem, given in (22b). A tenth-order approximation was used for the convective terms, requiring an 11-point stencil. To control non-linear instabilities, at the end of each time step the velocity components were filtered using a tenth-order filter (again using an 11-point stencil)(Lele, 1992; Steijl, 2001). After filtering to obtain $\mathbf{u}^{filtered}$, the velocity $\mathbf{u}$ was replaced by $(1-q)\mathbf{u} + q\mathbf{u}^{filtered}$, with $q = 0.05$.

The flow was initialized in the x -direction with linear variation in the y -direction. The upper wall at $y = 2$ was maintained at a velocity $u = 1$ and the lower wall at $y = -2$ was maintained at a velocity of $u = -1$, giving a dimensionless shear rate of $S = 0.5$. The no-slip

boundary condition was applied at both the top and bottom wall in the y -direction, and the flow was assumed to be periodic in the x - and z -directions. A layer of five ghost points in each direction surrounded the computational domain, so that no adjustment of the differentiation schemes was needed near the domain boundaries. The velocity on the ghost points was set at the upper and lower edges of the grid by linearly extrapolating the velocity from the point on the wall and the first point off of the wall. The velocity on the ghost points in the x - and z -directions were set so as to enforce periodicity. The fluid flow calculations were carried out on a Cartesian grid with equal spacing in each direction. The computations were performed on a 128^3 grid covering the interval $(-2,2)$ in each coordinate direction. The time step was held fixed at $\Delta t = 0.005$. The dimensionless fluid kinematic viscosity was set to $\nu = 0.0003$ for all computations.

3. Agglomerate Motion and Breakup in Shear Flow

In this section we examine the dynamics of a single particle agglomerate exposed to shear flow, with particular focus on examination of the particle-induced flow field associated with rotation of the agglomerate in the shear flow and on the conditions for agglomerate breakup. This section helps to set the stage for the study of agglomerate collision in shear flow in the next section. The problem of agglomerate dynamics in a shear flow has been previously examined by a number of authors. A series of experiments on this problem were reported by Sonntag and Russel (1986), who found that the average radius of gyration of the agglomerates could be expressed as a power law function of the shear rate as $R_g^3 \propto S^{-1.06}$. Since the average number of particles in the agglomerate, N , was related to radius of gyration by a power law expression of

the form (17), with $d_f \cong 2.48$ in their experiments, their expression for agglomerate size in the shear flow could alternatively be expressed as $N \propto S^{-0.878}$.

A number of DEM simulations of agglomerate dynamics in a shear flow have been reported (Potanin, 1993; Higashitani et al., 2001; Fanelli et al., 2006; Becker et al., 2009) based on the so-called free-draining approximation, which assumes that the particles do not influence the fluid flow (one-way coupling). Potanin (1993) and Becker et al. (2009) further assumed that particles did not influence fluid forces on each other (even under close packing in the agglomerate), whereas Higashitani et al. (2001) and Fanelli et al. (2006) assumed that fluid drag forces act only on particle surfaces on the outside of the agglomerate (i.e., that fluid does not penetrate into the agglomerate). Higashitani et al. (2001) observed that the average number of particles in broken agglomerate fragments, N , varies with the adhesion parameter as $N \propto Ad^{0.872}$, where Ad represents a ratio of adhesive to hydrodynamic force. Since Ad is inversely proportional to shear rate, this observation is consistent with the scaling found experimentally by Sonntag and Russel (1986). Becker et al. (2009) compared the DEM simulations using the free-draining approximation to a full finite-element simulation of the flow field and found that the free-draining approximation breaks down as the agglomerate size increases. This observation is consistent with that made in a recent CFD-DEM study of turbulent agglomeration by Dizaji and Marshall (2017), who compared results with one-way and two-way coupling and found significant deviance between the two as the agglomerate size increased. Becker et al. (2009) observed that small agglomerates rotate in an almost rigid-body fashion in the shear flow, large agglomerates break up into pieces, and agglomerates of an intermediate size undergo a restructuring process, in which they deform and change form as they rotate but do not break up.

A full CFD-DEM study of agglomerate dynamics in a shear flow was reported by Zeidan et al. (2007), but the computations were restricted to two dimensions and the models used for particle collision and adhesion forces were highly simplified. For instance, no tangential forces on the particles were included to resist rolling and sliding motions, which as noted by Becker et al. (2009) are important in modeling agglomerate deformation under the shear flow.

In the current section, we report on a three-dimensional CFD-DEM study of agglomerate dynamics in a shear flow using a complete and well-validated DEM approach, with a focus on resolving and understanding the flow field induced by the particles. In order to work with agglomerate structures typical of those found in turbulent agglomeration processes, the computations were initiated by extracting an agglomerate from the turbulent flow computation described in Section 2.2 and inserting it into an initially linear shear flow. The flow evolution is then computed using the CFD method described in Section 2.3 and the DEM model described in Section 2.1.

The shear flow acts to rotate and stretch the agglomerate, whereas the adhesion force acts to hold the agglomerate together as a rigid body. The competition between these two effects determines the agglomerate behavior in the shear flow. We let R_{g0} denote the initial radius of gyration of the agglomerate and S denote the ambient shear rate. The characteristic length, time and velocity scales of the flow were selected as R_{g0} , $1/S$, and SR_{g0} , respectively. The primary dimensionless parameter governing the agglomerate behavior in the shear flow is the adhesion parameter, which for current purposes is defined as the ratio of the adhesion force between individual particles ($O(\gamma d)$) to the viscous force ($O(\mu dU)$) imposed on a particle by the fluid flow. Using $U \sim SR_{g0}$ as the typical velocity scale, the adhesion parameter for this problem takes the form

$$\text{Ad} = \frac{\gamma}{\mu S R_{g0}}. \quad (23)$$

This measure is essentially the same as the inverse of the fragmentation number proposed by Hansen et al. (1998). A secondary parameter characterizing the particle motion is the particle Stokes number St , which is interpreted as the ratio of particle characteristic time scale $\tau_p = m / 3\pi\mu d$ to the fluid time scale $\tau_f = 1 / S$, giving

$$\text{St} = \frac{\rho_p d^2 S}{18\mu}. \quad (24)$$

The values of the adhesion parameter Ad , the initial number of particles N_0 , and the ratio R_{g0} / d of initial agglomerate gyration radius to particle diameter are given for all single-agglomerate runs in Table 1. All computations reported in the paper have Stokes number of $\text{St} = 1.4$ and density ratio of $\rho_p / \rho_f = 10$. The shear Reynolds number can be defined in terms of shear rate and radius of gyration as $\text{Re}_s = S R_{g0}^2 / \nu$, which is found to have a value ranging from 52-102 in the current computations, depending on which of the three extracted agglomerates are under consideration. In a turbulent flow, the parameters used in these computations would therefore be larger than the Kolmogorov scale and smaller than the integral scale, perhaps typical of the Taylor microscale of the turbulent motion.

Computations in this section were performed using three different agglomerates selected from the turbulent agglomeration simulation, and for four different adhesion parameter values

for each agglomerate. A time series of the particle positions during a typical run (Case A.4) for a case where the agglomerate rotates without breakup, but exhibits some restructuring during the rotation, is shown in Figure 3. The particles are immersed in a fluid flow, for which a velocity can be defined both outside and inside the particle agglomerate. The particles in Figure 3 are colored by the magnitude of the particle velocity relative to the fluid, which is called the *relative particle velocity* and defined by $\mathbf{w} \equiv \mathbf{v} - \mathbf{u}$, where $\mathbf{v}$ and $\mathbf{u}$ denote the particle velocity and fluid velocity at the particle centroid, respectively. We will also later refer to the *relative fluid velocity* $\mathbf{u}_{rel} = \mathbf{u} - S\mathbf{y}\mathbf{e}_x$, which is set equal to the computed fluid velocity $\mathbf{u}$ minus the velocity of the ambient shear flow ($S\mathbf{y}\mathbf{e}_x$).

In the reported computations, the initial velocity of the fluid was set equal to the shear flow velocity $S\mathbf{y}\mathbf{e}_x$. The initial velocity of the agglomerate particles is set equal to a rigid body rotation at the rotation rate $S/2$ of the shear flow, for which there exists a vertical y -component of velocity in addition to the x -component of velocity characteristic of the ambient shear. This initial rotation rate of the agglomerate gives rise to a linear variation of the relative particle velocity extending outward from the agglomerate center, as shown in Figure 3a. At later times, the size of the region of low relative particle velocity near the agglomerate center appears to grow and the particles with higher values of relative particle velocity are restricted to the outer parts of the agglomerate.

In the following, we shall examine in detail the results for Case A.4, which is typical of a case where the agglomerate does not break up in the shear flow. The particle coordination number for this computation remains nearly constant with time at a value of 3.9. The radius of gyration R_g and the particle concentration c_{agg} within the agglomerate oscillate in time, as shown in Figure 4b. The value of c_{agg} is computed by dividing the volume of all particles

associated with the agglomerate, $V_p = (\pi/6)Nd^3$, by the effective volume V_{eff} occupied by the agglomerate. The agglomerate effective volume is estimated by $V_{eff} = (4\pi/3)R_{eff,i}^3$, where the effective radius of the agglomerate R_{eff} is related to the radius of gyration by $R_{eff} = \sqrt{5/2} R_g$. This expression is based on the expression for radius of gyration of a solid sphere of uniform density. The particle volume fraction of the agglomerate can be related to the fractal dimension by (Jiang et al., 1991; Kusters et al., 1997)

$$c_{agg,i} = c_0 (R_{g,i} / d)^{d_f - 3}, \quad (25)$$

where c_0 is a constant. If the fractal dimension $d_f < 3$, an increase in agglomerate size results in a decrease in average particle volume fraction (Olfert et al., 2007). Both the radius of gyration and the particle concentration c_{agg} within the agglomerate oscillate during the computation as agglomerate restructuring occurs, with oscillation amplitude of about 3% of the mean radius of gyration and 9% of the mean particle concentration.

The time variation of the magnitude of the particle velocity $\mathbf{v}$ and the relative particle velocity $\mathbf{w}$ are plotted in Figure 5a. The particle velocity magnitude oscillates during the computation and the relative particle velocity exhibits a rapid initial decrease and then oscillates during the remainder of the computation. The latter result indicates that the fluid flow within the agglomerate responds quickly to changes in the particle velocity. While the relative velocity changes quickly in the time interval $0 < t < 1$, we do not observe significant deformation or breakup of the agglomerate during this interval. The fact that the relative particle velocity

magnitude is lower than the particle velocity magnitude for most of the computation is a result of the particle-induced flow, which acts to decrease the relative velocity.

The distribution of different quantities within the agglomerate is examined by computing the second-moment measure $\mu_i(F)$ of a given field $F(\mathbf{x})$ for each agglomerate i as

$$\mu_i(F) = \frac{N_i \left(\sum_{j=1}^{N_i} |\mathbf{x}_j - \bar{\mathbf{x}}_i|^2 F_j \right)}{\left(\sum_{j=1}^{N_i} |\mathbf{x}_j - \bar{\mathbf{x}}_i|^2 \right) \left(\sum_{j=1}^{N_i} F_j \right)}, \quad (26)$$

where $\bar{\mathbf{x}}_i$ is the centroid of agglomerate i and F_j is the value of the function $F(\mathbf{x})$ evaluated at the centroid $\mathbf{x}_j$ of the j^{th} particle within the agglomerate (Dizaji and Marshall, 2017). The second-moment measure is shown in Figure 5b for three different fields – the particle coordination number n_c , the magnitude of the relative particle velocity magnitude $\mathbf{w}$, and the magnitude of the relative particle rotation rate about the agglomerate center

$$\boldsymbol{\Omega}_{agg,rel} = (\mathbf{x} - \mathbf{x}_{c,agg}) \times \mathbf{w} / |\mathbf{x} - \mathbf{x}_{c,agg}|^2. \quad (27)$$

A value of the second moment $\mu(F)$ equal to unity indicates that the function $F(\mathbf{x})$ is uniform (or statistically randomly varying) across the agglomerate, whereas a value of $\mu_i(F)$ that is less (greater) than unity implies that particles with higher (lower) values of $F(\mathbf{x})$ are found near the center of the agglomerate compared to particles on the outer parts of the agglomerate. Figure 5b shows that the second moment measure for the coordination number is consistently less than

unity (close to 0.9), indicating that the agglomerate is more compact near its center than in its outer parts, as would be expected of a fractal agglomerate structure. The second moment of the relative velocity magnitude oscillates as the agglomerate restructures during rotation in the shear flow, but its value remains well above unity, varying from about 1.35 to 1.95. This observation supports the statement made earlier that small values of relative particle velocity are found near the center of the agglomerate and larger values are found only on the outermost particles. While this difference is related, in part, simply to the rotation of the agglomerate about its centroid, it is evident by comparison of Figure 3a and 3d that this effect becomes more pronounced with time, indicating that the particle-induced flow also plays a role. The relative particle rotation rate about the agglomerate centroid also oscillates in time, increasing from near unity at the start of the computation to an average value of about 1.2 in the second half of the computation. This quantity can be viewed as a measure of the effect of the particle-induced fluid flow - if there were no particle-induced flow the value of this quantity would remain at unity. The fact that this measure increases above unity is an indication that the particle-induced flow shields the inner parts of the agglomerate, resulting in a lower ratio of the relative velocity to radial distance in this region than in the outer part of the agglomerate. A somewhat similar observation of shielding of the center parts of agglomerates falling in a fluid was noted by Kusters et al. (1997).

The rotation frequency of a fluid element in the shear flow is equal to $f_{fluid} = (S/2)/2\pi \cong 0.0398$. The rotation period of the agglomerate was estimated by labeling each particle and observing the time required for one rotation. This measurement is necessarily somewhat imprecise since there is some restructuring of the agglomerate during the rotation, but we took care to also estimate the uncertainty in the estimate. Taking the inverse of the rotation period, our estimate of agglomerate rotation frequency for this computation is

$f_{agg} \cong 1/37.3 = 0.027 \pm 0.002$. Consequently, we observe that the particle agglomerate is rotating about 30% more slowly than would a fluid element in the shear flow. This observation is consistent with the findings of Li et al. (2016), who found that a porous circular particle in a two-dimensional shear flow rotates in the flow more slowly than a fluid element. In Figure 6a, we plot contours of the relative fluid velocity in the streamwise (x) direction, u_{rel} , at time $t = 20$, which is typical of the results observed throughout the computation. The relative fluid velocity is found to be oriented in a direction opposite to the ambient shear velocity, with negative value for $y > 0$ and positive value for $y < 0$. A profile of the relative fluid velocity along the y -axis ($x = z = 0$) is shown in Figure 6b as dots, with the ambient shear flow drawn as a solid line. We again see that the computed velocity in the region near the agglomerate ($|y| < 0.4$) lags behind the ambient shear velocity, which is due to the fact that the particle agglomerate is rotating more slowly than the fluid element so that the forces induced by the particles retard the fluid flow.

A series of plots in the three cross-sectional planes (x - y , x - z , and y - z) are shown in Figure 7, where for each plane we plot the in-plane streamlines (obtained by setting the normal velocity component to zero) and the contours of both the normal vorticity and velocity components. The plots do not include the entire computational domain, but instead focus on the central part of the domain near the agglomerate. In Fig. 7a, the streamlines in the x - y plane are seen to exhibit a vortex at the origin (i.e., at the center of the agglomerate); however, we note that the fluid velocity near the vortex center is very weak, and hence the normal vorticity magnitude at the vortex center is small. In all three cross-sectional planes, the normal vorticity component has a quadrupole structure, with four vorticity patches of alternating sign. From these cross-sectional plots, the velocity and vorticity fields associated with the rotating particle agglomerate appear to have the form of two tilted vortex rings with opposite circulation immersed in the shear flow.

To better illustrate this flow field, we compute the relative fluid vorticity $\boldsymbol{\omega}_{rel} = \boldsymbol{\omega} + S\mathbf{e}_z$, where we recall that the vorticity of the ambient shear flow is $-S\mathbf{e}_z$. The iso-surface $\omega_{rel} = 0.46$ of the magnitude of $\boldsymbol{\omega}_{rel}$ is plotted in Figure 8 in both the x - y plane (looking from the side) and the x - z plane (looking from the top). The same two views of this iso-surface are also shown in Figure 8 showing contours of ω_{rel} on a slice of the flow field in the normal plane. The ω_{rel} iso-surfaces clearly show that the particle-induced flow field for a single rotating agglomerate in a shear flow has the form of a pair of tilted vortex rings of opposite sign, with tilt angle of approximately 45° relative to the ambient shear flow (x -direction). As seen in the slices of the flow field in Figures 8c and 8d, each vortex ring is surrounded by stretched and reoriented vorticity from the ambient shear flow that trails behind the vortex rings in each direction. The dynamics of a single vortex ring in a linear shear flow was studied by Cheng et al. (2009), who found that the vortex ring becomes tilted relative to the shear and maintains a ring-like form while it drifts upward in the shear field (in the y -direction). This upward drift is negated in the current situation by the mutually-induced flow field when two rings of opposite sign co-exist, leading to a quasi-stationary flow with a quadrupole far-field structure (as is evident in the streamlines in Fig. 7c). For computations where the shear flow does not trigger breakup of the agglomerate, such as for Case A.4, this flow structure is observed to remain nearly constant with time as the agglomerate rotates in the shear flow.

As the adhesion parameter is varied in different computations, different behavior of the particle agglomerates in the shear flow is observed. For sufficiently low adhesion parameter values, some agglomerates are observed to break up into multiple fragments in the presence of the shear flow. A time series illustrating agglomerate breakup in the shear flow is shown in Figure 9 for Case A.1. We note from this example that while the fragments that shed from the

agglomerate are limited by the maximum size that the agglomerate can attain without breakup in the shear flow, there are also many agglomerates that are formed of a much smaller size. The set of fragments thus has a wide size distribution. A set of plots summarizing the computed agglomerate evolution for all of the single-agglomerate computations (Cases A.1 - A.12) is given in Figure 10. In Figure 10a, we plot the number of fragments N_{frag} into which the agglomerate breaks up as a function of the adhesion parameter Ad , defined in (22). The data are from three agglomerates extracted from the turbulent agglomeration computation, and different symbols are used in Figure 10 to denote the data from each agglomerate. For sufficiently high values of adhesion parameter, the agglomerate doesn't break up and the value of $N_{frag} = 1$ in Figure 10a. The number of particles N in each fragment at the end of the computation ($t = 30$) is plotted versus adhesion parameter in Figure 10b on a log-log plot. The power law expression $N \propto S^{-0.878}$ of Sonntag and Russel (1986) can be written in terms of the adhesion parameter as $N \propto Ad^{0.878}$. This expression is plotted as a dashed line in Figure 10b, where the coefficient of proportionality is fit to the data. The expression is found to be a reasonable fit for the maximum values of N , thus setting the largest size agglomerates that can survive without breakup in the shear flow.

4. Agglomerate Pair Collision in Shear Flow

In this section, we examine the collision of two agglomerates in a shear flow. As stated in Section 2, each particle in an agglomerate is in contact with at least one other particle in the agglomerate. Two agglomerates collide when at least one particle in each agglomerate come in contact with each other. For the computations of agglomerate collision reported in this section, three different agglomerates were extracted from the turbulent agglomeration computation

described in Section 2.2, which were used to conduct 30 computations of agglomerate collision, the parameters for which are listed in Table 2. For each computation, the agglomerates are initialized as shown in Figure 2b, with orientations of $\pm 45^\circ$ and displacement of the agglomerate centroid by an amount $\pm D_a$ in the y -direction. Each computation examines collision of an agglomerate with an exact copy, and we did not consider collisions of different size agglomerates. Consideration of the computational results indicates three different types of behaviors, which are illustrated in scatter plots in Figure 11. In these plots, each particle is colored either red or blue to indicate the agglomerate from which the particle originated. The first type of collision outcome is *merger* of the agglomerates into a single agglomerate, which then rotates in the shear flow. The second type of behavior, referred to as a *bouncing* collision, results in two separate agglomerates following the collision. As seen in Figure 11, it is common for some particles to be exchanged between the two colliding agglomerates during bouncing collisions. The third type of behavior is referred to as *fragmentation*, which describes collisions that result in three or more agglomerates. In the case shown in Figure 11, the collision results in three agglomerates - one composed entirely of red particles, one composed entirely of blue particles, and one composed of a combination of red and blue particles. In other cases, more than three agglomerates will form in a fragmentation collision, often yielding a wide variation in agglomerate sizes. Sometimes it is not clear whether a collision should be classified as a bouncing case or a fragmentation case; for instance, cases where two colliding agglomerates break away from each other but leave behind a very small third 'satellite' agglomerate composed of just a few particles can be regarded as somewhat in-between these two classifications. For purposes of this paper, collisions are classified as bouncing cases if only a single 'satellite' particle is separated from the two main agglomerates, and they are classified as fragmentation

cases if the satellite agglomerate consists of two or more particles. More typical fragmentation cases are similar to that shown in Figure 11, however, producing at least three large agglomerates and sometimes also several smaller agglomerates.

The question of whether a given collision will be of the merger, bouncing or fragmentation type depends primarily on the values of the adhesion parameter Ad and the ratio of the y -direction offset distance D_a to the initial radius of gyration R_{g0} of the two agglomerates. A plot identifying the type of collision for all computations conducted is shown in a mapping of Ad versus D_a / R_{g0} in Figure 12, and details of the number of particles in each agglomerate following collision are listed in Table 2. Regions of the map in Figure 12 are marked to provide a rough identification of values of Ad and D_a / R_{g0} for which the agglomerates individually break up in the shear flow (to the far left of the plot) and values resulting in merger, bouncing and fragmentation type collisions. The dashed line separating the merger and bouncing regimes is given by the line $D_a / R_{g0} = 0.45 + 0.0002Ad$. The numbers indicate the number of agglomerates present at the conclusion of the computation, where an agglomerate is defined as a group of two or more touching particles. In general, collisions resulting in mergers occurred for smaller values of dimensionless offset distance D_a / R_{g0} and values of Ad well above the critical value for breakup of the individual agglomerate in shear flow. Bouncing collisions occur for larger values of D_a / R_{g0} , resulting in glancing collisions of the agglomerates. Fragmentation occurs for moderate values of D_a / R_{g0} with adhesion parameter values just slightly larger than the critical value for breakup of a single agglomerate in the shear flow. Two cases in Figure 12 requiring special discussion are indicated with asterisks. One of these cases, indicated by 2^* , was identified as a bouncing collision because it resulted in

two agglomerates, but a much larger number of particles were exchanged between the two agglomerates than was the case for other bouncing collision cases. Indeed, 172 particles originating in the red agglomerate, out of an initial 577 particles, were torn off and captured by the blue agglomerate during the collision. The case indicated by 4* in Figure 12 was, on the other hand, a fairly typical fragmentation case, resulting in three fairly large agglomerates with 263, 315 and 69 particles and one smaller 'satellite' agglomerate with 8 particles. The presence of this fragmentation case in a region where we otherwise see a lot of bouncing cases is a reminder that each agglomerate has its own unique structure and each collision involves different parts of these unique agglomerates, so one must expect substantial variation from case to case. The plot in Figure 12 should therefore be regarded as providing only a rough indication of the conditions under which different types of collisions occur and not as a strict regime map.

For the problem of collision of two particles, the criterion for sticking or bouncing of the particles can usually be expressed as a critical value of the particle Stokes number St , which is a function of the adhesion parameter Ad . Applying this same idea for the problem of agglomerate collision, we can define an agglomerate Stokes number St_A as the ratio of an agglomerate time scale $\tau_A = m_A / 3\pi\mu R_{g0}$ and the fluid time scale $\tau_f = L / U = R_{g0} / SD_a$, giving

$$St_A = \frac{m_A S}{3\pi\mu R_{g0}} \frac{D_a}{R_{g0}}. \quad (28)$$

In this expression, m_A is the agglomerate mass, $U = SD_a$ is the characteristic velocity difference between the agglomerates, and R_{g0} is an agglomerate length scale. The agglomerate Stokes number is therefore found to vary linearly with the ratio D_a / R_{g0} used in Figure 12.

While we have used the term *bouncing collision* to be in conformity with terminology used in previous literature (e.g., Brisset et al., 2016), it is clear that the bouncing agglomerate collisions for the loosely-structured agglomerates examined in the current study differ substantially from the traditional bouncing collision of two elastic particles. In a traditional bouncing process, two colliding elastic bodies deform locally near the collision point, resulting in an elastic (or sometimes plastic) repulsion force pushing the two bodies away from each other. In a bouncing case, this repulsion force is sufficiently strong to overcome the adhesive force between the bodies, so that the two bodies will detach and continue to move away from each other. The bouncing collisions of two loosely-structured agglomerates observed in the current paper are characterized more by tearing away and eventual capture of particles from the opposing agglomerate by the particle adhesion force. It is not that the elastic force between the agglomerates overcomes the adhesive force between the bodies, but rather that the adhesion force imposed on the captured particles by one agglomerate overcomes the adhesion force from the agglomerate to which the captured particles were originally attached. A plot showing number of captured particles from both agglomerates during the different bouncing collisions computed is given in Figure 13. As we see from this plot, all bouncing collisions included captured particles. In some cases only one agglomerate captures particles, and in other cases both colliding agglomerates capture particles from the other agglomerate.

While exchange of particles was a characteristic feature of all bouncing collisions, this is not to say that there was no rebound force between the agglomerates. An examination of the rebound force is reported below for the bouncing collision in Case B.19, in which 28 particles originating in the red agglomerate are captured by the blue agglomerate and one blue particle is captured by the red agglomerate. The number of touching red-blue particles (i.e., touching

particles originating from opposite agglomerates) is plotted as a function of time in Figure 14a. This number is zero until $t = 5$, at which time the collision occurs, and then suddenly spikes up to a peak value of 18 at a time of about $t = 6.5$. After this point the number of touching red-blue particles decreases to 14 and remains there, with the exception of a small blip at $t = 10$ due to restructuring. The fact that the number of red-blue touching particles does not reduce to zero following the collision is due to the presence of captured particles. The total compressive force between the two agglomerates (which is characteristic of the elastic rebound force) is plotted as a function of time in Figure 14b. We again observe a sudden increase at collision onset at $t = 5$ and a peak value at $t = 6.5$, followed by a gradual decrease of the compression force as the two agglomerates tear away from each other.

The position of particles carrying the compressive load between the two colliding agglomerates is illustrated in Figure 15 at a time of $t = 7$, close to the peak time of the collision. In Figure 15a, we color the particle scatter plot with red or blue to identify the originating agglomerate for each particle. In Figure 15b, each particle is colored by the magnitude of the total compressive force acting on the particle. The highest compressive loads are borne by a core of particles on the inside of the agglomerate, shown in Figure 15c with the lower-compression particles removed, within a tube of force chains radiating outward from the collision point. The highest compressive load occurs on the particles just at the collision point, indicated by red or orange in Figure 15c. We have thus confirmed that a rebound force does occur in bouncing collisions, and it may be reasonable to characterize this aspect of the collision phenomenon by some type of effective elastic modulus assigned to an effective spherical body representing the agglomerate. However, this effective sphere representation does not include the important

phenomenon of particle capture during bouncing collisions, which in most of the cases that we have examined is very important to the agglomerate behavior during collision.

In Section 3, we discussed the observation that the particle-induced flow field from a single agglomerate in a shear flow has the form of two tilted vortex rings of opposite sign. In the event of a collision of two agglomerates, one naturally wonders what happens to the particle-induced flow during the collision. To examine this question, an iso-surface of the relative vorticity magnitude ω_{rel} is plotted at four different times during a collision resulting in merger (Figure 16 for Case B.15) and during a collision resulting in bouncing (Figure 17 for Case B.19). The relative-vorticity iso-surface for fragmentation cases depends on the number of fragments produced, and so are highly variable. The agglomerate centroids and initial radius of gyration are indicated in these figures by a black dot and a circle, respectively, for each agglomerate. Below each figure is given a scatter plot showing the particle positions at that time, with color used to identify the agglomerate of origin for each particle.

In Figure 16, the particle-induced flow field at time $t = 6$ (just before the collision) has the form of two opposite-sign tilted vortex rings for each agglomerate, hence four tilted vortex rings in all. At time $t = 8$ the agglomerates are in the midst of colliding and the innermost vortex rings of each agglomerate collide with one another. At $t = 10$, the inner vortex rings have significantly decayed while the outer vortex rings have grown in strength. The inner rings continue to break up and be swept downstream by $t = 12$, leaving the two strong outer vortex rings, which have opposite sign from each other. With the exception of the small-scale remnants of the inner rings, the particle-induced flow for the merged agglomerates at $t = 12$ thus appears similar to that for a single agglomerate in a shear flow, as discussed in the previous section, but the vortex rings are larger and stronger for the merged agglomerate.

In Figure 17, a time series of iso-relative vorticity magnitude surfaces are plotted for a case with bouncing agglomerate collision. The first two images in Figure 17 appear similar to those in Figure 16 for a merging collision. The two inner rings collide at time $t = 6$ and nearly extinguish each other by time $t = 8$ as the agglomerate collision occurs. However, as the agglomerates bounce and move away from each other, the inner rings reform, such that by $t = 12$ we see a pair of vortex rings for each agglomerate moving away from each other. A trail of vorticity connects these two vortex ring pairs, which is either left over from the collision or generated by stretching of the background shear vorticity.

5. Conclusions

A computational study was reported examining rotation and breakup of a single particle agglomerate and collision of two particle agglomerates in a shear flow. The agglomerates are extracted from a direct numerical simulation of turbulent agglomeration, and therefore have the characteristic loose fractal structure typical of turbulent agglomeration processes. Computations are performed with four-way coupling between the particles and the fluid and with sufficient resolution of the agglomerates to capture the details of the particle-induced flow field. Simulations of a single agglomerate rotating in the shear flow with high values of the adhesion parameter indicate that the agglomerate rotates more slowly than would an ambient fluid element in the shear flow. The flow field induced by the particles of a rotating agglomerate in a shear flow are found to exhibit a very distinctive form, characterized by a pair of tilted vortex rings with opposite sign circulation, surrounded by a sea of stretched vorticity from the ambient shear flow. To our knowledge, this is the first time that the particle-induced flow of an agglomerate in shear flow has been examined in detail and the first time that the interesting vortex ring pair

structure of this flow has been described. This vortex pair ring structure remains with constant orientation and strength as the particle agglomerate rotates. For sufficiently low values of the adhesion parameter, the agglomerate is observed to break up in the shear flow, where the exact value of adhesion parameter at breakup varies slightly with the specific choice of the agglomerate under examination.

The problem of collision of two agglomerates was found to result in either merger, bouncing or fragmentation, depending on the value of the adhesion parameter and the ratio of offset distance to agglomerate radius of gyration. In merger collisions, the inner vortex rings of the particle-induced flow from each agglomerate interact with each other and eventually break up into small scale structures, and the outer vortex rings grow stronger leading to development of the vortex ring pair structure typical of that observed for a single agglomerate. It was observed that bouncing collisions result both in repulsive force between the agglomerates due to elastic deformation as well as exchange of particles between agglomerates. The innermost vortex ring structures of the particle-induced flow for bouncing collisions similarly exhibit interaction of the two inner vortex rings, but these inner rings are found to quickly reform as the agglomerates bounce and move away from each other. Fragmentation collisions may result in three or more agglomerates with widely different sizes, many of which are formed of a combination of particles originating in different agglomerate structures.

Many theoretical and computation models of turbulent agglomeration processes make use of the common approximation that an agglomerate can be replaced by an 'effective particle', in which some effective elastic modulus of the agglomerate is assigned. The current study demonstrates that this effective particle approximation omits certain important physical phenomena associated with agglomerate collision, including fragmentation collisions (resulting

in three or more agglomerates) and exchange of particles between agglomerates in bouncing collisions. The particle-induced flow field is also quite different for a loosely-structured agglomerate than it is for an equivalent sphere due to the fact that the fluid flow can penetrate into the outer parts of the agglomerate. This penetration affects the rotation rate of an agglomerate in a shear flow and gives rise to the tilted vortex ring structure of the particle-induced flow.

The current study suggests the need for future work in a number of areas. The current study has focused on collision of particle agglomerates with exact copies of themselves, using a relatively small number of agglomerates in order to focus on the effect of the adhesion number and the spacing ratio D_a / R_{g0} . There is a need to conduct runs with larger number of agglomerates, including cases involving collisions of agglomerates of different sizes. The current study considered agglomerates suspended in a simple shear flow, whereas in practice agglomerates will also experience a mean drift relative to the surrounding flow, either from inertia or from a body force such as gravity. An agglomerate falling under gravity in a fluid is known to induce a single vortex ring within the agglomerate (Nitsche & Batchelor, 1997), in contrast to the pair of tilted vortex rings that we have observed to be induced by an agglomerate in a shear flow. The particle-induced flow for an agglomerate experiencing a combination of shear and mean drift would therefore be of interest for future study. Finally, resolution of the flow within the agglomerate using the CFD-DEM approach employed in the current paper requires that the agglomerate size is significantly larger than the particle size. It would be of interest to examine fluid flow effects and collision of smaller agglomerates, for which a computational method capable of resolving flow around individual particles would be necessary.

Acknowledgements

This research was supported by the U.S. National Science Foundation under grants CBET-1332472 and DGE-1144388.

References

- Adachi, Y. & Ooi, S. 1990 Geometrical structure of a floc. *Journal of Colloid and Interface Science* **135** (2), 374-384.
- Akiki, G., Moore, W. C. & Balachandar, S. 2017a Pairwise-interaction extended point - particle model for particle-laden flows. *Journal of Computational Physics* **351**, 329-357.
- Akiki, G., Jackson, T.L. & Balachandar, S. 2017b Pairwise interaction extended point-particle model for a random array of monodisperse spheres. *Journal of Fluid Mechanics* **813**, 882-928.
- Bagi, K. & Kuhn, M.R. 2004 A definition of particle rolling in a granular assembly in terms of particle translations and rotations. *Journal of Applied Mechanics* **71**, 493-501.
- Balakin, B., Hoffmann, A. C. & Kosinski, P. 2011 The collision efficiency in a shear flow. *Chemical Engineering Science* **68**, 305-312.
- Becker, V., Schlauch, E., Behr, M. & Briesen, H. 2009 Restructuring of colloidal aggregates in shear flows and limitations of the free-draining approximation. *Journal of Colloid and Interface Science* **339**, 362-372.
- Beitz, E., Güttler, C., Blum, J., Meisner, T., Teiser, J. & Wurm, G. 2011 Low-velocity collisions of centimeter-sized dust aggregates. *The Astrophysical Journal* **736** (1), 34.
- Bosse, T., Kleiser, L., Härtel, C. & Meiburg, E. 2005 Numerical simulation of finite Reynolds number suspension drops settling under gravity. *Physics of Fluids* **17**, 037101.
- Brasil, A. M., Farias, T. L., Carvalho, M. G. & Koçlu, U. O. 2001 Numerical characterization of the morphology of aggregated particles. *Journal of Aerosol Science* **32**, 489-508.
- Brisset, J., Heißelmann, D., Kothe, S., Weidling, R. & Blum, J. 2016 Submillimetre-sized dust aggregate collision and growth properties: Experimental study of a multi-particle system on a suborbital rocket. *Astronomy & Astrophysics* **593**, A3.
- Brunk, B. K., Koch, D. L. & Lion, L. W. 1998 Turbulent coagulation of colloidal particles. *Journal of Fluid Mechanics* **364**, 81-113.
- Chokshi, A., Tielens, A. G. G. M. & Hollenbach, D. 1993 Dust coagulation. *The Astrophysical Journal* **407**, 806-819.
- Chun, J. & Koch, D. L. 2005 Coagulation of monodisperse aerosol particles by isotropic turbulence. *Physics of Fluids* **17**, 027102.
- Cleary, P. W., Metcalfe, G. & Liffman, K. 1998 How well do discrete element granular flow models capture the essentials of mixing processes? *Applied Mathematical Modeling* **22**, 995-1008.

Crowe, C. T., Schwarzkopf, J. D., Sommerfeld, M. & Tsuji, Y. 2012 *Multiphase Flows with Droplets and Particles*, 2nd ed., CRC Press, Boca Raton, Florida.

Di Felice, R. 1994 The voidage function for fluid-particle interaction systems. *International Journal of Multiphase Flow* **20**, 153-159.

Ding, W., Zhang, H. & Cetinkaya, C. 2008 Rolling resistance moment-based adhesion characterization of microspheres. *Journal of Adhesion* **84**, 996-1006.

Dizaji, F. F. & Marshall, J. S. 2016 An accelerated stochastic vortex structure method for particle collision and agglomeration in homogeneous turbulence. *Physics of Fluids* **28**, 113301.

Dizaji, F. F. & Marshall, J. S. 2017 On the significance of two-way coupling in simulation of turbulent particle agglomeration. *Powder Technology* **318**, 83-94.

Dominik, C. & Tielens, A. G. G. M. 1995 Resistance to rolling in the adhesive contact of two elastic spheres. *Philosophical Magazine A* **92** (3), 783-803.

Eggersdorfer, M. L., Kadau, D., Hermann, H. J. & Pratsinis, S. E. 2011 Multiparticle sintering dynamics: From fractal-like aggregates to compact structures. *Langmuir* **27**, 6358-6367.

Fanelli, M., Feke, D. L. & Manas-Zloczower, I. 2006 Prediction of the dispersion of particle clusters in the nano-scale – Part I: Steady shearing responses. *Chemical Engineering Science* **61**, 473-488.

Gunkelmann, N., Ringl, C. & Urbassek, H. M. 2016 Influence of porosity on collisions between dust aggregates. *Astronomy & Astrophysics* **589**, A30.

Hansen, S., Khakhars, D. V. & Ottino, J.M. 1998 Dispersion of solids in nonhomogeneous viscous flows. *Chemical Engineering Science* **53**(10), 1803-1817.

Hatakeyama, N. & Kambe, T. 1997 Statistical laws of random strained vortices in turbulence. *Physical Review Letters* **79** (7), 1257-1260.

Higashitani, K., Iimura, K. & Sanda, H. 2001 Simulation of deformation and breakup of large aggregates in flows of viscous fluids. *Chemical Engineering Science* **56**, 2927-2938.

Ihalainen, M., Lind, T., Torvela, T., Lehtinen, K. E. J. & Jokiniemi, J. 2012 A method to study agglomerate breakup and bounce during impaction, *Aerosol Science and Technology* **46** (9), 990-1001.

Iimura, K., Suzuki, M., Hirota, M. & Higashitani, K. 2009a Simulation of dispersion of agglomerates in gas phase - acceleration field and impact on cylindrical obstacle. *Advanced Powder Technology* **20**, 210-215.

- Iimura, K., Yanagiuchi, M., Suzuki, M., Hirota, M. & Higashitani, K. 2009b Simulation of dispersion and collection process of agglomerated particles in collision with fibers using discrete element method. *Advanced Powder Technology* **20**, 582-587.
- Jiang, Q. & Logan, B. E. 1991 Fractal dimensions of aggregates determined from steady-state size distributions. *Environmental Science and Technology* **25**, 2031-2038.
- Johnson, K. L., Kendall, K. & Roberts, A. D. 1971 Surface energy and the contact of elastic solids. *Proceedings of the Royal Society of London A* **324**, 301-313.
- Joseph, G. G., Zenit, R., Hunt, M. L. & Rosenwinkel, A. M. 2001 Particle-wall collisions in a viscous fluid. *Journal of Fluid Mechanics* **433**, 329-346.
- Kafui, K. D. & Thornton, C. 2000 Numerical simulations of impact breakage of a spherical crystalline agglomerate. *Powder Technology* **109**, 113-132.
- Koch, D. L. & Pope, S. B. 2002 Coagulation-induced particle-concentration fluctuations in homogeneous, isotropic turbulence. *Physics of Fluids* **14**, 2447-2455.
- Kosinski, P. & Hoffmann, A. C. 2010 An extension of the hard-sphere particle-particle collision model to study agglomeration. *Chemical Engineering Science* **65** (10), 3231-3239.
- Kun, F. & Herrmann, H. J. 1999 Transition from damage to fragmentation in collision of solids. *Physical Review E* **59** (3), 2623-2632.
- Kusters, K. A., Wijers, J. G. & Thoenes, D. 1997 Aggregated kinetics of small particles in agitates vessels. *Chemical Engineering Science* **52** (1), 107-121.
- Lele, S. K. 1992 Compact finite difference schemes with spectral-like resolution. *Journal of Computational Physics* **103** (1), 16-42.
- Li, C., Ye, M. & Liu, Z. 2016 On the rotation of a circular porous particle in 2D simple shear flow with fluid inertia. *Journal of Fluid Mechanics* **808**, R3.
- Lian, G., Thornton, C. & Adams, M. J. 1998 Discrete particle simulation of agglomerate impact coalescence. *Chemical Engineering Science* **53** (19), 3381-3391.
- Lu, J. & Wang, J. K. 2006 Agglomeration, breakage, population balance, and crystallization kinetics of reactive precipitation process. *Chemical Engineering Communications* **193**, 891-902.
- Marangoni, A. G. & Narine, S. S. 2001 Elasticity of fractal aggregate networks: Mechanical arguments. In *Crystallization and Solidification Properties of Lipids*, Widlak, N., Hartel, R.W., and Narine, S., eds., The American Oil Chemists Society, Champaign, IL.
- Marshall, J. S. 2009 Discrete-element modeling of particulate aerosol flows. *Journal of Computational Physics* **228**, 1541-1561.

Marshall, J. S. & Li, S. 2014 *Adhesive Particle Flow: A Discrete Element Approach*, Cambridge University Press, New York.

Marshall, J. S. & Sala, K. 2013 Comparison of methods for computing the concentration field of a particulate flow. *International Journal of Multiphase Flow* **56**, 4-14.

Mindlin, R. D. 1949 Compliance of elastic bodies in contact. *Journal of Applied Mechanics* **16**, 259-268.

Moreno, R., Ghadiri, M. & Antony, S. J. 2003 Effect of the impact angle on the breakage of agglomerates: a numerical study using DEM. *Powder Technology* **130**, 132-137.

Moreno-Atanasio, R. & Ghadiri, M. 2006 Mechanistic analysis and computer simulation of impact breakage of agglomerates: Effect of surface energy. *Chemical Engineering Science* **61**, 2476-2481.

Nguyen, D., Rasmuson, A., Thalberg, K. & Björn, I. N. 2014 Numerical modelling of breakage and adhesion of loose fine-particle agglomerates. *Chemical Engineering Science* **116**, 91-98.

Ning, Z., Boerefijn, R., Ghadiri, M. & Thornton, C. 1997 Distinct element simulation of impact breakage of lactose agglomerates. *Advanced Powder Technology* **8** (1), 15-37.

Nitsche, J.M. & Batchelor, G.K. 1997 Break-up of a falling drop containing dispersed particles. *Journal of Fluid Mechanics* **340**, 161-175.

Olfert, J. S., Symonds, J. P. R. & Collings, N. 2007 The effective density and fractal dimension of particles emitted from a light-duty vehicle with a diesel oxidation catalyst. *Journal of Aerosol Science* **38**, 69-82.

Ormel, C. W., Paszun, D., Dominik, C. & Tielens, A. G. G. M. 2009 Dust coagulation and fragmentation in molecular clouds: I. How collisions between dust aggregates alter the dust size distribution. *Astronomy & Astrophysics* **502**, 845–869.

Ormel, C. W., Spaans, M. & Tielens, A. G. G. M. 2007 Dust coagulation in protoplanetary disks: porosity matters. *Astronomy & Astrophysics* **461**, 215-232.

Potatin, A. A. 1993 On the computer simulation of the deformation and breakup of colloidal aggregates in shear flow. *Journal of Colloid and Interface Science* **157**, 399-410.

Rai, M. & Moin, P. 1991 Direct simulation of turbulent flow using finite-difference schemes. *Journal of Computational Physics* **96**, 15–53.

Reinhold, A. & Briesen, H. 2012 Numerical behavior of a multiscale aggregation model - coupling population balances and discrete element models. *Chemical Engineering Science* **70**, 165-175.

- Rubinow, S. I. & Keller, J. B. 1961 The transverse force on a spinning sphere moving in a viscous fluid. *Journal of Fluid Mechanics* **11**, 447-459.
- Rumpf, H. 1962 The strength of granules and agglomerates. In: W.A. Knepper (Ed.) *Agglomeration*, John Wiley, New York, pp. 379-418.
- Saffman, P. G. 1965 The lift on a small sphere in a slow shear flow. *Journal of Fluid Mechanics* **22**, 385-400.
- Saffman, P. G. & Turner, J. S. 1956 On the collision of drops in turbulent clouds. *Journal of Fluid Mechanics* **1** (1), 16-30.
- Samimi, A., Moreno, R. & Ghadiri, M. 2004 Analysis of impact damage of agglomerates: effect of impact angle. *Powder Technology* **143-144**, 97-109.
- Sayvet, O. & Navard, P. 2000 Collision-induced dispersion of agglomerate suspensions in a shear flow. *Journal of Applied Polymer Science* **78**, 1130–1133.
- Schäfer, C., Speith, R. & Kley, W. 2007 Collisions between equal-sized ice grain agglomerates. *Astronomy and Astrophysics* **470**, 733-739.
- Schiller, L. & Naumann, A. 1933 Über die grundlegenden Berechnungen bei der Schwerkraftaufbereitung. *Zeitschrift des Vereines Deutscher Ingenieure* **77**, 318-320.
- Selomulya, C., Amal, R., Bushell, G. & Waite, T. D. 2001 Evidence of shear rate dependence on restructuring and breakup of latex aggregates. *Journal of Colloid and Interface Science* **236**, 67-77.
- Seizinger, A. & Kley, W. 2013 Bouncing behavior of microscopic dust aggregates. *Astronomy and Astrophysics* **551**, A65.
- Serra, T., Colomer, J. & Casamitjana, X. 1997 Aggregation and breakup of particles in a shear flow. *Journal of Colloid and Interface Science* **187**, 466-473.
- Smoluchowski, M. 1917 Versuch einer mathematischen Theorie der Koagulationskinetik kolloider Lösungen. *Zeitschrift für Physikalische Chemie* **92**, 129–168.
- Sonntag, R. C. & Russel, W. B. 1986 Structure and breakup of flocs subjected to fluid stresses. I. Shear experiments. *Journal of Colloid and Interface Science* **113**(2), 399-413.
- Steijl, R. 2001 Computational study of vortex pair dynamics. Ph.D. dissertation, University of Twente, Enschede, The Netherlands, p. 64.
- Thornton, C. 1991 Interparticle sliding in the presence of adhesion. *Journal Physics D: Applied Physics* **24**, 1942-1946.

Thornton, C., Ciomocos, M. T. & Adams, M. J. 1999 Numerical simulations of agglomerate impact breakage. *Powder Technology* **105**, 74–82.

Thornton, C. & Liu, L. 2004 How do agglomerates break? *Powder Technology* **143–144**, 110–116.

Tong, Z. B., Yang, R. Y., Chu, K. W., Yu, A. B., Adi, S. & Chan, H. K. 2010 Numerical study of the effects of particle size and polydispersity on the agglomerate dispersion in a cyclonic flow. *Chemical Engineering Journal* **164**, 432–441.

Tong, Z. B., Yang, R. Y., Yu, A. B., Adi, S. & Chan, H. K. 2009 Numerical modelling of the breakage of loose agglomerates of fine particles. *Powder Technology* **196**, 213–221.

Tong, Z. B., Zheng, B., Yang, R. Y., Yu, A. B. & Chan, H. K. 2013 CFD-DEM investigation of the dispersion mechanisms in commercial dry powder inhalers. *Powder Technology* **240**, 19–24.

Tong, Z. B., Zhong, W., Yu, A. B., Chan, H. K. & Yang, R. Y. 2016 CFD-DEM investigation of the effect of agglomerate–agglomerate collision on dry powder aerosolisation. *Journal of Aerosol Science* **92**, 109–121.

Tsuji, Y., Tanaka, T. & Ishida, T. 1992 Lagrangian numerical simulation of plug flow of cohesionless particles in a horizontal pipe. *Powder Technology* **71**, 239–250.

Uhlmann, M. 2005 An immersed boundary method with direct forcing for the simulation of particulate flows. *Journal of Computational Physics* **209**, 448–476.

Verzicco, R. & Orlandi, P. 1996 A finite-difference scheme for three-dimensional incompressible flows in cylindrical coordinates. *Journal of Computational Physics* **123**, 402–414.

Waldner, M. H., Sefcik, J., Soos, M. & Morbidelli, M. 2005 Initial growth kinetics of aggregates in turbulent coagulator. *Powder Technology* **156**, 226–234.

Wang, L. P., Wexler, A. S. & Zhou, Y. 1998 Statistical mechanical descriptions of turbulent coagulation. *Physics of Fluids* **10**, 2647–2651.

Wen, C. Y. & Yu, Y. H. 1966 Mechanics of fluidization. *Chemical Engineering Progress Symposium Series* **62** (62), 100–111.

Yang, J., Wu, C. Y. & Adams, M. 2014 Three-dimensional DEM-CFD analysis of air-flow-induced detachment of API particles from carrier particles in dry powder inhalers. *Acta Pharmaceutica Sinica B* **4** (1), 52–59.

Zeidan, M., Xu, B. H., Jia, X. & Williams, R. A. 2007 Simulation of aggregate deformation and breakup in simple shear flows using a combined continuum and discrete model. *Chemical Engineering Research and Design* **85** (A12), 1645–1654.

Zhu, H. P., Zhou, Z. Y., Yang, R. Y. & Yu, A. B. 2007 Discrete particle simulation of particulate systems: theoretical developments. *Chemical Engineering Science* **62**, 3378-3396.

Tables

Table 1. Listing of parameter values for cases examined with a single agglomerate in a shear flow, including adhesion parameter, initial number of particles, and ratio of initial gyration radius to particle diameter. For all cases examined $St = 1.4$ and $\rho_p / \rho_f = 10$.

Case Number	Ad	N_0	R_{g0} / d
A.1	133	328	4.81
A.2	333	328	4.81
A.3	666	328	4.81
A.4	999	328	4.81
A.5	146	269	4.40
A.6	364	269	4.40
A.7	728	269	4.40
A.8	1092	269	4.40
A.9	104	577	6.17
A.10	259	577	6.17
A.11	518	577	6.17
A.12	778	577	6.17

Table 2. Listing of parameter values for cases examined for collision of two agglomerates, including adhesion parameter, initial numbers of particles in each agglomerate (N), ratio of initial gyration radius (R_{g0}) of each agglomerate to particle diameter d , and ratio of initial offset distance D_a to R_{g0} . For each case examined $St = 1.4$ and $\rho_p / \rho_f = 10$. Also listed was the observed type of collision - merger (M), bouncing (B) or fragmentation (F) – and the number of particles in each remaining agglomerate (Aggl 1-5) after the collision.

Case Number	Ad	N_0	R_{g0} / d	D_a / R_{g0}	Collision Type	Aggl 1	Aggl 2	Aggl 3	Aggl 4	Aggl 5
B.1	333	328	4.81	0.52	F	213	392	51	-	-
B.2	666	328	4.81	0.52	M	656	-	-	-	-
B.3	999	328	4.81	0.52	M	656	-	-	-	-
B.4	1998	328	4.81	0.52	M	656	-	-	-	-
B.5	333	328	4.81	0.78	F	338	168	8	5	3
B.6	666	328	4.81	0.78	F	331	317	8	-	-
B.7	999	328	4.81	0.78	F	276	380	-	-	-
B.8	1998	328	4.81	0.78	M	656	-	-	-	-
B.9	333	328	4.81	1.04	F	262	315	69	8	-
B.10	666	328	4.81	1.04	B	350	305	-	-	-
B.11	999	328	4.81	1.04	B	358	298	-	-	-
B.12	1998	328	4.81	1.04	B	326	330	-	-	-
B.13	364	269	4.40	0.57	F	326	161	51	-	-
B.14	728	269	4.40	0.57	M	538	-	-	-	-
B.15	1092	269	4.40	0.57	M	538	-	-	-	-
B.16	2184	269	4.40	0.57	M	538	-	-	-	-
B.17	364	269	4.40	0.85	B	291	247	-	-	-
B.18	728	269	4.40	0.85	B	286	252	-	-	-
B.19	1092	269	4.40	0.85	B	296	242	-	-	-
B.20	2184	269	4.40	0.85	M	538	-	-	-	-
B.21	364	269	4.40	1.14	B	268	270	-	-	-
B.22	728	269	4.40	1.14	B	268	270	-	-	-
B.23	1092	269	4.40	1.14	B	268	270	-	-	-
B.24	2184	269	4.40	1.14	B	268	270	-	-	-
B.25	778	577	6.17	0.41	B	749	405	-	-	-
B.26	1556	577	6.17	0.41	M	1154	-	-	-	-
B.27	778	577	6.17	0.61	F	171	619	364	-	-
B.28	1556	577	6.17	0.61	M	1154	-	-	-	-
B.29	778	577	6.17	0.81	B	579	575	-	-	-
B.30	1556	577	6.17	0.81	B	607	547	-	-	-

Figure Captions

Figure 1. Plot of the number of particles in an agglomerate N versus the ratio of the radius and gyration of the agglomerate R_g and the individual particle radius r_p . The slope of the plot indicates the dimension $d_f = 2.12$ of the power law in Eq. (17).

Figure 2. Schematic diagram of the initial conditions for the problems of (a) a single agglomerate in a shear flow and (b) two-agglomerate collision in a shear flow. Circles indicate the radius of gyration R_g , and the offset distance D_a is indicated in (b) in both positive and negative directions.

Figure 3. Particle positions at times (a) $t = 0$, (b) 10, (c) 20, and (d) 30 for Case A.4. The particles are colored by the magnitude of the relative velocity vector. The agglomerate is rotating clockwise in the shear flow and completes approximately one rotation in the time interval shown.

Figure 4. Plot showing the time-variation of the gyration radius R_g (solid line, left-hand axis) and the particle concentration within the agglomerate c_{agg} (dashed line, right-hand axis) for Case A.4.

Figure 5. Plot showing the time-variation of (a) the average value of the magnitude of the particle velocity $\mathbf{v}$ (dashed line) and the relative particle velocity vector $\mathbf{w} \equiv \mathbf{v} - \mathbf{u}$ (solid line) and (b) the second-moment measure for particle coordination number (black line), relative rotation rate about the agglomerate centroid (blue line), and relative velocity magnitude (red line) for Case A.4.

Figure 6. (a) Contour plot of the x -component u_{rel} of the relative velocity in the x - y plane, for Case A.4 at $t = 26$. (b) Profile of the x -component of velocity u along the y -axis. The solid line denotes the ambient shear flow and the dots denote the computed velocity profile.

Figure 7. (Left) contours of normal vorticity and streamlines of the in-plane velocity field and (right) contours of normal component of the relative velocity $\mathbf{u}_{rel}$ in three orthogonal planes passing through the agglomerate, for Case A.4 at $t = 26$.

Figure 8. Iso-surface of the relative vorticity magnitude $\omega_{rel} = 0.46$ obtained from the velocity field for Case A.4 at $t = 26$, showing two tilted vortex rings generated by the particle-induced velocity field near the rotating agglomerate. The top two plots show iso-surfaces in the (a) x - y plane and (b) x - z plane. The bottom two plots, (c) and (d), show the same iso-surface views together with a slice showing ω_{rel} contours in the normal plane.

Figure 9. Time series showing breakup of single agglomerate in a shear flow, for Case A.1 at times (a) $t = 0$, (b) 5, (c) 10, (d) 15 and (e) 20.

Figure 10. Plots showing measures characterizing breakup of a single agglomerate in a shear flow. (a) Number of fragments that an agglomerate breaks up into versus adhesion parameter. When the agglomerate does not break up, $N_{frag} = 1$. (b) Number of particles N in agglomerates following breakup versus adhesion parameter. The dashed line is the experimental power-law fit $N \propto Ad^{0.879}$ from Sonntag and Russel (1986) for maximum number of particles, where the proportionality coefficient is fit to the data. The data is plotted for Cases A.1-A.4 (red deltas), A.5-A.8 (green circles), and A.9-A.12 (blue diamonds) from Table 1.

Figure 11. Scatter plots illustrating three types of agglomerate interactions: merger (Case B.15), bouncing (Case B.19) and fragmentation (Case B.13).

Figure 12. Summary of results for all agglomerate collision runs, showing the number of agglomerates (N_{agg}) remaining after collision as a function of adhesion parameter and the ratio $D_a / R_{g,ave}$ of offset distance to initial radius of gyration. Colors indicate results from different agglomerates. Numbers indicate cases with agglomerate merger ($N_{agg} = 1$), bounce ($N_{agg} = 2$), and fragmentation ($N_{agg} > 2$).

Figure 13. Plot indicating the number of captured particles in bouncing collisions versus the total number of particles in an agglomerate. The number of red particles captured by blue agglomerates is plotted in red, and the number of blue particles captured by red agglomerates is plotted in blue. Different symbols are used to indicate different computations, with one red and one blue symbol for each computation.

Figure 14. Time variation of (a) number of touching particles originating in different agglomerates and (b) total dimensionless compressive force between the agglomerates for a typical bouncing case (Case B.19). Collision onset occurs at approximately $t = 5$ and the agglomerates detach at $t = 14$.

Figure 15. Scatter plots during a bouncing agglomerate collision (Case B.19) at $t = 7$, with colors indicating (a) agglomerate from which each particle originated, (b) total compressive force acting on each particle, and (c) same plot as in (b) with the low-compression particles (with compressive force < 1.5) blanked out. High compression force chains occur in a particle core region spreading outward from the collision point.

Figure 16. Iso-surface of relative vorticity magnitude $\omega_{rel} = 0.3$ (top) and particle scatter plot colored by initial agglomerate identity (bottom) for a case where the particle agglomerates merge (Case B.15), at times (a) $t = 6$, (b) 8, (c) 10, and (d) 12 during which collision and merger of the agglomerates occurs. The agglomerate centroids and initial radius of gyration are indicated in the upper plots by a black dot and a circle, respectively, for each agglomerate.

Figure 17. Iso-surface of relative vorticity magnitude $\omega_{rel} = 0.3$ (top) and particle scatter plot colored by initial agglomerate identity (bottom) for a case where the particle agglomerates bounce (Case B.19), at times (a) $t = 6$, (b) 8, (c) 10, and (d) 12 during which collision of the

agglomerates occurs. The agglomerate centroids and initial radius of gyration are indicated in the upper plots by a black dot and a circle, respectively, for each agglomerate.




Figure 1. Plot of the number of particles in an agglomerate N versus the ratio of the radius and gyration of the agglomerate R_g and the individual particle radius r_p . The slope of the plot indicates the dimension $d_f = 2.12$ of the power law in Eq. (17).




Figure 2. Schematic diagram of the initial conditions for the problems of (a) a single agglomerate in a shear flow and (b) two-agglomerate collision in a shear flow. Circles indicate the radius of gyration R_g , and the offset distance D_a is indicated in (b) in both positive and negative directions.




Figure 3. Particle positions at times (a) $t = 0$, (b) 10, (c) 20, and (d) 30 for Case A.4. The particles are colored by the magnitude of the relative velocity vector. The agglomerate is rotating clockwise in the shear flow and completes approximately one rotation in the time interval shown.




Figure 4. Plot showing the time-variation of the gyration radius R_g (solid line, left-hand axis) and the particle concentration within the agglomerate c_{agg} (dashed line, right-hand axis) for Case A.4.




Figure 5. Plot showing the time-variation of (a) the average value of the magnitude of the particle velocity $\mathbf{v}$ (dashed line) and the relative particle velocity vector $\mathbf{w} \equiv \mathbf{v} - \mathbf{u}$ (solid line) and (b) the second-moment measure for particle coordination number (black line), relative rotation rate about the agglomerate centroid (blue line), and relative velocity magnitude (red line) for Case A.4.




Figure 6. (a) Contour plot of the x -component u_{rel} of the relative velocity in the x - y plane, for Case A.4 at $t = 26$. (b) Profile of the x -component of velocity u along the y -axis. The solid line denotes the ambient shear flow and the dots denote the computed velocity profile.




Figure 7. (Left) contours of normal vorticity and streamlines of the in-plane velocity field and (right) contours of normal component of the relative velocity $\mathbf{u}_{rel}$ in three orthogonal planes passing through the agglomerate, for Case A.4 at $t = 26$.




Figure 8. Iso-surface of the relative vorticity magnitude $\omega_{rel} = 0.46$ obtained from the velocity field for Case A.4 at $t = 26$, showing two tilted vortex rings generated by the particle-induced velocity field near the rotating agglomerate. The top two plots show iso-surfaces in the (a) x - y plane and (b) x - z plane. The bottom two plots, (c) and (d), show the same iso-surface views together with a slice showing ω_{rel} contours in the normal plane.




Figure 9. Time series showing breakup of single agglomerate in a shear flow, for Case A.1 at times (a) $t = 0$, (b) 5, (c) 10, (d) 15 and (e) 20.




Figure 10. Plots showing measures characterizing breakup of a single agglomerate in a shear flow. (a) Number of fragments that an agglomerate breaks up into versus adhesion parameter. When the agglomerate does not break up, $N_{frag} = 1$. (b) Number of particles N in agglomerates following breakup versus adhesion parameter. The dashed line is the experimental power-law fit $N \propto Ad^{0.879}$ from Sonntag and Russel (1986) for maximum number of particles, where the proportionality coefficient is fit to the data. The data is plotted for Cases A.1-A.4 (red deltas), A.5-A.8 (green circles), and A.9-A.12 (blue diamonds) from Table 1.




Figure 11. Scatter plots illustrating three types of agglomerate interactions: merger (Case B.15), bouncing (Case B.19) and fragmentation (Case B.13).




Figure 12. Summary of results for all agglomerate collision runs, showing the number of agglomerates (N_{agg}) remaining after collision as a function of adhesion parameter and the ratio D_a / R_{g0} of offset distance to initial radius of gyration. Colors indicate results from different agglomerates. Numbers indicate cases with agglomerate merger ($N_{agg} = 1$), bounce ($N_{agg} = 2$), and fragmentation ($N_{agg} > 2$).




Figure 13. Plot indicating the number of captured particles in bouncing collisions versus the total number of particles in an agglomerate. The number of red particles captured by blue agglomerates is plotted in red, and the number of blue particles captured by red agglomerates is plotted in blue. Different symbols are used to indicate different computations, with one red and one blue symbol for each computation.




Figure 14. Time variation of (a) number of touching particles originating in different agglomerates and (b) total dimensionless compressive force between the agglomerates for a typical bouncing case (Case B.19). Collision onset occurs at approximately $t = 5$ and the agglomerates detach at $t = 14$.




Figure 15. Scatter plots during a bouncing agglomerate collision (Case B.19) at $t = 7$, with colors indicating (a) agglomerate from which each particle originated, (b) total compressive force acting on each particle, and (c) same plot as in (b) with the low-compression particles (with compressive force < 1.5) blanked out. High compression force chains occur in a particle core region spreading outward from the collision point.




Figure 16. Iso-surface of relative vorticity magnitude $\omega_{rel} = 0.3$ (top) and particle scatter plot colored by initial agglomerate identity (bottom) for a case where the particle agglomerates merge (Case B.15), at times (a) $t = 6$, (b) 8, (c) 10, and (d) 12 during which collision and merger of the agglomerates occurs. The agglomerate centroids and initial radius of gyration are indicated in the upper plots by a black dot and a circle, respectively, for each agglomerate.




Figure 17. Iso-surface of relative vorticity magnitude $\omega_{rel} = 0.3$ (top) and particle scatter plot colored by initial agglomerate identity (bottom) for a case where the particle agglomerates bounce (Case B.19), at times (a) $t = 6$, (b) 8, (c) 10, and (d) 12 during which collision of the agglomerates occurs. The agglomerate centroids and initial radius of gyration are indicated in the upper plots by a black dot and a circle, respectively, for each agglomerate.