



Fourier Series of Circle Embeddings

Leonid V. Kovalev¹ · Xuerui Yang¹

Received: 11 August 2018 / Revised: 20 January 2019 / Accepted: 30 January 2019 /

Published online: 11 March 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

We study the Fourier series of circle homeomorphisms and circle embeddings, with an emphasis on the Blaschke product approximation and the vanishing of Fourier coefficients. The analytic properties of the Fourier series are related to the geometry of the circle embeddings, and have implications for the curvature of minimal surfaces.

Keywords Circle homeomorphisms · Fourier series · Circle embeddings · Harmonic maps · Blaschke products

Mathematics Subject Classification 31A05 · 30J10 · 42A16

1 Introduction

This paper concerns sense-preserving embeddings f of the unit circle $\mathbb{T} = \{z \in \mathbb{C}: |z| = 1\}$ into the complex plane \mathbb{C} , with the emphasis on the relation between the geometry of the image $f(\mathbb{T})$ and the behavior of the Fourier coefficients $\hat{f}: \mathbb{Z} \rightarrow \mathbb{C}$. The main questions addressed here are: how often does the Fourier series of a circle homeomorphism ($f(\mathbb{T}) = \mathbb{T}$) terminate in either positive or negative direction? And does a circle embedding have to have non-zero Fourier coefficients in some fixed finite subset of \mathbb{Z} ?

Communicated by Dmitry Khavinson.

L. V. Kovalev was supported by the National Science Foundation Grants DMS-1362453 and DMS-1764266. X. Yang was supported by an Undergraduate Summer Research Award from Syracuse University.

✉ Leonid V. Kovalev
lvkovale@syr.edu

Xuerui Yang
xyang20@syr.edu

¹ 215 Carnegie, Mathematics Department, Syracuse University, Syracuse, NY 13244, USA

Our main results are stated in Theorems 4.2, 5.1, and 5.2. Theorem 4.2 shows that functions of the form $\arg(B(\zeta)/\zeta^n)$ are dense in $C^1(\mathbb{T})$, improving a C^0 -approximation result due to Helson and Sarason [11]. As a corollary, circle diffeomorphisms of the form $\zeta \mapsto B(\zeta)/\zeta^n$ are uniformly dense among all sense-preserving circle homeomorphisms (Corollary 4.1). Theorem 5.1 shows that circle embeddings can have arbitrarily long gaps at the beginning of their Fourier series. In the opposite direction, Theorem 5.2 provides a Heinz-type estimate $|\hat{f}(-1)| + |\hat{f}(1)| \geq c > 0$ (cf. [10]) for circle embeddings with a horizontally convex image.

The properties of the Fourier coefficients $\hat{f}(n)$ of circle embeddings f are of interest for multiple reasons. First, $\hat{f}(n)$ are also the Taylor coefficients of the harmonic extension of f to the unit disk \mathbb{D} . This implies, for example, that the inequality $|\hat{f}(1)| \leq |\hat{f}(-1)|$ is an obstruction to the injectivity of this extension, by a theorem of Lewy [5, p. 20].

When $f(\mathbb{T}) = \mathbb{T}$, a lower bound on $|\hat{f}(1)|^2 + |\hat{f}(-1)|^2$ yields an upper bound on the Gaussian curvature of a minimal graph over the unit disk. This can be viewed as a quantitative form of the Bernstein theorem on minimal surfaces: every minimal graph over \mathbb{R}^2 is a plane. This relation motivated the Heinz inequality [10] with several subsequent improvements until the sharp form was achieved by Hall [7]. The optimal Gaussian curvature bound remains conjectural [5, Conj. 10.3.2].

Finally, the fact that $|\hat{f}(1)| > |\hat{f}(-1)|$ for every circle homeomorphism, in its quantitative form, is a key to the conformally natural extension of circle homeomorphisms devised by Douady and Earle [3].

This paper is organized as follows. Sections 3 and 4 concern the circle diffeomorphisms whose Fourier series terminates in one direction; they are closely related to Blaschke products. In Sect. 5, we consider the circle embeddings that lack low-frequency Fourier terms. Section 2 collects the necessary background results.

2 Preliminaries

An *embedding* is a map that is a homeomorphism onto its image. For maps $f: \mathbb{T} \rightarrow \mathbb{C}$, this property is equivalent to being continuous and injective. All circle embeddings considered in this paper are sense-preserving. In the special case $f(\mathbb{T}) = \mathbb{T}$, we have a circle homeomorphism.

The Fourier coefficients of an integrable function $f: \mathbb{T} \rightarrow \mathbb{C}$ are given by the following:

$$\hat{f}(n) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) e^{-in\theta} d\theta.$$

We write $\text{supp } \hat{f} = \{n \in \mathbb{Z}: \hat{f}(n) \neq 0\}$. The Lebesgue space $L^2(\mathbb{T})$ and the sequence space $\ell^2(\mathbb{Z})$ are equipped with inner products:

$$\langle f, g \rangle_{L^2(\mathbb{T})} = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) \overline{g(e^{i\theta})} d\theta$$

and

$$\langle a, b \rangle_{\ell^2(\mathbb{Z})} = \sum_{n \in \mathbb{Z}} a_n \overline{b_n}.$$

Parseval's theorem asserts that $f \mapsto \hat{f}$ is an isomorphism, with

$$\langle f, g \rangle_{L^2(\mathbb{T})} = \langle \hat{f}, \hat{g} \rangle_{\ell^2(\mathbb{Z})}. \quad (2.1)$$

Proposition 2.1 *For every measurable function $f: \mathbb{T} \rightarrow \mathbb{T}$, the set of shifted Fourier coefficients $\{\hat{f}(\cdot - k): k \in \mathbb{Z}\}$ is an orthonormal basis of $\ell^2(\mathbb{Z})$.*

Proof By virtue of (2.1), the statement is equivalent to $\{e^{ik\theta} f(e^{i\theta}): k \in \mathbb{Z}\}$ being an orthonormal basis of $L^2(\mathbb{T})$. The latter follows from the multiplication map $g \mapsto fg$ is a unitary operator on $L^2(\mathbb{T})$. \square

A complex-valued function is called harmonic if its real and imaginary parts are harmonic. For any continuous function on \mathbb{T} , the Poisson integral provides a continuous extension $F: \overline{\mathbb{D}} \rightarrow \mathbb{C}$, which is harmonic in \mathbb{D} [1, p. 169]. The continuity allow us to relate the Taylor coefficients of F to the Fourier coefficients of f , as follows:

$$\lim_{r \nearrow 1} \frac{1}{2\pi} \int_0^{2\pi} F(re^{i\theta}) e^{-in\theta} d\theta = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) e^{-in\theta} d\theta = \hat{f}(n).$$

Let us record this as a proposition.

Proposition 2.2 *If $f: \mathbb{T} \rightarrow \mathbb{C}$ is continuous, then the series*

$$F(z) = \sum_{n=0}^{\infty} \hat{f}(n) z^n + \sum_{n=1}^{\infty} \hat{f}(-n) \bar{z}^n$$

defines a harmonic function in \mathbb{D} , for which f provides a continuous boundary extension.

If $f: \mathbb{T} \rightarrow \mathbb{C} \setminus \{0\}$ is a continuous function, we denote by $\Delta_{\mathbb{T}} \arg f$ the change of the continuous argument of $f(z)$ as z travels around \mathbb{T} once in the positive direction. Note that $\Delta_{\mathbb{T}} \arg f$ is the winding number of f around 0, multiplied by 2π .

A (finite) *Blaschke product* is a function of the form:

$$B(z) = \sigma \prod_{k=1}^n \frac{z - z_k}{1 - \bar{z}_k z},$$

where $z_1, \dots, z_n \in \mathbb{D}$ and $\sigma \in \mathbb{T}$. Note that $\Delta_{\mathbb{T}} \arg B = 2\pi n$. We refer to n as the degree of B . A Blaschke product of degree 1 is a *Möbius transformation*. All Blaschke

products in this paper are finite. The recent book [2] is an excellent reference on such products. We will often use the identity [2, (3.4.7)], which states that

$$\zeta \frac{B'(\zeta)}{B(\zeta)} = \sum_{k=1}^n \frac{1 - |z_k|^2}{|\zeta - z_k|^2} = \sum_{k=1}^n P_{\mathbb{D}}(z_k, \zeta) \quad (\zeta \in \mathbb{T}), \quad (2.2)$$

where $P_{\mathbb{D}}$ is the Poisson kernel for \mathbb{D} [12, p. 8]:

$$P_{\mathbb{D}}(z, \zeta) = \operatorname{Re} \left(\frac{\zeta + z}{\zeta - z} \right) = \frac{1 - |z|^2}{|\zeta - z|^2}.$$

Note that, for $\zeta \in \mathbb{T}$, the quantity $\zeta B'(\zeta)/B(\zeta)$ is the derivative of $\arg B(\zeta)$ with respect to $\arg \zeta$. This derivative can be used for the following elementary characterization of circle homeomorphisms and diffeomorphisms.

Proposition 2.3 *Suppose that $f: \mathbb{T} \rightarrow \mathbb{T}$ is a C^1 -smooth map. Let $g(e^{i\theta}) = \frac{d \arg f(e^{i\theta})}{d\theta}$ be its derivative. Then*

- (a) *f is a homeomorphism if and only if $\Delta_{\mathbb{T}} \arg f = 2\pi$, $g \geq 0$ on \mathbb{T} and g is not identically zero on any non-trivial subarc of \mathbb{T} ;*
- (b) *f is a diffeomorphism if and only if $\Delta_{\mathbb{T}} \arg f = 2\pi$ and $g > 0$ on \mathbb{T} .*

The assumption of g not vanishing on any subarc holds if f is real-analytic.

3 Circle Homeomorphisms with a Terminating Fourier Series

In this section, $f: \mathbb{T} \rightarrow \mathbb{T}$ is a sense-preserving homeomorphism. Our goal is to identify all such homeomorphisms for which $\operatorname{supp} \hat{f}$ is bounded from above or from below. It is a simple observation that $\operatorname{supp} \hat{f}$ can be bounded from both sides only if it is exactly $\{1\}$.

Proposition 3.1 *If f is a circle homeomorphism with $\operatorname{supp} \hat{f}$ bounded, then f is a rotation.*

Proof Let $a = \min \operatorname{supp} \hat{f}$ and $b = \max \operatorname{supp} \hat{f}$. If $a < b$, then

$$\left\langle \hat{f}(a + \cdot), \hat{f}(b + \cdot) \right\rangle_{\ell^2(\mathbb{Z})} = \hat{f}(a) \overline{\hat{f}(b)} \neq 0,$$

contradicting Proposition 2.1. Hence, $a = b$, which means that $f(z) = cz^a$ for some constant c . This map is a homeomorphism only when $a = 1$, i.e., f is a rotation. \square

In contrast to the above, there are rich families of circle homeomorphisms whose Fourier series terminates in one direction only.

Proposition 3.2 *Suppose $f: \mathbb{T} \rightarrow \mathbb{T}$ is a circle homeomorphism. Then*

- (a) *$\operatorname{supp} f$ is bounded below if and only if $f(\zeta) = B(\zeta)/\zeta^n$ for some $n \in \mathbb{N} \cup \{0\}$, where B is a Blaschke product of degree $n + 1$;*

(b) *supp f is bounded above if and only if $f(\zeta) = \zeta^n/B(\zeta)$ for some $n \in \mathbb{N}$, where B is a Blaschke product of degree $n - 1$.*

Proof The “if” part is clear in both cases: the Fourier series of $B(\zeta)/\zeta^n$ is supported on $\mathbb{Z} \cap [-n, \infty)$, while the Fourier series of $\zeta^n/B(\zeta) = \zeta^n \overline{B(\zeta)}$ is supported on $\mathbb{Z} \cap (-\infty, n]$. We proceed to “only if”.

(a) Let $n = \max(0, -\min \text{supp } \hat{f})$ and $g(\zeta) = \zeta^n f(\zeta)$. By construction, $\hat{g}(k) = 0$ for all $k < 0$. By Proposition 2.2, the function g has a holomorphic extension to \mathbb{D} which is continuous on $\overline{\mathbb{D}}$. Since $|g| \equiv 1$ on \mathbb{T} , it follows that g is a Blaschke product [2, Thm. 3.5.2]. Let d be the degree of g . It follows that $\Delta_{\mathbb{T}} \arg g = 2\pi d$. On the other hand, $\Delta_{\mathbb{T}} \arg f = 2\pi$, because f is a homeomorphism. Hence, $\Delta_{\mathbb{T}} \arg(\zeta^n) = 2\pi(d - 1)$, which means $n = d - 1$. Part (a) is proved.

To prove (b), let $n = \max(0, \max \text{supp } \hat{f})$ and $g(\zeta) = \zeta^n \overline{f(\zeta)} = \zeta^n / f(\zeta)$. By construction $\hat{g}(k) = 0$ for all $k < 0$. As in part (a), we conclude that g is a Blaschke product of some degree d . This time, $\Delta_{\mathbb{T}} \arg g = 2\pi d$ together with the relation $g(\zeta) = \zeta^n / f(\zeta)$ yield $d = n - 1$, completing the proof. \square

Proposition 3.2 does not yet establish the above claim about having rich families of circle homeomorphisms, because not every quotient of the form stated in Proposition 3.2 is a homeomorphism of \mathbb{T} . As a warm-up, let us consider the special case $n = 2$ of Proposition 3.2 (b); that is

$$f(\zeta) = \sigma \zeta^2 \frac{1 - \overline{z_1} \zeta}{\zeta - z_1} \quad (z_1 \in \mathbb{D}, \sigma \in \mathbb{T}). \quad (3.1)$$

Letting $B(\zeta) = (\zeta - z_1)/(1 - \overline{z_1} \zeta)$, we obtain from (2.2) that

$$\frac{\zeta f'(\zeta)}{f(\zeta)} = 2 - \frac{\zeta B'(\zeta)}{B(\zeta)} = 2 - \frac{1 - |z_1|^2}{|\zeta - z_1|^2} \quad (\zeta \in \mathbb{T}). \quad (3.2)$$

The expression (3.2) is minimized when $|\zeta - z_1| = 1 - |z_1|$, and thus, its minimum value is as follows:

$$2 - \frac{1 + |z_1|}{1 - |z_1|} = \frac{1 - 3|z_1|}{1 - |z_1|}.$$

By Proposition 2.3, the function (3.1) is a circle homeomorphism if and only if $|z_1| \leq 1/3$, and is a diffeomorphism if and only if $|z_1| < 1/3$.

To treat both cases of Proposition 3.2 in a unified way, let us consider the quotients of two Blaschke products. These are precisely the rational functions that map \mathbb{T} to itself.

Lemma 3.1 *Suppose that B_1 and B_2 are finite Blaschke products; that is*

$$B_1(z) = \sigma_1 \prod_{k=1}^n \frac{z - z_k}{1 - \overline{z_k} z}, \quad B_2(z) = \sigma_2 \prod_{k=1}^m \frac{z - w_k}{1 - \overline{w_k} z}, \quad (3.3)$$

where $z_1, \dots, z_n, w_1, \dots, w_m \in \mathbb{D}$ and $\sigma_1, \sigma_2 \in \mathbb{T}$. The quotient $f = B_1/B_2$ is a circle homeomorphism if and only if $n - m = 1$ and

$$\sum_{k=1}^n P_{\mathbb{D}}(z_k, \zeta) \geq \sum_{k=1}^m P_{\mathbb{D}}(w_k, \zeta) \quad \text{for all } \zeta \in \mathbb{T}, \quad (3.4)$$

where $P_{\mathbb{D}}$ is the Poisson kernel. If, additionally, strict inequality holds in (3.4) for all $\zeta \in \mathbb{T}$, then f is a circle diffeomorphism.

Proof Using (2.2), we get

$$\frac{\zeta f'(\zeta)}{f(\zeta)} = \frac{\zeta B'_1(\zeta)}{B_1(\zeta)} - \frac{\zeta B'_2(\zeta)}{B_2(\zeta)} = \sum_{k=1}^n P_{\mathbb{D}}(z_k, \zeta) - \sum_{k=1}^m P_{\mathbb{D}}(w_k, \zeta),$$

and then, the conclusion follows from Proposition 2.3. \square

Lemma 3.1 raises the question of verifying the condition (3.4). This condition can be restated in two equivalent ways. First, it is equivalent to saying that the *balayage* of the signed measure:

$$\sum_{k=1}^n \delta_{z_k} - \sum_{k=1}^m \delta_{w_k}$$

onto the boundary of \mathbb{D} is a positive measure; see [13, p. 6]. Second, it is equivalent to the inequality

$$\sum_{k=1}^n h(z_k) \geq \sum_{k=1}^m h(w_k), \quad (3.5)$$

being true for every positive harmonic function h in \mathbb{D} ; this follows by expressing h as a Poisson integral. However, neither of these two interpretations is easier to verify in practice than the original condition (3.4). If the approach used for the special case (3.1) is applied to the analysis of (3.4), it leads to the sufficient condition:

$$\sum_{k=1}^n \frac{1 - |z_k|}{1 + |z_k|} \geq \sum_{k=1}^m \frac{1 + |w_k|}{1 - |w_k|}. \quad (3.6)$$

However, (3.6) is unsatisfactory as it lacks the Möbius invariance property that is inherent in condition (3.4) [this invariance is particularly clear from the form (3.5)].

Let $d(z, w)$ denote the pseudo-hyperbolic distance between z and w , namely

$$d(z, w) = \left| \frac{z - w}{1 - \bar{z}w} \right| \quad (z, w \in \mathbb{D}).$$

By construction, d is invariant under the Möbius transformations of \mathbb{D} .

Theorem 3.1 *If the points $z_0, \dots, z_n, w_1, \dots, w_n \in \mathbb{D}$ satisfy the condition:*

$$d(z_k, w_k) \leq \frac{(1 - d(z_k, z_0))(1 - d(w_k, z_0))}{4n}, \quad k = 1, \dots, n, \quad (3.7)$$

then the quotient

$$f(\zeta) = \prod_{k=0}^n \frac{\zeta - z_k}{1 - \bar{z}_k \zeta} \left(\prod_{k=1}^n \frac{\zeta - w_k}{1 - \bar{w}_k \zeta} \right)^{-1} \quad (3.8)$$

is a circle homeomorphism. Furthermore, if (3.7) is strict for some k , then f is a diffeomorphism of \mathbb{T} .

The proof requires the following “additive Harnack inequality” for positive harmonic functions in \mathbb{D} .

Lemma 3.2 *For every positive harmonic function h on \mathbb{D} and all $z, w \in \mathbb{D}$, we have the following:*

$$|h(z) - h(w)| \leq \frac{2|z - w|}{(1 - |z|)(1 - |w|)} h(0). \quad (3.9)$$

Proof Since h can be written as a Poisson integral:

$$h(z) = \int_{\mathbb{T}} P_{\mathbb{D}}(z, \zeta) d\mu(\zeta)$$

for some measure μ on \mathbb{T} [4, Thm. 1.1], it suffices to prove (3.9) for $h(z) = P(z, \zeta)$ with $\zeta \in \mathbb{T}$. For such h , we have the following:

$$\begin{aligned} |h(z) - h(w)| &= \left| \operatorname{Re} \left(\frac{\zeta + z}{\zeta - z} - \frac{\zeta + w}{\zeta - w} \right) \right| \\ &\leq \left| \frac{2(z - w)\zeta}{(\zeta - z)(\zeta - w)} \right| \leq \frac{2|z - w|}{(1 - |z|)(1 - |w|)}, \end{aligned}$$

which proves (3.9), as $h(0) = 1$. \square

Equality is sometimes attained in (3.9), for example, if $h(z) = P(z, 1)$, $w = 0$, and $0 < z < 1$. However, (3.9) is not sharp for general $z, w \in \mathbb{D}$. It may be of interest to obtain a sharp form of the additive Harnack inequality.

Proof of Theorem 3.1 By Lemma 3.1, we only need to estimate

$$\sum_{k=0}^n P_{\mathbb{D}}(z_k, \zeta) - \sum_{k=1}^n P_{\mathbb{D}}(w_k, \zeta) \quad (3.10)$$

from below. By composing f with a Möbius transformation, we can achieve $z_0 = 0$ without affecting the homeomorphism property of f . Then, (3.7) takes the form:

$$d(z_k, w_k) \leq \frac{(1 - |z_k|)(1 - |w_k|)}{4n}, \quad k = 1, \dots, n. \quad (3.11)$$

Using Lemma 3.2 and noting that $|z_k - w_k| \leq 2d(z_k, w_k)$, we arrive at

$$|P_{\mathbb{D}}(z_k, \zeta) - P_{\mathbb{D}}(w_k, \zeta)| \leq \frac{4d(z_k, w_k)}{(1 - |z_k|)(1 - |w_k|)} \leq \frac{1}{n}.$$

Since $P_{\mathbb{D}}(0, \zeta) = 1$ on \mathbb{T} , we conclude that (3.10) is bounded from below by $1 - n(1/n) = 0$. Furthermore, if 0 is attained, then equality must hold in (3.11) and, consequently, in (3.7) for all k . \square

We can finally show that the circle homeomorphisms of the two types identified by Proposition 3.2, indeed, exist for all values of n . The following corollary of Theorem 3.1 is obtained by specializing the theorem to the cases $w_k \equiv 0$ and $z_k \equiv 0$.

Corollary 3.1 *Suppose $z_1, \dots, z_n \in \mathbb{D}$.*

(a) *If, for $k = 1, \dots, n - 1$:*

$$|z_k| \leq \frac{1 - |z_n|}{4(n - 1)}(1 - d(z_k, z_n)),$$

then the function

$$f(\zeta) = \zeta^{1-n} \prod_{k=1}^n \frac{\zeta - z_k}{1 - \overline{z_k} \zeta} \quad (3.12)$$

is a circle homeomorphism.

(b) *If $|z_k| \leq 1/(4n + 1)$ for $k = 1, \dots, n$, then the function*

$$f(\zeta) = \zeta^{n+1} \prod_{k=1}^n \frac{1 - \overline{z_k} \zeta}{\zeta - z_k} \quad (3.13)$$

is a circle homeomorphism.

Proof (a) When $w_1 = \dots = w_n$ in Theorem 3.1, condition (3.7) becomes

$$|z_k| \leq \frac{(1 - d(z_k, z_0))(1 - |z_0|)}{4n}, \quad k = 1, \dots, n.$$

To align this with the condition in part (a), replace n by $n - 1$ and relabel z_0 as z_n .

The map (3.8) then becomes (3.12).

(b) When $z_0 = \dots = z_n$ in Theorem 3.1, condition (3.7) becomes

$$|w_k| \leq \frac{1 - |w_k|}{4n}, \quad k = 1, \dots, n,$$

which simplifies to $|w_k| \leq 1/(4n + 1)$. It remains to relabel w_1, \dots, w_n as z_1, \dots, z_n and observe that (3.8) reduces to (3.13). \square

4 Approximation by Rational Circle Homeomorphisms

The goal of this section is to show that the circle homeomorphisms with a terminating Fourier series (i.e., those identified by the Proposition 3.2) are dense in the set of all circle homeomorphisms, with respect to the uniform norm. A similar approximation result for continuous functions was obtained by Helson and Sarason [11], see also [2, Thm. 4.3.1].

Theorem 4.1 [11, p. 9] *Every continuous function $u: \mathbb{T} \rightarrow \mathbb{R}$ can be approximated uniformly by functions of the form $\arg(B(\zeta)/\zeta^n)$, where B is a Blaschke product and n is the degree of B .*

Given a circle homeomorphism $f: \mathbb{T} \rightarrow \mathbb{T}$, we can take a continuous branch of $u(\zeta) = \arg(f(\zeta)/\zeta)$ on \mathbb{T} and apply Lemma 4.1 to it. It follows that f can be uniformly approximated by functions of the form $B(\zeta)/\zeta^{n-1}$ with $n = \deg B$. The same argument, applied to $v(\zeta) = \arg(\zeta/f(\zeta))$, yields an approximation to f of the form $\zeta^{n+1}/B(\zeta)$. However, this does not achieve our goal stated above, since it is not guaranteed that the approximating function is a circle homeomorphism.

We need a stronger form of the Helson–Sarason theorem, with approximation in the C^1 norm instead of the uniform norm. A smooth function $u: \mathbb{T} \rightarrow \mathbb{R}$ can be interpreted as a smooth 2π -periodic function on \mathbb{R} , and we use this interpretation to define its derivative $u': \mathbb{T} \rightarrow \mathbb{R}$ and the C^1 norm $\|u\|_{C^1} = \sup_{\mathbb{T}} |u| + \sup_{\mathbb{T}} |u'|$.

Theorem 4.2 *Every C^1 -smooth function $u: \mathbb{T} \rightarrow \mathbb{R}$ can be approximated in the C^1 norm by functions of the form $\arg(B(\zeta)/\zeta^n)$, where B is a Blaschke product and n is the degree of B .*

Before proving Theorem 4.2, let us observe that it implies that circle diffeomorphisms with a terminating Fourier series are uniformly dense in the set of circle homeomorphisms.

Corollary 4.1 *Every circle homeomorphism $f: \mathbb{T} \rightarrow \mathbb{T}$ is a uniform limit of circle diffeomorphisms of the form $\zeta \mapsto B(\zeta)/\zeta^{n-1}$, where B is a Blaschke product of degree n . It is also a uniform limit of circle diffeomorphisms of the form $\zeta \mapsto \zeta^{n+1}/B(\zeta)$, where B is a Blaschke product of degree n .*

Proof It is straightforward to approximate f by a diffeomorphisms $g: \mathbb{T} \rightarrow \mathbb{T}$ in the uniform norm. Indeed, f lifts to a continuous strictly increasing function $F: \mathbb{R} \rightarrow \mathbb{R}$, such that $f(e^{i\theta}) = e^{iF(\theta)}$. It is easy to see that the convolution of F with a smooth bump function has strictly positive derivative, and, thus, descends to a circle diffeomorphism.

The function $u(\zeta) = \arg(g(\zeta)/\zeta)$ is C^1 -smooth, with $u' > -1$ on \mathbb{T} . Theorem 4.2 provides C^1 approximation to u of the form $v(\zeta) = \arg(B(\zeta)/\zeta^n)$. When $\|u - v\|_{C^1}$ is small enough, we have $v' > -1$, and therefore, the map $\zeta \mapsto B(\zeta)/\zeta^{n-1}$ is a circle diffeomorphism.

To prove the second statement, apply Theorem 4.2 to the function $u(\zeta) = \arg(\zeta/g(\zeta))$ for which $u' < 1$, and conclude as above. \square

Algebraic approximation of circle maps was also considered by Ebenfelt et al. [6]. Ebenfelt et al. [6, Thm. 2.3] assert that circle diffeomorphisms of the form $z \mapsto$

$B(z)^{1/n}$, where B is a Blaschke product of degree n , approximate all orientation-preserving circle diffeomorphisms in the C^1 norm. Our proof of Theorem 4.2 has some common elements with the proof of Ebenfelt et al. [6, Thm. 2.3]. There are also differences between these results, e.g., our approximation is by rational functions.

Our first step toward the proof of Theorem 4.2 is to approximate a continuous function with zero mean by a linear combination of Poisson kernels with integer coefficients.

Lemma 4.1 *Suppose that $h: \mathbb{T} \rightarrow \mathbb{R}$ is continuous and $\int_{\mathbb{T}} h = 0$. Then, for any $\epsilon > 0$, there exist $n \in \mathbb{N}$ and $z_k, w_k \in \mathbb{D}$ ($k = 1, \dots, n$), such that*

$$\left| h(\zeta) - \sum_{k=1}^n P_{\mathbb{D}}(z_k, \zeta) + \sum_{k=1}^n P_{\mathbb{D}}(w_k, \zeta) \right| < \epsilon \quad \text{for all } \zeta \in \mathbb{T}. \quad (4.1)$$

Proof Let H be the harmonic extension of h to the unit disk. We fix $r < 1$, such that

$$|H(r\zeta) - h(\zeta)| < \epsilon/3 \quad \text{for all } \zeta \in \mathbb{T}. \quad (4.2)$$

In what follows it will be convenient to consider h as a 2π -periodic function on \mathbb{R} , writing $h(\theta)$ instead of $h(e^{i\theta})$. Similarly, we write $P_r(\theta)$ for $P_{\mathbb{D}}(re^{i\theta}, 1)$.

Since h has zero mean, there exists a C^1 -smooth 2π -periodic function g , such that $g' = h$. The Poisson integral of h can be written in terms of g via integration by parts:

$$\begin{aligned} H(re^{i\theta}) &= \frac{1}{2\pi} \int_0^{2\pi} h(\phi) P_r(\phi - \theta) d\phi \\ &= -\frac{1}{2\pi} \int_0^{2\pi} g(\phi) \frac{\partial P_r(\phi - \theta)}{\partial \phi} d\phi. \end{aligned} \quad (4.3)$$

Denote

$$Q_r(\theta) = \frac{\partial P_r(\theta)}{\partial \theta} \quad \text{and} \quad R_r(\theta) = \frac{\partial^2 P_r(\theta)}{\partial \theta^2}.$$

For sufficiently large n , the last integral in (4.3) is well approximated by an n -point Riemann sum, meaning that

$$\left| H(re^{i\theta}) + \frac{1}{2\pi n} \sum_{k=1}^n g(\phi_k) Q_r(\phi_k - \theta) \right| < \frac{\epsilon}{3}, \quad (4.4)$$

where $\phi_k = \frac{2\pi k}{n}$.

For $1 \leq k \leq n$, let $a_k = -g(\phi_k)/n$. By Taylor's theorem,

$$P_r(\phi_k - \theta + a_k) - P_r(\phi_k - \theta) - a_k Q_r(\phi_k - \theta) = \frac{R_r(\psi_k)}{2} a_k^2 \quad (4.5)$$

for some ψ_k . Both g and R_r are bounded on \mathbb{R} , being continuous and periodic. Therefore, (4.5) implies that

$$|P_r(\phi_k - \theta + a_k) - P_r(\phi_k - \theta) - a_k Q_r(\phi_k - \theta)| \leq \frac{C}{n^2} \quad (4.6)$$

with a constant C independent of n . Summing over k and letting n be sufficiently large, we conclude that

$$\left| \sum_{k=1}^n \{P_r(\phi_k - \theta + a_k) - P_r(\phi_k - \theta) - a_k Q_r(\phi_k - \theta)\} \right| < \frac{\epsilon}{3}. \quad (4.7)$$

Finally, combine (4.2), (4.4), and (4.7) to obtain

$$\left| h(\theta) - \sum_{k=1}^n \{P_r(\phi_k - \theta + a_k) - P_r(\phi_k - \theta)\} \right| < \epsilon$$

from where the lemma follows by letting $z_k = r e^{i(\phi_k + a_k)}$ and $w_k = r e^{i\phi_k}$. \square

The approximation provided by Lemma 4.1 yields approximation in the C^1 norm by a quotient of Blaschke products, as shown below.

Lemma 4.2 *Suppose $u: \mathbb{T} \rightarrow \mathbb{R}$ is a C^1 smooth function, and the numbers $z_1, \dots, z_n, w_1, \dots, w_n \in \mathbb{D}$ are such that*

$$\left| u'(\zeta) - \sum_{k=1}^n P_{\mathbb{D}}(z_k, \zeta) + \sum_{k=1}^n P_{\mathbb{D}}(w_k, \zeta) \right| < \epsilon \quad (4.8)$$

for all $\zeta \in \mathbb{T}$. Then, there exists $\sigma \in \mathbb{T}$, such that the rational function

$$B(\zeta) = \sigma \prod_{k=1}^n \frac{\zeta - z_k}{1 - \overline{z_k} \zeta} \frac{1 - \overline{w_k} \zeta}{\zeta - w_k}$$

satisfies $\|u - \arg B\|_{C^1} < (\pi + 1)\epsilon$ for some continuous branch of $\arg B$.

Proof Recall from (2.2) that the derivative of $\arg B(e^{i\theta})$ with respect to θ is equal to

$$\sum_{k=1}^n P_{\mathbb{D}}(z_k, e^{i\theta}) - \sum_{k=1}^n P_{\mathbb{D}}(w_k, e^{i\theta}).$$

By (4.8), the derivative of the difference $u(e^{i\theta}) - \arg B(e^{i\theta})$ is less than ϵ in absolute value. By choosing σ , so that $\arg B(1) = u(1)$ and using the mean value theorem, we conclude that $|u(e^{i\theta}) - \arg B(e^{i\theta})| < \pi\epsilon$ for all $\theta \in [-\pi, \pi]$. The estimate for the C^1 norm of $u - \arg B$ follows. \square

The combination of Lemmas 4.1 and 4.2 yields an approximation result similar to Theorem 4.2 but with the quotient of two Blaschke products instead of a Blaschke product divided by a monomial. The following result will allow us to shift the poles of the quotient to 0.

Lemma 4.3 *For every $z_0 \in \mathbb{D}$ and every $\epsilon > 0$, there exist $n \in \mathbb{N}$ and z_1, \dots, z_{n-1} in \mathbb{D} , such that*

$$\left| \sum_{k=0}^{n-1} P(z_k, \zeta) - n \right| < \epsilon \quad \text{for all } \zeta \in \mathbb{T}. \quad (4.9)$$

Specifically, we can take $z_k = z_0 \exp(2\pi i k/n)$.

Proof Fix $\zeta \in \mathbb{T}$. Let $r = |z_0|$ and choose R , such that $r < R < 1$. The rational function $\psi(z) = (\zeta + z)/(\zeta - z)$ is holomorphic and bounded by $M = (1+R)/(1-R)$ on the disk $|z| \leq R$. Trefethen and Weideman [14, Thm. 2.1] asserts that uniform Riemann sums converge exponentially fast to the integral of ψ over $|z| = r$, namely

$$\left| \frac{2\pi}{n} \sum_{k=0}^{n-1} \psi(z_k) - \int_0^{2\pi} \psi(re^{i\theta}) d\theta \right| \leq \frac{2\pi M}{(R/r)^n - 1}, \quad (4.10)$$

where $z_k = z_0 \exp(2\pi i k/n)$. By the mean value property $\int_0^{2\pi} \psi(re^{i\theta}) d\theta = 2\pi\psi(0) = 2\pi$. Multiplying (4.10) by $n/(2\pi)$, we obtain the following:

$$\left| \sum_{k=0}^{n-1} \psi(z_k) - n \right| \leq \frac{2\pi M n}{(R/r)^n - 1}. \quad (4.11)$$

The right-hand side of (4.11) tends to 0 as $n \rightarrow \infty$. This fact and the identity $\operatorname{Re} \psi(z_k) = P_{\mathbb{D}}(z_k, \zeta)$ yield (4.9). \square

The following is a corollary of Lemmas 4.1 and 4.3.

Corollary 4.2 *Suppose that $h: \mathbb{T} \rightarrow \mathbb{R}$ is continuous and $\int_{\mathbb{T}} h = 0$. Then, for any $\epsilon > 0$, there exist $n \in \mathbb{N}$ and $z_1, \dots, z_n \in \mathbb{D}$, such that*

$$\left| h(\zeta) - \sum_{k=1}^n P_{\mathbb{D}}(z_k, \zeta) + n \right| < \epsilon \quad \text{for all } \zeta \in \mathbb{T}. \quad (4.12)$$

Proof Lemma 4.1 yields an approximation of the form:

$$\max_{\zeta \in \mathbb{T}} \left| h(\zeta) - \sum_{k=1}^n P_{\mathbb{D}}(z_k, \zeta) + \sum_{k=1}^n P_{\mathbb{D}}(w_k, \zeta) \right| < \frac{\epsilon}{2}. \quad (4.13)$$

Then, we use Lemma 4.3 to replace each term $P_{\mathbb{D}}(w_k, \zeta)$ in (4.13) by a sum of the form $n_k - \sum_{j=1}^{n_k-1} P(w_{kj}, \zeta)$, such that

$$\max_{\zeta \in \mathbb{T}} \left| P_{\mathbb{D}}(w_k, \zeta) - \left\{ n_k - \sum_{j=1}^{n_k-1} P(w_{kj}, \zeta) \right\} \right| < \frac{\epsilon}{2n},$$

thus, arriving at (4.12) with some larger value of n . \square

Proof of Theorem 4.2 First, apply Corollary 4.2 to the function $h = u'$. Then, use Lemma 4.2 with $w_1 = \dots = w_n = 0$. \square

5 Vanishing Fourier Coefficients

In the investigation of the Fourier coefficients of circle homeomorphisms, a special role is played by the first coefficient $\hat{f}(1)$. Indeed, the identity map $f(\zeta) = \zeta$ has all coefficients other than $\hat{f}(1)$ equal to zero. In contrast, Hall [7] proved that $\hat{f}(1)$ never vanishes for circle homeomorphisms. This result cannot be strengthened to a lower bound for $|\hat{f}(1)|$, as is shown by the Möbius transformation $f(\zeta) = (\zeta + a)/(1 + \bar{a}\zeta)$ which has $\hat{f}(0) = a$, and consequently, $|\hat{f}(1)|^2 \leq 1 - |a|^2$ by Parseval's theorem. However, having large $|\hat{f}(0)|$ is the only obstruction here: Hall [7, Thm. 2] gave a positive lower bound for $|\hat{f}(0)| + |\hat{f}(1)|$ among all circle homeomorphisms, which Weitsman [15] sharpened to $|\hat{f}(0)| + |\hat{f}(1)| > 2/\pi$, using [8].

The most notable estimate for the Fourier coefficients of circle homeomorphisms is $|\hat{f}(-1)|^2 + |\hat{f}(1)|^2 \geq 27/(4\pi^2)$, which is a sharp bound obtained by Hall [7] after 30 years of gradual improvements, starting with the paper [10] by Heinz. This line of investigation, motivated by curvature estimates for minimal surfaces, remains unfinished (see [5, §10.3] and [9]).

There are also non-vanishing results for more general circle embeddings. When $f(\mathbb{T})$ is convex, the Radó–Kneser–Choquet theorem [5, p. 29] states that the harmonic extension of f is a diffeomorphism, and therefore, $\hat{f}(1) \neq 0$; more precisely, $|\hat{f}(1)| > |\hat{f}(-1)|$. When $f(\mathbb{T})$ is star-shaped about 0, Hall [7, Thm. 2] proved that $|\hat{f}(0)| + |\hat{f}(1)| > 0$. The following proposition shows that the term $|\hat{f}(0)|$ is necessary here.

Proposition 5.1 *There exists an embedding $f : \mathbb{T} \rightarrow \mathbb{C}$, such that $f(\mathbb{T})$ is star-shaped about 0 and $\hat{f}(1) = 0$.*

Proof We look for f , such that $\overline{f(\zeta)} = f(\bar{\zeta})$ for all $\zeta \in \mathbb{T}$, which ensures that $f(\mathbb{T})$ is symmetric about the real axis, and that \hat{f} is real-valued. Let us write $f(e^{i\theta}) = F(\theta)$ for $\theta \in [0, \pi]$, and then:

$$\hat{f}(n) = \frac{1}{\pi} \operatorname{Re} \int_0^\pi F(\theta) e^{-in\theta} d\theta.$$

Our F will be piecewise linear, which justifies integration by parts:

$$\pi \hat{f}(1) = \operatorname{Re} i F(\theta) e^{-i\theta} \Big|_0^\pi - \operatorname{Re} \int_0^\pi i F'(\theta) e^{-i\theta} d\theta = \operatorname{Im} \int_0^\pi F'(\theta) e^{-i\theta} d\theta,$$

where the boundary term has zero contribution, because $F(0)$ and $F(\pi)$ are real.

We choose F piecewise linear with $F(0) = 1$, $F(\pi) = -1$, and $F(2\pi/3) = x + iy$, where $x, y > 0$ are to be chosen later. Thus:

$$F'(\theta) = \begin{cases} \frac{3}{2\pi}(x - 1 + iy), & 0 < \theta < 2\pi/3; \\ \frac{3}{\pi}(-1 - x - iy), & 2\pi/3 < \theta < \pi, \end{cases}$$

which yields

$$\begin{aligned} \operatorname{Im} \int_0^\pi F'(\theta) e^{-i\theta} d\theta &= -\frac{3(x-1)}{2\pi} \int_0^{2\pi/3} \sin \theta d\theta + \frac{3y}{2\pi} \int_0^{2\pi/3} \cos \theta d\theta \\ &\quad + \frac{3(x+1)}{\pi} \int_{2\pi/3}^\pi \sin \theta d\theta - \frac{3y}{\pi} \int_{2\pi/3}^\pi \cos \theta d\theta \\ &= -\frac{9(x-1)}{4\pi} + \frac{3\sqrt{3}y}{4\pi} + \frac{3(x+1)}{2\pi} + \frac{3\sqrt{3}y}{2\pi} \\ &= \frac{3}{4\pi} (-x + 3\sqrt{3}y + 5). \end{aligned}$$

For example, we achieve $\hat{f}(1) = 0$ with the choice $(x, y) = (8, 1/\sqrt{3})$. The curve $f(\mathbb{T})$ is a non-convex quadrilateral with vertices $1, x + iy, -1, x - iy$, which is obviously star-shaped about 0. \square

By subtracting a constant from f in Proposition 5.1, we can achieve $\hat{f}(0) = \hat{f}(1) = 0$; of course, $f(\mathbb{T})$ will no longer be star-shaped about 0 then. On the other hand, for every circle embedding $\hat{f}(n)$ must be non-zero for some positive n . Indeed, a computation with Green's formula shows that the area enclosed by $f(\mathbb{T})$ is $\pi \sum_{n \in \mathbb{Z}} n |\hat{f}(n)|^2$, which implies $\hat{f}(n) \neq 0$ for some $n > 0$. This raises the question: is there a fixed integer N , such that

$$\sum_{n=-N}^N |\hat{f}(n)| > 0, \tag{5.1}$$

for every circle embedding f ? By Hall's theorem, $N = 1$ suffices when $f(\mathbb{T})$ is star-shaped about 0. The main result of this section shows there is no universal N for general circle embeddings.

Theorem 5.1 *For every $N \in \mathbb{N}$, there exists an embedding $f: \mathbb{T} \rightarrow \mathbb{C}$, such that $\hat{f}(n) = 0$ whenever $|n| \leq N$.*

Proof Let $k = N+2$. We will consider embeddings $f: \mathbb{T} \rightarrow \mathbb{C}$ with k -fold symmetry; that is

$$f(e^{2\pi i/k} \zeta) = e^{2\pi i/k} f(\zeta), \quad \zeta \in \mathbb{T}. \tag{5.2}$$

As a consequence of (5.2), the Fourier coefficients of f satisfy that

$$e^{2\pi ni/k} \hat{f}(n) = e^{2\pi i/k} \hat{f}(n), \quad n \in \mathbb{Z},$$

which implies that $\hat{f}(n) = 0$ for all n , such that $n \not\equiv 1 \pmod{k}$. It remains to construct an embedding f , such that (5.2) holds and $\hat{f}(1) = 0$, which will ensure that $\hat{f}(n) = 0$ for $1 - k < n < 1 + k$.

For $\theta \in \mathbb{R}$, we define the following:

$$g(\theta) = \arccos(\cos \theta), \quad \rho(\theta) = 1 + \frac{2}{\pi} g(\theta), \quad h(\theta) = g(\theta) + \frac{1}{\pi} g(\theta)^2.$$

Since ρ and h are 2π -periodic and continuous, the function

$$f(e^{i\theta}) = \rho(k\theta) e^{i(\theta+h(k\theta))} \quad (5.3)$$

is well defined and continuous on \mathbb{T} . It has the k -fold symmetry (5.2) by construction.

Let us check that f is injective. Suppose that $f(e^{i\theta}) = f(e^{i\psi})$ for some $\theta, \psi \in \mathbb{R}$. Then, $\rho(k\theta) = \rho(k\psi)$, which, by the definition of ρ , implies $g(k\theta) = g(k\psi)$, and hence, $h(k\theta) = h(k\psi)$. Comparing the arguments of $f(e^{i\theta})$ and $f(e^{i\psi})$, we see that $\theta + h(k\theta) \equiv \psi + h(k\psi) \pmod{2\pi}$. Therefore, $\theta \equiv \psi \pmod{2\pi}$ as required.

Since g is 2π -periodic and even, it follows that:

$$\begin{aligned} \hat{f}(1) &= \frac{1}{2\pi} \int_0^{2\pi} \rho(k\theta) e^{ih(k\theta)} d\theta = \frac{k}{2\pi} \int_0^{2\pi/k} \rho(k\theta) e^{ih(k\theta)} d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \rho(t) e^{ih(t)} dt = \frac{1}{\pi} \int_0^\pi \rho(t) e^{ih(t)} dt. \end{aligned}$$

However, $g(t) = t$ for $t \in [0, \pi]$, which simplifies the above to

$$\begin{aligned} \pi \hat{f}(1) &= \int_0^\pi \left(1 + \frac{2}{\pi} t \right) e^{i(t+t^2/\pi)} dt \\ &= \int_0^{2\pi} e^{is} ds = 0 \end{aligned}$$

where we used $s = t + t^2/\pi$. The proof is complete. \square

The example constructed in Theorem 5.2 is highly non-convex and is not star-shaped with respect to any point (see Fig. 1 which illustrates the case $k = 3$).

We obtain a non-vanishing result under the assumption of the horizontal convexity of $f(\mathbb{T})$, which is substantially weaker than convexity. Horizontal convexity naturally appears in the studies of harmonic maps [5, §3.4].

Definition 5.1 A subset $A \subset \mathbb{R}^2$ is *horizontally convex* if its intersection with every horizontal line is connected (i.e., is an interval or empty set). Referring to a Jordan curve, we say that it is horizontally convex if its interior region is.




Fig. 1 Threefold symmetry with $\hat{f}(1) = 0$

Observe that $f(\mathbb{T})$ is horizontally convex if and only if \mathbb{T} is the union of two arcs on each of which $\text{Im } f$ is monotone. These arcs connect the global maximum of $\text{Im } f$ on \mathbb{T} to its global minimum; all the extrema of $\text{Im } f$ are global.

The map in Proposition 5.1 shows that the horizontal convexity of $f(\mathbb{T})$ allows for $\hat{f}(1) = 0$. We need to consider another Fourier coefficient to ensure at least one of them is non-zero.

Theorem 5.2 *If $f: \mathbb{T} \rightarrow \mathbb{C}$ is an embedding with horizontally convex image $f(\mathbb{T})$, then $|\hat{f}(-1)| + |\hat{f}(1)| > 0$. If, in addition, $\text{Im } f$ is Lipschitz continuous, then*

$$|\hat{f}(-1)| + |\hat{f}(1)| \geq \frac{\delta}{2\pi} \left(1 - \cos \frac{\delta}{4L}\right), \quad (5.4)$$

where $\delta = \max_{\mathbb{T}} \text{Im } f - \min_{\mathbb{T}} \text{Im } f$ and L is the Lipschitz constant of $\text{Im } f$.

Proof By the Borsuk–Ulam theorem (or just the intermediate value theorem), there exists $\zeta_0 \in \mathbb{T}$, such that $\text{Im } f(\zeta_0) = \text{Im } f(-\zeta_0)$. Replacing f with $f(\alpha\zeta) + \beta$ for suitable $\alpha \in \mathbb{T}$ and $\beta \in \mathbb{C}$, we can arrange that $\zeta_0 = 1$ and $\text{Im } f(\zeta_0) = 0$. Note that this replacement does not affect either side of (5.4).

By virtue of horizontal convexity, $\text{Im } f$ does not attain values of opposite sign on the upper half-circle; the same applies to the lower half-circle. Thus, $\text{Im } f(e^{i\theta}) \sin \theta$ does not attain values of opposite sign on \mathbb{T} . Since

$$\text{Re} \left(\hat{f}(1) - \hat{f}(-1) \right) = \frac{1}{\pi} \int_0^{2\pi} \text{Im } f(e^{i\theta}) \sin \theta \, d\theta, \quad (5.5)$$

we find that $\operatorname{Re} \left(\hat{f}(1) - \hat{f}(-1) \right) \neq 0$ unless the integrand in (5.5) is identically zero. However, the latter is impossible, because the Jordan curve $f(\mathbb{T})$ cannot be contained in a line.

It remains to prove (5.4). Pick $\zeta_1 \in \mathbb{T}$, such that $|\operatorname{Im} f| \geq \delta/2$. Let γ be the arc of length $\delta/(2L)$ centered at ζ_1 . On this arc, we have $|\operatorname{Im} f| \geq \delta/4$ by the Lipschitz condition. Therefore, the absolute value of the integral in (5.5) is at least

$$\frac{\delta}{4} \int_{\gamma} |\sin \theta| d\theta \geq \frac{\delta}{4} \int_{-\delta/(4L)}^{\delta/(4L)} |\sin \theta| d\theta = \frac{\delta}{2} \left(1 - \cos \frac{\delta}{4L} \right)$$

proving (5.4). \square

Let us record an application of Theorem 5.2 to minimal surfaces.

Corollary 5.1 *Let $F: \mathbb{D} \rightarrow \mathbb{R}^3$ be a conformally parameterized minimal surface with a continuous extension to \mathbb{T} . Let $f: \mathbb{T} \rightarrow \mathbb{R}^2$ be the composition of $F|_{\mathbb{T}}$ with an orthogonal projection $\mathbb{R}^3 \rightarrow \mathbb{R}^2$. If f satisfies the assumptions of Theorem 5.2, then the Gaussian curvature K of the minimal surface at $F(0)$ does not exceed*

$$\frac{32\pi^2}{\delta^2 \left(1 - \cos \frac{\delta}{4L} \right)^2},$$

where δ and L are as in Theorem 5.2.

Proof The computation in [5, p. 183] shows that

$$K \leq \frac{4}{|\hat{f}(-1)|^2 + |\hat{f}(1)|^2}.$$

On the other hand, Theorem 5.2 implies that

$$|\hat{f}(-1)|^2 + |\hat{f}(1)|^2 \geq \frac{1}{2} (|\hat{f}(-1)| + |\hat{f}(1)|)^2 \geq \frac{\delta^2}{8\pi^2} \left(1 - \cos \frac{\delta}{4L} \right)^2,$$

which proves the estimate, as claimed \square

Question 5.1 Is there a non-vanishing result of the form (5.1) for circle embeddings with a star-shaped image? On one hand, Hall's theorem [7, Thm. 2] gives $|\hat{f}(0)| + |\hat{f}(1)| > 0$ if $f(\mathbb{T})$ is star-shaped about 0; on the other, subtracting $\hat{f}(0)$ from the example in Proposition 5.1 shows that $|\hat{f}(0)| + |\hat{f}(1)|$ can vanish for general star-shaped embeddings.

Acknowledgements We thank the anonymous referee for suggesting several improvements and for bringing [6] to our attention.

References

1. Ahlfors, L.V.: Complex analysis, p. xi+331. McGraw-Hill Book Co., New York (1978)
2. Garcia, S.R., Mashreghi, J., Ross, W.T.: Finite Blaschke products and their connections. Springer, (2018)
3. Douady, A., Earle, C.J.: Conformally natural extension of homeomorphisms of the circle. *Acta Math.* **157**(1–2), 23–48 (1986)
4. Duren, P.L.: Theory of H^p spaces. Pure and Applied Mathematics, Vol. 38, Academic Press, New York-London, xii+258 (1970)
5. Duren, P.L.: Harmonic mappings in the plane. Cambridge Tracts in Mathematics, 156, Cambridge University Press, Cambridge, xii+212 (2004)
6. Ebenfält, P., Khavinson, D., Shapiro, H.S.: Two-dimensional shapes and lemniscates. In: Complex analysis and dynamical systems IV, Part 1, 45–59. Contemp. Math., 553, Israel Math. Conf. Proc., Amer. Math. Soc., Providence, RI (2011)
7. Hall, R.R.: On an inequality of E. Heinz. *J. Analyse Math.*, 42, 185–198 (1982/83)
8. Hall, R.R.: A class of isoperimetric inequalities. *J. Analyse Math.* **45**, 169–180 (1985)
9. Hall, R.R.: The Gaussian curvature of minimal surfaces and Heinz' constant. *J. Reine Angew. Math.* **502**, 19–28 (1998)
10. Heinz, E.: Über die Lösungen der Minimalflächengleichung. *Nachr. Akad. Wiss. Göttingen. Math.-Phys. Kl. Math.-Phys.-Chem. Abt.* **1952**, 51–56 (1952)
11. Helson, H., Sarason, D.: Past and future. *Math. Scand.* **21**, 5–16 (1967)
12. Ransford, Th.: Potential theory in the complex plane, London Mathematical Society Student Texts, 28, Cambridge University Press, Cambridge, x+232 (1995)
13. Totik, V.: Metric properties of harmonic measures. *Mem. Amer. Math. Soc.* **184**(867), vi+163 (2006)
14. Trefethen, L.N., Weideman, J.A.C.: The exponentially convergent trapezoidal rule. *SIAM Rev.* **56**(3), 385–458 (2014)
15. Weitsman, A.: On the Fourier coefficients of homeomorphisms of the circle. *Math. Res. Lett.* **5**(3), 383–390 (1998)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.