


Polariton Assisted Down-Conversion of Photons via Nonadiabatic Molecular Dynamics: A Molecular Dynamical Casimir Effect

Juan B. Pérez-Sánchez* and Joel Yuen-Zhou*

Department of Chemistry and Biochemistry, University of California San Diego, La Jolla, California 92093, United States

S Supporting Information

ABSTRACT: Quantum dynamics of the photoisomerization of a single 3,3'-diethyl-2,2'-thiacynine iodide molecule embedded in an optical microcavity was theoretically studied. The molecular model consisting of two electronic states and the reaction coordinate was coupled to a single cavity mode via the quantum Rabi Hamiltonian, and the corresponding time-dependent Schrödinger equation starting with a purely molecular excitation was solved using the Multiconfigurational Time-Dependent Hartree Method (MCTDH). We show that, for single-molecule strong coupling with the photon mode, nonadiabatic molecular dynamics produces mixing of polariton manifolds with differing number of excitations, without the need of counter-rotating light–matter coupling terms. Therefore, an electronic excitation of the molecule at the *cis* configuration is followed by the generation of two photons in the *trans* configuration upon isomerization. Conditions for this phenomenon to be operating in the collective strong light–matter coupling regime are discussed and found to be unfeasible for the present system, based on simulations of two molecules inside the microcavity. Yet, our finding suggests a new mechanism that, without ultrastrong coupling, achieves photon down-conversion by exploiting the emergent molecular dynamics arising in polaritonic architectures.



Most photochemical and spectroscopic processes studied at present correspond to the so-called weak light–matter coupling regime. In the latter, the effect of light is merely to produce transitions between molecular eigenstates. However, using confined electromagnetic media, such as those in optical microcavities¹ or nanostructures,² it has been possible to reach new regimes in which the light–matter interaction energy is comparable to electronic or vibrational energies, so that light has to be explicitly considered beyond a perturbative treatment.^{3–5} The new hybrid light–matter states in strong coupling (SC) and ultrastrong coupling (USC) regimes are called polaritons. These regimes do not necessarily require high intensity lasers, but occur despite the cavity modes being in their vacua or low-lying excitations.

In the case of a single molecule interacting with a single cavity mode, SC and USC regimes are commonly described by the Rabi model^{3,6,7} where the matter part is taken to be a two-level system with no internal structure, an insufficient description to study molecular processes where nuclear dynamics plays a major role.⁸ Theories that take into account the correlated nuclear–electronic–photonic dynamics have been recently developed to account for the rovibrational structure of molecules.^{9–11} In these new frameworks, molecular–photonic dynamics are described in dressed or polaritonic potential energy surfaces (PESs) and are governed by novel features such as light-induced avoided crossings (LIACs) and light-induced conical intersections (LICIs).^{12–16}

The idea of using light to drive chemical reactions with high selectivity has been one of the dreams of chemistry since the invention of the laser, and the ability to tune the characteristics

(polarization, mode volume, spatial profile) of the optical modes of a confined photonic media seems to offer new and versatile control knobs to manipulate molecular properties on demand, without invoking costly and time-consuming synthetic modifications.^{17,18} Indeed, most recent works focus on using strong light–matter coupling to change molecular processes such as photodissociation,^{19–21} photoisomerization,^{10,22–26} and charge and energy transfer.^{25,27} In this paper we focus on a less addressed complementary question: can the emergent molecular dynamics under SC be harnessed for photonic applications? Previous studies on atomic systems have shown that nonlinear optical effects can arise from the interplay of translational or vibrational motion and the coupling to a confined radiation field;^{28,29} however, as far as we are aware, no molecular analogues of these phenomena have been previously reported. By theoretically studying the photoisomerization of a single 3,3'-diethyl-2,2'-thiacynine iodide molecule that strongly interacts with a cavity, we find that, for specific cavity frequencies and sufficiently strong couplings, molecular photoexcitation into an electronic excited state can be followed by the spontaneous emission of two photons of a lower frequency via the cavity after isomerization, thus offering a new mechanism for photonic down-conversion using molecular polaritons. This phenomenon provides a molecular version of the dynamical Casimir (DC) effect, where

Received: September 28, 2019

Accepted: December 10, 2019

Published: December 10, 2019






Figure 1. Potential energy surfaces $V_a(\phi)$ and $V_b(\phi)$, diabatic coupling $V_{ab}(\phi)$, and transition dipole moment $\mu_{ab}(\phi)$ for 3,3'-diethyl-2,2'-thiacynine iodide; parameters from ref 32. *cis* and *trans* configurations are located around $\phi = \pm\pi$ and $\phi = 0$, respectively. Diabatic coupling $V_{ab}(\phi)$ generates molecular avoided crossings near $\phi = \pm 1.63$ rad.

photon pair creation arises from nonadiabatic modulation of the electromagnetic vacuum. As we shall show, molecular nonadiabatic effects mix states with different excitation numbers, without the need of the usually invoked counter-rotating light–matter coupling terms which are relevant in the standard realizations of the DC effect, which operate under USC conditions.^{3,30,31}

To begin with, the bare molecular Hamiltonian is given by

$$(\hbar = 1) \hat{H}_{\text{mol}} = \hat{T}_N + \hat{H}_{\text{el}}(\phi), \text{ where } \hat{T}_N = -\frac{1}{2m} \frac{\partial^2}{\partial \phi^2},$$

$$\hat{H}_{\text{el}}(\phi) = \begin{pmatrix} V_a(\phi) & V_{ab}(\phi) \\ V_{ab}(\phi) & V_b(\phi) \end{pmatrix} \quad (1)$$

ϕ is the torsional angle of the molecule (reaction coordinate), $V_a(\phi)$ and $V_b(\phi)$ are *diabatic* PESs, and $V_{ab}(\phi)$ is the diabatic coupling, responsible to produce transitions between the electronic states $|a\rangle$ and $|b\rangle$. Diagonalization of $\hat{H}_{\text{el}}$ as a function of ϕ produces adiabatic states of low energy $|g\rangle$ and high energy $|e\rangle$. These purely molecular quantities can be determined by quantum chemistry calculations and spectroscopic measurements. In this work, we take these properties from a previous model parametrized by Hoki and Brumer (see Figure 1).³²

On the other hand, the Hamiltonian of the cavity mode is given by

$$\hat{H}_{\text{cav}} = \omega_c(\hat{a}^\dagger \hat{a} + 1/2) \quad (2)$$

where ω_c is the cavity frequency and $\hat{a}$ is the photon annihilation operator. The Hamiltonian of the cavity–molecule system is the sum of those corresponding to the molecule, the cavity mode, and the interaction between them:

$$\hat{H} = \hat{H}_{\text{mol}} + \hat{H}_{\text{cav}} + \hat{H}_I \quad (3)$$

In this work, the cavity–molecule coupling is modeled as described in ref 13, where the photon is coupled to the electronic transition through the molecular transition dipole moment $\mu_{ab}(\phi)$ (see Figure 1). Equation 3 can then be re-expressed as $\hat{H} = \hat{T}_N + \hat{H}_{\text{e-p}}(\phi)$, with the adiabatic polaritonic BO Hamiltonian given by

$$\hat{H}_{\text{e-p}}(\phi) = \hat{H}_{\text{el}} + \omega_c(\hat{a}^\dagger \hat{a} + 1/2) + g(\phi)(\hat{a}^\dagger + \hat{a})\hat{\sigma}_x \quad (4)$$

Here, $g(\phi) = \epsilon \omega \mu_{ab}(\phi)$ and $\epsilon = 1/\sqrt{2V\omega_c\epsilon_0}$.

By conveniently using the Fock state basis for the cavity mode, $\hat{H}_{\text{e-p}}$ can be diagonalized to obtain adiabatic polaritonic PESs. These are shown in Figure 2 for specific values of cavity frequency and light–matter coupling. To establish a reference, the electronic energy gap at the *cis* configuration of the




Figure 2. Polaritonic PESs for $\omega_c = 0.5\omega_{ab}$. (a) Ignoring both light–matter and diabatic couplings. (b) Turning on diabatic coupling, generating adiabatic states $|g\rangle$ and $|e\rangle$. (c) Turning on both light–matter and diabatic couplings ($\epsilon = 0.04$ au $\rightarrow g \approx 0.21$ eV). (d) Mechanism of photon down-conversion is shown as a sequence of five steps. Light red: one photon $|g,1\rangle$. Dark red: two photons $|g,2\rangle$. Light blue: one exciton $|e,0\rangle$. Dark blue: one exciton with one photon $|e,1\rangle$. (d) Diagrammatic representation of mechanism in (c).

molecule is labeled as $\omega_{ab} \equiv V_a(\phi = -\pi) - V_b(\phi = -\pi) = 2.70$ eV. An experimental realization of single-molecule SC was recently carried out by Chikkaraddy and collaborators using a plasmonic nanocavity ($V < 40$ nm³) and an electronic transition of the methyl blue molecule.⁶ The coupling strength at the single molecule level was $g \approx 0.09$ eV with the transition dipole moment of the molecule being 3.8 D. For the present model, the transition dipole moment is 10 D, so we believe it is reasonable to consider $g = 0.21$ eV as the highest single-molecule coupling strength. Following that restriction, we varied ω_c from 25% to 100% of ω_{ab} , as well as the light–matter scaling parameter ϵ from 0.01 au to 0.04 au. This corresponds to a light–matter coupling ranging from 4% to 16% of ω_c . As an illustration, a coupling constant $g = 0.21$ eV

in our model corresponds to a single-photon electric field amplitude inside the optical microcavity of 1×10^9 V/m, corresponding to the free space electric field amplitude of a wave laser at the same frequency, but with an intensity $I = 1.4 \times 10^{11}$ W/cm².

Notice that polaritonic PESs (Figure 2c) at the *cis* and *trans* configurations resemble their counterparts in the absence of light–matter coupling (Figure 2b); this effect is a consequence of the cavity and the molecule being highly off-resonance at those configurations. However, as the torsional angle ϕ changes, so does the electronic energy gap, leading to resonances at $\phi = \pm 1.16, \pm 2.14$ rad. Light–matter coupling about these resonances generates LIACs. In this adiabatic polaritonic basis, the kinetic energy operator is not longer diagonal, and it generates nonadiabatic couplings among different polaritonic PES. By analogy with atomic systems or synthetic qubit systems, where different polariton manifolds can be couple through counter-rotating terms,^{3,30,31,33} coexistence of nonadiabatic and light–matter couplings is enough to mix polaritons with different number of excitations, with the difference that parity in excitation number need not be conserved (see the Supporting Information).

Figure 2c and d summarizes our proposal to achieve photon down-conversion using molecular polaritons:

(1) Resonant optical excitation of the molecule from state $|g,0\rangle$ to state $|e,0\rangle$. This transition occurs via direct interaction between the molecular dipole and a high frequency photon that is transparent (nonresonant) with respect to the cavity.

(2) Adiabatic dynamics across a LIAC converts electronic excitation into a cavity photon ($|e,0\rangle \rightarrow |g,1\rangle$).

(3) Nonadiabatic wavepacket dynamics across a molecular avoided crossing converts vibrational energy into electronic energy ($|g,1\rangle \rightarrow |e,1\rangle$).

(4) Adiabatic dynamics across a second LIAC converts electronic excitation into a second cavity photon ($|e,1\rangle \rightarrow |g,2\rangle$).

(5) Photons are spontaneously emitted into the electromagnetic bath through the cavity ($|g,2\rangle \rightarrow |g,0\rangle$).

We emphasize that steps (2) and (4) are possible only if light–matter coupling is strong enough to create a sizable LIAC that favors adiabatic nuclear dynamics. In other words, energy exchange between cavity photon and molecule must be fast compared to the instantaneous nuclear motion at the vicinity of the LIAC. Interestingly, steps 3 and 4 resemble a mechanism previously studied by Dobrovsky and Levine, who explored the possibility of light emission starting with a high-energy collision and subsequent nonadiabatic molecular dynamics.³⁴ Notice that our mechanism does not rely on counter-rotating light–matter coupling terms in any of its steps. To illustrate our proposal, we numerically solve the time-dependent Schrödinger equation using the Multi-Configurational Time-Dependent Hartree method (MCTDH).^{35,36} Even though we gained much conceptual insight by appealing to a Fock basis for the cavity mode, we will numerically deal with the latter in quadrature coordinates,³⁷

$$\hat{a} = \sqrt{\frac{\omega_c}{2}} \left(\hat{x} + \frac{i}{\omega} \hat{p} \right) \quad (5)$$

where $\hat{p} = -i\frac{\partial}{\partial x}$. With these identifications, eq 2 can be rewritten as

$$\hat{H} = \hat{T}_N + \frac{\hat{p}^2}{2} + \begin{pmatrix} V_a(\phi) + \frac{1}{2}\omega_c^2 x^2 & V_{ab}(\phi) + g(\phi)x \\ V_{ab}(\phi) + g(\phi)x & V_b(\phi) + \frac{1}{2}\omega_c^2 x^2 \end{pmatrix} \quad (6)$$

where the cavity mode appears as an additional “vibrational” coordinate, whose implementation in the MCTDH is straightforward.¹⁴ The wave function is expanded as a linear combination of diabatic electronic states $|k\rangle$:

$$\langle x, \phi | \Psi(t) \rangle = \sum_k \psi_k(x, \phi, t) |k\rangle, \quad \text{for } k = a, b \quad (7)$$

and the initial state is chosen to represent an impulsive Franck–Condon excitation of the molecule (directly via a high-energy photon that is transparent to the cavity), namely, a product state of the molecular ground state on top of the excited electronic state $|a\rangle$ at the *cis* configuration $\phi_b(\phi)$, accompanied by the vacuum state of the cavity mode $\chi(x)$:

$$\langle x, \phi | \Psi(0) \rangle = \phi_b(\phi) \chi(x) |a\rangle \quad (8)$$

To analyze the computational results, we calculate adiabatic populations of electronic and photonic states as a function of time:

$$P_{\kappa,n}(t) = \langle \Psi(t) | \kappa, n \rangle \langle \kappa, n | \Psi(t) \rangle, \quad \text{for } \kappa = g, e \text{ and } n = 0, 1, 2, \dots \quad (9)$$

In this expression, the torsional degree of freedom ϕ is traced out.

In Figure 3a, we present the adiabatic populations as a function of time. At short time, we can observe low amplitude and fast oscillations corresponding to off-resonance population transfer between $|e,0\rangle$ and $|g,1\rangle$ due to light–matter coupling. However, as the dynamics proceeds, those two states become




Figure 3. (a) Time-dependent adiabatic populations of photonic - electronic states for $\epsilon = 0.04$ au ($g = 0.21$ eV) and $\omega_c/\omega_{ab} = 0.5$. (b) Time average of two-photon population $\overline{P}_{g,2} = \frac{1}{T} \int_0^T P_{g,2}(t) dt$ ($T = 50$ fs) for different values of coupling strength ϵ and cavity frequency ω_c .

resonant, and there is a fast decay of the initial state $|e,0\rangle$ first into $|g,1\rangle$, then into $|e,1\rangle$, and finally into $|g,0\rangle$ and $|g,2\rangle$ by the end of the isomerization. The population of $|g,2\rangle$ at 30 fs evidences the photon pair generation. In Figure 3b, we see that if ω_c is too low or too large compared to ω_{ab} , the state with two photons is not significantly populated. In the first case, the LIAC lies near the region in which the transition dipole moment is drastically reduced, suppressing light–matter coupling. In addition, the polaritonic PES near the LIAC is too steep, implying nuclear dynamics that are too fast to be affected by the electronic–photonic coupling. In the second case, although light–matter coupling is not suppressed, the initial energy is not high enough to produce a nonadiabatic transition that would generate the second photon (see Supporting Information Figure S3). Those inconveniences are overcome if ω_c is near half of ω_{ab} and the light–matter coupling is sufficiently strong so that a large population of the state $|g,2\rangle$ is produced. It should also be noticed that the cavity frequency does not have to be exactly half of the exciton frequency, as vibrations can account for the remaining energy to form the two photons. For a better understanding of the mechanism, we calculate the time dependent probability density for each adiabatic state:

$$\rho_\kappa(x, \phi, t) = \langle \Psi(t) | x, \phi, \kappa \rangle \langle x, \phi, \kappa | \Psi(t) \rangle, \quad \kappa = g, e \quad (10)$$

Numerical simulations shown in Figure 4 support the mechanism proposed: at 0 fs the wavepacket corresponds to




Figure 4. Time dependent probability density along cavity (x) and nuclear (ϕ) degrees of freedom for adiabatic electronic states $|g\rangle$ (left) and $|e\rangle$ (right). The largest number of photons in the wave function is given by the number of nodes along the x coordinate.

the Franck–Condon excitation of the *cis* molecular configuration with the cavity in the vacuum state $|e,0\rangle$. At 10 fs there is population of the state $|g,1\rangle$ due to strong light–matter coupling. Subsequently, population at the state $|e,1\rangle$ is created at 20 fs, indicating that nonadiabatic molecular dynamics has occurred. Finally, at 30 fs, we clearly notice formation of a wavepacket in the lower adiabatic state with two photons, $|g,2\rangle$, in the *trans* configuration. Consistently, 30 fs is the time when the population of the $|g,2\rangle$ state reaches its maximum value (see Figure 3a). Other mechanisms that can be observed proceed as $|e,0\rangle \rightarrow |g,1\rangle \rightarrow |e,0\rangle$ (purely adiabatic dynamics and no photon down-conversion), $|e,0\rangle \rightarrow |g,1\rangle$ (adiabatic dynamics across one LIAC and no photon down-conversion), and $|e,0\rangle \rightarrow |g,0\rangle$ (purely nonadiabatic dynamics and no photon down-conversion). It is also worth noticing that states with more than two excitations (e.g., $|e,2\rangle$, $|g,3\rangle$) are not appreciably populated during the dynamics at any time, confirming that counter-rotating terms do not play a significant role in this phenomenon.

We shall briefly discuss the effects of various types of dissipation in our down-conversion mechanism. It is clear that any dissipative effect that operates within the time scale of the photon down-conversion mechanism above (~ 30 fs) would be detrimental. In particular, one may think that cavity leakage could play a significant constraint in the realization of our mechanism: single-molecule strong light–matter coupling utilizes plasmonic nanocavities with a very low quality factor ($Q \approx 12$),⁶ leading to photon leakage times less than 10 fs. However, we believe this time scale should not limit our mechanism, as it relies on the upper polariton being mostly molecular due to the large detuning between the cavity and the molecular transition at the Franck–Condon region. Hence, the dominant dissipative effects must be molecular, and they operate within a time scale of 40 fs, according to the study in ref 32. Importantly, this molecular dissipation time scale is chemical specific, and we suspect that some molecules will suffer important sources of down-conversion efficiency reduction due to nonadiabatic effects (e.g., nonradiative decay through conical intersections³⁸) from additional intramolecular vibrational modes that require explicit description. A more detailed account of such processes is beyond the scope of our work, but should be an important direction in future studies.

A complementary interpretation of the down-conversion mechanism can be provided from a time-independent perspective: for the molecule at the *cis* configuration, the upper polariton is mostly excitonic and accessible by means of a high-frequency photon. However, due to nonadiabatic couplings, the upper polariton is mixed with the lower polariton of the second excitation manifold at the *trans* configuration, which is accompanied by two photons. To further prove the two photon character of the eigenstates of $\hat{H}$ (eq 3) corresponding to the upper polariton PES, in Figure 5, we plot a variant of the absorption spectrum calculated as the spectral function of the initial wavepacket, and the spectral overlap of this wavepacket with the two-photon state $|2\rangle$. Spectral functions are computed as Fourier transforms of the following correlation functions (see the Supporting Information for further discussion of these calculations):^{39,40}

$$\sigma(\omega) = \int_0^T \int_0^T dt \, dt' \langle \Psi(t') | \Psi(t) \rangle e^{i(\omega + \omega_0)t} e^{-i(\omega + \omega_0)t'} \quad (11)$$




Figure 5. Transition probabilities into polaritonic eigenstates due to direct optical excitation of the molecule via the dipole (red) and their two-photon character (blue).

$$s(\omega) = \int_0^T \int_0^T dt' dt \langle \Psi(t') | 2 \rangle \langle 2 | \Psi(t) \rangle e^{i(\omega+\omega_0)t} e^{-i(\omega+\omega_0)t'} \quad (12)$$

where ω_0 is the energy of the ground state at the *cis* configuration, $T = 100$ fs, and we have employed the Condon approximation in light of the transition dipole moment being essentially constant at the Franck–Condon region (see Figure 1). The overlap of the two spectra evidences the two photon character of the polaritonic eigenstates in the absorption spectrum.

Preparation of a wavepacket in the polaritonic excited state can be done with a weak ultrafast laser pulse. The excited state populations can spontaneously decay via cavity leakage, part of the emission corresponding to the down-converted photons. Thus, our proposal is similar in spirit to the previously proposed spontaneous conversion Effect, in the context of conversion of virtual into real photons using counter-rotating light–matter coupling terms³⁰ (as explained, our proposal does not rely on such terms). A calculation that explicitly includes the laser pulse is shown in Figure 6, where we have used a




Figure 6. Time-dependent adiabatic populations of photonic–electronic states for $\epsilon = 0.04$ au and $\omega_c/\omega_{ab} = 0.5$ ($g = 0.21$ eV). Excitation was carried out by a Gaussian pulse (black) with $I_0 = 3.5 \times 10^{10}$ W/cm², $\omega = 2.7$ eV, and fwhm 25 fs.

Gaussian envelope with a maximum intensity $I_0 = 3.5 \times 10^{10}$ W/cm², $\omega = 2.7$ eV, and fwhm of 25 fs. Although the laser is strong enough to transfer about 15% of the population from the ground to the excited states during the first 50 fs, it is still weak enough that the population of the two-photon states is linear with respect to the pump intensity (see the Supporting Information), confirming that the mechanism relies on absorption of only one photon from the pump.

While the description of down-conversion above assumes it occurs spontaneously via photon-leakage, it should be enhanced by a parametric field passing through the cavity at

frequency $\omega \sim \omega_c$ which stimulates the emission of down-converted photons. Alternatively, the mechanism could be enhanced by using a chirped pulse that causes both the excitation at frequency ω_{ab} and the stimulated emission of the down-converted photons at ω_c .

The presented photon down-conversion phenomenon constitutes a molecular analogue of the dynamical Casimir effect (DCE),⁴¹ where a cavity mode can be excited out of the vacuum by means of nonadiabatic variations of the mirrors in an optical microcavity.^{42–45} Here, the kinetic energy is provided by the optical transition to the electronic excited state, forcing the nuclei to move out of the Franck–Condon region and undergo nonadiabatic molecular dynamics.

Experiments involving a single emitter strongly coupled to a confined electromagnetic field have been recently carried out by placing molecules on plasmonic nanocavities^{6,7,46–49} and Fabry–Pérot microcavities.⁵⁰ However, one of the most common setups to achieve SC involves the use of a macroscopic amount of molecules. The electromagnetic field interacts with an ensemble of N molecules to form 2 (upper and lower) polariton states, and $N - 1$ mostly molecular (dark) states that mix weakly with light either due to disorder or vibrational motion.^{4,10} Under these conditions, collective light–matter coupling scales as $\sim \sqrt{N}\epsilon$, where ϵ is the individual light–matter coupling and N is the number of particles in the microcavity.

To investigate the feasibility of achieving down-conversion using collective light–matter coupling, we performed calculations of two molecules embedded in an optical microcavity. Equation 6 can be generalized to N molecules,^{10,22} assuming molecules interact identically with the cavity mode and have no direct electrostatic interaction between them. The Hamiltonian is a generalization of the Dicke model⁵¹ that includes the nuclear degrees of freedom:

$$\hat{H} = \sum_{i=1}^N (\hat{T}_{N,i} + \hat{H}_{el,i}) + \omega_c(\hat{a}^\dagger \hat{a} + 1/2) + \sum_{i=1}^N g(\phi_i)(\hat{a}^\dagger + \hat{a})\hat{\sigma}_{x,i} \quad (13)$$

We study the two-photon generation at constant collective light–matter coupling for $N = 1, 2$ molecules (thus setting the individual light–matter coupling in each case at $\epsilon = 0.04/\sqrt{N}$ au). We first assume that only molecule 1 is initially excited, and calculate electronic state populations of each molecule at the *cis* ($|g_C\rangle, |e_C\rangle$) and *trans* ($|g_T\rangle, |e_T\rangle$) configurations using the projection operators

$$\hat{P}_{\kappa,C}^i = \int d\phi_i [\Theta(\phi_i - 1.63) + \Theta(-\phi_i - 1.63)] |\kappa_i, \phi_i\rangle \langle \kappa_i, \phi_i| \quad (14)$$

and

$$\hat{P}_{\kappa,T}^i = 1 - \hat{P}_{\kappa,C}^i \quad (15)$$

where $\hat{P}_{\kappa,C}^i$ is the projector of the i th molecule over the *cis* configuration and electronic state κ_i and $\Theta(\phi)$ is the Heaviside step function. For instance, the probability of both molecules to be at the *cis* configuration in the ground state, i.e., state $|g_C g_C\rangle$, is given by $\langle \hat{P}_{g_C}^1 \hat{P}_{g_C}^2 \rangle$. Furthermore, we calculate the population of two photon states as the expectation value of the

projection operator $|2\rangle\langle 2|$. Finally, we repeat the same analysis for an initial electronic state given by $\frac{1}{\sqrt{2}}(|ge\rangle + |eg\rangle)$.

As can be observed in Figure 7a, most dynamics during the first 100 fs involves the isomerization of the initially excited




Figure 7. Population dynamics for $\omega_c/\omega_{ab} = 0.5$ and $\epsilon = 0.03$ au, if only molecule 1 is initially excited. (a) States with significant population during the first 100 fs. (b) Population of states in which molecule 2 undergoes isomerization reaction. Notice the difference in vertical scale. (c) Comparison between populations of two-photon states for one and two molecules with the same *collective* light–matter coupling (blue vs red and green), and for two and one molecules with the same *individual* light–matter coupling (red and green vs dashed black).

molecule, while the second molecule remains at the *cis* configuration in the ground state. This path resembles that of the single-molecule scenario, in which the excited molecule isomerizes in around 30 fs, ending at the *trans* configuration in both the molecular ground and the excited state (see Figure 4). After 40 fs, the second molecule gets excited and also undergoes isomerization. However, this process seems to occur only if the first molecule is in its ground state at the *cis* configuration (Figure 7b). This can be understood by noticing that the molecules at the *cis* configuration are closer to resonance with the cavity mode. The individual coupling of each molecule with the cavity mode produces an effective coupling between them, causing the excitation of one molecule to be transferred to the other one (i.e., $|e_C g_C\rangle \rightarrow |g_C e_C\rangle$), as has been reported in previous works.^{22,52} Evidently, this process is not very likely in our setup and should not be observed in the limit where the single-molecule coupling is weak.

The mechanism involving two molecules can thus be summarized as follows:

- (1) Optical excitation of the first molecule ($|g_C g_C\rangle \rightarrow |e_C g_C\rangle$).
- (2) Isomerization of the first molecule ($|e_C g_C\rangle \rightarrow |e_T g_C\rangle$ and $|e_C g_C\rangle \rightarrow |g_T g_C\rangle$). As in the single molecule scenario, this can produce zero, one or two photons.
- (3) Cavity-mediated excitation-energy transfer from the first to the second molecule at the *cis* configuration ($|e_C g_C\rangle \rightarrow |g_C e_C\rangle$).
- (4) Isomerization of the second molecule ($|g_C e_C\rangle \rightarrow |g_C e_T\rangle$ and $|g_C e_C\rangle \rightarrow |g_C g_T\rangle$), producing zero, one, or two photons.

As one would expect based on the mechanism above, Figure 7c shows that having two molecules instead of one does not increase the likelihood of generation of two photons (see blue and red curves), so long as the *collective* light–matter coupling $\sqrt{N}\epsilon = 0.04$ au remains the same. To reinforce the observation that the observed effect is essentially a single-molecule one, we notice that, for fixed *individual* light–matter coupling $\epsilon = 0.028$ au, the two-photon state populations for two and one molecules (see red and black dashed curves) is very similar at short times. The reason we do not observe significant collective effects is that the molecules at the initial configuration are not in resonance with the cavity but become resonant as the isomerization proceeds. As a consequence, initial excitation of one molecule can lead to efficient energy exchange with the cavity only after sufficient nuclear dynamics ensues, while the other molecule remains off resonant. This was observed to be true even if the initial excitation is shared by all molecules in superposition. In other words, the Rabi splittings for the relevant LIACs relevant for this process mainly depend on the single-molecule coupling even if many molecules are present.

We believe that the fact that the cavity and the molecule are off-resonance at the initial configuration is not a vital characteristic of the down-conversion mechanism proposed here. If the molecules are resonant with the cavity at the Franck–Condon region, an excitation of the upper-polariton can still produce a molecular nonadiabatic transition that would generate a second excitation, which can afterward become a second photon. The Rabi splitting in that case would be of a collective nature, and the two-photon generation would be possible if the isomerization is faster than the decay from the upper-polariton to the dark states.³⁸ However, deleterious effects arising solely from collective coupling²⁰ may also complicate the down-conversion mechanism. The nonadiabatic molecular dynamics associated with molecules which are resonant at the Franck–Condon configuration could not be observed for the molecule studied here. Hence, the effectiveness of our down-conversion scheme in the collective regime is molecule-specific and will require additional investigations.

In summary, we have shown that strong light–matter coupling in conjunction with nonadiabatic molecular dynamics can lead to emerging nonlinear optical phenomena such as photon down-conversion. While much attention has been recently placed into the study novel chemical dynamics afforded by molecular polaritons, we wish to emphasize a complementary aspect of the problem that is equally rich and relevant: the use of molecular dynamics to generate new photonic phenomena. The elucidated effect operates at the single molecule SC regime, but we have provided plausible arguments that would allow to extend it to the collective regime, where many experiments are being currently performed.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jpclett.9b02870>.

Computational details; role of dipole self-energy term; polaritonic energy surfaces for cases where $\omega_c > \omega_{ab}/2$ and $\omega_c < \omega_{ab}/2$; details on the calculation of spectral functions; a more detailed explanation about the breaking of parity symmetry due to nonadiabatic coupling; two-photon population vs pump intensity (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: jperezsa@ucsd.edu.

*E-mail: joelyuen@ucsd.edu.

ORCID

Joel Yuen-Zhou: 0000-0002-8701-8793

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

J.B.P.S. and J.Y.Z. were funded with the NSF EAGER Award CHE 1836599.

■ REFERENCES

- (1) Vahala, K. J. Optical Microcavities. *Nature* **2003**, *424*, 839–846.
- (2) Vasa, P.; Lienau, C. Strong Light–Matter Interaction in Quantum Emitter/Metal Hybrid Nanostructures. *ACS Photonics* **2018**, *5*, 2–23.
- (3) Frisk Kockum, A.; Miranowicz, A.; De Liberato, S.; Savasta, S.; Nori, F. Ultrastrong Coupling Between Light and Matter. *Nat. Rev. Phys.* **2019**, *1*, 19–40.
- (4) Ribeiro, R. F.; Martínez-Martínez, L. A.; Du, M.; Campos-Gonzalez-Angulo, J.; Yuen-Zhou, J. Polariton Chemistry: Controlling Molecular Dynamics with Optical Cavities. *Chem. Sci.* **2018**, *9*, 6325–6339.
- (5) Chen, H.-T.; Li, T. E.; Nitzan, A.; Subotnik, J. E. Predictive Semiclassical Model for Coherent and Incoherent Emission in the Strong Field Regime: The Mollow Triplet Revisited. *J. Phys. Chem. Lett.* **2019**, *10*, 1331–1336.
- (6) Chikkaraddy, R.; de Nijs, B.; Benz, F.; Barrow, S. J.; Scherman, O. A.; Rosta, E.; Demetriadou, A.; Fox, P.; Hess, O.; Baumberg, J. J. Single-Molecule Strong Coupling at Room Temperature in Plasmonic Nanocavities. *Nature* **2016**, *535*, 127.
- (7) Santhosh, K.; Bitton, O.; Chuntanov, L.; Haran, G. Vacuum Rabi Splitting in a Plasmonic Cavity at the Single Quantum Emitter Limit. *Nat. Commun.* **2016**, *7*, 11823.
- (8) Flick, J.; Narang, P. Cavity-Correlated Electron-Nuclear Dynamics from First Principles. *Phys. Rev. Lett.* **2018**, *121*, 113002.
- (9) Schäfer, C.; Ruggenthaler, M.; Rubio, A. Ab Initio Non-relativistic Quantum Electrodynamics: Bridging Quantum Chemistry and Quantum Optics from Weak to Strong Coupling. *Phys. Rev. A: At, Mol., Opt. Phys.* **2018**, *98*, 043801.
- (10) Galego, J.; Garcia-Vidal, F. J.; Feist, J. Cavity-Induced Modifications of Molecular Structure in the Strong-Coupling Regime. *Phys. Rev. X* **2015**, *5*, 041022.
- (11) Lacombe, L.; Hoffmann, N. M.; Maitra, N. T. Exact Potential Energy Surface for Molecules in Cavities. *Phys. Rev. Lett.* **2019**, *123*, 083201.
- (12) Moiseyev, N.; Šindelka, M.; Cederbaum, L. S. Laser-Induced Conical Intersections in Molecular Optical Lattices. *J. Phys. B: At, Mol. Opt. Phys.* **2008**, *41*, 221001.
- (13) Kowalewski, M.; Bennett, K.; Mukamel, S. Cavity Femtochemistry: Manipulating Nonadiabatic Dynamics at Avoided Crossings. *J. Phys. Chem. Lett.* **2016**, *7*, 2050–2054.
- (14) Kowalewski, M.; Bennett, K.; Mukamel, S. Non-Adiabatic Dynamics of Molecules in Optical Cavities. *J. Chem. Phys.* **2016**, *144*, 054309.
- (15) Csehi, A.; Halász, G. J.; Vibók, Á. Collective Effect of Light-Induced and Natural Nonadiabatic Phenomena in the Dissociation Dynamics of the NaI Molecule. *Chem. Phys.* **2018**, *509*, 91–97.
- (16) Csehi, A.; Halász, G. J.; Cederbaum, L. S.; Vibók, g. Intrinsic and Light-Induced Nonadiabatic Phenomena in the NaI Molecule. *Phys. Chem. Chem. Phys.* **2017**, *19*, 19656–19664.
- (17) Thomas, A.; Lethuillier-Karl, L.; Nagarajan, K.; Vergauwe, R. M. A.; George, J.; Chervy, T.; Shalabney, A.; Devaux, E.; Genet, C.; Moran, J.; et al. Tilting a Ground-State Reactivity Landscape by Vibrational Strong Coupling. *Science* **2019**, *363*, 615–619.
- (18) Kéna-Cohen, S.; Yuen-Zhou, J. Polariton Chemistry: Action in the Dark. *ACS Cent. Sci.* **2019**, *5*, 386–388.
- (19) Triana, J. F.; Peláez, D.; Sanz-Vicario, J. L. Entangled Photonic-Nuclear Molecular Dynamics of LiF in Quantum Optical Cavities. *J. Phys. Chem. A* **2018**, *122*, 2266–2278.
- (20) Vendrell, O. Collective Jahn-Teller Interactions through Light-Matter Coupling in a Cavity. *Phys. Rev. Lett.* **2018**, *121*, 253001.
- (21) Kowalewski, M.; Mukamel, S. Manipulating Molecules with Quantum Light. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114*, 3278–3280.
- (22) Mandal, A.; Huo, P. Investigating New Reactivities Enabled by Polariton Photochemistry. *J. Phys. Chem. Lett.* **2019**, *10*, 5519–5529.
- (23) Feist, J.; Galego, J.; Garcia-Vidal, F. J. Polaritonic Chemistry with Organic Molecules. *ACS Photonics* **2018**, *5*, 205–216.
- (24) Galego, J.; Garcia-Vidal, F. J.; Feist, J. Suppressing Photochemical Reactions with Quantized Light Fields. *Nat. Commun.* **2016**, *7*, 13841.
- (25) Groenhof, G.; Toppari, J. J. Coherent Light Harvesting through Strong Coupling to Confined Light. *J. Phys. Chem. Lett.* **2018**, *9*, 4848–4851.
- (26) Fregoni, J.; Granucci, G.; Coccia, E.; Persico, M.; Corni, S. Manipulating Azobenzene Photoisomerization through Strong Light-Molecule Coupling. *Nat. Commun.* **2018**, *9*, 4688.
- (27) Du, M.; Martínez-Martínez, L. A.; Ribeiro, R. F.; Hu, Z.; Menon, V. M.; Yuen-Zhou, J. Theory for Polariton-Assisted Remote Energy Transfer. *Chem. Sci.* **2018**, *9*, 6659–6669.
- (28) Ritsch, H.; Domokos, P.; Brennecke, F.; Esslinger, T. Cold Atoms in Cavity-Generated Dynamical Optical Potentials. *Rev. Mod. Phys.* **2013**, *85*, 553–601.
- (29) Fernández-Vidal, S.; Zippilli, S.; Morigi, G. Nonlinear Optics with Two Trapped Atoms. *Phys. Rev. A: At, Mol., Opt. Phys.* **2007**, *76*, 053829.
- (30) Stassi, R.; Ridolfo, A.; Di Stefano, O.; Hartmann, M. J.; Savasta, S. Spontaneous Conversion from Virtual to Real Photons in the Ultrastrong-Coupling Regime. *Phys. Rev. Lett.* **2013**, *110*, 243601.
- (31) Stassi, R.; Savasta, S.; Garziano, L.; Spagnolo, B.; Nori, F. Output Field-Quadrature Measurements and Squeezing in Ultrastrong Cavity-QED. *New J. Phys.* **2016**, *18*, 123005.
- (32) Hoki, K.; Brumer, P. Mechanisms in Adaptive Feedback Control: Photoisomerization in a Liquid. *Phys. Rev. Lett.* **2005**, *95*, 168305.
- (33) Ashhab, S.; Nori, F. Qubit-Oscillator Systems in the Ultrastrong-Coupling Regime and their Potential for Preparing Nonclassical States. *Phys. Rev. A: At, Mol., Opt. Phys.* **2010**, *81*, 042311.
- (34) Dobrovsky, I.; Levine, R. Electronically Non-Adiabatic Transitions in the Evans-Polanyi Valence Bond Model. *Chem. Phys. Lett.* **1998**, *286*, 155–162.
- (35) Vendrell, O. Coherent Dynamics in Cavity Femtochemistry: Application of the Multi-Configuration Time-Dependent Hartree Method. *Chem. Phys.* **2018**, *509*, 55–65.
- (36) Worth, G. A.; Beck, M. H.; Jäckle, A.; Meyer, H.-D. *MCTDH Package*, version 8.2.; Heidelberg University, 2000.

- (37) Flick, J.; Appel, H.; Ruggenthaler, M.; Rubio, A. Cavity Born-Oppenheimer Approximation for Correlated Electron-Nuclear-Photon Systems. *J. Chem. Theory Comput.* **2017**, *13*, 1616–1625.
- (38) Ulusoy, I. S.; Gomez, J. A.; Vendrell, O. Modifying the Nonradiative Decay Dynamics through Conical Intersections via Collective Coupling to a Cavity Mode. *J. Phys. Chem. A* **2019**, *123*, 8832–8844.
- (39) Heller, E. J. *The Semiclassical Way to Dynamics and Spectroscopy*; Princeton University Press: Princeton, NJ, 2018.
- (40) Tannor, D. *Introduction to Quantum Mechanics: A Time-Dependent Perspective*; University Science Books: Sausalito, CA, 2007.
- (41) Triana, J. F.; Sanz-Vicario, J. L. Revealing the Presence of Potential Crossings in Diatomics Induced by Quantum Cavity Radiation. *Phys. Rev. Lett.* **2019**, *122*, 063603.
- (42) Wilson, C. M.; Johansson, G.; Pourkabirian, A.; Simoen, M.; Johansson, J. R.; Duty, T.; Nori, F.; Delsing, P. Observation of the Dynamical Casimir Effect in a Superconducting Circuit. *Nature* **2011**, *479*, 376–379.
- (43) de Melo e Souza, R.; Impens, F.; Maia Neto, P. A. Microscopic Dynamical Casimir Effect. *Phys. Rev. A: At., Mol., Opt. Phys.* **2018**, *97*, 032514.
- (44) Kim, W.-J.; Brownell, J. H.; Onofrio, R. Detectability of Dissipative Motion in Quantum Vacuum via Superradiance. *Phys. Rev. Lett.* **2006**, *96*, 200402.
- (45) Sánchez-Burillo, E.; Martín-Moreno, L.; García-Ripoll, J. J.; Zueco, D. Single Photons by Quenching the Vacuum. *Phys. Rev. Lett.* **2019**, *123*, 013601.
- (46) Reithmaier, J. P.; Søk, G.; Löffler, A.; Hofmann, C.; Kuhn, S.; Reitzenstein, S.; Keldysh, L.; Kulakovskii, V.; Reinecke, T.; Forchel, A. Strong Coupling in a Single Quantum Dot-Semiconductor Microcavity System. *Nature* **2004**, *432*, 197.
- (47) Ojambati, O. S.; Chikkaraddy, R.; Deacon, W. D.; Horton, M.; Kos, D.; Turek, V. A.; Keyser, U. F.; Baumberg, J. J. Quantum Electrodynamics at Room Temperature Coupling a Single Vibrating Molecule with a Plasmonic Nanocavity. *Nat. Commun.* **2019**, *10*, 1049.
- (48) Park, K.-D.; May, M. A.; Leng, H.; Wang, J.; Kropp, J. A.; Gougousi, T.; Pelton, M.; Raschke, M. B. Tip-Enhanced Strong Coupling Spectroscopy, Imaging, and Control of a Single Quantum Emitter. *Sci. Adv.* **2019**, *5*, eaav5931.
- (49) Chikkaraddy, R.; Turek, V.; Kongsuwan, N.; Benz, F.; Carnegie, C.; van de Goor, T.; de Nijs, B.; Demetriadou, A.; Hess, O.; Keyser, U. F.; et al. Mapping Nanoscale Hotspots with Single-Molecule Emitters Assembled into Plasmonic Nanocavities Using DNA Origami. *Nano Lett.* **2018**, *18*, 405–411.
- (50) Wang, D.; Kelkar, H.; Martin-Cano, D.; Rattenbacher, D.; Shkarin, A.; Utikal, T.; Göttinger, S.; Sandoghdar, V. Turning a Molecule into a Coherent Two-Level Quantum System. *Nat. Phys.* **2019**, *15*, 483.
- (51) Dicke, R. H. Coherence in Spontaneous Radiation Processes. *Phys. Rev.* **1954**, *93*, 99.
- (52) Galego, J.; Garcia-Vidal, F. J.; Feist, J. Many-Molecule Reaction Triggered by a Single Photon in Polaritonic Chemistry. *Phys. Rev. Lett.* **2017**, *119*, 136001.

Polariton Assisted Down-Conversion of Photons via Nonadiabatic Molecular Dynamics: A Molecular Dynamical Casimir Effect

Juan B. Pérez-Sánchez* and Joel Yuen-Zhou*

*Department of Chemistry and Biochemistry, University of California San Diego, La Jolla,
California 92093*

E-mail: jperezsa@ucsd.edu; joelyuen@ucsd.edu

Computational Details

For the numerical integration of the TDSE, we employ the MCTDH algorithm implemented in the Heidelberg package^{1,2}. In the multi-set formulation, The wavefunction is expanded in a set of electronic states ($k = a, b$):

$$\langle x, \phi | \Psi(t) \rangle = \sum_{(k=a)}^b \psi^{(k)}(x, \phi, t) |k\rangle, \quad (1)$$

where the wavefunction $\psi^{(k)}(x, \phi, t)$ is represented as a linear combination of Hartree products, each one of them consisting of a product of so-called single-particle functions (SPFs), namely

$$\psi^{(k)}(x, \phi, t) = \sum_{j_x=1}^{n_x} \sum_{j_\phi=1}^{n_\phi} A_{j_x j_\phi}^{(k)}(t) \varphi_{j_x}^{(k)}(x, t) \times \varphi_{j_\phi}^{(k)}(\phi, t), \quad (2)$$

where n_x and n_ϕ are the number of SPFs for the cavity and nuclear degrees of freedom respectively. For each SPF, in turn, a discrete variable representation (DVR) is used.

For an accurate representation of the wavefunction, convergence of the dynamics with respect to the number of SPFs and number of DVR points for each degree of freedom must be ensured. We used $n_x = n_\phi = 4$, and 199 and 150 grid points for the nuclear and photonic degrees of freedom respectively.

Dipole self-energy term

In general, the dipole self-energy term $\frac{g(\phi)^2}{\omega_c}$ should be included in the Hamiltonian (Eq. 4) in order to ensure a bounded ground state of the polaritonic system.³ In Figure S2, we compare the results of the quantum dynamics simulations with and without such term in the Hamiltonian.




Figure S1: Convergence of simulations for 2-photon population for $\omega_c/\omega_{ab} = 0.5$ and $\epsilon = 0.04$. a) With respect to the number of grid points. b) With respect to the number of Single Particle Functions (SPFs).




Figure S2: 2-photon population for $\omega_c/\omega_{ab} = 0.5$ and $\epsilon = 0.04$ with (green) and without (black) the correction dipole self-energy term.

It is evident that such term has almost no influence in the dynamics. This is because our model has no permanent dipole, light-matter coupling is not large enough, and the transition dipole moment is constant for most values of ϕ (see Figure 1).

Polaritonic energy surfaces for $\omega_c > \omega_{ab}/2$ and $\omega_c < \omega_{ab}/2$.

The following figure shows the polaritonic PESs and the adiabatic populations for $\omega_c/\omega_{ab} \neq 0.5$.




Figure S3: Polaritonic PES and population dynamics for numerical simulations at a) $\omega_c/\omega_{ab} = 0.25$, $\epsilon = 0.04$. b) $\omega_c/\omega_{ab} = 0.75$, $\epsilon = 0.04$.

For $\omega_c/\omega_{ab} < 0.5$, the LIAC is reduced because it lies on the region in which transition dipole moment decreases. In addition, the polaritonic PES is steep, which favors a deleterious non-adiabatic transition through the LIAC. The featured dynamics proves this by showing no population of any photonic state. For $\omega_c/\omega_{ab} > 0.5$, the LIAC is large enough to induce oscillation of excitonic and photonic state. However, the nuclear kinetic energy is not high enough to produce a molecular non-adiabatic transition.

Spectral functions

We are interested in the absorption spectrum of the polaritonic system via the molecular transition dipole moment $\hat{\mu}_{ab}$, but also in the two-photon character of the accessible polaritonic eigenstates of $\hat{H}$. The following procedure allows for such calculations.

The first-order perturbative state accessed by a weak laser field at frequency ω is proportional to⁴

$$|\Psi^{(1)}(t, \omega)\rangle = \int_0^t dt' e^{-i\hat{H}(t-t')} \hat{\mu} e^{-i\omega t'} e^{-i\hat{H}t'} |\phi(0)\rangle, \quad (3)$$

where $|\phi(0)\rangle$ is the ground state of the system (molecule + cavity).

Given that $\hat{H}|\phi(0)\rangle = \omega_0|\phi(0)\rangle$, we have

$$|\Psi^{(1)}(t, \omega)\rangle = \int_0^t dt' e^{-i\hat{H}(t-t')} |\Psi(0)\rangle e^{-i(\omega+\omega_0)t'}, \quad (4)$$

where $|\Psi(0)\rangle = \hat{\mu}|\phi(0)\rangle$ is the initial state used in the computational simulations.

Total absorption spectrum

Using Eq. 4, the change of population due to the weak interaction with the EM field out of the cavity is given by

$$\sigma_t(\omega) \equiv \langle \Psi^{(1)}(t, \omega) | \Psi^{(1)}(t, \omega) \rangle = \int_0^t \int_0^t dt' dt'' \langle \Psi(t') | \Psi(t'') \rangle e^{i(\omega+\omega_0)(t''-t')}. \quad (5)$$

Changing variables to $\tau = t'' - t'$ and $\tau' = t'' + t'$ we get

$$\sigma_t(\omega) = t \int_{-t}^t d\tau \langle \Psi(0) | \Psi(\tau) \rangle e^{i(\omega+\omega_0)\tau}, \quad (6)$$

where the integral in the right-hand side gives rise to the standard absorption spectrum:

$$\int_{-t}^t d\tau \langle \Psi(0) | \Psi(\tau) \rangle e^{i(\omega+\omega_0)\tau} = 2 \sum_n |c_n|^2 \frac{\sin(\omega_n - (\omega + \omega_0))t}{(\omega_n - (\omega + \omega_0))}, \quad (7)$$

where $c_n = \langle n | \Psi(0) \rangle$ and $|n\rangle$ is an eigenstate of $\hat{H}$. In the Condon approximation, c_n is the Frank-Condon factor.

Therefore, for long enough (yet finite) propagation times t , $\sigma_t(\omega)$ is proportional to the absorption spectrum

$$\sigma_t(\omega) = 2t \sum_n |c_n|^2 \frac{\sin(\omega_n - (\omega + \omega_0))t}{(\omega_n - (\omega + \omega_0))}. \quad (8)$$

Two photon character from absorption spectrum

Note that Eq. 5 can also be written as

$$\sigma_t(\omega) = \sum_i \int_0^t \int_0^t dt' dt'' \langle \Psi(t') | i \rangle \langle i | \Psi(t'') \rangle e^{i(\omega + \omega_0)(t'' - t')} \equiv \sum_i s_t^{(i)}(\omega), \quad (9)$$

where $\{|i\rangle\}$ is a complete basis set.

In this picture, the absorption spectrum $\sigma_t(\omega)$ is a sum of spectra $s_t^{(i)}(\omega)$. Based on this, for sufficiently large t values, a comparison of both spectra $s_t^{(i)}(\omega)$ and $\sigma_t(\omega)$ gives us an estimation of the overlap of interest $\frac{s_t^{(i)}(\omega)}{\sigma_t(\omega)} = |\langle i | n \rangle|^2 < 1$ via a time-dependent approach. We use these ideas to calculate the two-photon character of eigenstates accessible by one-photon excitation, using $|i\rangle = |2\rangle$ (See Figure 5 in the manuscript).

Breaking of parity symmetry due to non-adiabatic coupling

Products of electronic and photonic states form a convenient basis to represent the Hamiltonian of the system. Using *adiabatic* electronic states ($|g\rangle$ and $|e\rangle$) and Fock states for the cavity model, the Hamiltonian in Eq. 3 of the manuscript can be written as:

$$\mathbf{H} = \mathbf{T}_N + \begin{pmatrix} Vg(\phi) + 0.5\omega_c & \hat{C}_{ge}(\phi) & 0 & g(\phi) & 0 & 0 & \cdot & \cdot & \cdot \\ \hat{C}_{eg}(\phi) & Ve(\phi) + 0.5\omega_c & g(\phi) & 0 & 0 & 0 & \cdot & \cdot & \cdot \\ 0 & g(\phi) & Vg(\phi) + 1.5\omega_c & \hat{C}_{ge}(\phi) & 0 & g(\phi) & \cdot & \cdot & \cdot \\ g(\phi) & 0 & \hat{C}_{eg}(\phi) & Ve(\phi) + 1.5\omega_c & g(\phi) & 0 & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & g(\phi) & Vg(\phi) + 2.5\omega_c & \hat{C}_{ge}(\phi) & \cdot & \cdot & \cdot \\ 0 & 0 & g(\phi) & 0 & \hat{C}_{eg}(\phi) & Ve(\phi) + 2.5\omega_c & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix},$$

where $\mathbf{T}_N$ is a diagonal matrix consisting of the nuclear kinetic energy $-\frac{1}{2m_\phi} \frac{\partial^2}{\partial \phi^2}$, and the diagonal non-adiabatic terms $\hat{C}_{gg}(\phi)$ and $\hat{C}_{ee}(\phi)$ (see Ref. 5). By eliminating the molecular non-adiabatic couplings (red) and the counter-rotating light-matter coupling terms (blue), we recover the Jaynes-Cummings Hamiltonian where the total number of excitations, defined as the sum of electronic and cavity excitations, is conserved. Such Hamiltonian has been previously used in investigations on molecular polaritons such as Refs. 22,23, and 52. In the ultrastrong coupling regime, terms in blue must be considered, and even though the total number of excitations is no longer conserved, the parity in excitation number is. Finally, if molecular non-adiabatic couplings are taken into account, states with differing excitation numbers can couple and neither number of excitations nor parity are longer conserved.

Two-photon population vs pump intensity

Two-photon population vs pump intensity for calculations that involve explicit interaction of the polaritonic system with an external classical field.




Figure S4: Average the two-photon population $\overline{P_{g,2}} = \frac{1}{T} \int_0^T P_{g,2}(t) dt$ ($T = 100$ fs) as a function of the laser intensity I_0 . This calculation ensures that at $I_0 = 3.5 \times 10^{10} \text{ W/cm}^2$, the two-photon generation is linear with respect to the pump, so the (non-linear) down-conversion effect comes only from the polaritonic dynamics.

References

- (1) Worth, G. A.; Beck, M. H.; Jäckle, A.; Meyer, H.-D. The MCTDH Package, Version 8.2. 2000.
- (2) Vendrell, O. Coherent dynamics in cavity femtochemistry: application of the multi-configuration time-dependent Hartree method. *Chem. Phys.* **2018**, *509*, 55 – 65.
- (3) Rokaj, V.; Welakuh, D. M.; Ruggenthaler, M.; Rubio, A. Light–matter interaction in the long-wavelength limit: No ground-state without dipole self-energy. *J. Phys. B* **2018**, *51*, 034005.
- (4) Tannor, D. *Introduction to Quantum Mechanics*; University Science Books, 2007.
- (5) Triana, J. F.; Peláez, D.; Sanz-Vicario, J. L. Entangled Photonic-Nuclear Molecular Dynamics of LiF in Quantum Optical Cavities. *J. Phys. Chem. A* **2018**, *122*, 2266–2278.