

## Antibody-mediated immobilization of virions in mucus

Melanie A. Jensen · Ying-Ying Wang ·  
Samuel K. Lai · M. Gregory Forest ·  
Scott A. McKinley

Received: date / Accepted: date

**Abstract** Antibodies have been shown to hinder the movement of Herpes Simplex Virus (HSV) virions in cervicovaginal mucus (CVM), as well as other viruses in other mucus secretions. However, it has not been possible to directly observe the mechanisms underlying this phenomenon, so the nature of virion-antibody-mucin interactions remain poorly understood. In this work, we analyzed thousands of virion traces from single particle tracking experiments to explicate how antibodies must cooperate to immobilize virions for relatively long time periods. First, using a clustering analysis, we observed a clear separation between two classes of virion behavior: Freely Diffusing and Immobilized. While the proportion of Freely Diffusing virions decreased with antibody concentration, the magnitude of their diffusivity did not, implying an all-or-nothing dichotomy in the pathwise effect of the antibodies. Proceeding under the assumption that all binding events are reversible, we used a novel switch-point detection method to conclude that there are very few, if any, state-switches on the experimental time scale of twenty seconds. To understand this slow state-switching, we analyzed a recently proposed continuous-time Markov

---

NIH R01GM122082-01, R21AI093242, U19AI096398, NSF DMS-1462992, DMS-1517274,  
CAREER Award DMR-1151477

Melanie Jensen  
Department of Mathematics, Tulane University, New Orleans, LA.  
E-mail: mjensen1@tulane.edu

Y.Y Wang  
Department of Biophysics, Johns Hopkins University, Baltimore, Maryland.

S.K. Lai  
Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill,  
NC.

M.G. Forest  
Department of Mathematics, University of North Carolina at Chapel Hill, Chapel Hill, NC.

S.A McKinley  
Department of Mathematics, Tulane University, New Orleans, LA.

chain model for binding kinetics and virion movement. Model analysis implied that virion immobilization requires cooperation by multiple antibodies that are simultaneously bound to the virion and mucin matrix, and that there is an entanglement phenomenon that accelerates antibody-mucin binding when a virion is immobilized. In addition to developing a widely-applicable framework for analyzing multi-state particle behavior, this work substantially enhances our mechanistic understanding of how antibodies can reinforce a mucus barrier against passive invasive species.


**Keywords** Mucosal immunology · Particle tracking · Switching diffusion · Uncertainty quantification

**Mathematics Subject Classification (2010)** 92B05 · 62-07 · 60J70

## 1 Introduction


There are several mechanisms by which antibodies (Ab) produced by the immune system can interfere with and even prevent viral infection after an invasion. Antibodies have long been known to bind to surface epitopes on invading virions, rendering the pathogen ineffective either by blocking the epitope from binding to receptors on target cells, or signaling to other immune cells/molecules to inactivate the virus or destroy virus-infected cells. Recent experiments have revealed a previously under-appreciated mechanism: physical *hindrance* of virion motion and potentially the complete *immobilization* of virions in mucus secretions that lie on the epithelium [20,11]. Specifically, the presence of virion-binding, Immunoglobulin G (IgG) antibody, was shown to directly decrease the mobility of the Herpes Simplex Virus (HSV) virions in human cervicovaginal mucus (CVM) [20,17], as well as Influenza and Ebola virus-like particles in human airway mucus [24]. An example of the effect can be seen in Figure 1, where we display virion trajectories for two populations of HSV virions, originally studied in Wang et al. [20]. The left and right columns show virion movement in the presence of low and high Ab concentrations, respectively. The degree of activity in the low Ab concentration is notably higher.

The possibility of using IgG to hinder the motion of different viruses in mucus provides a novel strategy for immunologists to develop methods to prevent and/or treat viral infection [11,23]. **Population-scale experimental methods have shown that Ab are slightly less mobile in mucus than in phosphate-buffered saline [13].** The reduced diffusivity of Ab in mucus has been attributed to weak transient bonds between individual Ab and the polymeric microstructure of mucus, or “mucin mesh” [13]. Meanwhile many virions have been shown to diffuse unimpeded in mucus in the absence of a detectable Ab concentration [13,20]. **For this reason, the observation that virion mobility in CVM is impeded in the presence of Ab (even across the menstrual cycle) implies there must be some physico-chemical mechanism at work [20,17].**



**Fig. 1** The trajectories of HSV virions for Donor F17 at exogenous antibody concentrations  $0\mu\text{g/mL}$  (left) and  $1.0\mu\text{g/mL}$  (right). **Top Row:** The displacement of HSV virions in the  $x$ -direction. The time indicated in the horizontal axis is shifted for each path so that  $t = 0$  corresponds to the moment the path is first observed. **Bottom Row:** All two dimensional HSV virion trajectories overlaid and plotted in a single frame. For all sub-figures the trajectory frame-rates are 15 observations per second.

Recently the authors and collaborators have explored the possibility that Ab can work in tandem with the mucin mesh to hinder diffusing virions. (See Figure 2 for an idealized schematic of the interactions.) In theory, as a virion diffuses through mucus, an array of Ab can accumulate on its surface. When a sufficient number of virion-bound Ab form low affinity bonds to the mucin mesh, the virion can become tethered and essentially trapped. **This hypothesis was introduced by Olmsted et al. in 2001 [13] and confirmed by Wang et al. in 2014 [20], by Newby et al. in 2017 [11], and by Schroeder et al. in 2018 [17].** In 2014, Chen et al. [2] introduced a stochastic/deterministic hybrid model for the immobilization of Human Immunodeficiency Virus (HIV) by IgG in CVM, and demonstrated the potential impact of the tandem effect of Ab-virion binding and Ab-mucus transient binding on the ability of viral populations to cross, enter, and pass through a thin mucosal layer. Later, Wessler et al. [22] used numerical simulations to explore combinations of Ab-virion and Ab-mucus reaction kinetics that produce an optimal effect. Newby et al. [11] further demonstrated that very low affinity Ab-mucus bonds optimize trapping of diffusing nanoparticles using experimental and simulated data along with providing theoretical arguments.



**Fig. 2** A schematic depiction of the proposed immobilization process of virions, green circles, by antibodies, blue 'Y's, in a mucosal medium. Virions become immobilized when 'enough' antibodies are bound to the virions and the mucosal fibers, gray lines. Arrows indicate Ab interacting solely with the mucin fibers. Figure originally presented in [20].


Underlying these mathematical models is a *Switching Diffusion Hypothesis*: that the chemical reactions responsible for virion (or nanoparticle) immobilization are reversible and, as a consequence, virions should switch between diffusive and immobilized states. When compared to the experimentally observable timescale of 10-20 seconds, the Ab-mucin kinetic rates are expected to be fast while the Ab-virion kinetic rates are expected to be slow (see Table 1). It is not clear, however, whether the state-switching between diffusion and immobilization should be on a faster or slower timescale than the observable 10-20 seconds.

In recent modeling efforts, [2,22], the effect of surface bound Ab on the diffusivity of a virion was assumed to be incrementally multiplicative. That is to say, there is a constant  $\alpha \in (0, 1)$  such that the diffusivity has the following state-dependent form.

$$\text{Incremental Knockdown Hypothesis: } D(S(t), N(t)) = \alpha^{N(t)} D. \quad (1)$$

Here  $D$  is the diffusivity of the virion in mucus in the absence of Ab,  $N(t)$  is the number of virion-bound Ab, and  $S(t)$  is the subset of Ab simultaneously bound to the mucin mesh. This reduction in diffusivity is independent of  $S(t)$  because the number of simultaneously bound Ab changes so rapidly (relative to the number of bound Ab), the virion only feels the average effect of these changes, which is captured by the number of bound Ab,  $N(t)$ . The parameter  $\alpha$  can be expressed in terms of the Ab-mucin binding and unbinding rates ( $m_{\text{on}}$  and  $m_{\text{off}}$ , respectively) and the effective concentration  $[M]$  of binding sites on the surfaces of mucin fibers. If  $m_{\text{on}}[M]$  and  $m_{\text{off}}$  are very large, so that there are many on-and-off switches per second, then an effective diffusivity arises with a so-called "knockdown factor"  $\alpha = m_{\text{off}}/(m_{\text{on}}[M] + m_{\text{off}})$  [2]. In this way, we say that the Incremental Knockdown Hypothesis follows from assuming that the dynamics is in a *Fast Switching Regime*. That is to say, in this modeling regime, one assumes that [diffusion  $\rightleftharpoons$  immobilization] switching is faster than

the times between experimental observations and faster than what we plan to use as a simulation time step. We depict a typical trajectory of a virion under this hypothesis in Figure 3(a). A virion rapidly changes between the immobilized (red) state and freely diffusing states (green). The resulting path has a reduced *effective diffusivity* that is well-approximated by Equation 1, and the virion exhibits qualitatively less movement than a virion predominately in the freely diffusing state (seen in blue).



**Fig. 3** **Top row** (a)-(c): The path of a virion assuming it takes one (green trajectory) or ten (blue trajectory) simultaneously bound Ab for immobilization. Red intervals correspond to periods of immobilization. **Bottom Row** (d)-(f): The virion-Ab-mucin dynamics that govern the movement of the simulated virion directly above it. Within each frame, the number of bound Ab  $N(t)$  is shown by the purple trajectory and the subset of these Ab that are simultaneously bound to the mucin fibers  $S(t)$  assuming a low threshold,  $T = 1$ , and higher threshold,  $T = 10$ , shown by the green trajectory and blue trajectory, respectively. The binding rate cascade factor  $c$  increases from left to right:  $c = 1$ ,  $c = 20$  and  $c = 200$ , respectively. Other model parameters used in the simulation are  $([A]_0, [A]_{\text{exo}}, N_*) = (0.2\mu\text{g}/\text{mL}, 0.1\mu\text{g}/\text{mL}, 120)$ . The mathematical model is fully described in Section 2.4.

Recent particle tracking experiments now make it possible to analyze virion behavior as it is modulated by various concentrations of Ab [20]. In Figure 1, we display two populations of HSV virions diffusing in CVM with  $0\ \mu\text{g}/\text{mL}$  and  $1\ \mu\text{g}/\text{mL}$  concentrations of exogenous HSV-binding IgG. There is qualitatively less virion movement in CVM with higher concentrations of Ab, but, as we argue below using path-by-path analysis, the trajectories of individual virions appear to resemble either that of a strictly immobilized virion or a strictly freely diffusing virion. This absence of observable switches between immobilized and freely diffusing states might seem to ratify the fast switching hypothesis. However, closer analysis of the freely diffusing particles shows that the diffusivity of freely diffusing virions is essentially the same across all exogenous Ab concentrations. This contradicts the Incremental Knockdown

Hypothesis, which predicts the diffusivity should decrease with increasing Ab concentration. While there are essentially no observable switches, and the diffusivity of the free population is not incrementally affected by Ab concentration, we find that the proportion of completely immobilized virions is unmistakably increasing with Ab concentration. (See also [20].) This suggests an alternative hypothesis: we are in a *Slow Switching Regime* where switching takes place fast enough (less than the incubation period of thirty minutes) so that the experiments display different movement patterns, but slow enough (more than twenty seconds) so that we do not see switches in the observational time window.

In this work, we develop and implement the tools necessary for making the preceding claims. To be specific, we use clustering analysis to partition virion paths into a few distinct behavioral patterns. We implement a Bayesian switch-point detection algorithm to assess the prevalence of switches in mobile virions. We develop a Markov chain model for virion-Ab-mucin interactions for use in our characterization of the dependence of virion motility on Ab concentration. A critical feature of this model is the possibility that virion immobilization requires multiple simultaneously surface bound Ab, and that a single virion-Ab-mucin binding event might lead to a cascade of such binding events, which would serve to enhance trapping. Using uncertainty quantification techniques we explore the limitations of the available data, but argue there is a reasonable parameter regime that is fully consistent with experimental observations.

## 2 Data Collection, Statistical Methods, and Mathematical Model

### 2.1 Data collection

Single particle tracking data of HSV virions was collected from seven different CVM samples at five added doses of exogenously anti-HSV-1 IgG, (0, 0.033, 0.1, 0.333, 1.0)  $\mu\text{g}/\text{mL}$  with an incubation period of half an hour to one hour. For each sample, the virions were tracked for a duration of 20 seconds. The  $x$ -position and the  $y$ -position of all traces were observed at a time interval of  $\delta = 1/15\text{s}$ . **For all the experiments, the tracked particles were the HSV-1 mutant 166v, containing a VP22-GFP tegument protein. The translation motion of the fluorescent virions was captured with MetaMorph imaging software resulting in videos (512x512 pixels, 16-bit image depth) with temporal solution 66.7 ms and spatial resolution of 10nm for 20 seconds.** For a more detailed description of the collection process see the Methods section in [20].

### 2.2 Statistical tools for virion trajectory analysis

We used standard statistical techniques to assess whether the behavior of each virion is consistent with the defining properties of Brownian motion (stationarity with Gaussian independent increments) and to infer physical parameters.

### 2.2.1 Test for Gaussianity and independence of increments

We used normal quantile-quantile (**qqnorm**) plots to qualitatively verify that the path statistics are approximately Gaussian. The **qqnorm** plots for the increment processes had approximately linear relationships for all particles indicating the  $x$  and  $y$  increment processes for all particles could be described as Gaussian. **To construct such plots, we used the `qqnorm()` function in the R programming language found in the stats library.**

Noting that if a Gaussian process has uncorrelated increments then the increments are independent, we tested for independence of increments by quantifying the statistical significance of their correlation. Let  $\{U_i(k) := X_i(k\delta) - X_i((k-1)\delta)\}_{k=1}^n$  and  $\{V_i(k) := Y_i(k\delta) - Y_i((k-1)\delta)\}_{k=1}^n$  denote the  $i$ th particle's  $x$  and  $y$  increment processes, respectively. For the  $i$ th particle, we estimated the correlation between the  $x$  and  $y$  increment processes separated  $h$  time steps apart using the sample autocorrelation function,  $\mathcal{A}_i(h; U)$  and  $\mathcal{A}_i(h; V)$  used in the R programming language. If there are  $n$  increments of uniform duration  $\delta$  then for a time lag of  $h\delta$

$$\mathcal{A}_i(h; X) := \frac{\frac{1}{n} \sum_{k=1}^{n-h} (U_i((k+h)\delta) - \bar{U}_i)(U_i(k\delta) - \bar{U}_i)}{\frac{1}{n} \sum_{j=1}^n (U_i(j\delta) - \bar{U}_i)^2}, \quad (2)$$

where  $\bar{U}_i := \frac{1}{n} \sum_{k=1}^n U_i(k)$  [18]. We say the  $i$ th particle's increment processes are anti-persistent (persistent) if both  $\mathcal{A}_i(h = 1; X)$  and  $\mathcal{A}_i(h = 1; Y)$  are below (above) the critical value for a 95% significance level and independent otherwise.

### 2.2.2 Mean-Squared Displacement

The primary statistical tool for describing a population of microparticle paths is the so-called *ensemble* mean-squared displacement (MSD), which we denote  $\langle \mathcal{M}(t) \rangle$ . To calculate it, we first compute a *pathwise* MSD for each trajectory (denoted  $\mathcal{M}_i(t)$  for the  $i$ th path) and then take an average over these functions. If there are  $n$  steps that are uniform of duration  $\delta$ , then as defined in [15],

$$\mathcal{M}_i(k\delta) := \frac{1}{n-k+1} \sum_{j=0}^{n-k} |X_i((j+k)\delta) - X_i(j\delta)|^2.$$

For  $t$  between the time points  $\{k\delta\}$  we define  $\mathcal{M}_i(t)$  by linear interpolation. The slope of the MSD displayed on a log-log scale provides an estimate for each particle's diffusive exponent,  $\nu$ , in the large time regime ( $\mathcal{M}_i(t) \sim Ct^\nu$ ). Following standard particle tracking nomenclature, an individual path is said to be Brownian if  $\nu = 1$ , subdiffusive if  $\nu \in (0, 1)$ , and stationary if  $\nu = 0$ .

### 2.2.3 Effective Diffusivity

A fundamental quantity to measure for a Brownian path is its *diffusivity*  $D$ . If  $(X(t), Y(t))$  is the  $2d$  position of the particle at time  $t$ , then its diffusivity is defined to be  $D := \lim_{t \rightarrow \infty} \mathbb{E}(X^2(t) + Y^2(t))/4t$ . For a Brownian path with  $n$  steps of uniform duration  $\delta$ , the maximum likelihood estimator (MLE) for its diffusivity has the form

$$D_{\text{eff}} := \frac{1}{4\delta n} \sum_{j=1}^n (U(j\delta)^2 + V(j\delta)^2), \quad (3)$$

shown in Appendix A. We refer to  $D_{\text{eff}}$  as the path's *effective diffusivity*. We note that this effective diffusivity is only a consistent estimator for  $D$  if the path has all the characteristics of Brownian motion, namely stationary, independent, Gaussian increments.

However, as seen in Figure 4(a)-(c) there are many paths with anti-correlated increments. For such a process, “diffusivity” is not well-defined. **For those paths that can be described by Brownian motion,  $D_{\text{eff}}$  does not account for observational error. Using the method proposed by Vestergaard et al. [19], we found a typical variance of the localization error to be minor (of order  $0.01\mu\text{m}^2/\text{s}$ ) as compared to the effective diffusivity (of order  $1\mu\text{m}^2/\text{s}$ ) for such virions, see the SI Section 2 for further details and supporting figure.** Nevertheless we use  $D_{\text{eff}}$  as a descriptor for these paths because this serves the purpose to distinguish between the particles in two different states by the clustering methods described below.

For a given collection of  $N$  particles, the *ensemble effective diffusivity* is the weighted average effective diffusivity of the tracked particles in the sample, denoted  $\langle D_{\text{eff}} \rangle$ . When evaluating population statistics in particle tracking experiments, if particle paths are weighted independent of path length, then it has been shown that there is a bias toward highly mobile particles, further discussed in Section 2.3.1,[21]. Based on that analysis, we report the effective diffusivity of an ensemble by taking an average weighted by path lengths. Let  $D_{\text{eff}}^i$  denote the effective diffusivity of  $i$ th freely diffusing virion, which has path length  $n_i$ . Then

$$\langle D_{\text{eff}} \rangle := \sum_{i=1}^N \omega_i D_{\text{eff}}^i \quad \text{where } \omega_i = \frac{n_i}{\sum_{j=1}^N n_j}. \quad (4)$$

### 2.2.4 Bias-corrected and accelerated percentile ( $BC_a$ ) confidence interval method

We constructed confidence intervals for ensemble statistics based on the bootstrapping  $BC_a$  method due to its second order accuracy and invariance under transformations. See [4] for the formulation of confidence intervals using this method. **We used the `boot.ci()` function in the R programming language found in the in the `boot` library to obtain the  $BC_a$  confidence**

**intervals for the ensemble statistics as follows.** First, we simulated 10,000 booted samples (with replacement) from an ensemble of  $N$  tracked particles weighted by the particle path lengths. The  $BC_a$  confidence interval is then the usual confidence interval constructed using this population of (weighted) bootstrap samples.

### 2.3 Classification scheme for virion paths

For each donor and concentration, we employed a hierarchical clustering algorithm to separate the HSV virions into distinct clusters based on a set of pathwise statistics:  $x$ -increment ACF,  $y$ -increment ACF, and log 10 transform of the effective diffusivity. We defined the dissimilarity measure between pairs of virions  $i$  and  $j$  by a weighted Euclidean distance  $d(i, j)$  with weights of 1/4, 1/4, and 1/2 for the differences in  $\mathcal{A}_i(1; X)$ ,  $\mathcal{A}_i(1; Y)$ , and  $\log 10(D_{\text{eff}})$  respectively. The dissimilarity measure between clusters was set to be the average linkage. That is to say, the dissimilarity between clusters  $R$  and  $Q$  is defined to be

$$d(R, Q) = \frac{1}{|R||Q|} \sum_{i \in R, j \in Q} d(i, j). \quad (5)$$

Hierarchical clustering is an agglomerative clustering method [8]. The algorithm is initialized by setting each data point as a distinct cluster. During each iteration, clusters are merged together to minimize the dissimilarity between all clusters. The algorithm stops when all data points are in a single cluster. This process is depicted graphically through the dendrogram where clusters merge at a height equal to dissimilarity between them. We obtained the  $k$  cluster by cutting the resulting dendrogram at the uniform height yielding  $k$  clusters.

**Because hierarchical clustering is an unsupervised clustering method, in which the number of clusters is not known a priori, the number of clusters has to be chosen by the practitioner. One such method of obtaining the number of clusters is called the elbow method. In this approach, the within-sum-of-squares values of the clusters (WSS) is computed and plotted for a range of cluster numbers. The WSS decreases with the number of clusters, and typically there is a bend or “elbow” in the graph which guides the selection of an appropriate number of clusters.**

In all cases, there was a major drop in WSS from one to two clusters but it was rarely clear how to specify the elbow among  $k = 2, 3, 4$  or 5 clusters. We chose to use  $k = 4$  in almost all cases because the results were consistent with our bio-physical intuition that there might be Freely Diffusing, Immobilized, Subdiffusive, and Outlier states. A few examples of each class are displayed in the supplemental materials, Figure 1. Although we allowed for four clusters when labeling each cluster with a biological classification, clusters

were typically merged together as the subdiffusive and outlier class were few in number.

### 2.3.1 ‘Frame-by-frame’ method to compute empirical distribution of each cluster

It has been shown in [21] that fast moving particles are overestimated on shorter time scales in 2d particle tracking. This bias towards the fast moving particles arises due to individual fast particles leaving and reappearing in the focal plane as distinct traces and to new particles entering and leaving the focal plane throughout the duration of the experiment. To minimize overestimating the freely diffusing population, we employed the ‘frame-by-frame’ method developed in [21] to compute the fraction of each population present in the data. The ‘frame-by-frame’ method assigns each tracked particle a weight based on the number of frames the particle appears in the field of view, whereas in the conventional method each particle has the uniform weight of one. Under this weighting system, for a sample of size  $N$ , the weighted sample proportion of the  $i$ th state is given by

$$\hat{p}_i = \sum_{k=1}^N \omega_k \delta_{ik} \quad \text{for} \quad \omega_k = \frac{n_k}{\sum_{k=1}^N n_i} \quad (6)$$

where  $\delta_{ik}$  is the Kronecker delta function.

## 2.4 Mathematical model for asymptotic probability of immobilization

We mathematically model the dynamics of a virion under the Switching Diffusion Hypothesis by the following SDE:

$$dX(t) = \sqrt{2D(N(t), S(t))} dW(t) \quad (7)$$

where  $W(t)$  is standard 2d Brownian motion and the state-dependent diffusivity,  $D(N(t), S(t))$ , depends on two time-dependent processes:  $N(t)$ , the number of antibodies bound to the surface of a focal virion at time  $t$ , and  $S(t)$ , the subset of these antibodies simultaneously bound to mucin binding sites at time  $t$ . We establish a threshold parameter  $T$ . A virion is defined to be *immobilized* if there are at least  $T$  simultaneously bound antibodies,  $S(t) \geq T$ , and defined to be *freely diffusing* if there are fewer than  $T$  simultaneously bound antibodies,  $S(t) < T$ . Under this convention, the time-dependent diffusivity is given by

$$D(N(t), S(t)) = \begin{cases} D & 0 \leq S(t) < T \\ 0 & T \leq S(t) \leq N(t) \end{cases}$$

where the constant  $D$  is the diffusivity of the virion in mucus in the absence of Ab. In the following sections, we present a mathematical model that describes the asymptotic probability of the immobilized state when exposed to varying exogenous antibody concentrations.

#### 2.4.1 Model assumptions

Based on the initial population clustering analysis, there appears to a sub-population of virions that do not interact with the antibodies. We define  $q$  to be the probability that a given virion will interact with the Ab population. Second, for the sake of simplicity, we assume that Ab-virion binding sites operate independently from each other. However, we allow for **cooperativity** among the Ab in binding to the mucosal environment. Once the virion has  $T$  simultaneously bound Ab-mucin-virion interactions ( $S \geq T$ ) the surface bound antibodies might bind to the mucin fibers differently than if the virion was freely diffusing. We parametrize this by a multiplicative change in Ab-mucin binding rate through the introduction of the dimensionless parameter  $c$ . If  $c > 1$  the parameter has a cascade effect, aiding in the immobilization process [3, 6, 7, 9, 11].

#### 2.4.2 A Markov Chain model for virion-Ab-mucin dynamics

Let  $N_*$  denote the number of independent Ab binding sites on the surface of an HSV virion. Antibodies bind and unbind from these sites at rates  $k_{\text{on}}$  and  $k_{\text{off}}$ , respectively, with dissociation constant  $k_d := k_{\text{off}}/k_{\text{on}}$ .

Virion-surface-bound antibodies interact with the surrounding mucosal medium, binding to and unbinding from mucin binding sites, at rates  $m_{\text{on}}$  and  $m_{\text{off}}$ , with dissociation constant  $m_d := m_{\text{off}}/m_{\text{on}}$ . The total Ab concentration  $[A]$  is the sum of the exogenous  $[A]_{\text{exo}}$  and endogenous  $[A]_0$  Ab concentrations, and the total concentration of binding sites on mucin fibers is denoted  $[M]$ . See Table 1 for a comprehensive list of variables.

**Table 1** Parameters and known values incorporated in the model. \* indicates that the value has not been directly measured. The given value is chosen to be consistent with indirect observations.


Parameter	Symbol	Value	Reference
Cell Properties			
Initial Ab concentration in CVM	$[A]_0$	Model Parameter	
Concentration of Ab binding sites on mucin fibers in CVM	$[M]$	unknown	*
	$m_{\text{on}}[M]$	$11.1 s^{-1}$	*
Molecule Properties			
bnAb (IgG) Diameter		$0.011 \mu\text{m}$	[20]
HSV-1 Diameter		$\sim 0.180 \mu\text{m}$	[20]
Number of Ab binding sites on HSV-1	$N_*$	Model Parameter	*
Reaction Kinetics			
Ab-mucin affinity (Knockdown Factor)	$\alpha$	0.9	[13]
Ab-mucin binding rate	$m_{\text{on}}$	Unknown	*
Ab-mucin unbinding rate	$m_{\text{off}}$	$100 s^{-1}$	*
Ab-virion binding rate	$k_{\text{on}}$	$4.26 \times 10^4 [M]^{-1} s^{-1}$	[2]
Ab-virion unbinding rate	$k_{\text{off}}$	$2.87 \times 10^{-4} s^{-1}$	[2]
Change in (Ab-virion)-mucin binding rate after immobilization	$c$	Model Parameter	*
Number of Ab bond to mucus to immobilize a virion	$T$	Model Parameter	*

We model the the Ab-virion interactions using a continuous time Markov Chain (CTMC) assuming linear state transitions. If a given virion has  $n$  oc-

cupied (Ab-bound) surface binding sites at time  $t$ , then the CTMC transition rates are given by



If there are  $s$  simultaneously bound Ab cross-linking the virion to mucin fibers at time  $t$  and  $n$  occupied virion-surface-binding sites, then the conditional Ab-mucin dynamics are modeled by a CTMC with state transition rates



for  $s \leq n$ , where

$$g(s) = \begin{cases} 1 & s < T \\ c & s \geq T. \end{cases} \quad (10)$$

The function in Equation 10 quantifies the impact immobilization has on the rate at which additional antibodies crosslink to the mucin fibers, i.e. the binding cascade effect, and results in a non-linear transition rate when  $c \neq 1$ . We note that the transition  $(n, s) \rightarrow (n - 1, s - 1)$  is omitted from our analysis to facilitate with explicit likelihood calculations. This does not qualitatively affect our results **because of the time-scale separation between Ab-virion and Ab-mucin kinetics. Since we assume that (conditioned on number of Ab-virion bindings  $N$ ) the stationary distribution of  $S$  is achieved rapidly, the initial “error” using  $S = s$  instead of  $S = s - 1$  after an  $N = n \rightarrow n - 1$  transition does not affect long-term dynamics.**

We show the impact the immobilization threshold,  $T$ , and the cascade factor,  $c$ , have on the immobilization process in Figure 3. Within each frame, it can be seen that a higher immobilization threshold allows for longer freely diffusion periods, while across frames a higher cascade factor leads to longer immobilized periods. In Figure 3(d)-(f) we simulated realizations of the processes  $(N(t), S(t))$  for various combinations of  $T$  and  $c$ . The number of bound antibodies,  $N(t)$ , is displayed by the purple trajectory, and the number of simultaneously bound Ab with a low immobilization threshold,  $S(t)$  when  $T = 1$ , and with a higher immobilization threshold,  $S(t)$  when  $T = 10$ , are shown by the green and blue trajectory, respectively. Moving left to right, the factor by which the Ab-mucin binding rate changes after immobilization increases,  $c = 1, 20$ , and  $200$ , respectively. In Figure 3(a)-(c), we show how these processes dictate the movement of the virion. The virion with process  $(N(t), S(t))$  when  $T = 1$  is colored in green while  $(N(t), S(t))$  when  $T = 10$  is colored in blue. For both trajectories immobilized periods,  $S(t) \geq T$ , are colored in red.

When immobilization does not affect the Ab-mucin binding rate, Figure 3(d), the process  $S(t)$  rapidly crosses the immobilization threshold (dashed line) resulting in a virion transitioning between states faster than the experimental time step, Figure 3(a), for both  $T = 1$  and  $T = 10$ . By increasing the cascade factor, Figure 3(e)-(f),  $S(t)$  remains above the immobilization threshold, for observable periods. In this case, the simulated virions in Figure

3(b)-(c) change states on the experimental time scale of twenty seconds and longer than twenty seconds, respectively.

#### 2.4.3 Our approximation for the stationary probability of being immobilized

We assume that the antibody-virion dynamics are slow compared to the antibody-mucin dynamics. To approximate a virion's long-term probability of being immobilized, we use a product of two factors. The first is the steady-state distribution for the number of surface-bound Ab,  $N(t)$ . Then we compute the stationary distribution for the number of simultaneously bound Ab,  $S(t)$ , conditioned on each value  $N(t) = n$  (where  $n \in \{0, \dots, N_*\}$ ).

We introduce the notation  $b(x, n, p)$  for the binomial probability mass function. That is, if  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{P}\{X = x\} = b(x, n, p)$ . Our approximation to the stationary distribution of immobilization can be understood as an average over the transitions of the fast process  $S(t)$ . Let  $\sigma$  denote the time a particle spends in the immobilized state, and  $\tau$  the time a particle spends in the freely diffusing state. Then our approximation takes the form

$$\tilde{\pi}([A]_{\text{exo}}) = q \sum_{n=T}^{N_*} \frac{\mathbb{E}(\sigma; T, c, n)}{\mathbb{E}(\sigma; T, c, n) + \mathbb{E}(\tau; T, n)} b\left(n; N_*, \frac{[A]_0 + [A]_{\text{exo}}}{k_d + ([A]_0 + [A]_{\text{exo}})}\right) \quad (11)$$

where

$$\mathbb{E}(\sigma; T, c, n) = \frac{1}{T m_{\text{off}}} \frac{\sum_{s=T}^n b\left(s; n, \frac{c m_{\text{on}}[M]}{m_{\text{off}} + c m_{\text{on}}[M]}\right)}{b\left(T; n, \frac{c m_{\text{on}}[M]}{m_{\text{off}} + c m_{\text{on}}[M]}\right)}, \quad (12)$$

and  $\mathbb{E}(\tau; T, c, n) = \frac{1}{(n-T+1)m_{\text{on}}[M]} \frac{\sum_{s=0}^{T-1} b\left(s; n, \frac{m_{\text{on}}[M]}{m_{\text{off}} + m_{\text{on}}[M]}\right)}{b\left(T-1; n, \frac{m_{\text{on}}[M]}{m_{\text{off}} + m_{\text{on}}[M]}\right)}.$

The derivation of Equation 11 and Equation 12 rely on Markov Chain Theory and Renewal Theory and can be found in Appendix B.

It follows from the law of total expectation and the time-scale approximation, the expected time immobilized and expected time freely diffusing are respectively:

$$\begin{aligned} \mathbb{E}(\sigma) &= \sum_{n=T}^{N_*} \mathbb{E}(\sigma; T, c, n) b\left(n; N_*, \frac{[A]}{k_d + [A]}\right); \\ \mathbb{E}(\tau) &= \sum_{n=T}^{N_*} \mathbb{E}(\tau; T, c, n) b\left(n; N_*, \frac{[A]}{k_d + [A]}\right). \end{aligned} \quad (13)$$

We say that a parameter vector is in the *Slow Switching Regime* if, for all tested exogenous Ab concentrations, the average times spent in the immobilized and diffusing states are more than 20 seconds. To be precise, we define

$$\Theta_{\text{slow}} := \{\boldsymbol{\theta} : \mathbb{E}(\sigma; [A]_{\text{exo}}, \boldsymbol{\theta}) > 20 \text{ and } \mathbb{E}(\tau; [A]_{\text{exo}}, \boldsymbol{\theta}) > 20 \text{ for all } [A]_{\text{exo}} \in [0, 1]\}. \quad (14)$$

## 2.5 Switch point detection

We develop an algorithm for detecting whether there is a *single* switch from diffusion to immobilization or immobilization to diffusion. The mathematical model presented in Section 2.4, Equation 7, assumes complete immobilization but in fact immobilized virions exhibit spatial motion. Bernstein and Fricks in [1] account for this spatial motion by describing the bound state as a diffusing particle trapped in a potential well. Using an Expectation-Maximization algorithm they provide an evolving probability for each particle that it is in an immobilized or diffusing state. In contrast to the many-switch paths considered by Bernstein and Fricks, we argue in Section 3.1.2 that the virion paths in our data set have at most one or two switches. We therefore developed and implement a Bayesian algorithm that is designed to identify the presence of a single switch point.

To derive a likelihood function, we extend our SDE model Equation 7 to include a path-specific trapping potential well, similar to [1]. Our extended model for a [diffusion  $\rightarrow$  immobilization] switch is

$$d\mathbf{X}(t) = \begin{cases} \sqrt{2Dd}\mathbf{W}(t) & 0 \leq t \leq \tau \\ -\tilde{\kappa}(\mathbf{X}(t) - \mathbf{X}(\tau))dt + \sqrt{2Dd}\mathbf{W}(t) & \tau < t \end{cases} \quad (15)$$

$$\mathbf{X}(0) = 0$$

and for [immobilization  $\rightarrow$  diffusion], we have

$$d\mathbf{X}(t) = \begin{cases} -\tilde{\kappa}(\mathbf{X}(t) - \mathbf{X}(0))dt + \sqrt{2Dd}\mathbf{W}(t) & 0 \leq t \leq \tau \\ \sqrt{2Dd}\mathbf{W}(t) & \tau < t \end{cases} \quad (16)$$

$$\mathbf{X}(0) = 0$$

where  $\mathbf{X}(t) = (X(t), Y(t))^T$  and  $\mathbf{W}(t)$  is 2d Brownian Motion. These SDEs are derived from the Langevin equation for particles diffusing in a quadratic (Hookean spring) potential well. The constant  $\tilde{\kappa} = \kappa/\gamma$  where  $\kappa$  is the spring constant and  $\gamma$  is the viscous drag experienced by the particle. Due to the Fluctuation-Dissipation relationship,  $\gamma$  also appears in the diffusivity constant, which has the form  $D = k_B\mathcal{T}/\gamma$ , where  $k_B$  is Boltzmann's constant and  $\mathcal{T}$  is the temperature of the fluid. To obtain an analytically trackable likelihood function, we introduce simplifying assumptions that (1) the switch occurs exactly at an observation time point, and (2) there is no measurement error. We derive the likelihood function in Appendix C.

We take a Bayesian approach to jointly estimate  $D$ ,  $\tilde{\kappa}$ , and  $\tau$  under both switching scenarios using a Gibbs sampling algorithm. If the 95% credible region for  $\tau$  is completely contained within the interval  $[0.1T_{\text{final}}, 0.9T_{\text{final}}]$  where  $T_{\text{final}}$  is the duration of a path, then we say that path is a candidate for switching. For both switching scenarios we estimated a false discovery rate for this criterion by simulating freely diffusing particles and setting the false discovery rate to the percent of simulated Brownian particles that were labeled as candidates for switching for the given switching model, Equation 16 or 15.

Similarly, we estimated the power of criterion through simulation. For both scenarios we simulated particles that switched states once, and set the power to the fraction of paths that were candidates for switching. See Section 5 for more details on how these tests were constructed, and the results are presented in Section 3.1.2.

## 2.6 Uncertainty quantification

The model given by Equation 11 depends on the parameter vector  $\boldsymbol{\theta} = (T, c, N_*, q, [A]_0, k_d, m_{\text{off}}, \alpha)$ . In specifying the model to HSV-IgG data, Section 2.1, we set  $k_d = 0.8969$  [10] and  $\alpha = 0.90$  [13]. The Ab-mucin binding and unbinding rates have not been directly estimated. We assume they are fast compared to the experimental time scale and, for example, set  $m_{\text{off}} = 100s^{-1}$ . To assess the remaining parameters,  $\boldsymbol{\theta} = (T, c, N_*, q, [A]_0)$  – which are the immobilization threshold value, the binding cascade factor, the number of sites on the surface of virions, the virion-Ab interaction probability, and the endogenous Ab concentration – we employed the numerical method of profile likelihoods [5, 16]. We used the numerically obtained relationships among parameters to obtain conditions on  $\boldsymbol{\theta}$  such that the Switching Diffusion Hypothesis (in the Slow Switching Regime) is consistent with the data.

In order to quantify the model's error in predicting the immobilized fraction, for each donor  $i$ , we partitioned the paths according to exogenous Ab concentration  $\{[A]_j\}_{j=1}^5$ , and introduced the following residual function:

$$\chi_i^2(\boldsymbol{\theta}) = \sum_{j=1}^5 N_{ij} \frac{(\tilde{\pi}([A]_j; \boldsymbol{\theta}) - \hat{p}_{ij})^2}{\tilde{\pi}([A]_j; \boldsymbol{\theta})(1 - \tilde{\pi}([A]_j; \boldsymbol{\theta}))}, \quad (17)$$

where  $\tilde{\pi}([A]_j; \boldsymbol{\theta})$  denotes the model evaluated at  $[A]_j$  with parameters  $\boldsymbol{\theta}$  (as defined in Equation 11), while  $N_{ij}$  and  $\hat{p}_{ij}$  are, respectively, number of paths observed and the fraction that are immobilized in the  $j$ th subpopulation associated with donor  $i$ . Assuming a normal approximation to the binomial distribution, our residual function can be seen as the sum of five independent squared normal random variables, i.e. with a  $\chi^2$ -distribution with 5 degrees of freedom.

### 2.6.1 Numerical method of profile likelihoods to deduce parameter identifiability

Because we assume normal approximation to the binomial distribution, working with a residual function is equivalent to using a likelihood function to define confidence intervals [14, 16]. For ease of notation in this section, we will suppress the dependence on  $i$  when considering the residual function  $\chi^2(\boldsymbol{\theta})$  for donor  $i$ .

To discuss identifiability of our model parameters, we use the nomenclature introduced by Raue in [16]. Our minimum residual estimator is defined to be

$\hat{\theta} := \text{argmin}[\chi^2(\boldsymbol{\theta})]$ . The *likelihood-based confidence region of level  $\alpha$*  for  $\boldsymbol{\theta}$  is then defined to be

$$\Theta_{\alpha,df} := \{\boldsymbol{\theta} : \chi^2(\boldsymbol{\theta}) - \chi^2(\hat{\boldsymbol{\theta}}) < \chi^2(\alpha, df)\}, \quad (18)$$

where  $\chi^2(\alpha, df)$  is the  $\alpha$  quantile of the  $\chi^2$  distribution with  $df$  degrees of freedom. When establishing a confidence interval for one of the parameters, we set  $df = 1$ . When establish a confidence region for multiple parameters, we set  $df$  equal to the number of parameters [14].

A parameter  $\theta_k$  is said to be *structurally identifiable* when there is a unique minimum of  $\chi^2(\boldsymbol{\theta})$  with respect  $\theta_k$ , i.e., if there exists a unique  $\theta_k$  such that

$$\theta_k = (\text{argmin}_{\boldsymbol{\theta} \in \mathbb{R}^5} \{\chi(\boldsymbol{\theta})\})_k.$$

Alternatively,  $\theta_k$  can be unidentifiable due to the structure of the model or because the quality and quantity of the data is insufficient in estimating  $\theta_k$ . For the former case, we say  $\theta_k$  is *structurally unidentifiable* if the set

$$\theta_{\min} := \{\boldsymbol{\theta} : \chi(\boldsymbol{\theta}) = \min_{\vartheta \in \mathbb{R}} \chi(\vartheta)\}$$

is not unique and contains at least two elements whose  $\theta_k$  components are distinct. This often occurs when there is a functional relationship  $\phi$  among  $\theta_k$  and at least one other parameter, say  $\theta_j$  such that  $\chi$  can be expressed directly in terms of  $\phi(\theta_k, \theta_j)$ . As for the latter data-restricted type of unidentifiability, we say  $\theta_k$  is *practically unidentifiable* when a unique minimum exists of  $\chi^2(\boldsymbol{\theta})$  with respect  $\theta_k$  but the likelihood based confidence interval for  $\boldsymbol{\theta}$  extends infinitely in increasing and/or decreasing values of  $\theta_k$ .

These definitions can be interpreted graphically using profile likelihoods. For residual function  $\chi^2(\boldsymbol{\theta})$  the *profile likelihood* of the  $k$ -th parameter defined to be

$$\chi_{\text{PL}}^2(\theta_k) = \min_{\theta_{j \neq k}} [\chi^2(\boldsymbol{\theta})]. \quad (19)$$

If  $\theta_k$  is a structurally identifiable parameter then  $\chi_{\text{PL}}^2(\theta_k)$  exceeds the threshold  $\Delta_\alpha$  for both increasing and decreasing values of  $\theta_k$  forming a deep valley around  $\hat{\theta}_k$ . If  $\theta_k$  is structurally unidentifiable the profile likelihood is flat. Lastly, if  $\theta_k$  is practically unidentifiable,  $\chi_{\text{PL}}^2(\theta_k)$  obtains a unique minimum but does not exceed  $\Delta_\alpha$  in increasing and/or decreasing values of  $\theta_k$ , forming a shallow valley around  $\hat{\theta}_k$ .

We further investigate unidentifiable combinations of parameters by extending Equation 19 to profile parameter  $\theta_j$  and  $\theta_k$  simultaneously,

$$\chi_{\text{PL}}^2(\theta_j, \theta_k) := \min_{\theta_{i \notin \{j, k\}}} \chi^2(\boldsymbol{\theta}). \quad (20)$$

Structural relationships between the two profile parameters manifest as flat valleys extending infinitely along the functional relationship in the contour plots of  $\chi_{\text{PL}}^2(\theta_j, \theta_k)$ . We note this flat valley only traces out the functional relationship  $\theta_j$  and  $\theta_k$  when the dimension of the parameter space is larger than 2.

### 3 Results


3.1 Data do not support the Incremental Knockdown Hypothesis for a 20 second time frame

*3.1.1 No evidence of fast switching: ensemble effective diffusivities of the free subpopulation are the same regardless of exogenous Ab concentration*

For each Donor/Ab-concentration combination, the associated sample of virions contained a clear division among the tracked particles' MSD and ACF behavior. We used the classification scheme described in Section 2.3 to label each tracked virion as Immobilized, Freely-Diffusing, Subdiffusive or Outlier. The Immobilized class was characterized by low effective diffusivity ( $< 10^{-1} \mu\text{m}^2\text{s}^{-1}$ ) and either anti-persistent or uncorrelated increment processes. Meanwhile the Freely-Diffusing class had uncorrelated increment processes and effective diffusivities larger than  $0.2 \mu\text{m}^2/\text{s}$ . The Subdiffusive and Outlier classifications were rare and did not appear in all samples. For this reason, we removed these categories from the analysis but give a description of them in the SI. In Figure 4(a)-(c), we display the results of the classification for Donor F08 at 0, 0.1, and 1  $\mu\text{g}/\text{mL}$  added anti-HSV IgG in terms of  $D_{\text{eff}}$  and the average of the  $x$ - and  $y$ -ACF, as defined in Section 2.2.3 and Section 2.2.1 respectively. The clear separation of groups and locations of the clusters were qualitatively similar for the other donors (further figures included in the Supplemental Information).

The pathwise MSDs for Donor F08 virions are displayed in Figure 4(d)-(f), and we note the similarity of the Freely-Diffusing category of virions across all three panels. The Incremental Knockdown Hypothesis would predict that freely diffusing virions would be “slower and slower” in the presence of more and more Ab. However, we found that the diffusivities of the Freely-Diffusing classes are consistent across all exogenous Ab concentrations. In Figure 5 we display this fact in two ways. In the left panel, we display the ensemble MSD averaged over the Freely-Diffusing (green triangles) and Immobilized (red x's) populations for each Ab concentration. There is remarkable overlap within each group. Moreover, in the right panel, we display the ensemble effective diffusivity for the Freely-Diffusing class at the various exogenous Ab concentrations for all donors. While there is variation in the effective diffusivity, the overlapping  $BC_a$  confidence intervals indicate there is insufficient evidence to conclude the effective diffusivity decreases with antibody concentration. (We provide 95% weighted bootstrap confidence intervals for each estimate in the supplementary material Figure 13).

We can express this finding in terms of a statistical test by comparing the weighted ensemble effective diffusivity for the freely diffusing subpopulation at the two extreme Ab concentrations. We used a one-tailed paired difference



**Fig. 4 (a)-(c):** The unweighted composition of the tracked virions for Ab concentration 0, 0.1 and 1.0  $\mu\text{g/mL}$ , respectively for Donor F08. Each point corresponds to a tracked virion with the given estimated diffusivity on a log 10 scale and average-ACF value. The character of points denotes clusters prescribed by the hierarchical clustering algorithm and color of the point denotes the class of the cluster. **(d)-(f):** The path-wise MSD for all the tracked virions for Donor F08 at  $[A]_{\text{exo}} = 0, 0.1$ , and 1.00  $\mu\text{g/mL}$ . The colors, green, red, blue, denote the final clusters, freely diffusing, immobilized, and subdiffusive, respectively. Reference line with slope = 1, is denoted in black. (We note that the relative size of the different classes in this figure is not reweighted by path length as it is in the population counts reported in Figure 6.)


hypothesis test:

$$H_0 : \langle D_{\text{eff}}([A]_1) \rangle - \langle D_{\text{eff}}([A]_5) \rangle = 0, \quad H_A : \langle D_{\text{eff}}([A]_1) \rangle - \langle D_{\text{eff}}([A]_5) \rangle > 0. \quad (21)$$

for  $[A]_1 = 0.0 \mu\text{g/mL}$  and  $[A]_5 = 1.0 \mu\text{g/mL}$ . At an  $\alpha = 0.05$  level of significance, we failed to find significant evidence that the ensemble effective diffusivity of the freely diffusing population decreased when exogenous Ab concentration increased from zero exogenous Ab to the highest concentration ( $t_6 = 0.2567$ ,  $p\text{-value} = 0.4030$ ). We report the results of paired difference tests for all other combinations of the tested exogenous Ab concentration in the supplementary materials Table 10.

### 3.1.2 Little evidence of switching on the experimental time scale

We found little evidence that virions switch between states on the experimental time scale of 20 seconds. If tracked particles were typically experiencing many subtle switches, we expect that their computed effective diffusivities would be diminished by a factor determined by the time spent immobilized. Moreover,



**Fig. 5** (a) Ensemble MSD of the Freely Diffusing class and the immobilized class at various exogenous antibody concentrations represented by the green and red curves, respectively, for Donor F08. The black line refers to the ensemble MSD of Brownian particles, slope equal to 1. (b) The estimated ensemble effective diffusivity of the free population versus exogenous antibody concentration where the shade and point style of the curve corresponds Donor. See supplementary materials Figure13 for the ensemble effective diffusivity with 95%  $BC_a$  confidence intervals.

because there are distinct behavioral regimes, the distribution of the increment processes are essentially a mixture of two Gaussian distributions (one for the Immobilized state and one for the Freely Diffusing state). This would manifest itself as a violation of linearity in `qqnorm` plots, which we do not see for the vast majority of HSV virion paths.


While the `qqnorm` test can identify paths that might experience switches, they do not affirm the presence of a switch. To this end, we developed a Bayesian method for identifying whether there is a single switch point in a given virion path, described in the Section 2.5. We say a path of duration  $T_{\text{final}}$  is a candidate for switching if the 95% credible region for  $\tau$  was completely contained within the interval  $[0.1T_{\text{final}}, 0.9T_{\text{final}}]$ . The method was very effective on simulated data. When we applied the method to simulated Brownian motion (Freely-Diffusing), we found a 0.0119 and 0.0080 False Discovery Rate of [diffusion  $\rightarrow$  immobilization] switches and [immobilization  $\rightarrow$  diffusion] switches, respectively. On the other hand, 96.38% of the simulated [diffusion  $\rightarrow$  immobilization] paths were correctly identified as [diffusion  $\rightarrow$  immobilization] switches, while 94.37% of the simulated [immobilization  $\rightarrow$  diffusion] paths were identified as [immobilization  $\rightarrow$  diffusion] switches (Table 2). Under this method, we found that 1.12% of the Freely-Diffusing class (1689 total tracked virions) were identified as [diffusion  $\rightarrow$  immobilization] switch candidates and 1.24% of the free populations were [immobilization  $\rightarrow$  diffusion] switch candidates. We therefore concluded that state switches occurred relatively rarely on the experimental time scale.

**Table 2** Fraction of Freely Diffusing virions that possibly switched states once by Donor.

Model	Power	False Discovery Rate	Virion switch candidates
$D \rightarrow I$	0.9638	0.0119	0.0112
$I \rightarrow D$	0.9437	0.0080	0.0124

### 3.1.3 Fraction immobilized increases with exogenous antibody concentration

While Ab concentration did not seem to affect the behavior of virions labeled Freely-Diffusing, it did have a significant effect on the fraction of virions that were placed in this class. This is consistent with the findings reported in [20]. We computed the Immobilized fraction for each Donor/Ab-concentration sample using the method discussed in Section 2.3.1 and display the results in Figure 6, where each curve in the panel (b) corresponds to a different donor. While there is heterogeneity in the fraction of Immobilized virions across donors, there is a visible overall increase in proportion immobilized from 0 to 1  $\mu\text{g}/\text{mL}$ . This qualitative assessment is supported by statistical evidence provided by non-overlapping  $BC_a$  confidence intervals between the extreme exogenous Ab concentrations supplementary materials Figure 12.



**Fig. 6** (a) The weighted proportion of the 4 classes for Donor F08 at the various tested exogenous Ab concentrations. (b) Weighted proportions of Immobilized virions for each donor. See supplementary materials Figure 12 for plots with 95%  $BC_a$  confidence intervals.

For each donor, the fraction of Immobilized virions increased with Ab concentration in the 0 to 0.333  $\mu\text{g}/\text{mL}$  range and seemed to be saturated at higher Ab concentrations. We tested the significance of this observed trend by fitting a negative exponential growth model with predictors: exogenous antibody concentration and individual effect terms relative to Donor F08. Let  $\chi_k$  be the indicator function that a virion in the  $k$ th donor sample is in the immobilized

state. Our negative exponential growth model takes the form

$$P(\chi_k = 1) = (\beta_0 + \beta_k) - e^{-(\alpha_0 + \alpha_{\text{exo}}[A]_{\text{exo}}) + \alpha_k} \quad (22)$$


where  $\alpha_k$  and  $\beta_k$  are the effect terms for the  $k$ -th donor. We found the exogenous antibody concentration ( $\alpha_{\text{exo}} = 15.920$ ,  $p\text{-value} < 0.001$ ), the growth rate due to the baseline donor ( $\alpha_0 = -0.8427$ ,  $p\text{-value} = 0.0043$ ), and baseline saturation probability ( $\beta_0 = 0.9138$ ,  $p\text{-value} < 0.0001$ ) were statistically significant in predicting the immobilization probability, whereas the constants accounting for deviations from the baseline due to donor sample were not significant. The model was fit using the R command `nls()` with the minimization algorithm set to Gauss-Newton's method.

### 3.2 The Simple Linear Model predicts fast switching

The results from Section 3.1 provide evidence against the hypothesis that switching between the diffusing and immobilized states is fast relative to the experimental time scale. Our next goal was to determine whether there is a parameter regime that predicts slow switching while simultaneously being consistent with the exogenous Ab-dependent Immobilization data displayed in Figure 6. This analysis depends strongly on two assumptions: (1) whether one virion-bound Ab is sufficient to crosslink the virion to mucin, and (2) whether Ab-mucin binding rates increase when the virion is immobilized, the so-called cascade effect. We introduced two variables –  $T$ , the threshold number, and  $c$ , the cascade factor – in our general model to account for these possible effects. In recent works, it has been assumed either that  $T = c = 1$  [2,11] or that  $T = 1$  and  $c > 1$  [22]. We refer to  $T = c = 1$  as the Simple Linear Model (SLM) because all the CTMC transition rates are linear. By computing the expected durations of the immobilized and diffusing states (Equation 13, derivation in Appendix B.2), we were able to show that the data is not consistent with the SLM, or any case where  $T = 1$ .

We say a model is *consistent* with the observed data for a specified donor if there exists a parameter vector  $\boldsymbol{\theta}$  that is within the 95% confidence region for the Immobilized Fraction data (denoted  $\Theta_{\alpha,df}$ , defined in Equation 18) and also predicts expected state times larger than 20 seconds (denoted  $\Theta_{\text{slow}}$ , defined in Equation 14). In Figure 7, we demonstrate that the SLM is not consistent with the data for Donor F08. In the left panel, we show a 2d profile likelihood plot for the endogenous Ab concentration  $[A]_0$  and number of virion surface binding sites  $N_*$ . For each  $([A]_0, N_*)$  pair, we calculated the best fit for the remaining parameter  $q$ , the virion-Ab interaction probability, and display the residual value by the shading (darker means better fits). The black region represents the 95% confidence region for these two parameters. We uniformly sampled this confidence region,  $\Theta_{0.05,3}$ , and displayed the predicted Immobilized Fraction curves for these parameter samples in panel (b) and the Ab-concentration dependent expected state durations in panel (c). We note that all parameter combinations in  $\Theta_{0.05,3}$  had diffusing states that

lasted less than 0.1 seconds for all values of  $[A]_{\text{exo}}$ . We repeated this analysis for all donors and in each case found that  $\Theta_{0.05,3} \cap \Theta_{\text{slow}} = \emptyset$ .




**Fig. 7** (a),(d) Profile likelihood contour plots (Donor F08) for  $\chi^2_{\text{PL}}([A]_0, N_*)$  and  $\chi^2_{\text{PL}}(c, N_*)$  when  $T = c = 1$  and  $T = 19$ , respectively. Darker shades correspond to smaller profile likelihood values and the black region corresponds to the 95% confidence regions  $\Theta_{0.05,3}$  and  $\Theta_{0.05,5}$ . (b),(e) Predicted Immobilized Fraction curves (gray lines) for  $\theta$  sampled from  $\Theta_{0.05,3}$  and  $\Theta_{0.05,5}$ . The black curve is the prediction of the best fit in each case for Donor F08. The observed Immobilized Fraction is shown by the purple line with triangles. (c),(f) Expected duration of Immobilized (red curves) and Freely-Diffusing (green curves) states for  $\theta$  sampled from  $\Theta_{0.05,3}$  and  $\Theta_{0.05,5}$ . When  $T = c = 1$ , frame (c), none of predicted state times are above 20 seconds, horizontal black line. On the other hand, when  $T = 19$ , frame (f), there are some parameter combinations that do yield slow switching. These are marked in light blue as appropriate in Panels (d)-(f).

### 3.3 Threshold and binding cascade parameters allow slow switching

By allowing the immobilization process to require multiple cross-linking antibodies,  $T > 1$ , and for the Ab-mucin dynamics to be state-dependent,  $c \neq 1$ , we found both that (1) the subset of parameters that lead to slow switching is non-empty ( $\Theta_{\text{slow}} \neq \emptyset$ ), and (2) there is an overlap between slow-switching parameters and parameters that fit the Immobilized Fraction data well ( $\Theta_{0.05,5} \cap \Theta_{\text{slow}} \neq \emptyset$ ). For example, in Figure 7 panels (d)-(f) we demonstrate this fact assuming  $T = 19$  for Donor F08. The 2d profile likelihood plot in panel (d) shows an inverse relationship between  $N_*$  and the cascade factor  $c$ . Again the black region corresponds to all  $(c, N_*)$  pairs that appear in  $\Theta_{0.05,5}$ .

For a uniform sample of such pairs, in panel (e) we display the Immobilized Fraction predictions, and in panel (f) the corresponding expected immobilization and diffusion state durations. Only a small subset of  $\Theta_{0.05,5}$  allows for slow switches. We mark this subset in blue in all three panels. Notably, conditioned on  $T = 19$ , we have that  $N_* \leq 120$ , which is somewhat smaller than the typical estimate for  $N_*$ . In the next section we note that assuming higher values for  $T$  leads to higher allowable values for  $N_*$ . This type of result holds for all donors: for sufficiently high assumed  $T$ , the corresponding parameters sets  $\Theta_{0.05,5}$  and  $\Theta_{\text{slow}}$  overlap.



**Fig. 8** (a) Parameter combinations of  $T$  and  $c$  that predict expected immobilized times greater than 20s, red points, and predict expected freely-diffusing times greater than 20s, green points, assuming  $N_* = 300$  and  $[A]_0 = 0.1\mu\text{g/mL}$ . The overlapping  $(T, c)$  combinations (blue points) are those combinations that satisfy slow switching condition and subset  $(T, c_{\min})$  are denoted by black. (b) The minimal value of  $T$  required for our model to predict slow switching as a function of  $N_*$ , orange curve. Given an  $N_*$  and corresponding minimal  $T$  pair, the minimal value of  $c$  required for our model to predict slow switching is denoted by the brown curve. The endogenous Ab concentration is fixed at  $[A]_0 = 0.1\mu\text{g/mL}$ .

By testing over a range of  $\boldsymbol{\theta} = (T, c, N_*, [A]_0)$ , we uncovered some relationships among the components of the parameter vectors  $\boldsymbol{\theta}$  that yield slow switching  $\Theta_{\text{slow}}$ . We first investigated the relationship between  $T$  and  $c$  by fixing  $N_*$  and  $[A]_0$ . Noting that  $\mathbb{E}_{\theta}(\tau)$  is independent of  $c$  and  $\mathbb{E}_{\theta}(\sigma)$  is an increasing function in  $c$ , we calculated the minimal  $c$  required to satisfy the slow switching condition, labeling this value  $c_{\min}$ . Though we could not obtain an explicit relationship between  $T$  and  $c_{\min}$ , we found that virions with a large immobilization threshold  $T$  can only satisfy the slow switching condition if there is a corresponding large cascade effect, large  $c_{\min}$ . To visualize this, in Figure 8(a) we display the parameter combinations of  $(T, c, N_* = 300, [A]_0 = 0.1)$  that yield  $\mathbb{E}_{\theta}(\tau) > 20$  (green) and  $\mathbb{E}_{\theta}(\sigma) > 20$  (red) for all exogenous antibody concentrations between 0 and  $1\mu\text{g/mL}$ . The overlapping region (blue points) corresponds to  $\theta \in \Theta_{\text{slow}}$  and the combinations of interest  $(T, c_{\min})$  are shown in black.

We draw the conclusion that if  $N_* = 300$ , then  $T$  must be at least 34 and  $c$  must be at least 63. If we increase the assumption about  $N_*$  while keeping  $[A]_0$  fixed, then we found that the minimal allowable  $T$  and  $c$  for slow switching increase and decrease, respectively. We demonstrate this relationship in Figure 8(b). For  $[A]_0 = 0.1\mu\text{g/mL}$ , the orange (circles) curve corresponds to the minimal  $T$  value (left  $y$ -axis) for the given  $N_*$  ( $x$ -axis) required such that  $\theta \in \Theta_{\text{slow}}$  where  $[A]_0 = 0.1\mu\text{g/mL}$ . The brown (triangles) curve denotes the minimal  $c$  value (right  $y$  axis) required for the given  $N_*$ , minimal  $T$ , and  $[A]_0 = 0.1\mu\text{g/mL}$  to result in expected state times longer than 20 seconds.

### 3.4 Model with threshold and binding cascade parameter is unidentifiable


As implied by the results in the preceding section, we found that the introduction of  $T > 1$  and  $c \neq 1$  resulted in issues with identifiability. That is to say, it appears that the confidence region  $\Theta_{0.05,5}$  is infinite even when restricted to the subspace  $\Theta_{0.05,5} \cap \Theta_{\text{slow}}$ . We use the Immobilized Fraction data for Donor F08 to demonstrate this fact but provide information for each Donor in the Supplementary Information. Throughout this section we will use the terminology defined in Section 2.6.

Over the full parameter space  $\Theta$ , the 1d profile likelihoods revealed that all three of the parameters  $T$ ,  $c$ , and  $N_*$  are practically unidentifiable over the range we tested. The profile likelihoods are displayed in black in Figure 9(a)-(c). When we profiled the parameters  $T$ ,  $c$ , and  $N_*$  restricted to the Slow Switching Regime  $\Theta_{0.05,5} \cap \Theta_{\text{slow}}$ , we found  $T$  is still practically unidentifiable over the range  $T \geq 19$ , while  $c$  is practically unidentifiable a large range of positive values. The number of binding sites  $N_*$  does seem to be identifiable, with a deep valley centered around the unique minimum at approximately  $N_* = 120$ . These profile likelihoods are represented in blue in Figure 9(a)-(c). The dashed lines correspond to the 95% confidence interval boundaries for each parameter. Since the blue curves are below the confidence interval we can say that there exist parameter combinations in the Slow Switching Regime that reasonably fit the Immobilized Fraction data in Figure 9(e).

## 4 Discussion

We have developed mathematical models and statistical methods to analyze the behavior of HSV virions diffusing in CVM in the presence of various concentrations of cross-linking Ab. With a few exceptions, we found that particle paths can be partitioned into two basic categories: Freely Diffusing and Immobilized. While the fraction of Immobilized virions increases with Ab concentration, we found that the mobility of the Freely Diffusing class is not Ab-concentration dependent.

Because we expect all the individual bonds to be reversible, virions should switch between the Freely Diffusing and Immobilized states. Previously, it had



**Fig. 9** (a)-(c) The 1d profile likelihoods for the parameters: immobilization threshold  $T$ , cascade factor  $c$ , and number of Ab binding sites on the virion  $N_*$ , respectively over all tested parameter combinations (black curves) and when restricted to the slow switching regime (blue curves). The 95% confidence interval for each parameter consists of those parameter values with profile likelihood values below the dashed line.

been hypothesized that such switches are rapid with respect to the experimental time scale, but our analysis contradicts that assumption. This raises the question of whether or not it is possible for the basic kinetic model to produce “slow-switching” paths where switches occur on a time scale much larger than the experimental time window. We found that this is possible if the model allows for a lower bound on the number of Ab necessary to immobilize a virion and assuming a “cascade effect” in Ab-mucin binding that encourages entanglement.

Introducing these extra features leads to a fundamental issue with unidentifiability in the statistical analysis. We can make claims like “the minimum number of antibodies needed to immobilize a virion must be greater than 20 or so”, but we cannot be more specific. In order to do so, we would need to have access to time series that are much longer than what is currently experimentally feasible.

While we have shown that it is possible for reversible kinetics to be consistent with the path data, it might also be possible to explain the data with a model that assume all binding events are irreversible. Unfortunately the available data cannot distinguish between the two models. One possible resolution is to conduct experiments that explicitly control for the time between the introduction of Ab to the virion population and the observation of virion trajectories. Based on our model, in which we assume the immobilization process is reversible prior to the system reaching stationarity, switching should be more common when the number of antibodies bound to surface epitopes is low. Therefore, starting the tracking immediately enhances the probability of observing state switches before any long-lasting immobilization events occur.

On the other hand, observing virions at different time points long after Ab introduction will help determine whether or not the system reaches a stationary distribution. If so, there should be substantial information in analyzing how (or if) that stationary distribution depends on the Ab concentration, and the rate at which that stationary distribution is achieved.

From a biological point of view, this uncertainty about the true time scale of state switching prevents conclusive answers for the type of model presented by Chen et al. []. Namely, in order to characterize the percentage of virions that can pass through a mucosal layer within two hours, we must know whether state switching takes place on the order of minutes, days or longer. The analysis in this work simply served to eliminate the possibility of switching on a time scale of milliseconds or seconds, which was the standing assumption.

The statistical methods and mathematical model introduced here apply to a broad class of biological systems that are composed of distinct subpopulations. Our classification scheme based on path-by-path analysis detects subpopulation dynamics that can be masked when considering only overall ensemble behavior. Clustering and then analyzing subpopulation ensemble statistics provides insight on the way the proportion and dynamics of these subpopulation change in response to the environmental factors. The model proposed in Section 2.4, can be modified to describe the general scenario when nanoparticles work to entrap a diffusing pathogen by anchoring the pathogen to the surrounding environment.

## A Derivation of the MLE for D

From the defining properties of Brownian motion, the likelihood function of 2d Brownian motion defined by  $d\mathbf{X}(t) = \sqrt{2D}d\mathbf{W}(t)$  has form

$$L(D; u, v) = \left( \frac{1}{4\pi\delta D} \right)^n \exp \left( - \sum_{k=1}^n \frac{(X(k\delta) - X((k-1)\delta))^2}{4D\delta} \right) \times \exp \left( - \sum_{k=1}^n \frac{(Y(k\delta) - Y((k-1)\delta))^2}{4D\delta} \right). \quad (23)$$

In terms of the increment process,  $U(k\delta)$  and  $V(k\delta)$ , the loglikelihood is

$$\ell(D) = -n(\log(4\pi\delta) + \log(D)) - \frac{1}{4D\delta} \sum_{k=1}^n (U(k\delta)^2 + V(k\delta)^2). \quad (24)$$

Solving the likelihood equation  $\frac{d}{dD}\ell(D) = 0$ , the ML estimator for  $D$  is given by

$$\hat{D}_{\text{MLE}} = \frac{1}{4\delta n} \sum_{k=1}^n (U(k\delta)^2 + V(k\delta)^2). \quad (25)$$

## B Mathematical Model in Section 2.4

We arrive at our approximation to the probability of immobilization Equation 11 presented in Section 2.4 by averaging over the transitions of the number of antibodies simultaneously bound to the virion,  $S(t)$ . To do this, we consider a simplified model in which the Ab-mucin binding rate is the same for both an immobilized virion and freely diffusing virion. That is

the function defined in Equation 10 is constant,  $g(s) \equiv c$ . In this case, let  $S(t)|_{n,c}$  denote the Markov chain with transition rates:

$$s \xrightarrow[s m_{\text{off}}]{(n-s)cm_{\text{on}}[M]} s+1. \quad (26)$$

First, we derive the stationary distribution for the number of bound antibodies,  $N(t)$ , and the conditional number of simultaneously bound antibodies assuming  $g(s) \equiv c$ ,  $S(t)|_{n,c}$ . Then we compute the expected duration of the Immobilized state and the Freely-Diffusing state of a virion from those quantities assuming  $g(s) \equiv c$ . Finally, we obtain Equation 11 using the results of the previous two steps.

### B.1 Stationary distribution of the two processes $N(t)$ and $S(t)|_{n,c}$

We model the two processes  $N(t)$  and  $S(t)|_{n,c}$  as CTMC with transition rates given by Equation 8 and Equation 26, respectively. Because they are irreducible Markov chains with a finite state space, there exists a unique stationary distribution, and convergence is exponential. Under the assumption that the Ab-binding sites on the surface of a virion operate independently, the process  $N(t)$  follows a binomial distribution with  $N_*$  trials and a time dependent success probability. Evoking a classical result from Renewal Theory, the steady state success probability is given by the long run fraction of being in the bound state, so that

$$\lim_{t \rightarrow \infty} N(t) \sim \text{Binom}\left(\frac{k_{\text{on}}A}{k_{\text{off}} + k_{\text{on}}A}, N_*\right). \quad (27)$$

By assuming  $g(s) \equiv c$ , each antibody bound to the virion interacts with the mucin fibers independently, so that  $S(t)|_{n,c}$  is a binomial random variable with  $n$  trials and time dependent success probability. It follows from the same reasoning as above, that

$$\lim_{t \rightarrow \infty} S(t)|_{n,c} \sim \text{Binom}\left(\frac{cm_{\text{on}}M}{m_{\text{off}} + cm_{\text{on}}M}, n\right) \quad (28)$$

is the unique stationary distribution.

### B.2 Expected duration of the Freely-Diffusing and Immobilized states

We derive the expected duration of the Freely-Diffusing state,  $\tau$ , and Immobilized states,  $\sigma$ , of a virion by considering the simplified model when the number of simultaneous bound antibodies has transition rates Equation 26.

We introduce the following notation  $\tau_{T;c}$  and  $\sigma_{T;c}$  denote the expected time the process  $S_{n;c}$  spends in the freely diffusing state and immobilized state, respectively. The expected duration of the Freely-Diffusing state,  $S_{n;c}(t) < T$ , is simply the expected hitting time of state  $T$ , given  $S_{n;c}$  starts with  $T-1$  simultaneously bound antibodies. By solving a system of linear equations for the vector of expecting hitting times of state  $T$ , see [12], yields

$$\mathbb{E}(S_{n;c}(t) = T | S_{n;c}(0) = T-1) = \frac{1}{(n-T+1)m_{\text{on}}[M]b\left(T-1; n, \frac{m_{\text{on}}M}{m_{\text{off}} + m_{\text{on}}[M]}\right)} \sum_{s=0}^{T-1} b(s; n, \frac{m_{\text{on}}[M]}{m_{\text{off}} + m_{\text{on}}[M]}). \quad (29)$$

Similarly, the expected duration of the Immobilized state,  $S_{n;c}(t) \geq T$ , is the expected hitting time of state  $T-1$  given  $S_{n;c}$  starts in state  $T$ . By solving a system of linear equations for the vector of expecting hitting times of state  $T-1$ ,

$$\mathbb{E}(S_{n;c}(t) = T-1 | S_{n;c}(0) = T) = \frac{1}{Tm_{\text{off}}b\left(T; n, \frac{m_{\text{on}}[M]}{m_{\text{off}} + m_{\text{on}}[M]}\right)} \sum_{s=T}^n b(s; n, \frac{m_{\text{on}}[M]}{m_{\text{off}} + m_{\text{on}}[M]}). \quad (30)$$

We observe that the transition rates of a Freely Diffusing virion are the same as a virion modeled by the simplified transition rates given by Equation 26, specifically for  $c = 1$ . The transition rates of an immobilized virion are the same as a virion modeled by the simplified transition rates Equation 26. Hence, the following equalities hold

$$\mathbb{E}(\sigma; T, c, n) = \mathbb{E}(\sigma_{T;c}; T, c, n) \quad \text{and} \quad \mathbb{E}(\tau; T, c, n) = \mathbb{E}(\tau_{T;1}; T, 1, n).$$

Explicitly, the duration of the Freely-diffusing state and the Immobilized state of a virion are given by

$$\mathbb{E}(\tau_{T;c}; T, c, n) = \frac{1}{(n-T+1)m_{\text{on}}M} \left( \frac{\sum_{s=0}^{T-1} \binom{n}{s} \left( \frac{m_{\text{on}}M}{m_{\text{off}}+m_{\text{on}}M} \right)^s \left( \frac{m_{\text{off}}}{m_{\text{off}}+m_{\text{on}}M} \right)^{n-s}}{\binom{n}{T-1} \left( \frac{m_{\text{on}}M}{m_{\text{off}}+m_{\text{on}}M} \right)^{T-1} \left( \frac{m_{\text{off}}}{m_{\text{off}}+m_{\text{on}}M} \right)^{n-T+1}} \right) \quad (31)$$

$$\mathbb{E}(\sigma_{T;c}; T, c, n) = \frac{1}{Tm_{\text{off}}} \left( \frac{\sum_{s=T}^n \binom{n}{s} \left( \frac{cm_{\text{on}}M}{m_{\text{off}}+cm_{\text{on}}M} \right)^s \left( \frac{m_{\text{off}}}{m_{\text{off}}+cm_{\text{on}}M} \right)^{n-s}}{\binom{n}{T} \left( \frac{cm_{\text{on}}M}{m_{\text{off}}+cm_{\text{on}}M} \right)^T \left( \frac{m_{\text{off}}}{m_{\text{off}}+cm_{\text{on}}M} \right)^{n-T}} \right), \quad (32)$$

respectively.

### B.3 Asymptotic probability of number of bound antibodies

We are now ready to derive our time-scale approximation of the asymptotic probability of immobilization function, Equation 11. By conditioning on the slow process, the antibody-virion dynamics,

$$\begin{aligned} \hat{\pi}(A; \theta) &= \lim_{t \rightarrow \infty} \mathbb{P}\{q > 0 \cap \{S(t) \geq T\}\} \\ &= \mathbb{P}\{q > 0\} \lim_{t \rightarrow \infty} \mathbb{P}\{S(t) \geq T | q > 0\} \\ &= q \sum_{n=T}^{N_*} \left( \lim_{t \rightarrow \infty} \mathbb{P}\{S(t) \geq T | N(t) = n, q > 0\} \right) \cdot \left( \lim_{t \rightarrow \infty} \mathbb{P}\{N(t) = n\} \right). \end{aligned} \quad (33)$$

An application in Renewal Theory leads to the stationary probability of immobilization for the conditional process  $S(t)$  of the form

$$\lim_{t \rightarrow \infty} \mathbb{P}\{S(t) \geq T | N(t) = n, q > 0\} = \frac{\mathbb{E}(\sigma; T, c, n)}{\mathbb{E}(\sigma; T, c, n) + \mathbb{E}(\tau; T, n)}. \quad (34)$$

Plugging in the results from Appendix B.1 and B.2, gives Equation 11.

## C Derivation of likelihoods

We derive the likelihood of Equation 15 and Equation 16 from the exact solution of the SDEs under the assumptions that the switch point,  $\tau$ , occurs at a time measurement, and the true 2d position of the particle is  $\mathbf{X}(t) = (X(t), Y(t))$ . We denote the time measurement  $t_k = k\delta$  for  $k = 1, \dots, n$ , where  $t_0 = 0$  and  $t_n = T$ , and  $\mathbf{X}(t_k) = \mathbf{X}_k$ .

For the [diffusion  $\rightarrow$  immobilization] model, when  $t > \tau$  the SDE is linear with additive noise. Hence a conditional exact solution can be expressed using Duhamel's formula,

$$\mathbf{X}_k | \mathbf{x}_{k-1} = \begin{cases} \mathbf{x}_{k-1} + \sqrt{2D}(\mathbf{W}_k - \mathbf{W}_{k-1}) & 0 < t_k \leq \tau \\ \mathbf{x}_{k-1} e^{-\tilde{\kappa}\Delta} + (1 - e^{-\tilde{\kappa}\delta})\mathbf{x}_\tau + \sqrt{2D} \int_{t_{k-1}}^{t_k} e^{-\tilde{\kappa}(t_k-s)} d\mathbf{W}(s) & \tau < t_k \leq T. \end{cases} \quad (35)$$

It follows immediately from the definition of Brownian motion and from an application of Ito's Isometry,

$$\mathbf{X}_n |_{\mathbf{x}_{n-1}} \sim \begin{cases} N(\mathbf{x}_{k-1}, 2\delta D) & 0 < t_k \leq \tau \\ N(\rho \mathbf{x}_{k-1} + (1-\rho)\mathbf{x}_\tau, \frac{D}{\tilde{\kappa}}(1-\rho^2)) & \tau < t_k \leq T. \end{cases} \quad (36)$$

where  $\rho = e^{-\tilde{\kappa}\delta}$ . Because the solutions to SDEs satisfy the Markov Property,

$$\begin{aligned} L((\mathbf{x}_1, \dots, \mathbf{x}_n)) &= \left( \prod_{k=1}^{\tau} \mathbb{P}_{0, \mathbf{x}_\tau}(\mathbf{X}_k |_{\mathbf{x}_{k-1}} = \mathbf{x}_k) \right) \left( \prod_{k=\tau+1}^n \mathbb{P}_{0, \mathbf{x}_\tau}(\mathbf{X}_k |_{\mathbf{x}_{k-1}} = \mathbf{x}_k) \right) \\ &= \left( \frac{1}{4\pi\delta D} \right)^\tau \left( \frac{\tilde{\kappa}}{2\pi D(1-\rho^2)} \right)^{n-\tau} \exp \left( \frac{-1}{4\delta D} \sum_{k=1}^{\tau} ((X_k - X_{k-1})^2 + (Y_k - Y_{k-1})^2) \right) \\ &\times \exp \left( \frac{-\tilde{\kappa}}{2D(1-\rho^2)} \sum_{k=\tau+1}^n ((X_k - \rho X_{k-1} - (1-\rho)x_\tau)^2 + (Y_k - \rho Y_{k-1} - (1-\rho)y_\tau)^2) \right). \end{aligned} \quad (37)$$

The likelihood equation for the [immobilization  $\rightarrow$  diffusion] switching model derivation is similar to that [diffusion  $\rightarrow$  immobilization] switching model but now we assume the immobilized particle is centered around the origin. Under the same reasoning as above

$$\begin{aligned} L((\mathbf{x}_1, \dots, \mathbf{x}_n)) &= \left( \prod_{i=1}^{\tau} \mathbb{P}_{0, \mathbf{x}_\tau}(\mathbf{X}_k |_{\mathbf{x}_{k-1}} = \mathbf{x}_k) \right) \left( \prod_{i=\tau+1}^n \mathbb{P}_{0, \mathbf{x}_\tau}(\mathbf{X}_k |_{\mathbf{x}_{k-1}} = \mathbf{x}_k) \right) \\ &= \left( \frac{\tilde{\kappa}}{2\pi D(1-\rho^2)} \right)^\tau \exp \left( \frac{-\tilde{\kappa}}{2D(1-\rho^2)} \sum_{n=1}^{\tau} ((X_k - \rho X_{k-1})^2 + (Y_k - \rho Y_{k-1})^2) \right) \\ &\times \left( \frac{1}{4\pi\delta D} \right)^{n-\tau} \exp \left( \frac{-1}{4\delta D} \sum_{k=\tau+1}^n ((X_k - X_{k-1})^2 + (Y_k - Y_{k-1})^2) \right). \end{aligned} \quad (38)$$

**Acknowledgements** The authors would like to thank Jay Newby, John Fricks, and Michelle Lacey for the helpful conversations that contributed to the development of this work.


## References

1. Bernstein, J., Fricks, J.: Analysis of single particle diffusion with transient binding using particle filtering. *Journal of Theoretical Biology* **401**, 109–121 (2016)
2. Chen, A., McKinley, S.A., Wang, S., Shi, F., Mucha, P.J., Forest, M.G., Lai, S.K.: Transient antibody-mucin interactions produce a dynamic molecular shield against viral invasion. *Biophysical Journal* **106**(9), 2028–2036 (2014)
3. D'Orsogna, M.R., Chou, T.: Optimal cytoplasmic transport in viral infections. *PLoS One* **4**(12), e8165 (2009)
4. Efron, B., Tibshirani, R.J.: An introduction to the bootstrap. CRC press (1994)
5. Eisenberg, M.C., Hayashi, M.A.: Determining identifiable parameter combinations using subset profiling. *Mathematical Biosciences* **256**, 116–126 (2014)
6. Goychuk, I., Kharchenko, V.O., Metzler, R.: How molecular motors work in the crowded environment of living cells: coexistence and efficiency of normal and anomalous transport. *PLoS One* **9**(3), e91700 (2014)
7. Holcman, D., Schuss, Z.: Time scale of diffusion in molecular and cellular biology. *Journal of Physics A: Mathematical and Theoretical* **47**(17), 173001 (2014)
8. Kaufman, L., Rousseeuw, P.J.: Finding groups in data: an introduction to cluster analysis, vol. 344. John Wiley & Sons (2009)

9. Matsuda, H., Putzel, G.G., Backman, V., Szleifer, I.: Macromolecular crowding as a regulator of gene transcription. *Biophysical Journal* **106**(8), 1801–1810 (2014)
10. McKinley, S.A., Chen, A., Shi, F., Wang, S., Mucha, P.J., Forest, M.G., Lai, S.K.: Modeling neutralization kinetics of hiv by broadly neutralizing monoclonal antibodies in genital secretions coating the cervicovaginal mucosa. *PloS One* **9**(6), e100598 (2014)
11. Newby, J., Schiller, J.L., Wessler, T., Edelstein, J., Forest, M.G., Lai, S.K.: A blueprint for robust crosslinking of mobile species in biogels with weakly adhesive molecular anchors. *Nature Communications* **8**(1), 833 (2017)
12. Norris, J.R.: Markov chains, vol. 2. Cambridge university press (1998)
13. Olmsted, S.S., Padgett, J.L., Yudin, A.I., Whaley, K.J., Moench, T.R., Cone, R.A.: Diffusion of macromolecules and virus-like particles in human cervical mucus. *Biophysical Journal* **81**(4), 1930–1937 (2001)
14. Press, W.H., Teukolsky, S.A., Vetterling, W.T., Flannery, B.P.: Numerical recipes in Fortran 90, vol. 2. Cambridge university press Cambridge (1996)
15. Qian, H., Sheetz, M.P., Elson, E.L.: Single particle tracking: analysis of diffusion and flow in two-dimensional systems. *Biophysical Journal* **60**(4), 910–921 (1991)
16. Raue, A., Kreutz, C., Maiwald, T., Bachmann, J., Schilling, M., Klingmüller, U., Timmer, J.: Structural and practical identifiability analysis of partially observed dynamical models by exploiting the profile likelihood. *Bioinformatics* **25**(15), 1923–1929 (2009)
17. Schroeder, H.A., Nunn, K.L., Schaefer, A., Henry, C.E., Lam, F., Pauly, M.H., Whaley, K.J., Zeitlin, L., Humphrys, M.S., Ravel, J., et al.: Herpes simplex virus-binding IgG traps HSV in human cervicovaginal mucus across the menstrual cycle and diverse vaginal microbial composition. *Mucosal Immunology* **11**(5), 1477 (2018)
18. Venables, W.N., Ripley, B.D.: Modern applied statistics with S-PLUS. Springer Science & Business Media (2013)
19. Vestergaard, C.L., Blainey, P.C., Flyvbjerg, H.: Optimal estimation of diffusion coefficients from single-particle trajectories. *Physical Review E* **89**(2), 022726 (2014)
20. Wang, Y.Y., Kannan, A., Nunn, K.L., Murphy, M.A., Subramani, D.B., Moench, T., Cone, R., Lai, S.K.: IgG in cervicovaginal mucus traps HSV and prevents vaginal herpes infections. *Mucosal Immunology* **7**(5), 1036–1044 (2014)
21. Wang, Y.Y., Nunn, K.L., Harit, D., McKinley, S.A., Lai, S.K.: Minimizing biases associated with tracking analysis of submicron particles in heterogeneous biological fluids. *Journal of Controlled Release* **220**, 37–43 (2015)
22. Wessler, T., Chen, A., McKinley, S.A., Cone, R., Forest, M.G., Lai, S.K.: Using computational modeling to optimize the design of antibodies that trap viruses in mucus. *ACS Infectious Diseases* **2**(1), 82–92 (2015)
23. Witten, J., Ribbeck, K.: The particle in the spider's web: transport through biological hydrogels. *Nanoscale* **9**(24), 8080–8095 (2017)
24. Yang, B., Schaefer, A., Wang, Y.Y., McCallen, J., Lee, P., Newby, J.M., Arora, H., Kumar, P.A., Zeitlin, L., Whaley, K.J., et al.: Zmapp reinforces the airway mucosal barrier against ebola virus. *The Journal of infectious diseases* p. jiy230 (2018)

## 1 Typical virion trajectories

We display the typical trajectories of the four different types of classified particles: Freely Diffusing, Immobilized, Subdiffusive, and Outlier in Figure 1. The Freely Diffusing class made up 45.5246% of the paths (1689/3707), Immobilized class made up 53.0618% of the paths (1967/3707), Subdiffusive class made up 0.3507% of the paths (13/3707), and the outlier class made up 1.0251% of the paths (38/3707),




**Fig. 1** (a)-(f): A typical 2d-trajectory of a Virion classified as Freely Diffusing, Immobilized, Subdiffusive, Freely-Diffusing switch candidate, Outlier (Type I) and Outlier (Type 2), respectively.

## 2 Localization error and effective diffusivity

We used the method provided by Vestergaard et al. [1] (Section III.C) to estimate the variance of the localization error and the diffusivity of a freely diffusing particles for all tracked virions, under the assumption of maximal exposure time (motion blur coefficient = 1/6). In some cases where the virion has a correlated first-increment process (not freely diffusing), we obtained negative estimates for both the variance of the localization error and the diffusivity using the covariance based estimator. This was more common in the Immobilized class and the Freely Diffusing class had strictly positive diffusivity estimates using both methods.

The empirical distribution of the localization errors are provided in Figure 2(a) for the Immobilized and Freely Diffusing class in red and green, respectively. For those Freely Diffusing virions with positive localization error estimates, the estimate for the diffusivity using the covariance based estimator estimator was typical larger than the estimate for localization error by two orders of magnitude, as depicted in Figure 2(b). In Figure 2(c), we compare the estimate for the diffusivity using the maximum likelihood estimator ( $x$ -axis) and the estimator in [1], covariance based estimator, ( $y$ -axis).



**Fig. 2** (a): The empirical density of the variance of the localization error for the Immobilized (red) and Freely Diffusing (green) sub-population. (b): The diffusivity covariance based estimator (CVE) versus the estimate for the variance of the localization error for the Freely Diffusing class with positive estimates for the variance of the localization error on a log 10 vs. log 10 scale. (c): The covariance based estimator (CVE) versus the maximum likelihood estimator (MLE) for the diffusivity for the Immobilized (red) and Freely Diffusing (green) class. The dashed black line denotes the line  $x = y$ .

### 3 Simulations of the immobilization process

We simulate the immobilization process, depicted in Figure 3 using a Gillespie algorithm. First, we simulate the number of bound Ab to the surface of a virion,  $N(t)$ . To do this, we use the transition rates given in Equation 8 to randomly select if a bound Ab is gained or lost and randomly sample the time the virion remains in the current state.

Then for the given value of  $N(t) = n$ , we simulate the number of simultaneously bound Ab,  $S(t)$  for the duration of time  $N(t)$  remains in given state. Because we assume the Ab-mucin dynamics are fast compared to Ab-virion dynamics, we approximate  $S(t)$  by a continuous space process using the stochastic differential equation

$$d\tilde{S}(t) = (\lambda(\tilde{S}(t)) - \mu(\tilde{S}(t)))dt + \sqrt{(\lambda(\tilde{S}(t)) + \mu(\tilde{S}(t)))}dW(t) \quad (1)$$

where  $\lambda(\tilde{S}(t)) = (N(t) - \tilde{S}(t))g(\tilde{S}(t))m_{\text{on}}[M]$  (the rate of gaining a simultaneously bound Ab) and  $\mu(\tilde{S}(t)) = \tilde{S}(t)m_{\text{off}}$  (the rate of losing a simultaneously bound Ab). Then we use the Euler–Maruyama method to simulate the process  $\tilde{S}(t)$  for the duration of time the number of bound Ab remains fixed.

### 4 Clustering Algorithm Decision

For each donor we give a table of the classification of each cluster at each tested exogenous antibody concentration and graphically display the grouping of clusters based on the path-wise statistics. We provide a description of each of the four subpopulations in Table 1.

**Table 1** Ensemble statistics for each of the four biological subpopulations.


	Diffusing	Immobilized		Subdiffusive	Outlier
$\langle D_{\text{eff}} \rangle$	$\geq 10^{-1}$	$< 10^{-1}$		$\geq 10^{-1}$	-
$\mathcal{A}(1; X), \mathcal{A}(1; Y)$	Independent	Independent	Anti-persistent	Anti-persistent	-
Motion	Brownian	Hindered Brownian	Stationary	Subdiffusive	-
Color	Green	Red		Blue	Brown

#### 4.1 Donor F02

The dendrogram for Donor F02 was cut at a uniform height to yield four cluster for each tested exogenous Ab concentration except for  $[A]_{\text{exo}} = 1.0 \mu\text{g/mL}$ , which was cut at a uniform height to yield five clusters because one particle was separated into its own cluster. The biological label that we assign to each cluster is reported in Table 2. We display the clusters based on the path-wise statistics in Figure 3, where the number corresponds to the hierarchical cluster and the color corresponds to the biological interpretation of the cluster.

**Table 2** The classification of the clusters separated by the hierarchical clustering algorithm described in Section 2.3. Short hand notation:  $I$  = immobilized,  $F$  = Freely Diffusing,  $S$  = Subdiffusive, and  $O$  = Outlier. We indicate a member of a cluster was removed and classified as Outlier by  $-/O$ .

$[A]_{\text{exo}}(\mu\text{g/mL})$	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5
0	I	I	F	F	-
0.033	I	I	I	F	-
0.100	I	I	F/O	F	-
0.333	I	I	I	F	-
1.0	I	O	S/O	S	F




**Fig. 3 (a)-(e)** A depiction of the classification system, where each point, corresponds to a tracked virion with estimated diffusivity on a log 10 scale and average-ACF value, for Ab concentration 0, 0.033, 0.1, 0.333, and 1.0  $\mu\text{g/mL}$ , respectively. The numerical value of each point corresponds to the prescribed cluster by the hierarchical clustering algorithm and the color of the point represents the biological class.

#### 4.2 Donor F05

The dendrogram for Donor F05 was cut at a uniform height to yield four cluster for each tested exogenous Ab concentration. The biological label that we assign to each cluster is reported in Table 3. We display the clusters based on the path-wise statistics in Figure 4, where the number corresponds to the hierarchical cluster and the color corresponds to the biological interpretation of the cluster.

**Table 3** The classification of the clusters separated by the hierarchical clustering algorithm described in Section 2.3. Short hand notation:  $I$  = immobilized,  $F$  = Freely Diffusing,  $S$  = Subdiffusive, and  $O$  = Outlier. We indicate a member of a cluster was removed and classified as Outlier by  $-/O$ .

$[A]_{\text{exo}}(\mu\text{g/mL})$	Cluster 1	Cluster 2	Cluster 3	Cluster 4
0	I	I	F	O
0.033	I	I	S	F
0.100	I	I	I	F
0.333	I	I	I	F
1.0	I	I	I	F



**Fig. 4 (a)-(e)** A depiction of the classification system, where each point, corresponds to a tracked virion with estimated diffusivity on a log 10 scale and average-ACF value, for Ab concentration 0, 0.033, 0.1, 0.333, and 1.0 $\mu\text{g/mL}$ , respectively. The numerical value of each point corresponds to the prescribed cluster by the hierarchical clustering algorithm and the color of the point represents the biological class.

#### 4.3 Donor F08

The dendrogram for Donor F08 was cut at a uniform height to yield four cluster for each tested exogenous Ab concentration. The biological label that we assign to each cluster is reported in Table 4. We display the clusters based on the path-wise statistics in Figure 5, where the number corresponds to the hierarchical cluster and the color corresponds to the biological interpretation of the cluster.

**Table 4** The classification of the clusters separated by the hierarchical clustering algorithm described in Section 2.3. Short hand notation:  $I$  = immobilized,  $F$  = Freely Diffusing,  $S$  = Subdiffusive, and  $O$  = Outlier. We indicate a member of a cluster was removed and classified as Outlier by  $-/O$ .

$[A]_{\text{exo}}(\mu\text{g/mL})$	Cluster 1	Cluster 2	Cluster 3	Cluster 4
0	I	F	F	F
0.033	I	I	F	F
0.100	I	I	S	F
0.333	I	I	F	F
1.0	I	I	S	F

**Fig. 5 (a)-(e)** A depiction of the classification system, where each point, corresponds to a tracked virion with estimated diffusivity on a log 10 scale and average-ACF value, for Ab concentration 0, 0.033, 0.1, 0.333, and  $1.0\mu\text{g/mL}$ , respectively. The numerical value of each point corresponds to the prescribed cluster by the hierarchical clustering algorithm and the color of the point represents the biological class.

#### 4.4 Donor F13


The dendrogram for Donor F13 was cut at a uniform height to yield four cluster for each tested exogenous Ab concentration except for  $[A]_{\text{exo}} = 1.0\mu\text{g/mL}$ , which was cut at a uniform height to yield five clusters because one particle was separated into its own cluster. The biological label that we assign to each cluster is reported in Table 5. We display the clusters based on the path-wise statistics in Figure 6, where the number corresponds to the hierarchical cluster and the color corresponds to the biological interpretation of the cluster.

**Table 5** The classification of the clusters separated by the hierarchical clustering algorithm described in Section 2.3. Short hand notation:  $I$  = immobilized,  $F$  = Freely Diffusing,  $S$  = Subdiffusive, and  $O$  = Outlier. We indicate a member of a cluster was removed and classified as Outlier by  $-/O$ .

$[A]_{\text{exo}}(\mu\text{g/mL})$	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5
0	I	I	S	F	F
0.033	I	I	F	F	F
0.100	I	I	O	S	F
0.333	I	I	I	F/O	F
1.0	I	I	I	F	F

#### 4.5 Donor F15

The dendrogram for Donor F15 was cut at a uniform height to yield four cluster for each tested exogenous Ab concentration. The biological label that we assign to each cluster is reported in Table 6. We display the clusters based on the path-wise statistics in Figure 7, where the number corresponds to the hierarchical cluster and the color corresponds to the biological interpretation of the cluster.




**Fig. 6 (a)-(e)** A depiction of the classification system, where each point, corresponds to a tracked virion with estimated diffusivity on a log 10 scale and average-ACF value, for Ab concentration 0, 0.033, 0.1, 0.333, and 1.0  $\mu\text{g/mL}$ , respectively. The numerical value of each point corresponds to the prescribed cluster by the hierarchical clustering algorithm and the color of the point represents the biological class.

**Table 6** The classification of the clusters separated by the hierarchical clustering algorithm described in Section 2.3. Short hand notation:  $I$  = immobilized,  $F$  = Freely Diffusing,  $S$  = Subdiffusive, and  $O$  = Outlier. We indicate a member of a cluster was removed and classified as Outlier by  $-O$ .

$[A]_{\text{exo}}(\mu\text{g/mL})$	Cluster 1	Cluster 2	Cluster 3	Cluster 4
0	I	I	I	F
0.033	I	I	I	F
0.100	I	I	F	F
0.333	I	I	F	F
1.0	I	I	I	F

#### 4.6 Donor F17

The dendrogram for Donor F17 was cut at a uniform height to yield four cluster for each tested exogenous Ab concentration. The biological label that we assign to each cluster is reported in Table 7. We display the clusters based on the path-wise statistics in Figure 8, where the number corresponds to the hierarchical cluster and the color corresponds to the biological interpretation of the cluster.




**Fig. 7 (a)-(e)** A depiction of the classification system, where each point corresponds to a tracked virion with estimated diffusivity on a log 10 scale and average-ACF value, for Ab concentration 0, 0.033, 0.1, 0.333, and 1.0  $\mu\text{g/mL}$ , respectively. The numerical value of each point corresponds to the prescribed cluster by the hierarchical clustering algorithm and the color of the point represents the biological class.

**Table 7** The classification of the clusters separated by the hierarchical clustering algorithm described in Section 2.3.. Short hand notation:  $I$  = immobilized,  $F$  = Freely Diffusing,  $S$  = Subdiffusive, and  $O$  = Outlier. We indicate a member of a cluster was removed and classified as Outlier by  $-O$ .

$[A]_{\text{exo}}(\mu\text{g/mL})$	Cluster 1	Cluster 2	Cluster 3	Cluster 4
0	I	I	F	F
0.033	I	I	I	F
0.100	I	I	F	F
0.333	I	I	I	F
1.0	I	I	I	F

#### 4.7 Donor F21

The dendrogram for Donor 21 was cut at a uniform height to yield four cluster for each tested exogenous Ab concentration. The biological label that we assign to each cluster is reported in Table 8. We display the clusters based on the path-wise statistics in Figure 9, where the number corresponds to the hierarchical cluster and the color corresponds to the biological interpretation of the cluster.



**Fig. 8 (a)-(e)** A depiction of the classification system, where each point corresponds to a tracked virion with estimated diffusivity on a log 10 scale and average-ACF value, for Ab concentration 0, 0.033, 0.1, 0.333, and 1.0  $\mu\text{g/mL}$ , respectively. The numerical value of each point corresponds to the prescribed cluster by the hierarchical clustering algorithm and the color of the point represents the biological class.

**Table 8** The classification of the clusters separated by the hierarchical clustering algorithm described in Section 2.3. Short hand notation:  $I$  = immobilized,  $F$  = Freely Diffusing,  $S$  = Subdiffusive, and  $O$  = Outlier. We indicate a member of a cluster was removed and classified as Outlier by  $-/O$ .

$[A]_{\text{exo}}(\mu\text{g/mL})$	Cluster 1	Cluster 2	Cluster 3	Cluster 4
0	I	I	F	F
0.033	I	I	F	F
0.100	I	I	I	F
0.333	I	I	F	F
1.0	I	I	I	F


## 5 Switch Point Detection Algorithm

### 5.1 Hamiltonian Monte Carlo (HMC) Sampling Algorithm

We used an HMC sampling algorithm to jointly estimate the parameters  $(D, \tilde{\kappa}, \tau)$ . Hamiltonian Monte Carlo is a form of Markov Chain Monte Carlo sampling that relies on Bayes Rule to sample from the joint posterior distribution of  $(D, \tilde{\kappa}, \tau)$ ,

$$p(D, \tilde{\kappa}, \tau | \mathbf{x}, \mathbf{y}) \stackrel{c}{=} L(\mathbf{x}, \mathbf{y}; D, \tilde{\kappa}, \tau) p(D, \tilde{\kappa}, \tau) \quad (2)$$

where  $L(\mathbf{x}, \mathbf{y}; (D, \tilde{\kappa}, \tau))$  is the likelihood of seeing the data given the parameters  $(D, \tilde{\kappa}, \tau)$  and  $p(D, \tilde{\kappa}, \tau)$  is the prior joint distribution of  $(D, \tilde{\kappa}, \tau)$ . In our model,



**Fig. 9 (a)-(e)** A depiction of the classification system, where each point corresponds to a tracked virion with estimated diffusivity on a log 10 scale and average-ACF value, for Ab concentration 0, 0.033, 0.1, 0.333, and 1.0  $\mu\text{g/mL}$ , respectively. The numerical value of each point corresponds to the prescribed cluster by the hierarchical clustering algorithm and the color of the point represents the biological class.

we assume the parameters  $D$ ,  $\tilde{\kappa}$ , and  $\tau$  are independent so that the joint prior distribution factors into the product of priors,  $p(D, \tilde{\kappa}, \tau) = p(D)p(\tilde{\kappa})p(\tau)$ .

The likelihood function,  $L(\mathbf{x}, \mathbf{y}; D, \tilde{\kappa}, \tau)$  is a consequence of the model. For the [diffusing  $\rightarrow$  immobilized] switching scenario and [immobilized  $\rightarrow$  diffusing] switching scenario  $L(\mathbf{x}, \mathbf{y}; (D, \tilde{\kappa}, \tau))$  is given by Equation 16 and 15, respectively.

As for the priors, we placed a loosely informative prior on  $D$

$$\pi(D) \sim \text{Gamma}(D_{\text{eff}} * 0.1, 0.1)$$

and chose a discrete uniform distribution over the observation times  $\{t_n\}_{n=1}^{N-1}$  for the prior distribution on  $\tau$ . Rather than directly sampling the parameter  $\tilde{\kappa}$ , we sampled the transform  $\rho = \exp(-\Delta\tilde{\kappa})$  and then used the inverse transformation  $\tilde{\kappa} = -\log(\rho)/\Delta$  to obtain posterior samples for  $\tilde{\kappa}$ . In doing this, we expressed our uncertainty of the order of  $\tilde{\kappa}$  by placing a lognormal prior on  $\rho$ ,

$$\pi(\rho) \sim \text{lognormal}\left(\text{loc} = \log\left(\frac{0.5^2}{\sqrt{0.5^2+2}}\right), \text{shape} = \log(1 + \frac{2}{0.5^2})^{1/2}\right).$$

We note that because  $\tau$  is a discrete parameter taking values in  $\{t_n\}_{n=1}^{N-1}$  whereas  $D, \tilde{\kappa}$  are continuous real-valued parameters, posterior samples of  $\tau|D, \tilde{\kappa}$  were first drawn from  $\{t_n\}_{n=1}^{N-1}$  with probability

$$p(\tau = t_j | \mathbf{x}, \mathbf{y}, D, \tilde{\kappa}) = \frac{L(\mathbf{x}, \mathbf{y}; D, \tilde{\kappa}, \tau = t_j)p(\tau = t_j)}{\sum_{k=1}^N L(\mathbf{x}, \mathbf{y}; D, \tilde{\kappa}, \tau = t_k)p(\tau = t_k)} \quad \text{for } j = 1, \dots, N-1. \quad (3)$$

Then, we sampled  $D$  and  $\tilde{\kappa}$  using the marginal posterior distribution of  $D$  and  $\tilde{\kappa}$ ,

$$p(D, \tilde{\kappa} | \mathbf{x}, \mathbf{y}) \stackrel{c}{=} p(D)p(\tilde{\kappa}) \sum_{k=1}^N L(\mathbf{x}, \mathbf{y}; D, \tilde{\kappa}, \tau = t_k)p(\tau = t_k). \quad (4)$$

The HMC sampling algorithm was run using the R package rstan, and computations were done using high performance computing (HPC) resources and services provided by Technology Services at Tulane University, New Orleans, LA.

## 5.2 Testing Freely Diffusing population for switch candidates


Recall, we say a path is a candidate for switching if the 95% credible region for  $\tau | D, \tilde{\kappa}$  is completely contained within the interval  $[0.1T_{\text{final}}, 0.9T_{\text{final}}]$  where  $T_{\text{final}}$  is the duration of a path. We tested if the  $i$ -th Freely Diffusing virion of path duration  $T_{\text{final}}$ , was a [immobilization  $\rightarrow$  diffusion] switching candidate using the following procedure. Assuming the [immobilization  $\rightarrow$  diffusion] switching model, we obtained posterior samples of  $\tau | D, \tilde{\kappa}$  using our HMC sampling algorithm, in which we drew 2000 posterior samples and kept the last 1000 samples for 4 different runs of the algorithm. The resulting 4000 posterior samples of  $\tau | D, \tilde{\kappa}$  were used to construct the 95% credible interval for  $\tau$ . If this interval was contained in  $[0.1T_{\text{final}}, 0.9T_{\text{final}}]$ , then we marked the virion as a candidate for [immobilization  $\rightarrow$  diffusion] switching. We repeated this for all virions in the Freely Diffusing diffusing subpopulation to obtain the fraction of candidate for [immobilization  $\rightarrow$  diffusion] switching. We obtained the fraction of candidate for [diffusion  $\rightarrow$  immobilization] switching, following the same method described in the paragraph above but assuming the [diffusion  $\rightarrow$  immobilization] switching model.

The Gaussian kernel density estimate of the posterior samples of  $\tau$  within the 95% credible region for [immobilization  $\rightarrow$  diffusion] switch candidates, and [diffusion  $\rightarrow$  immobilization] switch candidates are displayed in top Figure 10 (a) and (b), respectively, for Donor F15. The path duration for each switch candidate is marked by a red  $x$ . In Figure 10 (c) and (d), we display the trajectory of an [immobilization  $\rightarrow$  diffusion] switch candidates and [diffusion  $\rightarrow$  immobilization] switch candidate, where the start of the trajectory by the green dot and the maximum a posteriori estimate for  $\tau$  is marked by the blue star.

## 5.3 Estimating the false discovery rate

To estimate the false discovery rate of our switch detection test, we simulated Feely diffusing trajectories using parameters obtained from the class of Freely Diffusing virions in our data. To do this, the number of simulated Brownian paths was set to the number of virions classified as Freely diffusing, 1689 virions. Let  $\mathbf{n}_{\text{free}}$  and  $\mathbf{D}_{\text{free}}$  denote all the path lengths and the effective diffusivities of the Freely Diffusing virions, respectively. For the  $i$ -th simulated Brownian path, the path length  $n_i$  was sampled from  $\mathbf{n}_{\text{free}}$  and the diffusivity constant  $D_i$  was sampled from  $\mathbf{D}_{\text{free}}$ .

Using our switch point criterion, we set the false discovery rate of [Immobilization  $\rightarrow$  Diffusion] switches to the percent of simulated Brownian particles that



**Fig. 10** (a)-(b) Violin plots for the posterior samples of  $\tau$  within the 95% credible interval for those Freely-Diffusing virions from Donor F15 considered [immobilization  $\rightarrow$  diffusion] switch candidates, and [Diffusion  $\rightarrow$  Immobilization] switch candidate, respectively. (c) The 2d trajectory [immobilization  $\rightarrow$  diffusion] switch candidate, (tracked particle 211 for Donor F15 at  $[A]_{\text{exo}} = 0.333\mu\text{g/mL}$ ) (d) The 2d trajectory [diffusion  $\rightarrow$  immobilization] switch candidate, (tracked particle 226 for Donor F15 at  $[A]_{\text{exo}} = 1.0\mu\text{g/mL}$ ) In Panels (c) and (d) the green circle denotes the initial position of the virion and the blue star corresponds to the estimated switch point.

were labeled as candidates for [Immobilization  $\rightarrow$  Diffusion] switching for a likelihood function given by Equation 16. Specifically, the  $i$ -th simulated Brownian path and [Immobilization  $\rightarrow$  Diffusion] switching model, the 95% credible region for  $\tau$  was constructed using the 4000 posterior samples of  $\tau|_{D,\tilde{\kappa}}$  obtained by the last 1000 (out of 2000) posterior samples of 4 runs of our HMC algorithm. We obtained an estimate for the false discovery rate of [Diffusion  $\rightarrow$  Immobilization] switches in the same manner but assuming a [Diffusion  $\rightarrow$  Immobilization] model.

#### 5.4 Estimating the power of the candidate switching test

In estimating the power of our test, we assumed that the switch point occurs within the middle 80% of the path. We believe this was a reasonable assumption because on average a Freely Diffusing virion had a path length of 80, and the first or last eight observations would be too noisy to capture the true dynamics of the virion. First, we created a simulated data set of 1689 [immobilization  $\rightarrow$

diffusion] paths from the explicit solution of SDE given in Equation 15. For the  $i$ th [immobilization  $\rightarrow$  diffusion] path, the model parameters  $(n_i, D_i, \tilde{\kappa}_i, \tau_i)$  were sampled from the summary statistics of the tracked virions:

$$n_i \sim \text{Unif}\{\mathbf{n}_{\text{free}}\}, \quad D_i \sim \text{Unif}\{\mathbf{D}_{\text{free}}\}, \quad \tilde{\kappa}_i \sim \text{Exp}(\lambda = 1/500), \text{ and } \tau_i \sim \text{Unif}\{[0.1n_i, 0.9n_i]\}.$$

We ran our sampling algorithm assuming the model given by Equation 16 on the simulated [immobilization  $\rightarrow$  diffusion] data. The sampling algorithm was run 4 times each for 2000 iterations where the last 1000 were kept as posterior samples. We then set the power of our [immobilization  $\rightarrow$  diffusion] switch point detecting test to the percentage of them have credible regions for  $\tau$  within  $[0.1T_{\text{final}}, 0.9T_{\text{final}}]$ . The test failed to detect switches in the following two cases: (1) The switch was located near the endpoints of the interval  $[0.1T_{\text{final}}, 0.9T_{\text{final}}]$ , resulting in a credible region outside the region allowed by the test. (2) When  $\tilde{\kappa}$  is approximately less than 10. In this case, the immobilized state was indistinguishable from Brownian Motion because the deterministic component of the SDE given in Equation 16 was relatively small with respect to the random component. In Figure 11(a)-(b), we display the relative position of the change point and  $\tilde{\kappa}$  for simulated paths that were undetected by our proposed test. The trajectory of the simulated path marked by the pink diamond is shown in Figure 11(c)-(d).

We estimated the power of [diffusion  $\rightarrow$  immobilization] switch point detecting test following the same procedure as above but assumed the model given by Equation 15. For this switching scenario, switches went undetected due to the same reasons mentioned above.

## 6 Statistical Evidence for trend in proportion immobilized


In Figure 12, we display the 95%  $BC_a$  confidence interval. for the observed proportion immobilized for each donor. We remark that for all donors, the proportion immobilized at the lowest tested exogenous Ab concentration ( $0\mu\text{g/mL}$ ) and the highest tested exogenous Ab concentration ( $1\mu\text{g/mL}$ ) have non-overlapping confidence intervals. In Table 9, we report the estimates coefficient and associated  $p$ -values for a negative exponential growth for the observed proportion of virions immobilized.

**Table 9** Negative exponential growth coefficient estimates and associated  $p$ -values for the observed fraction of virions immobilized.

	Estimate	$p$ -value
$\beta_0$	0.9138	< 0.0001
$\alpha_0$	-0.8427	0.0043
$\alpha_{\text{exo}}$	15.9201	< 0.0001

	Estimate	$p$ -value
$\beta_1$	0.0642	0.4547
$\beta_2$	-0.1084	0.2123
$\beta_4$	-0.0700	0.4156
$\beta_5$	0.1140	0.1913
$\beta_6$	0.0050	0.9534
$\beta_7$	0.0347	0.6844

	Estimate	$p$ -value
$\alpha_1$	0.2846	0.3952
$\alpha_2$	0.2272	0.5049
$\alpha_4$	0.4286	0.1852
$\alpha_5$	0.4385	0.1745
$\alpha_6$	0.1901	0.5819
$\alpha_7$	-0.2406	0.5763



**Fig. 11** (a)-(b) The true parameter  $\bar{\kappa}$  and the relative position of the switch point,  $\tau$ , for the simulated switching particles, [immobilization  $\rightarrow$  diffusion] and [diffusion  $\rightarrow$  immobilized], respectively, whose switch point was undetected by our switching detection test. The dashed navy vertical lines mark the tenth percentile and ninetieth percentile of the trajectory. (c)-(d) The trajectory of a simulated switching particles marked by the pink diamond panels (a) and (b), respectively.

## 7 Statistical Evidence for Effective Diffusivity


In Figure 13, we display the 95%  $BC_a$  confidence interval for the ensemble effective diffusivity for the Freely-Diffusing subpopulation. We remark that for all donors, the ensemble effective diffusivity at the lowest tested exogenous Ab concentration (0  $\mu\text{g/mL}$ ) and the highest tested exogenous Ab concentration (1  $\mu\text{g/mL}$ ) have overlapping confidence intervals. We report the results of all the paired difference tests of the form,

$$H_0 : \langle D_{\text{eff}}([A]_i) \rangle - \langle D_{\text{eff}}([A]_j) \rangle = 0, \quad H_A : \langle D_{\text{eff}}([A]_i) \rangle - \langle D_{\text{eff}}([A]_j) \rangle > 0. \quad (5)$$


for  $1 \leq i < j \leq 5$  in Table 10.

**Table 10** The  $t$ -values of all possible combination of paired-difference tests of the form Equation 5. The critical value  $t_{\alpha=0.05,6} = 1.943$ .

	(1,2)	(1,3)	(1,4)	(1,5)	(2,3)	(2,4)	(2,5)	(3,4)	(3,5)	(4,5)
$t$ -value	-0.7592	0.6669	0.0879	0.2325	1.2058	1.124	1.492	-0.2887	-0.3973	0.0931



**Fig. 12** (a)-(g) For each donor, the observed proportion immobilized with 95%  $BC_a$  confidence interval.



**Fig. 13** (a)-(g) For each donor, the ensemble effective diffusivity of the free subpopulation with 95%  $BC_a$  confidence interval.

## 8 Simple Linear Model

In 11, we report the range of each parameter for each donor and whether it was identifiable, practically unidentifiable, or structurally unidentifiable, under the assumption it takes one simultaneously bound Ab to immobilize a virion ( $T = 1$ ) and the Ab-mucin binding rate is not affected by immobilization ( $c = 1$ ).

**Table 11** Parameter Identifiable assuming that  $T = 1$  and  $c = 1$  where ID indicates a structurally identifiable parameter and PU refers to a practically unidentifiable parameters. The confidence interval are given for each parameter where  $\theta_{\alpha=0.95,3} = 7.814728$ .

	$[A]_0$ ( $\mu\text{g/mL}$ )	$N_*$	$q$	$\min(\chi^2(\boldsymbol{\theta}))$
Donor F02	ID, $[0.006, 0.04]$	ID, $[140, 560]$	ID, $[0.89, 0.969]$	5.2106
Donor F05	ID, $[0.015, 0.085]$	ID, $[60, 260]$	ID, $[0.76, 0.908]$	1.6292
Donor F08	ID, $[0.020, 0.115]$	ID, $[80, 360]$	ID, $[0.86, 0.954]$	0.2755
Donor F13	ID, $[0.020, 0.075]$	ID, $[50, 100]$	PU, $[0.93, 1)$	15.3627
Donor F15	ID, $[0.003, 0.01]$	ID, $[310, 780]$	ID, $[0.93, .963]$	4.6863
Donor F17	ID, $[0.025, 0.095]$	ID, $[80, 210]$	ID, $[0.90, .969]$	3.661
Donor F21	ID, $[0.035, 0.195]$	ID, $[70, 250]$	ID, $[0.92, .989]$	1.277

## 9 Full Model


Let  $\hat{\boldsymbol{\theta}}$  denote the numeric estimate for  $\boldsymbol{\theta} \in \Theta$  and  $\hat{\boldsymbol{\theta}}_{\text{slow}}$  denote the numeric estimate for  $\boldsymbol{\theta}$  in the restricted parameter space  $\Theta_{0.05,5} \cap \Theta_{\text{slow}}$ . In 12, we report the range of each parameter for each donor within  $\Theta_{0.05,5} \cap \Theta_{\text{slow}}$  and whether it was identifiable, practically unidentifiable, or structurally unidentifiable. We display the proportion immobilized predicted by our model assuming  $\hat{\boldsymbol{\theta}}_{\text{slow}}$  is the numeric estimate of the parameters restricted to the subspace  $\Theta_{0.05,5} \cap \Theta_{\text{slow}}$  for each Donor in Figure 14.

**Table 12** The values of  $T$ ,  $c$ , and  $N_*$  that can permit  $\boldsymbol{\theta} \in \Theta_{0.05,5} \cap \Theta_{\text{slow}}$ . Lower and upper bounds of the tested parameter range are underlined. Note that  $\chi^2(\alpha = 0.95, 5) = 11.0705$ .

	$T$	$c$	$N_*$	$\chi^2(\hat{\boldsymbol{\theta}})$	$\chi^2(\hat{\boldsymbol{\theta}}_{\text{slow}})$
Donor F02	<u>19,60</u>	$[44.67, 10^4]$	$[120, 620]$	0.7070	9.1772
Donor F05	<u>16,60</u>	$[79.43, 10^4]$	$[90, 370]$	0.3912	4.1879
Donor F08	<u>17,60</u>	$[56.23, 10^4]$	$[100, 540]$	0.1562	1.2343
Donor F13	{13}	$[2818.38, 10^4]$	{70}	10.0055	18.0788
Donor F15	<u>30,60</u>	$[44.68, 10^4]$	$[250, 630]$	4.2209	4.2209
Donor F17	17,34	$[281.84, 10^4]$	$[100, 200]$	1.6432	3.0156
Donor F21	<u>19,27</u>	$[10^3, 10^4]$	$[120, 170]$	0.5974	8.1838

## References

1. Vestergaard, C.L., Blainey, P.C., Flyvbjerg, H.: Optimal estimation of diffusion coefficients from single-particle trajectories. *Physical Review E* **89**(2), 022726 (2014)



**Fig. 14 (a)-(g)** For each donor the observed proportion immobilized (dashed lines), proportion immobilized predicted by  $\tilde{\pi}([A]; \hat{\theta})$ , (solid black line), where  $\hat{\theta}$  is the numeric estimate for  $\theta$  and the proportion immobilized predicted by  $\tilde{\pi}([A]; \hat{\theta}_{\text{slow}})$  (solid blue line), where  $\hat{\theta}_{\text{slow}}$  is the numeric estimate restricted to the subspace  $\Theta_{0.05,5} \cap \Theta_{\text{slow}}$ . Note in Figure (e)  $\hat{\theta}_{\text{slow}} = \hat{\theta}$ .