

Perspective

Stable Carbenes, Nitrenes, Phosphinidenes, and Borylenes: Past and Future

Michele Soleilhavoup^{1,*} and Guy Bertrand^{1,*}

The different modes of stabilization for carbenes are briefly discussed in the context of the discovery of their first stable representatives. Largely because a diversity of stable carbenes are available today, their use has spanned across the chemical sciences, including medicinal and material applications. Much less is known about the isolobal group 15 cousins of carbenes, namely nitrenes and phosphinidenes, given that only one example of each has been isolated. The difficulties associated with their applications are discussed, and possible solutions are presented. As for group 13 element carbenoids, two types are considered: (1) mono(Lewis base)-stabilized borylenes that have been recently isolated and (2) borylenes, the only carbenoids discussed in this article that have eluded the synthetic skills of investigators. This Perspective describes potentially attainable targets, such as monocoordinated aminocarbanions, aminocarbynes, and aminocarbocations, which feature a carbon atom with six, five, and four valence electrons, respectively.

The octet rule, one of the fundamental principles of organic chemistry, dates back to Mendeleev at the end of the 19th century and Abegg, Lewis, and Langmuir at the beginning of the 20th century.¹ Scientists love challenging rules, and one can argue that this motivated Gomberg to prepare the first stable radical, a species in which a carbon atom has only seven valence electrons. Similarly, Curtius and Staudinger attempted to synthesize carbenes, compounds in which a carbon atom features a six-valence-electron shell. In agreement with the octet rule, they found that carbenes were only transient intermediates but were involved in several important chemical transformations. Then, for many years, the quest for stable carbenes became an unreasonable target until the isolation of a distillable (phosphino)(silyl) carbene (**1**) in 1988² and a crystalline imidazol-2-ylidene (**2**) in 1991³ (Figure 1). Interestingly, the mode of stabilization for singlet carbenes **1** and **2** is totally different. In agreement with Pauling's prediction,⁴ our group believed that a π -donor and a π -acceptor substituent (a push-pull mesomeric substitution pattern) was necessary to decrease both the Lewis acidity and basicity of the carbene center of **1**. However, the isolation of Arduengo's carbene **2**, which features two π -donor substituents (a push-push mesomeric substitution pattern), showed that the instability of singlet carbenes was mainly due to their Lewis acid character, although one could argue that the amino groups have an inductive pull effect, decreasing the basicity of the lone pair.⁵ During the next decade, after the discovery of **1** and **2**, it was believed that two electronic active substituents were required to allow for the isolation of carbenes. However, at the beginning of the 21st century, we showed that a single π -donor phosphino⁶ or amino^{7,8} substituent, as exemplified by **3**, was enough to extend the lifetime of carbenes to weeks, provided that the other substituent was sufficiently bulky.

The Bigger Picture

Challenges and opportunities:

- Further development of carbenes to enable new and important chemical transformations and translation into medicinal and material chemistry applications.
- Stable nitrenes, phosphinidenes, (Lewis base)-stabilized borylenes, and their applications remain underdeveloped.
- Free borylenes have still eluded the synthetic skills of investigators, and species featuring a carbon atom with six, five, or four valence electrons should be exciting and attainable targets.






Figure 1. Stable Group 14 Carbenoids and Selected Complexes

Some of the first isolated carbenes (1–3), catalytically active complexes (4 and 5), an organocatalyst (6), some of the most popular carbenes (7–11), an efficient blue emitter for OLEDs (12), and the first stable silylene (13).

Very often in chemistry, when a novel type of compound is discovered, it is first considered a laboratory curiosity, and it sometimes takes decades for the first applications to appear. For example, phosphorus ylides were first synthesized in 1894 by Michaelis and Gimborn, but it was only in 1953 that Wittig and Geissler reported the so-called Wittig reaction, which was soon after used by BASF for the synthesis of vitamin A. This is not the case for carbenes given that already in 1995, Herrmann et al. reported that the palladium complex **4** bearing imidazol-2-ylidenes, similar to **2** as ancillary ligands, promoted the Mizoroki-Heck reaction.⁹ After this discovery, transition-metal complexes bearing carbenes—such as **2** and their saturated version, i.e., imidazolin-2-ylidene,¹⁰ the so-called N-heterocyclic carbenes (NHCs)—have been used for a variety of catalytic chemical transformations,¹¹ the most recognizable of which is the Grubbs second-generation olefin metathesis catalyst **5**.¹² In parallel, the 1,2,4-triazolin-5-ylidene **6**, reported by Enders,¹³ has proven to be an excellent organocatalyst on its own, in line with the pioneering work by Breslow on the thiazolidene-catalyzed benzoin condensation.¹⁴

Nowadays, aside from the carbenes mentioned above, a myriad of stable carbenes with distinct steric and electronic properties are available. Arguably, the most popular are the cyclic (alkyl)(amino)carbenes (CAACs) **7**,^{15,16} N,N'-diamidocarbenes (DACs) **8**,^{17,18} benzimidazolylidenes **9**,¹⁹ mesoionic carbenes **10a**^{20–22} and **10b**,^{23,24} and the cyclopropenylidene **11**.²⁵ As a consequence of this diversity, the use of carbenes disseminated across chemical sciences at large, including medicinal²⁶ and material²⁷ applications. Furthermore, thanks in part to carbene's

¹UCSD-CNRS Joint Research Laboratory, UMI 3555, Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, CA 92093-0358, USA

*Correspondence:
msoleilhavoup@ucsd.edu (M.S.),
gbertrand@ucsd.edu (G.B.)

<https://doi.org/10.1016/j.chempr.2020.04.015>




Figure 2. Group 15 Carbenoids

The isolated nitrene **14** and its reactivity with CO_2 exemplify the non-innocence of the phosphorus center. Aminonitrene **15** is an attractive target. The isolated phosphinidene **16** behaves as an electrophile, as shown by the formation of **17**, and to some extent as a transition metal given that it reacts with CO to give **19**, which undergoes ligand exchange, as shown by **17** and **18**.

pronounced ability to reversibly activate enthalpically strong bonds and small molecules,^{28,29} there has been significant development toward their use as transition-metal surrogates. They can also stabilize low-coordinate main-group compounds^{30–32} and organic and inorganic paramagnetic species³³ and can be used for the functionalization of nanoscale and bulk surfaces.³⁴ They even enabled the preparation of organic light-emitting diodes (OLEDs) based on earth-abundant metals,^{35,36} which achieve performance comparable to that of the state-of-the-art luminescent complexes of iridium, platinum, and ruthenium. As an example, the two-coordinate copper(I) complex **12** achieved photoluminescence efficiencies > 99% and microsecond lifetimes, which lead to an efficient blue-emitting OLED.³⁷

The discovery of stable carbenes was followed by the isolation of their heavier analogs, e.g., silylenes. Indeed, in 1994 Denk et al. reported the isolation of **13**,³⁸ the design of which was clearly derived from the corresponding NHC **2**. Although the applications of silylenes are still very limited, some encouraging results were obtained when they were used as ligands in transition-metal catalysis.³⁹

Much less is known about the isolobal group 15 cousins of carbenes and silylenes, namely nitrenes and phosphinidenes (Figure 2). In 2012, our group isolated the first nitrene (**14**).⁴⁰ The bonding between phosphorus and nitrogen is analogous to that observed for metallonitrenes, which are postulated as key catalytic species in the industrial Haber-Bosch hydrogenation of N_2 into NH_3 . However, since its discovery, no striking developments have occurred with **14**. This is mainly due to the presence of the phosphino substituent. Indeed, although nitrene **14** can activate small




Figure 3. Group 13 Carbenoids

The first attempts to prepare mono(Lewis base)-stabilized borylenes (**21a** and **21b**) and their isolated counterparts **21c** and **21d** (metallomimics), as shown by their reactivity with H₂ and CO. The elimination of CO under irradiation from **26** leads to a transient mono(Lewis base)-stabilized borylene (**21e**), which can be trapped by an isonitrile to give the bis(Lewis base)-stabilized borylene **27**. The transient formation of borylenes of type **28** has been postulated on the basis their chemical reactivity, as shown by **29**.

molecules,⁴¹ the involvement of phosphorus in the reaction, as exemplified with CO₂, might preclude the use of this nitrene in catalytic processes. We believe, however, that a stable amino nitrene, such as that derived from Dervan's scaffold (**15**),⁴² would be a more promising candidate. The only isolated phosphinidene (**16**) to date also features a bulky and strong π -donor phosphino substituent.⁴³ Of particular interest, despite its phosphorus-phosphorus (PP) multiple-bond character and the presence of a partial negative charge on the terminal phosphorus, **16** behaves as an electrophile, as expected for an atom featuring a formal valence sextet structure. Indeed, **16** reacts with Lewis bases, such as phosphines, and prefers the more basic PCy₃ over PPh₃.⁴⁴ Importantly, phosphinidene **16** resembles a transition metal to some extent given that it reacts with CO, giving **19**, and undergoes ligand exchange.⁴⁵ For example, **19** reacts with phosphines to afford the corresponding adduct **17**. Similarly, the latter reacts with isonitriles to give **18**. It is important to note that these ligand-exchange reactions can be performed starting from non-hindered phosphinidene-CO adducts, which are readily available from the reaction of the corresponding chlorophosphine with NaPCO.⁴⁶ Just as for nitrenes, amino-substituted phosphinidenes would be attractive targets because they might give the opportunity to isolate both the singlet and the triplet states of a given compound, which has never been done with carbenoid species.^{47,48} Indeed, calculations by Nguyen et al.⁴⁹ predicted that the singlet-triplet gap for amino phosphinidenes would be below ± 3 kcal/mol; in other words, the ground state would have two degenerate orbitals.⁵⁰

As for group 13 element carbenoids, two species have to be considered: mono(Lewis base)-stabilized borylenes (**21**) and borylenes (**28**) (Figure 3).⁵¹ The former are isoelectronic with carbenes with a six-electron valence shell, whereas the latter have only a four-electron valence shell. The first attempt to isolate a compound of type **21** was reported in 2007 by Robinson and co-workers⁵² through reduction of the (NHC)BBr₃ adduct **20**. They isolated the diborene **22**, which can be regarded as a dimer of the desired mono(Lewis base)-stabilized borylene **21a**. As already stated, carbenes and related species, featuring a sextet of electrons, need a π -donor substituent to be isolated. Although the (NHC)aminoborylene **21b** fulfilled this criterion, it was not isolable because an intramolecular CH insertion gave **23**.³¹ Because of its electropositivity, boron was reluctant to be electron rich,⁵³ and thus more electrophilic carbenes than NHCs seemed more appropriate to stabilize a compound of type **21**. Indeed, reduction of the DAC adduct of the diisopropylaminodichloroborane **24** and of the CAAC adduct of the bis(trimethylsilyl)aminodichloroborane **25** afforded the desired mono(Lewis base)-stabilized borylenes **21c** and **21d**.^{54,55} Unsurprisingly, both compounds feature an almost linear allenic structure ($>C=B=NR_2$), but their flexibility gives rise to a highly electrophilic boron center that has carbene-like behavior. Akin to electrophilic singlet carbenes, **21d** can activate small molecules, such as H₂, and coordinate an additional ligand, such as CO; in other words, compounds **21** are boron metallomimics.^{56–58} Along this line, Braunschweig et al.⁵⁹ have shown that the CO ligand of **26** can be removed by photolysis to afford the transient mono(Lewis base)-stabilized borylenes **21e**, which can be trapped by Lewis bases to afford novel bis(Lewis base)-stabilized borylenes **27**.^{60,61}




Figure 4. The Stable Mono-substituted Carbene 30 and Future Targets 31–33

In contrast to the other carbenoids, borylenes **28** have two vacant orbitals. These compounds are well known in the coordination sphere of transition metals, and Braunschweig et al. have shown that these complexes display a rich chemistry.^{62,63} Until now, metal-free borylenes have been spectroscopically characterized only in the gas phase or in inert matrices at very low temperatures.⁶⁴ In addition, the transient formation of borylenes **28** has been postulated on the basis of their chemical reactivity,⁶⁵ as exemplified with the formation of **29**.⁶⁶

In conclusion, it appears that borylenes **28** are the only carbenoids we discuss that have eluded the synthetic skills of investigators. However, contrary to carbenes, nitrenes, and phosphinidenes, which feature either a singlet or a triplet ground state, all computationally studied borylenes **28** have a singlet ground state;⁶⁷ therefore, we believe that with the right substituent, they should be isolable. Hope for a stable borylene **28** comes from the recent isolation of the mono-substituted carbene **30** (Figure 4), which demonstrates that a single bulky amino substituent could single handedly tame the intrinsic tendency of carbenes toward dimerization.⁶⁸ Along this line, we believe that monocoordinated aminocarbanions (**31**) could be isolated. They have six valence electrons, similar to carbenes, and two lone pairs and a vacant orbital, like phosphinidenes **16**. It might even be possible to undress carbon even more. What about aminocarbynes (**32**) and aminocarbocations (**33**),⁶⁹ compounds in which the carbon center formally features only five and four valence electrons, respectively? Not only are these molecules exciting synthetic challenges, but their unique properties should also unveil new applications. As an illustration, note that carbenes can simultaneously form two sigma bonds, whereas monocoordinated compounds could form three,⁷⁰ which opens a new mode of reactivity.

ACKNOWLEDGMENTS

This work was supported by the National Science Foundation (CHE-1661518), the ACS Petroleum Research Fund (60776-ND1), and the Catalysis Science program of the Basic Energy Sciences program of the US Department of Energy's Office of Science (DE-SC0009376).

AUTHOR CONTRIBUTIONS

M.S. and G.B. wrote the manuscript.

REFERENCES

- Jensen, W.B. (1984). Abegg, Lewis, Langmuir, and the octet rule. *J. Chem. Educ.* **61**, 191–200.
- Igau, A., Grutzmacher, H., Baceiredo, A., and Bertrand, G. (1988). Analogous $\alpha\alpha'$ bis carbenoid triply bonded species: synthesis of a stable λ^3 -phosphinocarbene – λ^5 -phosphaacetylene. *J. Am. Chem. Soc.* **110**, 6463–6466.
- Arduengo, A.J., III, Harlow, R.L., and Kline, M. (1991). A stable crystalline carbene. *J. Am. Chem. Soc.* **113**, 361–363.
- Pauling, L. (1980). The structure of singlet carbene molecules. *J. Chem. Soc. Chem. Commun.* **1980**, 688–689.
- Sauers, R. (2014). Singlet carbene stability: linear free-energy analysis of substituent effects. *Arkivoc* **2014**, 376–383.

6. Buron, C., Gornitzka, H., Romanenko, V., and Bertrand, G. (2000). Stable versions of transient push-pull carbenes: extending lifetimes from nanoseconds to weeks. *Science* 288, 834–836.
7. Sole, S., Gornitzka, H., Schoeller, W.W., Bourissou, D., and Bertrand, G. (2001). (Amino)(aryl)carbenes: stable singlet carbenes featuring a spectator substituent. *Science* 292, 1901–1903.
8. Lavallo, V., Mafhouz, J., Canac, Y., Donnadiou, B., Schoeller, W.W., and Bertrand, G. (2004). Synthesis, reactivity, and ligand properties of a stable alkyl carbene. *J. Am. Chem. Soc.* 126, 8670–8671.
9. Herrmann, W.A., Elison, M., Fischer, J., Köcher, C., and Artus, G.R.J. (1995). Metal complexes of N-heterocyclic carbenes—a new structural principle for catalysts in homogeneous catalysis. *Angew. Chem. Int. Ed.* 34, 2371–2374.
10. Arduengo, A.J., III, Goerlich, J.R., and Marshall, W.J. (1995). A stable diaminocarbene. *J. Am. Chem. Soc.* 117, 11027–11028.
11. Díez-González, S. (2016). N-Heterocyclic Carbenes: from Laboratory Curiosities to Efficient Synthetic Tools (Royal Society of Chemistry).
12. Scholl, M., Ding, S., Lee, C.W., and Grubbs, R.H. (1999). Synthesis and activity of a new generation of ruthenium-based olefin metathesis catalysts coordinated with 1,3-dimesityl-4,5-dihydroimidazol-2-ylidene ligands. *Org. Lett.* 1, 953–956.
13. Enders, D., Breuer, K., Raabe, G., Runsink, J., Teles, J.H., Melder, J.P., Ebel, K., and Brode, S. (1995). Preparation, structure, and reactivity of 1,3,4-triphenyl-4,5-dihydro-1*H*-1,2,4-triazol-5-ylidene, a new stable carbene. *Angew. Chem. Int. Ed.* 34, 1021–1023.
14. Breslow, R. (1958). On the mechanism of thiamine action. IV.¹ Evidence from studies on model systems. *J. Am. Chem. Soc.* 80, 3719–3726.
15. Lavallo, V., Canac, Y., Präsang, C., Donnadiou, B., and Bertrand, G. (2005). Stable cyclic (alkyl)(amino)carbenes as rigid or flexible, bulky, electron-rich ligands for transition-metal catalysts: a quaternary carbon atom makes the difference. *Angew. Chem. Int. Ed.* 44, 5705–5709.
16. Melaimi, M., Jazzar, R., Soleilhavoup, M., and Bertrand, G. (2017). Cyclic (alkyl)(amino)carbenes (CAACs): recent developments. *Angew. Chem. Int. Ed.* 56, 10046–10068.
17. Moerdyk, J.P., and Bielawski, C.W. (2012). Diamidocarbenes as versatile and reversible [2+1] cycloaddition reagents. *Nat. Chem.* 4, 275–280.
18. Moerdyk, J.P., Schilter, D., and Bielawski, C.W. (2016). N,N'-diamidocarbenes: isolable divalent carbons with bona fide carbene reactivity. *Acc. Chem. Res.* 49, 1458–1468.
19. Hahn, F.E., Wittenbecher, L., Boese, R., and Bläser, D. (1999). N-N'-bis(2,2-dimethylpropyl)benzimidazolin-2-ylidene: a stable nucleophilic carbene derived from benzimidazole. *Chem. Eur. J.* 5, 1931–1935.
20. Aldeco-Perez, E., Rosenthal, A.J., Donnadiou, B., Parameswaran, P., Frenking, G., and Bertrand, G. (2009). Isolation of a C5-deprotonated imidazolium, a crystalline “abnormal” N-heterocyclic carbene. *Science* 326, 556–559.
21. Vivancos, Á., Segarra, C., and Albrecht, M. (2018). Mesoionic and related less heteroatom-stabilized N-heterocyclic carbene complexes: synthesis, catalysis, and other applications. *Chem. Rev.* 118, 9493–9586.
22. Sau, S.C., Hota, P.K., Mandal, S.K., Soleilhavoup, M., and Bertrand, G. (2020). Stable abnormal N-heterocyclic carbenes and their applications. *Chem. Soc. Rev.* 49, 1233–1252.
23. Guisado-Barrios, G., Bouffard, J., Donnadiou, B., and Bertrand, G. (2010). Crystalline 1*H*-1,2,3-triazol-5-ylidenes: new stable mesoionic carbenes (MICs). *Angew. Chem. Int. Ed.* 49, 4759–4762.
24. Guisado-Barrios, G., Soleilhavoup, M., and Bertrand, G. (2018). 1*H*-1,2,3-triazol-5-ylidenes: readily available mesoionic carbenes. *Acc. Chem. Res.* 51, 3236–3244.
25. Lavallo, V., Canac, Y., Donnadiou, B., Schoeller, W.W., and Bertrand, G. (2006). Cyclopropenylidenes: from interstellar space to an isolated derivative in the laboratory. *Science* 312, 722–724.
26. Johnson, N.A., Southerland, M.R., and Youngs, W.J. (2017). Recent developments in the medicinal applications of silver-NHC complexes and imidazolium salts. *Molecules* 22, 1263.
27. For an excellent review on the applications of stable carbenes, see: Hopkinson, M.N., Richter, C., Schedler, M., and Glorius, F. (2014). An overview of N-heterocyclic carbenes. *Nature* 510, 485–496.
28. Tolentino, D.R., Neale, S.E., Isaac, C.J., Macgregor, S.A., Whittlesey, M.K., Jazzar, R., and Bertrand, G. (2019). Reductive elimination at carbon under steric control. *J. Am. Chem. Soc.* 141, 9823–9826.
29. Ryu, Y., Ahumada, G., and Bielawski, C.W. (2019). Redox- and light-switchable N-heterocyclic carbenes: a “soup-to-nuts” course on contemporary structure-activity relationships. *Chem. Commun.* 55, 4451–4466.
30. Khan, S., and Roesky, H.W. (2019). Carbene-stabilized exceptional silicon halides. *Chemistry* 25, 1636–1648.
31. Wang, Y., and Robinson, G.H. (2011). Carbene stabilization of highly reactive main-group molecules. *Inorg. Chem.* 50, 12326–12337.
32. Nesterov, V., Reiter, D., Bag, P., Frisch, P., Holzner, R., Porzelt, A., and Inoue, S. (2018). NHCs in Main Group Chemistry. *Chem. Rev.* 118, 9678–9842.
33. Kundu, S., Sinhababu, S., Chandrasekhar, V., and Roesky, H.W. (2019). Stable cyclic (alkyl)(amino)carbene (CAAC) radicals with main group substituents. *Chem. Sci.* 10, 4727–4741.
34. Zhukhovitskiy, A.V., MacLeod, M.J., and Johnson, J.A. (2015). Carbene ligands in surface chemistry: from stabilization of discrete elemental allotropes to modification of nanoscale and bulk substrates. *Chem. Rev.* 115, 11503–11532.
35. Di, D., Romanov, A.S., Yang, L., Richter, J.M., Rivett, J.P.H., Jones, S., Thomas, T.H., Abdi Jalebi, M.A., Friend, R.H., Linnolahti, M., et al. (2017). High-performance light-emitting diodes based on carbene-metal-amides. *Science* 356, 159–163.
36. Hamze, R., Shi, S., Kapper, S.C., Muthiah Ravinson, D.S., Estergreen, L., Jung, M.C., Tadde, A.C., Haiges, R., Djurovich, P.I., Peltier, J.L., et al. (2019). “Quick-Silver” from a systematic study of highly luminescent, 2-coordinate, d10 coinage metal complexes. *J. Am. Chem. Soc.* 141, 8616–8626.
37. Hamze, R., Peltier, J.L., Sylvinson, D., Jung, M., Cardenas, J., Haiges, R., Soleilhavoup, M., Jazzar, R., Djurovich, P.I., Bertrand, G., and Thompson, M.E. (2019). Eliminating nonradiative decay in Cu(I) emitters: >99% quantum efficiency and microsecond lifetime. *Science* 363, 601–606.
38. Denk, M., Lennon, R., Hayashi, R., West, R., Belyakov, A.V., Verne, H.P., Haaland, A., Wagner, M., and Metzler, N. (1994). Synthesis and structure of a stable silylene. *J. Am. Chem. Soc.* 116, 2691–2692.
39. Zhou, Y.P., and Driess, M. (2019). Isolable silylene ligands can boost efficiencies and selectivities in metal-mediated catalysis. *Angew. Chem. Int. Ed.* 58, 3715–3728.
40. Dielmann, F., Back, O., Henry-Ellinger, M., Jerabek, P., Frenking, G., and Bertrand, G. (2012). A crystalline singlet phosphinonitrene: a nitrogen atom-transfer agent. *Science* 337, 1526–1528.
41. Dielmann, F., and Bertrand, G. (2015). Reactivity of a stable phosphinonitrene towards small molecules. *Chemistry* 21, 191–198.
42. Hinsberg, W.D., Schultz, P.G., and Dervan, P.B. (1982). Direct studies of 1,1-diazenes. Syntheses, infrared and electronic spectra, and kinetics of the thermal decomposition of N-(2,2,6,6-tetramethylpiperidyl)nitrene and N-(2,2,5,5-tetramethylpyrrolidyl)nitrene. *J. Am. Chem. Soc.* 104, 766–773.
43. Liu, L., Ruiz, D.A., Munz, D., and Bertrand, G. (2016). A room temperature stable singlet phosphinidene. *Chem* 1, 147–153.
44. Hansmann, M.M., Jazzar, R., and Bertrand, G. (2016). Singlet (phosphino)phosphinidenes are electrophilic. *J. Am. Chem. Soc.* 138, 8356–8359.
45. Hansmann, M.M., and Bertrand, G. (2016). The transition-metal-like behavior of main group elements: ligand exchange at a phosphinidene. *J. Am. Chem. Soc.* 138, 15885–15888.
46. Goicoechea, J.M., and Grützmacher, H. (2018). The chemistry of the 2-phosphaethynolate anion. *Angew. Chem. Int. Ed.* 57, 16968–16994.
47. Perera, T.A., Reinheimer, E.W., and Hudnall, T.W. (2017). Photochemically switching diamidocarbene spin states leads to reversible Büchner ring expansions. *J. Am. Chem. Soc.* 139, 14807–14814.
48. Sultane, P.R., Ahumada, G., Janssen-Müller, D., and Bielawski, C.W. (2019). Cyclic(aryl)(amido)carbenes: NHCs with triplet-like reactivity. *Angew. Chem. Int. Ed.* 58, 16320–16325.

49. Nguyen, M.T., Van Keer, A.V., and Vanquickenborne, L.G. (1996). In search of singlet phosphinidenes. *J. Org. Chem.* 61, 7077–7084.
50. Skell, P.S. (1985). The beginnings of modern carbene chemistry triplets and singlets. *Tetrahedron* 41, 1427–1428.
51. Soleilhavoup, M., and Bertrand, G. (2017). Borylenes: an emerging class of compounds. *Angew. Chem. Int. Ed.* 56, 10282–10292.
52. Wang, Y., Quillian, B., Wei, P., Wannere, C.S., Xie, Y., King, R.B., Schaefer, H.F., III, Schleyer, P.V., and Robinson, G.H. (2007). A stable, neutral diborene containing a B=B double bond. *J. Am. Chem. Soc.* 129, 12412–12413.
53. Segawa, Y., Yamashita, M., and Nozaki, K. (2006). Boryllithium: isolation, characterization, and reactivity as a boryl anion. *Science* 314, 113–115.
54. Ledet, A.D., and Hudnall, T.W. (2016). Reduction of a diamidocarbene-supported borenium cation: isolation of a neutral boryl-substituted radical and a carbene-stabilized aminoborylene. *Dalton Trans.* 45, 9820–9826.
55. Dahchegh, F., Martin, D., Stephan, D.W., and Bertrand, G. (2014). Synthesis and reactivity of a CAAC-aminoborylene adduct: a hetero-allene or an organoboron isoelectronic with singlet carbenes? *Angew. Chem. Int. Ed.* 53, 13159–13163.
56. Braunschweig, H., Dewhurst, R.D., Hupp, F., Nutz, M., Radacki, K., Tate, C.W., Vargas, A., and Ye, Q. (2015). Multiple complexation of CO and related ligands to a main-group element. *Nature* 522, 327–330.
57. Légaré, M.A., Bélanger-Chabot, G., Dewhurst, R.D., Welz, E., Krummenacher, I., Engels, B., and Braunschweig, H. (2018). Nitrogen fixation and reduction at boron. *Science* 359, 896–900.
58. Légaré, M.A., Prankevicus, C., and Braunschweig, H. (2019). Metallomimetic chemistry of boron. *Chem. Rev.* 119, 8231–8261.
59. Braunschweig, H., Krummenacher, I., Légaré, M.A., Matler, A., Radacki, K., and Ye, Q. (2017). Main-group metallomimetics: transition metal-like photolytic CO substitution at boron. *J. Am. Chem. Soc.* 139, 1802–1805.
60. Kinjo, R., Donnadieu, B., Celik, M.A., Frenking, G., and Bertrand, G. (2011). Synthesis and characterization of a neutral tricoordinate organoboron isoelectronic with amines. *Science* 333, 610–613.
61. Kong, L., Li, Y., Ganguly, R., Vidovic, D., and Kinjo, R. (2014). Isolation of a bis(oxazol-2-ylidene)-phenylborylene adduct and its reactivity as a boron-centered nucleophile. *Angew. Chem. Int. Ed.* 53, 9280–9283.
62. Goettel, J.T., and Braunschweig, H. (2019). Recent advances in boron-centered ligands and their transition metal complexes. *Coord. Chem. Rev.* 380, 184–200.
63. Braunschweig, H., Dewhurst, R.D., and Gessner, V.H. (2013). Transition metal borylene complexes. *Chem. Soc. Rev.* 42, 3197–3208.
64. See, for example: Bettinger, H.F. (2006). Phenylborylene: direct spectroscopic characterization in inert gas matrices. *J. Am. Chem. Soc.* 128, 2534–2535.
65. Krasowska, M., and Bettinger, H.F. (2016). Ring Enlargement of three-membered boron heterocycles upon reaction with organic π systems: implications for the trapping of borylenes. *Chem. Eur. J.* 22, 10661–10670.
66. Grigsby, W.J., and Power, P.P. (1996). Isolation and reduction of sterically encumbered arylboron dihalides: novel boranediyl insertion into C–C σ -bonds. *J. Am. Chem. Soc.* 118, 7981–7988.
67. Krasowska, M., Edelmann, M., and Bettinger, H.F. (2016). Electronically excited states of borylenes. *J. Phys. Chem. A* 120, 6332–6341.
68. Nakano, R., Jazzar, R., and Bertrand, G. (2018). A crystalline monosubstituted carbene. *Nat. Chem.* 10, 1196–1200.
69. Note that silicon analogues, namely silyliumylidene ions, have been investigated: Powley, S.L., and Inoue, S. (2019). NHC-Stabilized Silyliumylidene Ions. *Chem. Rec.* 19, 2179–2188.
70. Wang, Z., Herraiz, A.G., del Hoyo, A.M., and Suero, M.G. (2018). Generating carbyne equivalents with photoredox catalysis. *Nature* 554, 86–91.