


# Hyperconjugative Antiaromaticity Activates 4H-Pyrazoles as Inverse-Electron-Demand Diels–Alder Dienes

Brian J. Levandowski,<sup>†,‡,§,ID</sup> Nile S. Abularrage,<sup>†,§,ID</sup> K. N. Houk,<sup>\*,‡,§,ID</sup> and Ronald T. Raines<sup>\*,†,ID</sup><sup>†</sup>Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States<sup>‡</sup>Department of Chemistry and Biochemistry, University of California, Los Angeles, California 90095, United States**S Supporting Information**


**ABSTRACT:** The Diels–Alder reactivity of 4,4-difluoro-3,5-diphenyl-4H-pyrazole was investigated experimentally and computationally with *endo*-bicyclo[6.1.0]non-4-yne. The computationally predicted rate enhancement from hyperconjugative antiaromaticity induced by fluorination of cyclopentadienes at the 5-position extends to five-membered heterocyclic dienes containing a saturated center. 4,4-Difluoro-4H-pyrazoles are new electron-deficient dienes with rapid reactivities toward strained alkynes.



Cyclopentadienes have been explored as a new class of bio-orthogonal dienes with rapid reactivities and robust biological stabilities.<sup>1</sup> Tetrachlorocyclopentadiene ketal (TCK) does not dimerize<sup>2</sup> and can be synthesized in one step from readily available starting materials. As an ambiphilic diene, TCK reacts with both electron-deficient and electron-rich dienophiles and is stable toward biological nucleophiles. The rate constant for the Diels–Alder reaction of TCK with *endo*-bicyclo[6.1.0]non-4-yne (BCN) at room temperature in methanol is  $0.26 \text{ M}^{-1} \text{ s}^{-1}$ . TCK cycloadditions are as fast as the most rapid azide–alkyne cycloadditions<sup>3</sup> but slower than the inverse-electron-demand Diels–Alder reactions of 3,6-diphenyl- and 3,6-dipyridyl-1,2,4,5-tetrazines with BCN, which have rate constants in MeOH at room temperature of 3.3 and  $44.8 \text{ M}^{-1} \text{ s}^{-1}$ , respectively.<sup>4</sup>

The rapid Diels–Alder reactivity of cyclopentadienes is the result of the minimal distortion of the cyclopentadiene ground-state geometry required to achieve the transition-state geometry.<sup>5</sup> The Diels–Alder reactivities of cyclopentadienes can be tuned by substitution at the 5-position.<sup>6</sup> Negative hyperconjugation at the saturated center of a cyclopentadiene enhances the Diels–Alder reactivity of the diene by producing antiaromatic  $4\pi$ -electron destabilization.<sup>7</sup> This destabilization, which is especially strong when a fluoro group is the substituent, attracted our attention to the 4,4-difluoro-4H-pyrazole scaffold. Hünig<sup>8</sup> and Adam<sup>9</sup> have studied the Diels–Alder reactions of 3,5-dimethyl- and 3,5-diaryl-substituted 4,4-dimethyl-4H-pyrazoles extensively and found that acid catalysis is needed to promote the reaction, even with highly strained and electron-rich dienophiles. As shown in **Scheme 1**, the Diels–Alder reaction of 4,4-dimethyl-3,5-dimethyl-4H-pyrazole with cyclopentadiene (Cp) proceeds only in the presence of an acid catalyst.<sup>8a</sup> Though Diels–Alder reactions of 4,4-difluoro-4H-

**Scheme 1. Diels–Alder Reaction of 4,4-Dimethyl-3,5-diphenyl-4H-pyrazole and Cyclopentadiene (Cp) with or without an Acid Catalyst<sup>9</sup>**



pyrazoles have yet to be studied, the destabilizing effects of hyperconjugative antiaromaticity invoked by the geminal fluoro groups suggest that 4,4-difluoro-4H-pyrazoles would be highly reactive as Diels–Alder dienes.


We have studied computationally and experimentally the Diels–Alder reactivity of 4,4-dimethyl-3,5-diphenyl-4H-pyrazole (DMP), 4,4-difluoro-3,5-diphenyl-4H-pyrazole (DFP), and 3,6-diphenyl-1,2,4,5-tetrazine (Tz) with BCN (**Scheme 2**). We find that fluorine substitution enhances the Diels–Alder reactivity of DFP, making it more reactive than Tz toward BCN.

We began by evaluating the aromaticity of the 4H-pyrazole scaffold by using nuclear independent chemical shift (NICS)<sup>10</sup> calculations. Our intent was to determine if the hyperconjugative aromaticity and antiaromaticity observed in cyclopentadienes extends to 4H-pyrazoles. NICS(0) calculations measure the magnetic shielding at the center of the ring and have been shown to correlate well with the Diels–Alder reactivities of

**Received:** September 21, 2019

**Published:** October 7, 2019

**Scheme 2. Structures of 4,4-Dimethyl-3,5-diphenyl-4*H*-pyrazole (DMP), 4,4-Difluoro-3,5-diphenyl-4*H*-pyrazole (DFP), 3,6-Diphenyl-1,2,4,5-tetrazine (Tz), and *endo*-Bicyclo[6.1.0]non-4-yne (BCN)**



heteroaromatic azadienes.<sup>11</sup> Negative NICS values are associated with a diatropic (aromatic) ring current, whereas positive NICS values are indicative of a paratropic (antiaromatic) ring current. The NICS(0) value of 0.5 ppm for the unsubstituted 4*H*-pyrazole suggests that the 4*H*-pyrazole scaffold is not aromatic (Table S1). Fluorination of the 4*H*-pyrazole leads to antiaromatic electron delocalization (NICS(0) = 6.2 ppm), whereas silylation induces aromatic electron delocalization (NICS(0) = -3.3 ppm).

The antiaromaticity of 4,4-difluoro-4*H*-pyrazoles led us to believe that they would be highly reactive as inverse-electron-demand Diels–Alder dienes. We computationally surveyed the Diels–Alder reactivities of DFP, DMP, and Tz toward BCN at the M06-2X/6-31G(d)//M06-2X/6-311++(d,p)-SMD(H<sub>2</sub>O) level of theory.<sup>12</sup> Gas-phase energies are listed in Table S2. Our calculations reveal that fluorine substitution significantly enhances the reactivity of the 4*H*-pyrazole scaffold. The reaction of DFP with BCN is predicted to be  $5.2 \times 10^5$  faster and 12.5 kcal/mol more exergonic than the cycloaddition of DMP with BCN (Figure 1). The increased reactivity and exergonicity of the DFP cycloaddition are consistent with DFP being antiaromatic. The lowest unoccupied molecular orbital (LUMO) energies of DFP and DMP were calculated to be -2.1 and -1.3 eV, respectively. Decreasing the LUMO results in more favorable frontier molecular orbital interactions and is an additional factor that contributes to the increased reactivity of DFP as an inverse-electron-demand diene.

We also compared the reactivity of the 4,4-difluoro-4*H*-pyrazole scaffold to the highly reactive tetrazine scaffold. We found that the 4,4-difluoro-4*H*-pyrazole and tetrazine scaffolds have similar predicted reactivities toward BCN, with the


computed activation free energies differing by only 0.1 kcal/mol in favor of DFP.

We synthesized DFP in one step and high yield (93%) from commercial starting materials by a route reported previously.<sup>13</sup> To evaluate how the reactivity of the 4,4-difluoro-4*H*-pyrazole scaffold compares to that of the tetrazine scaffold, we measured pseudo-first-order kinetics for the Diels–Alder reactions of DFP and Tz with BCN in 9:1 methanol/water at room temperature. A plot of the observed rate constants with respect to the BCN concentration is shown in Figure 2. The 4,4-difluoro-4*H*-pyrazole scaffold is more reactive than is the tetrazine scaffold toward BCN, with second-order rate constants of 5.2 and 3.2 M<sup>-1</sup> s<sup>-1</sup> for DFP and Tz, respectively. The rapid reactivity of DFP arises from the antiaromatic 4*π*-electron delocalization invoked by negative hyperconjugation of the fluoro substituents.<sup>7</sup> To provide confirmation for hyperconjugative antiaromaticity as the source of the high reactivity of DFP, we stirred DMP with BCN; we detected no reaction after 2 h by NMR spectroscopy.



To discern if hyperconjugative antiaromaticity compromises the stability of DFP in a biological context, we incubated DMP and DFP in phosphate-buffered saline (PBS) containing fetal bovine serum (FBS; 10% v/v) at 37 °C. After 8 h, 42% of the DFP and 98% of the DMP remained intact (Figure S1). Thus, hyperconjugative antiaromaticity also increases the reactivity of the diene toward biological nucleophiles but not to a substantial detriment.

The DFP–BCN cycloadduct loses N<sub>2</sub>(g) spontaneously to yield a 5,5-difluorocyclopentadiene product that we characterized by NMR spectroscopy and mass spectrometry. The geminal fluoro groups at the saturated cyclopentadiene center of the 5,5-difluorocyclopentadiene product induce hyperconjugative antiaromaticity.<sup>6</sup> To see if this hyperconjugative antiaromaticity compromises stability, we incubated the cycloadduct in PBS containing FBS (10% v/v) at 37 °C. After 8 h, no decomposition of the cycloadduct was apparent by UV–vis spectroscopy (Figure S1).

In conclusion, we have experimentally confirmed the computationally predicted rate-enhancing effects of hyperconjugative antiaromaticity on the Diels–Alder reactivity of five-membered cyclic dienes. The use of hyperconjugative antiaromaticity to tune the Diels–Alder reactivity of five-membered cyclic dienes and three-membered cyclic dieno-



**Figure 1.** Transition-state geometries and calculated Gibbs free energies of activation ( $\Delta G^\ddagger$ ) and reaction ( $\Delta G_{\text{rxn}}$ ) computed at the M06-2X/6-31G(d)//M06-2X/6-311++G(d,p)-SMD(H<sub>2</sub>O) level of theory for the Diels–Alder reactions of 4,4-dimethyl-3,5-diphenyl-4*H*-pyrazole (DMP), 4,4-difluoro-3,5-diphenyl-4*H*-pyrazole (DFP), and 3,6-diphenyl-1,2,4,5-tetrazine (Tz) toward BCN. The lengths (Å) of forming bonds are shown. Values of  $k_{\text{rel}}$  were calculated at 298 K with the Arrhenius equation.



**Figure 2.** Reaction of BCN (1, 5, or 10 mM) with DFP (0.1 mM) or Tz (0.1 mM) in 9:1 methanol/water. Values of  $k_{\text{obs}}$  are the mean  $\pm$  SEM for reactions performed in triplicate. Second-order rate constants as calculated from a linear fit of the data were  $k_2 = 5.2 \text{ M}^{-1} \text{ s}^{-1}$  for DFP ( $R^2 = 0.97$ ) and  $k_2 = 3.2 \text{ M}^{-1} \text{ s}^{-1}$  for Tz ( $R^2 = 0.99$ ).

philes<sup>14</sup> will be useful to click chemistry applications that require rapid and tunable kinetics.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the [ACS Publications website](#) at DOI: [10.1021/acs.orglett.9b03351](https://doi.org/10.1021/acs.orglett.9b03351).

Tables S1 and S2; Figure S1; computational methods and additional computational data, Cartesian coordinates, and energies; and synthetic methods and analytical data ([PDF](#))

## ■ AUTHOR INFORMATION

### Corresponding Authors

\*E-mail: [houk@chem.ucla.edu](mailto:houk@chem.ucla.edu).

\*E-mail: [rtraines@mit.edu](mailto:rtraines@mit.edu).

### ORCID

Brian J. Levandowski: [0000-0002-8139-9417](https://orcid.org/0000-0002-8139-9417)

Nile S. Abularrage: [0000-0002-3112-2591](https://orcid.org/0000-0002-3112-2591)

K. N. Houk: [0000-0002-8387-5261](https://orcid.org/0000-0002-8387-5261)

Ronald T. Raines: [0000-0001-7164-1719](https://orcid.org/0000-0001-7164-1719)

### Author Contributions

§B.J.L. and N.S.A. contributed equally to this work.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This research was supported by Grants R01 GM044783 (NIH) and CHE-1764328 (NSF). Computational resources were provided by the UCLA Institute for Digital Research and Education (IDRE) and the Extreme Science and Engineering Discovery Environment (XSEDE),<sup>15</sup> which is supported by Grant ACI-1548562 (NSF).

## ■ REFERENCES

- (1) Levandowski, B. J.; Gamache, R. F.; Murphy, J. M.; Houk, K. N. Readily Accessible Ambiphilic Cyclopentadienes for Bioorthogonal Labeling. *J. Am. Chem. Soc.* **2018**, *140*, 6426–6431.
- (2) (a) Eibler, E.; Burgemeister, T.; Höcht, P.; Prantl, B.; Roßmaier, H.; Schuhbauer, H. M.; Wiest, H.; Sauer, J. Polyhalogenated
- Cyclopentadienes in [4 + 2] Cycloadditions: Preparative Aspects. *Liebigs Ann.* **1997**, *1997*, 2451–2469. (b) Eibler, E.; Höcht, P.; Prantl, B.; Roßmaier, H.; Schuhbauer, H. M.; Wiest, H.; Sauer, J. Transitions of Electron Demand in Pericyclic Reactions: Normal, Neutral, and Inverse Diels–Alder Reactions of Polyhalogenated Cyclopentadienes. *Liebigs Ann.* **1997**, *1997*, 2471–2484.
- (3) Dommerholt, J.; Rutjes, F. P. J. T.; van Delft, F. L. Strain-Promoted 1,3-Dipolar Cycloaddition of Cycloalkynes and Organic Azides. *Top. Curr. Chem.* **2016**, *374*, 16.
- (4) Chen, W.; Wang, D.; Dai, C.; Hamelberg, D.; Wang, B. Clicking 1,2,4,5-Tetrazine and Cyclooctynes with Tunable Reaction Rates. *Chem. Commun.* **2012**, *48*, 1736–1738.
- (5) Levandowski, B. J.; Houk, K. N. Theoretical Analysis of Reactivity Patterns in Diels–Alder Reactions of Cyclopentadiene, Cyclohexadiene, and Cycloheptadiene with Symmetrical and Unsymmetrical Dienophiles. *J. Org. Chem.* **2015**, *80*, 3530–3537.
- (6) (a) Nyulászi, L.; Schleyer, P. v. R. Hyperconjugative  $\pi$ -Aromaticity: How To Make Cyclopentadiene Aromatic. *J. Am. Chem. Soc.* **1999**, *121*, 6872–6875. (b) Fernández, I.; Wu, J. I.; von Ragüé Schleyer, P. Substituent Effects on “Hyperconjugative” Aromaticity and Antiaromaticity in Planar Cyclopolyenes. *Org. Lett.* **2013**, *15*, 2990–2993.
- (7) (a) Levandowski, B. J.; Zou, L.; Houk, K. N. Schleyer hyperconjugative aromaticity and Diels–Alder reactivity of 5-substituted cyclopentadienes. *J. Comput. Chem.* **2016**, *37*, 117–123. (b) Levandowski, B. J.; Zou, L.; Houk, K. N. Hyperconjugative Aromaticity and Antiaromaticity Control the Reactivities and  $\pi$ -Facial Stereoselectivities of 5-Substituted Cyclopentadiene Diels–Alder Cycloadditions. *J. Org. Chem.* **2018**, *83*, 14658–14666.
- (8) (a) Beck, K.; Höhn, A.; Hünig, S.; Prokschy, F. Azobrücke aus Azinen, I. Isopyrazole als elektronenarme Diene zur Synthese von 2,3-Diazabicyclo[2.2.1]heptenen. *Chem. Ber.* **1984**, *117*, 517–533. (b) Beck, K.; Hünig, S.; Klärner, F.-G.; Kraft, P.; Artschwager-Perl, U. Azobrücke aus Azinen, VII. Diels–Alder-Reaktionen mit inversem Elektronenbedarf zwischen Isopyrazolen und Cycloalkenen sowie Cycloalkadienen.—Ein Vergleich von Säure-Katalyse und Beschleunigung durch Druck. *Chem. Ber.* **1987**, *120*, 2041–2051.
- (9) (a) Adam, W.; Harrer, H. M.; Nau, W. M.; Peters, K. Electronic Substituent Effects on the Acid-Catalyzed [4<sup>+</sup> + 2] Cycloaddition of Isopyrazoles with Cyclopentadiene and the Photochemical and Thermal Denitrogenation of the Resulting 1,4-Diaryl-7,7-dimethyl-2,3-diazabicyclo[2.2.1]hept-2-ene Azoalkanes to Bicyclo[2.1.0]-pentanes. *J. Org. Chem.* **1994**, *59*, 3786–3797. (b) Adam, W.; Ammon, H.; Nau, W. M.; Peters, K. 4-Halo-4H-pyrazoles: Cycloaddition with Cyclopentadiene to Azoalkanes of the 2,3-Diazabicyclo[2.2.1]hept-2-ene Type versus Electrophilic Addition with Cyclopentene. *J. Org. Chem.* **1994**, *59*, 7067–7071.

(10) (a) Chen, Z.; Wannere, C. S.; Corminboeuf, C.; Puchta, R.; Schleyer, P. v. R. Nucleus-Independent Chemical Shifts (NICS) as an Aromaticity Criterion. *Chem. Rev.* **2005**, *105*, 3842–3888. (b) Fallah-Bagher-Shaidaei, H.; Wannere, C. S.; Corminboeuf, C.; Puchta, R.; Schleyer, P. v. R. Which NICS Aromaticity Index for Planar  $\pi$  Rings Is Best? *Org. Lett.* **2006**, *8*, 863–866. (c) Gershoni-Poranne, R.; Stanger, A. Magnetic Criteria of Aromaticity. *Chem. Soc. Rev.* **2015**, *44*, 6597–6615.

(11) Yang, Y.-F.; Liang, Y.; Liu, F.; Houk, K. N. Diels–Alder Reactivities of Benzene, Pyridine, and Di-, Tri-, and Tetrazines: The Roles of Geometrical Distortions and Orbital Interactions. *J. Am. Chem. Soc.* **2016**, *138*, 1660–1667.

(12) (a) Zhao, Y.; Truhlar, D. G. The M06 Suite of Density Functionals for Main Group Thermochemistry, Thermochemical Kinetics, Noncovalent Interactions, Excited States, and Transition Elements: Two New Functionals and Systematic Testing of Four M06-class Functionals and 12 Other Functionals. *Theor. Chem. Acc.* **2008**, *120*, 215–241. (b) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. Universal Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions. *J. Phys. Chem. B* **2009**, *113*, 6378–6396.

(13) (a) Walton, L.; Duplain, H. R.; Knapp, A. L.; Eidell, C. K.; Bacsa, J.; Stephens, C. E. Synthesis and Charge Density Determination of 4,4-Difluoro-3,5-Diphenyl-4*H*-pyrazole and a Hydrate Derivative. *J. Fluorine Chem.* **2015**, *173*, 12–17. (b) Breen, J. R.; Sandford, G.; Patel, B.; Fray, J. Synthesis of 4,4-Difluoro-1*H*-pyrazole Derivatives. *Synlett* **2014**, *26*, 51–54.

(14) (a) Levandowski, B. J.; Houk, K. N. Hyperconjugative, Secondary Orbital, Electrostatic, and Steric Effects on the Reactivities and Endo and Exo Stereoselectivities of Cyclopropene Diels–Alder Reactions. *J. Am. Chem. Soc.* **2016**, *138*, 16731–16736. (b) Rivasco, J. M. J. M.; Monteiro, C. M.; Trindade, A. F. Cyclopropenes: A New Tool for the Study of Biological Systems. *Org. Chem. Front.* **2017**, *4*, 1167–1198.

(15) Towns, J.; Cockerill, T.; Dahan, M.; Foster, I.; Gaither, K.; Grimshaw, A.; Hazlewood, V.; Lathrop, S.; Lifka, D.; Peterson, G. D.; Roskies, R.; Scott, J. R.; Wilkins-Diehr, N. XSEDE: Accelerating Scientific Discovery. *Comput. Sci. Eng.* **2014**, *16*, 62–74.