

LETTER

The symmetric representation of the generalized rigid body equations and symplectic reduction

To cite this article: Tomoki Ohsawa 2019 *J. Phys. A: Math. Theor.* **52** 36LT01

View the [article online](#) for updates and enhancements.



IOP | **ebooks**TM

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Letter

The symmetric representation of the generalized rigid body equations and symplectic reduction

Tomoki Ohsawa 

Department of Mathematical Sciences, The University of Texas at Dallas, 800 W Campbell Rd, Richardson, TX 75080-3021, United States of America

E-mail: tomoki@utdallas.edu

Received 6 November 2018, revised 1 March 2019

Accepted for publication 10 May 2019

Published 13 August 2019



CrossMark

Abstract

We show that a symplectic reduction of the symmetric representation of the generalized n -dimensional rigid body equations yields the n -dimensional Euler equation. This result provides an alternative to the more elaborate relationship between these equations established by Bloch, Crouch, Marsden, and Ratiu (Bloch *et al* 2002 *Nonlinearity* **15** 1309–41). Specifically, we exploit the inherent $\text{Sp}(2n, \mathbb{R})$ -symmetry in the symmetric representation to present its relationship with the Euler equation via symplectic reduction facilitated by the dual pair recently developed by Skerritt and Vizman (Skerritt and Vizman 2019 *J. Geom. Mech.* **11** 255–75).

Keywords: rigid body dynamics, symplectic reduction, dual pair

1. Introduction

1.1. Euler's equation for rigid body dynamics

The configuration space for the rotational dynamics of a rigid body in \mathbb{R}^3 about its center of mass is the set of all three-dimensional (3D) rotations

$$\text{SO}(3) := \{Q \in \mathbf{M}_3(\mathbb{R}) \mid Q^T Q = I, \det Q = 1\},$$

where $\mathbf{M}_n(\mathbb{R})$ is the vector space of real $n \times n$ matrices. Hence one can describe the dynamics as a Hamiltonian system on the cotangent bundle (or the phase space) $T^*\text{SO}(3)$. However, the presence of the $\text{SO}(3)$ -symmetry helps us reduce the system to the dual $\mathfrak{so}(3)^* \cong (\mathbb{R}^3)^*$ of the Lie algebra $\mathfrak{so}(3) \cong \mathbb{R}^3$ of $\text{SO}(3)$; one may think of $\mathfrak{so}(3)$ and $\mathfrak{so}(3)^*$ as the space of all possible values of the body angular velocity and momentum, respectively, seen in the frame attached to the body. This yields Euler's equation for the body angular momentum Π in $\mathfrak{so}(3)^*$ (identified with \mathbb{R}^3):

$$\dot{\Pi} = \Pi \times \mathbb{I}^{-1} \Pi,$$

where $\mathbb{I} := \text{diag}(I_1, I_2, I_3)$ is the inertia tensor with respect to the principal axes of the body, and $\Omega := \mathbb{I}^{-1} \Pi$ is the body angular velocity in $\mathfrak{so}(3)$ (again identified with \mathbb{R}^3).

1.2. Generalized rigid body equations

Manakov [8] and Ratiu [13] generalized the 3D rigid body equations to n dimensions as follows: let

$$\text{SO}(n) := \{Q \in \text{M}_n(\mathbb{R}) \mid Q^T Q = I, \det Q = 1\}$$

and $\mathfrak{so}(n)$ be its Lie algebra, and equip $\mathfrak{so}(n)$ with the inner product $\langle \cdot, \cdot \rangle: \mathfrak{so}(n) \times \mathfrak{so}(n) \rightarrow \mathbb{R}$ defined as

$$\langle A, B \rangle := \frac{1}{2} \text{tr}(A^T B). \quad (1)$$

So we may identify the dual $\mathfrak{so}(n)^*$ with $\mathfrak{so}(n)$. Under this identification, we define the inertia operator $\mathcal{I}: \mathfrak{so}(n) \rightarrow \mathfrak{so}(n)^* \cong \mathfrak{so}(n)$ of the generalized n -dimensional rigid body as

$$\mathcal{I}(\Omega) := \Lambda \Omega + \Omega \Lambda, \quad (2)$$

where $\Lambda := \text{diag}(\lambda_1, \dots, \lambda_n)$ with $\lambda_i + \lambda_j > 0$ for all $i \neq j$. This is a generalization of the inertia tensor \mathbb{I} defined above. In fact, setting $I_1 = \lambda_2 + \lambda_3$, $I_2 = \lambda_3 + \lambda_1$, and $I_3 = \lambda_1 + \lambda_2$ for $n = 3$ yields

$$\Pi = \mathcal{I}(\Omega) = \begin{bmatrix} 0 & -I_3 \Omega_3 & I_2 \Omega_2 \\ I_3 \Omega_3 & 0 & -I_1 \Omega_1 \\ -I_2 \Omega_2 & I_1 \Omega_1 & 0 \end{bmatrix} \leftrightarrow \begin{bmatrix} I_1 \Omega_1 \\ I_2 \Omega_2 \\ I_3 \Omega_3 \end{bmatrix},$$

which is the body angular momentum in $\mathfrak{so}(3)^* \cong \mathbb{R}^3$. Let $Q \in \text{SO}(n)$ be the rotational configuration of the generalized rigid body, $\Omega := Q^{-1} \dot{Q} \in \mathfrak{so}(n)$ be the body angular velocity, and $\Pi := \mathcal{I}(\Omega) \in \mathfrak{so}(n)^* \cong \mathfrak{so}(n)$ be the body angular momentum. Then the generalized rigid body or Euler–Poisson equations on $\text{SO}(n) \times \mathfrak{so}(n)^*$ are given by

$$\dot{Q} = Q \Omega, \quad (3a)$$

$$\dot{\Pi} = [\Pi, \Omega], \quad (3b)$$

where $[\cdot, \cdot]: \mathfrak{so}(n) \times \mathfrak{so}(n) \rightarrow \mathfrak{so}(n)$ is the standard commutator. The n -dimensional Euler equation (3b) is the Hamiltonian system on a coadjoint orbit in $\mathfrak{so}(n)^*$ with the Hamiltonian (see Ratiu [13, theorem 3.1]):

$$h(\Pi) := \frac{1}{2} \langle \Pi, \mathcal{I}^{-1}(\Pi) \rangle. \quad (4)$$

1.3. Symmetric representation

The so-called *symmetric representation* of the generalized rigid body equations was originally discovered by Bloch and Crouch [3] as a necessary condition of optimality for the following optimal control problem: let $T > 0$ be fixed, and consider

$$\min_U \int_0^T \frac{1}{2} \langle \mathcal{I}(U), U \rangle dt \quad \text{subject to} \quad \dot{Q} = QU \text{ and } Q(0) \in \text{SO}(n),$$

where \mathcal{I} is the inertia operator (2) and $U: [0, T] \rightarrow \mathfrak{so}(n)$. Formally, the configuration space is $\mathbf{M}_n(\mathbb{R})$, and thus $Q: [0, T] \rightarrow \mathbf{M}_n(\mathbb{R})$. However, we impose the initial condition $Q(0) \in \text{SO}(n)$ so that $Q(t) \in \text{SO}(n)$ for any $t \in [0, T]$ because of the constraint $Q(t)^{-1}\dot{Q}(t) = U(t) \in \mathfrak{so}(n)$. This is a special case of the so-called *embedded optimal control problem* considered in Bloch *et al* [6].

The control Hamiltonian $H_c: T^*\mathbf{M}_n(\mathbb{R}) \times \mathfrak{so}(n) \rightarrow \mathbb{R}$ is then defined as

$$H_c(Q, P, U) := P \cdot (QU) - \frac{1}{2} \langle \mathcal{I}(U), U \rangle = \text{tr}(P^T QU) - \frac{1}{2} \langle \mathcal{I}(U), U \rangle,$$

where the ‘ \cdot ’ above stands for the natural dual pairing

$$T_Q^*\mathbf{M}_n(\mathbb{R}) \times T_Q\mathbf{M}_n(\mathbb{R}) \rightarrow \mathbb{R}; \quad (P, \dot{Q}) \mapsto \text{tr}(P^T \dot{Q}) =: P \cdot \dot{Q}.$$

By the Pontryagin maximum principle [12], the optimal control U^* necessarily maximizes the Hamiltonian, i.e. for any $\delta U \in \mathfrak{so}(n)$,

$$\frac{d}{ds} H_c(Q, P, U^* + s\delta U) \Big|_{s=0} = 0.$$

However,

$$\begin{aligned} \frac{d}{ds} H_c(Q, P, U^* + s\delta U) \Big|_{s=0} &= \text{tr}(P^T Q \delta U) - \langle \mathcal{I}(U^*), \delta U \rangle \\ &= \frac{1}{2} \text{tr}((Q^T P - P^T Q)^T \delta U) - \langle \mathcal{I}(U^*), \delta U \rangle \\ &= \langle Q^T P - P^T Q - \mathcal{I}(U^*), \delta U \rangle. \end{aligned}$$

Note that $Q^T P - P^T Q - \mathcal{I}(U^*) \in \mathfrak{so}(n)$. Since $\delta U \in \mathfrak{so}(n)$ is arbitrary, we have $Q^T P - P^T Q = \mathcal{I}(U^*)$, and thus obtain the optimal control U^* as follows:

$$U^*(Q, P) = \mathcal{I}^{-1}(Q^T P - P^T Q).$$

In what follows, we write $\Omega := U^*(Q, P)$ for short; this quantity in fact coincides with the angular velocity Ω defined in section 1.2 as we shall see below. As a result, the (optimal) Hamiltonian $H: T^*\mathbf{M}_n(\mathbb{R}) \rightarrow \mathbb{R}$ is given by

$$H(Q, P) := H_c(Q, P, U^*(Q, P)) = \frac{1}{2} \langle Q^T P - P^T Q, \mathcal{I}^{-1}(Q^T P - P^T Q) \rangle, \quad (5)$$

whereas the standard symplectic form ω on $T^*\mathbf{M}_n(\mathbb{R})$ is written as

$$\omega((\dot{Q}_1, \dot{P}_1), (\dot{Q}_2, \dot{P}_2)) = \text{tr}(\dot{Q}_1^T \dot{P}_2 - \dot{P}_1^T \dot{Q}_2). \quad (6)$$

The optimal solution is necessarily an integral curve of the Hamiltonian vector field X_H on $T^*\mathbf{M}_n(\mathbb{R})$ defined by Hamilton's equation $\mathbf{i}_{X_H}\omega = \mathbf{d}H$, or in coordinates,

$$\dot{Q} = Q\Omega, \quad \dot{P} = P\Omega. \quad (7)$$

These equations are called the *symmetric representation* of the generalized rigid body equations (3). In fact, if we set

$$\Pi(t) := \mathcal{I}(\Omega(t)) = Q(t)^T P(t) - P(t)^T Q(t)$$

for $t \in [0, T]$, then $\Pi: [0, T] \rightarrow \mathfrak{so}(n)^* \cong \mathfrak{so}(n)$ satisfies the n -dimensional Euler equation (3b); see also Bloch *et al* [5, 6] for its generalization to other matrix Lie groups.

Although it is a straightforward calculation to show that the generalized rigid body equations (3) follow from the symmetric representation (7), the question remains as to how these two Hamiltonian systems are related to each other from the symplectic-geometric point of view. Bloch *et al* [4] gave one such relationship: specifically, they constructed symplectic submanifolds $S \subset \mathrm{SO}(n) \times \mathrm{SO}(n)$ of $T^* \mathrm{M}_n(\mathbb{R}) \cong \mathrm{M}_n(\mathbb{R}) \times \mathrm{M}_n(\mathbb{R})$ and S_M of $T^* \mathrm{SO}(n) \cong \mathrm{SO}(n) \times \mathfrak{so}(n)^*$ as well as a diffeomorphism between S and S_M in a rather elaborate manner to establish an equivalence between the symmetric representation (7) and the Euler–Poisson equations (3).

1.4. Main result

We present an alternative connection between the symmetric representation (7) and the Euler equation (3b) by showing that the two are related via symplectic reduction. Although this is not an equivalence, this connection exploits an inherent symmetry of the symmetric representation that seems to have been overlooked, and provides a geometrically natural and appealing alternative to the result of Bloch *et al* [4]. More specifically, we show the following:

Theorem. *The symmetric representation (7) of the n -dimensional generalized rigid body equation possesses an $\mathrm{Sp}(2n, \mathbb{R})$ -symmetry, and the Marsden–Weinstein reduction [10] of (7) restricted to an open subset of $T^* \mathrm{M}_n(\mathbb{R})$ by the symmetry at a certain level set of the associated momentum map yields the n -dimensional Euler equation (3b) in a coadjoint orbit in $\mathfrak{so}(n)^*$.*

2. Symmetry and conservation law in the symmetric representation

2.1. $\mathrm{Sp}(2n, \mathbb{R})$ -symmetry of the symmetric representation

Let us first identify the cotangent bundle $T^* \mathrm{M}_n(\mathbb{R})$ with the vector space of real $2n \times n$ matrices:

$$T^* \mathrm{M}_n(\mathbb{R}) \cong \mathrm{M}_{2n \times n}(\mathbb{R}) = \left\{ Z = \begin{bmatrix} Q \\ P \end{bmatrix} \mid Q, P \in \mathrm{M}_n(\mathbb{R}) \right\}.$$

Then we may write the symplectic form (6) in a more succinct form: for any $Z \in \mathrm{M}_{2n \times n}(\mathbb{R})$ and any $X, Y \in T_Z \mathrm{M}_{2n \times n}(\mathbb{R})$,

$$\omega(X, Y) = \mathrm{tr}(X^T \mathbb{J} Y) \quad \text{with} \quad \mathbb{J} := \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix} \quad (8)$$

where I is the $n \times n$ identity matrix. Notice also that we may rewrite the Hamiltonian $H: \mathrm{M}_{2n \times n}(\mathbb{R}) \rightarrow \mathbb{R}$ defined in (5) in the following more concise form:

$$H(Z) = \frac{1}{2} \langle Z^T \mathbb{J} Z, \mathcal{I}^{-1}(Z^T \mathbb{J} Z) \rangle. \quad (9)$$

Let $\mathrm{Sp}(2n, \mathbb{R})$ be the symplectic group

$$\mathrm{Sp}(2n, \mathbb{R}) := \{S \in \mathrm{M}_{2n}(\mathbb{R}) \mid S^T \mathbb{J} S = \mathbb{J}\},$$

and consider the $\mathrm{Sp}(2n, \mathbb{R})$ -action on $\mathrm{M}_{2n \times n}(\mathbb{R})$ by left multiplication, i.e.

$$\Phi: \mathrm{Sp}(2n, \mathbb{R}) \times \mathrm{M}_{2n \times n}(\mathbb{R}) \rightarrow \mathrm{M}_{2n \times n}(\mathbb{R}); \quad (S, Z) \mapsto SZ =: \Phi_S(Z). \quad (10)$$

Then one easily sees that this action is symplectic as well as that the Hamiltonian is invariant under the action, i.e. $\Phi_S^* \omega = \omega$ and $H \circ \Phi_S = H$ for any $S \in \mathrm{Sp}(2n, \mathbb{R})$.

2.2. $\mathrm{Sp}(2n, \mathbb{R})$ -momentum map

Let us find the momentum map associated with the above action Φ . First notice that the symplectic form (8) is written as $\omega = -\mathbf{d}\theta$ with the one-form θ on $\mathrm{M}_{2n \times n}(\mathbb{R})$ defined as follows:

for any $Z = \begin{bmatrix} Q \\ P \end{bmatrix} \in \mathrm{M}_{2n \times n}(\mathbb{R})$ and any $\dot{Z} = \begin{bmatrix} \dot{Q} \\ \dot{P} \end{bmatrix} \in T_Z \mathrm{M}_{2n \times n}(\mathbb{R})$,

$$\theta(Z) \cdot \dot{Z} := -\frac{1}{2} \mathrm{tr}(Z^T \mathbb{J} \dot{Z}) = \frac{1}{2}(P^T \dot{Q} - Q^T \dot{P}).$$

It is clear that θ is invariant under the $\mathrm{Sp}(2n, \mathbb{R})$ -action, i.e. $\Phi_S^* \theta = \theta$ for any $S \in \mathrm{Sp}(2n, \mathbb{R})$.

Let $\mathfrak{sp}(2n, \mathbb{R})$ be the Lie algebra of $\mathrm{Sp}(2n, \mathbb{R})$, i.e.

$$\mathfrak{sp}(2n, \mathbb{R}) = \{\xi \in \mathrm{M}_{2n}(\mathbb{R}) \mid \xi^T \mathbb{J} + \mathbb{J} \xi = 0\}.$$

We equip $\mathfrak{sp}(2n, \mathbb{R})$ with the inner product $\langle \cdot, \cdot \rangle: \mathfrak{sp}(2n, \mathbb{R}) \times \mathfrak{sp}(2n, \mathbb{R}) \rightarrow \mathbb{R}$ defined as

$$\langle \xi, \eta \rangle := \frac{1}{2} \mathrm{tr}(\xi^T \eta)$$

just as in (1), and thus we may identify the dual $\mathfrak{sp}(2n, \mathbb{R})^*$ with $\mathfrak{sp}(2n, \mathbb{R})$. The infinitesimal generator of the above action Φ is, for any $\xi \in \mathfrak{sp}(2n, \mathbb{R})$,

$$\xi_{\mathrm{M}_{2n \times n}(\mathbb{R})}(Z) := \left. \frac{d}{ds} \Phi_{\exp(s\xi)}(Z) \right|_{s=0} = \xi Z.$$

Since $T^* \mathrm{M}_n(\mathbb{R}) \cong \mathrm{M}_{2n \times n}(\mathbb{R})$ is an exact symplectic manifold with $\omega = -\mathbf{d}\theta$ and Φ leaves θ invariant, the associated momentum map $\mathbf{J}: \mathrm{M}_{2n \times n}(\mathbb{R}) \rightarrow \mathfrak{sp}(2n, \mathbb{R})^* \cong \mathfrak{sp}(2n, \mathbb{R})$ satisfies the following (see, e.g. Abraham and Marsden [1, theorem 4.2.10 on p 282]): for any $\xi \in \mathfrak{sp}(2n, \mathbb{R})$,

$$\begin{aligned} \langle \mathbf{J}(Z), \xi \rangle &= \theta(Z) \cdot \xi_{\mathrm{M}_{2n \times n}(\mathbb{R})}(Z) \\ &= -\frac{1}{2} \mathrm{tr}(Z^T \mathbb{J} \xi Z) \\ &= \frac{1}{2} \mathrm{tr}((\mathbb{J} Z Z^T)^T \xi) \\ &= \langle \mathbb{J} Z Z^T, \xi \rangle, \end{aligned}$$

and so we obtain

$$\mathbf{J}(Z) = \mathbb{J} Z Z^T = \begin{bmatrix} P Q^T & P P^T \\ -Q Q^T & -Q P^T \end{bmatrix}. \quad (11)$$

This is the special case with $m = n$ of Skerritt and Vizman [15, proposition 4.1]. It is also easy to see that \mathbf{J} is equivariant: for any $S \in \mathrm{Sp}(2n, \mathbb{R})$,

$$\mathbf{J} \circ \Phi_S = \text{Ad}_{S^{-1}}^* \circ \mathbf{J}.$$

By Noether's theorem (see, e.g. Marsden and Ratiu [9, theorem 11.4.1 on p 372]), \mathbf{J} is a conserved quantity of the symmetric representation (7) due to the $\text{Sp}(2n, \mathbb{R})$ -symmetry. That each block of this matrix is a conserved quantity is also pointed out by Bloch *et al* [4] via direct computations.

3. Symplectic reduction of the symmetric representation

3.1. Symplectic reduction

Let P be a symplectic manifold with symplectic form ω , and suppose in addition that there is a symplectic action of a Lie group G on P , \mathfrak{g}^* be the dual of the Lie algebra \mathfrak{g} of G , and $\mathbf{J}: P \rightarrow \mathfrak{g}^*$ be the momentum map associated with the action. The Marsden–Weinstein reduction [10] (see also [11, sections 1.1 and 1.2]) states that, if either (i) the G -action on P is free and proper, or (ii) $\mu \in \mathfrak{g}^*$ is a regular value of \mathbf{J} and the action of the isotropy group

$$G_\mu := \{g \in G \mid \text{Ad}_{g^{-1}}^* \mu = \mu\}$$

on the level set $\mathbf{J}^{-1}(\mu)$ is free and proper, then the quotient space $\mathbf{J}^{-1}(\mu)/G_\mu$ is also a symplectic manifold with symplectic structure $\bar{\omega}_\mu$ that is naturally induced by ω and the geometric setting; see below for more details. Now, given a Hamiltonian $H: P \rightarrow \mathbb{R}$, one may define the Hamiltonian vector field X_H on P by setting $\mathbf{i}_{X_H} \omega = dH$. If H is invariant under the G -action, it gives rise to the reduced Hamiltonian \bar{H}_μ on $\mathbf{J}^{-1}(\mu)/G_\mu$, and then the Hamiltonian dynamics in P is reduced to the Hamiltonian dynamics in $\mathbf{J}^{-1}(\mu)/G_\mu$ defined in terms of \bar{H}_μ and $\bar{\omega}_\mu$. In other words, one can reduce a Hamiltonian system with a Lie-group symmetry to a lower-dimensional Hamiltonian system.

3.2. Some technical issues of symplectic reduction

We would like to perform the Marsden–Weinstein reduction of the symmetric representation (7); here we have $P = T^* \mathbf{M}_n(\mathbb{R}) \cong \mathbf{M}_{2n \times n}(\mathbb{R})$, $G = \text{Sp}(2n, \mathbb{R})$ with the action Φ defined in (10), and the momentum map \mathbf{J} from (11). However, condition (i) clearly does not hold because Φ is not a free action on $\mathbf{M}_{2n \times n}(\mathbb{R})$, and so one either needs to remedy this or check (ii). Otherwise, the quotient $\mathbf{J}^{-1}(\mu)/\text{Sp}(2n, \mathbb{R})_\mu$ may not be a manifold. The other issue is how to characterize the quotient space $\mathbf{J}^{-1}(\mu)/\text{Sp}(2n, \mathbb{R})_\mu$ explicitly in order to describe the symplectic structure and the reduced dynamics there in an explicit manner.

Fortunately, the recent work by Skerritt and Vizman [15] provides a geometric setting that is tailor-made for circumventing these issues. More specifically, we consider the following pair of momentum maps defined on $\mathbf{M}_{2n \times n}(\mathbb{R})$:

$$\mathfrak{sp}(2n, \mathbb{R})^* \xleftarrow{\mathbf{J}} \mathbf{M}_{2n \times n}(\mathbb{R}) \xrightarrow{\mathbf{M}} \mathfrak{o}(n)^*,$$

where \mathbf{J} is defined above in (11) and \mathbf{M} is the momentum map associated with the action of the orthogonal group $\text{O}(n)$ on $\mathbf{M}_{2n \times n}(\mathbb{R})$ to be described below; $\mathfrak{o}(n)$ is the Lie algebra of $\text{O}(n)$. What they show is that, by considering an open subset \mathcal{Z} of $\mathbf{M}_{2n \times n}(\mathbb{R})$ and restricting the actions and the momentum maps there, one may identify the Marsden–Weinstein quotient $\mathbf{J}^{-1}(\mu)/\text{Sp}(2n, \mathbb{R})_\mu$ with a coadjoint orbit in $\mathfrak{o}(n)^*$. We note that their result is slightly more

general than this: they have the result with $M_{2n \times m}(\mathbb{R})$ and $O(m)$ with $n, m \in \mathbb{N}$ instead, and so our setting is the special case of theirs with $m = n$.

3.3. $O(n)$ -action and momentum map

Consider the action of the orthogonal group $O(n)$ on $M_{2n \times n}(\mathbb{R})$ defined by right multiplication, i.e.

$$\Psi: O(n) \times M_{2n \times n}(\mathbb{R}) \rightarrow M_{2n \times n}(\mathbb{R}); \quad (R, Z) \mapsto ZR = \Psi_R(Z).$$

It is a straightforward calculation to see that Ψ leaves the one-form θ invariant and hence is a symplectic action with respect to the symplectic form ω defined in (8), i.e. $\Psi_R^* \theta = \theta$ and hence $\Psi_R^* \omega = \omega$ for any $R \in O(n)$.

Since $\mathfrak{o}(n) = \mathfrak{so}(n)$, we identify the dual $\mathfrak{o}(n)^*$ with $\mathfrak{o}(n)$ via the inner product (1). Then, following a similar calculation as the one for \mathbf{J} from above (see also Skerritt and Vizman [15, proposition 4.1]), we obtain the associated momentum map $\mathbf{M}: M_{2n \times n}(\mathbb{R}) \rightarrow \mathfrak{o}(n)^* \cong \mathfrak{o}(n)$ as follows:

$$\mathbf{M}(Z) = Z^T \mathbb{J} Z = Q^T P - P^T Q. \quad (12)$$

Again, it is a straightforward calculation to see that \mathbf{M} is equivariant, i.e. for any $R \in O(n)$,

$$\mathbf{M} \circ \Psi_R = \text{Ad}_R^* \circ \mathbf{M}.$$

3.4. Symplectic reduction and dual pair

Following Skerritt and Vizman [15], let us consider the subset of $M_{2n \times n}(\mathbb{R})$ that is consisting of the full-rank elements, i.e.

$$\mathcal{Z} := \{Z \in M_{2n \times n}(\mathbb{R}) \mid \text{rank } Z = n\}. \quad (13)$$

As shown in [15], \mathcal{Z} is an open subset of $M_{2n \times n}(\mathbb{R})$, and the actions Φ and Ψ preserve \mathcal{Z} . Hence we may restrict the symplectic form ω and the momentum maps \mathbf{J} and \mathbf{M} to \mathcal{Z} ; we denote these restrictions by the same symbols for simplicity of notation:

$$\mathfrak{sp}(2n, \mathbb{R})^* \xleftarrow{\mathbf{J}} \mathcal{Z} \xrightarrow{\mathbf{M}} \mathfrak{o}(n)^*.$$

Skerritt and Vizman [15, proposition 4.2] proved that Φ and Ψ define *mutually transitive actions* on \mathcal{Z} in the following sense: (i) The $\text{Sp}(2n, \mathbb{R})$ -action Φ and the $O(n)$ -action Ψ commute; (ii) Φ and Ψ are symplectic actions; (iii) the momentum maps \mathbf{J} and \mathbf{M} are equivariant; (iv) each level set of \mathbf{J} is an $O(n)$ -orbit, and each level set of \mathbf{M} is an $\text{Sp}(2n, \mathbb{R})$ -orbit.

The mutual transitivity has the following important consequence (see also Balleier and Wurzbacher [2, theorem 2.9 (iii)] and Skerritt [14, proposition 3.5]): let $Z_0 \in \mathcal{Z}$ and $\mu_0 := \mathbf{J}(Z_0)$ and $\Pi_0 := \mathbf{M}(Z_0)$. Then one can identify the Marsden–Weinstein quotient

$$\overline{\mathcal{Z}}_{\mu_0} := \mathbf{J}^{-1}(\mu_0) / \text{Sp}(2n, \mathbb{R})_{\mu_0}$$

with the coadjoint orbit \mathcal{O}_{Π_0} through Π_0 in $\mathfrak{o}(n)^*$.

Remark 1. We do not have to check that the condition (mentioned in section 3.2) that the $\text{Sp}(n, \mathbb{R})$ -action Φ or the $\text{Sp}(n, \mathbb{R})_{\mu_0}$ action on $\mathbf{J}^{-1}(\mu_0)$ is free and proper. In fact, the smooth structure on the reduced space $\overline{\mathcal{Z}}_{\mu_0}$ is induced by that of the coadjoint orbit \mathcal{O}_{Π_0} . See the proof of proposition 2.8 in [15] for details.

More specifically, let $i_{\mu_0}: \mathbf{J}^{-1}(\mu_0) \hookrightarrow \mathcal{Z}$ be the inclusion and $\pi_{\mu_0}: \mathbf{J}^{-1}(\mu_0) \rightarrow \overline{\mathcal{Z}}_{\mu_0}$ be the quotient map. Then the reduced symplectic form $\overline{\omega}_{\mu_0}$ on $\overline{\mathcal{Z}}_{\mu_0}$ is uniquely determined by

$$i_{\mu_0}^* \omega = \pi_{\mu_0}^* \overline{\omega}_{\mu_0};$$

see [10] and Marsden *et al* [11, sections 1.1 and 1.2]. Also, let $\omega_{\mathcal{O}_{\Pi_0}}$ be the $(-)$ -Kirillov–Kostant–Souriau symplectic structure, i.e. for any $\Pi \in \mathcal{O}_{\Pi_0}$ and $A, B \in \mathfrak{o}(n)$,

$$\omega_{\mathcal{O}_{\Pi_0}}(-\text{ad}_A^* \Pi, -\text{ad}_B^* \Pi) := -\langle \Pi, [A, B] \rangle,$$

where $[\cdot, \cdot]$ is the commutator on $\mathfrak{o}(n)$; see, e.g. Kirillov [7, chapter 1] and Marsden and Ratiu [9, chapter 14]. Then the momentum map \mathbf{M} restricted to the level set $\mathbf{J}^{-1}(\mu_0)$ gives rise to a diffeomorphism $\overline{\mathbf{M}}: \overline{\mathcal{Z}}_{\mu_0} \rightarrow \mathcal{O}_{\Pi_0}$; moreover this map is symplectic with respect to the above symplectic forms, i.e.

$$\overline{\mathbf{M}}^* \omega_{\mathcal{O}_{\Pi_0}} = \overline{\omega}_{\mu_0}.$$

The diagram below gives an overview of this result.

$$\begin{array}{ccc} & \mathcal{Z} & \\ i_{\mu_0} \uparrow & & \\ \mathbf{J}^{-1}(\mu_0) & & \\ \pi_{\mu_0} \downarrow & \searrow \mathbf{M}|_{\mathbf{J}^{-1}(\mu_0)} & \\ \overline{\mathcal{Z}}_{\mu_0} & \xrightarrow[\overline{\mathbf{M}}]{} & \mathcal{O}_{\Pi_0} \end{array}.$$

3.5. Reduction of symmetric representation

Let $Q(0) = Q_0 \in \text{SO}(n)$ and $\Pi_0 \in \mathfrak{o}(n)^*$ be the initial rotational configuration and the initial body angular momentum of the rigid body, and fix $P_0 \in \text{SO}(n)$ so that

$$Q_0^T P_0 - P_0^T Q_0 = \Pi_0.$$

See Bloch and Crouch [3] and Bloch *et al* [4] for the condition under which this is possible. Then clearly $Z_0 := (Q_0, P_0)$ is in the open subset $\mathcal{Z} \subset \mathbf{M}_{2n \times n}(\mathbb{R})$ defined in (13) and $\Pi_0 = \mathbf{M}(Z_0)$. Now, setting

$$\mu_0 := \mathbf{J}(Z_0) = \begin{bmatrix} P_0 Q_0^T & I \\ -I & -Q_0 P_0^T \end{bmatrix} \in \mathfrak{sp}(2n, \mathbb{R})^*,$$

the level set

$$\mathbf{J}^{-1}(\mu_0) = \left\{ \begin{bmatrix} Q \\ P \end{bmatrix} \in \mathcal{Z} \mid Q Q^T = I, P P^T = I, P Q^T = P_0 Q_0^T \right\}$$

is an invariant submanifold of the symmetric representation (7).

Let $h: \mathfrak{o}(n)^* \rightarrow \mathbb{R}$ be a collective Hamiltonian, i.e. $h \circ \mathbf{M} = H$. From the expressions (9) and (12) of H and \mathbf{M} , we find

$$h(\Pi) = \frac{1}{2} \langle \Pi, \mathcal{I}^{-1}(\Pi) \rangle,$$

which is the Hamiltonian (4) of the generalized rigid body in the body representation. Then the result from the previous subsection implies that the $\mathrm{Sp}(2n, \mathbb{R})$ -reduced dynamics in $\overline{\mathcal{Z}}_{\mu_0}$ is equivalent to the Lie–Poisson equation

$$\dot{\Pi} = \mathrm{ad}_{Dh(\Pi)}^* \Pi$$

in the coadjoint orbit $\mathcal{O}_{\Pi_0} \subset \mathfrak{o}(n)^*$, where $Dh(\Pi) \in \mathfrak{o}(n)$ is defined so that, for any $\delta\Pi \in \mathfrak{o}(n)^*$,

$$\langle \delta\Pi, Dh(\Pi) \rangle = \frac{d}{ds} h(\Pi + s\delta\Pi) \Big|_{s=0} = \langle \delta\Pi, \mathcal{I}^{-1}(\Pi) \rangle,$$

that is, $Dh(\Pi) = \mathcal{I}^{-1}(\Pi)$. However, under the identification $\mathfrak{o}(n)^* \cong \mathfrak{o}(n)$, $\mathrm{ad}_A^* \Pi = [\Pi, A]$ for any $A \in \mathfrak{o}(n)$ and $\Pi \in \mathfrak{o}(n)^*$, and thus we obtain

$$\dot{\Pi} = [\Pi, \mathcal{I}^{-1}(\Pi)],$$

which is the n -dimensional Euler equation (3b).

Acknowledgments

I would like to thank Paul Skerritt for helpful discussions on dual pairs. This work was partially supported by NSF grant CMMI-1824798.

ORCID iDs

Tomoki Ohsawa  <https://orcid.org/0000-0001-9406-132X>

References

- [1] Abraham R and Marsden J E 1978 *Foundations of Mechanics* 2nd edn (Reading, MA: Addison-Wesley)
- [2] Balleier C and Wurzbacher T 2012 On the geometry and quantization of symplectic Howe pairs *Math. Z.* **271** 577–91
- [3] Bloch A M and Crouch P E 1996 Optimal control and geodesic flows *Syst. Control Lett.* **28** 65–72
- [4] Bloch A M, Crouch P E, Marsden J E and Ratiu T S 2002 The symmetric representation of the rigid body equations and their discretization *Nonlinearity* **15** 1309–41
- [5] Bloch A M, Crouch P E, Marsden J E and Sanyal A K 2008 Optimal control and geodesics on quadratic matrix Lie groups *Found. Comput. Math.* **8** 469–500
- [6] Bloch A M, Crouch P E, Nordkvist N and Sanyal A K 2011 Embedded geodesic problems and optimal control for matrix Lie groups *J. Geom. Mech.* **3** 197–223
- [7] Kirillov A A 2004 *Lectures on the Orbit Method* (Graduate Studies in Mathematics) (Providence, RI: American Mathematical Society)
- [8] Manakov S V 1976 Note on the integration of Euler's equations of the dynamics of an n -dimensional rigid body *Funct. Anal. Appl.* **10** 328–9
- [9] Marsden J E and Ratiu T S 1999 *Introduction to Mechanics and Symmetry* (Berlin: Springer)
- [10] Marsden J E and Weinstein A 1974 Reduction of symplectic manifolds with symmetry *Rep. Math. Phys.* **5** 121–30

- [11] Marsden J E, Misiolek G, Ortega J P, Perlmutter M and Ratiu T S 2007 *Hamiltonian Reduction by Stages (Lecture Notes in Mathematics)* (Berlin, Heidelberg: Springer-Verlag) (<https://doi.org/10.1007/978-3-540-72470-4>)
- [12] Pontryagin L S, Boltyanskii V G, Gamkrelidze R V and Mishchenko E F 1962 *The Mathematical Theory of Optimal Processes* (New York: Wiley)
- [13] Ratiu T S 1980 The motion of the free n -dimensional rigid body *Indiana Univ. Math. J.* **29** 609–29
- [14] Skerritt P 2018 The frame bundle picture of Gaussian wave packet dynamics in semiclassical mechanics (arXiv:[1802.04362](https://arxiv.org/abs/1802.04362))
- [15] Skerritt P and Vizman C 2019 Dual pairs for matrix groups *J. Geom. Mech.* **11** 255–75