

LINEARIZATION OF NONLINEAR FOKKER-PLANCK EQUATIONS AND APPLICATIONS^{*,†}

BY PANPAN REN, MICHAEL RÖCKNER[†] AND FENG-YU WANG^{*}

Tianjin University, Bielefeld University, Chinese Academy of Science, University of Oxford and Swansea University.

Let $\mathcal{P}$ be the space of probability measures on $\mathbb{R}^d$. We associate a coupled nonlinear Fokker-Planck equation on $\mathbb{R}^d$, i.e. with solution paths in $\mathcal{P}$, to a linear Fokker-Planck equation for probability measures on the product space $\mathbb{R}^d \times \mathcal{P}$, i.e. with solution paths in $\mathcal{P}(\mathbb{R}^d \times \mathcal{P})$. We explicitly determine the corresponding linear Kolmogorov operator $\mathbf{L}_t$ using the natural tangent bundle over $\mathcal{P}$ with corresponding gradient operator $\nabla^{\mathcal{P}}$. Then it is proved that the diffusion process generated by $\mathbf{L}_t$ on $\mathbb{R}^d \times \mathcal{P}$ is intrinsically related to the solution of a McKean-Vlasov stochastic differential equation (SDE). We also characterize the ergodicity of the diffusion process generated by $\mathbf{L}_t$ in terms of asymptotic properties of the coupled nonlinear Fokker-Planck equation. Another main result of the paper is that the restricted well-posedness of the non-linear Fokker-Planck equation and its linearized version imply the (restricted) well-posedness of the McKean-Vlasov equation and that in this case the laws of the solutions have the Markov property. As applications, we prove the restricted weak well-posedness and the Markov property of the so-called nonlinear distorted Brownian motion, whose associated nonlinear Fokker-Planck equation is a porous media equation perturbed by a nonlinear transport term. As a further application we obtain a probabilistic representation of solutions to Schrödinger type PDEs on $\mathbb{R}^d \times \mathcal{P}_2$, through the Feynman-Kac formula for the corresponding diffusion processes.

1. Introduction. As a first result of this paper (see Section 3) we identify the continuity equation corresponding to a non-linear Fokker-Planck equation on $\mathbb{R}^d$ with weakly continuous solution paths in $\mathcal{P}$, i.e. the space of probability measures on $\mathbb{R}^d$ equipped with the weak topology. We determine explicitly the vector field, defining the continuity equation, as a section in the natural tangent bundle over $\mathcal{P}$. More precisely, let for $m, d \in \mathbb{N}$

$$(1.1) \quad \begin{aligned} b &= (b_i)_{1 \leq i \leq d}: [0, \infty) \times \mathbb{R}^d \times \mathcal{P} \rightarrow \mathbb{R}^d, \\ \sigma &= (\sigma_{ij})_{1 \leq i, j \leq d}: [0, \infty) \times \mathbb{R}^d \times \mathcal{P} \rightarrow \mathbb{R}^d \otimes \mathbb{R}^m \end{aligned}$$

be Borel measurable maps and consider the corresponding nonlinear Kolmogorov operator

$$(1.2) \quad \begin{aligned} L_{t,\mu} h(x) &= \frac{1}{2} \sum_{i,j=1}^d (\sigma \sigma^*)_{i,j}(t, x, \mu) \partial_i \partial_j h(x) + \sum_{i=1}^d b_i(t, x, \mu) \partial_i h(x) \\ &= \frac{1}{2} ((\sigma \sigma^*)(t, x, \mu) \nabla) \cdot \nabla h(x) + b(t, x, \mu) \cdot \nabla h(x), \end{aligned}$$

^{*}Supported in part by NNSFC (11771326, 11831014) and the DFG through the CRC 1283.

[†]Supported in part by the DFG through the CRC 1283.

MSC 2010 subject classifications: Primary 60J60; secondary 58J65.

Keywords and phrases: Nonlinear Fokker-Planck equation, McKean-Vlasov stochastic differential equation, diffusion process, ergodicity, Feynman-Kac formula.

where $(t, x, \mu) \in [0, \infty) \times \mathbb{R}^d \times \mathcal{P}$, $h \in C_0^2(\mathbb{R}^d)$, “ $\cdot$ ” denotes inner product in $\mathbb{R}^d$ and $L_{t,\mu}$ determines the nonlinear Fokker-Planck equation

$$(1.3) \quad \partial_t \mu_t = L_{t,\mu_t}^* \mu_t,$$

meant in the weak sense with test function space $C_0^\infty(\mathbb{R}^d)$ (see Definition 2.2 below for details). By the recipe suggested in [1], [2] (see also [3], [4], [22], [25], [26], [30] and [31]), which we briefly recall in the Appendix of this paper, one then finds the well-known tangent bundle over $\mathcal{P}$, namely

$$(1.4) \quad (L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu))_{\mu \in \mathcal{P}},$$

and a corresponding intrinsic gradient $\nabla^{\mathcal{P}}$ for a suitable and sufficiently large class $\mathcal{F}C_b^2(\mathcal{P})$ of functions $F: \mathcal{P} \rightarrow \mathbb{R}^d$ (defined in (2.4) below), which maps such F into sections in this tangent bundle, i.e.

$$(1.5) \quad \nabla^{\mathcal{P}} F(\mu) \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu), \quad \mu \in \mathcal{P}$$

(for details see the Appendix). $\nabla^{\mathcal{P}}$ is, of course, the well-known Otto-gradient introduced in [24]. For the case where the space of $\mathbb{Z}_+$ -valued Radon measures on $\mathbb{R}^d$ replaces $\mathcal{P}$, it was, however, already introduced in [1], [2], even with a Riemannian manifold M replacing $\mathbb{R}^d$. Then the continuity equation corresponding to the non-linear evolution equation (1.3), which is *linear* and by definition an equation, whose solutions are weakly continuous paths of measure Γ_t , $t \geq 0$, in $\mathcal{P}(\mathcal{P})$ (the space of probability measures on $\mathcal{P}$ again equipped with the weak topology), is then given as

$$(1.6) \quad \partial_t \Gamma_t = (\tilde{\mathbf{L}}_t)^* \Gamma_t$$

in the weak sense with test function space $\mathcal{F}C_b^2(\mathcal{P})$ (see Section 3 for details). The corresponding Kolmogorov operator $\tilde{\mathbf{L}}_t$ is of first order and determined by a vector field in the tangent bundle $(L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu))_{\mu \in \mathcal{P}}$ as follows:

$$(1.7) \quad \tilde{\mathbf{L}}_t F(\mu) = \left\langle \frac{1}{2}(\sigma\sigma^*)(t, \cdot, \mu) \nabla + b(t, \cdot, \mu), \nabla^{\mathcal{P}} F(\mu) \right\rangle_{L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)},$$

where $\mu \in \mathcal{P}$, $F \in \mathcal{F}C_b^2(\mathcal{P})$ and $\langle \cdot, \cdot \rangle_{L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)}$ denotes the inner product in $L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)$ (see Proposition 3.1). So, every solution μ_t^ζ , $t \geq 0$, to (1.3) with initial condition $\zeta \in \mathcal{P}$ gives rise to a solution to (1.5) with initial condition the Dirac measure δ_ζ in $\mathcal{P}(\mathcal{P})$. Mixing these initial conditions ζ according a measure $\Gamma \in \mathcal{P}(\mathcal{P})$, i.e., looking at the push forward measure under the flow generated by (1.3) one obtains a solution path Γ_t^Γ , $t \geq 0$, in $\mathcal{P}(\mathcal{P})$ of (1.6). Interestingly, it turns out the operator $\partial_t + \tilde{\mathbf{L}}_t$ with a suitable domain $\tilde{\mathcal{F}}$ is dissipative, hence closable on $L^1([0, T] \times \mathcal{P}, \Gamma_t^\Gamma dt)$. Thus $\partial_t + \tilde{\mathbf{L}}_t$ extends uniquely to a much larger space of functions on $[0, T] \times \mathcal{P}$ (including e.g. certain functions which are Lipschitz with respect to metrics generating the weak topology on $\mathcal{P}$). Thus, through (1.6) we can construct an abundance of measures of type $\Gamma_t^\Gamma dt$ on $[0, T] \times \mathcal{P}$ for which the first order operator $\partial_t + \tilde{\mathbf{L}}_t$, which is given by a vector field over $[0, T] \times \mathcal{P}(\mathbb{R}^d)$, i.e. a section in $(L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu))_{\mu \in \mathcal{P}(\mathbb{R}^d)}$, and which in turn is canonically determined by the nonlinear Kolmogorov operator (1.2), is closable on $L^1([0, T] \times \mathcal{P}(\mathbb{R}^d); \Gamma_t^\Gamma dt)$ (see Remark 3.2 below for details).

In the second main result of this paper (see Section 4.1) we obtain weak uniqueness in law and Markov properties for solutions to McKean-Vlasov equations from uniqueness of their corresponding non-linear Fokker-Planck equation and their “freed” linear version. More precisely, for b and σ

as in (1.1) consider the McKean-Vlasov stochastic differential equation (SDE) on $\mathbb{R}^d$ (see [12] and the references therein)

$$(1.8) \quad dX_t = b(t, X_t, \mathcal{L}_{X_t})dt + \sigma(t, X_t, \mathcal{L}_{X_t})dW_t, \quad t \geq 0,$$

where W_t , $t \geq 0$, is an $(\mathcal{F}_t)$ -Brownian motion in $\mathbb{R}^m$ defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with normal filtration $(\mathcal{F}_t)_{t \geq 0}$ and the solution X_t , $t \geq 0$, is an $(\mathcal{F}_t)$ -adapted stochastic process on $(\Omega, \mathcal{F})$ with $\mathbb{P}$ -a.s. continuous sample paths in $\mathbb{R}^d$ and time marginal laws $\mathcal{L}_{X_t} := \mathbb{P} \circ X_t^{-1}$, $t \geq 0$. Then obviously by Itô's formula for any (probabilistically) weak solution to (1.8) its time marginals $\mu_t := \mathcal{L}_{X_t}$, $t \geq 0$, solve the nonlinear Fokker-Planck equation (1.3). Recently, it was proved in [5], [6] that under a natural integrability condition on σ and b the converse is also true. Hence in this sense weak solutions to McKean-Vlasov SDE and nonlinear Fokker-Planck equations are equivalent. Once one has this equivalence, it is fairly straightforward to get a sufficient condition for weak uniqueness for (1.8). For this we consider the “frozen” linear version of the non-linear Fokker-Planck equation (1.3), i.e. for a fixed solution μ_t , $t \geq 0$, of (1.3)

$$(1.9) \quad \partial_t \nu_t = L_{t, \mu_t}^* \nu_t$$

and look at the pair of Fokker-Planck equations

$$(1.10) \quad \begin{cases} \partial_t \mu_t &= L_{t, \mu_t} \mu_t \\ \partial_t \nu_t &= L_{t, \mu_t}^* \nu_t. \end{cases}$$

Then we introduce a notion of “restricted” well-posedness for (1.10) (see Definition 2.2(2) below). It is “restricted” in the sense that we restrict to subclasses of probability measures as initial data and to subclasses of solutions with certain properties. This restriction causes major technical complications, but is necessary, because the equation is only well solved on a subspace of $\mathcal{P}$ according to specific conditions on coefficients, and when the distribution density function is concerned it is natural to consider the class of absolutely continuous probability measures (see for instance Section 5). Indeed, when $b(t, x, \cdot)$ and $\sigma(t, x, \cdot)$ are Lipschitz continuous in the Wasserstein distance $\mathbb{W}_2$, (1.8) is solved for initial distributions having finite second moment, see [14, 20, 34] and references within. Theorem 4.1 below provides the equivalence of the restricted well-posedness of (1.10) with that of the corresponding SDE-version, i.e.,

$$(1.11) \quad \begin{cases} \partial_t \mu_t &= L_{t, \mu_t} \mu_t \\ dX(t) &= b(t, X(t), \mu_t)dt + \sigma(t, X(t), \mu_t)dW(t). \end{cases}$$

From Theorem 4.1 we then deduce that the restricted well-posedness of (1.10) implies that of the McKean-Vlasov SDE (1.8). Under the assumption of restricted well-posedness we also prove that the laws of the solutions to both the second equation in (1.11) and to (1.8) have the Markov property (see Theorem 4.5 and Corollary 4.6 in Section 4.3). The proof, however, is much more involved in comparison to the case of well-posedness for all initial conditions. It turns out that the same results hold, if in the second equations of (1.10) and (1.11), we change the operator L_{t, μ_t} , i.e. the coefficients b and σ , to another Kolmogorov operator $\bar{L}_{t, \mu_t}$ with coefficients $\bar{b}, \bar{\sigma}$. Therefore, we formulate our results in this more general case.

In Section 5 we apply these results to an interesting example, where the above nonlinear Fokker-Planck equation is a porous medium equation perturbed by a first order (transport) term recently studied in [7]. The solution to the corresponding McKean-Vlasov SDE can be considered as a nonlinear distorted Brownian motion (see [7] for details).

The third main result of this paper is about transforming the non-linear coupled Fokker-Planck equation (1.10) into a *linear* Fokker-Planck equation on $\mathbb{R}^d \times \mathcal{P}$, i.e. with solution paths in $\mathcal{P}(\mathbb{R}^d \times \mathcal{P})$ (see Section 4.2). The motivation comes from the hope that for the understanding of McKean-Vlasov SDEs it could be useful to study the pair process $(X(t), \mathcal{L}_{X(t)})$, $t \geq 0$, on the state space $\mathbb{R}^d \times \mathcal{P}$. By the Appendix of this paper we know that the corresponding tangent bundle is

$$(1.12) \quad \mathbb{R}^d \oplus (L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d; \mu))_{\mu \in \mathcal{P}}.$$

As a consequence of this and Section 3 we derive the Fokker-Planck equation associated to the process $(X(t), \mathcal{L}_{X(t)})_{t \geq 0}$ on $\mathbb{R}^d \times \mathcal{P}$ which turns out to be a *linear* Fokker-Planck equation, namely (see Definition 2.2 below)

$$(1.13) \quad \partial_t \Lambda_t = \mathbf{L}_t^* \Lambda_t, \quad t \geq 0,$$

where $(\Lambda_t)_{t \geq 0}$ is a weakly continuous path of probability measures on $\mathbb{R}^d \times \mathcal{P}$, i.e. in $\mathcal{P}(\mathbb{R}^d \times \mathcal{P})$, and with the corresponding *linear* Kolmogorov operator $\mathbf{L}_t$ (of course, first order in μ) on $\mathbb{R}^d \times \mathcal{P}$ being given explicitly on a reasonably rich class $\mathcal{C}$ of functions $G : \mathbb{R}^d \times \mathcal{P} \rightarrow \mathbb{R}$ as

$$(1.14) \quad \mathbf{L}_t G = \mathbf{L}_t^{(1)} G + \mathbf{L}_t^{(2)} G$$

where

$$(1.15) \quad \mathbf{L}_t^{(1)} G(x, \mu) = L_t(G(\cdot, \mu))(x)$$

$$(1.16) \quad \mathbf{L}_t^{(2)} G(x, \mu) = \left\langle \frac{1}{2}(\sigma \sigma^*)(t, \cdot, \mu) \nabla + b(t, \cdot, \mu), \nabla^{\mathbb{R}^d \times \mathcal{P}} G(x, \mu) \right\rangle_{L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)}$$

(see Section 4.2 for details). The exact relation between solutions to (1.10) and (1.13) is given in Theorem 4.3(i) below. In particular, an explicit formula for solutions of (1.13) is given through probability kernels $\mathbf{P}_{s,t}(x, \zeta; dy, d\nu) \in \mathcal{P}(\mathbb{R}^d \times \mathcal{P})$, $(x, \zeta \in \mathbb{R}^d \times \mathcal{P}, s \leq t$, which satisfy the Chapman-Kolmogorov equations (see Theorem 4.3(ii)).

As another consequence we prove the Markov property of the law of the solution $(X(t), \mathcal{L}_{X(t)})$ of the McKean-Vlasov SDE (1.8) in a stronger form under the stronger condition (4.8) (see Theorem 4.11 in Section 4.4 below for the precise formulation of this result).

In the time homogeneous case a further application of the results in Section 4 is a characterization of the ergodicity for solutions to (1.11) in terms of the ergodicity of $(\mathbf{P}_{s,t})_{s \leq t}$ (see Theorem 4.12 in Section 4.5). An application of this result is presented in Section 6 and concerns a case with more regular coefficients b and σ (see Theorem 6.1 below) where $\mathcal{P}(\mathbb{R}^d)$ is replaced by $\mathcal{P}_2(\mathbb{R}^d)$, i.e. all elements in $\mathcal{P}(\mathbb{R}^d)$ with finite second moments (equipped with the Wasserstein metric). In this case even exponential ergodicity is proved.

As a further consequence, in Section 7 we then prove a Feynman-Kac formula for the above diffusion process on $\mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d)$, from which we derive a probabilistic representation for solutions of Schrödinger type PDE on $\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)$ of the following form

$$(1.17) \quad \partial_t u(t, x, \zeta) + \mathbf{L}_t u(t, \cdot, \cdot)(x, \zeta) + (\mathbf{V}u)(t, x, \zeta) + \mathbf{f}(t, x, \zeta) = 0, \quad t \in [0, T],$$

where $T > 0$ is fixed, $(x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)$, and $\mathbf{V}, \mathbf{f}$ are measurable functions on $[0, T] \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)$. This generalizes some known results from the literature (see [10, 13, 15, 19, 21]).

Finally we would like to emphasize again that the main motivation of this paper is to contribute to the theory of nonlinear Fokker-Planck equations on one side and McKean-Vlasov SDEs on the other. The literature on both parts of the theory is overwhelming, so that we apologize that an overview of the known results is beyond the scope of this paper. Therefore, we confine ourselves to refer the reader to the monographs [17] and [9] concerning Fokker-Planck(-Kolmogorov) equations and to [12] concerning McKean-Vlasov SDEs as well as the references therein and e.g. the very recent papers [5, 6, 11, 14, 18, 19, 20, 23, 27, 28, 29, 34]. Furthermore, we would like to stress that according to general Markov process theory via the martingale problem given by the underlying generator (see e.g. [32] in the classical case) there is a close connection of our results with the very nice recent works ([13], [16]) on the Itô-formula for the process $(X_t, \mathcal{L}_{X(t)})$, $t \geq 0$, coming from the McKean-Vlasov SDE (1.8) above. The connection is obvious, since one can show that our intrinsic gradient $\nabla^{\mathcal{P}}$ on functions in $\mathcal{FC}_b^2(\mathcal{P})$ (see Appendix A) is the same as the Lions-derivative from [11]. This fact was proved in [29]. Since, however, our approach is more analytic and based more on nonlinear Fokker-Planck equations, we do not need this Itô-formula.

2. Preliminaries and notation. Let $\mathcal{P}$ denote the set of all probability measures on $\mathbb{R}^d$ equipped with the weak (= narrow) topology and corresponding Borel- σ -algebra $\mathcal{B}(\mathcal{P})$. Likewise $\mathcal{P}(\mathcal{P})$ and $\mathcal{P}(\mathbb{R}^d \times \mathcal{P})$ denote the set of all probability measures on $\mathcal{P}$ and $\mathbb{R}^d \times \mathcal{P}$ respectively and both are considered with the weak topology and corresponding Borel- σ -algebras. Let

$$(2.1) \quad b, \bar{b} : [0, \infty) \times \mathbb{R}^d \times \mathcal{P} \rightarrow \mathbb{R}^d; \quad \sigma, \bar{\sigma} : [0, \infty) \times \mathbb{R}^d \times \mathcal{P} \rightarrow \mathbb{R}^d \otimes \mathbb{R}^m,$$

be Borel-measurable maps, where $m, d \in \mathbb{N}$. Furthermore, besides $L_{t,\mu}$ in (1.2) we define the following measure-dependent Kolmogorov operator on $\mathbb{R}^d$:

$$(2.2) \quad \bar{L}_{t,\mu} h(x) := \frac{1}{2} \sum_{i,j=1}^d (\bar{\sigma} \bar{\sigma}^*)_{ij}(t, x, \mu) \partial_i \partial_j h(x) + \sum_{i=1}^d \bar{b}_i(t, x, \mu) \partial_i h(x), \quad h \in C_0^2(\mathbb{R}^d).$$

Let us define the following test function spaces:

$$(2.3) \quad \mathcal{FC}_b^2(\mathcal{P}) := \{F(\mu) := g(\mu(h_1), \dots, \mu(h_n)) : n \geq 1, g \in C_b^1(\mathbb{R}^n), h_i \in C_0^2(\mathbb{R}^d)\}.$$

on $\mathcal{P}$ and

$$(2.4) \quad \mathcal{C} := \{(x, \mu) \mapsto h_0(x)F(\mu) : h_0 \in C_0^2(\mathbb{R}^d), F \in \mathcal{FC}_b^2(\mathcal{P})\}.$$

Next, consider the time-dependent differentiable operator $\mathbf{L}_t$ defined by (1.14), (1.15) and (1.16), where $\nabla^{\mathbb{R}^d \times \mathcal{P}}$ is as defined in (A.4) in the Appendix, but with $\mathbb{R}^d \times \mathcal{P}$ replacing $\mathcal{P}$ (see Subsection 4.2 below). Consider the coupled nonlinear Fokker-Planck equation (1.10) meant in the weak sense (see Definition 2.2(2) below). In Section 4 we are going to establish a correspondence between solutions $(\mu_t, \nu_t)_{t \geq 0}$ of (1.10) and those to the *linear* Fokker-Planck equation (1.13) on $\mathbb{R}^d \times \mathcal{P}$ again meant in the weak sense (see Definition 2.2(3) below). Let $C([s, \infty) \rightarrow \mathcal{P})$ denote the set of all weakly continuous paths in $\mathcal{P}$ starting from $s \in [0, \infty)$ and $C([s, \infty) \rightarrow \mathcal{P}(\mathcal{P}))$, $C([s, \infty) \rightarrow \mathcal{P}(\mathbb{R}^d \times \mathcal{P}))$ are defined likewise.

REMARK 2.1. *For nonlinear Fokker-Planck equations (1.3) typically one cannot expect to have a unique solution for every initial condition $\zeta \in \mathcal{P}$ at time $s \in [0, \infty)$, but only for $\zeta \in \mathcal{P}_0$, where $\mathcal{P}_0 \in \mathcal{B}(\mathcal{P})$. In addition, even for such restricted initial conditions $\zeta \in \mathcal{P}_0$ generally one does not have a unique solution to (1.3) in all of $C([s, \infty) \rightarrow \mathcal{P})$, but rather in a subset thereof whose paths in particular leave $\mathcal{P}_0$ invariant. Therefore, we introduce property (P) below.*

For $\mathcal{P}_0 \in \mathcal{B}(\mathcal{P})$ and $\mathcal{A} \subset C([0, \infty) \rightarrow \mathcal{P}_0)$ we need the following property for the pair $(\mathcal{P}_0, \mathcal{A})$:

- (P) If $\tilde{\mu} \in \mathcal{P}$ such that $\tilde{\mu} \leq C\mu$ for some $\mu \in \mathcal{P}_0$, $C \in (0, \infty)$, then $\tilde{\mu} \in \mathcal{P}_0$. If $(\lambda_t)_{t \geq 0} \in \mathcal{A}$, then $(\lambda_{s+t})_{t \geq 0} \in \mathcal{A}$ for all $s \geq 0$, and $(\tilde{\lambda}_t)_{t \geq 0} \in \mathcal{A}$, provided $(\tilde{\lambda}_t)_{t \geq 0} \in C([0, \infty) \rightarrow \mathcal{P}_0)$ and $\tilde{\lambda}_t \leq C\lambda_t$ for all $t \geq 0$ and some $C \in (0, \infty)$.

DEFINITION 2.2 (Solution to Fokker-Planck equations, see [9]). *Let $s \geq 0$ be fixed.*

- (1) $(\mu_t)_{t \geq s} \in C([s, \infty) \rightarrow \mathcal{P})$ is called a solution to (1.3) from time s , if for all $t \in [s, \infty)$

$$(2.5) \quad \int_s^t dr \int_{\mathbb{R}^d} (|b| + \|\sigma\|^2)(r, x, \mu_r) \mu_r(dx) < \infty,$$

and

$$(2.6) \quad \int_{\mathbb{R}^d} h d\mu_t = \int_{\mathbb{R}^d} h d\mu_s + \int_s^t dr \int_{\mathbb{R}^d} L_{r, \mu_r} h d\mu_r, \quad h \in C_0^\infty(\mathbb{R}^d).$$

- (2) A pair $(\mu_t, \nu_t)_{t \geq s}$ with $(\mu_t)_{t \geq s}, (\nu_t)_{t \geq s} \in C([s, \infty) \rightarrow \mathcal{P})$ is called a solution to (1.10) from time s , if $(\mu_t)_{t \geq s}$ is a solution of (1.3) from time s and for all $t \in [s, \infty)$

$$(2.7) \quad \int_s^t dr \int_{\mathbb{R}^d} (|\bar{b}| + \|\bar{\sigma}\|^2)(r, x, \mu_r) \nu_r(dx) < \infty$$

and

$$(2.8) \quad \int_{\mathbb{R}^d} h d\nu_t = \int_{\mathbb{R}^d} h d\nu_s + \int_s^t dr \int_{\mathbb{R}^d} \bar{L}_{r, \mu_r} h d\nu_r, \quad h \in C_0^\infty(\mathbb{R}^d).$$

For two pairs $(\mathcal{P}_0, \mathcal{A}), (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}})$ with property (P) we call (1.10) well-posed in

$((\mathcal{P}_0, \mathcal{A}), (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}}))$ if the following holds:

- (a) For every $(s, \zeta) \in [0, \infty) \times \mathcal{P}_0$ there exists a unique solution $(\mu_{s,t}^\zeta)_{s,t \geq 0}$ to (1.3) starting from s with $\mu_{s,s}^\zeta = \zeta$ and $(\mu_{s,s+t}^\zeta)_{t \geq 0} \in \mathcal{A}$ such that $\zeta \mapsto \mu_{s,t}^\zeta$ is Borel measurable for all s, t .
- (b) For every $(s, \theta, \zeta) \in [0, \infty) \times \tilde{\mathcal{P}}_0 \times \mathcal{P}_0$ and $(\mu_{s,t}^\zeta)_{t \geq 0}$ as in (a) there exists a unique solution $(\nu_{s,t}^{\zeta, \theta})_{t \geq s}$ to (2.8) with $\mu_{s,r}^\zeta$ replacing μ_r , $r \geq s$, starting from s with $\nu_{s,s}^{\zeta, \theta} = \theta$ and $(\nu_{s,s+t}^{\zeta, \theta})_{t \geq 0} \in \tilde{\mathcal{A}}$ such that $(\theta, \zeta) \mapsto \nu_{s,t}^{\zeta, \theta}$ is Borel measurable for all s, t .
- (3) $(\Lambda_t)_{t \geq s} \in C([s, \infty) \rightarrow \mathcal{P}(\mathbb{R}^d \times \mathcal{P}))$ is called a solution to (1.13) from time s , if for all $t \in [s, \infty)$

$$(2.9) \quad \int_s^t dr \int_{\mathbb{R}^d} (|b| + \|\sigma\|^2)(r, \cdot, \mu) \big|_{L^1(\mathbb{R}^d, \mu)} + (|\bar{b}| + \|\bar{\sigma}\|^2)(r, x, \mu) \Lambda_r(dx, d\mu) < \infty,$$

and for any $G \in \mathcal{C}$,

$$(2.10) \quad \int_{\mathbb{R}^d \times \mathcal{P}} G d\Lambda_t = \int_{\mathbb{R}^d \times \mathcal{P}} G d\Lambda_s + \int_s^t dr \int_{\mathbb{R}^d \times \mathcal{P}} \mathbf{L}_r G d\Lambda_r.$$

(4) $(\Gamma_t)_{t \geq s} \in C([s, \infty) \rightarrow \mathcal{P}(\mathcal{P}))$ is called a solution to (1.6) from time s , if for all $t \in [s, \infty)$

$$(2.11) \quad \int_s^t dr \int_{\mathbb{R}^d} |(|b| + \|\sigma\|^2)(r, \cdot, \mu)|_{L^1(\mathbb{R}^d, \mu)} \Gamma_r(d\mu) < \infty,$$

and for any $F \in \mathcal{F}C_b^2(\mathcal{P})$

$$(2.12) \quad \int_{\mathcal{P}} F d\Gamma_t = \int_{\mathcal{P}} F d\Gamma_s + \int_s^t dr \int_{\mathcal{P}} \tilde{L}_r F d\Gamma_r.$$

REMARK 2.3. If (1.10) is well-posed, $b = \bar{b}$, $\sigma = \bar{\sigma}$ and $\mathcal{P}_0 \subset \tilde{\mathcal{P}}_0$, we have for all $(s, \zeta) \in [0, \infty) \times \mathcal{P}_0$

$$\nu_{s,t}^{\zeta, \zeta} = \mu_{s,t}^{\zeta}, \quad t \geq 0$$

because both solve (1.3) with the same initial condition ζ .

Let us consider another coupled equation involving the first equation of (1.10) and the stochastic equation corresponding to the linear Kolmogorov operator $\bar{L}_{\mu,t}$ in (1.2), i.e.

$$(2.13) \quad \begin{cases} \partial_t \mu_t = L_{t, \mu_t}^* \mu_t, \\ dX_t = \bar{b}(t, X_t, \mu_t) dt + \bar{\sigma}(t, X_t, \mu_t) dW_t, \end{cases}$$

with solution paths $(X_t, \mu_t)_{t \geq 0}$ in $\mathbb{R}^d \times \mathcal{P}$. Here $(W_t)_{t \geq 0}$ is an $(\mathcal{F}_t)$ -Brownian motion on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with normal filtration $(\mathcal{F}_t)_{t \geq 0}$ and values in $\mathbb{R}^m$. Below we set $\mathcal{L}_{X_t} := \mathbb{P} \circ X_t^{-1}$ (“time marginal law” at t). We now introduce the notion of weak solution and weak well-posedness for (2.13).

DEFINITION 2.4. (i) Let $(s, \theta, \zeta) \in [0, \infty) \times \mathcal{P} \times \mathcal{P}$. If (1.3) has a solution $(\mu_t)_{t \geq s}$ with $\mu_s = \zeta$, and the SDE

$$(2.14) \quad dX_{s,t}^{\zeta, \theta} = \bar{b}(t, X_{s,t}^{\zeta, \theta}, \mu_t) dt + \bar{\sigma}(t, X_{s,t}^{\zeta, \theta}, \mu_t) dW_t, \quad t \geq s, \mathcal{L}_{X_{s,s}^{\zeta, \theta}} = \theta,$$

has a pathwise continuous weak solution, then $(X_{s,t}^{\zeta, \theta}, \mu_t)_{t \geq s}$ is called a weak solution to (2.13) with initial value (θ, ζ) at time s .

(ii) Let $(\mathcal{P}_0, \mathcal{A}), (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}})$ be two pairs with property **(P)**. We call (2.13) well-posed in $((\mathcal{P}_0, \mathcal{A}), (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}}))$, if (a) in Definition 2.2 (2) holds and if for every $(s, \theta, \zeta) \in [0, \infty) \times \tilde{\mathcal{P}}_0 \times \mathcal{P}_0$ and for the unique solution $(\mu_{s,t}^{\zeta})_{s \geq t}$ to (1.3) there exists a unique weak solution $(X_{s,t}^{\zeta, \theta})_{t \geq s}$ to (2.14) with $\mu_{s,t}^{\zeta}$ replacing μ_t , $t \geq s$, such that for all s, t the law of $X_{s,t}^{\zeta, \theta}$ is Borel-measurable in (ζ, θ) .

REMARK 2.5. Obviously, in the situation of Definition 2.4(i) we have by Itô’s formula that $\nu_{s,t}^{\zeta, \theta} := \mathcal{L}_{X_{s,t}^{\zeta, \theta}}, t \geq s$, is a solution to (2.8).

Now we consider the McKean-Vlasov SDE (1.8)

DEFINITION 2.6. (i) Let $(s, \zeta) \in [0, \infty) \times \mathcal{P}$. The McKean-Vlasov SDE (1.8) is said to have a weak solution starting from s with time marginal law ζ , if there exist an $(\mathcal{F}_t)$ -Brownian

motion $W_t, t \geq 0$, on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with normal filtration $(\mathcal{F}_t)_{t \geq 0}$ and values in $\mathbb{R}^m$ and an $(\mathcal{F}_t)$ -adapted continuous process $(X_t)_{t \geq s}$ in $\mathbb{R}^d$ satisfying (1.8), such that $\mathcal{L}_{X_s} = \zeta$,

$$\int_s^t dr \int_{\mathbb{R}^d} (|b| + \|\sigma\|^2)(r, x, \mathcal{L}_{X_r}) \mathcal{L}_{X_r}(dx) < \infty, \quad t \geq s,$$

and $\mathbb{P}$ -a.s.

$$X_t = X_s + \int_s^t b(r, X_r, \mathcal{L}_{X_r}) dr + \int_s^t \sigma(r, X_r, \mathcal{L}_{X_r}) dW_r, \quad t \geq s.$$

(ii) Let $(\mathcal{P}_0, \mathcal{A})$ be a pair with property **(P)**. We call (1.8) weakly well-posed in $(\mathcal{P}_0, \mathcal{A})$ if for every $(s, \zeta) \in [0, \infty) \times \mathcal{P}_0$ and any two weak solutions with time marginal laws $(\mathcal{L}_{X_t})_{t \geq s}$, $(\mathcal{L}_{\tilde{X}_t})_{t \geq s} \in \mathcal{A}$ and $\mathcal{L}_{X_s} = \mathcal{L}_{\tilde{X}_s} = \zeta$, we have that their laws coincide.

We close this section with recalling the following result from [5, 6].

THEOREM 2.7. *Let $(s, \zeta) \in [0, \infty) \times \mathcal{P}$. Then the McKean-Vlasov SDE (1.8) has a weak solution starting from s with time marginal law ζ , if and only if (1.3) has a solution $(\mu_t)_{t \geq s}$ starting from s with $\mu_s = \zeta$. In this case $\mu_t = \mathcal{L}_{X_t}$, $t \geq s$.*

PROOF. If (1.8) has a weak solution starting from s with time marginal law ζ then by Remark 2.5 it follows that $\mu_t := \mathcal{L}_{X_t}$, $t \geq s$, is a solution of (1.3) starting from s with $\mu_s = \zeta$. The converse is proved in Section 2 of [6] (and [5]). $\square$

3. Linearization of nonlinear Fokker-Planck equations and construction of probability measures on $[s, \infty) \times \mathcal{P}$. Let $s \geq 0$ be fixed. The nonlinear Fokker-Planck equation (1.3) can be considered as an evolution equation in $\mathcal{P}$. For evolution equations there is a standard way to linearize them by transforming them to an evolution equation on the space of probability measures over their state space, hence in our case onto $\mathcal{P}(\mathcal{P})$, i.e. the space of probability measures on $\mathcal{P}$ equipped with the Borel- σ -algebra generated by the weak topology on $\mathcal{P}$. More precisely, for weakly continuous solution paths $(\mu_t)_{t \geq s}$ for (1.3) in $\mathcal{P}$, one derives an equation for the paths $(\delta_{\mu_t})_{t \geq s}$ in $\mathcal{P}(\mathcal{P})$ as follows:

For all $F \in \mathcal{F}C_b^2(\mathcal{P})$,

$$(3.1) \quad F(\mu) = f(\mu(h_1), \dots, \mu(h_n)), \quad n \in \mathbb{N}, \quad h_1, \dots, h_n \in C_0^2(\mathbb{R}^d), \quad f \in C_b^1(\mathbb{R}^n),$$

we have for any path $(\mu_t)_{t \geq s}$, in $\mathcal{P}$, such that $t \mapsto \mu_t(h)$ is weakly differentiable for all $h \in C_0^2(\mathbb{R}^d)$, by the chain rule and (A.4) in the Appendix

$$\begin{aligned} \frac{d}{dt} \delta_{\mu_t}(F) &= \frac{d}{dt} F(\mu_t) = \sum_{i=1}^n \partial_i f(\mu_t(h_1), \dots, \mu_t(h_n)) \partial_t \mu_t(h_i) \\ &= \sum_{i=1}^n \partial_i f(\mu_t(h_1), \dots, \mu_t(h_n)) \int_{\mathbb{R}^d} L_{t, \mu_t} h_i d\mu_t \\ &= \sum_{i=1}^n \left(\partial_i f(\mu_t(h_1), \dots, \mu_t(h_n)) \right. \\ &\quad \cdot \int_{\mathbb{R}^d} \left(\frac{1}{2} (\sigma \sigma^*)(t, x, \mu_t) \nabla \cdot \nabla h_i(x) + b(t, x, \mu_t) \cdot \nabla h_i(x) \right) \mu_t(dx) \Big) \\ (3.2) \quad &= \delta_{\mu_t} \left(\left\langle \frac{1}{2} (\sigma \sigma^*)(t, \cdot) \nabla + b(t, \cdot), \nabla^{\mathcal{P}} F(\cdot) \right\rangle_{L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \cdot)} \right), \end{aligned}$$

where $(\sigma\sigma^*)(t, \mu_t)$, $b(t, \mu_t)$ denote the maps

$$\begin{aligned}\mathbb{R}^d \ni x &\rightarrow \sigma\sigma^*(t, x, \mu_t), \\ \mathbb{R}^d \ni x &\mapsto b(t, x, \mu_t),\end{aligned}$$

which by assumption (2.5) (as part of the definition of the solution to (1.3)) are μ_t -integrable. Furthermore, here we set $\delta_{\mu_t}(g(\cdot)) := \int g(\nu)\delta_{\mu_t}(d\nu) = g(\mu_t)$ for a Borel measurable map $g: \mathcal{P} \rightarrow \mathbb{R}$. Hence we obtain

$$(3.3) \quad \partial_t \delta_{\mu_t}(F) = \delta_{\mu_t}(\langle \frac{1}{2}(\sigma\sigma^*)(t, \cdot)\nabla + b(t, \cdot), \nabla^{\mathcal{P}} F(\cdot) \rangle_{L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \cdot)}) \quad \text{for all } F \in \mathcal{F}C_b^2(\mathcal{P}),$$

which clearly is a linear equation for δ_{μ_t} , $t \geq s$, in $\mathcal{M}(\mathcal{P})$, i.e. the space of all bounded variation measures on $\mathcal{P}$. Vice versa, by the above derivation, (3.3) implies (1.3) by just taking $F_l \in \mathcal{F}C_b^2(\mathcal{P})$, $l \in \mathbb{N}$, such that for $l \in \mathbb{N}$

$$F(\mu) = f_l(\mu(h)), \quad h \in C_b^2(\mathbb{R}^d),$$

and $f_l \in C_b^1(\mathbb{R}^1)$ such that $\frac{d}{dx} f_l \rightarrow 1$ as $l \rightarrow \infty$. Hence we have proved the following result.

PROPOSITION 3.1. *A weakly continuous $\mathcal{P}$ -valued path $(\mu_t)_{t \geq s}$ satisfies the nonlinear F-P equation (1.3) in the sense of Definition 2.2(1), if and only if the $\mathcal{P}$ -valued path $(\delta_{\mu_t})_{t \geq s}$ satisfies the linear first order F-P equation (3.3).*

Now we proceed in the canonical way: For $\zeta \in \mathcal{P}$ let us denote a solution $(\mu_t)_{t \geq s}$ to (1.3) with initial condition $\mu_s = \zeta$ by $\mu(t, \zeta)$, $t \geq s$. Then consider $\mathcal{P}$ equipped with the σ -algebra $\tilde{\mathcal{B}}$ generated by the maps

$$\mathcal{P} \ni \zeta \mapsto \mu(t, \zeta) \in \mathcal{P}, \quad t \geq s,$$

where the image space is considered with the Borel σ -algebra $\mathcal{B}(\mathcal{P})$. Then for any probability measure Γ on $(\mathcal{P}, \tilde{\mathcal{B}})$ define

$$\Gamma_t := \Gamma \circ \mu(t, \cdot)^{-1}.$$

Then by (3.2) for all $F \in \mathcal{F}C_b^2(\mathcal{P})$

$$\begin{aligned}(3.4) \quad & \frac{d}{dt} \int F(\mu) \Gamma_t(d\mu) \\ &= \int \frac{d}{dt} F(\mu(t, \zeta)) \Gamma(d\zeta) \\ &= \int \langle \frac{1}{2}(\sigma\sigma^*)(t, \mu)\nabla + b(t, \mu), \nabla^{\mathcal{P}} F(\mu) \rangle_{L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)} \Gamma_t(d\mu),\end{aligned}$$

and rewriting this in the weak sense (with test function space $\mathcal{F}C_b^2(\mathcal{P})$, see Definition 2.2(4)) we obtain

$$(3.5) \quad \frac{d}{dt} \Gamma_t = \tilde{\mathbf{L}}_t \Gamma_t, \quad \Gamma_0 = \Gamma,$$

where $\tilde{\mathbf{L}}_t$ is defined in (1.7). Whether the above σ -algebra $\tilde{\mathcal{B}}$ on $\mathcal{P}$ coincides with $\mathcal{B}(\mathcal{P})$ has to be checked in every particular case and is, of course, fulfilled if the solution to (1.2) is continuous in its initial condition ζ with respect to a suitable topology, as is e.g. the case for our main examples below.

REMARK 3.2. By the product rule (3.5) implies that for every Γ as above, $T \in (0, \infty)$ and for all $F \in \mathcal{F}C_b^2(\mathcal{P})$, $G \in C^1([0, T]; \mathbb{R}^d)$ with $G(T) = 0$, and $\tilde{F}(t, \mu) := G(t)F(\mu)$, $(t, \mu) \in [0, T] \times \mathcal{P}$

$$\int_0^T \int_{\mathcal{P}} \left(\frac{\partial}{\partial t} + \tilde{\mathbf{L}}_t \right) \tilde{F} \, d\Gamma_t \, dt = - \int \tilde{F}(0, \mu) \, \Gamma(d\mu).$$

Letting $\tilde{\mathcal{F}}$ denote the linear space of all such functions $\tilde{F}$, we obtain that for all nonnegative $\tilde{F} \in \tilde{\mathcal{F}}$

$$\int_0^T \int_{\mathcal{P}} \left(\frac{\partial}{\partial t} + \tilde{\mathbf{L}}_t \right) \tilde{F} \, d\Gamma_t \, dt \leq 0,$$

i.e. the operator $\frac{\partial}{\partial t} + \tilde{\mathbf{L}}_t$ with domain $\tilde{\mathcal{F}}$ is dissipative on $L^1([0, T] \times \mathcal{P}, \Gamma_t dt)$, hence in particular closable. So, by the above we have a means to construct a whole class of probability measures on $[0, T] \times \mathcal{P}$ for which $(\frac{\partial}{\partial t} + \tilde{\mathbf{L}}_t, \tilde{\mathcal{F}})$ is dissipative (hence closable) on the corresponding L^1 space and $\tilde{\mathbf{L}}_t$ is given by a time dependent vector field in the tangent bundle $(L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu))_{\mu \in \mathcal{P}}$ over $\mathcal{P}$, determined by b and σ , where b, σ are arbitrary as in (2.1).

4. Correspondences and their consequences. Consider the situation of Section 2.

4.1. Correspondence of (1.10) and (2.13) and weak well-posedness for McKean-Vlasov SDEs.

THEOREM 4.1. Let $(\mathcal{P}_0, \mathcal{A})$, $(\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}})$ be two pairs having property **(P)**. Then (1.10) is well-posed in $((\mathcal{P}_0, \mathcal{A}), (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}}))$ (in the sense of Definition 2.2(2)) if and only if (2.13) is well-posed in $((\mathcal{P}_0, \mathcal{A}), (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}}))$ (in the sense of Definition 2.4(ii)). In this case $\nu_{s,t}^{\zeta, \theta} = \mathcal{L}_{X_{s,t}^{\zeta, \theta}}$ for all $(s, \theta, \zeta) \in [0, \infty) \times \tilde{\mathcal{P}}_0 \times \mathcal{P}_0$, $t \in [s, \infty)$.

PROOF. This is an immediate consequence of Remark 2.5 and Lemma 2.12 in [33]. The latter lemma is indeed applicable by property **(P)** above. $\square$

COROLLARY 4.2. Let $\sigma = \bar{\sigma}$, $b = \bar{b}$ and let $(\mathcal{P}_0, \mathcal{A})$ be a pair with property **(P)**. If (1.10) is well-posed in $((\mathcal{P}_0, \mathcal{A}), (\mathcal{P}_0, \mathcal{A}))$ then (1.8) is weakly well-posed in $(\mathcal{P}_0, \mathcal{A})$.

PROOF. This is a direct consequence of Theorem 4.1. $\square$

4.2. Correspondence of (1.10) and its linearization (1.13). In this section we modify the situation considered in Appendix A in the following way. Instead of $\mathcal{P}$, curves $\gamma_\phi^\mu(t)$, $t \geq 0$, $\mu \in \mathcal{P}$, $\phi \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)$, (see (A.2) below), test functions $\mathcal{F} := \mathcal{F}C_b^2(\mathcal{P})$ (see (2.3)) and resulting tangent bundle $(L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu))_{\mu \in \mathcal{P}}$ and gradient $\nabla^{\mathcal{P}}$ (see (A.5) below), we consider $\mathbb{R}^d \times \mathcal{P}$, curves $(\gamma^x(t), \gamma_\phi^\mu(t))$, $t \geq 0$, $(x, \mu) \in \mathbb{R}^d \times \mathcal{P}$, with γ^x any smooth curve in $\mathbb{R}^d$ with $\gamma^x(0) = x$ and γ_ϕ^μ as in (A.2) and test functions $\mathcal{F} = \mathcal{C}$ (see (2.4)). The resulting tangent bundle and gradient (using the procedure presented in Appendix A) one obtains is $(\mathbb{R}^d \oplus L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu))_{(x, \mu) \in \mathbb{R}^d \times \mathcal{P}}$ and

$$\nabla^{\mathbb{R}^d \times \mathcal{P}} G(x, \mu) = F(\mu) \nabla h_0(x) + h_0(x) \nabla^{\mathcal{P}} F(\mu), \quad (x, \mu) \in \mathbb{R}^d \times \mathcal{P},$$

where $G \in \mathcal{C}$, $G(x, \mu) = h_0(x)F(\mu)$.

THEOREM 4.3. (i) Let $s \geq 0$ and $(\mu_t)_{t \geq s}$, $(\nu_t)_{t \geq s} \in C([s, \infty) \rightarrow \mathcal{P})$. Then $(\mu_t, \nu_t)_{t \geq s}$ solves (1.10) if and only if $\Lambda_t := \nu_t \times \delta_{\mu_t}$, $t \geq s$, solves (1.13).

- (ii) Let $s \geq 0$ and let $\mathcal{P}_0, \tilde{\mathcal{P}}_0 \in \mathcal{B}(\mathcal{P})$ such that for each $\zeta \in \mathcal{P}_0$ there exists a solution $(\mu_{s,t}^\zeta)_{t \geq s}$ to (1.3) with $\mu_{s,s}^\zeta = \zeta$ and such that for each $\theta \in \tilde{\mathcal{P}}_0$ there exists a solution $(\nu_{s,t}^{\zeta,\theta})_{t \geq s}$ to (2.8) with $\mu_{s,r}^\zeta$ replacing μ_r , $r \geq s$, such that $\nu_{s,s}^{\zeta,\theta} = \theta$. Suppose that for every $t \in [s, \infty)$

$$\mathcal{P}_0 \ni \zeta \mapsto \mu_{s,t}^\zeta \in \mathcal{P}, \quad \mathcal{P}_0 \times \tilde{\mathcal{P}}_0 \ni (\zeta, \theta) \mapsto \nu_{s,t}^{\zeta,\theta} \in \mathcal{P}$$

are Borel measurable. Then for every $\Lambda \in \mathcal{P}(\mathcal{P}_0 \times \tilde{\mathcal{P}}_0)$ (= all probability measures on $\mathcal{P}_0 \times \tilde{\mathcal{P}}_0$)

$$\Lambda_{s,t}^\Lambda := \int_{\tilde{\mathcal{P}}_0 \times \mathcal{P}_0} (\nu_{s,t}^{\zeta,\theta} \times \delta_{\mu_{s,t}^\zeta}) \Lambda(d\theta, d\zeta), \quad t \in [0, \infty),$$

is a solution to (1.13).

PROOF. (i): The proof is entirely analogous to the proof of Proposition 3.1.

(ii): The proof follows by (i) since (1.13) is a linear Fokker-Planck equation (cf. the last paragraph in Section 3). $\square$

REMARK 4.4. We conjecture that if (1.10) is well-posed then so is (1.13). But for the proof one needs that any solution to (1.13) is of the type as Λ_t , $t \geq s$, is in assertion (i) of Theorem 4.3 above, which we did not succeed to prove yet.

4.3. *Markov property of weak solutions to McKean-Vlasov equations.* In this section we fix two pairs $(\mathcal{P}_0, \mathcal{A})$ and $(\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}})$ with property **(P)** and assume that

$$(4.1) \quad (1.10) \text{ is well-posed in } \left((\mathcal{P}_0, \mathcal{A}), (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}}) \right).$$

For $(s, \theta, \zeta) \in [0, \infty) \times \tilde{\mathcal{P}}_0 \times \mathcal{P}_0$ we define the laws

$$(4.2) \quad \begin{aligned} \mathbb{P}_{\zeta, (r, \theta)} &:= \mathbb{P} \circ (X_{r, \cdot}^{\mu_{s,r}^\zeta, \theta})^{-1}, \quad r \in [s, \infty), \\ \mathbb{P}_{\zeta, (r, x)} &:= \mathbb{P}_{\zeta, (r, \delta_x)}, \quad x \in \mathbb{R}^d, \text{ provided } \delta_x \in \tilde{\mathcal{P}}_0 \end{aligned}$$

on $C([r, \infty) \rightarrow \mathbb{R}^d)$, equipped with the σ -algebra $\mathcal{G}$ generated by all maps π_t , $t \geq r$, where π_t is the evaluation map at t . In addition, for $t \in [s, \infty)$ we define

$$(4.3) \quad \mathcal{G}_{s,t} := \sigma(\pi_u : u \in [s, t]).$$

Furthermore, we denote the corresponding expectations by $\mathbb{E}_{\zeta, (s, \theta)}$.

As mentioned before, in general it is not possible to prove uniqueness of linear (or more so, non-linear) Fokker-Planck equations for all initial probability measures θ , in particular not for Dirac measures. Therefore, we only assume a restricted well-posedness in (4.1). Hence (see [33, Lemma 2.12]) also the martingale problems corresponding to linear Fokker-Planck equations are only restricted well-posed. Therefore, the standard fact, that well-posedness (i.e. for all Dirac, hence all probability measures) implies that the corresponding family of probability measures $\mathbb{P}_x$ (= solution with initial marginal δ_x), $x \in \mathbb{R}^d$, form a Markov process, is not applicable. Nevertheless, we shall prove that under condition (4.1) we have the Markov property for our laws $\mathbb{P}_{\zeta, (s, \theta)}$ defined above and as a consequence also for the laws of the solutions to the McKean-Vlasov equation (1.8). And this holds just assuming the integrability conditions (2.5) and (2.7) on our coefficients.

As preparation for fixed $(s, \theta, \zeta) \in [0, \infty) \times \tilde{\mathcal{P}}_0 \times \mathcal{P}_0$ and $r \in [0, \infty)$ we disintegrate the measure $\mathbb{P}_{\zeta, (r, \nu_{s,r}^{\zeta, \theta})}$ with $\nu_{s,r}^{\zeta, \theta}$ as in Definition 2.2(2)(b) with respect to the map $\pi_r: C([r, \infty) \rightarrow \mathbb{R}^d) \rightarrow \mathbb{R}^d$ as follows

$$(4.4) \quad \mathbb{P}_{\zeta, (r, \nu_{s,r}^{\zeta, \theta})}(dw) = p(x, dw) \nu_{s,r}^{\zeta, \theta}(dx),$$

where p is a probability kernel from $\mathbb{R}^d$ to $C([r, \infty) \rightarrow \mathbb{R}^d)$ such that $p(x, \{\pi_r = x\}) = 1$ for all $x \in \mathbb{R}^d$. The existence of such a kernel follows by standard results on disintegration of measures.

THEOREM 4.5. *Assume that (4.1) holds, let p be as in (4.4) and let $(s, \theta, \zeta) \in [0, \infty) \times \tilde{\mathcal{P}}_0 \times \mathcal{P}_0$ and $r \in [s, \infty)$. Then for every $g \in \mathcal{B}_b(\mathbb{R}^d)$ and $t \in [r, \infty)$*

$$(4.5) \quad \mathbb{E}_{\zeta, (s, \theta)}[g(\pi_t) | \mathcal{G}_{s,r}] = \int g(\pi_t(w)) p(\pi_r, dw) \quad \mathbb{P}_{\zeta, (s, \theta)}\text{-a.e.}$$

PROOF. Since for all $\varphi \in C_0^\infty(\mathbb{R}^d)$

$$\varphi(\pi_t) - \varphi(\pi_r) - \int_r^t L_{u, \mu_{s,u}^\zeta} \varphi \, du, \quad t \geq r,$$

is a $(\mathcal{G}_{r,t})_{t \geq r}$ -martingale under $\mathbb{P}_{\zeta, (r, \nu_{s,r}^{\zeta, \theta})}$, it is elementary to check that for $\nu_{s,r}^{\zeta, \theta}$ -a.e. $x \in \mathbb{R}^d$ this is also true under $p(x, dw)$ defined in (4.4). Hence this is also true for the measure

$$(4.6) \quad \mathbb{P}_\rho(dw) := \int p(x, dw) \rho(x) \, d\nu_{s,r}^{\zeta, \theta}(dx)$$

for every probability density ρ with respect to $\nu_{s,r}^{\zeta, \theta}$, i.e. $\mathbb{P}_\rho$ satisfies the martingale problem on $C([r, \infty) \rightarrow \mathbb{R}^d)$ for $L_{u, \mu_{s,u}^\zeta}$ with initial measure $\rho \nu_{s,r}^{\zeta, \theta}$. The latter follows from the fact that $p(x, dw)$ is supported by $\{\pi_r = x\}$.

Now let $n \in \mathbb{N}$, $s \leq u_1 \leq \dots \leq u_n \leq r$, $h \in \mathcal{B}_b((\mathbb{R}^d)^n)$, $h \geq 0$ and $g \in \mathcal{B}_b(\mathbb{R}^d)$. Define the factorized conditional expectation

$$\rho(x) := c \mathbb{E}_{\zeta, (s, \theta)}[h(\pi_{u_1}, \dots, \pi_{u_n}) | \pi_r] |_{\pi_r = x}, \quad x \in \mathbb{R}^d,$$

where $c \in (0, \infty)$ so that its integral w.r.t. $\nu_{s,r}^{\zeta, \theta}$ is equal to 1. ρ is uniquely defined $\nu_{s,r}^{\zeta, \theta}$ -a.e. Now consider $\mathbb{P}_\rho$ defined in (4.6) for this ρ . Furthermore, let

$$\bar{\rho} := c h(\pi_{u_1}, \dots, \pi_{u_n}).$$

Then $\bar{\rho}$ is a probability density w.r.t. $\mathbb{P}_{\zeta, (s, \theta)}$. We denote the image measure of $\bar{\rho} \cdot \mathbb{P}_{\zeta, (s, \theta)}$ under the natural projection of $C([s, \infty) \rightarrow \mathbb{R}^d)$ onto $C((r, \infty) \rightarrow \mathbb{R}^d)$ by $\mathbb{P}_{\bar{\rho}}$.

Claim: $\mathbb{P}_{\bar{\rho}} = \mathbb{P}_\rho$.

It is easy to check that also $\mathbb{P}_{\bar{\rho}}$ satisfies the above martingale problem. Furthermore, also under $\mathbb{P}_{\bar{\rho}}$ the law of π_r is equal to $\rho \cdot \nu_{s,r}^{\zeta, \theta}$. Therefore, by (4.1) and [33, Lemma 2.12] the claim follows.

By the Claim we have for $g \in \mathcal{B}_b(\mathbb{R}^d)$

$$\begin{aligned}
& \mathbb{E}_{\zeta, (s, \theta)}[h(\pi_{u_1}, \dots, \pi_{u_n})g(\pi_t)] \\
&= \frac{1}{c} \int g(\pi_t) \, d\mathbb{P}_{\bar{\rho}} \\
&= \frac{1}{c} \int g(\pi_t) \, d\mathbb{P}_{\rho} = \frac{1}{c} \mathbb{E}_{\zeta, (s, \theta)} \left[\int_{\mathbb{R}^d} g(\pi_t(w)) \, p(\pi_r, dw) \, \rho(\pi_r) \right] \\
&= \mathbb{E}_{\zeta, (s, \theta)} \left[\int_{\mathbb{R}^d} g(\pi_t(w)) \, p(\pi_r, dw) \, h(\pi_{u_1}, \dots, \pi_{u_n}) \right].
\end{aligned}$$

Now (4.5) follows by a monotone class argument. $\square$

COROLLARY 4.6. *Let $\sigma = \bar{\sigma}$, $b = \bar{b}$ and assume that (4.1) holds with $\mathcal{P}_0 = \tilde{\mathcal{P}}_0$. Then for every $(s, \zeta) \in [0, \infty) \times \mathcal{P}_0$ the law $\mathbb{P}_{\zeta, (s, \zeta)}$ of the solution of the McKean-Vlasov equation (1.8) started from s at ζ is Markov, i.e. satisfies (4.5) with p as in (4.4).*

REMARK 4.7. *We note that in the situation of Corollary 4.6 we obviously have that $\nu_{s,t}^{\zeta, \zeta} = \mu_{s,t}^{\zeta}$ for all $0 \leq s \leq t$ and $\zeta \in \mathcal{P}_0$.*

4.4. *The Markov process on $\mathbb{R}^d \times \mathcal{P}$.* In this and the next subsection we fix a pair $(\mathcal{P}_0, \mathcal{A})$ with property **(P)** (see Section 2) and set

$$(4.7) \quad \tilde{\mathcal{P}}_0 := \mathcal{P}, \quad \tilde{\mathcal{A}} := C([0, \infty) \rightarrow \mathcal{P}),$$

so obviously $(\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}})$ is also a pair with property **(P)**. We also assume throughout this and the next subsection

$$(4.8) \quad (1.10) \text{ is well-posed in } ((\mathcal{P}_0, \mathcal{A}), (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}})) \text{ with } (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}}) \text{ defined as in (4.7).}$$

For $(s, x, \zeta) \in [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_0$, as in Definition 2.2(2)(a), (b), we denote the corresponding solutions by $(\mu_{s,t}^{\zeta}, \nu_{s,t}^{\zeta, \delta_x})_{t \geq s}$. For $x \in \mathbb{R}^d$, $\zeta \in \mathcal{P}_0$, $s \in [0, \infty)$ define for $t \in [s, \infty)$

$$(4.9) \quad \mathbf{P}_{s,t}(x, \zeta; dy d\mu) = (\nu_{s,t}^{\zeta, \delta_x} \times \delta_{\mu_{s,t}^{\zeta}})(dy d\mu),$$

i.e. for $G : \mathbb{R}^d \times \mathcal{P}_0 \rightarrow [0, \infty)$ Borel-measurable

$$\int_{\mathbb{R}^d} \int_{\mathcal{P}_0} G(y, \mu) \mathbf{P}_{s,t}(x, \zeta; dy d\mu) = \int_{\mathbb{R}^d} G(y, \mu_{s,t}^{\zeta}) \nu_{s,t}^{\zeta, \delta_x}(dy).$$

Then by Theorem 4.3(i) it follows that $\mathbf{P}_{s,t}(x, \zeta; dy d\mu)$ solves (1.13) starting from time s and furthermore

$$(4.10) \quad \mathbf{P}_{s,s}(x, \zeta; dy d\mu) = (\delta_x \times \delta_{\zeta})(dy d\mu).$$

PROPOSITION 4.8. *Suppose (4.8) holds. Then the family of probability measures $\mathbf{P}_{s,t}(x, \zeta; \cdot)$, $0 \leq s \leq t$, $(x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_0$, is a Markov transition kernel on $\mathbb{R}^d \times \mathcal{P}_0$, i.e., it satisfies the following properties:*

(C₁) $\mathbf{P}_{s,t}(\cdot; A)$ is Borel-measurable in (x, ζ) for any $0 \leq s \leq t$, $A \in \mathcal{P}(\mathbb{R}^d \times \mathcal{P}_0)$ and $\mathbf{P}_{s,s}(x, \zeta; \cdot) = \delta_{(x, \zeta)}$, the Dirac measure at (x, ζ) , for any $s \geq 0$ and $(x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_0$.

(C₂) The Chapman-Kolmogorov equations hold, i.e. for all $0 \leq s < r < t$ and $(x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_0$

$$(4.11) \quad \mathbf{P}_{s,t}(x, \zeta; \cdot) = \int_{\mathbb{R}^d \times \mathcal{P}_0} \mathbf{P}_{r,t}(y, \mu; \cdot) \mathbf{P}_{s,r}(x, \zeta; dy, d\mu).$$

PROOF. (C₁) is obvious, so it only remains to prove the Chapman-Kolmogorov equations. For $0 \leq s \leq r \leq t$ and $(x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_0$ the right hand side of (4.11) is equal to

$$\begin{aligned} & \int_{\mathbb{R}^d} \mathbf{P}_{r,t}(y, \mu_{s,r}^\zeta; \cdot) \nu_{s,r}^{\zeta, \delta_x}(dy) \\ &= \int_{\mathbb{R}^d} \left(\nu_{r,t}^{\mu_{s,r}^\zeta, \delta_y} \times \delta_{\mu_{s,r}^\zeta} \right) \nu_{s,r}^{\zeta, \delta_x}(dy) \\ &= \left(\int_{\mathbb{R}^d} \nu_{r,t}^{\mu_{s,r}^\zeta, \delta_y} \nu_{s,r}^{\zeta, \delta_x}(dy) \right) \times \delta_{\mu_{s,t}^\zeta} \\ &= \nu_{s,t}^{\zeta, \delta_x} \times \delta_{\mu_{s,t}^\zeta} = \mathbf{P}_{s,t}(x, \zeta; \cdot) \end{aligned}$$

where we used the well-posedness of (1.10), more precisely Definition 2.2(2)(a), (b) in the second and third equality respectively. $\square$

A Markov transition kernel on $\mathbb{R}^d \times \mathcal{P}_0$ determines a family of Markov operators $\{\mathbf{P}_{s,t} : 0 \leq s \leq t\}$ on $\mathcal{B}_b(\mathbb{R}^d \times \mathcal{P}_0)$, the Banach space of bounded Borel-measurable functions on $\mathbb{R}^d \times \mathcal{P}_0$:

$$(4.12) \quad \mathbf{P}_{s,t}f(x, \zeta) := \int_{\mathbb{R}^d \times \mathcal{P}_0} f(\xi) \mathbf{P}_{s,t}(x, \zeta; d\xi), \quad (x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_0, f \in \mathcal{B}_b(\mathbb{R}^d \times \mathcal{P}_0).$$

Conditions (C₁) and (C₂) are equivalent to $\mathbf{P}_{s,s}f = f$ and the semigroup property

$$\mathbf{P}_{s,t} = \mathbf{P}_{s,r} \mathbf{P}_{r,t}, \quad 0 \leq s \leq r \leq t.$$

By (4.8) and Theorem 4.1 we have that (2.13) is well-posed in $((\mathcal{P}_0, \mathcal{A}), (\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}}))$, with $(\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}})$ as in (4.7). For $(s, x, \zeta) \in [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_0$, as in Definition 2.4(ii) we denote the corresponding solutions by $(X_{s,t}^{\zeta, \delta_x}, \mu_{s,t}^\zeta)_{t \geq s}$ on $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ with $(\mathcal{F}_t)$ -Brownian motion $(W_t)_{t \geq 0}$ (see Section 2). We note that the stochastic basis and the Brownian motion depend on (s, x, ζ) , but for simplicity we do not express this in the notation.

Our next aim is to prove that the laws of $(X_{s,t}^{\zeta, \delta_x}, \mu_{s,t}^\zeta)_{t \geq s}$, $(s, x, \zeta) \in [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_0$, form a Markov process with Markov transition kernel, $\mathbf{P}_{s,t}$, $0 \leq s \leq t$, defined in (4.9). For $(s, \theta, \zeta) \in [0, \infty) \times \mathcal{P} \times \mathcal{P}_0$ we define the laws

$$(4.13) \quad \begin{aligned} \mathbb{P}_{(s, \theta, \zeta)} &:= \mathbb{P} \circ (X_{s, \cdot}^{\zeta, \theta}, \mu_{s, \cdot}^\zeta)^{-1} = \left(\mathbb{P} \circ (X_{s, \cdot}^{\zeta, \theta})^{-1} \right) \times \delta_{\mu_{s, \cdot}^\zeta} = \mathbb{P}_{\zeta, (s, \theta)} \times \delta_{\mu_{s, \cdot}^\zeta} \quad \text{and} \\ \mathbb{P}_{(s, x, \zeta)} &:= \mathbb{P}_{(s, \delta_x, \zeta)}, \quad x \in \mathbb{R}^d, \end{aligned}$$

on $C([s, \infty) \rightarrow \mathbb{R}^d \times \mathcal{P}_0) = C([s, \infty) \rightarrow \mathbb{R}^d) \times C([s, \infty) \rightarrow \mathcal{P}_0)$. We denote the canonical projection on this path space by π_t^0 , $t \in [s, \infty)$, and equip it with the σ -algebra $\mathcal{G}^0$ generated by these projections. In addition, we define for $t \in [s, \infty)$

$$\mathcal{G}_{s,t}^0 := \sigma(\pi_u^0 : u \in [s, t]).$$

Furthermore, we denote the corresponding expectations by $\mathbb{E}_{(s, \theta, \zeta)}$.

THEOREM 4.9. Suppose that (4.8) holds. Let $(\mathbf{P}_{s,t})_{t \geq s \geq 0}$ be in (4.12). Then for any $(s, x, \zeta) \in [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_0$, $t \in [s, \infty)$, and $r \in [s, t]$:

- (i) $\mathbb{P}_{(s,x,\zeta)} \circ \pi_t^{-1} = \mathbf{P}_{s,t}(x, \zeta; \cdot)$.
- (ii) $\mathbb{P}_{(s,x,\zeta)}$ -a.s. we have for every $A \in \mathcal{B}(\mathbb{R}^d \times \mathcal{P}_0)$

$$(4.14) \quad \mathbb{P}_{(s,x,\zeta)}[\pi_t^0 \in A | \mathcal{G}_r^0] = \mathbb{P}_{(r,\pi_r^0)}[\pi_t^0 \in A] = \mathbf{P}_{r,t}(\pi_r^0; A),$$

i.e., $\mathbb{P}_{(s,x,\zeta)}$, $(s, x, \zeta) \in [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_0$, form a Markov process with Markov transition kernel $\mathbf{P}_{s,t}$, $0 \leq s \leq t$.

PROOF. (i) is obvious from the definitions.

(ii) Since by our assumptions we have weak uniqueness for the second equation in (2.13), by [32, Theorem 6.2.2] and Theorem 4.1 above we know that for every $s \leq r \leq t$, $g \in \mathcal{B}_b(\mathbb{R}^d)$

$$(4.15) \quad \mathbb{E}_{\zeta,(s,x)}[g(\pi_t) | \mathcal{G}_{s,r}] = \mathbb{E}_{\zeta,(r,\pi_r)}[g(\pi_t)] = \int_{\mathbb{R}^d} g \, d\nu_{r,t}^{\mu_{s,r}^{\zeta}, \delta_{\pi_r}} \quad \mathbb{P}_{\zeta,(s,x)}\text{-a.s.}$$

For $n \in \mathbb{N}$; $s \leq u_1 \leq \dots \leq u_n \leq r$, and $H \in \mathcal{B}((\mathbb{R}^d \times \mathcal{P}_0)^n)$, we have that $\mathbb{P}_{(s,x,\zeta)}$ -a.s. with $G := 1_A$

$$\begin{aligned} & \mathbb{E}_{(s,x,\zeta)}[H(\pi_{u_1}^0, \dots, \pi_{u_n}^0)G(\pi_t^0)] \\ &= \mathbb{E}[H((X_{s,u_1}^{\zeta, \delta_x}, \mu_{s,u_1}^{\zeta}), \dots, (X_{s,u_n}^{\zeta, \delta_x}, \mu_{s,u_n}^{\zeta}))G(X_{s,t}^{\zeta, \delta_x}, \mu_{s,t}^{\zeta})] \\ &= \mathbb{E}_{\zeta,(s,x)}[H((\pi_{u_1}, \mu_{s,u_1}^{\zeta}), \dots, (\pi_{u_n}, \mu_{s,u_n}^{\zeta}))G(\pi_t, \mu_{s,t}^{\zeta})] \\ &\stackrel{(4.15)}{=} \mathbb{E}[H((X_{s,u_1}^{\zeta, \delta_x}, \mu_{s,u_1}^{\zeta}), \dots, (X_{s,u_n}^{\zeta, \delta_x}, \mu_{s,u_n}^{\zeta}))\mathbb{E}_{\zeta,(r,X_{s,r}^{\zeta, \delta_x})}[G(\pi_t, \mu_{s,t}^{\zeta})]]. \end{aligned}$$

But since by the well-posedness of the first equation in (2.13) in $(\mathcal{P}_0, \mathcal{A})$ we have the flow property

$$\mu_{s,t}^{\zeta} = \mu_{r,t}^{\mu_{s,r}^{\zeta}},$$

(4.2) and (4.13) imply that for $\mathbb{P}$ -a.e. $\omega \in \Omega$

$$\begin{aligned} & \mathbb{E}_{\zeta,(r,X_{s,r}^{\zeta, \delta_x}(\omega))}[G(\pi_t, \mu_{s,t}^{\zeta})] \\ &= \mathbb{E}[G(X_{r,t}^{\mu_{s,r}^{\zeta}, \delta_{X_{s,r}^{\zeta, \delta_x}(\omega)}}, \mu_{r,t}^{\mu_{s,r}^{\zeta}})] \\ &= \mathbb{E}_{(r,X_{s,r}^{\zeta, \delta_x}(\omega), \mu_{s,r}^{\zeta})}[G(\pi_t^0)]. \end{aligned}$$

Hence altogether

$$\begin{aligned} & \mathbb{E}_{(s,x,\zeta)}[H(\pi_{u_1}^0, \dots, \pi_{u_n}^0)G(\pi_t^0)] \\ &= \mathbb{E}_{(s,x,\zeta)}[G(\pi_{u_1}^0, \dots, \pi_{u_n}^0)\mathbb{E}_{(r,\pi_r^0)}[G(\pi_t^0)]]]. \end{aligned}$$

Hence the first inequality in (4.14) follows by a monotone class argument and the second is obvious by (i). $\square$

We note that since (4.8) is a stronger assumption than (4.1) we get a more explicit way to formulate the Markov property for the laws of the (unique) weak solution to the McKean-Vlasov equation (1.8) (see (4.17), (4.18) below).

$$(4.16) \quad b = \bar{b}, \quad \sigma = \bar{\sigma}.$$

Then for $(s, \zeta) \in [0, \infty) \times \mathcal{P}_0$ by (4.8) we have that (using our notations above) for every $\theta \in \mathcal{P}$ there exists a unique in law weak solution $(X_{s,t}^{\zeta, \theta})_{t \geq s}$ of the second equation in (2.13) with b, σ replacing $\bar{b}$ and $\bar{\sigma}$ respectively on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with normal filtration $(\mathcal{F}_t)_{t \geq 0}$ and $(\mathcal{F}_t)$ -Brownian motion $(W_t)_{t \geq 0}$ (see Section 2). Again all these three quantities depend on (s, θ, ζ) , but for simplicity we do not express this in the notation. In particular, for $\theta := \zeta$ we obtain a weak solution to (1.8) with marginal law ζ at time s .

LEMMA 4.10. *For all $(s, \theta, \zeta) \in [0, \infty) \times \mathcal{P} \times \mathcal{P}_0$ we have*

$$\mathbb{P}_{\zeta, (s, \theta)} = \int_{\mathbb{R}^d} \mathbb{P}_{\zeta, (s, x)} \theta(dx).$$

PROOF. The proof is standard, but we repeat the argument here: The two probability measures in the assertion solve the martingale problem with initial condition ζ for the Kolmogorov operator $L_{t, \zeta}$ defined in (1.2) in the sense of [32]. But by (4.8) and Theorem 4.1 this martingale problem has a unique solution. So, both measures coincide. $\square$

THEOREM 4.11. *Let (4.8) hold and let $(s, \zeta) \in [0, \infty) \times \mathcal{P}_0$. Then:*

- (i) *The unique weak solution to (1.8) with marginal law ζ at s has the Markov property, i.e. for every $s \leq r \leq t$, $g \in \mathcal{B}_b(\mathbb{R}^d)$,*

$$(4.17) \quad \mathbb{E}_{\zeta, (s, \zeta)}[g(\pi_t) | \mathcal{G}_{s, r}] = \mathbb{E}_{\zeta, (r, \pi_r)}[g(\pi_t)] = \int_{\mathbb{R}^d} g \, d\nu_{r, t}^{\mu_{s, r}^{\zeta}, \delta_{\pi_r}} \quad \mathbb{P}_{\zeta, (s, \zeta)}\text{-a.s.}$$

- (ii) *$\mathbb{P}_{(s, \zeta, \zeta)}$ is Markov. More precisely, for $s \leq r \leq t$ and $A \in \mathcal{B}(\mathbb{R}^d \times \mathcal{P}_0)$*

$$(4.18) \quad \mathbb{P}_{(s, \zeta, \zeta)}[\pi_t^0 \in A | \mathcal{G}_r^0] = \mathbf{P}_{r, t}(\pi_r^0; A) \quad \mathbb{P}_{(s, \zeta, \zeta)}\text{-a.s.}$$

PROOF. (i): The assertion immediately follows by (4.15) and Lemma 4.10.
(ii): As an easy consequence of Lemma 4.10 we get that

$$\mathbb{P}_{(s, \zeta, \zeta)} = \int \mathbb{P}_{(s, x, \zeta)} \zeta(dx).$$

Then the assertion immediately follows from Theorem 4.9(ii). $\square$

4.5. *Ergodicity.* We recall that we still assume (4.8) with $(\tilde{\mathcal{P}}_0, \tilde{\mathcal{A}})$ as in (4.7). In this subsection we assume additionally that for our coefficients from (2.1) we have

$$(4.19) \quad b, \bar{b}, \sigma \text{ and } \bar{\sigma} \text{ do not depend on } t \in [0, \infty).$$

Then due to (4.8) for our Markov transition kernel $\mathbf{P}_{s, t}$, $0 \leq s \leq t$, defined in (4.9) we have that $\mathbf{P}_{s, t} = \mathbf{P}_{0, t-s}$, i.e. it is time-homogeneous. Set

$$(4.20) \quad \mathbf{P}_t := \mathbf{P}_{0, t}, \quad t \geq 0.$$

Then by (C_2) this is a semigroup of probability kernels on $\mathbb{R}^d \times \mathcal{P}_0$ or equivalently (see (4.12)) of operators on $\mathcal{B}_b(\mathbb{R}^d \times \mathcal{P}_0)$. We recall that $\Lambda \in \mathcal{P}(\mathbb{R}^d \times \mathcal{P}_0)$ is called $(\mathbf{P}_t)$ -invariant if

$$(4.21) \quad \int_{\mathbb{R}^d \times \mathcal{P}_0} \mathbf{P}_t G \, d\Lambda = \int_{\mathbb{R}^d \times \mathcal{P}_0} G \, d\Lambda,$$

for all $G \in \mathcal{B}_b(\mathbb{R}^d \times \mathcal{P}_0)$, $t > 0$, and that $(\mathbf{P}_t)_{t>0}$ is called ergodic if there exists $\Lambda \in \mathcal{P}(\mathbb{R}^d \times \mathcal{P}_0)$ such that for all $\tilde{\Lambda} \in \mathcal{P}(\mathbb{R}^d \times \mathcal{P}_0)$ (or equivalently for all $\tilde{\Lambda} = \delta_x \times \delta_\zeta$, $(x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_0$)

$$(4.22) \quad \lim_{t \rightarrow \infty} \int_{\mathbb{R}^d \times \mathcal{P}_0} \mathbf{P}_t G d\tilde{\Lambda} = \int_{\mathbb{R}^d \times \mathcal{P}_0} G d\Lambda,$$

for all $G \in C_b(\mathbb{R}^d \times \mathcal{P}_0)$, where the latter denotes the set of all bounded continuous functions on $\mathbb{R}^d \times \mathcal{P}_0$. In this case Λ is then obviously the unique $(\mathbf{P}_t)$ -invariant measure. From the definition of $\mathbf{P}_t$ we now obtain the following characterization.

THEOREM 4.12. *Assume (4.8) and (4.19) hold. Then the following assertions are equivalent:*

- (i) $(\mathbf{P}_t)_{t>0}$ is ergodic.
- (ii) There exist $\nu_\infty \in \mathcal{P}$, $\mu_\infty \in \mathcal{P}_0$ such that for every $(x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_0$

$$\mu_{0,t}^\zeta \rightarrow \mu_\infty \text{ and } \nu_{0,t}^{\zeta, \delta_x} \rightarrow \nu_\infty \text{ weakly as } t \rightarrow \infty.$$

In this case μ_∞ and ν_∞ are uniquely determined and the unique $(\mathbf{P}_t)$ -invariant measure Λ is given by $\Lambda = \nu_\infty \times \delta_{\mu_\infty}$. Furthermore, if $b = \bar{b}$ and $\sigma = \bar{\sigma}$, then $\mu_\infty = \nu_\infty$.

PROOF. (ii) $\Rightarrow$ (i): This is easy to see from the definition of $\mathbf{P}_t$ in (4.20) and (4.9), and the unique $(\mathbf{P}_t)$ -invariant measure is $\Lambda := \nu_\infty \times \delta_{\mu_\infty}$.

(i) $\Rightarrow$ (ii): Let Λ be the $(\mathbf{P}_t)$ -invariant measure such that (4.22) holds. Let $\Pi_1: \mathbb{R}^d \times \mathcal{P}_0 \rightarrow \mathbb{R}^d$, and $\Pi_2: \mathbb{R}^d \times \mathcal{P}_0 \rightarrow \mathcal{P}_0$ be the canonical projections and define

$$\nu_\infty := \Lambda \circ \Pi_1^{-1}(\in \mathcal{P}) \text{ and } \Lambda_2 := \Lambda \circ \Pi_2^{-1}(\in \mathcal{P}(\mathcal{P}_0)).$$

Then for every $(x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_0$ as $t \rightarrow \infty$

$$(4.23) \quad \nu_{0,t}^{\zeta, \delta_x} \rightarrow \nu_\infty \text{ weakly in } \mathcal{P}$$

and

$$(4.24) \quad \delta_{\mu_{0,t}^\zeta} \rightarrow \Lambda_2 \text{ weakly in } \mathcal{P}(\mathcal{P}_0) \text{ (hence also in } \mathcal{P}(\mathcal{P}) \text{ by extending } \Lambda_2 \text{ by zero)}.$$

We claim that (4.24) implies $\Lambda_2 = \delta_{\mu_\infty}$ for some $\mu_\infty \in \mathcal{P}_0$. Combining this with (4.22) for $\tilde{\Lambda} = \zeta \times \delta_x$ and using (4.23), we conclude that $\Lambda = \nu_\infty \times \delta_{\mu_\infty}$ as desired.

To prove the claim, let $\mu_\infty \in \mathcal{P}$ be defined by

$$\mu_\infty(A) := \int_{\mathcal{P}_0} \mu(A) \Lambda_2(d\mu) = \int_{\mathcal{P}} \mu(A) \Lambda_2(d\mu), \quad A \in \mathcal{B}(\mathbb{R}^d),$$

where Λ_2 is extended to $\mathcal{P}$ by zero, i.e. $\Lambda_2(\mathcal{P} \setminus \mathcal{P}_0) = 0$. For any $h \in C_b(\mathbb{R}^d)$ we take $F(\mu) = \mu(h) := \int h d\mu$, $\mu \in \mathcal{P}$. Then $F \in C_b(\mathcal{P})$. Since $\delta_{\mu_{0,t}^\zeta} \rightarrow \Lambda_2$ weakly in $\mathcal{P}(\mathcal{P})$, we have

$$\lim_{t \rightarrow \infty} \mu_{0,t}^\zeta(h) = \lim_{t \rightarrow \infty} \delta_{\mu_{0,t}^\zeta}(F) = \Lambda_2(F) = \mu_\infty(h).$$

So, $\mu_{0,t}^\zeta \rightarrow \mu_\infty$ weakly in $\mathcal{P}$, and hence $\delta_{\mu_{0,t}^\zeta} \rightarrow \delta_{\mu_\infty}$ weakly in $\mathcal{P}(\mathcal{P})$ as $t \rightarrow \infty$. Combining this with (4.24) we prove $\Lambda_2 = \delta_{\mu_\infty}$ and $\mu_\infty \in \mathcal{P}_0$ since Λ_2 is supported on $\mathcal{P}_0$.

We note that for every $(\zeta, \theta) \in \mathcal{P}_0 \times \mathcal{P}$

$$\nu_{0,t}^{\zeta,\theta} = \int \nu_{0,t}^{\zeta,\delta_x} \theta(dx), \quad t \geq 0,$$

since the second equation in (1.10) is linear and well-posed. Hence from (ii) we have

$$\nu_{0,t}^{\zeta,\theta} \longrightarrow \nu_\infty \quad \text{weakly as } t \rightarrow \infty.$$

Now assume that $b = \bar{b}$ and $\sigma = \bar{\sigma}$. Then for all $\zeta \in \mathcal{P}_0$

$$\begin{aligned} \nu_\infty &= w - \lim_{t \rightarrow \infty} \nu_{0,t}^{\zeta,\zeta} \\ &= w - \lim_{t \rightarrow \infty} \mu_{0,t}^\zeta \\ &= \mu_\infty, \end{aligned}$$

where we used Remark 2.3 in the second equality. This completes the proof. $\square$

5. Application to nonlinear distorted Brownian motion. In this section we want to apply our results to the so-called nonlinear distorted Brownian motion (NLDBM) in which case the nonlinear Fokker-Planck equation (1.3) is a porous media equation perturbed by a nonlinear transport term. We shall give details below, but want to stress already now that in this case the solutions of (1.3) are absolutely continuous with respect to Lebesgue measure dx , if so is the initial condition, i.e. , $u(dx) = u(t, x)dx$, $t \geq 0$. Furthermore, in the case of NLDBM the coefficients b , σ in (2.1) (to be introduced below explicitly) depend "Nemytski type" on μ_t , more precisely for $(t, x, \mu) \in [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_0$, where

$$(5.1) \quad \mathcal{P}_0 := \left\{ \mu \in \mathcal{P} \mid u_\mu := \frac{d\mu}{dx} \in L^\infty(\mathbb{R}^d, dx) \right\},$$

we have

$$(5.2) \quad \begin{aligned} b(t, x, \mu) &= \tilde{b}\left(t, x, \frac{d\mu}{dx}(x)\right), \\ \sigma(t, x, \mu) &= \tilde{\sigma}\left(t, x, \frac{d\mu}{dx}(x)\right) \end{aligned}$$

for some Borel measurable functions

$$\tilde{b}: [0, \infty) \times \mathbb{R}^d \times \mathbb{R} \longrightarrow \mathbb{R}^d,$$

and

$$\tilde{\sigma}: [0, \infty) \times \mathbb{R}^d \times \mathbb{R} \longrightarrow \mathbb{R}^d \otimes \mathbb{R}^m.$$

Of course, for $x \in \mathbb{R}^d$ we have to choose the dx -version of $\frac{d\mu}{dx}$ in such a way that the maps

$$\begin{aligned} [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_0 \ni (t, x, \mu) &\longmapsto \tilde{b}\left(t, x, \frac{d\mu}{dx}(x)\right), \\ [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_0 \ni (t, x, \mu) &\longmapsto \tilde{\sigma}\left(t, x, \frac{d\mu}{dx}(x)\right) \end{aligned}$$

are Borel-measurable. But this is easily achieved by looking at the dx -version obtained by defining $\frac{d\mu}{dx} \equiv 0$ on the complement of its Lebesgue points. Below we shall always take this version without further mentioning it.

To introduce $\tilde{b}$ and $\tilde{\sigma}$ concretely in the case of NLDBM we consider maps $\beta: \mathbb{R} \rightarrow \mathbb{R}$, $D: \mathbb{R}^d \rightarrow \mathbb{R}^d$ and $b: \mathbb{R} \rightarrow \mathbb{R}$ satisfying the following hypotheses:

- (H) (i) $\beta \in C^1(\mathbb{R})$, $\beta(0) = 0$, $\gamma \leq \beta'(r) \leq \gamma_1$, $\forall r \in \mathbb{R}$, for $0 < \gamma < \gamma_1 < \infty$.
- (ii) $b \in C_b(\mathbb{R}) \cap C^1(\mathbb{R})$.
- (iii) $D \in C_b(\mathbb{R}^d; \mathbb{R}^d) \cap W^{1,\infty}(\mathbb{R}^d; \mathbb{R}^d)$.
- (iv) $D = -\nabla \Phi$, where $\Phi \in C^1(\mathbb{R}^d)$, $\Phi \geq 1$, $\lim_{|x|_d \rightarrow \infty} \Phi(x) = +\infty$ and there exists $m \in [2, \infty)$ such that $\Phi^{-m} \in L^1(\mathbb{R}^d)$.

A typical example for Φ is

$$(5.3) \quad \Phi(x) = C(1 + |x|^2)^\alpha, \quad x \in \mathbb{R}^d,$$

with $\alpha \in (0, \frac{1}{2}]$.

Then we have for the corresponding (t -independent) Kolmogorov operator (1.3) for $\mu \in \mathcal{P}_0$:

$$(5.4) \quad L_\mu h(x) := \frac{1}{2} \frac{\beta(u_\mu(x))}{u_\mu(x)} \Delta h(x) + b(u_\mu(x)) D(x) \cdot \nabla h(x), \quad x \in \mathbb{R}^d, \quad h \in C_0^2(\mathbb{R}^d),$$

where we set $\frac{\beta(0)}{0} := \beta'(0)$, i.e. compared to (1.3) we have for $(t, x, \mu) \in [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_0$

$$(5.5) \quad \begin{aligned} \sigma(t, x, \mu) &= \frac{\beta(u_\mu(x))}{u_\mu(x)} \text{Id}, \\ b(t, x, \mu) &= b(u_\mu(x)) D(x). \end{aligned}$$

Here Id denotes the identity matrix on $\mathbb{R}^d$. Hence (1.2) becomes an equation for density $\mu(t, \cdot) = \frac{d\mu_t}{dx}$ and then reads as

$$(5.6) \quad \partial_t u(t, x) = \frac{1}{2} \Delta \beta(u(t, x)) - \text{div}(D(x) b(u(t, x)) u(t, x)).$$

In this section we consider the case $b = \bar{b}$, $\sigma = \bar{\sigma}$. So, the first equation in (1.10) is just (5.6) and the second reads

$$(5.7) \quad \partial_t \nu_t = \frac{1}{2} \Delta \left(\frac{\beta(u(t, \cdot))}{u(t, \cdot)} \nu_t \right) - \text{div}(D(\cdot) b(u(t, \cdot)) \nu_t),$$

(as always in this paper) meant in the weak sense.

In particular, if $\zeta := u_0 dx \in \mathcal{P}_0$ is the initial condition for the solution of (5.6), then according to our notation in Definition 2.2(2)(b) the solution to (5.7) starting from $s \in [0, \infty)$ with $\theta \in \mathcal{P}_0$ is denoted by $\nu_{s,t}^{\zeta, \theta}$, $t \geq s$.

The corresponding McKean-Vlasov equation reads:

$$(5.8) \quad dX_t = b(\mathcal{L}_{X_t}(X_t)) D(X_t) dt + \frac{1}{2} \frac{\beta(\mathcal{L}_{X_t}(X_t))}{\mathcal{L}_{X_t}(X_t)} dW_t.$$

We recall that by Theorem 2.7 we have that (5.6) and (5.8) are equivalent (with $\mu_t = \mathcal{L}_{X_t}$, $t \geq 0$). (5.6) and consequently weak solutions of (5.8) have been analyzed in [7], [8] and the solution process to (5.8) has been named *nonlinear distorted Brownian motion*, because in the linear case, i.e. $b = \text{constant}$ and $\beta = \text{identity}$, (5.8) reduces to the SDE for distorted Brownian and (5.4) to the corresponding linear Kolmogorov operator.

Now let

$$\mathcal{A} := C([0, \infty) \rightarrow \mathcal{P}_0) \cap L^\infty([0, \infty) \times \mathbb{R}^d).$$

Then obviously condition **(P)** in Section 2 holds. The following is our main result on NLDBM.

THEOREM 5.1. *Assume (H)(i)-(iv) hold. Then:*

- (i) (5.8) is weakly well-posed in $(\mathcal{P}_0, \mathcal{A})$.
- (ii) Nonlinear distorted Brownian motion has the Markov property. More precisely, for every $(s, \zeta) \in [0, \infty) \times \mathcal{P}_0$ the law $\mathbb{P}_{\zeta, (s, \zeta)}$ of the (unique weak) solution of (5.8) started from s at ζ satisfies (4.5) with p as in (4.4).

PROOF. (i): By [7, Proposition 2.2] and [8, Theorems 2.1 and 3.1] in the case of NLDBM we have well-posedness for (1.10) in $(\mathcal{P}_0, \mathcal{A})$ (as defined above). Hence the assertion follows by Corollary 4.2 above.

(ii) is then a consequence of (i) and Corollary 4.6. $\square$

REMARK 5.2. *If additionally we make the following assumptions:*

- (H) (v) $b(r) \geq b_0 > 0$ for all $r \in \mathbb{R}$
- (vi) $\gamma_1 \Delta \Phi - b_0 |\nabla \Phi|^2 \leq 0$,

then it has been proved in [7] (see Theorems 6.1 and 6.5 in there) that (5.6) has a stationary solution which is unique in a slightly restricted class of probability densities in $L^1(\mathbb{R}^d, dx)$. Hence nonlinear distorted Brownian motion as a unique invariant measure in this class.

6. Exponential ergodicity of $\mathbf{P}_t$. In this section, we let $b(t, x, \mu) = b(x, \mu)$ and $\sigma(t, x, \mu) = \sigma(x, \mu)$ do not depend on t , and consider the exponential convergence of the Markov process generated by $\mathbf{L} = \mathbf{L}_t$ on the Wasserstein space

$$\mathcal{P}_2 := \left\{ \mu \in \mathcal{P} : \|\mu\|_2 := \left(\int_{\mathbb{R}^d} |x|^2 \mu(dx) \right)^{\frac{1}{2}} < \infty \right\}.$$

To this end, we will take $\mathcal{P}_0 = \tilde{\mathcal{P}}_0 = \mathcal{P}_2$, which is a Polish space under the Wasserstein distance

$$\mathbb{W}_2(\mu, \nu) := \inf_{\pi \in \mathcal{C}(\mu, \nu)} \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} |x - y|^2 \pi(dx, dy) \right)^{\frac{1}{2}},$$

where $\mathcal{C}(\mu, \nu)$ is the set of couplings for μ and ν .

We will need the following linear growth and monotone conditions.

- (A)** $b, \bar{b}, \sigma, \bar{\sigma}$ are continuous on $\mathbb{R}^d \times \mathcal{P}_2$ and there exist constants $K, \lambda, \kappa, \bar{\lambda}, \bar{\kappa} \geq 0$ such that for any $(x, \mu), (y, \nu) \in \mathbb{R}^d \times \mathcal{P}_2$, we have

$$(6.1) \quad \{|b| + \|\sigma\| + |\bar{b}| + \|\bar{\sigma}\|\}(x, \mu) \leq K(1 + |x| + \|\mu\|_2),$$

$$(6.2) \quad 2\langle b(x, \mu) - b(y, \nu), x - y \rangle + \|\sigma(x, \mu) - \sigma(y, \nu)\|_{HS}^2 \leq \kappa \mathbb{W}_2(\mu, \nu)^2 - \lambda |x - y|^2,$$

$$(6.3) \quad 2\langle \bar{b}(x, \mu) - \bar{b}(y, \nu), x - y \rangle + \|\bar{\sigma}(x, \mu) - \bar{\sigma}(y, \nu)\|_{HS}^2 \leq \bar{\kappa} \mathbb{W}_2(\mu, \nu)^2 - \bar{\lambda} |x - y|^2.$$

According to [28, Theorem 2.1], under **(A)** both the SDE (1.8) and the second equation in (2.13) are well-posed for initial distributions in $\mathcal{P}_2$. We denote $P_t^* \zeta = \mathcal{L}_{X_t}$ for X_t solving (1.8) with $\mathcal{L}_{X_0} = \zeta \in \mathcal{P}_2$. Then by Theorem 2.7, the coupled equation (2.13) is well-posed for

$$\mathcal{P}_0 = \tilde{\mathcal{P}}_0 = \mathcal{P}_2, \quad \mathcal{A} = \tilde{\mathcal{A}} = C([0, \infty) \rightarrow \mathcal{P}_2),$$

with $\mu_t = P_t^* \mu_0$ for $\mu_0 = \zeta, t \geq 0$. We denote by $(X_t^{\zeta, x})_{t \geq 0}$ the solution to the second equation in (2.13) starting at x . Then we have

$$(6.4) \quad dX_t^{\zeta, x} = \bar{b}(X_t^{\zeta, x}, P_t^* \zeta) dt + \bar{\sigma}(X_t^{\zeta, x}, P_t^* \zeta) dW_t, \quad X_0^{\zeta, x} = x.$$

Thus, Theorem 4.1 implies condition (4.8), so that by Proposition 4.8 we see that (4.9) gives a time-homogenous Markov transition kernel

$$(6.5) \quad \mathbf{P}_t(x, \zeta; \cdot) := \mathbf{P}_{0,t}(x, \zeta; \cdot) = \mathcal{L}_{X_t^{\zeta, x}} \times \delta_{P_t^* \zeta}, \quad t \geq 0, (x, \zeta) \in \mathbb{R}^d \times \mathcal{P}_2,$$

generated by $\mathbf{L}(= \mathbf{L}_t, t \geq 0)$. For any $\nu, \mu \in \mathcal{P}_2$, the distribution of the $\mathbf{L}$ -diffusion process (X_t, μ_t) at time t with $\mathcal{L}_{X_0} = \theta$ and $\mu_0 = \zeta$ is given by

$$(6.6) \quad \mathbf{P}_t(\theta, \zeta; \cdot) := \int_{\mathbb{R}^d} \mathbf{P}_t(x, \zeta; \cdot) \theta(dx) = \mathcal{L}_{X_t^{\zeta, \theta}} \times \delta_{P_t^* \zeta}, \quad \zeta, \theta \in \mathcal{P}_2, t \geq 0,$$

where $X_t^{\zeta, \theta}$ solves (2.14) for $s = 0$, i.e.

$$(6.7) \quad dX_t^{\zeta, \theta} = \bar{b}(X_t^{\zeta, \theta}, P_t^* \zeta) dt + \bar{\sigma}(X_t^{\zeta, \theta}, P_t^* \zeta) dW_t, \quad \mathcal{L}_{X_0^{\zeta, \theta}} = \theta.$$

Let $\mathbb{W}_2^\rho$ be the L^2 -Wasserstein distance induced by the following metric on $\mathbb{R}^d \times \mathcal{P}_2$:

$$\rho((x, \mu), (y, \nu)) := \sqrt{|x - y|^2 + \mathbb{W}_2(\mu, \nu)^2}.$$

Then for any two probability measures Λ_1, Λ_2 on $E := \mathbb{R}^d \times \mathcal{P}_2$,

$$\mathbb{W}_2^\rho(\Lambda_1, \Lambda_2)^2 := \inf_{\Pi \in \mathcal{C}(\Lambda_1, \Lambda_2)} \int_{E \times E} \rho((x, \mu), (y, \nu))^2 \Pi((dx, d\mu); (dy, d\nu)),$$

where $\mathcal{C}(\Lambda_1, \Lambda_2)$ is the set of all couplings of Λ_1 and Λ_2 .

THEOREM 6.1. *Assume **(A)**. If $\lambda > \kappa \geq 0$, then $\mathbf{P}_t$ has a unique invariant probability measure $\Lambda = \nu_\infty \times \delta_{\mu_\infty}$ for some $\mu_\infty, \nu_\infty \in \mathcal{P}_2$, such that for any $\zeta, \theta \in \mathcal{P}_2$,*

$$\mathbb{W}_2^\rho(\mathbf{P}_t(\theta, \zeta; \cdot), \Lambda)^2 \leq \mathbb{W}_2(\zeta, \mu_\infty)^2 \left(e^{-(\lambda - \kappa)t} + \frac{\bar{\kappa}(e^{-(\lambda - \kappa)t} - e^{-\bar{\lambda}t})}{\kappa + \bar{\lambda} - \lambda} \right) + \mathbb{W}_2(\theta, \nu_\infty)^2 e^{-\bar{\lambda}t}, \quad t \geq 0,$$

where when $\kappa + \bar{\lambda} = \lambda$,

$$\frac{e^{-(\lambda - \kappa)t} - e^{-\bar{\lambda}t}}{\kappa + \bar{\lambda} - \lambda} := te^{-\bar{\lambda}t}, \quad t \geq 0.$$

Consequently, the unique solution $(\mu_t, \nu_t)_{t \geq 0}$ to (1.10) with $\mu_0, \nu_0 \in \mathcal{P}_2$ satisfies

$$\begin{aligned} & \mathbb{W}_2(\mu_t, \mu_\infty)^2 + \mathbb{W}_2(\nu_t, \nu_\infty)^2 \\ & \leq \mathbb{W}_2(\mu_0, \mu_\infty)^2 \left(e^{-(\lambda - \kappa)t} + \frac{\bar{\kappa}(e^{-(\lambda - \kappa)t} - e^{-\bar{\lambda}t})}{\kappa + \bar{\lambda} - \lambda} \right) + \mathbb{W}_2(\nu_0, \nu_\infty)^2 e^{-\bar{\lambda}t}, \quad t \geq 0. \end{aligned}$$

PROOF. It suffices to prove the first assertion.

Firstly, by [34, Theorem 3.1], (A) implies that P_t^* has a unique invariant probability measure μ_∞ such that

$$(6.8) \quad \mathbb{W}_2(P_t^* \zeta, \mu_\infty)^2 \leq e^{-(\lambda-\kappa)t} \mathbb{W}_2(\zeta, \mu_\infty)^2, \quad t \geq 0, \zeta \in \mathcal{P}_2.$$

Next, by (6.3), there exist constants $c_1, c_2 > 0$ such that

$$2\langle \bar{b}(x, \mu_\infty), x \rangle + \|\bar{\sigma}(x, \mu_\infty)\|_{HS}^2 \leq c_1 - c_2|x|^2, \quad x \in \mathbb{R}^d.$$

It is standard that this implies the existence of an invariant probability measure ν_∞ of the diffusion process $\bar{X}_t$ associated with the SDE

$$(6.9) \quad d\bar{X}_t = \bar{b}(\bar{X}_t, \mu_\infty)dt + \bar{\sigma}(\bar{X}_t, \mu_\infty)dW_t.$$

Take an $\mathcal{F}_0$ -measurable random variable $(\bar{X}_0, X_0^{\zeta, \theta})$ on $\mathbb{R}^d \times \mathbb{R}^d$, such that $\mathcal{L}_{(\bar{X}_0, X_0^{\zeta, \theta})} \in \mathcal{C}(\nu_\infty, \theta)$ and

$$(6.10) \quad \mathbb{E}|\bar{X}_0 - X_0^{\zeta, \theta}|^2 = \mathbb{W}_2(\theta, \nu_\infty)^2.$$

Let $(\bar{X}_t)_{t \geq 0}$ solve (6.9) with initial value $\bar{X}_0$ which has law ν_∞ . Since ν_∞ is the invariant probability measure for the solution, we have

$$(6.11) \quad \mathcal{L}_{\bar{X}_t} = \nu_\infty, \quad t \geq 0.$$

So,

$$(6.12) \quad \mathbb{W}_2(\mathcal{L}_{X_t^{\zeta, \theta}}, \nu_\infty)^2 \leq \mathbb{E}|X_t^{\zeta, \theta} - \bar{X}_t|^2, \quad t \geq 0.$$

On the other hand, let $\Lambda = \nu_\infty \times \delta_{\mu_\infty}$. Then (6.6) implies

$$(6.13) \quad \mathbb{W}_2(\mathbf{P}_t(\theta, \zeta; \cdot), \Lambda)^2 \leq \mathbb{W}_2(P_t^* \zeta, \mu_\infty)^2 + \mathbb{W}_2(\mathcal{L}_{X_t^{\zeta, \theta}}, \nu_\infty)^2, \quad t \geq 0.$$

By (6.3), (6.7), (6.9) and Itô's formula, we obtain

$$\begin{aligned} & d|X_t^{\zeta, \theta} - \bar{X}_t|^2 \\ &= \left\{ 2\langle \bar{b}(X_t^{\zeta, \theta}, P_t^* \zeta) - \bar{b}(\bar{X}_t, \mu_\infty), X_t^{\zeta, \theta} - \bar{X}_t \rangle + \|\bar{\sigma}(X_t^{\zeta, \theta}, P_t^* \zeta) - \bar{\sigma}(\bar{X}_t, \mu_\infty)\|_{HS}^2 \right\} dt + dM_t \\ &\leq \{ \bar{\kappa} \mathbb{W}_2(P_t^* \zeta, \mu_\infty)^2 - \bar{\lambda} |X_t^{\zeta, \theta} - \bar{X}_t|^2 \} dt + dM_t, \quad t \geq 0, \end{aligned}$$

where

$$dM_t := 2\langle \bar{\sigma}(X_t^{\zeta, \theta}, P_t^* \zeta) - \bar{\sigma}(\bar{X}_t, \mu_\infty) dW_t, X_t^{\zeta, \theta} - \bar{X}_t \rangle$$

is a martingale. Combining this with (6.8), (6.11) and (6.10), we get

$$\begin{aligned} (6.14) \quad \mathbb{E}|X_t^{\zeta, \theta} - \bar{X}_t|^2 &\leq e^{-\bar{\lambda}t} \mathbb{E}|\bar{X}_0 - X_0^{\zeta, \theta}|^2 + \bar{\kappa} \mathbb{W}_2(\zeta, \mu_\infty)^2 \int_0^t e^{-\bar{\lambda}(t-s) - (\lambda-\kappa)s} ds \\ &= e^{-\bar{\lambda}t} \mathbb{W}_2(\nu, \nu_\infty)^2 + \frac{\bar{\kappa} \mathbb{W}_2(\zeta, \mu_\infty)^2}{\kappa + \bar{\lambda} - \lambda} (e^{-(\lambda-\kappa)t} - e^{-\bar{\lambda}t}), \end{aligned}$$

where when $\kappa + \bar{\lambda} = \lambda$,

$$\frac{e^{-(\lambda-\kappa)t} - e^{-\bar{\lambda}t}}{\kappa + \bar{\lambda} - \lambda} := e^{-\bar{\lambda}t} \lim_{s \rightarrow \bar{\lambda}} \frac{e^{(\bar{\lambda}-s)t} - 1}{\bar{\lambda} - s} = te^{-\bar{\lambda}t}.$$

Combining (6.14) with (6.8), (6.12) and (6.13), we derive

$$\mathbb{W}_2(\mathbf{P}_t(\theta, \zeta; \cdot), \Lambda)^2 \leq \mathbb{W}_2(\zeta, \mu_\infty)^2 \left(e^{-(\lambda-\kappa)t} + \frac{\bar{\kappa}(e^{-(\lambda-\kappa)t} - e^{-\bar{\lambda}t})}{\kappa + \bar{\lambda} - \lambda} \right) + e^{-\bar{\lambda}t} \mathbb{W}_2(\theta, \nu_\infty)^2, \quad t \geq 0.$$

As a consequence, Λ is the unique invariant probability measure for the $\mathbf{L}$ -diffusion process. $\square$

7. Feynman-Kac formula for PDEs on $\mathbb{R}^d \times \mathcal{P}_2$. In this section we aim to solve the PDE (1.17) by using the diffusion process generated by $\mathbf{L}_t$. To this end, we first recall the notion of intrinsic/ L -derivative, and present some classes of differentiable functions on $\mathcal{P}_2$, see Appendix below for a geometry explanation which implies that the class of cylindrical functions $\mathcal{F}C_b^2(\mathcal{P})$ is included in $C_b^{(1,1)}(\mathcal{P}_2)$.

DEFINITION 7.1. Let f be a real function on $\mathcal{P}_2$, and let $\text{Id} : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be the identity map.

(1) We call f intrinsically differentiable at $\mu \in \mathcal{P}_2$, if

$$L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d; \mu) \ni \phi \mapsto \nabla_\phi^\mathcal{P} f(\mu) := \lim_{\varepsilon \downarrow 0} \frac{f(\mu \circ (\text{Id} + \varepsilon \phi)^{-1}) - f(\mu)}{\varepsilon}$$

is a well defined bounded linear functional. In this case, the intrinsic derivative is the unique element $\nabla^\mathcal{P} f(\mu) \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d; \mu)$ such that

$$\langle \nabla^\mathcal{P} f(\mu), \phi \rangle_{L^2(\mu)} = \nabla_\phi^\mathcal{P} f(\mu), \quad \phi \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d; \mu).$$

f is called intrinsically differentiable if it is intrinsically differentiable at any $\mu \in \mathcal{P}_2$.

(2) We call f L -differentiable on $\mathcal{P}_2$ if it is intrinsically differentiable and

$$\lim_{\|\phi\|_{L^2(\mu)} \downarrow 0} \frac{|f(\mu \circ (\text{Id} + \phi)^{-1}) - f(\mu) - \nabla_\phi^\mathcal{P} f(\mu)|}{\|\phi\|_{L^2(\mu)}} = 0, \quad \mu \in \mathcal{P}_2.$$

Let $C^1(\mathcal{P}_2)$ be the set of all L -differentiable functions $f : \mathcal{P}_2 \rightarrow \mathbb{R}$ with $\nabla^\mathcal{P} f(\mu)(y)$ having a jointly continuous version in $(\mu, y) \in \mathcal{P}_2 \times \mathbb{R}^d$, and denote $f \in C_b^1(\mathcal{P}_2)$ if moreover $\nabla^\mathcal{P} f(\mu)(y)$ is bounded.

(3) We write $f \in C^{(1,1)}(\mathcal{P}_2)$ if $f \in C^1(\mathcal{P}_2)$ and $\nabla^\mathcal{P} f(\mu)(y)$ is differentiable in y , such that $\nabla\{\nabla^\mathcal{P} f(\mu)(\cdot)\}(y)$ is jointly continuous in $(\mu, y) \in \mathcal{P}_2 \times \mathbb{R}^d$. If moreover $f \in C_b^1(\mathcal{P}_2)$ and $\nabla\{\nabla^\mathcal{P} f(\mu)(\cdot)\}(y)$ is bounded, we denote $f \in C_b^{(1,1)}(\mathcal{P}_2)$.

7.1. Main result. We will work with the following class $C_b^{0,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$.

DEFINITION 7.2. We write $f \in C_b^{0,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$, if $f(t, x, \mu)$ is continuous in $(t, x, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}_2$, C^2 in $x \in \mathbb{R}^d$, $C^{(1,1)}$ in $\mu \in \mathcal{P}_2$, such that the derivatives

$$\nabla f(t, x, \mu), \quad \nabla^2 f(t, x, \mu), \quad \nabla^\mathcal{P} f(t, x, \mu)(y), \quad \nabla\{\nabla^\mathcal{P} f(t, x, \mu)(\cdot)\}(y)$$

are bounded and jointly continuous in $(t, x, \mu, y) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}_2 \times \mathbb{R}^d$. If moreover $\partial_t f(t, x, \mu)$ is continuous in $(t, x, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}_2$, we denote $f \in C_b^{1,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$; and write $f \in C_b^{2,(1,1)}(\mathbb{R}^d \times \mathcal{P}_2)$ if $f(t, x, \mu)$ does not depend on t .

According to Section 4, for any $(x, \mu) \in \mathbb{R}^d \times \mathcal{P}_2$, the $\mathbf{L}_t$ -diffusion process $(X_{s,t}^{\mu,x}, \mu_t)_{t \geq s}$ starting at (x, μ) generated by $\mathbf{L}_t$ on $\mathbb{R}^d \times \mathcal{P}_2$ can be constructed as follows: $\mu_t = P_{s,t}^* \mu$ is the law of X_t which is the unique solution of (1.8) from time s with $\mathcal{L}_{X_s} = \mu$, and $(X_{s,t}^{\mu,x})_{t \geq s}$ is the unique solution to the second equation in (2.13) with $X_{s,s}^{\mu,x} = x$.

The main result in this section is the following.

THEOREM 7.3. Assume that $b, \sigma, \bar{b}, \bar{\sigma} \in C_b^{0,2,2}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$. Then for any $\mathbf{V}, \mathbf{f} \in C_b^{0,2,(1,1)}(\mathbb{R}^d \times \mathcal{P}_2)$ where $\mathbf{V}$ is bounded, and for any $\Phi \in C_b^{2,(1,1)}(\mathbb{R}^d \times \mathcal{P}_2)$, the PDE (1.17) with $u(T, \cdot, \cdot) = \Phi$ has a unique solution $u \in C_b^{1,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$, and the solution is given by

$$(7.1) \quad \begin{aligned} & u(t, x, \mu) \\ &= \mathbb{E} \left[\Phi(X_{t,T}^{\mu,x}, P_{t,T}^* \mu) e^{\int_t^T \mathbf{V}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr} + \int_t^T \mathbf{f}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) e^{\int_t^r \mathbf{V}(\theta, X_{t,\theta}^{\mu,x}, P_{t,\theta}^* \mu) d\theta} dr \right] \end{aligned}$$

for $(t, x, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}_2$.

When $b = \bar{b}$ and $\sigma = \bar{\sigma}$ do not depend on t and $\mathbf{V} = 0$, this result is included by [21, Theorem 9.2] under slightly stronger conditions where the class $C_b^{0,2,(1,1)}$ is replaced by $C_{b,Lip}^{0,2,(1,1)}$: $f \in C_{b,Lip}^{0,2,(1,1)}$ means it is in $C_b^{0,2,(1,1)}$ such that

$$\nabla f(t, x, \mu), \quad \nabla^2 f(t, x, \mu), \quad \nabla^{\mathcal{P}} f(t, x, \mu)(y), \quad \nabla \{ \nabla^{\mathcal{P}} f(t, x, \mu)(\cdot) \}(y)$$

are Lipschitz continuous in $(x, \mu) \in \mathbb{R}^d \times \mathcal{P}_2$. Moreover, [21, Theorem 9.2] generalizes (with jump) and improves (under weaker conditions) the corresponding earlier results in [10, 13, 15].

7.2. Proof of Theorem 7.3. We first recall the following result taken from [21] (see also [10, 19]), which was proved for time independent coefficients $b = \bar{b}$ and $\sigma = \bar{\sigma}$, but the proof obviously works for the present time dependent coefficients, since all calculations therein only rely on the regularity of coefficients in the space-distribution variables (x, μ) but has nothing to do with derivatives in time.

LEMMA 7.4. Let $b, \sigma, \bar{b}, \bar{\sigma} \in C_b^{0,2,2}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$. Then

$$\nabla X_{s,t}^{\mu,\cdot}(x), \quad \nabla^2 X_{s,t}^{\mu,\cdot}(x), \quad \nabla^{\mathcal{P}} X_{s,t}^{\cdot,x}(\mu)(y), \quad \nabla \{ \nabla^{\mathcal{P}} X_{s,t}^{\cdot,x}(\mu)(\cdot) \}(y)$$

are jointly continuous in $(t, x, \mu, y) \in [s, T] \times \mathbb{R}^d \times \mathcal{P}_2 \times \mathbb{R}^d$, and there exists a constant $c > 0$ such that

$$\mathbb{E}(\|\nabla X_{s,t}^{\mu,\cdot}(x)\|^2 + \|\nabla^2 X_{s,t}^{\mu,\cdot}(x)\|^2 + \|\nabla^{\mathcal{P}} X_{s,t}^{\cdot,x}(\mu)\|_{L^2(\mu)}^2, \|\nabla \{ \nabla^{\mathcal{P}} X_{s,t}^{\cdot,x}(\mu) \}\|_{L^2(\mu)}^2) \leq c,$$

holds for all $0 \leq s \leq T$ and $(x, \mu) \in \mathbb{R}^d \times \mathcal{P}_2$.

Replacing $(\bar{b}, \bar{\sigma})$ in (2.14) by (b, σ) , we consider the SDE

$$(7.2) \quad dY_{s,t}^{\mu,x} = b(t, Y_{s,t}^{\mu,x}, P_{s,t}^* \mu) dt + \sigma(t, Y_{s,t}^{\mu,x}, P_{s,t}^* \mu) dW_t, \quad Y_{s,s}^{\mu,x} = x.$$

Then the Markov property of the solution implies

$$(7.3) \quad P_{s,t}^* \mu = \int_{\mathbb{R}^d} \mathcal{L}_{Y_{s,t}^{\mu,x}} \mu(dx),$$

where $P_{s,t}^* \mu := \mathcal{L}_{X_t}$ for X_t solving (1.8) from time s with $\mathcal{L}_{X_s} = \mu$. Combining this with Lemma 7.4, which applies to $Y_{s,t}^{\mu,x}$ replacing $X_{s,t}^{\mu,x}$ by taking $(\bar{b}, \bar{\sigma}) = (b, \sigma)$, we have the following result.

LEMMA 7.5. Let $b, \sigma \in C_b^{0,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$. Then the assertion in Lemma 7.4 holds for $Y_{s,t}^{\mu,x}$ replacing $X_{s,t}^{\mu,x}$, and for any $f \in C_b^{(1,0)}(\mathcal{P}_2)$,

$$(7.4) \quad \begin{aligned} \nabla^{\mathcal{P}} f(P_{s,t}^* \cdot)(\mu)(y) &= \int_{\mathbb{R}^d} \mathbb{E} \left[\{ \nabla^{\mathcal{P}} Y_{s,t}^{\mu,z}(\mu)(y) \} (\nabla^{\mathcal{P}} f)(P_{s,t}^* \mu)(Y_{s,t}^{\mu,z}) \right] \mu(dz) \\ &\quad + \mathbb{E} \left[\{ \nabla Y_{s,t}^{\mu,\cdot}(y) \} (\nabla^{\mathcal{P}} f)(P_{s,t}^* \mu)(Y_{s,t}^{\mu,y}) \right], \end{aligned}$$

where for vectors $v_1, v_2 \in \mathbb{R}^d$,

$$\langle \{ \nabla Y_{s,t}^{\mu,\cdot}(z) \} v_1, v_2 \rangle := \langle \nabla_{v_2} Y_{s,t}^{\mu,\cdot}(z), v_1 \rangle,$$

and $\{ \nabla^{\mathcal{P}} Y_{s,t}^{\mu,z}(\mu)(\cdot) \} v_1 \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d; \mu)$ is defined by

$$\langle \{ \nabla^{\mathcal{P}} Y_{s,t}^{\mu,z}(\mu)(\cdot) \} v_1, \phi \rangle_{L^2(\mu)} := \langle \nabla_{\phi}^{\mathcal{P}} \{ Y_{s,t}^{\mu,z}(\mu)(\cdot) \}, v_1 \rangle, \quad \phi \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d; \mu).$$

PROOF. Let $f \in C_b^{(1,0)}(\mathcal{P}_2)$, i.e. f is L -differentiable with $\nabla^{\mathcal{P}} f(\mu)(y)$ having a bounded jointly continuous version. For a family of random variables $\{ \xi^\varepsilon : \varepsilon \in [0, 1] \}$ with $\dot{\xi}^0 := \frac{d\xi^\varepsilon}{d\varepsilon} \big|_{\varepsilon=0}$ existing in $L^1(\mathbb{P})$, we have

$$(7.5) \quad \lim_{\varepsilon \downarrow 0} \frac{f(\mathcal{L}_{\xi(\varepsilon)}) - f(\mathcal{L}_{X^0})}{\varepsilon} = \mathbb{E} \langle \nabla^{\mathcal{P}} f(\mathcal{L}_{\xi(0)})(\xi(0)), \dot{\xi}^0 \rangle.$$

See for instance [27, Proposition 3.1], which is slightly extended from [18, Proposition A.2]. Since $P_{s,t}^* \mu = \int_{\mathbb{R}^d} \mathcal{L}_{Y_{s,t}^{\mu,z}} \mu(dz)$, it follows from (7.5) that for any $\phi \in L^2(\mu)$,

$$\begin{aligned} \nabla_{\phi}^{\mathcal{P}} f(P_{s,t}^* \cdot)(\mu) &= \frac{d}{d\varepsilon} \left\{ f \left(\int_{\mathbb{R}^d} \mathcal{L}_{Y_{s,t}^{\mu \circ (\text{Id} + \varepsilon \phi)} - 1, z} \mu(dz) \right) + \frac{d}{d\varepsilon} f \left(\int_{\mathbb{R}^d} \mathcal{L}_{Y_{s,t}^{\mu, z + \varepsilon \phi(z)}} \mu(dz) \right) \right\} \Big|_{\varepsilon=0} \\ &= \int_{\mathbb{R}^d} \mathbb{E} \langle (\nabla^{\mathcal{P}} f)(P_{s,t}^* \mu)(Y_{s,t}^{\mu,z}), \nabla_{\phi}^{\mathcal{P}} Y_{s,t}^{\mu,\cdot}(\mu) + \nabla_{\phi(z)} Y_{s,t}^{\mu,\cdot}(z) \rangle \mu(dz) \\ &= \mathbb{E} \int_{\mathbb{R}^d \times \mathbb{R}^d} \left[\langle \{ \nabla^{\mathcal{P}} Y_{s,t}^{\mu,z}(\mu)(\cdot) \} (\nabla^{\mathcal{P}} f)(P_{s,t}^* \mu)(Y_{s,t}^{\mu,z}), \phi \rangle_{L^2(\mu)} \right. \\ &\quad \left. + \langle \{ \nabla Y_{s,t}^{\mu,\cdot}(z) \} (\nabla^{\mathcal{P}} f)(P_{s,t}^* \mu)(Y_{s,t}^{\mu,z}), \phi(z) \rangle \right] \mu(dz). \end{aligned}$$

Therefore, $f(P_{s,t}^* \cdot)$ is intrinsic differentiable such that (7.4) holds. $\square$

LEMMA 7.6. Assume that $b, \sigma, \bar{b}, \bar{\sigma} \in C_b^{0,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$. For $\mathbf{V}, \mathbf{f} \in C_b^{0,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$, let u be in (7.1). Then $u \in C_b^{0,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$.

PROOF. The proof is more or less standard, but for completeness we present below a brief proof.

Obviously, $u(t, x, \mu)$ is joint continuous in (t, x, μ) . Next, by making derivatives in (x, μ) for u in (7.1), we obtain

$$\begin{aligned} \nabla^i u(t, \cdot, \mu)(x) &= \mathbb{E} \left[e^{\int_t^T \mathbf{V}(X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr} \left(\langle \nabla^i \Phi(\cdot, P_{s,T}^* \mu)(X_{t,T}^{\mu,x}), \nabla^i X_{t,T}^{\mu,\cdot}(x) \rangle \right. \right. \\ &\quad \left. \left. + \Phi(X_{t,T}^{\mu,x}, P_{s,T}^* \mu) \int_t^T \langle \nabla^i \mathbf{V}(\cdot, P_{s,r}^* \mu)(X_{t,r}^{\mu,x}), \nabla^i X_{t,r}^{\mu,\cdot}(x) \rangle dr \right) \right. \\ &\quad \left. + \int_t^T e^{\int_t^r \mathbf{V}(\theta, X_{t,\theta}^{\mu,x}, P_{t,\theta}^* \mu) d\theta} \left(\langle \nabla^i \mathbf{f}(\cdot, P_{s,r}^* \mu)(X_{t,r}^{\mu,x}), \nabla^i X_{t,r}^{\mu,\cdot}(x) \rangle dr \right. \right. \\ &\quad \left. \left. + \mathbf{f}(X_{t,r}^{\mu,x}, P_{s,r}^* \mu) \int_t^r \langle \nabla^i \mathbf{V}(\theta, \cdot, P_{t,\theta}^* \mu)(X_{t,\theta}^{\mu,x}), \nabla^i X_{t,\theta}^{\mu,\cdot}(x) \rangle d\theta \right) dr \right], \quad i = 1, 2. \end{aligned}$$

and

$$\begin{aligned}
& \nabla^{\mathcal{P}} u(t, x, \cdot)(\mu)(y) \\
&= \mathbb{E} \left[e^{\int_t^T \mathbf{V}(X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr} \left(\langle \nabla^{\mathcal{P}} \Phi(X_{t,T}^{\cdot,x}, P_{t,T}^* \mu)(\mu)(y), \nabla^{\mathcal{P}} X_{t,T}^{\cdot,x}(\mu)(y) \rangle \right. \right. \\
&+ \Phi(X_{t,T}^{\mu,x}, P_{s,T}^* \mu) \int_t^T \langle \nabla^{\mathcal{P}} \mathbf{V}(X_{t,r}^{\cdot,x}, P_{s,r}^* \mu)(\mu)(y), \nabla^{\mathcal{P}} X_{t,r}^{\cdot,x}(\mu)(y) \rangle dr \\
&+ \nabla^{\mathcal{P}} \Phi(X_{s,t}^{x,\mu}, P_{s,t}^* \cdot)(\mu)(y) + \Phi(X_{t,T}^{x,\mu}, P_{s,T}^* \mu) \int_t^T \nabla^{\mathcal{P}} \mathbf{V}(X_{t,r}^{x,\mu}, P_{s,r}^* \cdot)(\mu)(y) dr \Big) \\
&+ \int_t^T e^{\int_t^r \mathbf{V}(\theta, X_{t,\theta}^{\mu,x}, P_{t,\theta}^* \mu) d\theta} \left(\langle \nabla^{\mathcal{P}} \mathbf{f}(X_{t,r}^{\cdot,x}, P_{s,r}^* \mu)(\mu), \nabla^{\mathcal{P}} X_{t,r}^{\cdot,x}(\mu) \rangle(y) + \nabla^{\mathcal{P}} \mathbf{f}(r, X_{s,r}^{\mu,x}, P_{s,r}^* \cdot)(\mu)(y) \right. \\
&\left. \left. + \mathbf{f}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) \int_t^r \{ \nabla^{\mathcal{P}} \mathbf{V}(\theta, X_{t,\theta}^{\cdot,x}, P_{t,\theta}^* \mu)(\mu)(y) + \nabla^{\mathcal{P}} \mathbf{V}(\theta, X_{t,\theta}^{\mu,x}, P_{s,\theta}^* \cdot)(\mu)(y) \} d\theta \right) dr \right].
\end{aligned}$$

Combining these with Lemma 7.5, we finish the proof. $\square$

We will need the following lemma, which follows from [13, Theorem 3.3] or [18, Proposition A.6] under the stronger condition

$$(7.6) \quad \int_s^T \mathbb{E}(|\alpha_t|^2 + \|\beta_t\|^4) dt < \infty, \quad T \in (s, \infty),$$

where $\|\cdot\|$ is the operator norm of matrices.

LEMMA 7.7. *Let $\alpha : [s, \infty) \rightarrow \mathbb{R}^d$ and $\beta : [s, \infty) \rightarrow \mathbb{R}^{d \otimes m}$ be progressively measurable with*

$$(7.7) \quad \mathbb{E} \left(\int_s^T |\alpha_t| dt \right)^2 + \mathbb{E} \int_s^T \|\beta_t\|^2 dt < \infty, \quad T \in (s, \infty).$$

For $X_s \in L^2(\Omega \rightarrow \mathcal{F}_s; \mathbb{P})$, let $\mu_t = \mathcal{L}_{X_t}$ for

$$X_t := X_s + \int_s^t \alpha_r dr + \int_s^t \beta_r dW_r, \quad t \geq s.$$

Then $\mu \in C([s, \infty) \rightarrow \mathcal{P}_2)$ and for any $f \in C_b^{(1,1)}(\mathcal{P}_2)$,

$$\frac{df(\mu_t)}{dt} = \mathbb{E} \left[\frac{1}{2} \langle \beta_t \beta_t^*, \nabla \{ \nabla^{\mathcal{P}} f(\mu_t)(\cdot) \}(X_t) \rangle_{HS} + \langle \alpha_t, (\nabla^{\mathcal{P}} f(\mu_t))(X_t) \rangle \right], \quad t \geq s.$$

PROOF. Since $\mu_s \in \mathcal{P}_2$ and (7.7) holds, it is easy to see that $\mu \in C([s, \infty) \rightarrow \mathcal{P}_2)$. For any $n \geq 1$, let $\alpha_t^n = \alpha_t 1_{\{|\alpha_t| \leq n\}}$, $\beta_t^n = \beta_t 1_{\{\|\beta_t\| \leq n\}}$, and let $\mu_t^n = \mathcal{L}_{X_t^n}$ for

$$X_t^n := X_s + \int_s^t \alpha_r^n dr + \int_s^t \beta_r^n dW_r, \quad t \geq s.$$

Then

$$\lim_{n \rightarrow \infty} \sup_{t \in [s, T]} W_2(\mu_t^n, \mu_t) = 0$$

and (7.6) holds for (α_t^n, β_t^n) replacing (α_t, β_t) . So, by [13, Theorem 3.3] or [18, Proposition A.6], we obtain

$$f(\mu_t^n) = f(\mu_s) + \mathbb{E} \int_s^t \left[\frac{1}{2} \langle \beta_r^n \{\beta_r^n\}^*, \nabla \{ \nabla^{\mathcal{P}} f(\mu_r^n)(\cdot) \} (X_r^n) \rangle_{HS} + \langle \alpha_r^n, (\nabla^{\mathcal{P}} f(\mu_r^n)(X_r^n)) \rangle \right] dr$$

for $t \geq s$. Since $\nabla^{\mathcal{P}} f(\mu)(y)$ and $\nabla \{ \nabla^{\mathcal{P}} f(\mu)(\cdot) \}(y)$ are bounded and continuous in (μ, y) , by (7.7) we may apply the dominated convergence theorem with $n \rightarrow \infty$ to derive

$$f(\mu_t) = f(\mu_s) + \mathbb{E} \int_s^t \left[\frac{1}{2} \langle \beta_r \beta_r^*, \nabla \{ \nabla^{\mathcal{P}} f(\mu_r)(\cdot) \} (X_r) \rangle_{HS} + \langle \alpha_r, (\nabla^{\mathcal{P}} f(\mu_r)(X_r)) \rangle \right] dr, \quad t \geq s.$$

Then the proof is finished. $\square$

We now apply Lemma 7.7 to prove the following Itô's formula for $(X_{s,t}^{x,\mu}, P_{s,t}^* \mu)$.

LEMMA 7.8. *Assume that there exists an increasing function $K : [0, \infty) \rightarrow (0, \infty)$ such that*

$$(7.8) \quad \{ |b| + \|\sigma\| + |\bar{b}| + \|\bar{\sigma}\| \}(t, x, \mu) \leq K(t)(1 + |x| + \|\mu\|_2), \quad (t, x, \mu) \in [0, \infty) \times \mathbb{R}^d \times \mathcal{P}_2.$$

Then for any $f \in C^{2,(1,1)}(\mathbb{R}^d \times \mathcal{P}_2)$ and $s \in [0, T)$,

$$df(X_{s,t}^{\mu,x}, P_{s,t}^* \mu) = \mathbf{L}_t f(X_{s,t}^{\mu,x}, P_{s,t}^* \mu) dt + \langle \nabla f(\cdot, P_{s,t}^* \mu)(X_{s,t}^{\mu,x}), \sigma(t, X_{s,t}^{\mu,x}, P_{s,t}^* \mu) dW_t \rangle, \quad t \in [s, T].$$

PROOF. It is easy to see that (7.8) implies (7.7) for

$$\alpha_t := b(t, X_t, P_{s,t}^* \mu), \quad \beta_t := \sigma(t, X_t, P_{s,t}^* \mu)$$

for X_t solving (1.8) from time s with $\mathcal{L}_{X_s} = \mu$. By Lemma 7.7 and the definition of $\mathbf{L}_t^{(2)}$, for any $z \in \mathbb{R}^d$ we have

$$df(z, P_{s,t}^* \mu) = \mathbf{L}_t^{(2)} f(z, P_{s,t}^* \mu) dt, \quad t \geq s.$$

Combining this with Itô's formula for $X_{s,t}^{x,\mu}$ in (2.14), we obtain

$$\begin{aligned} df(X_{s,t}^{\mu,x}, P_{s,t}^* \mu) &= \{ df(z, P_{s,t}^* \mu) \}_{|_{z=X_{s,t}^{\mu,x}}} + \{ df(X_{s,t}^{\mu,x}, \nu) \}_{|_{\nu=P_{s,t}^* \mu}} \\ &= \mathbf{L}_t f(X_{s,t}^{\mu,x}, P_{s,t}^* \mu) dt + \langle \nabla f(\cdot, P_{s,t}^* \mu)(X_{s,t}^{\mu,x}), \sigma(t, X_{s,t}^{\mu,x}, P_{s,t}^* \mu) dW_t \rangle. \end{aligned}$$

$\square$

We are now ready to prove Theorem 7.3 as follows.

PROOF OF THEOREM 7.3. (a) We first prove that u in (7.1) solves (1.17). Let $(t, x, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}_2$. For any $\varepsilon \in (0, T - t)$ we have

$$\begin{aligned} u(t, x, \mu) &:= \mathbb{E} \left[\Phi(X_{t,T}^{x,\mu}, P_{t,T}^* \mu) e^{\int_t^T \mathbf{V}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr} + \int_t^T e^{\int_t^r \mathbf{V}(\theta, X_{t,\theta}^{\mu,x}, P_{t,\theta}^* \mu) d\theta} \mathbf{f}(r, X_{t,r}^{x,\mu}, P_{t,r}^* \mu) dr \right] \\ &= I_1 + I_2, \end{aligned}$$

where

$$\begin{aligned}
I_1 &:= \mathbb{E} \left[\Phi(X_{t,T}^{\mu,x}, P_{t,T}^* \mu) e^{\int_{t+\varepsilon}^T \mathbf{V}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr} + \int_{t+\varepsilon}^T e^{\int_{t+\varepsilon}^r \mathbf{V}(\theta, X_{t,\theta}^{\mu,x}, P_{t,\theta}^* \mu) d\theta} \mathbf{f}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr \right], \\
I_2 &:= \mathbb{E} \left[\Phi(X_{t,T}^{\mu,x}, P_{t,T}^* \mu) \left\{ e^{\int_t^T \mathbf{V}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr} - e^{\int_{t+\varepsilon}^T \mathbf{V}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr} \right\} \right. \\
&\quad + \int_t^{t+\varepsilon} e^{\int_t^r \mathbf{V}(\theta, X_{t,\theta}^{\mu,x}, P_{t,\theta}^* \mu) d\theta} \mathbf{f}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr \\
&\quad \left. + \int_{t+\varepsilon}^T \left\{ e^{\int_t^r \mathbf{V}(\theta, X_{t,\theta}^{\mu,x}, P_{t,\theta}^* \mu) d\theta} - e^{\int_{t+\varepsilon}^r \mathbf{V}(\theta, X_{t,\theta}^{\mu,x}, P_{t,\theta}^* \mu) d\theta} \right\} \mathbf{f}(r, X_{t,r}^{\mu,x}, P_{t,r}^* \mu) dr \right].
\end{aligned}$$

By the Markov property of $(X_{t,r}^{\mu,x}, P_{t,r}^* \mu)_{r \in [t, T]}$, we obtain

$$\begin{aligned}
I_1 &= \mathbb{E} \left\{ \mathbb{E} \left[\Phi(X_{t+\varepsilon, T}^{\nu, y}, P_{t+\varepsilon, T}^* \nu) e^{\int_{t+\varepsilon}^T \mathbf{V}(r, X_{t+\varepsilon, r}^{\nu, y}, P_{t+\varepsilon, r}^* \nu) dr} \right. \right. \\
&\quad \left. \left. + \int_{t+\varepsilon}^T \mathbf{f}(r, X_{t+\varepsilon, r}^{\nu, y}, P_{t+\varepsilon, r}^* \nu) e^{\int_{t+\varepsilon}^r \mathbf{V}(\theta, X_{t+\varepsilon, \theta}^{\nu, y}, P_{t+\varepsilon, \theta}^* \nu) d\theta} dr \right] \middle| (y, \nu) = (X_{t, t+\varepsilon}^{\mu, x}, P_{t, t+\varepsilon}^* \mu) \right\} \\
&= \mathbb{E} u(t + \varepsilon, X_{t, t+\varepsilon}^{\mu, x}, P_{t, t+\varepsilon}^* \mu).
\end{aligned}$$

Combining this with Lemma 7.6 and Lemma 7.8, we arrive at

$$I_1 = u(t + \varepsilon, x, \mu) + \mathbb{E} \int_t^{t+\varepsilon} \mathbf{L}_r u(t + \varepsilon, \cdot, \cdot)(X_{t,r}^{\mu, x}, P_{t,r}^* \mu) dr.$$

Noting that $u(t, x, \mu) = I_1 + I_2$, b, σ, Φ and $\mathbf{f}$ are continuous with linear growth, $\mathbf{V}$ is continuous and bounded, and $u \in C_b^{0,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$, we may apply the dominated convergence theorem to derive

$$\begin{aligned}
&\lim_{\varepsilon \downarrow 0} \frac{u(t, x, \mu) - u(t + \varepsilon, x, \mu)}{\varepsilon} \\
&= \lim_{\varepsilon \downarrow 0} \frac{1}{\varepsilon} \mathbb{E} \left\{ \int_t^{t+\varepsilon} \mathbf{L}_r u(t + \varepsilon, \cdot, \cdot)(X_{t,r}^{\mu, x}, P_{t,r}^* \mu) dr + I_2 \right\} \\
&= \mathbf{L}_t u(t, \cdot, \cdot)(x, \mu) + (u \mathbf{V})(t, x, \mu) + \mathbf{f}(t, x, \mu).
\end{aligned}$$

Therefore, u solves (1.17), and $\partial_t u$ is continuous on $[0, T] \times \mathbb{R}^d \times \mathcal{P}_2$, so that by Lemma 7.6 and the definition, we have $u \in C_b^{1,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$.

(b) Let $u \in C^{1,2,(1,1)}([0, T] \times \mathbb{R}^d \times \mathcal{P}_2)$ be a solution to (1.17), we prove that it satisfies (7.1). Indeed, let

$$\eta_t = u(t, X_{s,t}^{\mu, x}, P_{s,t}^* \mu) e^{\int_s^t \mathbf{V}(r, X_{s,r}^{\mu, x}, P_{s,r}^* \mu) dr} + \int_s^t \mathbf{f}(r, X_{s,r}^{\mu, x}, P_{s,r}^* \mu) e^{\int_s^r \mathbf{V}(\theta, X_{s,\theta}^{\mu, x}, P_{s,\theta}^* \mu) d\theta} dr, \quad t \in [0, T].$$

By Lemma 7.8 and (1.17), for any $s \in [0, T)$, we have

$$\begin{aligned}
d\eta_t &= \left\{ (\partial_t + \mathbf{L}_t) u(t, \cdot, \cdot)(X_{s,t}^{\mu, x}, P_{s,t}^* \mu) + (u \mathbf{V})(t, X_{s,t}^{\mu, x}, P_{s,t}^* \mu) + \mathbf{f}(t, X_{s,t}^{\mu, x}, P_{s,t}^* \mu) \right. \\
&\quad \left. + \langle \nabla u(t, \cdot, P_{s,t}^* \mu)(X_{s,t}^{\mu, x}), \sigma(t, X_{s,t}^{\mu, x}, P_{s,t}^* \mu) dW_t \rangle \right\} e^{\int_s^t \mathbf{V}(r, X_{s,r}^{\mu, x}, P_{s,r}^* \mu) dr} \\
&= e^{\int_s^t \mathbf{V}(r, X_{s,r}^{\mu, x}, P_{s,r}^* \mu) dr} \langle \nabla u(t, \cdot, P_{s,t}^* \mu)(X_{s,t}^{\mu, x}), \sigma(t, X_{s,t}^{\mu, x}, P_{s,t}^* \mu) dW_t \rangle, \quad t \in [s, T].
\end{aligned}$$

Therefore, for any $s \in [0, T]$,

$$\begin{aligned} u(s, x, \mu) &= \mathbb{E}\eta_s = \mathbb{E}\eta_T \\ &= \mathbb{E}\left\{u(T, X_{s,T}^{\mu,x}, P_{s,T}^*\mu) e^{\int_s^T \mathbf{V}(r, X_{s,r}^{\mu,x}, P_{s,r}^*\mu) dr} + \int_s^T \mathbf{f}(r, X_{s,r}^{\mu,x}, P_{s,r}^*\mu) e^{\int_s^r \mathbf{V}(\theta, X_{s,\theta}^{\mu,x}, P_{s,\theta}^*\mu) d\theta} dr\right\}, \end{aligned}$$

that is, u satisfies (7.1). $\square$

APPENDIX A: A NATURAL TANGENT BUNDLE OVER $\mathcal{P}$

It is well-known how to identify natural tangent bundles over “manifold-like” state spaces M and their corresponding gradients. For this one has to fix a large enough space of “smooth” real-valued functions $\mathcal{F}$ (“test functions”) on M and for each $x \in M$ a set of “suitable curves” $\gamma^x : [0, 1] \rightarrow M$, $\gamma(0) = x$ along which we can differentiate $t \mapsto f(\gamma^x(t))$ at $t = 0$ for all $f \in \mathcal{F}$. This construction has been performed in [1], [2] (see also [3], [4], [22], [25], [26], [30], [31]) with the space M being the space Γ of all $\mathbb{Z}_+$ -valued Radon-measures on a Riemannian manifold, i.e. Γ is the *configuration space* over this manifold. The space $\mathcal{F}$ there consists of so-called finitely based smooth functions on Γ . It turns out that in this case the resulting tangent bundle $(T_\mu\Gamma)_{\mu \in \Gamma}$ consists of linear spaces $T_\mu\Gamma$ given by μ -square integrable sections over the underlying Riemannian manifold. Let us now present the completely analogous construction in the case we are concerned with in this paper, where the underlying Riemannian manifold is $\mathbb{R}^d$ and Γ is replaced by $\mathcal{P} =: \mathcal{P}$. Define (see (2.3))

$$\mathcal{F} := \mathcal{F}C_b^2(\mathcal{P})$$

and for $\mu \in \mathcal{P}$, $\phi \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)$:

$$(A.1) \quad \gamma_\phi^\mu(t) := \mu \circ (\text{Id} + t\phi)^{-1}, \quad t \geq 0.$$

So, in our case the set of “suitable curves” starting at $\mu \in \mathcal{P}$ are labelled by $\phi \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)$.

Claim: The resulting tangent bundle over $\mathcal{P}$ is $(T_\mu\mathcal{P})_{\mu \in \mathcal{P}} := (L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu))_{\mu \in \mathcal{P}}$

PROOF. Let $F \in \mathcal{F} = \mathcal{F}C_b^2(\mathcal{P})$ and $\mu \in \mathcal{P}$. Then

$$(A.2) \quad F(\mu) = f(\mu(h_1), \dots, \mu(h_n)),$$

for some $n \in \mathbb{N}$, $h_1, \dots, h_n \in C_b^2(\mathbb{R}^d)$, $f \in C_b^1(\mathbb{R}^n)$ and thus by the chain rule for all $\phi \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)$

$$\begin{aligned} (A.3) \quad & \frac{d}{dt} F(\gamma_\phi^\mu(t))|_{t=0} \\ &= \sum_{i=1}^n (\partial_i f)(\mu(h_1), \dots, \mu(h_n)) \mu(\langle \nabla h_i, \phi \rangle_{\mathbb{R}^d}) \\ &= \left\langle \sum_{i=1}^n (\partial_i f)(\mu(h_1), \dots, \mu(h_n)) \nabla h_i, \phi \right\rangle_{L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)} \end{aligned}$$

$\square$

Define the corresponding gradient for $F \in \mathcal{FC}_b^2(\mathcal{P})$ by

$$(A.4) \quad \nabla^{\mathcal{P}} F(\mu) := \sum_{i=1}^n (\partial_i f)(\mu(h_1), \dots, \mu(h_n)) \nabla h_i \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu).$$

Then by (A.3) we have for all $\phi \in L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)$

$$(A.5) \quad \frac{d}{dt} F(\gamma_\phi^\mu(t))|_{t=0} = \langle \nabla^{\mathcal{P}} F(\mu), \phi \rangle_{L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)}.$$

In particular, the definition of $\nabla^{\mathcal{P}} F$ is independent of the particular representation of F in (A.3).

REMARK A.1. We note that for $F \in \mathcal{FC}_b^2(\mathcal{P})$ we have that $\nabla^{\mathcal{P}} F$ is bounded on $\mathbb{R}^d$. So, the right hand side is well-defined also if merely $\phi \in L^1(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)$. For simplicity of notation and extending the inner product we keep the notation

$$\int \langle \nabla^{\mathcal{P}} F(\mu), \phi \rangle_{\mathbb{R}^d} d\mu =: \langle \nabla^{\mathcal{P}} F(\mu), \phi \rangle_{L^2(\mathbb{R}^d \rightarrow \mathbb{R}^d, \mu)}$$

also in this case in the entire paper.

REFERENCES

- [1] S. Albeverio, Y. G. Kondratiev, M. Röckner, *Differential geometry of Poisson spaces*, C. R. Acad. Sci. Paris Sér. I Math. 323(1996), 1129–1134.
- [2] S. Albeverio, Y. G. Kondratiev, M. Röckner, *Canonical Dirichlet operator and distorted Brownian motion on Poisson spaces* C. R. Acad. Sci. Paris Sér. I Math. 323(1996), 1179–1184.
- [3] S. Albeverio, Y. G. Kondratiev, M. Röckner, *Analysis and geometry on configuration spaces*, J. Funct. Anal. 154(1998), 444–500.
- [4] S. Albeverio, Y. G. Kondratiev, M. Röckner, *Analysis and geometry on configuration spaces: the Gibbsian case*, J. Funct. Anal. 157(1998), 242–291.
- [5] V. Barbu, M. Röckner, *Probabilistic representation for solutions to nonlinear Fokker-Planck equations*, SIAM J. Math. Anal. 50(2018), 4246–4260.
- [6] V. Barbu, M. Röckner, *From nonlinear Fokker-Planck equations to solutions of distribution dependent SDE*, arXiv:1808.10706v2, to appear in Annal. Probab.
- [7] V. Barbu, M. Röckner, *The evolution to equilibrium of solutions to nonlinear Fokker-Planck equation*, arXiv:1904.08291v3.
- [8] V. Barbu, M. Röckner, *Uniqueness for nonlinear Fokker-Planck equations and weak uniqueness for McKean-Vlasov SDEs*, arXiv:1909.04464.
- [9] V. I. Bogachev, N. V. Krylov, M. Röckner, S.V. Shaposhnikov, *Fokker-Planck-Kolmogorov equations*, Mathematical Surveys and Monographs 207, American Mathematical Society, Providence, RI, 2015.
- [10] R. Buckdahn, J. Li, S. Peng, C. Rainer, *Mean-field stochastic differential equations and associated PDEs*, Annal. Probab. 2(2017), 824–878.
- [11] P. Cardaliaguet, *Notes on mean field games*, P.-L. Lions lectures at College de France, <https://www.ceremade.dauphine.fr/~cardaliaguet/MFG20130420.pdf>.
- [12] R. Carmona, F. Delarue, *Probabilistic Theory of Mean Field Games with Applications I & II*, Springer, 2018.
- [13] J.-F. Chassagneux, D. Crisan, F. Delarue, *A Probabilistic approach to classical solutions of the master equation for large population equilibria*, arXiv:1411.3009v2, 2015.
- [14] P. E. Chaudru de Raynal, *Strong well-posedness of McKean-Vlasov stochastic differential equation with Hölder drift*, arXiv: 1512.08096; 2019: hal-01248324v3.
- [15] D. Crisan, E. McMurray, *Smoothing properties of McKean-Vlasov SDEs*, Probab. Theory Relat. Fields 171(2018), 97–148.
- [16] A. Figalli, *Existence and uniqueness of martingale solutions for SDEs with rough or degenerate coefficients*, J. Func. Anal. 254(2008), 109–153.
- [17] T. D. Frank, *Nonlinear Fokker-Planck Equations*, Springer, 2005.

- [18] W. Hammersley, D. Šiška, L. Szpruch, *McKean-Vlasov SDE under measure dependent Lyapunov conditions*, arXiv:1802.03974v1
- [19] T. Hao, J. Li, *Mean-field SDEs with jumps and nonlocal integral-PDEs*, Nonlin. Diff. Equat. Appl. 23(2016), 1–51.
- [20] X. Huang, F.-Y. Wang, *Distribution dependent SDEs with singular coefficients*, Stoch. Proc. Appl. 129(2019), 4747–4770.
- [21] J. Li, *Mean-field forward and backward SDEs with jumps and associated nonlocal quasi-linear integral-PDEs*, Stoch. Proc. Appl. 128(2018), 3118–3180.
- [22] Z.-M. Ma, M. Röckner, *Construction of diffusions on configuration spaces*, Osaka J. Math. 37(2000), 273–314.
- [23] Yu. S. Mishura, A. Yu. Veretennikov, *Existence and uniqueness theorems for solutions of McKean-Vlasov stochastic equations*, arXiv:1603.02212.
- [24] F. Otto, *The geometry of dissipative evolution equations: the porous medium equation*, Comm. Part. Diff. Eq. 26(2001), 101–174.
- [25] L. Overbeck, M. Röckner, B. Schmuland, *An analytic approach to Fleming-Viot processes with interactive selection*, Ann. Probab. 23(1995), 1–36.
- [26] L. Overbeck, M. Röckner, *Geometric aspects of finite- and infinite-dimensional Fleming-Viot processes*, Random Oper. Stochastic Equations 5(1997), 35–58.
- [27] P. Ren, F.-Y. Wang, *Bismut formula for Lions derivative of distribution dependent SDEs and applications*, J. Diff. Equat. 267(2019), 4745–4777.
- [28] P. Ren, F.-Y. Wang, *Space-distribution PDEs for path independent additive functionals of McKean-Vlasov SDEs*, arXiv:1805.10841.
- [29] P. Ren, F.-Y. Wang, *Derivative formulas in measure on Riemannian manifolds*, arXiv:1908.03711.
- [30] M. Röckner, *Stochastic analysis on configuration spaces: basic ideas and recent results*, New directions in Dirichlet forms (J. Jost et al., eds.), Studies in Advanced Mathematics, International Press, 1998, pp. 157–231.
- [31] M. Röckner, *Dirichlet forms on infinite-dimensional “manifold-like” state spaces: a survey of recent results and some prospects for the future*, Probability towards 2000 (L. Accardi et al., eds.), Lecture Notes in Statistics, Springer, Berlin, 1998, pp. 287–306.
- [32] D.W. Stroock, S.R.S. Varadhan, *Multidimensional diffusion processes*, Springer-Verlag, New York 1979.
- [33] D. Trevisan, *Well-posedness of multidimensional diffusion processes*, Elect. J. Probab. 21(2016), 1–41.
- [34] F.-Y. Wang, *Distribution dependent SDEs for Landau type equations*, Stoch. Proc. Appl. 128(2018), 595–621.

ADDRESS OF THE FIRST AUTHOR
 MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD
 WOODSTOCK ROAD, OX2 6GG, UNITED KINGDOM
 AND
 DEPARTMENT OF MATHEMATICS, SWANSEA UNIVERSITY
 BAY CAMPUS, SA1 8EN, UNITED KINGDOM
 E-MAIL: Panpan.ren@maths.ox.ac.uk

ADDRESS OF THE SECOND AUTHOR
 DEPARTMENT OF MATHEMATICS, BIELEFELD UNIVERSITY
 D-33501 BIELEFELD, GERMANY
 AND
 ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE
 CHINESE ACADEMY OF SCIENCE, BEIJING, CHINA
 E-MAIL: roeckner@math.uni-bielefeld.de

ADDRESS OF THE THIRD AUTHOR
 CENTER FOR APPLIED MATHEMATICS, TIANJIN UNIVERSITY
 TIANJIN 300072, CHINA
 AND
 DEPARTMENT OF MATHEMATICS, SWANSEA UNIVERSITY
 BAY CAMPUS, SA1 8EN, UNITED KINGDOM
 E-MAIL: wangfy@tju.edu.cn