

Cite this: *Org. Biomol. Chem.*, 2020, **18**, 3690

Received 29th March 2020,
Accepted 17th April 2020

DOI: 10.1039/d0ob00664e
rsc.li/obc

N-Hydroxy peptides: solid-phase synthesis and β -sheet propensity†

Matthew P. Sarnowski and Juan R. Del Valle  *

Peptide backbone amide substitution can dramatically alter the conformational and physiochemical properties of native sequences. Although uncommon relative to *N*-alkyl substituents, peptides harboring main-chain *N*-hydroxy groups exhibit unique conformational preferences and biological activities. Here, we describe a versatile method to prepare *N*-hydroxy peptide on solid support and evaluate the impact of backbone *N*-hydroxylation on secondary structure stability. Based on previous work demonstrating the β -sheet-stabilizing effect of α -hydrazino acids, we carried out an analogous study with *N*-hydroxy- α -amino acids using a model β -hairpin fold. In contrast to *N*-methyl substituents, backbone *N*-hydroxy groups are accommodated in the β -strand region of the hairpin without energetic penalty. An enhancement in β -hairpin stability was observed for a di-*N*-hydroxylated variant. Our results facilitate access to this class of peptide derivatives and inform the use of backbone *N*-hydroxylation as a tool in the design of constrained peptidomimetics.

Introduction

N-Hydroxy- α -amino acid residues are encountered in several natural metabolites and are biosynthetic precursors to structurally complex non-ribosomal peptides.¹ Several *N*-hydroxy peptide (NHP) natural products exhibit potent antibacterial properties, cancer cell cytotoxicity, or activity against endogenous hormone receptors.² The introduction of a hydroxamate group into the peptide backbone also serves to enhance metal chelating properties,³ stability toward some proteases,⁴ and the potential for enhanced hydrogen-bonding interactions with target proteins. For these reasons, several designed peptidomimetics harboring *N*-hydroxy- α -amino acid residues have found utility as biologically active analogues of native peptides.^{4,5}

Studies on the conformational impact of peptide *N*-hydroxylation or *N*-alkoxylation have been mostly limited to glycine derivatives.⁶ Although these residues lack side chain substituents, their *N*-acyl derivatives were found to exhibit a strong preference for the *trans* amide conformation. The hydroxamate group can also engage in both intra- and intermolecular hydrogen bonding interactions to stabilize extended or turn geometries. Despite these initial reports, the effect of backbone amide hydroxylation on peptide secondary structure remains largely unexplored.⁷ Conformational studies on NHPs featuring α -substituted *N*-hydroxy residues could provide design principles relevant to the stabilization of folded pep-

tides. Although solution-phase syntheses of several α -substituted NHPs have been described,^{3–4,5b,6b,7,8} access to wider array of diastereopure derivatives *via* solid-phase methods would enable further investigation.

We previously reported the solid-phase synthesis and conformational analysis of several *N*-amino peptides (NAPs) as novel β -strand mimics.⁹ Amide *N*-amination restricts backbone dihedral angles through cooperative non-covalent interactions as shown in Fig. 1. A key feature of NAPs is their ability to




Fig. 1 Structures and properties of *N*-heteroatom-substituted peptides.

Department of Chemistry & Biochemistry, University of Notre Dame, Notre Dame, Indiana 46556, USA. E-mail: jdelvalle@nd.edu


† Electronic supplementary information (ESI) available. See DOI: [10.1039/d0ob00664e](https://doi.org/10.1039/d0ob00664e)

engage in an intraresidue (C6) H-bond analogous to that observed in conformationally extended peptides and β -sheets (C5).¹⁰ Due to lone pair repulsion across the hydrazide bond, NAPs also avoid the significant *cis* amide rotamer propensity observed with *N*-alkylated peptides (peptoids, prolyl-containing peptides, and *N*-methyl peptides).¹¹ These factors result in stabilization of β -sheet-like conformation as determined by mutational studies using β -hairpin model systems.⁹ We envisioned that *N*-hydroxy- α -amino acid residues would exhibit similar or even enhanced β -sheet propensities given the greater electron density and hydrogen bond donor capacity of the OH group relative to NH₂. Here, we describe the solid-phase synthesis of diastereopure NHPs and examine the impact of *N*-hydroxy- α -amino acids on β -hairpin folding. Our results provide insights into the structure and stability of this underexplored class of *N*-heteroatom-substituted peptides.

Results and discussion

As shown in Fig. 1, *N*-heteroatom substitution introduces electron repulsion that may partially counteract the *cis* amide rotamer propensity of peptide tertiary amides. To quantify the impact of backbone *N*-hydroxylation on *trans/cis* amide bias we employed a Gly-Ala-Xaa-Gly tetrapeptide model system originally developed by Raleigh and co-workers.¹² This peptide has previously been used to determine the rotamer preferences of proline, pipecolic acid, ϵ -oxapipecolic acid, and other cyclic residues in place of Xaa.^{12,13} We adapted this sequence in order to directly compare a series of acyclic residues in the Xaa position, including those with *N*-heteroatom substituents. Following Fukuyama's method,¹⁴ we prepared the *N*-hydroxyalanine (hAla) variant in solution starting from *O*-benzyl-protected *L*-alanine 1 as shown in Scheme 1. *N*-Alkylation was effected with bromoacetonitrile to give secondary amine 2. Oxidation with *m*CPBA then provided the nitronate intermediate, which was subjected to aminolysis to give hAla benzyl ester 3. Selective *N*-acylation was achieved in the presence of pre-formed Fmoc alanine acid chloride and NaHCO₃.¹⁵ Elaboration of the dipeptide using standard protocols then provided tetrapeptide 7. Notably, we observed diacylated derivative 5 as the major product during condensation with Boc-Gly-OH. However, deacylation of the backbone *N*-OH occurred upon treatment with a 20% solution of piperidine in DMF, providing tetrapeptide 6 in good yield.

With NHP 7 in hand, we synthesized alanine, *N*-methylalanine, and *N*-aminoalanine variants for comparison by ¹H NMR (Table 1). Analysis of the spectra taken in D₂O revealed that *N*-heteroatom substituted analogues 7 and 8 exhibit significantly higher *trans* amide populations relative to *N*-methyl peptide 9. While NAP 8 did exhibit a small but detectable population of *cis* rotamer (~6%), NHP 7 exclusively adopts a *trans* amide conformation in solution. These results are consistent with lone pair repulsion in the *cis* conformation and the greater electron density of O relative to N. As expected, control peptide 10 exhibits a *trans*-only conformation in D₂O.





Scheme 1 Solution-phase synthesis of model NHP 8.

Table 1 Amide *trans/cis* equilibrium constants for 7–10 determined by NMR (rt in D₂O)

Peptide	R	<i>K</i> _{trans/cis}
7	OH	>99
8	NH ₂	16
9	CH ₃	2.8
10	H	>99

In order to evaluate the β -strand propensity of *N*-hydroxy- α -amino acid residues, we utilized a β -hairpin model developed by Horne and co-workers for mutational analysis.¹⁶ This 16-residue sequence is derived from the B1 domain of *Streptococcal* protein G (GB1, Fig. 2) and exhibits moderate folding in aqueous media. Its folded population can be quantified by ¹H NMR on the basis of diastereotopic separation of the Gly₁₀ H α signals in comparison to cyclic (fully folded) and truncated (random coil) controls. We have also employed this model system to assess the impact of backbone *N*-amination



Target dipeptide building blocks:





Fig. 2 β -Hairpin model system and dipeptide building blocks for Fmoc-SPPS.

on β -sheet stability.^{9b} In that study, elongation of *N*-amino peptides was achieved by on-resin acylation of α -hydrazino acid residues using Fmoc-protected acid chlorides. Here, we chose to simplify the SPPS protocol by pre-forming the hydroxamate bond in solution and incorporating modified building blocks as dipeptide fragments. This approach allows us to avoid the use of labile acid chlorides and the generation of insoluble salts during the automated phase of peptide synthesis. Because diacylated derivatives exemplified by 5 exhibit poor stability,^{8e} we chose to circumvent over-acylation during SPPS through the use of *O*-protected hydroxamates. Our target building blocks were thus the protected *N*-hydroxy dipeptides shown in Fig. 2, corresponding to two outer edge amide mutations within the model hairpin sequence.

The required Lys-hPhe dipeptide derivative was prepared as shown in Scheme 2, starting from methyl *D*-phenyllactate. Activation and displacement of the triflate intermediate with




Scheme 3 Synthesis of Ala-hVal building block 22.

allyloxamine proceeded smoothly to give 13. Dipeptide bond formation was then achieved through reaction with Fmoc-Lys (Boc)-Cl, which was generated *in situ* using Ghosez' reagent. Lithium iodide-promoted deesterification then provided building block 15 in 76% yield.

Efforts to employ a similar *S*_N2 displacement strategy for the synthesis of the analogous Ala-hVal dipeptide were unsuccessful due to the additional steric bulk of the valine side chain (Scheme 3). As an alternative, we prepared *N*-hydroxyvaline *t*-butyl ester using the nitroline methodology described above (see Scheme 1). Condensation of 21 with Fmoc-Ala-Cl was followed by Mitsunobu reaction with allyl alcohol and final TFA deprotection to give 22 in 60% yield over three steps.

Condensation of dipeptides or larger fragments can often lead to significant C-terminal epimerization. With access to *N*-allyloxy dipeptide acids 15 and 22, we sought to ensure that stereochemical integrity would not be compromised under standard coupling conditions during SPPS. Compound 15 was thus converted to (*S*)- α -methylbenzamide derivative 23 with HCTU and NMM at elevated temperature. A diastereomeric amide derivative differing in configuration at the (*O*-All)hPhe residue was synthesized for comparison by HPLC. Crude analytical traces revealed a $>99:1$ dr for 23 (Fig. 3). This level of diastereopurity demonstrates a lack of epimerization during activation of the dipeptide building block with HCTU/NMM. We attribute this configurational stability to suppressed oxazolionium formation in the case of *N*-alkoxy peptide fragments. Fig. 4 depicts the mechanism of peptide fragment epimerization upon activation of the carboxyl group.¹⁷ The *N*-allyloxy group is strongly electron withdrawing and presumably destabilizes the transition state leading to the labile oxazolonium intermediate.

Assembly of peptides on solid support employed standard Fmoc chemistry using HCTU/NMM activation (Table 2). Where appropriate, allyl deprotection with $Pd(Ph_3)_4$ was performed prior to global deprotection with TFA. Table 2 lists the peptides synthesized for the current study in addition to previously prepared peptides for comparative analysis. All amide



Scheme 2 Synthesis of Lys-hPhe building block 15.




Fig. 3 Determination of fragment epimerization during C-terminal condensation by RP-HPLC.





Fig. 4 Mechanism and suppression of oxazolone/oxazolonium-mediated epimerization.

substitutions are positioned on the outer edge of the putative hairpin so as not to disrupt interstrand H-bonding interactions. Peptides **11b–c** are mono-*N*-hydroxylated peptides (mono-NHPs) substituted at either the Phe or Val residue mutation sites. Peptides **11d–e** retain the allyl protecting group to provide comparison between *N*-hydroxy and *N*-alkoxy substitutions. To assess the impact of multiple backbone substitutions, peptides **11f–g** were synthesized. Entries **11h–j** from our previous work^{9b} are included here for comparison between *N*-hydroxy and *N*-amino substitution. Lastly, entries **11k–l** serve as a point of comparison between *N*-heteroatom and *N*-methyl substitutions.

Detailed NMR studies with **11b–11l** were carried in D_2O at 4 °C (1 mM peptide in 50 mM aq. $NaHPO_4$, pH 6.8). Full 1H NMR assignments were made on the basis of TOCSY and ROESY spectra. Several inter-strand NOE correlations observed for *N*-hydroxy and *N*-alkoxy peptides **11b–11g** were in agreement with those observed in the fully folded cyclic peptide, indicating the presence of β -hairpin structures.¹⁸ Quantitative assessment of folding was accomplished by calculating the

Table 2 Solid-phase assembly of β -hairpin peptides and list of synthesized sequences



Peptide	R ¹	R ²	Sequence
11a	H	H	H-GEWAYNPATGKFAVTE-NH ₂
11b	OH	H	H-GEWAYNPATGK- hPhe -AVTE-NH ₂
11c	H	OH	H-GEWAYNPATGKFA- hVal -TE-NH ₂
11d	OAll	H	H-GEWAYNPATGK-(<i>O</i> -All) hPhe -AVTE-NH ₂
11e	H	OAll	H-GEWAYNPATGKFA-(<i>O</i> -All) hVal -TE-NH ₂
11f	OH	OH	H-GEWAYNPATGK- hPhe -A- hVal -TE-NH ₂
11g	OAll	OAll	H-GEWAYNPATGK-(<i>O</i> -All) hPhe -A-(<i>O</i> -All) hVal -TE-NH ₂
11h	NH ₂	H	H-GEWAYNPATGK- aPhe -AVTE-NH ₂
11i	H	NH ₂	H-GEWAYNPATGKFA- aVal -TE-NH ₂
11j	NH ₂	NH ₂	H-GEWAYNPATGK- aPhe -A- aVal -TE-NH ₂
11k	Me	H	H-GEWAYNPATGK-(<i>N</i> -Me) Phe -AVTE-NH ₂
11l	H	Me	H-GEWAYNPATGKFA-(<i>N</i> -Me) Val -TE-NH ₂

Gly_{10} $\text{H}\alpha$ chemical shift separation, which was then converted to fraction folded and $\Delta\Delta G$ relative to **11a** (Table 3). The folded fraction of hPhe derivative **11b** was found to be 86%, corresponding to a stabilization of -0.5 kcal mol⁻¹ relative to **11a**. This increase in stability is comparable to NAPs **11h** and **11i**. *N*-Hydroxylation at Val **11c** did not enhance hairpin stability, but was still tolerated without energetic penalty. Entry **11f** shows that multiple *N*-hydroxylations are well supported within the hairpin, but did not provide the same gain in stability as diamination (**11j**). This may be due in part to the negligible effect of Val *N*-hydroxylation relative to *N*-amination (**11c** vs. **11i**). The folded fraction of *N*-alkoxy derivatives **11d–e**, lacking the C6 H-bonding capability of the *N*-hydroxy mutants, were found to be only 67%, corresponding to a slight destabilization. This trend is also seen in the *N*-alkoxy double mutant **11g**. While backbone *N*-methylation in **11k** resulted in significant destabilization ($+1.1$ kcal mol⁻¹), a less pronounced effect was observed in the case of (*N*-Me)Val mutant **11l**. The Val₁₄ residue appeared to be less sensitive to mutation in the case of the *N*-hydroxy, *N*-alkoxy, and *N*-methyl variants. Taken together, these results show that backbone *N*-hydroxylation is well tolerated in the strand region of a β -hairpin, and in some cases, provides enhancement of stability. Given that *O*-allyl variant **11g** exhibited only a single amide rotational isomer by NMR, it is likely that the loss of intraresidue H-bonds contributes to its lower folded population relative to di-NHP **11f**. As with *N*-amino peptides, a significant increase in folded population was observed for NHPs relative to their *N*-methylated analogues.


Table 3 Folding thermodynamics of NHPs and analogous peptides determined by NMR (D_2O , pH 6.8, 4 °C)

Peptide	Mutation	$\Delta\delta\text{Gly}_{10}$ (ppm)	% folded	ΔG_{fold} (kcal mol ⁻¹)	$\Delta\Delta G_{\text{fold}}$ (kcal mol ⁻¹)
11a	None	0.224	73	-0.5	
11b	hPhe ₁₂	0.255	86	-1.0	-0.5
11c	hVal ₁₄	0.221	72	-0.5	0.0
11d	(O-All)hPhe ₁₂	0.208	67	-0.4	+0.1
11e	(O-All)hVal ₁₄	0.209	67	-0.4	+0.1
11f	hPhe ₁₂ ,hVal ₁₄	0.260	88	-1.1	-0.6
11g	(O-All)hPhe ₁₂ ,(O-All)hVal ₁₄	0.203	65	-0.3	+0.2
11h	aPhe ₁₂	0.261	88	-1.1	-0.6
11i	aVal ₁₄	0.252	84	-0.9	-0.4
11j	aPhe ₁₂ ,aVal ₁₄	0.301	99 ^a	-2.5 ^a	-2.0
11k	(N-Me)Phe ₁₂		24	+0.6	+1.1
	trans	0.096	37	+0.3	+0.8
	cis	0.000	0	0	
11l	(N-Me)Val ₁₄		56	-0.1	+0.4
	trans	0.213	69	-0.4	+0.1
	cis	0.000	0	0	

^a A value of 99% was used for calculations as the derived %fold from $\Delta\delta\text{Gly}_{10}$ exceeded 100%.

Finally, it is worth noting that we observed reduced levels of crude purity for NHPs relative to their NAP counterparts. During mid-sequence LCMS analysis, we identified a mass fragment corresponding to loss of allyl alcohol. This prompted us to investigate the stability of *N*-allyloxy peptides towards basic conditions. Peptide 11d was thus dissolved in a 20% piperidine/MeOH solution at 5 mM concentration and analyzed by HPLC (Fig. 5). After 30 min exposure to base, a small amount of deallylated peptide was detected by LCMS.

At 2 h, a prominent peak corresponding to the N-terminal fragment of imine hydrolysis was identified. Interestingly, we observed a series of peaks that featured the mass of the imine species. We reasoned that these could represent the intact *N*-acyl imine as well as dehydrophenylalanine tautomers. These results suggest that *N*-alkoxy peptides are prone to base-promoted elimination across the $\text{N}\alpha$ -C α bond and that buffered piperidine solutions or lower reaction temperatures may improve the crude purity of NHPs prepared on solid support.

**Fig. 5** Proposed mechanism of base-mediated decomposition of 11d.

Conclusions

In summary, we describe the synthesis of *N*-hydroxy dipeptide building blocks, their incorporation into NHPs, and initial assessment of β -sheet propensity by NMR. Furthermore, a tetrapeptide model was synthesized in solution and used to demonstrate that NHPs exhibit an unusually high *trans* amide propensity relative to *N*-amino- and *N*-methyl peptides. *N*-Hydroxy- α -amino acid residues were accommodated into the strand region of a β -hairpin fold without energetic penalty. In the case of 11b and 11f, mono- and di-*N*-hydroxylation was found to enhance β -hairpin stability. Capping of the OH with an alkyl group compromised stability, suggesting that an intraresidue H-bond involving the hydroxamate may play an important role in structural preorganization. In addition to the conformational effects explored here, the current work provides a useful method to access various C α -substituted NHPs by SPPS. We anticipate that these results will enable further structural and biological studies on this intriguing class of *N*-heteroatom-substituted peptides.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

We gratefully acknowledge Dr Evgenii Kovrigin (Notre Dame Magnetic Resonance Research Center) and Dr Edwin Rivera (USF) for assistance with NMR acquisitions. This work was supported by a grant from the National Science Foundation (CHE1709927).

Notes and references

- (a) H. C. J. Ottenheijm and J. D. M. Herscheid, N-hydroxy . alpha.-amino acids in organic chemistry, *Chem. Rev.*, 1986, **86**, 697–707; (b) A. H. Cook and C. A. Slater, The structure of pulcherrimin, *J. Chem. Soc.*, 1956, 4133–4135; (c) J. C. MacDonald and G. P. Slater, Biosynthesis of gliotoxin and mycelianamide, *Can. J. Biochem.*, 1975, **53**, 475–478; (d) W. G. McCullough and R. S. Merkal, Iron-chelating compound from *Mycobacterium avium*, *J. Bacteriol.*, 1976, **128**, 15–20; (e) M. J. Cryle, S. G. Bell and I. Schlichting, Structural and Biochemical Characterization of the Cytochrome P450 CypX (CYP134A1) from *Bacillus subtilis*: A Cyclo-L-leucyl-L-leucyl Dipeptide Oxidase, *Biochemistry*, 2010, **49**, 7282–7296.
- For selected examples see: (a) H. Maehr, C. M. Liu, N. J. Palleroni, J. Smallheer, L. Todaro, T. H. Williams and J. F. Blount, Microbial products. VIII. Azinothricin, a novel hexadepsipeptide antibiotic, *J. Antibiot.*, 1986, **39**, 17–25; (b) K. Umezawa, K. Nakazawa, T. Uemura, Y. Ikeda, S. Kondo, H. Naganawa, N. Kinoshita, H. Hashizume, M. Hamada, T. Takeuchi and S. Ohba, Polyoxypeptin isolated from Streptomyces: A bioactive cyclic depsipeptide containing the novel amino acid 3-hydroxy-3-methylproline, *Tetrahedron Lett.*, 1998, **39**, 1389–1392; (c) D. J. Pettibone, B. V. Clineschmidt, P. S. Anderson, R. M. Freidinger, G. F. Lundell, L. R. Koupal, C. D. Schwartz, J. M. Williamson, M. A. Goetz, O. D. Hensens, J. M. Liesch and J. P. Springer, A Structurally Unique, Potent, and Selective Oxytocin Antagonist Derived from Streptomyces silvensis, *Endocrinology*, 1989, **125**, 217–222; (d) M. Igarashi, R. Sawa, N. Kinoshita, H. Hashizume, N. Nakagawa, Y. Homma, Y. Nishimura and Y. Akamatsu, Pargamicin A, a Novel Cyclic Peptide Antibiotic from Amycolatopsis sp, *J. Antibiot.*, 2008, **61**, 387; (e) S. I. Maffioli, Y. Zhang, D. Degen, T. Carzaniga, G. Del Gatto, S. Serina, P. Monciardini, C. Mazzetti, P. Guglierame, G. Candiani, A. I. Chiriac, G. Facchetti, P. Kaltofen, H.-G. Sahl, G. Deho, S. Donadio and R. H. Ebright, Antibacterial Nucleoside-Analog Inhibitor of Bacterial RNA Polymerase, *Cell*, 2017, **169**, 1240–1248.
- (a) Y. Ye, M. Liu, J. L. K. Kao and G. R. Marshall, Peptide-bond modification for metal coordination: Peptides containing two hydroxamate groups, *Biopolymers*, 2003, **71**, 489–515; (b) Y. Ye, M. Liu, J. L. F. Kao and G. R. Marshall, Novel trihydroxamate-containing peptides: design, synthesis, and metal coordination, *Biopolymers*, 2006, **84**, 472–489.
- (a) M. Marastoni, M. Bazzaro, S. Salvadori, F. Bortolotti and R. Tomatis, HIV-1 protease inhibitors containing an N-Hydroxyamino acid core structure, *Bioorg. Med. Chem.*, 2001, **9**, 939–945; (b) A. Bianco, C. Zabel, P. Walden and G. Jung, N-hydroxy-amide analogs of MHC-class I peptide ligands with nanomolar binding affinities, *J. Pept. Sci.*, 1998, **4**, 471–478.
- (a) M. Akiyama, A. Katoh, K. Iesaki and K. Shimizu, Synthesis and properties of enkephalin analogs containing an N-hydroxyamino acid, *Pept. Chem.*, 1985, **22nd**, 271–276; (b) M. Akiyama, K. Iesaki, A. Katoh and K. Shimizu, N-Hydroxy, amides. Part 5. Synthesis and properties of N-hydroxy peptides having leucine enkephalin sequences, *J. Chem. Soc., Perkin Trans. 1*, 1986, 851–855; (c) E. Ghidini, M. Delcanale, V. Servadio, C. Pietra, M. Bergamaschi, G. Villetti, E. Redenti, P. Ventura and L. Merlini, Synthesis of a new series of N-hydroxy, N-alkylamides of amino acids as ligands of NMDA glycine site, *Eur. J. Med. Chem.*, 1999, **34**, 711–717; (d) Y. Jin and D. H. Kim, Inactivation of thermolysin with N-chloroacetyl-N-hydroxy- β -L-phenylalanine N-methylamide, *Bull. Korean Chem. Soc.*, 1999, **20**, 777–780; (e) S. Hin, C. Zabel, A. Bianco, G. Jung and P. Walden, Cutting edge: N-hydroxy peptides: a new class of TCR antagonists, *J. Immunol.*, 1999, **163**, 2363–2367; (f) X. Hu, J. Zhu, S. Srivathsan and D. Pei, Peptidyl hydroxamic acids as methionine aminopeptidase inhibitors, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 77–79.
- (a) V. Dupont, A. Lecoq, J. P. Mangeot, A. Aubry, G. Boussard and M. Marraud, Conformational perturbations induced by N-amination and N-hydroxylation of peptides, *J. Am. Chem. Soc.*, 1993, **115**, 8898–8906; (b) M. Marraud, V. Dupont, V. Grand, S. Zerkout, A. Lecoq, G. Boussard, J. Vidal, A. Collet and A. Aubry, Modifications of the amide bond and conformational constraints in pseudopeptide analogs, *Biopolymers*, 1993, **33**, 1135–1148; (c) Y. Takeuchi and G. R. Marshall, Conformational Analysis of Reverse-Turn Constraints by N-Methylation and N-Hydroxylation of Amide Bonds in Peptides and Non-Peptide Mimetics, *J. Am. Chem. Soc.*, 1998, **120**, 5363–5372; (d) J. A. Crapster, J. R. Stringer, I. A. Guzei and H. E. Blackwell, Design and conformational analysis of peptoids containing N-hydroxy amides reveals a unique sheet-like secondary structure, *Biopolymers*, 2011, **96**, 604–616; (e) P. A. Jordan, B. Paul, G. L. Butterfoss, P. D. Renfrew, R. Bonneau and K. Kirshenbaum, Oligo(N-alkoxy glycines): trans substantiating peptoid conformations, *Biopolymers*, 2011, **96**, 617–626.
- For an example of α -helical constraint based on peptide N-alkoxylation see: J. Lawrence, M. Jourdan, Y. Vallee and V. Blandin, Peptide cyclization via ring-closing metathesis: the N-alkenoxy peptide approach, *Org. Biomol. Chem.*, 2008, **6**, 4575–4581.
- (a) G. Zvilichovsky and L. Heller, Synthesis of N-hydroxy peptides, *Tetrahedron Lett.*, 1969, 1159–1160;

(b) K. Shimizu, K. Nakayama and M. Akiyama, N-Hydroxy, amides. II. N-Benzylxy α -amino acid N-hydroxysuccinimide esters and synthesis of a hexapeptide having an alternating N-hydroxy amide-amide sequence, *Bull. Chem. Soc. Jpn.*, 1984, **57**, 2456–2462;

(c) P. L. Durette, F. Baker, P. L. Barker, J. Boger, S. S. Bondy, M. L. Hammond, T. J. Lanza, A. A. Pessolano and C. G. Caldwell, Total synthesis of L-156,602, a novel cyclic hexadepsipeptide antibiotic, *Tetrahedron Lett.*, 1990, **31**, 1237–1240; (d) P. Maire, V. Blandin, M. Lopez and Y. Vallee, Straightforward synthesis of N-hydroxy peptides, *Synlett*, 2003, 671–674; (e) J. Lawrence, L. Cointeaux, P. Maire, Y. Vallee and V. Blandin, N-Hydroxy and, N-acyloxy peptides: synthesis and chemical modifications, *Org. Biomol. Chem.*, 2006, **4**, 3125–3141; (f) K. J. Hale, S. Manaviazar and J. George, Total synthesis of (+)-A83586C, (+)-kettapeptin and (+)-azinothricin: powerful new inhibitors of β -catenin/TCF4- and E2F-mediated gene transcription, *Chem. Commun.*, 2010, **46**, 4021–4042; (g) Y. M. Elbatrawi, C. W. Kang and J. R. Del Valle, Total, synthesis of L-156,373 and an oxoPiz analog via a submonomer approach, *Org. Lett.*, 2018, **20**, 2707–2710.

9 (a) M. P. Sarnowski, C. W. Kang, Y. M. Elbatrawi, L. Wojtas and J. R. Del Valle, Peptide, N-Amination Supports β -Sheet Conformations, *Angew. Chem., Int. Ed.*, 2017, **56**, 2083–2086; (b) M. P. Sarnowski, K. P. Pedretty, N. Giddings, H. L. Woodcock and J. R. Del Valle, Synthesis and, beta-sheet propensity of constrained N-amino peptides, *Bioorg. Med. Chem.*, 2018, **26**, 1162–1166.

10 (a) C. Toniolo and E. Benedetti, Intramolecularly Hydrogen-Bonded Peptide Conformation, *Crit. Rev. Biochem.*, 1980, **9**, 1–44; (b) R. W. Newberry and R. T. Raines, A prevalent intraresidue hydrogen bond stabilizes proteins, *Nat. Chem. Biol.*, 2016, **12**, 1084–1088.

11 (a) R. J. Simon, R. S. Kania, R. N. Zuckermann, V. D. Huebner, D. A. Jewell, S. Banville, S. Ng, L. Wang, S. Rosenberg, C. K. Marlowe, *et al.*, Peptoids: a modular approach to drug discovery, *Proc. Natl. Acad. Sci. U. S. A.*, 1992, **89**, 9367–9371; (b) M. W. MacArthur and J. M. Thornton, Influence of proline residues on protein conformation, *J. Mol. Biol.*, 1991, **218**, 397–412; (c) J. Chatterjee, F. Rechenmacher and H. Kessler, N-Methylation, of Peptides and Proteins: An Important Element for Modulating Biological Functions, *Angew. Chem., Int. Ed.*, 2013, **52**, 254–269.

12 W.-J. Wu and D. P. Raleigh, Conformational Heterogeneity about Pipecolic Acid Peptide Bonds: Conformational, Thermodynamic, and Kinetic Aspects, *J. Org. Chem.*, 1998, **63**, 6689–6698.

13 (a) E. H. Howard, C. F. Cain, C. Kang and J. R. Del Valle, Synthesis, of Enantiopure ϵ -Oxapipecolic Acid, *J. Org. Chem.*, 2020, **85**, 1680–1686; (b) L. Ricci, L. Sernissi, D. Scarpi, F. Bianchini, A. Contini and E. G. Occhiato, Synthesis and conformational analysis of peptides embodying 2,3-methanopipeolic acids, *Org. Biomol. Chem.*, 2017, **15**, 6826–6836.

14 (a) H. Tokuyama, T. Kuboyama, A. Amano, T. Yamashita and T. Fukuyama, A Novel Transformation of Primary Amines to N-Monoalkylhydroxylamines, *Synthesis*, 2000, 1299–1304; (b) S. I. Medina, J. Wu and J. W. Bode, Nitrone protecting groups for enantiopure N-hydroxyamino acids: synthesis of N-terminal peptide hydroxylamines for chemoselective ligations, *Org. Biomol. Chem.*, 2010, **8**, 3405–3417.

15 (a) G. Prabhu, Basavaprabhu, N. Narendra, T. M. Vishwanatha and V. V. Sureshbabu, Amino acid chlorides: a journey from instability and racemization toward broader utility in organic synthesis including peptides and their mimetics, *Tetrahedron*, 2015, **71**, 2785–2832; (b) L. A. Carpino, B. J. Cohen, K. E. Stephens, S. Y. Sadat-Aalaee, J. H. Tien and D. C. Langridge, (Fluoren-9-ylmethoxy)carbonyl (Fmoc) amino acid chlorides. Synthesis, characterization, and application to the rapid synthesis of short peptide segments, *J. Org. Chem.*, 1986, **51**, 3732–3734.

16 (a) G. A. Lengyel and W. S. Horne, Design Strategies for the Sequence-Based Mimicry of Side-Chain Display in Protein β -Sheets by α/β -Peptides, *J. Am. Chem. Soc.*, 2012, **134**, 15906–15913; (b) F. J. Blanco, G. Rivas and L. Serrano, A short linear peptide that folds into a native stable beta-hairpin in aqueous solution, *Nat. Struct. Biol.*, 1994, **1**, 584–590.

17 M. Goodman and W. J. McGahren, Optically Active Peptide Oxazolones. Preliminary Racemization Studies under Peptide-Coupling Conditions, *J. Am. Chem. Soc.*, 1965, **87**, 3028–3029.

18 See the ESI† for details.