

Fatal arrhythmias: Another reason why doctors remain cautious about chloroquine/hydroxychloroquine for treating COVID-19

Ilija Uzelac, PhD,* Shahriar Iravani, MD, PhD,† Hiroshi Ashikaga, MD, PhD,‡
 Neal K. Bhatia, MD,† Conner Herndon, MS,* Abouzar Kaboudian, PhD,*
 James C. Gumbart, PhD,* Elizabeth M. Cherry, PhD,§ Flavio H. Fenton, PhD*

From the *School of Physics, Georgia Institute of Technology, Atlanta, Georgia, †Division of Cardiology, Section of Electrophysiology, Emory University Hospital, Atlanta, Georgia, ‡Cardiac Arrhythmia Service, Johns Hopkins University School of Medicine, Baltimore, Maryland, and §School of Computational Science and Engineering, Georgia Institute of Technology, Atlanta, Georgia.

BACKGROUND Early during the current coronavirus disease 19 (COVID-19) pandemic, hydroxychloroquine (HCQ) received a significant amount of attention as a potential antiviral treatment, such that it became one of the most commonly prescribed medications for COVID-19 patients. However, not only has the effectiveness of HCQ remained questionable, but mainly based on preclinical and a few small clinical studies, HCQ is known to be potentially arrhythmogenic, especially as a result of QT prolongation.

OBJECTIVE The purpose of this study was to investigate the arrhythmic effects of HCQ, as the heightened risk is especially relevant to COVID-19 patients, who are at higher risk for cardiac complications and arrhythmias at baseline.

METHODS An optical mapping technique utilizing voltage-sensitive fluorescent dyes was used to determine the arrhythmic

effects of HCQ in *ex vivo* guinea pig and rabbit hearts perfused with the upper therapeutic serum dose of HCQ (1000 ng/mL).

RESULTS HCQ markedly increased action potential dispersion, resulted in development of repolarization alternans, and initiated polymorphic ventricular tachycardia.

CONCLUSION The study results further highlight the proarrhythmic effects of HCQ.

KEYWORDS Action potential duration dispersion; Arrhythmias; Experimental optical mapping; Hydroxychloroquine; Long QT syndrome; T-wave alternans

(Heart Rhythm 2020; ■:1-7) © 2020 Heart Rhythm Society. All rights reserved.

Introduction

The drugs chloroquine (CQ) and hydroxychloroquine (HCQ), a less toxic derivative of CQ, are used to treat malaria and autoimmune conditions. They now have been proposed to have antiviral activity against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which is responsible for the coronavirus disease 2019 (COVID-19) pandemic. Recent *in vitro* studies^{1,2} and a small nonrandomized clinical trial of 36 patients from France³ had promising results and initiated the trend of using CQ/HCQ to treat COVID-19. However, the integrity of the nonrandomized clinical trial has been questioned by the International Society of Antimicrobial Chemotherapy for the trial's unclear inclusion criteria and triage of patients.⁴ Although a subsequent smaller randomized clinical trial of 30 patients showed little to no

effect,⁵ a larger randomized clinical trial of 62 patients showed that HCQ significantly reduced the incidence and duration of COVID-19 pneumonia.⁶ These studies do not have sufficient statistical power to unequivocally prove the positive effects of HCQ on COVID-19. Nevertheless, the urgency of the pandemic has resulted in (1) the United States Food and Drug Administration (FDA) issuing an emergency use authorization for CQ/HCQ as treatment of COVID-19, an action that has been criticized by former FDA leaders; and (2) a call by the World Health Organization (WHO) for rapid, large, global CQ/HCQ clinical trials.

Although CQ and HCQ have become the focus as treatment of COVID-19, they remain unendorsed by many physicians because of (1) limited clinical outcome data; (2) availability of other potentially more effective antiviral and interleukin inhibitors, such as remdesivir and tocilizumab, respectively; and (3) potential risk of malignant arrhythmia and sudden cardiac death (SCD) due to QT prolongation.^{7,8} In response to the trend of using CQ/HCQ for treatment of COVID-19, the Mayo Clinic,⁹ the Heart Rhythm Society

This study was supported by grants NIH 1R01HL143450-01 and NSF 1446675. **Address reprint requests and correspondence:** Dr Flavio H. Fenton, Georgia Institute of Technology, 837 State St NW, Atlanta, GA 30332. E-mail address: ffenton@physics.gatech.edu.

COVID-19 Task Force in conjunction with the American College of Cardiology and American Heart Association,¹⁰ and other clinicians¹¹ all independently called for urgently needed guidance when using HCQ alone and in combination with other drugs with regard to arrhythmias.

The most pressing issue is determining the dosage of CQ/HCQ that is effective yet safe, particularly in patients at risk for drug-induced long QT syndrome (LQTS) and SCD.⁹⁻¹² This is especially important because HCQ is now being used in combination with other antivirals and antibiotics such as ritonavir, lopinavir, and azithromycin, all of which are associated with drug-induced LQTS.¹¹ Recent reports have shown direct myocardial involvement in many COVID-19 patients,¹³ presumably due to the abundance of angiotensin-converting enzyme 2 receptors on the cardiac cell surface, which is the main entry point of SARS-CoV-2. Evidence of myocardial damage has been described in the form of elevated troponin levels and high inflammatory burden, which can induce vascular inflammation and myocarditis.^{13,14} Computational models have proposed that the SARS-CoV-2 envelope protein E' forms a pentameric ion channel, and incorporation of this channel into the cell membrane of cardiac myocytes may contribute to the reported arrhythmic events in COVID-19. Clinically, some patients with severe COVID-19 develop left ventricular dysfunction, cardiogenic shock, and various cardiac arrhythmias.^{15,16} Consequently, many COVID-19 patients who already are susceptible to arrhythmias may be at even higher risk for deadly arrhythmogenic effects from COVID-19 drug combination treatments.

A recent clinical trial of 81 patients in Brazil that tested 2 different doses of CQ was prematurely halted because several patients in the high-dose group developed LQTS and experienced malignant arrhythmias, including ventricular tachycardia (VT), within the first 3 days; 11 patients later died within 6 days.¹⁷ Although the precise mechanisms by which QC/HCQ initiates arrhythmia remain unclear, the electrophysiological effects of HCQ have been investigated in isolated mouse and guinea pig (GP) hearts.¹⁸ HCQ primarily blocks the funny current I_f , the L-type calcium current, and the rapid delayed rectifier potassium current I_{Kr} . As a result, the main electrophysiological manifestations of HCQ are sinus bradycardia (Online Figure 1) and repolarization abnormalities (QT prolongation, increased repolarization dispersion), which could lead to torsades de pointes¹⁹ (TdP) and polymorphic VT.¹⁷

In this study we used optical mapping to demonstrate the arrhythmic effects of HCQ on GP and rabbit hearts perfused with the upper therapeutic serum dose of HCQ (1000 ng/mL).

Methods

Tissue preparation

All experiments conformed to the current Guide for Care and Use of Laboratory Animals, published by the National Institutes of Health (NIH Publication No. 85-23, revised 1996) and were approved by the Office of Research and Integrity Assurance at Georgia Institute of Technology. Animals were lightly anesthetized with ketamine/xylazine/acepromazine (17/9/0.9 mg/kg), then deeply anesthetized with 3%-




Figure 1 Arrhythmic effects of hydroxychloroquine on rabbit and guinea pig hearts. **A: Top:** Transmembrane voltage signal in time from the region in the heart indicated with a *white dot* in the bottom frames. The voltage signal shows 5 regular activations at 250 ms, followed by initiation of fibrillation, characterized by a fast, disorganized electrical activity signal. **Bottom:** Two frames of the voltage profile on the heart's surface showing multiple fractionated waves driving fibrillation. **B:** Example of arrhythmic ectopic beats following a protocol of two S1 stimuli and a second shorter S2 originating from the apex. **Top:** Voltage signal showing the first 2 beats from a stimulation protocol that emulates a quick change of rate, which creates 2 arrhythmic ectopic (premature P1 and P2) beats. **Bottom:** Optical images showing the initiation of the S2 beat at the apex and then initiation of the first ectopic beat originating from the top center. Color bar indicates tissue polarized (-80 mV) in *blue* and depolarized ($+20$ mV) in *red*.

273 5% isoflurane mixed with oxygen for induction and approx-
 274 imately 1%–3% for maintenance of deep anesthesia. Five mi-
 275 nutes before euthanasia, heparin 300 U/kg was injected
 276 intravenously to prevent vessel blood clots. Hearts were
 277 excised via left thoracotomy under deep anesthesia. After
 278 excision, hearts were perfused retrogradely via the aorta
 279 with warm Tyrode solution ($37^\circ \pm 0.5^\circ\text{C}$) for 3 minutes
 280 and gassed with a mixture of 95% O₂/5% CO₂ to expel all
 281 blood and prepare the hearts for perfusion with cold cardio-
 282 plegic solution, also gassed with a mixture of 95% O₂/5%
 283 CO₂. The Tyrode solution consisted of the following (in
 284 mmol/L): for GP: NaCl 124, KCl 4.0, NaHCO₃ 24, NaH₂-
 285 PO₄·H₂O 1.2, MgCl₂·6H₂O 1.0, CaCl₂ 1.8, dextrose 4.0;
 286 and for rabbit: NaCl 124, KCl 4.0, NaHCO₃ 24, NaH₂-
 287 PO₄·H₂O 0.9, MgCl₂·6H₂O 0.7, CaCl₂ 2.0, dextrose 4.0.
 288 Cardioplegic cold solution contained the following (in
 289 mmol/L): NaCl 110, KCl 16, NaHCO₃ 10, MgCl₂·6H₂O
 290 16, and CaCl₂ 1.2; and was used to induce cardiac stasis to
 291 protect the myocardium while the hearts were transferred to
 292 the optical mapping laboratory (within 15 minutes). For GP
 293 hearts, a modified Tyrode solution was used for hypokalemia
 294 as described in the Results section.
 295

296 In the laboratory, the hearts were immersed in a heated
 297 oval chamber kept at $38^\circ \pm 0.5^\circ\text{C}$ and Langendorff-
 298 perfused with Tyrode solution gassed with a mixture of
 299 95% O₂/5% CO₂. Contraction motion was suppressed by
 300 (\pm)-Blebbistatin (Cayman Chemicals, Ann Arbor, MI) at Ty-
 301 rode concentration of 1.5–2 μM , previously prepared as a
 302 stock solution dissolved with dimethylsulfoxide at the ratio
 303 of 5 mg/mL. Voltage (Vm)-sensitive dye JPW-6003 (Dr Le-
 304 Leslie Loew's group, University at Connecticut, Farmington,
 305 CT) was previously prepared as a stock solution dissolved
 306 in ethanol at the ratio of 20 mg/mL. For the whole GP and
 307 rabbit hearts, 0.4 mg of Vm dye was used. The Vm dye
 308 was injected as a bolus in the cannulated aorta over a course
 309 of 3 minutes in small injection intervals 15 seconds apart so
 310 as not to interfere with the flow rate. The total accommoda-
 311 tion time for the heart, including dye staining before the
 312 initial measurements were taken, was 30 minutes. Forty-
 313 five minutes after cannulation, HCQ was added into the
 314 Tyrode solution at a concentration of 1000 ng/mL (upper
 315 therapeutic dose of HCQ), previously dissolved in a small
 316 volume of Tyrode solution free of organic solvents and
 317 stirred for 15 minutes on a hot plate kept at 35°C.
 318

319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 3




Figure 2 Dispersion of action potential duration (APD) as a result of hydroxychloroquine (HCQ) in rabbit and guinea pig (GP) hearts. Histograms were obtained from optical mapping after steady state was reached for a constant cycle length of 250 ms for rabbit and 220 ms for GP with and without HCQ. As alternans appeared with HCQ, 2 histograms are shown (1 for even beats and 1 for odd beats). **A:** APD distribution in rabbit ventricles with normal Tyrode solution and with HCQ showing an increase in APD_{max} and alternans from beat to beat. Optical action potentials also show APD_{max} increase and alternans in duration. **B:** APD distribution in GP ventricles with normal Tyrode solution and with HCQ. Optical mapping signals show similar effect as seen in rabbit. PCL = pacing cycle length.

control and HCQ. The histograms show how APD dispersion increased with HCQ, with APD_{max} prolongation from 155 to 205 ms, and median APD changing from 124 ms to alternating APD values with medians of 125 and 132 ms. Morphologic changes in action potential (AP) also can be observed in the optical mapping voltage trace comparing the normal constant AP with the increased and alternating APs with HCQ. Online [Video 3](#) shows the difference in wave propagation by periodic stimulation between control and HCQ. Without HCQ, continuous pacing is possible even at very short CL of 140 ms with propagation still smooth, whereas with HCQ at even relatively longer CL of 380 ms, large heterogeneities in wave propagation are present. These dynamic heterogeneities promote arrhythmia initiation at CL <250 ms and in some cases complex propagation at CL <350 ms (Online [Video 4](#)).

For GP, APD dispersion increased with HCQ, resulting in conduction blocks and arrhythmias initiated at CL <220 ms. [Figure 2B](#) shows APD dispersion at this CL without HCQ and the development of alternans with HCQ. With HCQ, alternans changed the normal APD_{min} of 80 ms and APD_{max} of 120 ms to 70 ms and 145 ms, respectively, on alternating beats. Mean APD also changed, alternating between 96 and 112 ms every other beat, representing a decrease and increase in the short and long APD, respectively, compared to mean APD of 104 ms without HCQ. [Figure 2B](#) shows the voltage trace and increase in AP duration and alternans with HCQ.

Although there was a large dispersion of APD across the GP heart, spatial alternans mostly developed as concordant alternans ([Figure 3](#)), in which APD alternates from long to short values from beat to beat. Only a small region of the tissue experienced discordant alternans. However, in rabbit,




Figure 3 Spatial dispersion of action potential duration (APD) as a result of hydroxychloroquine (HCQ) in rabbit and guinea pig (GP) hearts. Maps were obtained from steady-state activations at a pacing cycle length of 220 ms for GP and 250 ms for rabbit. **Top row:** Distribution of APD for even and odd beats. **Bottom row:** Change in action potential duration (Δ APD) for even and odd beats. **A:** GP displays mostly concordant alternans in which most of the tissue experiences a short action potential, followed by a long action potential. Only a small region of discordant alternans (DA) with the opposite pattern (long–short) appears in the lower right corner and edges of the tissue. **B:** Rabbit displays marked DA with different distribution of long and short APD on alternating beats. Note that during DA, regions that have a long APD will have a short APD on the next beat, and vice versa. Black bar indicates location of the stimulus electrode; data underneath could not be collected.

although the tissue showed a similar range of APDs across the tissue, a more complex spatial pattern developed as alternans was discordant, with some regions of the tissue alternating with long–short APDs and others alternating short–long. The large beat-to-beat changes in APD dispersion due to discordant alternans can be seen in the APD maps and Δ APD maps (Figure 3).

Although ectopic beats were not observed in rabbit hearts, ectopic beats were seen in GP hearts. Figure 1B and Online Video 5 show 2 ectopic beats generated after 3 stimulations in GP heart. Online Video 6 shows an episode of VT initiation in rabbit heart.

Because hypokalemia has been shown to be prevalent in COVID-19 patients, we tested the effect of HCQ under hypokalemia by switching the Tyrode solution in the GP experiment from normal to lower potassium (from 4 to 2.5 mmol/L) and lower magnesium (from 1 to 0.8 mmol/L). We perfused with hypokalemia Tyrode solution twice, with normal Tyrode solution (washout) in between. In both instances, fibrillation occurred spontaneously under the hypokalemic conditions (the heart was defibrillated when back in normal Tyrode solution), highlighting a possible higher risk of HCQ in patients with hypokalemia and thus the importance of monitoring and treating low potassium levels because they can increase the probability of arrhythmic events.¹⁹

Discussion

These experiments show that as QT interval/APD increased with HCQ, arrhythmias could be induced at CLs closer to physiological heart rates, thus demonstrating the potential dangers of HCQ. In particular, immediate arrhythmia initiation was observed when hypokalemia was induced with HCQ, but not only in the hypokalemia case. This is important because of particular concern about hypokalemia in COVID-19 disease.²¹ Hypokalemia seems to be present in almost all COVID-19 patients,²² likely due to the interaction of SARS-CoV2 with the renin-angiotensin system (RAS), binding to angiotensin I converting enzyme 2 receptor of RAS and causing prevalent hypokalemia.

With HCQ alone, spatial alternans was observed at longer CLs, including those of normal sinus heart rates without the drug. Although HCQ produces bradycardia that, in principle, may not allow these CLs to be reachable, sympathetic surges can easily lead to sudden increased heart rates that induce pause/bradycardia arrhythmias such as TdP. Furthermore, the increased APD heterogeneity induced by HCQ at bradycardic CLs also showed dramatic anisotropic conduction slowing even before alternans developed. This phenomenon can be seen clearly in Online Video 3. At CL of 380 ms, a marked slowing in conduction velocity is observed in a region of the heart that produces a nonsmooth propagation compared to the case without HCQ at a very short CL of

681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748

140 ms. Thus, HCQ induced complex repolarization heterogeneities and a substrate for arrhythmias such as TdP.

Study limitations

The most important limitation is the low number of experiments for rabbit and GP, and we are aware that our results are more descriptive because statistical analysis cannot be performed. However, because our results in normal rabbit and GP hearts compare well with those of other studies in both species,^{23,24} and in general when healthy these hearts do not go into fibrillation when slowly paced down (until conduction block) but did so under HCQ, we believe that our study has provided some mechanistic insights. Nevertheless, this study needs to be extended to allow for complete statistical analysis that can solidify the proarrhythmic mechanisms of HCQ as well as the possibility of any differences related to sex, given that female sex is known to be an independent risk factor for developing TdP.

Another limitation of our study is the use of animal models with small hearts and electrophysiologies different from human hearts. A major problem with using small hearts is the difficulty in inducing reentrant tachycardias due to the size mismatch between ventricles and reentrant wavelength. Therefore, a negative result in small hearts is not reassuring for the absence of arrhythmogenicity in humans. However, we observed polymorphic VT with characteristics similar to clinical TdP, which is a finding of concern because typically induction of TdP is easier in humans than in these models.

The most important ion channels in the genesis of TdP are the delayed rectifier potassium channels, especially I_{Kr} . Both rabbit and GP ventricles have dofetilide-sensitive I_{Kr} and are prone to exhibiting repolarization alternans as a result of potassium channel blockade.^{25–27} Therefore, these species are commonly used to test for drug-induced repolarization abnormalities^{28,29} that could affect humans.

Conclusion

A challenging aspect of COVID-19 treatment with respect to heart problems is the use of unusual drug combinations, such as HCQ with antibiotics such as azithromycin, which before the COVID-19 pandemic were rarely used together. The risk profiles of individual drugs with regard to QT prolongation have been characterized, but data on the combinations of those drugs are lacking. Only a few studies, just becoming available,³⁰ indicate prolongation of LQTS. Although current trials have seldom required discontinuation of therapy, it remains imperative to investigate the safety of potentially effective drugs such as CQ/HCQ and their combinations before their widespread clinical use as treatment against SARS-CoV-2.

Based on the limited and contradictory evidence reported to date, the efficacy of CQ/HCQ against COVID-19 is questionable at best, and the safety is unclear due to the propensity for fatal arrhythmia. Until the results of large, well-designed clinical trials (such as the WHO Solidarity Trial) are

available, clinicians have compelling reasons not to treat COVID-19 patients with CQ/HCQ.

Appendix

Supplementary data

Supplementary data associated with this article can be found in the online version at <https://doi.org/10.1016/j.hrthm.2020.05.030>.

References

1. Liu J, Cao R, Xu M, et al. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discov* 2020; 6:16.
2. Yao X, Ye F, Zhang M, et al. In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Published online March 9, 2020. *Clin Infect Dis* <https://doi.org/10.1093/cid/ciaa237>.
3. Gautret P, Lagier J-C, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. Published online March 20, 2020. *Int J Antimicrob Agents* <https://doi.org/10.1016/j.ijantimicag.2020.105949>.
4. Statement on IJAA Paper. Official Statement from International Society of Antimicrobial Chemotherapy.
5. Efficacy and Safety of Hydroxychloroquine for Treatment of Pneumonia Caused by 2019-nCoV (HC-nCoV). ClinicalTrials.gov.
6. Chen Z, Hu J, Zhang Z, et al. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. Preprint. Posted online April 10, 2020. *medRxiv* 2020:2020.03.22.20040758; <https://doi.org/10.1101/2020.03.22.20040758>.
7. Yazdany J, Kim AHJ. Use of hydroxychloroquine and chloroquine during the COVID-19 pandemic: what every clinician should know. *Ann Intern Med* 2020;172:754–755.
8. Kim AHJ, Sparks JA, Liew JW, et al; COVID-19 Global Rheumatology Alliance. A rush to judgment? Rapid reporting and dissemination of results and its consequences regarding the use of hydroxychloroquine for COVID-19. Published online March 30, 2020. *Ann Intern Med* <https://doi.org/10.7326/M20-1223>.
9. Mayo Clinic. Urgent guidance, approach to identify patients at risk of drug-induced sudden cardiac death from use of off-label COVID-19 treatments. March 25, 2020. *ScienceDaily*. www.sciencedaily.com/releases/2020/03/200325122009.htm.
10. Lakkireddy DR, Chung MK, Gopinathannair R, et al. Guidance for cardiac electrophysiology during the coronavirus (COVID-19) pandemic from the Heart Rhythm Society COVID-19 Task Force; Electrophysiology Section of the American College of Cardiology; and the Electrocardiography and Arrhythmias Committee of the Council on Clinical Cardiology, American Heart Association. Published online April 1, 2020. *Heart Rhythm* <https://doi.org/10.1016/j.hrthm.2020.03.028>.
11. Wu C-I, Postema PG, Arbelo, et al. SARS-CoV-2, COVID-19 and inherited arrhythmia syndromes. Published online March 31, 2020. *Heart Rhythm* <https://doi.org/10.1016/j.hrthm.2020.03.024>.
12. Joyce E, Fabre A, Mahon N. Hydroxychloroquine cardiotoxicity presenting as a rapidly evolving biventricular cardiomyopathy: key diagnostic features and literature review. *Eur Heart J Acute Cardiovasc Care* 2013;2:77–83.
13. Madjid M, Safavi-Naeini P, Solomon SD, Vardeny O. Potential effects of coronaviruses on the cardiovascular system: a review. Published online March 27, 2020. *JAMA Cardiol* <https://doi.org/10.1001/jamacardio.2020.1286>.
14. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–1062.
15. Hu H, Ma F, Wei X, Fang Y. Coronavirus fulminant myocarditis saved with glucocorticoid and human immunoglobulin. Published online March 16, 2020. *Eur Heart J* <https://doi.org/10.1093/eurheartj/ehaa190>.
16. Driggin E, Madhavan MV, Bikdeli B, et al. Cardiovascular considerations for patients, health care workers, and health systems during the coronavirus disease 2019 (COVID-19) pandemic. *J Am Coll Cardiol* 2020;75:2352–2371.
17. Borba M, Val F de A, Sampaio VS, et al. Chloroquine diphosphate in two different dosages as adjunctive therapy of hospitalized patients with severe respiratory syndrome in the context of coronavirus (SARS-CoV-2) infection: preliminary safety results of a randomized, double-blinded, phase IIb clinical trial (CloroCovid-19 Study). Preprint. Posted online April 16, 2020. *medRxiv* 2020 2020; <https://doi.org/10.1101/2020.04.07.20056424>.

817 18. Capel RA, Herring N, Kalla M, et al. Hydroxychloroquine reduces heart rate by
818 modulating the hyperpolarization-activated current I_f : novel electrophysiological
819 insights and therapeutic potential. *Heart Rhythm* 2015;12:2186–2194.

820 19. Szekely Y, Lichter Y, Shrikie BA, Bruck H, Oster HS, Viskin S. Chloroquine-
821 induced torsades de pointes in a patient with coronavirus disease 19. Published online
822 May 5, 2020. *Heart Rhythm* <https://doi.org/10.1016/j.hrthm.2020.04.046>.

823 20. Ji YC, Uzelac I, Otani N, et al. Synchronization as a mechanism for low-energy
824 anti-fibrillation pacing. *Heart Rhythm* 2017;14:1254–1262.

825 21. Basu-Ray I, Almaddah NK, Adeboye A, Soos MP. Cardiac manifestations of
826 coronavirus (COVID-19). In: StatPearls. Treasure Island, FL: StatPearls Publishing;
827 2020.

828 22. Chen D Jr, Li X, Song Q Sr, Hu C Jr, Su F, Dai J. Hypokalemia and clinical im-
829 plications in patients with coronavirus disease 2019 (COVID-19). Preprint. Pub-
830 lished online February 29, 2020. medRxiv 2020.02.27.20028530. <https://doi.org/10.1101/2020.02.27.20028530>.

831 23. Kanai A, Salama G. Optical mapping reveals that repolarization spreads aniso-
832 tropically and is guided by fiber orientation in guinea pig hearts. *Circ Res*
833 1995;77:784–802.

834 24. Uzelac I, Ji YC, Hornung D, et al. Simultaneous quantification of spatially dis-
835 cordant alternans in voltage and intracellular calcium in Langendorff-perfused rabbit
836 hearts and inconsistencies with models of cardiac action potentials and ca trans-
837 sients. *Front Physiol* 2017;8:819.

838 25. Osadchii OE. Dofetilide promotes repolarization abnormalities in perfused
839 guinea-pig heart. *Cardiovasc Drugs Ther* 2012;26:489–500.

840 26. Osadchii OE. Quinidine elicits proarrhythmic changes in ventricular repolariza-
841 tion and refractoriness in guinea-pig. *Can J Physiol Pharmacol* 2013;
842 91:306–315.

843 27. Zhang Y, Xiao J, Lin H, et al. Ionic mechanisms underlying abnormal QT pro-
844 longation and the associated arrhythmias in diabetic rabbits: a role of rapid delayed
845 rectifier K^+ current. *Cell Physiol Biochem* 2007;19:225–238.

846 28. Cheng HC, Incardona J, McCullough B. Isolated perfused and paced guinea pig
847 heart to test for drug-induced changes of the QT interval. *J Pharmacol Toxicol
848 Meth* 2006;54:278–287.

849 29. Cheng HC, Incardona J. Models of torsades de pointes: effects of FPL64176,
850 DPI201106, dofetilide, and chromanol 293B in isolated rabbit and guinea pig
851 hearts. *J Pharmacol Toxicol Meth* 2009;60:174–184.

852 30. Saleh M, Gabriels J, Chang D, et al. The effect of chloroquine, hydroxychloro-
853 quine and azithromycin on the corrected QT interval in patients with SARS-
854 CoV-2 infection. Published online April 29, 2020. *Circ Arrhythm Electrophysiol*
855 <https://doi.org/10.1161/CIRCEP.120.008662>.

856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952