

SoilTemp: A global database of near-surface temperature

Jonas J. Lembrechts¹  | Juha Aalto^{2,3}  | Michael B. Ashcroft^{4,5}  | Pieter De Frenne⁶  |
 Martin Kopecký^{7,8}  | Jonathan Lenoir⁹  | Miska Luoto³ | Ilya M. D. Maclean¹⁰  |
 Olivier Roupsard^{11,12}  | Eduardo Fuentes-Lillo^{13,14,15} | Rafael A. García^{13,14} |
 Loïc Pellissier^{16,17} | Camille Pitteloud^{16,17} | Juha M. Alatalo^{18,19}  | Stuart W. Smith^{20,21}  |
 Robert G. Björk^{22,23}  | Lena Muffler^{24,25}  | Amanda Ratier Backes^{26,27} |
 Simone Cesarz^{27,28} | Felix Gottschall^{27,28}  | Joseph Okello^{29,30} | Josef Urban^{31,32}  |
 Roman Plichta³¹ | Martin Svátek³¹  | Shyam S. Phartyal^{33,34}  | Sonja Wipf^{35,36}  |
 Nico Eisenhauer^{27,28}  | Mihai Pușcaș³⁷  | Pavel D. Turtureanu³⁸  |
 Andrej Varlagin³⁹  | Romina D. Dimarco⁴⁰ | Alistair S. Jump⁴¹  | Krystal Randall⁴²  |
 Ellen Dorrepaal⁴³  | Keith Larson⁴³ | Josefina Walz⁴³  | Luca Vitale⁴⁴ |
 Miroslav Svoboda⁸  | Rebecca Finger Higgins⁴⁵ | Aud H. Halbritter⁴⁶ |
 Salvatore R. Curasi⁴⁷  | Ian Klupar⁴⁷ | Austin Koontz⁴⁸  | William D. Pearse^{48,49}  |
 Elizabeth Simpson⁴⁸ | Michael Stemkovski⁴⁸ | Bente Jessen Graae²⁰  |
 Mia Vedel Sørensen²⁰ | Toke T. Høye⁵⁰  | M. Rosa Fernández Calzado⁵¹ | Juan Lorite⁵¹  |
 Michele Carbognani⁵²  | Marcello Tomaselli⁵²  | T'ai G. W. Forte⁵²  |
 Alessandro Petraglia⁵² | Stef Haesen⁵³  | Ben Somers⁵³  | Koenraad Van Meerbeek⁵³  |
 Mats P. Björkman^{22,23} | Kristoffer Hylander⁵⁴ | Sonia Merinero⁵⁴ | Mana Gharun⁵⁵  |
 Nina Buchmann⁵⁵  | Jiri Dolezal^{7,56}  | Radim Matula⁸ | Andrew D. Thomas⁵⁷  |
 Joseph J. Bailey⁵⁸ | Dany Ghosn⁵⁹  | George Kazakis⁵⁹ | Miguel A. de Pablo⁶⁰  |
 Julia Kemppinen³  | Pekka Niittynen³  | Lisa Rew⁶¹  | Tim Seipel⁶¹ |
 Christian Larson⁶¹  | James D. M. Speed⁶²  | Jonas Ardö⁶³  | Nicoletta Cannone⁶⁴ |
 Mauro Guglielmin⁶⁵ | Francesco Malfasi⁶⁵ | Maaïke Y. Bader⁶⁶  | Rafaella Canessa⁶⁶  |
 Angela Stanisci⁶⁷  | Juergen Kreyling²⁴  | Jonas Schmeddes²⁴ | Laurenz Teuber²⁴ |
 Valeria Aschero^{68,69}  | Marek Čiliak⁷⁰  | František Máliš⁷¹  | Pallieter De Smedt⁶  |
 Sanne Govaert⁶  | Camille Meeussen⁶  | Pieter Vangansbeke⁶  | Khatuna Gigauri⁷² |
 Andrea Lamprecht⁷³  | Harald Pauli⁷³  | Klaus Steinbauer⁷³  | Manuela Winkler⁷³  |
 Masahito Ueyama⁷⁴  | Martin A. Nuñez⁷⁵ | Tudor-Mihai Ursu⁷⁶  | Sylvia Haider^{26,27}  |
 Ronja E. M. Wedegärtner²⁰  | Marko Smiljanic⁷⁷ | Mario Trouillier⁷⁷ | Martin Wilmking⁷⁷  |
 Jan Altman⁷  | Josef Brůna⁷  | Lucia Hederová⁷ | Martin Macek⁷  | Matěj Man⁷  |
 Jan Wild⁷  | Pascal Vittoz⁷⁸  | Meelis Pärtel⁷⁹ | Peter Barančok⁸⁰  |
 Róbert Kanka⁸⁰  | Jozef Kollár⁸⁰ | Andrej Palaj⁸⁰  | Agustina Barros⁶⁹ |
 Ana C. Mazzolari⁶⁹ | Marijn Bauters²⁹  | Pascal Boeckx²⁹ | José-Luis Benito Alonso⁸¹  |

Shengwei Zong⁸² | Valter Di Cecco⁸³  | Zuzana Sitková⁸⁴  | Katja Tielbörger⁸⁵ |
 Liesbeth van den Brink⁸⁵  | Robert Weigel²⁵  | Jürgen Homeier²⁵ |
 C. Johan Dahlberg^{54,86}  | Sergiy Medinets⁸⁷  | Volodymyr Medinets⁸⁷  |
 Hans J. De Boeck¹ | Miguel Portillo-Estrada¹  | Lore T. Verryckt¹  | Ann Milbau⁸⁸  |
 Gergana N. Daskalova⁸⁹  | Haydn J. D. Thomas⁸⁹  | Isla H. Myers-Smith⁸⁹  |
 Benjamin Blonder^{90,91} | Jörg G. Stephan⁹²  | Patrice Descombes^{16,17,93}  |
 Florian Zellweger⁹³  | Esther R. Frei^{35,93}  | Bernard Heinesch⁹⁴  |
 Christopher Andrews⁹⁵  | Jan Dick⁹⁵ | Lukas Siebicke⁹⁶ | Adrian Rocha⁹⁷ |
 Rebecca A. Senior⁹⁸  | Christian Rixen³⁵ | Juan J. Jimenez⁹⁹  | Julia Boike^{100,101}  |
 Aníbal Pauchard^{13,14} | Thomas Scholten¹⁰²  | Brett Scheffers¹⁰³  | David Klings¹⁰⁴  |
 Edmund W. Basham¹⁰⁴  | Jian Zhang¹⁰⁵  | Zhaochen Zhang¹⁰⁵ | Charly Geron^{1,106}  |
 Fatih Fazlioglu¹⁰⁷  | Onur Candan¹⁰⁷  | Jhonatan Sallo Bravo¹⁰⁸  | Filip Hrbacek¹⁰⁹  |
 Kamil Laska¹⁰⁹  | Edoardo Cremonese¹¹⁰  | Peter Haase^{111,112}  |
 Fernando E. Moyano⁹⁶  | Christian Rossi^{36,113,114}  | Ivan Nijs¹ 

¹Research Group PLECO (Plants and Ecosystems), University of Antwerp, Wilrijk, Belgium

²Finnish Meteorological Institute, Helsinki, Finland

³Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland

⁴Centre for Sustainable Ecosystem Solutions, School of Biological Sciences, University of Wollongong, Wollongong, NSW, Australia

⁵Australian Museum, Sydney, NSW, Australia

⁶Forest & Nature Lab, Department of Environment, Ghent University, Melle-Gontrode, Belgium

⁷Institute of Botany of the Czech Academy of Sciences, Průhonice, Czech Republic

⁸Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Prague 6 - Suchbát, Czech Republic

⁹UR 'Ecologie et Dynamique des Systèmes Anthropisés' (EDYSAN, UMR 7058 CNRS-UPJV), Univ. de Picardie Jules Verne, Amiens, France

¹⁰Environment and Sustainability Institute, University of Exeter, Penryn, UK

¹¹CIRAD, UMR Eco&Sols, Dakar, Senegal

¹²Eco&Sols, Univ Montpellier, CIRAD, INRAE, IRD, Institut Agro, Montpellier, France

¹³Laboratorio de Invasiones Biológicas (LIB), Facultad de Ciencias Forestales, Universidad de Concepción, Concepción, Chile

¹⁴Instituto de Ecología y Biodiversidad (IEB), Santiago, Chile

¹⁵School of Education and Social Sciences, Adventist University of Chile, Chile

¹⁶Landscape Ecology, Institute of Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zürich, Zürich, Switzerland

¹⁷Unit of Land Change Science, Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

¹⁸Department of Biological and Environmental Sciences, Qatar University, Doha, Qatar

¹⁹Environmental Science Center, Qatar University, Doha, Qatar

²⁰Department of Biology, Norwegian University of Science and Technology, Trondheim, Norway

²¹Asian School of Environment, Nanyang Technological University, Singapore, Singapore

²²Department of Earth Sciences, University of Gothenburg, Gothenburg, Sweden

²³Gothenburg Global Biodiversity Centre, Gothenburg, Sweden

²⁴Experimental Plant Ecology, Institute of Botany and Landscape Ecology, University of Greifswald, Greifswald, Germany

²⁵Plant Ecology, Albrecht-von-Haller-Institute for Plant Sciences, University of Goettingen, Goettingen, Germany

²⁶Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

²⁷German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

²⁸Institute of Biology, Leipzig University, Leipzig, Germany

²⁹Isotope Bioscience Laboratory – ISOFYS, Ghent University, Gent, Belgium

³⁰Mountains of the Moon University, Fort Portal, Uganda

³¹Department of Forest Botany, Dendrology and Geobiocoenology, Mendel University, Brno, Czech Republic

³²Siberian Federal University, Krasnoyarsk, Russia

- ³³School of Ecology and Environment Studies, Nalanda University, Rajgir, India
- ³⁴Department of Forestry and NR, H.N.B. Garhwal University, Srinagar-Garhwal, India
- ³⁵WSL Institute for Snow and Avalanche Research SLF, Davos, Switzerland
- ³⁶Swiss National Park, Chastè Planta-Wildenberg, Zerne, Switzerland
- ³⁷A. Borza Botanical Garden and Department of Taxonomy and Ecology, Faculty of Biology and Geology, Babeş-Bolyai University, Cluj-Napoca, Romania
- ³⁸A. Borza Botanical Garden, Babes-Bolyai University, Cluj-Napoca, Romania
- ³⁹A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia
- ⁴⁰Grupo de Ecología de Poblaciones de Insectos, IFAB (INTA - CONICET), Bariloche, Argentina
- ⁴¹Biological and Environmental Sciences, Faculty of Natural Sciences, University of Stirling, Stirling, UK
- ⁴²Centre for Sustainable Ecosystem Solutions, School of Earth, Atmospheric and Life Sciences, University of Wollongong, Wollongong, NSW, Australia
- ⁴³Climate Impacts Research Centre, Department of Ecology and Environmental Sciences, Umeå University, Abisko, Sweden
- ⁴⁴CNR - Institute for Mediterranean Agricultural and Forest Systems, Ercolano (Napoli), Italy
- ⁴⁵Dartmouth College, Hanover, NH, USA
- ⁴⁶Department of Biological Sciences and Bjerknes Centre for Climate Research, University of Bergen, Bergen, Norway
- ⁴⁷Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA
- ⁴⁸Department of Biology and Ecology Center, Utah State University, Logan, UT, USA
- ⁴⁹Department of Life Sciences, Imperial College London, Ascot, UK
- ⁵⁰Department of Bioscience and Arctic Research Centre, Rønde, Denmark
- ⁵¹Department of Botany, University of Granada, Granada, Spain
- ⁵²Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Parma, Italy
- ⁵³Department of Earth and Environmental Sciences, Leuven, Belgium
- ⁵⁴Department of Ecology, Environment and Plant Sciences and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden
- ⁵⁵Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland
- ⁵⁶Faculty of Science, Department of Botany, University of South Bohemia, České Budějovice, Czech Republic
- ⁵⁷Department of Geography and Earth Sciences, Aberystwyth University, Wales, UK
- ⁵⁸Department of Geography, York St John University, York, UK
- ⁵⁹Department of Geo-information in Environmental Management, Mediterranean Agronomic Institute of Chania, Chania, Greece
- ⁶⁰Department of Geology, Geography and Environment, University of Alcalá, Madrid, Spain
- ⁶¹Department of Land Resources and Environmental Sciences, Montana State University, Bozeman, MT, USA
- ⁶²Department of Natural History, NTNU University Museum, Norwegian University of Science and Technology, Trondheim, Norway
- ⁶³Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden
- ⁶⁴Department of Science and High Technology, Insubria University, Como, Italy
- ⁶⁵Department of Theoretical and Applied Sciences, Insubria University, Varese, Italy
- ⁶⁶Ecological Plant Geography, Faculty of Geography, University of Marburg, Marburg, Germany
- ⁶⁷EnvixLab, Dipartimento di Bioscienze e Territorio, Università degli Studi del Molise, Termoli, Italy
- ⁶⁸Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Cuyo, Cuyo, Argentina
- ⁶⁹Instituto Argentino de Nivología, Glaciología y Ciencias Ambientales (IANIGLA), CONICET, CCT-Mendoza, Mendoza, Argentina
- ⁷⁰Faculty of Ecology and Environmental Sciences, Technical University in Zvolen, Zvolen, Slovakia
- ⁷¹Faculty of Forestry, Technical University in Zvolen, Zvolen, Slovakia
- ⁷²Georgian Institute of Public Affairs, Tbilisi, Georgia
- ⁷³GLORIA Coordination, Institute for Interdisciplinary Mountain Research, Austrian Academy of Sciences (ÖAW) & Department of Integrative Biology and Biodiversity Research, University of Natural Resources and Life Sciences Vienna (BOKU), Vienna, Austria
- ⁷⁴Graduate School of Life and Environmental Sciences, Osaka Prefecture University, Osaka, Japan
- ⁷⁵Grupo de Ecología de Invasiones, INIBIOMA, CONICET/Universidad Nacional del Comahue, Bariloche, Argentina
- ⁷⁶Institute of Biological Research Cluj-Napoca, National Institute of Research and Development for Biological Sciences, Bucharest, Romania
- ⁷⁷Institute of Botany and Landscape Ecology, University Greifswald, Greifswald, Germany
- ⁷⁸Institute of Earth Surface Dynamics, Faculty of Geosciences and Environment, University of Lausanne, Lausanne, Switzerland
- ⁷⁹Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia
- ⁸⁰Institute of Landscape Ecology, Slovak Academy of Sciences, Bratislava, Slovakia
- ⁸¹Jolube Consultor Botánico, Jaca, Spain
- ⁸²Key Laboratory of Geographical Processes and Ecological Security in Changbai Mountains, Ministry of Education, School of Geographical Sciences, Northeast Normal University, Changchun, China
- ⁸³Majella Seed Bank, Majella National Park, Lama dei Peligni, Italy

- ⁸⁴National Forest Centre, Forest Research Institute Zvolen, Zvolen, Slovakia
- ⁸⁵Plant Ecology Group, Department of Evolution and Ecology, University of Tübingen, Tübingen, Germany
- ⁸⁶County Administrative Board of Västra Götaland, Gothenburg, Sweden
- ⁸⁷Regional Centre for Integrated Environmental Monitoring, Odesa National I.I. Mechnikov University, Odesa, Ukraine
- ⁸⁸Research Institute for Nature and Forest (INBO), Brussels, Belgium
- ⁸⁹School of GeoSciences, University of Edinburgh, Edinburgh, UK
- ⁹⁰School of Life Sciences, Arizona State University, Tempe, AZ, USA
- ⁹¹Department of Environmental Science, Policy, and Management, University of California, Berkeley, Berkeley, CA, USA
- ⁹²Swedish Species Information Centre, Swedish University of Agricultural Sciences, Uppsala, Sweden
- ⁹³Swiss Federal Research Institute WSL, Birmensdorf, Switzerland
- ⁹⁴TERRA Teaching and Research Center, Faculty of Gembloux Agro-Bio Tech, University of Liege, Gembloux, Belgium
- ⁹⁵UK Centre for Ecology & Hydrology, Midlothian, UK
- ⁹⁶Bioclimatology, University of Goettingen, Göttingen, Germany
- ⁹⁷Department of Biological Sciences and the Environmental Change Initiative, University of Notre Dame, Notre Dame, IN, USA
- ⁹⁸Woodrow Wilson School of Public and International Affairs, Princeton University, Princeton, NJ, USA
- ⁹⁹ARAIID Research and Development, Zaragoza, Spain
- ¹⁰⁰Alfred Wegener Institute Helmholtz Center for Polar and Marine Research, Potsdam, Germany
- ¹⁰¹Geography Department, Humboldt-Universität zu Berlin, Berlin, Germany
- ¹⁰²Chair of Soil Science and Geomorphology, Department of Geosciences, University of Tuebingen, Tuebingen, Germany
- ¹⁰³Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL, USA
- ¹⁰⁴School of Natural Resources and Environment, University of Florida, Gainesville, FL, USA
- ¹⁰⁵Zhejiang Tiantong Forest Ecosystem National Observation and Research Station, School of Ecological and Environmental Sciences, East China Normal University, Shanghai, China
- ¹⁰⁶Biodiversity and Landscape, TERRA Research Centre, University of Liège, Gembloux Agro-Bio Tech, Gembloux, Belgium
- ¹⁰⁷Faculty of Arts and Sciences, Department of Molecular Biology and Genetics, Ordu University, Ordu, Turkey
- ¹⁰⁸Universidad Nacional de San Antonio Abad del Cusco, Cusco, Peru
- ¹⁰⁹Department of Geography, Masaryk University, Brno, Czech Republic
- ¹¹⁰Climate Change Unit, Environmental Protection Agency of Aosta Valley, Aosta, Italy
- ¹¹¹Senckenberg Research Institute and Natural History Museum Frankfurt, Gelnhausen, Germany
- ¹¹²Faculty of Biology, University of Duisburg-Essen, Essen, Germany
- ¹¹³Remote Sensing Laboratories, Department of Geography, University of Zurich, Zurich, Switzerland
- ¹¹⁴Research Unit Community Ecology, Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland

Correspondence

Jonas J. Lembrechts, Research Group
PLECO (Plants and Ecosystems), University
of Antwerp, 2610 Wilrijk, Belgium.
Email: jonas.lembrechts@uantwerpen.be

Funding information

Fonds Wetenschappelijk Onderzoek, Grant/
Award Number: 12P1819N and WOG
W001919N; European Union FP-5 project
GLORIA-Europe, Grant/Award Number:
EVK2-CT-2000-0006; Swiss MAVA
Foundation project; Swiss Federal Office
for the Environment (FOEN); Foundation Dr.
Joachim de Giamoni; Research Commission
and Staff of the Swiss National Park;
Flexible Pool project, Grant/Award Number:
W47014118; German Centre for Integrative
Biodiversity Research (iDiv) Halle-Jena-
Leipzig; University of Alcalá; Fondation
Mariétan, Société académique de Genève;
Swiss Federal Office of Education and
Science; Czech Science Foundation, Grant/
Award Number: 17-13998S, 17-07378S,
20-05840Y and 17-19376S; Czech Academy

Abstract

Current analyses and predictions of spatially explicit patterns and processes in ecology most often rely on climate data interpolated from standardized weather stations. This interpolated climate data represents long-term average thermal conditions at coarse spatial resolutions only. Hence, many climate-forcing factors that operate at fine spatiotemporal resolutions are overlooked. This is particularly important in relation to effects of observation height (e.g. vegetation, snow and soil characteristics) and in habitats varying in their exposure to radiation, moisture and wind (e.g. topography, radiative forcing or cold-air pooling). Since organisms living close to the ground relate more strongly to these microclimatic conditions than to free-air temperatures, microclimatic ground and near-surface data are needed to provide realistic forecasts of the fate of such organisms under anthropogenic climate change, as well as of the functioning of the ecosystems they live in. To fill this critical gap, we highlight a call for temperature time series submissions to SoilTemp, a geospatial database initiative compiling soil and near-surface temperature data from all over the world. Currently, this database contains time series from 7,538 temperature sensors from 51 countries

of Sciences, Grant/Award Number: RVO 67985939; Estonian Research Council, Grant/Award Number: PRG609; European Regional Development Fund; DFG GraKo 2010 Response; Qatar Petroleum, Grant/Award Number: QUEX-CAS-QP-RD-18/19; Ministry of Education and Science of Ukraine; Toward INMS; Swiss National Science Foundation, Grant/Award Number: 172198; Slovak Scientific Grant Agency, Grant/Award Number: VEGA 2/0132/18; Lund University; University of Helsinki; Ministry of Education, Youth and Sport of the Czech Republic, program Inter-Excellence, subprogram Inter-Action, Grant/Award Number: LTAUSA19137 and LTAUSA18007; Carlsberg Foundation, Grant/Award Number: CF16-0896; Villum Foundation, Grant/Award Number: 17523; German Research Foundation, Grant/Award Number: FZT 118; EU Horizon 2020, Grant/Award Number: 641918; Natural Environmental Research Council, Grant/Award Number: NE/L002558/1 and NE/M016323/1; UK Natural Environmental Research Council ShrubTundra, Grant/Award Number: NE/M016323/1; Ministry of Education, Youth and Sports of the Czech Republic, Grant/Award Number: INTER-TRANSFER LTT17017; National Institute of Food and Agriculture, Grant/Award Number: MONB00363 and 2017-70006-27272; Slovak Research and Development Agency; National Geographic Society, Grant/Award Number: 9480-14 and WW-240R-17; Research Council of Norway, Grant/Award Number: 262064; National Science Foundation, Grant/Award Number: ABI-1759965 and EF-1802605; Leverhulme Trust Research Fellowship; National Natural Science Foundation of China, Grant/Award Number: 41971124; Mendel University; Ministry of Youth and Sports of the Czech Republic; Ministry of Research and Innovation; RFBR, Grant/Award Number: 19-04-01234-a; European Research Council, Grant/Award Number: ERC-2562013-SyG-610028; Agence Nationale de la Recherche (ANR), Grant/Award Number: ANR-19-CE32-0005-01

across all key biomes. The database will pave the way toward an improved global understanding of microclimate and bridge the gap between the available climate data and the climate at fine spatiotemporal resolutions relevant to most organisms and ecosystem processes.

KEYWORDS

climate change, database, ecosystem processes, microclimate, soil climate, species distributions, temperature, topoclimate

1 | INTRODUCTION

Current ecological research increasingly deals with large-scale patterns and processes, with global databases of species distributions and traits becoming increasingly available (Bruehlheide et al., 2018; Kattge et al., 2019; Kissling et al., 2018). Analyses of these patterns and processes—and their predictions under anthropogenic climate change—often rely on global climatic grids at coarse spatial resolutions interpolated from standardized weather stations that represent long-term average atmospheric conditions (Lembrechts, Nijs, & Lenoir, 2019). Moreover, sensors in these weather stations

are shielded from direct solar radiation and located at ~2 m above a frequently mown lawn (free-air temperature or “macroclimate,” Jarraud, 2008). These climatic grids thus ignore many climate-forcing processes that operate near the ground surface, at fine spatiotemporal resolutions, and in environments that vary in their exposure to winds, radiation and moisture (“microclimate,” Bramer et al., 2018; Daly, 2006; Körner & Hiltbrunner, 2018). Importantly, while these microclimatic processes often operate at fine spatiotemporal resolutions, they can affect ecological relations both at the local and the global scales (Ashcroft, Cavanagh, Eldridge, & Gollan, 2014; De Frenne et al., 2013; Lembrechts, Lenoir et al., 2019). For example,

they can potentially protect ground-dwelling biota against long-term climate variability, providing microrefugia for these species to survive in locations deemed unsuitable in models using climate data at coarse spatial resolutions, or buffer organisms against short-term extreme events (Bramer et al., 2018; De Frenne et al., 2013; Lenoir, Hattab, & Pierre, 2017; Suggitt et al., 2018). Microclimates can however also expose organisms to more extreme temperatures, in which case distribution models that ignore such microclimates may erroneously predict species survival instead of extinction (Pincebourde & Casas, 2019). To provide realistic forecasts of species distributions and performance, as well as of the functioning of the ecosystems they operate in, climate data that incorporate microclimatic processes, ideally measured *in situ*, are thus urgently needed (Körner & Hiltbrunner, 2018).

2 | HORIZONTAL AND VERTICAL FEATURES DRIVING MICROCLIMATE

The offset between micro- and macroclimate is particularly pronounced around the soil surface, as temperatures measured at 2 m above the ground can differ substantially from those at ground level, or in the layers just above and below it (Geiger, 1950; Lembrechts, Lenoir et al., 2019). This offset can result from both “horizontal” and “vertical” features (Figure 1), and can exceed several degrees centigrade in annual averages. For example, Kearney (2019) modeled coarse-scale soil temperatures at various depths considering the vertical features affecting the radiation balance. These vertical features include the effects of vegetation characteristics (e.g. structure and cover), snow cover and soil characteristics (e.g. moisture content, geological types, texture and bulk density; Lembrechts, Lenoir et al., 2019; Li, 1926; Zhang, Wang, Barr, & Black, 2008). The result of these vertical features is not only an instantaneous temperature offset between air and soil temperatures but also a buffering effect, that is, the temporal variability in temperature changes is lower in the soil than in the air (Ashcroft & Gollan, 2013; Geiger, 1950). Horizontal processes, on the other hand, relate more to the spatial resolution of the climatic data. They can be broken up into those that require only fine-resolution environmental information for specific sites (e.g. effects of slope and aspect on radiation balances; Bennie, Huntley, Wiltshire, Hill, & Baxter, 2008), and those where temperatures are also affected by neighboring locations (e.g. topographic shading, cold-air drainage and atmospheric temperature inversions, which are landscape context dependent; Ashcroft & Gollan, 2012; Whiteman, 1982).

How horizontal and vertical features interact to define differences between soil and air temperature may differ with the biome, season and day time. For example, in grasslands during summer, incoming short-wave solar radiation is usually the dominant factor determining daytime soil surface temperatures, which, in turn, result in higher air temperatures through convective heating (Geiger, 1950). However, during winter, horizontal processes such as cold-air drainage and coastal buffering can have larger effects, especially on overnight air temperatures, when air temperatures may be driving

soil temperatures rather than vice-versa (Vitasse, Klein, Kirchner, & Rebetez, 2017). In dense forests, the situation is even more complex: upper canopies block the bulk of short-wave solar radiation such that sub-canopy temperatures are determined by convective heat transfer between the air surrounding the canopy and direct conductance through physical contact of different parts of the canopy layer, in addition to the limited radiation that does permeate the canopy (Körner & Paulsen, 2004; Lenoir et al., 2017; Zellweger, De Frenne, Lenoir, Rocchini, & Coomes, 2019). As a result, horizontal processes such as passing fronts, and winds blowing in hotter or colder air from outside the forest, will in large part define the—dampened—temperature patterns under forest canopies (Ashcroft, Chisholm, & French, 2008).

3 | THE NEED FOR MICROCLIMATE DATA ACROSS THE FIELD OF ECOLOGY

Many organisms living in the soil and close to the soil surface (e.g. soil micro-organisms like fungi, ground arthropods, herbs, mosses, tree seedlings and small vertebrates) only experience fine-scale soil and/or near-surface temperatures, and thus likely relate less strongly to free-air temperatures (Lembrechts, Lenoir et al., 2019; Niittynen & Luoto, 2017; Randin, Vuissoz, Liston, Vittoz, & Guisan, 2009). This may be reflected in a species' distribution but also their morphology, physiology and behavior (de Boeck, Velde, Groote, & Nijs, 2016; Kearney, Shine, & Porter, 2009; Körner & Paulsen, 2004; Opedal, Armbruster, & Graae, 2015). Many species indeed survive, live and reproduce where average background climate appears unsuitable, and equally may be gone from sites within apparently suitable areas where microclimatic extremes exceed their limits (Suggitt et al., 2011). Without microclimate data, we not only lack information on the potential thermal heterogeneity that is available for species to thermoregulate *in situ* but also on the true magnitude of climate change that species will be exposed to (Maclean, Suggitt, Wilson, Duffy, & Bennie, 2017; Pincebourde, Murdock, Vickers, & Sears, 2016). Accurately predicting how species' ranges will shift under climate change requires a good understanding of the variety of climate niches truly available to them (Lenoir et al., 2017; Maclean, Hopkins, Bennie, Lawson, & Wilson, 2015). The latter requires both a good understanding of what defines current microclimates and of how climate change will interact with the drivers of microclimatic conditions (Maclean, 2019). Additionally, it is the soil temperature rather than the air temperature that defines many ecosystem functions in and close to the soil, like evapotranspiration, decomposition, root growth, biogeochemical cycling and soil respiration (Gottschall et al., 2019; Hursh et al., 2017; Medinets, Gasche, Kiese, Rennenberg, & Butterbach-Bahl, 2019; Pleim & Gilliam, 2009; Portillo-Estrada et al., 2016). Given the repeatedly proven sensitivity of many of these processes to temperatures (Coûteaux, Bottner, & Berg, 1995; Rosenberg, Kimball, Martin, & Cooper, 1990; Schimel et al., 1996), here again having accurate measurements will be of utmost importance. The carbon balance in boreal forests, for example, is largely dependent on soil thaw and thus soil rather than air temperatures (Goulden et al., 1998).




FIGURE 1 The horizontal and vertical drivers of the offset between in-situ soil and free-air temperatures. Conceptually, there are two different sets of features responsible for the offset between coarse-scale free air temperatures (top left, e.g. WorldClim, Fick & Hijmans, 2017) and fine-scale soil temperatures (bottom right, e.g. Ashcroft & Gollan, 2012; Lembrechts, Lenoir et al., 2019). First, one can incorporate fine-scale horizontal climate-forcing factors such as topography and terrain-related features, land cover types and distance to water bodies to go from coarse-scaled to finer resolutions (top right, e.g. Aalto, Riihimäki, Meineri, Hylander, & Luoto, 2017; Macek, Kopecký, & Wild, 2019). Second, one can consider observation height, and the effects of vegetation characteristics (like structure and cover), snow cover and soil characteristics (like moisture, geological types, texture and bulk density) on the radiation balance to convert from free-air to soil temperatures (e.g. Kearney, 2019). Both horizontal and vertical features can introduce positive or negative differences (offset values) between soil and air temperatures through their effects on processes related to the radiation balance, like wind, convective heat transfer and surface albedo. The complexities of these horizontal and vertical processes can vary with biome, season and time of day. Temperatures are represented here using an unspecified temperature range from cold (blue) to warm (red)

These realizations highlight the urgency to start using soil and near-surface microclimate data when modeling the ecology and biogeography of surface and soil-dwelling organisms, as well as the functioning of soil ecosystems, instead of readily available coarse-scaled free-air climate data (from e.g. CHELSA [Karger et al., 2017],

TerraClimate [Abatzoglou, Dobrowski, Parks, & Hegewisch, 2018] or WorldClim [Fick & Hijmans, 2017]). While a suit of models now exist that produce fine-scale climate data (Bramer et al., 2018; Lembrechts, Nijs et al., 2019), we do not yet fully understand whether models using data that represent average conditions over

large areas provide adequate “mean field approximations” of (i.e. are representative for) more complex spatiotemporal effects driven by the climatic conditions that organisms experience (Bennie, Wilson, Maclean, & Suggitt, 2014). To accomplish the latter, global in-situ data are needed for large-scale fine-resolution calibration and validation of these models. However, while the quality and resolution of free-air temperature data and models at the global scale are rapidly improving (Bramer et al., 2018), soil temperature datasets used in biogeography and biogeochemistry are still largely restricted to the landscape or regional scale, at best, and from intensively studied regions only (Aalto, Scherrer, Lenoir, Guisan, & Luoto, 2018; Ashcroft et al., 2008; Ashcroft, Chisholm, & French, 2009; Carter et al., 2015), or they are derived from models lacking fine-grained ground-truthing data (e.g. Copernicus Climate Change Service (C3s), 2019). Land surface temperatures as obtained from satellite data, on the other hand, are hampered by their inability to measure below the vegetation cover (Bramer et al., 2018).

To accurately describe and predict the (future) distribution and/or traits of surface and soil-dwelling species at larger scales, we need to improve our general knowledge of the offsets and spatiotemporal changes in variability between soil-level and free-air temperatures (Aalto et al., 2018; Lembrechts, Lenoir et al., 2019). There is an urgent need to work toward globally available soil and near-surface temperature data based on in-situ measurements and at relevant spatiotemporal resolutions (Ashcroft & Gollan, 2012; Meineri & Hylander, 2017; Opedal et al., 2015; Pradervand, Dubuis, Pellissier, Guisan, & Randin, 2014; Slavich, Warton, Ashcroft, Gollan, & Ramp, 2014).

4 | LAUNCH OF THE SoilTemp DATABASE

To tackle these issues, we launch an ambitious database initiative, compiling soil and near-surface temperature data from all over the world into a global geospatial database: SoilTemp. At the time of writing, we brought together temperature data from 7,538 sensors placed both below, at and above (up to 2 m) the soil surface (Figure 2a), which is an accumulation of over 180,000 months of temperature data with measurement intervals between 1 and 240 min (>30% every 60 min). The database hosts loggers from 51 different countries spread across all continents, with a broad distribution across the world's climatic space (Figure 2b). There is a dominance of time series from Europe and areas below 1,500 m a.s.l. (Figure 2c,d). More than 75% of sensor measurements occurred within the last decade, but the database does contain several time series covering longer time periods as well, with a maximum of 42 years (Figure 2d).

When the remaining critical gaps in our spatial coverage will be filled (see below), this database will allow global assessments of the long-established theories on boundary layer climatology in heterogeneous environments (Geiger, 1950), which has so far been lacking. The growing database provides a unique opportunity to disentangle the role of the different horizontal and vertical features influencing soil and near-surface temperature across all biomes of the world,

with high spatial and temporal resolutions. It will allow relating patterns in soil temperature to processes in the lower air layers and calibrate and validate global models of soil temperature and (micro) climate (Carter et al., 2015; Kearney, Isaac, & Porter, 2014; Kearney, Shamakh, et al., 2014; Maclean et al., 2017). It will also allow us to create global maps of a wide array of general and microclimate-specific bioclimatic variables (e.g. growing degree days, growing season length) at relevant spatiotemporal resolutions (Körner & Hiltbrunner, 2018).

Ultimately, this joint global effort and the resulting global microclimatic products will enable us to improve analyses of the relationships between species' macroecology and the microclimate they experience, identify microrefugia and stepping stones and improve global models of ecosystem functioning and element cycling. Indeed, replacing the coarse-scaled free-air temperature averages used traditionally in models in all fields of ecology with these more relevant soil-specific data products is likely to increase their descriptive and predictive power, as the countless above-mentioned regional studies exemplify (Lembrechts, Lenoir et al., 2019). Additionally, this first global effort to combine and collect in-situ measurements will help solve long-standing issues regarding sensor comparability and data collection variability (Bramer et al., 2018), as well as address the question at what spatial scale microclimate data can prove most informative for ecological modeling (Jucker et al., 2020). The temperature time series in the database, many of which are covering increasingly long time periods of up to a decade or more, will also allow fine-tuning forecasts of microclimate data into the future by deepening our understanding of the link between microclimatic dynamics in the soil and the air (Bramer et al., 2018; Lenoir et al., 2017; Maclean, 2019; Wason, Bevilacqua, & Dovciak, 2017), improving our predictions of biodiversity and ecosystem functioning under climate change.

5 | DIG OUT YOUR LOGGERS! A CALL FOR CONTRIBUTIONS

To reach these goals, we encourage scientists owning in-situ measured temperature data to submit these to the growing SoilTemp database. All time series spanning 1 month or more, with temperature measurements a maximum of 4 hr apart, all soil depths, all heights above the ground up till 2 m, all biomes, and all sensor types and brands will be accepted. Note that both spatially dense and sparse logger networks, as well as single loggers are accepted. The achieved spatial resolution is dependent on the provision of spatially precise coordinates to achieve a good relationship with potential explanatory variables (e.g. high-resolution remotely sensed environmental data). If we have these coordinates and thus the location and distance between loggers, we can effectively obtain the extent and spacing for each logger network (Western, Grayson, & Blöschl, 2002).

We include data from both observational and experimental plots, yet sensors have to be measuring in-situ and not in pots, and experiments manipulating the local climate (e.g. open-top chambers, rain-out shelters or vegetation-removal experiments) are excluded




FIGURE 2 Overview of the status of the SoilTemp-database as of March 2020. Spatial (a), climatic (b), elevational (c) and temporal (d) distribution of sensors in the SoilTemp-database as of March 2020. (a) Background world map in WGS1984, hexagons with a resolution of approximately 70,000 km² using the *dggridR*-package in R. (b) Colors of hexagons indicate the number of sensors at each climatic location, with a 40 × 40 bin resolution. Small dots in the background represent the global variation in climatic space (obtained by sampling 1,000,000 random locations from the CHELSA world maps at a spatial resolution of 2.5 arc minutes). Overlay with dotted lines and numbers (from 1 to 9) depict a delineation of Whittaker biomes (adapted from Whittaker, 1975): (1) tundra and ice, (2) boreal forest, (3) temperate seasonal forest, (4) temperate rainforest, (5) tropical rainforest, (6) tropical seasonal forest/savanna, (7) subtropical desert, (8) temperate grassland/desert, (9) woodland/shrubland. (c) Number of sensors in each elevation class. (d) Time span covered by each sensor in the database, ranked by starting date. Data showed from 1992 onwards, note that the time period covered by four loggers with starting dates in 1976 is truncated

(Table 1). Given currently less well-represented climate regions, we especially encourage submissions from extreme cold and hot environments to fill the remaining gaps in our global coverage. More specifically, hot tropical climates (both tropical rainforests and tropical seasonal forests and savannas) and cold and hot deserts are currently still largely underrepresented (Figure 2b), in particular from

Africa, Asia, Antarctica and the Americas (Figure 2a). Data contributors will be invited as co-authors on the main global papers resulting from this database (see Supporting Information for details on terms of use and data ownership).

By encouraging sampling and submissions from remote areas, we aim to help solve the global sampling bias in soil ecological

TABLE 1 Minimal data requirements and obligatory metadata for submission to the database. For more details, see Supporting Information

Minimum data requirements	Obligatory metadata
Minimum one consecutive month of in-situ measured temperature time series	Accurate (handheld GPS or finer) spatial coordinates of the loggers (+ estimated accuracy)
Maximum time interval between measurements: 4 hr	Height/depth of the sensor relative to the soil surface
No climate manipulation experiments (only control plots of those experiments or observational studies)	Type or brand of temperature sensor used, and type of shelter (e.g. no shelter, home-made shelter, Stevenson screen, etc.)
No modeling studies (only empirical data)	Temporal resolution of the sensor
	Habitat classification

data (Cameron et al., 2018; Guerra et al., 2019), and we hope to build a truly global network representing—and actively engaging—scientists from a wide diversity of cultural backgrounds (Maestre & Eisenhauer, 2019). More information is available on the SoilTemp website, accessible via Figshare (<https://doi.org/10.6084/m9.figshare.12126516>).

When fully established, the SoilTemp database and its derivative products (e.g. bioclimatic variables) will be made freely available to facilitate the analysis of global patterns in microclimates, increase the comparability between regional studies and simplify the use of accurate microclimatic data in ecology (Bramer et al., 2018). At the moment, critical metadata are already freely accessible via Figshare (<https://doi.org/10.6084/m9.figshare.12126516>). Given the absence of and the need for globally available soil microclimate data products at relevant spatial resolutions for use in ecological analyses, we believe that SoilTemp has the potential to become a highly important resource that will enable a step change in ecological modeling.

ACKNOWLEDGEMENTS

We thank Sylvain Pincebourde and an anonymous reviewer for their critical evaluation of our manuscript. This work was supported by the Research Foundation Flanders (FWO) through a postdoctoral fellowship to Jonas J. Lembrechts (12P1819N) and a Research Network Grant (WOG W001919N). We gratefully acknowledge all data contributors, all staff of the author institutions engaged in field measurements and equipment maintenance (namely Erik Herberg, Iris Hamersveld, Ida Westman, Fredrik Brounes, Pernille Eidsen, Eleanor Walker and the teachers participating in the Tepåseförsöket 2015) and the ILTER-network, and thank local peoples for permission to collect data on their lands. Temperature data collection on European GLORIA summits was funded by European Union FP-5 project GLORIA-Europe (EVK2-CT-2000-0006) and the Swiss MAVA Foundation project “Climate change impacts in protected areas of the Alps and high mountains of Eastern Europe and the Mediterranean region,” on the Eastern Swiss GLORIA summits by the Swiss Federal Office for the Environment (FOEN), the Research

Commission and Staff of the Swiss National Park, and the Foundation Dr. Joachim de Giamoni, on Tenerife in the framework of the Flexible Pool project (W47014118) of Sylvia Haider funded by the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, on Livingston Island, Antarctica by different research projects of the Gubern of Spain (PERMAPLANET CTM2009-10165-E; ANTARPERMA CTM2011-15565-E; PERMASNOW CTM2014-52 021-R), and the PERMATHERMAL arrangement between the University of Alcalá and the Spanish Polar Committee and on the Western Swiss GLORIA summits by Département de la culture et des sports du Valais, Fondation Mariétan, Société académique de Genève, Swiss Federal Office of Education and Science and Swiss Federal Office for the Environment. Jan Wild, Martin Macek, Martin Kopecký, Lucia Hederová, Matěj Man and Josef Brůna were supported by the Czech Science Foundation (project 17-13998S) and the Czech Academy of Sciences (project RVO 67985939), Meelis Pärtel by an Estonian Research Council grant (PRG609) and by the European Regional Development Fund (Centre of Excellence EcolChange), Lena Muffler, Juergen Kreyling, Robert Weigel, Mario Trouillier, Martin Wilmking and Jonas Schmeddes by DFG GraKo 2010 Response, Juha M. Alatalo by Qatar Petroleum (QUEX-CAS-QP-RD-18/19), the authors from Odesa National I. I. Mechnikov University (Sergiy Medinets and Volodymyr Medinets) by EU FP6 The nitrogen cycle and its influence on the European greenhouse gas balance (NitroEurope), EU FP7 Effects of Climate Change on Air Pollution Impacts and Response Strategies for European Ecosystems (ÉCLAIRE), Ukrainian national research projects (No. 505, 550, 574) funded by Ministry of Education and Science of Ukraine and GEF-UNEP funded “Toward INMS” project, see www.inms.international for more details. Florian Zellweger was supported by the Swiss National Science Foundation (grant no. 172198), Peter Barančok, Róbert Kanka, Jozef Kollár and Andrej Palaj by the Slovak Scientific Grant Agency (project VEGA 2/0132/18), Jonas Ardö by a infrastructure grant from the Faculty of Science, Lund University, Julia Kempinen by the Doctoral Programme in Geosciences at the University of Helsinki, Jan Altman by the Czech Science Foundation (projects 17-07378S and 20-05840Y), the Czech Academy of Sciences (project RVO 67985939) and Ministry of Education, Youth and Sport of the Czech Republic, program Inter-Excellence, subprogram Inter-Action (project LTAUSA19137), Toke Thomas Høye by the Carlsberg Foundation (grant no. CF16-0896) and the Villum Foundation (grant no. 17523), Jiri Dolezal by the Czech Science Foundation (project 17-19376S), and Ministry of Education, Youth and Sport of the Czech Republic, program Inter-Excellence, subprogram Inter-Action (project LTAUSA18007), Nico Eisenhauer, Felix Gottschall and Simone Cesarz by the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation (FZT 118), Stuart W. Smith by AfricanBioServices project funded by the EU Horizon 2020 grant number 641918, Haydn Thomas by a K Natural Environmental Research Council doctoral training partnership grant NE/L002558/1, Isla H. Myers-Smith by the UK Natural Environmental Research Council ShrubTundra Project NE/M016323/1, Anibal Pauchard,

Rafael Garcia and Eduardo Fuentes-Lillo by the projects Fondecyt 1180205, Fondecyt 11170516 and CONICYT PIA APOYO CTE AFB170008, Rafaella Canessa, Maaïke Y. Bader, Liesbeth van den Brink, and Katja Tielbörger by the DFG Priority Programme 1803 EarthShape (projects 1 (BA 3843/6-1) and 11 (TI 338/14-1&2)), Martin Svátek by a grant from the Ministry of Education, Youth and Sports of the Czech Republic (grant number: INTER-TRANSFER LTT17017), Mihai Pușcaș by ODYSSEE project (ANR-13-ISV7-0004 France, PN-II-ID-JRP-RO-FR-2012 UEFISCDI Romania), Pavel Dan Turtureanu by UEFISCDI in Romania, MEMOIRE grant no. PN-III-P1-1.1-PD2016-0925, Jonathan Lenoir by the Agence Nationale de la Recherche (ANR) within the framework of the IMPRINT project "IMpacts des PRocessus microclimatiques sur la redistribution de la biodiversité forestière en contexte de réchauffement du macroclimat" (grant number: ANR-19-CE32-0005-01), Radim Matula and Roman Plichta by a grant Inter-Excellence (project: INTER-TRANSFER LTT17033) from the Ministry of Education, Youth and Sports of the Czech Republic, Lisa Rew by the National Institute of Food and Agriculture, U.S. Department of Agriculture Hatch MONB00363, Tim Seipel and Christian Larson by a grant from the United States National Institute of Food and Agriculture grant 2017-70006-27272, Nina Buchmann by the SNF (projects M4P 40FA40_154245, ICOS-CH 20FI21_148992, 20FI20_173691, InnoFarm 407340_172433) and the EU (SUPER-G contract no. 774124) for the Swiss FluxNet, Mana Gharun by the SNF project ICOS-CH Phase 2 20FI20_173691, Sanne Govaert by the Research Foundation Flanders (FWO; project GOH1517N), Pieter De Frenne, Camille Meeussen, and Pieter Van Gansbeke by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC Starting Grant FORMICA 757833), Olivier Roupsard by EU-LEAP-Agri (RAMSES II), Agropolis and Total Foundation (DSCATT), CGIAR (GLDC) and EU-DESIRA (CASSECS), Zuzana Sitková by the Slovak Research and Development Agency under the project No. APVV-16-0325 and project ITMS 26220220066 co-funded by the ERDF, Brett Ryan Scheffers by National Geographic Society (grant no. 9480-14 and WW-240R-17), James D. M. Speed by the Research Council of Norway (262064), William D. Pearse and the Pearse Lab by National Science Foundation ABI-1759965, NSF EF-1802605 and United States Department of Agriculture Forest Service agreement 18-CS-11046000-041, Isla H. Myers-Smith by the UK Natural Environmental Research Council ShrubTundra Project NE/M016323/1, Andrew D Thomas by a Leverhulme Trust Research Fellowship under Government of Botswana permit EWT8/36/4 VIII(4), Shengwei Zong by National Natural Science Foundation of China (no. 41971124), Roman Plichta by the post-doc project 7.3 of Institutional plan of Mendel University in Brno 2019–2020, František Máliš by the Slovak Research and Development Agency project APVV-15-0270, Filip Hrbacek and Kamil Laska by the projects LM2015078 and CZ.02.01/0.0/0.0/16_013/0001708 of Ministry of Youth and Sports of the Czech Republic, T-M. Ursu was supported by the Ministry of Research and Innovation through Projects for Excellence Financing in RDI: Contract no. 22 PFE/2018 and PN2019-2022/19270201

– Ctr. 25N BIODIVERS 3-BIOSERV and Andrej Varlagin by RFBR project number 19-04-01234-a. Lore T. Verryckt is funded by a PhD fellowship from the Research Foundation Flanders (FWO) and acknowledges support from the European Research Council Synergy Grant; ERC-2562013-SyG-610028 IMBALANCE-P and Pallieter De Smedt holds a postdoctoral fellowship of the Research Foundation-Flanders (FWO) and The Kreinitz Experiment is a cooperative research project initiated by the Helmholtz Centre for Environmental Research - UFZ. We also acknowledge project 18-74-10048 from the Russian Science Foundation, the Dirección General de Cambio Climático del Gobierno de Aragón, the Ordesa y Monte Perdido National Park and the Servicio de Medio Ambiente de Soria de la Junta de Castilla y León, the National Swiss Fund for research (SNSF, project "Lif3web", no. 162604).

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

AUTHOR CONTRIBUTION

J.J.L. performed the analyses and wrote the manuscript, J.J.L., J.A., M.B.A., P.D.F., M.K., J.L., M.L., I.M.D.M. and I.N. led the consortium and contributed to the writing; all authors contributed to the consortium and provided editorial advice.

ORCID

Jonas J. Lembrechts  <https://orcid.org/0000-0002-1933-0750>
 Juha Aalto  <https://orcid.org/0000-0001-6819-4911>
 Michael B. Ashcroft  <https://orcid.org/0000-0003-2157-5965>
 Pieter De Frenne  <https://orcid.org/0000-0002-8613-0943>
 Martin Kopecký  <https://orcid.org/0000-0002-1018-9316>
 Jonathan Lenoir  <https://orcid.org/0000-0003-0638-9582>
 Ilya M. D. Maclean  <https://orcid.org/0000-0001-8030-9136>
 Olivier Roupsard  <https://orcid.org/0000-0002-1319-142X>
 Juha M. Alatalo  <https://orcid.org/0000-0001-5084-850X>
 Stuart W. Smith  <https://orcid.org/0000-0001-9396-6610>
 Robert G. Björk  <https://orcid.org/0000-0001-7346-666X>
 Lena Muffler  <https://orcid.org/0000-0001-8227-7297>
 Felix Gottschall  <https://orcid.org/0000-0002-1247-8728>
 Josef Urban  <https://orcid.org/0000-0003-1730-947X>
 Martin Svátek  <https://orcid.org/0000-0003-2328-4627>
 Shyam S. Phartyal  <https://orcid.org/0000-0003-3266-6619>
 Sonja Wipf  <https://orcid.org/0000-0002-3492-1399>
 Nico Eisenhauer  <https://orcid.org/0000-0002-0371-6720>
 Mihai Pușcaș  <https://orcid.org/0000-0002-2632-640X>
 Pavel D. Turtureanu  <https://orcid.org/0000-0002-7422-3106>
 Andrej Varlagin  <https://orcid.org/0000-0002-2549-5236>
 Alistair S. Jump  <https://orcid.org/0000-0002-2167-6451>
 Krystal Randall  <https://orcid.org/0000-0003-2507-1000>
 Ellen Dorrepaal  <https://orcid.org/0000-0002-0523-2471>
 Josefina Walz  <https://orcid.org/0000-0002-0715-8738>
 Miroslav Svoboda  <https://orcid.org/0000-0003-4050-3422>
 Salvatore R. Curasi  <https://orcid.org/0000-0002-4534-3344>
 Austin Koontz  <https://orcid.org/0000-0002-6103-5894>

- William D. Pearce  <https://orcid.org/0000-0002-6241-3164>
- Bente Jessen Graae  <https://orcid.org/0000-0002-5568-4759>
- Toke T. Høye  <https://orcid.org/0000-0001-5387-3284>
- Juan Lorite  <https://orcid.org/0000-0003-4617-8069>
- Michele Carbognani  <https://orcid.org/0000-0001-7701-9859>
- Marcello Tomaselli  <https://orcid.org/0000-0003-4208-3433>
- T'ai G. W. Forte  <https://orcid.org/0000-0002-8685-5872>
- Stef Haesen  <https://orcid.org/0000-0002-4491-4213>
- Ben Somers  <https://orcid.org/0000-0002-7875-107X>
- Koenraad Van Meerbeek  <https://orcid.org/0000-0002-9260-3815>
- Mana Gharun  <https://orcid.org/0000-0003-0337-7367>
- Nina Buchmann  <https://orcid.org/0000-0003-0826-2980>
- Jiri Dolezal  <https://orcid.org/0000-0002-5829-4051>
- Andrew D. Thomas  <https://orcid.org/0000-0002-1360-1687>
- Dany Ghosn  <https://orcid.org/0000-0003-1898-9681>
- Miguel A. de Pablo  <https://orcid.org/0000-0002-4496-2741>
- Julia Kemppinen  <https://orcid.org/0000-0001-7521-7229>
- Pekka Niittynen  <https://orcid.org/0000-0002-7290-029X>
- Lisa Rew  <https://orcid.org/0000-0002-2818-3991>
- Christian Larson  <https://orcid.org/0000-0002-7567-4953>
- James D. M. Speed  <https://orcid.org/0000-0002-0633-5595>
- Jonas Ardö  <https://orcid.org/0000-0002-9318-0973>
- Maaikje Y. Bader  <https://orcid.org/0000-0003-4300-7598>
- Rafaella Canessa  <https://orcid.org/0000-0002-6979-9880>
- Angela Stanisci  <https://orcid.org/0000-0002-5302-0932>
- Juergen Kreyling  <https://orcid.org/0000-0001-8489-7289>
- Valeria Aschero  <https://orcid.org/0000-0003-3865-4133>
- Marek Čiliak  <https://orcid.org/0000-0002-6720-9365>
- František Máliš  <https://orcid.org/0000-0003-2760-6988>
- Pallietter De Smedt  <https://orcid.org/0000-0002-3073-6751>
- Sanne Govaert  <https://orcid.org/0000-0002-8939-1305>
- Camille Meeussen  <https://orcid.org/0000-0002-5869-4936>
- Pieter Vangansbeke  <https://orcid.org/0000-0002-6356-2858>
- Andrea Lamprecht  <https://orcid.org/0000-0002-8719-026X>
- Harald Pauli  <https://orcid.org/0000-0002-9842-9934>
- Klaus Steinbauer  <https://orcid.org/0000-0002-3730-9920>
- Manuela Winkler  <https://orcid.org/0000-0002-8655-9555>
- Masahito Ueyama  <https://orcid.org/0000-0002-4000-4888>
- Tudor-Mihai Ursu  <https://orcid.org/0000-0002-4898-6345>
- Sylvia Haider  <https://orcid.org/0000-0002-2966-0534>
- Ronja E. M. Wedegärtner  <https://orcid.org/0000-0003-4633-755X>
- Martin Wilmking  <https://orcid.org/0000-0003-4964-2402>
- Jan Altman  <https://orcid.org/0000-0003-4879-5773>
- Josef Brůna  <https://orcid.org/0000-0002-4839-4593>
- Martin Macek  <https://orcid.org/0000-0002-5609-5921>
- Matěj Man  <https://orcid.org/0000-0002-4557-8768>
- Jan Wild  <https://orcid.org/0000-0003-3007-4070>
- Pascal Vittoz  <https://orcid.org/0000-0003-4218-4517>
- Peter Barančok <https://orcid.org/0000-0003-1171-2524>
- Róbert Kanka <https://orcid.org/0000-0002-7071-7280>
- Andrej Palaj <https://orcid.org/0000-0001-7054-4183>
- Marijn Bauters <https://orcid.org/0000-0003-0978-6639>
- José-Luis Benito Alonso <https://orcid.org/0000-0003-1086-8834>
- Valter Di Cecco  <https://orcid.org/0000-0001-9862-1267>
- Zuzana Sitková  <https://orcid.org/0000-0001-6354-6105>
- Liesbeth van den Brink  <https://orcid.org/0000-0003-0313-8147>
- Robert Weigel  <https://orcid.org/0000-0001-9685-6783>
- C. Johan Dahlberg  <https://orcid.org/0000-0003-0271-3306>
- Sergiy Medinets  <https://orcid.org/0000-0001-5980-1054>
- Volodymyr Medinets  <https://orcid.org/0000-0001-7543-7504>
- Miguel Portillo-Estrada  <https://orcid.org/0000-0002-0348-7446>
- Lore T. Verryckt  <https://orcid.org/0000-0002-9452-5216>
- Ann Milbau  <https://orcid.org/0000-0003-3555-8883>
- Gergana N. Daskalova  <https://orcid.org/0000-0002-5674-5322>
- Haydn J. D. Thomas  <https://orcid.org/0000-0001-9099-6304>
- Isla H. Myers-Smith  <https://orcid.org/0000-0002-8417-6112>
- Jörg G. Stephan  <https://orcid.org/0000-0001-6195-7867>
- Patrice Descombes  <https://orcid.org/0000-0002-3760-9907>
- Florian Zellweger  <https://orcid.org/0000-0003-1265-9147>
- Esther R. Frei  <https://orcid.org/0000-0003-1910-7900>
- Bernard Heinesch  <https://orcid.org/0000-0001-7594-6341>
- Christopher Andrews  <https://orcid.org/0000-0003-2428-272X>
- Rebecca A. Senior  <https://orcid.org/0000-0002-8208-736X>
- Juan J. Jimenez  <https://orcid.org/0000-0003-2398-0796>
- Julia Boike  <https://orcid.org/0000-0002-5875-2112>
- Thomas Scholten  <https://orcid.org/0000-0002-4875-2602>
- Brett Scheffers  <https://orcid.org/0000-0003-2423-3821>
- Edmund W. Basham  <https://orcid.org/0000-0002-0167-7908>
- Jian Zhang  <https://orcid.org/0000-0003-0589-6267>
- Charly Géron  <https://orcid.org/0000-0001-7912-4708>
- Fatih Fazlioglu  <https://orcid.org/0000-0002-4723-3640>
- Onur Candan  <https://orcid.org/0000-0002-9254-4122>
- Jhonatan Sallo Bravo  <https://orcid.org/0000-0001-9007-4959>
- Filip Hrbacek  <https://orcid.org/0000-0001-5032-9216>
- Kamil Laska  <https://orcid.org/0000-0002-5199-9737>
- Edoardo Cremonese  <https://orcid.org/0000-0002-6708-8532>
- Peter Haase  <https://orcid.org/0000-0002-9340-0438>
- Fernando E. Moyano  <https://orcid.org/0000-0002-4090-5838>
- Christian Rossi  <https://orcid.org/0000-0001-9983-8898>
- Ivan Nijs  <https://orcid.org/0000-0003-3111-680X>

REFERENCES

- Aalto, J., Riihimäki, H., Meineri, E., Hylander, K., & Luoto, M. (2017). Revealing topoclimatic heterogeneity using meteorological station data. *International Journal of Climatology*, 37, 544–556. <https://doi.org/10.1002/joc.5020>
- Aalto, J., Scherrer, D., Lenoir, J., Guisan, A., & Luoto, M. (2018). Biogeophysical controls on soil-atmosphere thermal differences: Implications on warming Arctic ecosystems. *Environmental Research Letters*, 13, 074003. <https://doi.org/10.1088/1748-9326/aac83e>
- Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., & Hegewisch, K. C. (2018). TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015. *Scientific Data*, 5, 1958–2015. <https://doi.org/10.1038/sdata.2017.191>
- Ashcroft, M. B., Cavanagh, M., Eldridge, M. D. B., & Gollan, J. R. (2014). Testing the ability of topoclimatic grids of extreme temperatures to explain the distribution of the endangered brush-tailed rock-wallaby (*Petrogale penicillata*). *Journal of Biogeography*, 41, 1402–1413.

- Ashcroft, M. B., Chisholm, L. A., & French, K. O. (2008). The effect of exposure on landscape scale soil surface temperatures and species distribution models. *Landscape Ecology*, 23, 211–225. <https://doi.org/10.1007/s10980-007-9181-8>
- Ashcroft, M. B., Chisholm, L. A., & French, K. O. (2009). Climate change at the landscape scale: Predicting fine-grained spatial heterogeneity in warming and potential refugia for vegetation. *Global Change Biology*, 15, 656–667. <https://doi.org/10.1111/j.1365-2486.2008.01762.x>
- Ashcroft, M. B., & Gollan, J. R. (2012). Fine-resolution (25 m) topoclimatic grids of near-surface (5 cm) extreme temperatures and humidities across various habitats in a large (200 × 300 km) and diverse region. *International Journal of Climatology*, 32, 2134–2148.
- Ashcroft, M. B., & Gollan, J. R. (2013). Moisture, thermal inertia, and the spatial distributions of near-surface soil and air temperatures: Understanding factors that promote microrefugia. *Agricultural and Forest Meteorology*, 176, 77–89. <https://doi.org/10.1016/j.agrfor.2013.03.008>
- Bennie, J., Huntley, B., Wiltshire, A., Hill, M. O., & Baxter, R. (2008). Slope, aspect and climate: Spatially explicit and implicit models of topographic microclimate in chalk grassland. *Ecological Modelling*, 216, 47–59. <https://doi.org/10.1016/j.ecolmodel.2008.04.010>
- Bennie, J., Wilson, R. J., Maclean, I. M. D., & Suggitt, A. J. (2014). Seeing the woods for the trees – When is microclimate important in species distribution models? *Global Change Biology*, 20, 2699–2700. <https://doi.org/10.1111/gcb.12525>
- Bramer, I., Anderson, B., Bennie, J., Bladon, A., De Frenne, P., Hemming, D., ... Gillingham, P. K. (2018). Advances in monitoring and modelling climate at ecologically relevant scales. *Advances in Ecological Research*, 58, 101–161.
- Bruelheide, H., Dengler, J., Purschke, O., Lenoir, J., Jiménez-Alfaro, B., Hennekens, S. M., ... Jandt, U. (2018). Global trait-environment relationships of plant communities. *Nature Ecology & Evolution*, 2, 1906–1917. <https://doi.org/10.1038/s41559-018-0699-8>
- Cameron, E. K., Martins, I. S., Lavelle, P., Mathieu, J., Tedersoo, L., Gottschall, F., ... Eisenhauer, N. (2018). Global gaps in soil biodiversity data. *Nature Ecology & Evolution*, 2, 1042–1043. <https://doi.org/10.1038/s41559-018-0573-8>
- Carter, A., Kearney, M., Mitchell, N., Hartley, S., Porter, W., & Nelson, N. (2015). Modelling the soil microclimate: Does the spatial or temporal resolution of input parameters matter? *Frontiers of Biogeography*, 7, 138–154. <https://doi.org/10.21425/F5FBG27849>
- Copernicus Climate Change Service (C3s). (2019). *C3S ERA5-Land reanalysis* (Ed. Copernicus Climate Change Service).
- Coûteaux, M.-M., Bottner, P., & Berg, B. (1995). Litter decomposition, climate and litter quality. *Trends in Ecology & Evolution*, 10, 63–66.
- Daly, C. (2006). Guidelines for assessing the suitability of spatial climate data sets. *International Journal of Climatology*, 26, 707–721. <https://doi.org/10.1002/joc.1322>
- De Boeck, H. J., Van De Velde, H., De Groote, T., & Nijs, I. (2016). Ideas and perspectives: Heat stress: More than hot air. *Biogeosciences*, 13, 5821–5825. <https://doi.org/10.5194/bg-13-5821-2016>
- De Frenne, P., Rodriguez-Sanchez, F., Coomes, D. A., Baeten, L., Verstraeten, G., Vellend, M., ... Verheyen, K. (2013). Microclimate moderates plant responses to macroclimate warming. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 18561–18565. <https://doi.org/10.1073/pnas.1311190110>
- Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate climate surfaces for global land areas. *International Journal of Climatology*, 37, 4302–4315. <https://doi.org/10.1002/joc.5086>
- Geiger, R. (1950). *The climate near the ground*. Cambridge, MA: Harvard University Press.
- Gottschall, F., Davids, S., Newiger-Dous, T. E., Auge, H., Cesarz, S., & Eisenhauer, N. (2019). Tree species identity determines wood decomposition via microclimatic effects. *Ecology and Evolution*, 9, 12113–12127. <https://doi.org/10.1002/ece3.5665>
- Goulden, M. L., Wofsy, S. C., Harden, J. W., Trumbore, S. E., Crill, P. M., Gower, S. T., ... Munger, J. W. (1998). Sensitivity of boreal forest carbon balance to soil thaw. *Science*, 279, 214–217. <https://doi.org/10.1126/science.279.5348.214>
- Guerra, C. A., Heintz-Buschart, A., Sikorski, J., Chatzinotas, A., Guerrero-Ramírez, N., Cesarz, S., ... Delgado-Baquerizo, M. (2019). Blind spots in global soil biodiversity and ecosystem function research. *bioRxiv*, 774356. <https://doi.org/10.1101/774356>
- Hursh, A., Ballantyne, A., Cooper, L., Maneta, M., Kimball, J., & Watts, J. (2017). The sensitivity of soil respiration to soil temperature, moisture, and carbon supply at the global scale. *Global Change Biology*, 23, 2090–2103. <https://doi.org/10.1111/gcb.13489>
- Jarraud, M. (2008). *Guide to meteorological instruments and methods of observation* (WMO-No. 8). Geneva, Switzerland: World Meteorological Organisation.
- Jucker, T., Jackson, T., Zellweger, F., Swinfield, T., Gregory, N., Williamson, J., ... Coomes, D. (2020). A research agenda for microclimate ecology in human-modified tropical forests. *Frontiers in Forests and Global Change*, 2. <https://doi.org/10.3389/ffgc.2019.00092>
- Karger, D. N., Conrad, O., Böhrer, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., ... Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Scientific Data*, 4, 170122. <https://doi.org/10.1038/sdata.2017.122>
- Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., ... Wirth, C. (2019). TRY plant trait database-enhanced coverage and open access. *Global Change Biology*, 26, 119–188. <https://doi.org/10.1111/gcb.14904>
- Kearney, M. R. (2019). MicroclimOz – A microclimate data set for Australia, with example applications. *Austral Ecology*, 44, 534–544. <https://doi.org/10.1111/aec.12689>
- Kearney, M. R., Isaac, A. P., & Porter, W. P. (2014). microclim: Global estimates of hourly microclimate based on long-term monthly climate averages. *Scientific Data*, 1, 140006. <https://doi.org/10.1038/sdata.2014.6>
- Kearney, M. R., Shamakh, A., Tingley, R., Karoly, D. J., Hoffmann, A. A., Briggs, P. R., & Porter, W. P. (2014). Microclimate modelling at macro scales: A test of a general microclimate model integrated with gridded continental-scale soil and weather data. *Methods in Ecology and Evolution*, 5, 273–286. <https://doi.org/10.1111/2041-210X.12148>
- Kearney, M., Shine, R., & Porter, W. P. (2009). The potential for behavioural thermoregulation to buffer “cold-blooded” animals against climate warming. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 3835–3840. <https://doi.org/10.1073/pnas.0808913106>
- Kissling, W. D., Walls, R., Bowser, A., Jones, M. O., Kattge, J., Agosti, D., ... Guralnick, R. P. (2018). Towards global data products of essential biodiversity variables on species traits. *Nature Ecology & Evolution*, 2, 1531–1540. <https://doi.org/10.1038/s41559-018-0667-3>
- Körner, C., & Hiltbrunner, E. (2018). The 90 ways to describe plant temperature. *Perspectives in Plant Ecology, Evolution and Systematics*, 30, 16–21. <https://doi.org/10.1016/j.ppees.2017.04.004>
- Körner, C., & Paulsen, J. (2004). A world-wide study of high altitude treeline temperatures. *Journal of Biogeography*, 31, 713–732. <https://doi.org/10.1111/j.1365-2699.2003.01043.x>
- Lembrechts, J. J., Lenoir, J., Roth, N., Hattab, T., Milbau, A., Haider, S., ... Nijs, I. (2019). Comparing temperature data sources for use in species distribution models: From in-situ logging to remote sensing. *Global Ecology and Biogeography*, 28, 1578–1596. <https://doi.org/10.1111/geb.12974>
- Lembrechts, J., Nijs, I., & Lenoir, J. (2019). Incorporating microclimate into species distribution models. *Ecography*, 42, 1267–1279. <https://doi.org/10.1111/ecog.03947>
- Lenoir, J., Hattab, T., & Pierre, G. (2017). Climatic microrefugia under anthropogenic climate change: Implications for species redistribution. *Ecography*, 40, 253–266. <https://doi.org/10.1111/ecog.02788>

- Li, T.-T. (1926). *Soil temperature as influenced by forest cover*. New Haven, CT: Yale University - School of Forestry.
- Macek, M., Kopecký, M., & Wild, J. (2019). Maximum air temperature controlled by landscape topography affects plant species composition in temperate forests. *Landscape Ecology*, 34, 2541–2556. <https://doi.org/10.1007/s10980-019-00903-x>
- Maclean, I. M. (2019). Predicting future climate at high spatial and temporal resolution. *Global Change Biology*, 26, 1003–1011. <https://doi.org/10.1111/gcb.14876>
- Maclean, I. M. D., Hopkins, J. J., Bennie, J., Lawson, C. R., & Wilson, R. J. (2015). Microclimates buffer the responses of plant communities to climate change. *Global Ecology and Biogeography*, 24, 1340–1350. <https://doi.org/10.1111/geb.12359>
- Maclean, I. M. D., Suggitt, A. J., Wilson, R. J., Duffy, J. P., & Bennie, J. J. (2017). Fine-scale climate change: Modelling spatial variation in biologically meaningful rates of warming. *Global Change Biology*, 23, 256–268. <https://doi.org/10.1111/gcb.13343>
- Maestre, F. T., & Eisenhauer, N. (2019). Recommendations for establishing global collaborative networks in soil ecology. *Soil Organisms*, 91, 73.
- Medinets, S., Gasche, R., Kiese, R., Rennenberg, H., & Butterbach-Bahl, K. (2019). Seasonal dynamics and profiles of soil NO concentrations in a temperate forest. *Plant and Soil*, 445, 335–348. <https://doi.org/10.1007/s11104-019-04305-5>
- Meineri, E., & Hylander, K. (2017). Fine-grain, large-domain climate models based on climate station and comprehensive topographic information improve microrefugia detection. *Ecography*, 40, 1003–1013. <https://doi.org/10.1111/ecog.02494>
- Niittynen, P., & Luoto, M. (2017). The importance of snow in species distribution models of arctic vegetation. *Ecography*, 41, 1024–1037. <https://doi.org/10.1111/ecog.03348>
- Opedal, O. H., Armbruster, W. S., & Graae, B. J. (2015). Linking small-scale topography with microclimate, plant species diversity and intra-specific trait variation in an alpine landscape. *Plant Ecology & Diversity*, 8, 305–315. <https://doi.org/10.1080/17550874.2014.987330>
- Pincebourde, S., & Casas, J. (2019). Narrow safety margin in the phyllosphere during thermal extremes. *Proceedings of the National Academy of Sciences of the United States of America*, 116, 5588–5596. <https://doi.org/10.1073/pnas.1815828116>
- Pincebourde, S., Murdock, C. C., Vickers, M., & Sears, M. W. (2016). Fine-scale microclimatic variation can shape the responses of organisms to global change in both natural and urban environments. *Integrative and Comparative Biology*, 56, 45–61. <https://doi.org/10.1093/icb/icw016>
- Pleim, J. E., & Gilliam, R. (2009). An indirect data assimilation scheme for deep soil temperature in the Pleim-Xiu land surface model. *Journal of Applied Meteorology and Climatology*, 48, 1362–1376. <https://doi.org/10.1175/2009JAMC2053.1>
- Portillo-Estrada, M., Pihlatie, M., Korhonen, J. F. J., Levula, J., Frumau, A. K. F., Ibrom, A., ... Niinemets, U. (2016). Climatic controls on leaf litter decomposition across European forests and grasslands revealed by reciprocal litter transplantation experiments. *Biogeosciences*, 13, 1621–1633. <https://doi.org/10.5194/bg-13-1621-2016>
- Pradervand, J.-N., Dubuis, A., Pellissier, L., Guisan, A., & Randin, C. (2014). Very high resolution environmental predictors in species distribution models: Moving beyond topography? *Progress in Physical Geography*, 38, 79–96. <https://doi.org/10.1177/0309133313512667>
- Randin, C. F., Vuissoz, G., Liston, G. E., Vittoz, P., & Guisan, A. (2009). Introduction of snow and geomorphic disturbance variables into predictive models of alpine plant distribution in the Western Swiss Alps. *Arctic, Antarctic, and Alpine Research*, 41, 347–361. <https://doi.org/10.1657/1938-4246-41.3.347>
- Rosenberg, N. J., Kimball, B., Martin, P., & Cooper, C. (1990). From climate and CO₂ enrichment to evapotranspiration. In P. Wagoner (Ed.), *Climate change and US water resources* (pp. 151–175). New York, NY: John Wiley and Sons Inc.
- Schimel, D. S., Braswell, B., Mckeown, R., Ojima, D. S., Parton, W., & Pulliam, W. (1996). Climate and nitrogen controls on the geography and timescales of terrestrial biogeochemical cycling. *Global Biogeochemical Cycles*, 10, 677–692. <https://doi.org/10.1029/96GB01524>
- Slavich, E., Warton, D. I., Ashcroft, M. B., Gollan, J. R., & Ramp, D. (2014). Topoclimate versus macroclimate: How does climate mapping methodology affect species distribution models and climate change projections? *Diversity and Distributions*, 20, 952–963. <https://doi.org/10.1111/ddi.12216>
- Suggitt, A. J., Gillingham, P. K., Hill, J. K., Huntley, B., Kunin, W. E., Roy, D. B., & Thomas, C. D. (2011). Habitat microclimates drive fine-scale variation in extreme temperatures. *Oikos*, 120, 1–8. <https://doi.org/10.1111/j.1600-0706.2010.18270.x>
- Suggitt, A. J., Wilson, R. J., Isaac, N. J. B., Beale, C. M., Auffret, A. G., August, T., ... Maclean, I. M. D. (2018). Extinction risk from climate change is reduced by microclimatic buffering. *Nature Climate Change*, 8, 713. <https://doi.org/10.1038/s41558-018-0231-9>
- Vitasse, Y., Klein, G., Kirchner, J. W., & Rebetez, M. (2017). Intensity, frequency and spatial configuration of winter temperature inversions in the closed La Brevine valley, Switzerland. *Theoretical and Applied Climatology*, 130, 1073–1083. <https://doi.org/10.1007/s00704-016-1944-1>
- Wason, J. W., Bevilacqua, E., & Dovciak, M. (2017). Climates on the move: Implications of climate warming for species distributions in mountains of the northeastern United States. *Agricultural and Forest Meteorology*, 246, 272–280. <https://doi.org/10.1016/j.agrformet.2017.05.019>
- Western, A. W., Grayson, R. B., & Blöschl, G. (2002). Scaling of soil moisture: A hydrologic perspective. *Annual Review of Earth and Planetary Sciences*, 30, 149–180. <https://doi.org/10.1146/annurev.earth.30.091201.140434>
- Whiteman, C. D. (1982). Breakup of temperature inversions in deep mountain valleys: Part I. Observations. *Journal of Applied Meteorology*, 21, 270–289. [https://doi.org/10.1175/1520-0450\(1982\)021<0270:BOTIID>2.0.CO;2](https://doi.org/10.1175/1520-0450(1982)021<0270:BOTIID>2.0.CO;2)
- Whittaker, R. H. (1975). *Communities and ecosystems*. Communities and ecosystems (2nd ed). New York, NY: Macmillan.
- Zellweger, F., De Frenne, P., Lenoir, J., Rocchini, D., & Coomes, D. (2019). Advances in microclimate ecology arising from remote sensing. *Trends in Ecology & Evolution*, 34, 327–341. <https://doi.org/10.1016/j.tree.2018.12.012>
- Zhang, Y., Wang, S., Barr, A. G., & Black, T. (2008). Impact of snow cover on soil temperature and its simulation in a boreal aspen forest. *Cold Regions Science and Technology*, 52, 355–370. <https://doi.org/10.1016/j.coldregions.2007.07.001>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Lembrechts JJ, Aalto J, Ashcroft MB, et al. SoilTemp: A global database of near-surface temperature. *Glob Change Biol*. 2020;00:1–14. <https://doi.org/10.1111/gcb.15123>