

NONHEXAGONAL LATTICES FROM A TWO SPECIES INTERACTING SYSTEM*

SENPING LUO[†], XIAOFENG REN[‡], AND JUNCHENG WEI[†]

Abstract. A two species interacting system motivated by the density functional theory for triblock copolymers contains a long range interaction that affects the two species differently. In a two species periodic assembly of discs, the two species appear alternately on a lattice. A minimal two species periodic assembly is one with the least energy per area. There is a parameter b in $[0, 1]$ and the type of the lattice associated with a minimal assembly varies depending on b . There are several thresholds defined by a number $B = 0.1867\dots$. If $b \in [0, B)$, a minimal assembly is associated with a rectangular lattice whose ratio of the longer side to the shorter side is in $[\sqrt{3}, 1)$; if $b \in [B, 1 - B]$, a minimal assembly is associated with a square lattice; if $b \in (1 - B, 1]$, a minimal assembly is associated with a rhombic lattice with an acute angle in $[\frac{\pi}{3}, \frac{\pi}{2})$. Only when $b = 1$ is this rhombic lattice a hexagonal lattice. None of the other values of b yield a hexagonal lattice, a sharp contrast to the situation for one species interacting systems, where hexagonal lattices are ubiquitously observed.

Key words. two species interacting system, triblock copolymer, two species periodic assembly of discs, rectangular lattice, square lattice, rhombic lattice, hexagonal lattice, duality property

AMS subject classifications. 82B20, 82D60, 92C15, 11F20

DOI. 10.1137/19M1245980

1. Introduction. From honeycomb to chicken wire fence, from graphene to carbon nanotube, the hexagonal pattern is ubiquitous in nature. The honeycomb conjecture states that the hexagonal tiling is the best way to divide a surface into regions of equal area with the least total perimeter [12]. The Fekete problem minimizes an interaction energy of points on a sphere and obtains a hexagonal arrangement of minimizing points (with some defects due to a topological reason) [4].

Against this conventional wisdom, we present a problem where the hexagonal pattern is *generally not* the most favored structure. Our study is motivated by Nakazawa and Ohta's theory for triblock copolymer morphology [14, 20]. In an ABC triblock copolymer a molecule is a subchain of type A monomers connected to a subchain of type B monomers which in turn is connected to a subchain of type C monomers. Because of the repulsion between the unlike monomers, the different type subchains tend to segregate. However since subchains are chemically bonded in molecules, segregation cannot lead to a macroscopic phase separation; only microdomains rich in individual type monomers emerge, forming morphological phases. Bonding of distinct monomer subchains provides an inhibition mechanism in block copolymers.

The mathematical study of the triblock copolymer problem is still in the early stage. There are existence theorems about stationary assemblies of core shells [17], double bubbles [22], and discs [18], with the last work being the most relevant to this paper. Here we treat two of the three monomer types of a triblock copolymer as

*Received by the editors February 20, 2019; accepted for publication (in revised form) November 25, 2019; published electronically April 21, 2020.

<https://doi.org/10.1137/19M1245980>

Funding: The work of the first and third authors was partially supported by the NSERC grant NSERC-RGPIN-2018-03773. The work of the second author was partially supported by the National Science Foundation grant DMS-1714371.

[†]Department of Mathematics, University of British Columbia, Vancouver, BC, Canada V6T 1Z2 (spluo@math.ubc.ca, jwei@math.ubc.ca).

[‡]Corresponding author. Department of Mathematics, The George Washington University, Washington, DC 20052 (ren@gwu.edu).

species and view the third type as the surrounding environment, dependent on the two species. This way a triblock copolymer is a two species interacting system.

The definition of our two species interacting system starts with a lattice Λ on the complex plane generated by two nonzero complex numbers α_1 and α_2 with $\text{Im}(\alpha_2/\alpha_1) > 0$:

$$(1.1) \quad \Lambda = \{j_1\alpha_1 + j_2\alpha_2 : j_1, j_2 \in \mathbb{Z}\}.$$

Define by D_α the period parallelogram

$$(1.2) \quad D_\alpha = \{t_1\alpha_1 + t_2\alpha_2 : t_1, t_2 \in (0, 1)\}$$

associated with a basis $\alpha = (\alpha_1, \alpha_2)$ of the lattice Λ . The lattice Λ defines an equivalence relation on $\mathbb{C}$ where two complex numbers are equivalent if their difference is in Λ . The resulting space of equivalent classes is denoted $\mathbb{C}/\Lambda$. It can be represented by D_α , where the opposite edges of D_α are identified.

There are two sets of parameters in our model. The first consists of two numbers ω_1 and ω_2 satisfying

$$(1.3) \quad 0 < \omega_1, \omega_2 < 1, \text{ and } \omega_1 + \omega_2 < 1.$$

The second is a two by two symmetric matrix γ ,

$$(1.4) \quad \gamma = \begin{bmatrix} \gamma_{11} & \gamma_{12} \\ \gamma_{21} & \gamma_{22} \end{bmatrix}, \quad \gamma_{12} = \gamma_{21}.$$

Furthermore, in this paper we assume that

$$(1.5) \quad \gamma_{11} > 0, \gamma_{22} > 0, \gamma_{12} \geq 0, \gamma_{11}\gamma_{22} - \gamma_{12}^2 \geq 0.$$

Our model is a variational problem defined on pairs of Λ -periodic sets with prescribed average size. More specifically, a pair (Ω_1, Ω_2) of two subsets of $\mathbb{C}$ is admissible if the following conditions hold. Both Ω_1 and Ω_2 are Λ -periodic, i.e.,

$$(1.6) \quad \Omega_j + \lambda = \Omega_j \text{ for all } \lambda \in \Lambda, j = 1, 2;$$

Ω_1 and Ω_2 are disjoint in the sense that

$$(1.7) \quad |\Omega_1 \cap \Omega_2| = 0;$$

the average size of Ω_1 and Ω_2 are fixed at $\omega_1, \omega_2 \in (0, 1)$, respectively, i.e.,

$$(1.8) \quad \frac{|\Omega_j \cap D_\alpha|}{|D_\alpha|} = \omega_j, \quad j = 1, 2.$$

In (1.7) and (1.8), $|\cdot|$ denotes the two-dimensional Lebesgue measure. Although it can be given in terms of α_1 and α_2 , $|D_\alpha|$ actually depends on the lattice Λ , not the particular basis α , and therefore we alternatively write it as $|\Lambda|$,

$$(1.9) \quad |\Lambda| = |D_\alpha| = \text{Im}(\overline{\alpha_1}\alpha_2).$$

Given an admissible pair (Ω_1, Ω_2) , let $\Omega_3 = \mathbb{C} \setminus (\Omega_1 \cup \Omega_2)$. Again Λ imposes an equivalent relation on Ω_j and the resulting space of equivalence classes is denoted

Ω_j/Λ , $j = 1, 2, 3$. Define a functional $\mathcal{J}_\Lambda$ to be the free energy of (Ω_1, Ω_2) on a period parallelogram given by

$$(1.10) \quad \mathcal{J}_\Lambda(\Omega_1, \Omega_2) = \frac{1}{2} \sum_{j=1}^3 \mathcal{P}_{\mathbb{C}/\Lambda}(\Omega_j/\Lambda) + \sum_{j,k=1}^2 \frac{\gamma_{jk}}{2} \int_{D_\alpha} \nabla I_\Lambda(\Omega_j)(\zeta) \cdot \nabla I_\Lambda(\Omega_k)(\zeta) d\zeta.$$

In (1.10), $\mathcal{P}_{\mathbb{C}/\Lambda}(\Omega_j/\Lambda)$, $j = 1, 2$, is the perimeter of Ω_j/Λ in $\mathbb{C}/\Lambda$. One can take a representation D_α of $\mathbb{C}/\Lambda$, with its opposite sides identified, and treat $\Omega_j \cap D_\alpha$, also with points on opposite sides identified, as a subset of D_α . Then $\mathcal{P}_{\mathbb{C}/\Lambda}(\Omega_j/\Lambda)$ is the perimeter of $\Omega_j \cap D_\alpha$. If $\Omega_j \cap D_\alpha$ is bounded by C^1 curves, then the perimeter is just the total length of the curves. More generally, for a merely measurable Λ -periodic Ω_j ,

$$(1.11) \quad \mathcal{P}_{\mathbb{C}/\Lambda}(\Omega_j/\Lambda) = \sup_g \left\{ \int_{\Omega_j \cap D_\alpha} \operatorname{div} g(x) dx : g \in C^1(\mathbb{C}/\Lambda, \mathbb{R}^2), |g(x)| \leq 1 \ \forall x \in \mathbb{C} \right\}.$$

Here $g \in C^1(\mathbb{C}/\Lambda, \mathbb{R}^2)$ means that g is a continuously differential, Λ -periodic vector field on $\mathbb{C}$; $|g(x)|$ is the geometric norm of the vector $g(x) \in \mathbb{R}^2$.

In $\sum_{j=1}^3 \mathcal{P}_{\mathbb{C}/\Lambda}(\Omega_j/\Lambda)$ each boundary curve separating a Ω_j/Λ from a Ω_k/Λ , $j, k = 1, 2, 3$, $j \neq k$, is counted exactly twice. The constant $\frac{1}{2}$ in the front takes care of the double counting.

The function $I_\Lambda(\Omega_j)$ is the Λ -periodic solution of Poisson's equation

$$(1.12) \quad -\Delta I_\Lambda(\Omega_j)(\zeta) = \chi_{\Omega_j}(\zeta) - \omega_j \text{ in } \mathbb{C}, \quad \int_{D_\alpha} I_\Lambda(\Omega_j)(\zeta) d\zeta = 0,$$

where χ_{Ω_j} is the characteristic function of Ω_j . Despite the appearance, the functional $\mathcal{J}_\Lambda$ depends on the lattice Λ instead of the particular basis α .

A stationary point (Ω_1, Ω_2) of $\mathcal{J}_\Lambda$ is a solution to the following equations of a free boundary problem:

$$(1.13) \quad \kappa_{13} + \gamma_{11} I_\Lambda(\Omega_1) + \gamma_{12} I_\Lambda(\Omega_2) = \mu_1 \text{ on } \partial\Omega_1 \cap \partial\Omega_3,$$

$$(1.14) \quad \kappa_{23} + \gamma_{12} I_\Lambda(\Omega_1) + \gamma_{22} I_\Lambda(\Omega_2) = \mu_2 \text{ on } \partial\Omega_2 \cap \partial\Omega_3,$$

$$(1.15) \quad \kappa_{12} + (\gamma_{11} - \gamma_{12}) I_\Lambda(\Omega_1) + (\gamma_{12} - \gamma_{22}) I_\Lambda(\Omega_2) = \mu_1 - \mu_2 \text{ on } \partial\Omega_1 \cap \partial\Omega_2,$$

$$(1.16) \quad T_{13} + T_{23} + T_{12} = \vec{0} \text{ at } \partial\Omega_1 \cap \partial\Omega_2 \cap \partial\Omega_3.$$

In (1.13)–(1.15) κ_{13} , κ_{23} , and κ_{12} are the curvatures of the curves $\partial\Omega_1 \cap \partial\Omega_3$, $\partial\Omega_2 \cap \partial\Omega_3$, and $\partial\Omega_1 \cap \partial\Omega_2$, respectively. The unknown constants μ_1 and μ_2 are Lagrange multipliers associated with the constraints (1.8) for Ω_1 and Ω_2 , respectively. The three interfaces, $\partial\Omega_1 \cap \partial\Omega_3$, $\partial\Omega_2 \cap \partial\Omega_3$ and $\partial\Omega_1 \cap \partial\Omega_2$, may meet at a common point in D , which is termed a triple junction point. In (1.16), T_{13} , T_{23} , and T_{12} are, respectively, the unit tangent vectors of these curves at triple junction points. This equation simply says that at a triple junction point three curves meet at 120 degree angles.

In this paper, we only consider a special type of (Ω_1, Ω_2) , termed two species periodic assemblies of discs, denoted by $(\Omega_{\alpha,1}, \Omega_{\alpha,2})$, with

$$(1.17) \quad \Omega_{\alpha,1} = \bigcup_{\lambda \in \Lambda} \left\{ B(\xi, r_1) \cup B(\xi', r_1) : \xi = \frac{3}{4}\alpha_1 + \frac{1}{4}\alpha_2 + \lambda, \xi' = \frac{1}{4}\alpha_1 + \frac{3}{4}\alpha_2 + \lambda \right\},$$

$$(1.18) \quad \Omega_{\alpha,2} = \bigcup_{\lambda \in \Lambda} \left\{ B(\xi, r_2) \cup B(\xi', r_2) : \xi = \frac{1}{4}\alpha_1 + \frac{1}{4}\alpha_2 + \lambda, \xi' = \frac{3}{4}\alpha_1 + \frac{3}{4}\alpha_2 + \lambda \right\}.$$




FIG. 1. A two species periodic assembly of discs given by (1.17) and (1.18), and a shift of the assembly with disc centers at the lattice points and the half-lattice points.

In (1.17) and (1.18), $B(\xi, r_j)$, or $B(\xi', r_j)$, is the closed disc centered at ξ (or ξ') of radius r_j ; the r_j 's are given by

$$(1.19) \quad \omega_j = \frac{2\pi r_j^2}{|D_\alpha|}, \quad j = 1, 2.$$

Be aware that $(\Omega_{\alpha,1}, \Omega_{\alpha,2})$ defined this way depends on the basis α , not the lattice Λ generated by α . One may have two different bases that generate the same lattice, but they define two distinct assemblies.

Shifting $(\Omega_{\alpha,1}, \Omega_{\alpha,2})$ does not change its energy, so our choice for the centers of the discs in (1.17) and (1.18) is not the only one. Another aesthetically pleasing placement is to put the disc centers on the lattice points and half lattice points; see Figure 1. Nevertheless we prefer not to have discs on the boundary of the period parallelogram D_α .

A two species periodic assembly $(\Omega_{\alpha,1}, \Omega_{\alpha,2})$ is not a stationary point of the energy functional $\mathcal{J}_\Lambda$. However Ren and Wang have shown the existence of stationary points that are unions of perturbed discs in a bounded domain with the Neumann boundary condition [18]. Numerical evidence strongly suggests the existence of stationary points similar to two species assemblies [25].

In this paper we determine, in terms of α , which $(\Omega_{\alpha,1}, \Omega_{\alpha,2})$ is the most energetically favored. For this purpose, it is more appropriate to consider the energy per area instead of the energy on a period parallelogram. Namely, consider

$$(1.20) \quad \tilde{\mathcal{J}}_\Lambda(\Omega_1, \Omega_2) = \frac{1}{|\Lambda|} \mathcal{J}_\Lambda(\Omega_1, \Omega_2),$$

take (Ω_1, Ω_2) to be a two species periodic assembly, and minimize energy per area among all such assemblies with respect to α , i.e.,

$$(1.21) \quad \min_{\alpha} \left\{ \tilde{\mathcal{J}}_\Lambda(\Omega_{\alpha,1}, \Omega_{\alpha,2}) : \alpha = (\alpha_1, \alpha_2), \alpha_1, \alpha_2 \in \mathbb{C} \setminus \{0\}, \operatorname{Im} \frac{\alpha_2}{\alpha_1} > 0, \Lambda \text{ is generated by } \alpha \right\}.$$

Several lattices will appear as the most favored structures. They are illustrated in Figure 2. A rectangular lattice has a basis α whose parallelogram D_α is a rectangle.




FIG. 2. Two species periodic assemblies of discs with their associated lattices. First row from left to right: a rectangular lattice and a square lattice. Second row from left to right: a rhombic lattice and a hexagonal lattice.

A square lattice has a square as a parallelogram. A rhombic lattice has a rhombus parallelogram, i.e., a parallelogram whose four sides have the same length. Finally a hexagonal lattice has a parallelogram with four equal length sides and an angle of $\frac{\pi}{3}$ between two sides. If we let

$$(1.22) \quad \tau = \frac{\alpha_2}{\alpha_1},$$

then in terms of τ , Λ is rectangular if $\text{Re } \tau = 0$, Λ is square if $\tau = i$, Λ is rhombic if $|\tau| = 1$, and Λ is hexagonal if $\tau = e^{\frac{\pi i}{3}}$. Note that these classes of lattices are not mutually exclusive. A hexagonal lattice is a rhombic lattice; a square lattice is both a rectangular lattice and a rhombic lattice.

The reason that a rhombic lattice with a $\frac{\pi}{3}$ angle is termed a hexagonal lattice comes from its Voronoi cells. At each lattice point, the Voronoi cell of this lattice point consists of points in $\mathbb{C}$ that are closer to this lattice point than any other lattice points. For the rhombic lattice with a $\frac{\pi}{3}$ angle, the Voronoi cell at each lattice point is a regular hexagon. With Voronoi cells at all lattice points, the hexagonal lattice gives rise to a honeycomb pattern.

A two species periodic assembly of discs that minimizes the energy per area is called a minimal assembly and its associated lattice is called a minimal lattice. The

main result of this paper asserts that a minimal lattice must belong to one of three lattice classes:

1. it is a rectangular lattice with a rectangle period parallelogram whose ratio of the longer side to the shorter side lies in $(1, \sqrt{3}]$;
2. it is a square lattice;
3. it is a rhombic lattice with a rhombus period parallelogram whose acute angle is in $[\frac{\pi}{3}, \frac{\pi}{2})$.

Note that the rhombic lattice in the third scenario becomes a hexagonal lattice if the angle is $\frac{\pi}{3}$.

The parameter that determines which class of lattices a minimal lattice belongs to is

$$(1.23) \quad b = \frac{2\gamma_{12}\omega_1\omega_2}{\gamma_{11}\omega_1^2 + \gamma_{22}\omega_2^2},$$

given in terms of ω_j and γ_{jk} . Conditions (1.3), (1.4), and (1.5) on ω_j and γ_{jk} imply that

$$(1.24) \quad b \in [0, 1].$$

To ensure the disjoint condition (1.7) for potential minimal assemblies we assume that ω_1 and ω_2 are sufficiently small. Namely, let $\omega_0 > 0$ be small enough so that if

$$(1.25) \quad \omega_j < \omega_0, \quad j = 1, 2,$$

and $(\Omega_{\alpha,1}, \Omega_{\alpha,2})$ is a two species periodic assembly of discs whose basis (α_1, α_2) satisfies

$$(1.26) \quad \operatorname{Re} \tau = 0 \text{ and } |\tau| \in [1, \sqrt{3}]$$

or

$$(1.27) \quad |\tau| = 1 \text{ and } \arg \tau \in \left[\frac{\pi}{3}, \frac{\pi}{2}\right],$$

then $(\Omega_{\alpha,1}, \Omega_{\alpha,2})$ is disjoint in the sense of (1.7). The line segment (1.26) and the arc (1.27) are illustrated in the first plot of Figure 3. Now we state our theorem.

THEOREM 1.1. *Let the parameters ω_j , $j = 1, 2$, and γ_{jk} , $j, k = 1, 2$, satisfy the conditions (1.3), (1.4), (1.5), and (1.25). The minimization problem (1.21) always admits a minimum. Let $\alpha_* = (\alpha_{*,1}, \alpha_{*,2})$ be a minimum of (1.21), Λ_* be the lattice determined by α_* . Then there exists $B = 0.1867\dots$ such that the following statements hold.*

1. *If $b = 0$, then Λ_* is a rectangular lattice. It has a rectangle period parallelogram whose ratio of the longer side to the shorter side is $\sqrt{3}$.*
2. *If $b \in (0, B)$, then Λ_* is a rectangular lattice. It has a rectangle period parallelogram whose ratio of the longer side to the shorter side is in $(1, \sqrt{3})$. As b increases from 0 to B , this ratio decreases from $\sqrt{3}$ to 1.*
3. *If $b \in [B, 1 - B]$, then Λ_* is a square lattice.*
4. *If $b \in (1 - B, 1)$, then Λ_* is a nonsquare, nonhexagonal rhombic lattice. It has a rhombus period parallelogram with an acute angle in $(\frac{\pi}{3}, \frac{\pi}{2})$. As b increases from $1 - B$ to 1, this angle decreases from $\frac{\pi}{2}$ to $\frac{\pi}{3}$.*
5. *If $b = 1$, then Λ_* is a hexagonal lattice.*




FIG. 3. The left plot shows the set W . In $\overline{W}_{\mathbb{H}}$, f_b attain the maximum at a point either on the thick line segment or on the thick arc. In the right plot, with $b \in (0, B)$, f_b increases in the directions of the arrows. The dot on the imaginary axis is $q_b i$.

The threshold B is defined precisely in (4.40) by two infinite series, from which one finds its numerical value.

Only in the case $b = 1$, $\tilde{\mathcal{J}}_{\Lambda}(\Omega_{\alpha,1}, \Omega_{\alpha,2})$ is minimized by a hexagonal lattice. In all other cases minimal lattices are not hexagonal. As a matter of fact, our assumption on γ in this paper is a bit different from the conditions for γ in a triblock copolymer. In a triblock copolymer, instead of (1.5), γ needs to be positive definite. In [18], where Ren and Wang found assemblies of perturbed discs as stationary points, γ_{12} is positive. In terms of b , γ being positive definite and $\gamma_{12} > 0$ mean that $b \in (0, 1)$.

In this paper we include both the $b = 0$ case and the $b = 1$ case for good reasons. The case $b = 1$ corresponds to $\gamma_{11}\gamma_{22} - \gamma_{12}^2 = 0$, i.e., γ has a nontrivial kernel spanned by $(-\omega_1, \omega_2)$. This case is actually very special. It is equivalent to a problem studied by Chen and Oshita in [6], a simpler one species analogy of the two species problem studied here. The motivation of that problem comes from the study of diblock copolymers where a molecule is a subchain of type A monomers connected to a subchain of type B monomers. With one type treated as a species and the other as the surrounding environment, a diblock copolymer is a one species interacting system.

The recent years have seen active work on the diblock copolymer problem; see [19, 21, 1, 7, 13, 11] and the references therein. Based on a density functional theory of Ohta and Kawasaki [15], the free energy of a diblock copolymer system on a Λ -periodic domain is

$$(1.28) \quad \mathcal{E}_{\Lambda}(\Omega) = \mathcal{P}_{\mathbb{C}/\Lambda}(\Omega/\Lambda) + \frac{\gamma_d}{2} \int_{D_{\alpha}} |\nabla I_{\Lambda}(\Omega)(\zeta)|^2 d\zeta.$$

Here, analogously to the two species problem, Ω is a Λ -periodic subset of $\mathbb{C}$ under the average area constraint $\frac{|\Omega \cap D_{\alpha}|}{|D_{\alpha}|} = \omega$, where $\omega \in (0, 1)$ is one of the two given parameters. The other parameter is the number $\gamma_d > 0$. Now take Ω to be Ω_{Λ}^d , the union of discs centered at $\frac{\alpha_1 + \alpha_2}{2} + \lambda$, $\lambda \in \Lambda$, of radius $\sqrt{\frac{\omega|D_{\alpha}|}{\pi}}$, and minimize the energy per area with respect to Λ :

$$(1.29) \quad \min_{\Lambda} \frac{1}{|\Lambda|} \mathcal{E}_{\Lambda}(\Omega_{\Lambda}^d).$$

This time, unlike in the two species problem, Ω_Λ^d depends on the lattice Λ , not the basis (α_1, α_2) . Chen and Oshita showed that (1.29) is minimized by a hexagonal lattice.

In [23] Sandier and Serfaty studied the Ginzburg–Landau problem with magnetic field and arrived at a reduced energy. Minimization of this energy turns out to be the same as the minimization problem (1.29).

There is also a connection between problem (1.29) and the crystallization problem pointed out by Petrache and Serfaty in [16]. It is related to the Cohn–Kumar conjecture in two dimensions which states that the hexagonal lattice is universally minimizing with respect to all interaction potentials that are completely monotone functions of square distance [8, 9]. Regarding the crystallization problem, Theil showed that the hexagonal lattice is minimizing for a class of Lennard–Jones type potentials [24]. On the other hand, Bétermin, De Luca, and Petrache found a potential (not allowed in the Cohn–Kumar conjecture) whose crystallization is achieved by a square lattice [5].

In our two species problem (1.21), the condition $b = 1$ actually makes the two species indistinguishable as far as interaction is concerned. It means that the two species function as one species, hence the equivalence to the one species problem (1.29). The case $b = 0$ is *dual* to the $b = 1$ case, a point explained below. It is therefore natural to include both cases.

Our work starts with Lemma 2.1, which states that for us to solve (1.21) it suffices to minimize the energy among two species periodic assemblies of unit period parallelogram area. Then in Lemma 2.4 it is shown that the latter problem is equivalent to maximizing a function,

$$(1.30) \quad f_b(z) = b \log |\operatorname{Im}(z)\eta(z)| + (1-b) \log \left| \operatorname{Im}\left(\frac{z+1}{2}\right)\eta\left(\frac{z+1}{2}\right) \right|$$

with respect to z in the set $\{z \in \mathbb{C} : \operatorname{Im} z > 0, |z| \geq 1, 0 \leq \operatorname{Re} z \leq 1\}$. Here

$$(1.31) \quad \eta(z) = e^{\frac{\pi}{3}zi} \prod_{n=1}^{\infty} (1 - e^{2\pi nzi})^4$$

is the fourth power of the Dedekind eta function.

If $b = 1$, then $f_b = f_1$ and we are looking at the problem studied by Chen and Oshita [6], and Sandier and Serfaty [23]. In this case, f_1 is maximized in a smaller set, $\{z \in \mathbb{C} : |z| \geq 1, 0 \leq \operatorname{Re} z \leq 1/2\}$. Using a maximum principle argument, Chen and Oshita showed that f_1 is maximized at $z = e^{\frac{\pi i}{3}}$, which corresponds to the hexagonal lattice. Sandier and Serfaty used a relation between the Dedekind eta function and the Epstein zeta function, and a property of the Jacobi theta function to arrive at the same conclusion.

Neither method seems to be applicable to the two species system with $b \neq 1$. Instead we rely on a duality principle, Lemma 3.5, which shows that maximizing f_b is equivalent to maximizing f_{1-b} . This allows us to only consider $b \in [0, 1/2]$, and there we are able to show that $f_b(z)$ attains the maximum on the imaginary axis above i , i.e., $\operatorname{Re} z = 0$ and $\operatorname{Im} z \geq 1$.

So we turn to maximize $f_b(yi)$ with respect to $y \geq 1$. The most technical part of this work, Lemma 4.4, shows that when $b = 0$, $f_0(yi)$ is maximized at $y = \sqrt{3}$; when $b \in (0, B)$, $f_b(yi)$ is maximized at some $y = q_b \in (1, \sqrt{3})$; when $b \in [B, 1]$, $f_b(yi)$ is maximized at $y = 1$. The theorem then follows readily. The key step in the proof of Lemma 4.4 is to establish a monotonicity property for the ratio of the derivatives of

$f_0(yi)$ and $f_1(yi)$ with respect to $y \in (1, \sqrt{3})$. This piece of argument is placed in the appendix so a reader who is more interested in the overall strategy of this work may skip it at the first reading.

2. Derivation of f_b . The size and the shape of a two species periodic assembly play different roles in its energy. To separate the two factors write the basis of a two species periodic assembly as $t\alpha = (t\alpha_1, t\alpha_2)$, where $t \in (0, \infty)$ and the parallelogram generated by $\alpha = (\alpha_1, \alpha_2)$ has the unit area, i.e., $|D_\alpha| = 1$. This way the assembly is now denoted by $(\Omega_{t\alpha,1}, \Omega_{t\alpha,2})$ with t measuring the size of the assembly (note $|D_{t\alpha}| = t^2$), and D_α describing the shape of the assembly. The lattice generated by α is denoted Λ and the lattice generated by $t\alpha$ is $t\Lambda$.

LEMMA 2.1. Fix $\alpha_1, \alpha_2 \in \mathbb{C} \setminus \{0\}$, $\text{Im}(\alpha_2/\alpha_1) > 0$, and $|D_\alpha| = 1$. Among all two species periodic assemblies $\Omega_{t\alpha}$, $t \in (0, \infty)$, the energy per area is minimized by the one with $t = t_\alpha$, where

$$(2.1) \quad t_\alpha = \left(\frac{2\sqrt{2\pi\omega_1} + 2\sqrt{2\pi\omega_2}}{\sum_{j,k=1}^2 \gamma_{jk} \int_{D_\alpha} \nabla I_\Lambda(\Omega_{\alpha,j})(\zeta) \cdot \nabla I_\Lambda(\Omega_{\alpha,k})(\zeta) d\zeta} \right)^{1/3}.$$

The energy per area of this assembly is

$$(2.2) \quad \begin{aligned} & \tilde{\mathcal{J}}_{t_\alpha\Lambda}(\Omega_{t_\alpha\alpha,1}, \Omega_{t_\alpha\alpha,2}) \\ &= 3 \left(\sqrt{2\pi\omega_1} + \sqrt{2\pi\omega_2} \right)^{2/3} \left(\sum_{j,k=1}^2 \frac{\gamma_{jk}}{2} \int_{D_\alpha} \nabla I_\Lambda(\Omega_{\alpha,j})(\zeta) \cdot \nabla I_\Lambda(\Omega_{\alpha,k})(\zeta) d\zeta \right)^{1/3}. \end{aligned}$$

Consequently minimization of (1.21) is reduced to minimizing

$$(2.3) \quad \mathcal{F}(\alpha) = \sum_{j,k=1}^2 \frac{\gamma_{jk}}{2} \int_{D_\alpha} \nabla I_\Lambda(\Omega_{\alpha,j})(\zeta) \cdot \nabla I_\Lambda(\Omega_{\alpha,k})(\zeta) d\zeta, \quad |D_\alpha| = 1,$$

with respect to α of unit area.

Proof. Between the two lattices, the functions $I_{t\Lambda}(\Omega_{t\alpha,j})$ and $I_\Lambda(\Omega_{\alpha,j})$ are related by

$$(2.4) \quad I_{t\Lambda}(\Omega_{t\alpha,j})(\chi) = t^2 I_\Lambda(\Omega_{\alpha,j})(\zeta), \quad t\zeta = \chi, \quad \zeta, \chi \in \mathbb{C},$$

because of (1.12). Then

$$\begin{aligned} & \mathcal{J}_{t\Lambda}(\Omega_{t\alpha,1}, \Omega_{t\alpha,2}) \\ &= t \left(2\sqrt{2\pi\omega_1} + 2\sqrt{2\pi\omega_2} \right) + \sum_{j,k=1}^2 \frac{\gamma_{jk}}{2} \int_{D_{t\alpha}} \nabla I_{t\Lambda}(\Omega_{t\alpha,j})(\chi) \cdot \nabla I_{t\Lambda}(\Omega_{t\alpha,k})(\chi) d\chi \\ &= t \left(2\sqrt{2\pi\omega_1} + 2\sqrt{2\pi\omega_2} \right) + t^4 \sum_{j,k=1}^2 \frac{\gamma_{jk}}{2} \int_{D_\alpha} \nabla I_\Lambda(\Omega_{\alpha,j})(\zeta) \cdot \nabla I_\Lambda(\Omega_{\alpha,k})(\zeta) d\zeta. \end{aligned}$$

The energy per area is

$$\begin{aligned} & \tilde{\mathcal{J}}_{t\Lambda}(\Omega_{t\alpha,1}, \Omega_{t\alpha,2}) \\ &= \left(\frac{1}{t}\right)^2 \mathcal{J}_{t\Lambda}(\Omega_{t\alpha}) \\ &= \frac{1}{t} \left(2\sqrt{2\pi\omega_1} + 2\sqrt{2\pi\omega_2}\right) + t^2 \sum_{j,k=1}^2 \frac{\gamma_{jk}}{2} \int_{D_\alpha} \nabla I_\Lambda(\Omega_{\alpha,j})(\zeta) \cdot \nabla I_\Lambda(\Omega_{\alpha,k})(\zeta) d\zeta. \end{aligned}$$

With respect to t , the last quantity is minimized at $t = t_\alpha$ given in (2.1), and the minimum value is given in (2.2).

Later one needs to minimize the right side of (2.2) with respect to α , $|D_\alpha| = 1$. This is equivalent to minimizing $\mathcal{F}(\alpha)$ with respect to α , $|D_\alpha| = 1$. Once a minimum, say α_* , is found, then compute t_{α_*} from (2.1) and make the assembly $\Omega_{t_{\alpha_*}\alpha_*}$ with the basis $t_{\alpha_*}\alpha_*$. This assembly minimizes (1.21). $\square$

Now that the minimization problem (1.21) is reduced to minimizing $\mathcal{F}$, we proceed to simplify $\mathcal{F}(\alpha)$ to a more amenable form. To this end, one expresses the solution of (1.12) in terms of the Green function G_Λ of the $-\Delta$ operator as

$$(2.5) \quad I_\Lambda(\Omega_j)(\zeta) = \int_{\Omega_j \cap D_\alpha} G_\Lambda(\zeta - \chi) d\chi.$$

Here G_Λ is the Λ -periodic solution of

$$(2.6) \quad -\Delta G_\Lambda(\zeta) = \sum_{\lambda \in \Lambda} \delta_\lambda(\zeta) - \frac{1}{|\Lambda|}, \quad \int_{D_\alpha} G_\Lambda(\zeta) d\zeta = 0$$

in $\mathbb{C}$. In (2.6) δ_λ is the delta measure at λ , and D_α is a period parallelogram of Λ . It is known that

$$(2.7) \quad \begin{aligned} G_\Lambda(\zeta) &= \frac{|\zeta|^2}{4|\Lambda|} - \frac{1}{2\pi} \log \left| e \left(\frac{\zeta^2 \overline{\alpha_1}}{4i|\Lambda|\alpha_1} - \frac{\zeta}{2\alpha_1} + \frac{\alpha_2}{12\alpha_1} \right) \left(1 - e \left(\frac{\zeta}{\alpha_1} \right) \right) \right. \\ &\quad \cdot \left. \prod_{n=1}^{\infty} \left(\left(1 - e \left(n\tau + \frac{\zeta}{\alpha_1} \right) \right) \left(1 - e \left(n\tau - \frac{\zeta}{\alpha_1} \right) \right) \right) \right|, \end{aligned}$$

where

$$(2.8) \quad e(z) = e^{2\pi iz}$$

and

$$(2.9) \quad \tau = \frac{\alpha_2}{\alpha_1}.$$

A simple proof of this fact can be found in [6]. Sometimes one singles out the singularity of G_Λ at 0 and decomposes G_Λ into

$$(2.10) \quad G_\Lambda(\zeta) = -\frac{1}{2\pi} \log \frac{2\pi|\zeta|}{\sqrt{|\Lambda|}} + \frac{|\zeta|^2}{4|\Lambda|} + H_\Lambda(\zeta),$$

where

$$(2.11) \quad \begin{aligned} H_\Lambda(\zeta) &= -\frac{1}{2\pi} \log \left| e \left(\frac{\zeta^2 \overline{\alpha_1}}{4i|\Lambda|\alpha_1} - \frac{\zeta}{2\alpha_1} + \frac{\alpha_2}{12\alpha_1} \right) \frac{\sqrt{|\Lambda|}}{2\pi\zeta} \left(1 - e \left(\frac{\zeta}{\alpha_1} \right) \right) \right. \\ &\quad \cdot \left. \prod_{n=1}^{\infty} \left(\left(1 - e \left(n\tau + \frac{\zeta}{\alpha_1} \right) \right) \left(1 - e \left(n\tau - \frac{\zeta}{\alpha_1} \right) \right) \right) \right| \end{aligned}$$

is a harmonic function on $(\mathbb{C} \setminus \Lambda) \cup \{0\}$.

The integral term in (1.10) can be written in several ways:

$$(2.12) \quad \begin{aligned} \int_{D_\Lambda} \nabla I_\Lambda(\Omega_j)(\zeta) \cdot \nabla I_\Lambda(\Omega_k)(\zeta) d\zeta &= \int_{\Omega_k} I_\Lambda(\Omega_j)(\zeta) d\zeta = \int_{\Omega_j} I_\Lambda(\Omega_k)(\chi) d\chi \\ &= \int_{\Omega_k} \int_{\Omega_j} G_\Lambda(\zeta - \chi) d\chi d\zeta. \end{aligned}$$

LEMMA 2.2. *Minimizing $\mathcal{F}(\alpha)$ with respect to α of unit area is equivalent to minimizing*

$$(2.13) \quad \tilde{\mathcal{F}}(\alpha) = H_\Lambda(0) + G_\Lambda\left(\frac{\alpha_1 + \alpha_2}{2}\right) + b\left(G_\Lambda\left(\frac{\alpha_1}{2}\right) + G_\Lambda\left(\frac{\alpha_2}{2}\right)\right), \quad |D_\alpha| = 1,$$

where Λ is the lattice generated by α and

$$(2.14) \quad b = \frac{2\gamma_{12}r_1^2r_2^2}{\gamma_{11}r_1^4 + \gamma_{22}r_2^4} = \frac{2\gamma_{12}\omega_1\omega_2}{\gamma_{11}\omega_1^2 + \gamma_{22}\omega_2^2}.$$

Proof. Given a disc $B(\xi_j, r_j)$ one finds

$$(2.15) \quad \begin{aligned} I_\Lambda(B(\xi_j, r_j))(\zeta) &= \begin{cases} -\frac{|\zeta - \xi_j|^2}{4} + \frac{r_j^2}{4} - \frac{r_j^2}{2} \log r_j & \text{if } |\zeta - \xi_j| \in [0, r_j], \\ -\frac{r_j^2}{2} \log |\zeta - \xi_j| & \text{if } |\zeta - \xi_j| > r_j, \end{cases} \\ &\quad - \frac{r_j^2}{2} \log \frac{2\pi}{\sqrt{|\Lambda|}} + \frac{1}{4|\Lambda|} \left(\pi r_j^2 |\zeta - \xi_j|^2 + \frac{\pi r_j^4}{2} \right) + \pi r_j^2 H_\Lambda(\zeta - \xi_j) \end{aligned}$$

by (2.10) and the mean value property of the harmonic function H_Λ . Then

$$(2.16) \quad \begin{aligned} \int_{B(\xi_j, r_j)} \int_{B(\xi_j, r_j)} G_\Lambda(\zeta - \chi) d\chi d\zeta &= \int_{B(\xi_j, r_j)} I_\Lambda(B(\xi_j, r_j))(\zeta) d\zeta \\ &= \pi^2 r_j^4 H_\Lambda(0) + \frac{\pi r_j^4}{8} - \frac{\pi r_j^4}{2} \log \frac{2\pi r_j}{\sqrt{|\Lambda|}} + \frac{\pi^2 r_j^6}{4|\Lambda|}. \end{aligned}$$

When $j \neq k$ and $B(\xi_k, r_k) \cap B(\xi_j, r_j) = \emptyset$,

$$(2.17) \quad \begin{aligned} \int_{B(\xi_k, r_k)} \int_{B(\xi_j, r_j)} G_\Lambda(\zeta - \chi) d\chi d\zeta &= \int_{B(\xi_k, r_k)} I_\Lambda(B(\xi_j, r_j))(\zeta) d\zeta \\ &= \pi^2 r_j^2 r_k^2 G_\Lambda(\xi_j - \xi_k) + \frac{\pi^2 (r_j^2 r_k^4 + r_j^4 r_k^2)}{8|\Lambda|}. \end{aligned}$$

Since only the role played by the lattice basis α is of interest, let

$$(2.18) \quad c_{jj} = \frac{\pi r_j^4}{8} - \frac{\pi r_j^4}{2} \log \frac{2\pi r_j}{\sqrt{|\Lambda|}} + \frac{\pi^2 r_j^6}{4|\Lambda|}, \quad c_{jk} = \frac{\pi^2 (r_j^2 r_k^4 + r_j^4 r_k^2)}{8|\Lambda|}, \quad j \neq k,$$

which are independent of α when $|\Lambda| = 1$. Then

$$(2.19) \quad \int_{B(\xi_j, r_j)} \int_{B(\xi_j, r_j)} G_\Lambda(\zeta - \chi) d\chi d\zeta = \pi^2 r_j^4 H_\Lambda(0) + c_{jj},$$

$$(2.20) \quad \int_{B(\xi_k, r_k)} \int_{B(\xi_j, r_j)} G_\Lambda(\zeta - \chi) d\chi d\zeta = \pi^2 r_j^2 r_k^2 G_\Lambda(\xi_j - \xi_k) + c_{jk}.$$

Similarly,

$$(2.21) \quad \int_{B(\xi'_j, r_j)} \int_{B(\xi'_j, r_j)} G_\Lambda(\zeta - \chi) d\chi d\zeta = \pi^2 r_j^4 H_\Lambda(0) + c_{jj},$$

$$(2.22) \quad \int_{B(\xi'_k, r_k)} \int_{B(\xi'_j, r_j)} G_\Lambda(\zeta - \chi) d\chi d\zeta = \pi^2 r_j^2 r_k^2 G_\Lambda(\xi'_j - \xi'_k) + c_{jk}, \quad j \neq k,$$

$$(2.23) \quad \int_{B(\xi_k, r_k)} \int_{B(\xi'_j, r_j)} G_\Lambda(\zeta - \chi) d\chi d\zeta = \pi^2 r_j^2 r_k^2 G_\Lambda(\xi_j - \xi'_k) + c'_{jk}, \quad j, k = 1, 2,$$

where

$$(2.24) \quad c'_{jk} = \frac{\pi^2(r_j^2 r_k^4 + r_j^4 r_k^2)}{8|\Lambda|}, \quad j, k = 1, 2.$$

Note that in (2.23) and (2.24) j may be equal to k .

To complete the computation, note that

$$\begin{aligned} & \int_{\Omega_{\alpha,1}} \int_{\Omega_{\alpha,1}} G_\Lambda(\zeta - \chi) d\chi d\zeta \\ &= \int_{B(\xi_1, r_1) \cup B(\xi'_1, r_1)} \int_{B(\xi_1, r_1) \cup B(\xi'_1, r_1)} G_\Lambda(\zeta - \chi) d\chi d\zeta \\ &= \int_{B(\xi_1, r_1)} \int_{B(\xi_1, r_1)} G_\Lambda(\zeta - \chi) d\chi d\zeta + \int_{B(\xi'_1, r_1)} \int_{B(\xi'_1, r_1)} G_\Lambda(\zeta - \chi) d\chi d\zeta \\ &\quad + \int_{B(\xi_1, r_1)} \int_{B(\xi'_1, r_1)} G_\Lambda(\zeta - \chi) d\chi d\zeta + \int_{B(\xi'_1, r_1)} \int_{B(\xi_1, r_1)} G_\Lambda(\zeta - \chi) d\chi d\zeta \\ &= 2(\pi^2 r_1^4 H_\Lambda(0) + c_{11}) + 2(\pi^2 r_1^4 G_\Lambda(\xi_1 - \xi'_1) + c'_{11}) \int_{\Omega_{\alpha,2}} \int_{\Omega_{\alpha,2}} G_\Lambda(\zeta - \chi) d\chi d\zeta \\ &= 2(\pi^2 r_2^4 H_\Lambda(0) + c_{22}) + 2(\pi^2 r_2^4 G_\Lambda(\xi_2 - \xi'_2) + c'_{22}) \int_{\Omega_{\alpha,1}} \int_{\Omega_{\alpha,2}} G_\Lambda(\zeta - \chi) d\chi d\zeta \\ &= \pi^2 r_1^2 r_2^2 G_\Lambda(\xi_1 - \xi_2) + c_{12} + \pi^2 r_1^2 r_2^2 G_\Lambda(\xi'_1 - \xi'_2) + c_{12} \\ &\quad + \pi^2 r_1^2 r_2^2 G_\Lambda(\xi_1 - \xi'_2) + c'_{12} + \pi^2 r_1^2 r_2^2 G_\Lambda(\xi'_1 - \xi_2) + c'_{12}. \end{aligned}$$

In accordance with (1.17) and (1.18) choose

$$(2.25) \quad \xi_1 = \frac{3}{4}\alpha_1 + \frac{1}{4}\alpha_2, \quad \xi'_1 = \frac{1}{4}\alpha_1 + \frac{3}{4}\alpha_2,$$

$$(2.26) \quad \xi_2 = \frac{1}{4}\alpha_1 + \frac{1}{4}\alpha_2, \quad \xi'_2 = \frac{3}{4}\alpha_1 + \frac{3}{4}\alpha_2,$$

to derive

$$\begin{aligned}
 \mathcal{F}(\alpha) &= \sum_{j,k=1,2} \frac{\gamma_{jk}}{2} \int_{\Omega_{\alpha,j}} \int_{\Omega_{\alpha,k}} G_{\Lambda}(\zeta - \chi) d\chi d\zeta \\
 &= \gamma_{11} \left(\pi^2 r_1^4 H_{\Lambda}(0) + \pi^2 r_1^4 G_{\Lambda}\left(\frac{\alpha_1 - \alpha_2}{2}\right) + c_{11} + c'_{11} \right) \\
 &\quad + \gamma_{22} \left(\pi^2 r_2^4 H_{\Lambda}(0) + \pi^2 r_2^4 G_{\Lambda}\left(\frac{\alpha_1 + \alpha_2}{2}\right) + c_{22} + c'_{22} \right) \\
 &\quad + 2\gamma_{12} \left(\pi^2 r_1^2 r_2^2 \left(G_{\Lambda}\left(\frac{\alpha_1}{2}\right) + G_{\Lambda}\left(\frac{\alpha_2}{2}\right) \right) + c_{12} + c'_{12} \right) \\
 &= (\gamma_{11} \pi^2 r_1^4 + \gamma_{22} \pi^2 r_2^4) \left(H_{\Lambda}(0) + G_{\Lambda}\left(\frac{\alpha_1 + \alpha_2}{2}\right) \right) \\
 &\quad + 2\gamma_{12} \pi^2 r_1^2 r_2^2 \left(G_{\Lambda}\left(\frac{\alpha_1}{2}\right) + G_{\Lambda}\left(\frac{\alpha_2}{2}\right) \right) \\
 &\quad + \gamma_{11}(c_{11} + c'_{11}) + \gamma_{22}(c_{22} + c'_{22}) + 2\gamma_{12}(c_{12} + c'_{12}) \\
 &= (\gamma_{11} \pi^2 r_1^4 + \gamma_{22} \pi^2 r_2^4) \\
 &\quad \cdot \left[H_{\Lambda}(0) + G_{\Lambda}\left(\frac{\alpha_1 + \alpha_2}{2}\right) + \frac{2\gamma_{12} r_1^2 r_2^2}{\gamma_{11} r_1^4 + \gamma_{22} r_2^4} \left(G_{\Lambda}\left(\frac{\alpha_1}{2}\right) + G_{\Lambda}\left(\frac{\alpha_2}{2}\right) \right) \right] \\
 &\quad + \gamma_{11}(c_{11} + c'_{11}) + \gamma_{22}(c_{22} + c'_{22}) + 2\gamma_{12}(c_{12} + c'_{12}).
 \end{aligned}$$

Here $G_{\Lambda}\left(\frac{\alpha_1 + \alpha_2}{2}\right) = G_{\Lambda}\left(\frac{\alpha_1 - \alpha_2}{2}\right)$ follows from the Λ -periodicity of G_{Λ} . $\square$

Calculations based on (2.11) show

$$(2.27) \quad H_{\Lambda}(0) = -\frac{1}{2\pi} \log \left| \sqrt{\operatorname{Im} \tau} e \left(\frac{\tau}{12} \right) \prod_{n=1}^{\infty} (1 - e(n\tau))^2 \right|,$$

$$(2.28) \quad G_{\Lambda}\left(\frac{\alpha_1}{2} + \frac{\alpha_2}{2}\right) = -\frac{1}{2\pi} \log \left| e \left(-\frac{\tau}{24} \right) \prod_{n=1}^{\infty} \left(1 + e \left(\left(n - \frac{1}{2} \right) \tau \right) \right)^2 \right|$$

$$(2.29) \quad G_{\Lambda}\left(\frac{\alpha_1}{2}\right) = -\frac{1}{2\pi} \log \left| 2 e \left(\frac{\tau}{12} \right) \prod_{n=1}^{\infty} (1 + e(n\tau))^2 \right|$$

$$(2.30) \quad G_{\Lambda}\left(\frac{\alpha_2}{2}\right) = -\frac{1}{2\pi} \log \left| e \left(-\frac{\tau}{24} \right) \prod_{n=1}^{\infty} \left(1 - e \left(\left(n - \frac{1}{2} \right) \tau \right) \right)^2 \right|.$$

To derive (2.27) we have used $\frac{1}{|\alpha_1|} = \sqrt{\operatorname{Im} \tau}$, which follows from $1 = |D_{\Lambda}| = \operatorname{Im}(\overline{\alpha_1} \alpha_2)$.

Regarding the four infinite products in (2.27) through (2.30), one has the following formulas.

LEMMA 2.3.

$$\begin{aligned}
 &\prod_{n=1}^{\infty} (1 - e(n\tau)) \prod_{n=1}^{\infty} \left(1 + e \left(\left(n - \frac{1}{2} \right) \tau \right) \right) = \prod_{n=1}^{\infty} \left(1 - e \left(n \frac{\tau + 1}{2} \right) \right) \\
 &\prod_{n=1}^{\infty} \left(1 + e \left(\left(n - \frac{1}{2} \right) \tau \right) \right) \prod_{n=1}^{\infty} (1 + e(n\tau)) \prod_{n=1}^{\infty} \left(1 - e \left(\left(n - \frac{1}{2} \right) \tau \right) \right) = 1.
 \end{aligned}$$

Proof. To prove the first formula, rewrite and rearrange the terms as follows:

$$\begin{aligned} & \prod_{n=1}^{\infty} (1 - e(n\tau)) \prod_{n=1}^{\infty} \left(1 + e\left(\left(n - \frac{1}{2}\right)\tau\right)\right) \\ &= \prod_{n=1}^{\infty} \left(1 - e\left(2n\frac{\tau+1}{2}\right)\right) \prod_{n=1}^{\infty} \left(1 - e\left((2n-1)\frac{\tau+1}{2}\right)\right) \\ &= \prod_{n=1}^{\infty} \left(1 - e\left(n\frac{\tau+1}{2}\right)\right). \end{aligned}$$

For the second formula, consider

$$\begin{aligned} & \prod_{n=1}^{\infty} (1 - e(n\tau)) \prod_{n=1}^{\infty} \left(1 + e\left(\left(n - \frac{1}{2}\right)\tau\right)\right) \prod_{n=1}^{\infty} (1 + e(n\tau)) \prod_{n=1}^{\infty} \left(1 - e\left(\left(n - \frac{1}{2}\right)\tau\right)\right) \\ &= \prod_{n=1}^{\infty} (1 - e(2n\tau)) \prod_{n=1}^{\infty} (1 - e((2n-1)\tau)) \\ &= \prod_{n=1}^{\infty} (1 - e(n\tau)). \end{aligned}$$

The second formula follows after one divides out $\prod_{n=1}^{\infty} (1 - e(n\tau))$. $\square$

These identities will allow us to further simplify $\tilde{\mathcal{F}}(\alpha)$. Let

$$(2.31) \quad \mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$$

be the upper half of the complex plane. Define

$$(2.32) \quad f_b(z) = b \log |\operatorname{Im}(z)\eta(z)| + (1-b) \log \left| \operatorname{Im} \left(\frac{z+1}{2} \right) \eta \left(\frac{z+1}{2} \right) \right|, \quad z \in \mathbb{H},$$

where

$$(2.33) \quad \eta(z) = e\left(\frac{z}{6}\right) \prod_{n=1}^{\infty} (1 - e(nz))^4.$$

One often writes f_b as

$$(2.34) \quad f_b(z) = bf_1(z) + (1-b)f_0(z),$$

where

$$(2.35) \quad f_1(z) = \log |\operatorname{Im}(z)\eta(z)|, \quad f_0(z) = \log \left| \operatorname{Im} \left(\frac{z+1}{2} \right) \eta \left(\frac{z+1}{2} \right) \right|.$$

LEMMA 2.4. *Minimizing $\tilde{\mathcal{F}}(\alpha)$ with respect to α of unit area is equivalent to maximizing $f_b(z)$ with respect to z in $\mathbb{H}$.*

Proof. By the first formula in Lemma 2.3,

$$\begin{aligned} & H_{\Lambda}(0) + G_{\Lambda}\left(\frac{\alpha_1 + \alpha_2}{2}\right) = -\frac{1}{2\pi} \log \left| \sqrt{\operatorname{Im} \tau} e\left(\frac{\tau}{24}\right) \prod_{n=1}^{\infty} \left(1 - e\left(n\frac{\tau+1}{2}\right)\right)^2 \right| \\ (2.36) \quad & = -\frac{1}{4\pi} \log \left| \operatorname{Im} \left(\frac{\tau+1}{2} \right) \eta \left(\frac{\tau+1}{2} \right) \right| - \frac{1}{4\pi} \log 2. \end{aligned}$$

Using both formulas in Lemma 2.3, one deduces

$$\begin{aligned}
 G_{\Lambda}\left(\frac{\alpha_1}{2}\right) + G_{\Lambda}\left(\frac{\alpha_2}{2}\right) &= -\frac{1}{2\pi} \log \left| 2e\left(\frac{\tau}{24}\right) \prod_{n=1}^{\infty} (1 + e(n\tau))^2 \prod_{n=1}^{\infty} \left(1 - e\left(\left(n - \frac{1}{2}\right)\tau\right)\right)^2 \right| \\
 &= -\frac{1}{2\pi} \log \left| 2e\left(\frac{\tau}{24}\right) \prod_{n=1}^{\infty} (1 - e(n\tau))^2 / \prod_{n=1}^{\infty} \left(1 - e\left(n\frac{\tau+1}{2}\right)\right)^2 \right| \\
 (2.37) \quad &= -\frac{1}{4\pi} \left(\log |\operatorname{Im}(\tau)\eta(\tau)| - \log \left| \operatorname{Im}\left(\frac{\tau+1}{2}\right)\eta\left(\frac{\tau+1}{2}\right) \right| \right) - \frac{1}{4\pi} \log 2.
 \end{aligned}$$

By (2.36) and (2.37), $\tilde{\mathcal{F}}(\alpha)$ of (2.13) is reduced to

$$(2.38) \quad \tilde{\mathcal{F}}(\alpha) = -\frac{1}{4\pi} \left(b \log |\operatorname{Im}(\tau)\eta(\tau)| + (1-b) \log \left| \operatorname{Im}\left(\frac{\tau+1}{2}\right)\eta\left(\frac{\tau+1}{2}\right) \right| \right) - \frac{1+b}{4\pi} \log 2,$$

from which the lemma follows. $\square$

3. Duality property of f_b . The function η in the definition of f_b satisfies two functional equations.

LEMMA 3.1. *For all $z \in \mathbb{H}$,*

$$(3.1) \quad \eta(z+1) = e^{\frac{\pi i}{3}} \eta(z),$$

$$(3.2) \quad \eta\left(-\frac{1}{z}\right) = -z^2 \eta(z).$$

Proof. The function η in (2.33) is the fourth power of the Dedekind eta function which is

$$\eta_D(z) = e\left(\frac{z}{24}\right) \prod_{n=1}^{\infty} (1 - e(nz))$$

so

$$\eta(z) = \eta_D^4(z), \quad z \in \mathbb{H}.$$

For the Dedekind eta function, it is known [3, Chapter 2] that

$$(3.3) \quad \eta_D(z+1) = e^{\frac{\pi i}{12}} \eta_D(z),$$

$$(3.4) \quad \eta_D\left(-\frac{1}{z}\right) = \sqrt{-iz} \eta_D(z),$$

where $\sqrt{\cdot}$ stands for the principal branch of the square root. $\square$

These functional equations lead to invariance properties.

LEMMA 3.2.

1. $|\operatorname{Im}(z)\eta(z)|$ and, consequently, $f_1(z)$, are invariant under the transforms

$$z \rightarrow z+1 \quad \text{and} \quad z \rightarrow -\frac{1}{z}.$$

2. $|\operatorname{Im}\left(\frac{z+1}{2}\right)\eta\left(\frac{z+1}{2}\right)|$, and, consequently, $f_0(z)$, are invariant under the transforms

$$z \rightarrow z+2 \quad \text{and} \quad z \rightarrow -\frac{1}{z}.$$

Proof. The invariance of $|\operatorname{Im}(z)\eta(z)|$ under $z \rightarrow z+1$ and the invariance of $|\operatorname{Im}(\frac{z+1}{2})\eta(\frac{z+1}{2})|$ under $z \rightarrow z+2$ follow from (3.1). By (3.2) it is easy to see that

$$(3.5) \quad \left| \operatorname{Im}\left(-\frac{1}{z}\right) \eta\left(-\frac{1}{z}\right) \right| = |\operatorname{Im}(z)\eta(z)|,$$

so $|\operatorname{Im}(z)\eta(z)|$ is invariant under $z \rightarrow -\frac{1}{z}$.

The invariance of $|\operatorname{Im}(z)\eta(z)|$ under $z \rightarrow z+1$ implies its invariance under $z \rightarrow z+k$ for any integer k . Now one deduces

$$(3.6) \quad \begin{aligned} \left| \operatorname{Im}\left(\frac{(-\frac{1}{z})+1}{2}\right) \eta\left(\frac{(-\frac{1}{z})+1}{2}\right) \right| &= \left| \operatorname{Im}\left(\frac{z-1}{2z}\right) \eta\left(\frac{z-1}{2z}\right) \right| \\ &= \left| \operatorname{Im}\left(\frac{-z-1}{2z}\right) \eta\left(\frac{-z-1}{2z}\right) \right| \\ &= \left| \operatorname{Im}\left(\frac{2z}{z+1}\right) \eta\left(\frac{2z}{z+1}\right) \right| \\ &= \left| \operatorname{Im}\left(\frac{-2}{z+1}\right) \eta\left(\frac{-2}{z+1}\right) \right| \\ &= \left| \operatorname{Im}\left(\frac{z+1}{2}\right) \eta\left(\frac{z+1}{2}\right) \right| \end{aligned}$$

by applying the invariance of $|\operatorname{Im}(z)\eta(z)|$ under $z \rightarrow z-1$, $z \rightarrow -\frac{1}{z}$, $z \rightarrow z-2$, and $z \rightarrow -\frac{1}{z}$ successively. This proves invariance of $|\operatorname{Im}(\frac{z+1}{2})\eta(\frac{z+1}{2})|$ under $z \rightarrow -\frac{1}{z}$. $\square$

There is another invariance that is not a linear fractional transform: the reflection about the imaginary axis.

LEMMA 3.3. *Both $|\operatorname{Im}(z)\eta(z)|$ and $|\operatorname{Im}(\frac{z+1}{2})\eta(\frac{z+1}{2})|$ and, consequently, $f_1(z)$ and $f_0(z)$, are invariant under $z \rightarrow -\bar{z}$.*

Proof. These follow easily from the infinite product definition (2.33) of η . $\square$

The transforms $z \rightarrow z+1$ and $z \rightarrow -\frac{1}{z}$ generate the modular group Γ ,

$$\Gamma = \left\{ z \rightarrow \frac{az+b}{cz+d} : a, b, c, d \in \mathbb{Z}, ad-bc=1 \right\}$$

and Γ has

$$(3.7) \quad F_\Gamma = \{z \in \mathbb{H} : |z| > 1, -1/2 < \operatorname{Re} z < 1/2\}$$

as a fundamental region. It means that every orbit under this group has one element in $\overline{F_\Gamma}$, the closure of F_Γ in $\mathbb{H}$, and no two points in F_Γ belong to the same orbit [3].

The transforms $z \rightarrow z+2$ and $z \rightarrow -\frac{1}{z}$ generate a subgroup Γ' of Γ ,

$$(3.8) \quad \begin{aligned} \Gamma' = \left\{ z \rightarrow \frac{az+b}{cz+d} \in \Gamma : \begin{aligned} &a \equiv d \equiv 1 \pmod{2} \quad \text{and} \quad b \equiv c \equiv 0 \pmod{2}, \\ &\text{or } a \equiv d \equiv 0 \pmod{2} \quad \text{and} \quad b \equiv c \equiv 1 \pmod{2} \end{aligned} \right\}. \end{aligned}$$

It is known in number theory that this group has

$$(3.9) \quad F_{\Gamma'} = \{z \in \mathbb{H} : |z| > 1, -1 < \operatorname{Re} z < 1\}$$

as a fundamental region [10].

Denote by $\mathcal{G}$ the group of diffeomorphisms of $\mathbb{H}$ generated by

$$z \rightarrow z + 2, \quad z \rightarrow -\frac{1}{z}, \quad z \rightarrow -\bar{z}.$$

Note that Γ' is a subgroup of $\mathcal{G}$ but Γ is not a subgroup of $\mathcal{G}$.

With the group $\mathcal{G}$, maximizing f_b need not be carried out in $\mathbb{H}$, but in a smaller set which contains at least one element from each orbit of $\mathcal{G}$. Let

$$(3.10) \quad W = \{z \in \mathbb{H} : 0 < \operatorname{Re} z < 1, |z| > 1\}$$

and

$$(3.11) \quad \overline{W}_{\mathbb{H}} = \{z \in \mathbb{H} : 0 \leq \operatorname{Re} z \leq 1, |z| \geq 1\};$$

see Figure 3. Note that $\overline{W}_{\mathbb{H}}$ is the closure of W in $\mathbb{H}$ so $1 \notin \overline{W}_{\mathbb{H}}$.

LEMMA 3.4.

1. $|\operatorname{Im}(z)\eta(z)|$ and $f_1(z)$ are invariant under the group Γ and the transform $z \rightarrow -\bar{z}$.
2. $|\operatorname{Im}(\frac{z+1}{2})\eta(\frac{z+1}{2})|$, $f_0(z)$, and $f_b(z)$ for $b \in \mathbb{R}$, are invariant under the groups Γ' and $\mathcal{G}$.
3. As the group $\mathcal{G}$ acts on $\mathbb{H}$, each orbit of $\mathcal{G}$ has at least one element in $\overline{W}_{\mathbb{H}}$.

Proof. Parts 1 and 2 follow from Lemmas 3.2 and 3.3. Part 3 follows from $F_{\Gamma'}$ being the fundamental region of Γ' and the transform $z \rightarrow -\bar{z} \in \mathcal{G}$. $\square$

It is instructive to understand transforms from the viewpoint of bases. Let $\alpha = (\alpha_1, \alpha_2)$ and $\alpha' = (\alpha'_1, \alpha'_2)$ be two bases of unit area that define lattices Λ and Λ' , and two species periodic assemblies $(\Omega_{\alpha,1}, \Omega_{\alpha,2})$ and $(\Omega_{\alpha',1}, \Omega_{\alpha',2})$. Set $z = \frac{\alpha_2}{\alpha_1}$ and $z' = \frac{\alpha'_2}{\alpha'_1}$.

If α and α' are related by the transform $z \rightarrow z' = z + 1$, then

$$\frac{\alpha'_2}{\alpha'_1} = \frac{\alpha_1 + \alpha_2}{\alpha_1}$$

and, consequently, there exists $\kappa \in \mathbb{C}$ such that $\alpha'_1 = \kappa\alpha_1$ and $\alpha'_2 = \kappa(\alpha_1 + \alpha_2)$. Since both bases have unit area,

$$1 = \operatorname{Im}(\overline{\alpha'_1}\alpha'_2) = |\kappa|^2 \operatorname{Im}(\overline{\alpha_1}(\alpha_1 + \alpha_2)) = |\kappa|^2 \operatorname{Im}(\overline{\alpha_1}\alpha_2) = |\kappa|^2.$$

So there exists $\theta \in [0, 2\pi)$ such that $\kappa = e^{\theta i}$ and $\alpha' = e^{\theta i}(\alpha_1, \alpha_1 + \alpha_2)$. Let $\alpha'' = e^{\theta i}\alpha = e^{\theta i}(\alpha_1, \alpha_2)$. Then α' and α'' generate the same lattice Λ' and, consequently, Λ and Λ' are isometric in the sense that a period parallelogram D_{α} of Λ is just a rotation of a period parallelogram $D_{\alpha''}$ of Λ' . However the assemblies $(\Omega_{\alpha,1}, \Omega_{\alpha,2})$ and $(\Omega_{\alpha',1}, \Omega_{\alpha',2})$ are usually quite different since in general $\mathcal{J}_{\Lambda}(\Omega_{\alpha,1}, \Omega_{\alpha,2}) \neq \mathcal{J}_{\Lambda'}(\Omega_{\alpha',1}, \Omega_{\alpha',2})$.

The story changes if α and α' are related by the transform $z \rightarrow z' = z + 2$. This time not only Λ and Λ' are isometric, but $\mathcal{J}_{\Lambda}(\Omega_{\alpha,1}, \Omega_{\alpha,2}) = \mathcal{J}_{\Lambda'}(\Omega_{\alpha',1}, \Omega_{\alpha',2})$ as well by Lemma 3.4.2.

If α and α' are related by $z \rightarrow z' = -\frac{1}{z}$, one can show that $(\alpha'_1, \alpha'_2) = e^{\theta i}(-\alpha_2, \alpha_1)$, $\theta \in [0, 2\pi)$. Then Λ and Λ' are isometric since D_{α} and $D_{\alpha'}$ differ by a translation and rotation. Moreover $\mathcal{J}_{\Lambda}(\Omega_{\alpha,1}, \Omega_{\alpha,2}) = \mathcal{J}_{\Lambda'}(\Omega_{\alpha',1}, \Omega_{\alpha',2})$ by Lemma 3.4.2.

Finally if α and α' are related by $z \rightarrow z' = -\bar{z}$, then $(\alpha'_1, \alpha'_2) = e^{\theta i}(\overline{\alpha_1}, -\overline{\alpha_2})$, $\theta \in [0, 2\pi)$. Therefore D_{α} and $D_{\alpha'}$ differ by a mirror reflection, a translation and

a rotation, so Λ and Λ' are again isometric. By Lemma 3.4.2, $\mathcal{J}_\Lambda(\Omega_{\alpha,1}, \Omega_{\alpha,2}) = \mathcal{J}_{\Lambda'}(\Omega_{\alpha',1}, \Omega_{\alpha',2})$.

In summary if z and z' are related by an element $g \in \mathcal{G}$, i.e., $z' = gz$, then Λ and Λ' are isometric and $\mathcal{J}_\Lambda(\Omega_{\alpha,1}, \Omega_{\alpha,2}) = \mathcal{J}_{\Lambda'}(\Omega_{\alpha',1}, \Omega_{\alpha',2})$. For isometric Λ and Λ' , if Λ is a rectangular lattice, then Λ' is also a rectangular lattice of the same ratio of longer side to shorter side; if Λ is a rhombic lattice, then Λ' is also a rhombic lattice of the same acute angle. Therefore to prove Theorem 1.1, it suffices to find every minimum of f_b in $\overline{W}_\mathbb{H}$ and identify its associated lattice.

In the exceptional case $b = 1$, since f_1 is invariant under Γ , it suffices to maximize f_1 in a smaller set

$$(3.12) \quad \overline{U}_\mathbb{H} = \{z \in \mathbb{H} : 0 \leq \operatorname{Re} z \leq 1/2, |z| \geq 1\}$$

which is the closure of

$$(3.13) \quad U = \left\{z \in \mathbb{C} : |z| > 1, 0 < \operatorname{Re} z < \frac{1}{2}\right\}.$$

This fact was used critically in [6], but it is not valid if $b \neq 1$. The approach of this paper works for all $b \in [0, 1]$, giving a different proof for the $b = 1$ case as well.

One of the most important properties of f_b is the following duality relation between f_b and f_{1-b} .

LEMMA 3.5. *Under the transform $z \rightarrow w = \frac{z-1}{z+1}$ of $\mathbb{H}$,*

$$f_1(z) = f_0(w) \quad \text{and} \quad f_0(z) = f_1(w), \quad z \in \mathbb{H}, \quad \text{and} \quad w = \frac{z-1}{z+1} \in \mathbb{H}.$$

Consequently, for all $b \in \mathbb{R}$,

$$f_b(w) = f_{1-b}(z), \quad z \in \mathbb{H}, \quad \text{and} \quad w = \frac{z-1}{z+1} \in \mathbb{H}.$$

More generally, if $h : z' \rightarrow w' = \frac{z'-1}{z'+1}$ and $g_1 : z \rightarrow z'$, $g_2 : w' \rightarrow w$ are transforms in $\mathcal{G}$, then

$$f_b(w) = f_{1-b}(z), \quad z \in \mathbb{H}, \quad \text{and} \quad w = g_2 \circ h \circ g_1(z) \in \mathbb{H}.$$

Proof. The transform $z \rightarrow w' = \frac{z-1}{z+1}$ is in Γ , so

$$f_1(z) = f_1(w')$$

by the invariance of f_1 under Γ . On the other hand substitution shows

$$f_0(z) = f_0\left(\frac{w'}{-w'+1}\right) = \log \left| \operatorname{Im} \left(\frac{1}{-2w'+2} \right) \eta \left(\frac{1}{-2w'+2} \right) \right| = f_1\left(\frac{1}{-2w'+2}\right).$$

Now apply another transform $w' \rightarrow w = 2w' - 1$ which is not in Γ to find

$$f_1(z) = f_1\left(\frac{w+1}{2}\right) = f_0(w)$$

and

$$f_0(z) = f_1\left(\frac{1}{-w+1}\right) = f_1(w),$$

where the last equation follows from the invariance of f_1 under $w \rightarrow \frac{1}{-w+1} \in \Gamma$.

The composition of the two transforms is $z \rightarrow w' = \frac{z-1}{z+1} \rightarrow w = 2w' - 1 = \frac{z-1}{z+1}$. $\square$

Although b is supposed to be in $[0, 1]$ throughout this paper, some properties of f_b , like Lemma 3.5, hold for $b \in \mathbb{R}$. If this is the case we state so explicitly.

Let us write $z = x + yi$ henceforth, and set

$$(3.14) \quad X_b(z) = \frac{\partial f_b(z)}{\partial x} = bX_1(z) + (1-b)X_0(z), \quad Y_b(z) = \frac{\partial f_b(z)}{\partial y} = bY_1(z) + (1-b)Y_0(z),$$

where

$$(3.15) \quad X_1(z) = \frac{\partial}{\partial x} \log |\operatorname{Im}(z)\eta(z)|, \quad Y_1(z) = \frac{\partial}{\partial y} \log |\operatorname{Im}(z)\eta(z)|,$$

$$(3.16) \quad X_0(z) = \frac{\partial}{\partial x} \log \left| \operatorname{Im} \left(\frac{z+1}{2} \right) \eta \left(\frac{z+1}{2} \right) \right|, \quad Y_0(z) = \frac{\partial}{\partial y} \log \left| \operatorname{Im} \left(\frac{z+1}{2} \right) \eta \left(\frac{z+1}{2} \right) \right|.$$

These functions can be written as the following series:

$$(3.17) \quad X_1(z) = \sum_{n=1}^{\infty} \frac{8\pi n \sin 2\pi n x}{e^{2\pi n y} + e^{-2\pi n y} - 2 \cos 2\pi n x},$$

$$(3.18) \quad Y_1(z) = \frac{1}{y} - \frac{\pi}{3} + \sum_{n=1}^{\infty} \frac{-8\pi n e^{-2\pi n y} + 8\pi n \cos 2\pi n x}{e^{2\pi n y} + e^{-2\pi n y} - 2 \cos 2\pi n x},$$

$$(3.19) \quad X_0(z) = \sum_{n=1}^{\infty} \frac{4\pi n \sin \pi n(x+1)}{e^{\pi n y} + e^{-\pi n y} - 2 \cos \pi n(x+1)},$$

$$(3.20) \quad Y_0(z) = \frac{1}{y} - \frac{\pi}{6} + \sum_{n=1}^{\infty} \frac{-4\pi n e^{-\pi n y} + 4\pi n \cos \pi n(x+1)}{e^{\pi n y} + e^{-\pi n y} - 2 \cos \pi n(x+1)}.$$

We end this section with two formulas that relate f_b on the upper half of the unit circle to f_{1-b} on the upper half of the imaginary axis.

LEMMA 3.6. *Let the upper half of the unit circle be parametrized by $u + i\sqrt{1-u^2}$, $u \in (-1, 1)$. Then $\frac{\sqrt{1-u^2}}{1-u}i$ parametrizes the upper half of the imaginary axis, and*

$$(3.21) \quad X_b(u + i\sqrt{1-u^2}) = \frac{\sqrt{1-u^2}}{1-u} Y_{1-b} \left(\frac{\sqrt{1-u^2}}{1-u} i \right),$$

$$(3.22) \quad Y_b(u + i\sqrt{1-u^2}) = \frac{-u}{1-u} Y_{1-b} \left(\frac{\sqrt{1-u^2}}{1-u} i \right),$$

hold for $u \in (-1, 1)$.

Proof. Consider the transform in Lemma 3.5, $z \rightarrow w = \frac{z-1}{z+1}$. With $z = x + yi$ and $w = u + vi$,

$$(3.23) \quad u = \frac{x^2 + y^2 - 1}{(x+1)^2 + y^2}, \quad v = \frac{2y}{(x+1)^2 + y^2}.$$

Conversely,

$$(3.24) \quad x = \frac{1 - u^2 - v^2}{(1-u)^2 + v^2}, \quad y = \frac{2v}{(1-u)^2 + v^2}.$$

Differentiate $f_b(w) = f_{1-b}(z)$ with respect to u and v to find

$$\begin{aligned} X_b(w) &= X_{1-b}(z) \frac{\partial x}{\partial u} + Y_{1-b}(z) \frac{\partial y}{\partial u}, \\ Y_b(w) &= X_{1-b}(z) \frac{\partial x}{\partial v} + Y_{1-b}(z) \frac{\partial y}{\partial v}. \end{aligned}$$

When w is on the unit circle, z is on the imaginary axis. Since f_{1-b} is invariant under the reflection about the imaginary axis, $X_{1-b}(z) = 0$ on the imaginary axis. Also

$$(3.25) \quad \frac{\partial y}{\partial u} \Big|_{|w|=1} = \frac{\sqrt{1-u^2}}{1-u}, \quad \frac{\partial y}{\partial v} \Big|_{|w|=1} = \frac{-u}{1-u}$$

from which the lemma follows. $\square$

4. f_b on the imaginary axis. The behavior of f_b on the imaginary axis is studied in this section. Let us record some of the derivatives of f_1 and f_0 on the imaginary axis for easy reference. Let

$$(4.1) \quad r = e^{-\pi y}.$$

Then by (3.18) and (3.20),

$$(4.2) \quad Y_1(yi) = \frac{1}{y} - \frac{\pi}{3} + \sum_{n=1}^{\infty} \frac{8\pi n r^{2n}}{1-r^{2n}},$$

$$(4.3) \quad \frac{\partial Y_1(yi)}{\partial y} = -\frac{1}{y^2} - \sum_{n=1}^{\infty} \frac{16\pi^2 n^2 r^{2n}}{(1-r^{2n})^2},$$

$$(4.4) \quad \frac{\partial^2 Y_1(yi)}{\partial y^2} = \frac{2}{y^3} + \sum_{n=1}^{\infty} \frac{32\pi^3 n^3 (r^{2n} + r^{4n})}{(1-r^{2n})^3},$$

$$(4.5) \quad \frac{\partial^3 Y_1(yi)}{\partial y^3} = -\frac{6}{y^4} - \sum_{n=1}^{\infty} \frac{64\pi^4 n^4 (r^{2n} + 4r^{4n} + r^{6n})}{(1-r^{2n})^4},$$

$$(4.6) \quad Y_0(yi) = \frac{1}{y} - \frac{\pi}{6} + \sum_{n=1}^{\infty} \frac{4\pi n (-r)^n}{1-(-r)^n},$$

$$(4.7) \quad \frac{\partial Y_0(yi)}{\partial y} = -\frac{1}{y^2} - \sum_{n=1}^{\infty} \frac{4\pi^2 n^2 (-r)^n}{(1-(-r)^n)^2},$$

$$(4.8) \quad \frac{\partial^2 Y_0(yi)}{\partial y^2} = \frac{2}{y^3} + \sum_{n=1}^{\infty} \frac{4\pi^3 n^3 ((-r)^n + r^{2n})}{(1-(-r)^n)^3},$$

$$(4.9) \quad \frac{\partial^3 Y_0(yi)}{\partial y^3} = -\frac{6}{y^4} - \sum_{n=1}^{\infty} \frac{4\pi^4 n^4 ((-r)^n + 4r^{2n} + (-r)^{3n})}{(1-(-r)^n)^4}.$$

LEMMA 4.1. For all $b \in \mathbb{R}$,

$$f_b(yi) = f_b\left(\frac{i}{y}\right), \quad y > 0.$$

Consequently,

$$Y_b(yi) = \left(-\frac{1}{y^2}\right) Y_b\left(\frac{i}{y}\right), \quad y > 0.$$

In particular

$$Y_b(i) = 0.$$

Proof. Apply the invariance of $f_b(z)$ under $z \rightarrow -\frac{1}{z}$ with $z = yi$, $y > 0$. Then differentiate with respect to y and set $y = 1$. $\square$

LEMMA 4.2. *The function $y \rightarrow f_1(yi)$, $y > 0$, has one critical point at $y = 1$. Moreover*

$$\begin{aligned} Y_1(yi) &> 0 \text{ if } y \in (0, 1), \\ Y_1(yi) &< 0 \text{ if } y \in (1, \infty). \end{aligned}$$

Proof. Lemma 4.1 asserts that $y = 1$ is a critical point of $y \rightarrow f_1(yi)$, $y > 0$, i.e., $Y_1(i) = 0$. Define

$$(4.10) \quad A(z) = \arg(z\eta(z)) = \arg(z) + \arg(\eta(z)).$$

Note

$$(4.11) \quad \operatorname{Re}(\log(z\eta(z))) = \log|z\eta(z)|, \quad \operatorname{Im}(\log(z\eta(z))) = A(z).$$

Hence A is a harmonic function. We consider $A(z)$ in U and its closure $\bar{U}_{\mathbb{H}}$ given in (3.13) and (3.12), respectively.

On the imaginary axis, for $y > 0$, since $\eta(yi)$ is real and positive,

$$(4.12) \quad A(yi) = \arg(yi) + \arg(\eta(yi)) = \frac{\pi}{2} + 0 = \frac{\pi}{2},$$

On the line $x = \frac{1}{2}$, $\arg(\eta(z)) = \frac{\pi}{6}$ since $e^{2\pi n(\frac{1}{2} + yi)i}$ is real, and

$$(4.13) \quad A\left(\frac{1}{2} + yi\right) = \arctan(2y) + \frac{\pi}{6}.$$

In particular

$$(4.14) \quad A\left(\frac{1}{2} + yi\right) > \frac{\pi}{2} \quad \text{if } y > \frac{\sqrt{3}}{2}.$$

As $y \rightarrow \infty$ in $z = x + yi$,

$$(4.15) \quad \lim_{y \rightarrow \infty} A(x + yi) = \frac{\pi}{2} + \frac{\pi x}{3} \quad \text{uniformly with respect to } x \in \left[0, \frac{1}{2}\right].$$

Now consider A on the unit circle. By the functional equation (3.2) one has, in polar coordinates $z = re^{i\theta}$,

$$\log(r|\eta(re^{i\theta})|) = \log\left(\frac{1}{r}\left|\eta\left(-\frac{1}{r}e^{-i\theta}\right)\right|\right).$$

By the definition of η , one sees that $|\eta(-\bar{\zeta})| = |\eta(\zeta)|$ for all $\zeta \in \mathbb{H}$. Therefore

$$\log(r|\eta(re^{i\theta})|) = \log\left(\frac{1}{r}\left|\eta\left(\frac{1}{r}e^{i\theta}\right)\right|\right).$$

Differentiating the last equation with respect to r and setting $r = 1$ afterwards, one derives

$$\frac{\partial}{\partial r}\bigg|_{r=1} \log(r|\eta(re^{i\theta})|) = 0.$$

One of the Cauchy–Riemann equations in polar coordinates for $\log(z\eta(z))$ is

$$\frac{\partial}{\partial r} \operatorname{Re}(\log(z\eta(z))) = \frac{1}{r} \frac{\partial}{\partial \theta} \operatorname{Im}(\log(z\eta(z))).$$

By (4.11)

$$\frac{\partial}{\partial \theta} A(e^{i\theta}) = 0,$$

namely, A is constant on the unit circle. Since $\eta(i)$ is real and positive, $A(i) = \frac{\pi}{2}$. Hence

$$(4.16) \quad A(z) = \frac{\pi}{2} \quad \text{if } |z| = 1 \text{ and } \frac{\pi}{3} \leq \arg z \leq \frac{\pi}{2}.$$

By (4.12), (4.14), (4.15), (4.16), and the maximum principle,

$$(4.17) \quad A(z) > \frac{\pi}{2}, \quad z \in U;$$

by the Hopf lemma,

$$(4.18) \quad \frac{\partial}{\partial x} \Big|_{x=0, y>1} A(z) > 0.$$

By a Cauchy–Riemann equation

$$(4.19) \quad Y_1(yi) = -\frac{\partial}{\partial x} \Big|_{x=0, y>1} A(z) < 0, \quad y \in (1, \infty).$$

For $y \in (0, 1)$, by Lemma 4.1,

$$(4.20) \quad Y_1(yi) = \left(-\frac{1}{y^2}\right) Y_1\left(\frac{i}{y}\right) > 0, \quad y \in (0, 1).$$

This completes the proof. $\square$

LEMMA 4.3. *The function $y \rightarrow f_0(yi)$, $y > 0$, has three critical points at $\frac{\sqrt{3}}{3}$, 1, and $\sqrt{3}$. Moreover*

$$\begin{aligned} Y_0(yi) &> 0 \text{ if } y \in (0, \sqrt{3}/3), \\ Y_0(yi) &< 0 \text{ if } y \in (\sqrt{3}/3, 1), \\ Y_0(yi) &> 0 \text{ if } y \in (1, \sqrt{3}), \\ Y_0(yi) &< 0 \text{ if } y \in (\sqrt{3}, \infty). \end{aligned}$$

Proof. The transform $z \rightarrow \frac{z-1}{2z-1} \in \Gamma$ maps $\frac{1}{2} + yi$ to $\frac{1}{2} + \frac{i}{4y}$, so

$$\log \left| y \eta \left(\frac{1}{2} + yi \right) \right| = \log \left| \frac{1}{4y} \eta \left(\frac{1}{2} + \frac{i}{4y} \right) \right|$$

Differentiation with respect to y shows that

$$(4.21) \quad Y_1\left(\frac{1}{2} + yi\right) = -\frac{1}{4y^2} Y_1\left(\frac{1}{2} + \frac{i}{4y}\right).$$

One consequence of (4.21) is that

$$(4.22) \quad Y_0(i) = \frac{1}{2} Y_1\left(\frac{1}{2} + \frac{i}{2}\right) = 0;$$

namely, that 1 is a critical point of $y \rightarrow f_0(yi)$.

The combined transform of $z \rightarrow w = -\frac{1}{z} \in \Gamma$ and $w \rightarrow -\bar{w}$ maps the line $\frac{1}{2} + yi$ to $\frac{2}{4y^2+1} + \frac{4y}{4y^2+1}i$, the unit circle centered at 1. The invariance of f_1 under this transform yields

$$f_1\left(\frac{1}{2} + yi\right) = f_1\left(\frac{2}{4y^2+1} + \frac{4yi}{4y^2+1}\right).$$

Differentiation with respect to y shows that

$$\begin{aligned} Y_1\left(\frac{1}{2} + yi\right) &= X_1\left(\frac{2}{4y^2+1} + \frac{4yi}{4y^2+1}\right) \frac{\partial}{\partial y} \left(\frac{2}{4y^2+1}\right) \\ &\quad + Y_1\left(\frac{2}{4y^2+1} + \frac{4yi}{4y^2+1}\right) \frac{\partial}{\partial y} \left(\frac{4y}{4y^2+1}\right). \end{aligned}$$

By (3.17)

$$X_1\left(\frac{1}{2} + vi\right) = 0, \quad v > 0.$$

Hence, with $y = \frac{\sqrt{3}}{2}$, one deduces

$$(4.23) \quad Y_0(\sqrt{3}i) = \frac{1}{2}Y_1\left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) = 0,$$

i.e., $\sqrt{3}$ is a critical point of $y \rightarrow f_0(yi)$. By (4.21), $\frac{\sqrt{3}}{3}$ is also a critical point of $y \rightarrow f_0(yi)$.

Now show that

$$(4.24) \quad Y_0(yi) < 0 \text{ if } y \in (\sqrt{3}, \infty).$$

This fact was established by Chen and Oshita in [6]. Here we give a more direct alternative proof.

Consider the expression for $\frac{\partial Y_0(yi)}{\partial y}$ in (4.7). Note that the series

$$(4.25) \quad \sum_{n=1}^{\infty} \frac{n^2(-r)^n}{(1-(-r)^n)^2}$$

is alternating. The only nontrivial property to verify is that the absolute values of the terms decrease, and this follows from the following estimate:

$$\begin{aligned} \frac{n^2 r^n}{(1-(-r)^n)^2} - \frac{(n+1)^2 r^{n+1}}{(1-(-r)^{n+1})^2} &= \frac{(n+1)^2 r^{n+1}}{(1-(-r)^n)^2} \left(\frac{n^2}{(n+1)^2 r} - \frac{(1-(-r)^n)^2}{(1-(-r)^{n+1})^2} \right) \\ &\geq \frac{(n+1)^2 r^{n+1}}{(1-(-r)^n)^2} \left(\frac{e^{\sqrt{3}\pi}}{4} - \frac{(1+e^{-\sqrt{3}\pi})^2}{(1-e^{-2\sqrt{3}\pi})^2} \right) \\ (4.26) \quad &= \frac{(n+1)^2 r^{n+1}}{(1-(-r)^n)^2} \times 56.68 \dots > 0. \end{aligned}$$

This allows us to estimate $\frac{\partial Y_0(yi)}{\partial y}$ as follows:

$$\begin{aligned} \frac{\partial Y_0(yi)}{\partial y} &< -\frac{1}{y^2} + \frac{4\pi^2 e^{-\pi y}}{(1+e^{-\pi y})^2} < -\frac{1}{y^2} + 4\pi^2 e^{-\pi y} \\ &= \frac{1}{y^2} \left(-1 + 4\pi y^2 e^{-\pi y} \right) \\ &\leq \frac{1}{y^2} \left(-1 + 4\pi(\sqrt{3})^2 e^{-\pi\sqrt{3}} \right) \\ (4.27) \quad &\leq \frac{1}{y^2} \times (-0.8388 \dots) < 0. \end{aligned}$$

Here, to reach the fourth line, one notes that

$$(-1 + 4\pi y^2 e^{-\pi y})' = 4\pi e^{-\pi y} y(2 - \pi y) < 0 \quad \text{if } y > \sqrt{3}.$$

Since $Y_0(\sqrt{3}i) = 0$, (4.27) implies (4.24).

By (4.24) and Lemma 4.1, one deduces

$$(4.28) \quad Y_0(yi) > 0 \quad \text{if } y \in \left(0, \frac{\sqrt{3}}{3}\right).$$

Next consider $Y_0(yi)$ for $y \in (1, \sqrt{3})$. By (4.8)

$$(4.29) \quad \frac{\partial^2 Y_0(yi)}{\partial y^2} = \frac{2}{y^3} + \sum_{n=1}^{\infty} \frac{4\pi^3 n^3 ((-r)^n + r^{2n})}{(1 - (-r)^n)^3}, \quad r = e^{-\pi y}.$$

It turns out that the series

$$(4.30) \quad \sum_{n=1}^{\infty} \frac{n^3 (-r)^n}{(1 - (-r)^n)^3}$$

which is part of (4.29) is alternating. To see that the absolute values of the terms in (4.30) decrease, note

$$\frac{n^3 r^n}{(1 - (-r)^n)^3} - \frac{(n+1)^3 r^{n+1}}{(1 - (-r)^{n+1})^3} = \frac{(n+1)^3 r^{n+1}}{(1 - (-r)^n)^3} \left[\frac{n^3}{(n+1)^3 r} - \frac{(1 - (-r)^n)^3}{(1 - (-r)^{n+1})^3} \right],$$

and it suffices to show that the quantity in the brackets is positive. For $y > 1$,

$$\frac{n^3}{(n+1)^3 r} - \frac{(1 - (-r)^n)^3}{(1 - (-r)^{n+1})^3} > \frac{e^\pi}{8} - \frac{(1 + e^{-\pi})^3}{(1 - e^{-2\pi})^3} = 2.8925 \dots - 1.1417 \dots > 0.$$

An upper bound for (4.30) is available if one chooses two terms from the series:

$$(4.31) \quad \sum_{n=1}^{\infty} \frac{n^3 (-r)^n}{(1 - (-r)^n)^3} < \frac{-r}{(1+r)^3} + \frac{8r^2}{(1-r^2)^3}.$$

Then (4.29) becomes

$$\begin{aligned} \frac{\partial^2 Y_0(yi)}{\partial y^2} &< \frac{2}{y^3} - \frac{4\pi^3 r}{(1+r)^3} + \frac{32\pi^3 r^2}{(1-r^2)^3} + 4\pi^3 \sum_{n=1}^{\infty} \frac{n^3 r^{2n}}{(1 - (-r)^n)^3} \\ &< \frac{2}{y^3} - \frac{4\pi^3 r}{(1+r)^3} + \frac{32\pi^3 r^2}{(1-r^2)^3} + \frac{4\pi^3}{(1-r)^3} \sum_{n=1}^{\infty} n^3 r^{2n} \\ &= \frac{2}{y^3} - \frac{4\pi^3 r}{(1+r)^3} + \frac{32\pi^3 r^2}{(1-r^2)^3} + \frac{4\pi^3 r^2 (1 + 4r^2 + r^4)}{(1-r)^3 (1-r^2)^4} \\ &< \frac{2}{y^3} - \left[\frac{4\pi^3}{(1+e^{-\pi})^3} \right] r + \left[\frac{32\pi^3}{(1-e^{-2\pi})^3} + \frac{4\pi^3 (1 + 4e^{-2\pi} + e^{-4\pi})}{(1-e^{-\pi})^3 (1-e^{-2\pi})^4} \right] s^2 \\ &= \frac{2}{y^3} - A_1 r + A_2 r^2 \\ (4.32) \quad &= r \kappa(y), \end{aligned}$$

where we have used the summation formula

$$(4.33) \quad \sum_{n=1}^{\infty} n^3 t^n = t \left(t \left[t \left(\frac{1}{1-t} \right) \right]_t \right)_t = \frac{t(1+4t+t^2)}{(1-t)^4}, \quad |t| < 1,$$

to reach the third line, A_1 and A_2 are given by

$$(4.34) \quad A_1 = \frac{4\pi^3}{(1+e^{-\pi})^3} = 109.24\dots,$$

$$A_2 = \frac{32\pi^3}{(1-e^{-2\pi})^3} + \frac{4\pi^3(1+4e^{-2\pi}+e^{-4\pi})}{(1-e^{-\pi})^3(1-e^{-2\pi})^4} = 1,141.50\dots,$$

and κ is

$$(4.35) \quad \kappa(y) = \frac{2}{y^3 r} - A_1 + A_2 r = \frac{2e^{\pi y}}{y^3} - A_1 + A_2 e^{-\pi y}.$$

Regarding κ , one finds

$$\begin{aligned} \kappa''(y) &= e^{\pi y} (2\pi^2 y^{-3} - 12\pi y^{-4} + 24\pi y^{-5}) + \pi^2 A_2 e^{-\pi y} \\ &= 2e^{\pi y} y^{-5} ((\pi y - 3)^2 + 3) + \pi^2 A_2 e^{-\pi y} > 0 \end{aligned}$$

and

$$\kappa(1) = -13.63\dots < 0, \quad \kappa(\sqrt{3}) = -15.47\dots < 0.$$

Hence

$$(4.36) \quad k(y) < 0, \quad y \in [1, \sqrt{3}],$$

and by (4.32)

$$(4.37) \quad \frac{\partial^2 Y_0(yi)}{\partial y^2} < 0, \quad y \in [1, \sqrt{3}].$$

Since $Y_0(1) = Y_0(\sqrt{3}) = 0$ by (4.22) and (4.23), (4.37) implies

$$(4.38) \quad Y_0(yi) > 0 \quad \text{if } y \in (1, \sqrt{3}).$$

By (4.21), (4.38) implies

$$(4.39) \quad Y_0(yi) < 0 \quad \text{if } y \in \left(\frac{\sqrt{3}}{3}, 1 \right).$$

The lemma follows from (4.28), (4.39), (4.38), and (4.24). $\square$

For b between $(0, 1)$, the next lemma shows that the shape of f_b is similar to f_0 if b is small and similar to f_1 if b is large. The borderline is B given by

$$(4.40) \quad B = \frac{\frac{\partial Y_0(i)}{\partial y}}{\frac{\partial Y_0(i)}{\partial y} - \frac{\partial Y_1(i)}{\partial y}} = \frac{0.2982\dots}{0.2982\dots - (-1.298\dots)} = 0.1867\dots$$

The numerical values in (4.40) are computed from the series (4.3) and (4.7). One interpretation of B is that if $b = B$, the second derivative of $y \rightarrow f_B(yi)$ vanishes at $y = 1$, i.e.,

$$(4.41) \quad \frac{\partial Y_B(i)}{\partial y} = 0.$$

LEMMA 4.4. *The following properties hold for $y \rightarrow f_b(yi)$, $y \in (0, \infty)$.*

1. *When $b \in [0, B)$, the function $y \rightarrow f_b(yi)$, $y > 0$, has exactly three critical points at $\frac{1}{q_b}$, 1, and q_b , where $q_b \in (1, \sqrt{3}]$. Moreover*
 - (a) $Y_b(yi) > 0$ if $y \in (0, \frac{1}{q_b})$,
 - (b) $Y_b(yi) < 0$ if $y \in (\frac{1}{q_b}, 1)$,
 - (c) $Y_b(yi) > 0$ if $y \in (1, q_b)$,
 - (d) $Y_b(yi) < 0$ if $y \in (q_b, \infty)$.*As b increases from 0 to B , q_b decreases from $\sqrt{3}$ towards 1.*
2. *When $b \in [B, 1]$, the function $y \rightarrow f_b(0, y)$, $y > 0$, has only one critical point at 1, and*
 - (a) $Y_b(yi) > 0$ if $y \in (0, 1)$,
 - (b) $Y_b(yi) < 0$ if $y \in (1, \infty)$.

Proof. The shapes of f_1 and f_0 are already established in Lemmas 4.2 and 4.3. These lemmas imply that $y = 1$ is a critical point of $y \rightarrow f_b(yi)$, $y > 0$, i.e.,

$$(4.42) \quad Y_b(i) = 0 \quad \text{for all } b \in [0, 1]$$

and, moreover,

$$(4.43) \quad Y_b(yi) < 0 \quad \text{if } b \in (0, 1] \text{ and } y \geq \sqrt{3}.$$

To study $Y_b(yi)$ for $y \in (1, \sqrt{3})$ and $b \in (0, 1)$, write $Y_b = bY_1 + (1 - b)Y_0$ as

$$(4.44) \quad Y_b(yi) = bY_1(yi) \left(1 + \left(\frac{1-b}{b} \right) \frac{Y_0(yi)}{Y_1(yi)} \right).$$

Recall $Y_1(yi) < 0$ for $y > 1$ from Lemma 4.2. Regarding the quotient $\frac{Y_0(yi)}{Y_1(yi)}$, since $Y_0(i) = Y_1(i) = 0$, $\frac{Y_0(i)}{Y_1(i)}$ is understood as the limit

$$(4.45) \quad \frac{Y_0(i)}{Y_1(i)} = \lim_{y \rightarrow 1} \frac{Y_0(yi)}{Y_1(yi)} = \frac{\frac{\partial Y_0(i)}{\partial y}}{\frac{\partial Y_1(i)}{\partial y}} = \frac{0.2982 \dots}{-1.298 \dots} = -0.2297 \dots < 0$$

evaluated by L'Hôpital's rule. Since $Y_0(\sqrt{3}i) = 0$ by Lemma 4.3,

$$(4.46) \quad \frac{Y_0(\sqrt{3}i)}{Y_1(\sqrt{3}i)} = 0.$$

Lemmas 4.2 and 4.3 also assert that $Y_1(yi) < 0$ and $Y_0(yi) > 0$ if $y \in (1, \sqrt{3})$, so

$$(4.47) \quad \frac{Y_0(yi)}{Y_1(yi)} < 0, \quad y \in (1, \sqrt{3}).$$

However the most important property of this quotient is its monotonicity on $(1, \sqrt{3})$, namely,

$$(4.48) \quad \frac{\partial}{\partial y} \left(\frac{Y_0(yi)}{Y_1(yi)} \right) > 0, \quad y \in (1, \sqrt{3}).$$

The proof of (4.48) is long and brute force. We leave it to the appendix. The first time reader may wish to skip this part.

Return to (4.44). Since $Y_1(yi) < 0$ on $(1, \infty)$ and $\frac{1-b}{b} \in (0, \infty)$ when $b \in (0, 1)$, (4.48) implies that $Y_b(yi)$ can have at most one zero in $(1, \sqrt{3})$ at which $Y_b(yi)$ changes sign. Because of (4.46),

$$(4.49) \quad 1 + \left(\frac{1-b}{b}\right) \frac{Y_0(\sqrt{3}i)}{Y_1(\sqrt{3}i)} = 1 + 0 > 0, \quad b \in (0, 1).$$

Hence $Y_b(yi)$ admits a zero in $(1, \sqrt{3})$ if and only if

$$(4.50) \quad 1 + \left(\frac{1-b}{b}\right) \frac{Y_0(i)}{Y_1(i)} < 0.$$

The condition (4.50) is equivalent to

$$(4.51) \quad b < B$$

by (4.45). We denote this zero in $(1, \sqrt{3})$ of $Y_b(yi)$ by q_b when $b \in (0, B)$. It is also clear from (4.44) and (4.48) that as b increases from 0 to B , q_b decreases monotonically from $\sqrt{3}$ towards 1. This proves parts 1(c), 1(d), and 2(b) of the lemma. The remaining parts follow from Lemma 4.1. $\square$

5. f_b on upper half-plane. We start with a study of the singular point $z = 1$. Recall the set W from (3.10).

LEMMA 5.1.

$$\limsup_{W \ni z \rightarrow 1} X_b(z) = 0.$$

Proof. Note that if $z = x + yi \in W$, then, when $y < 1$,

$$(5.1) \quad 0 < 1 - x < 1 - \sqrt{1 - y^2}.$$

So $W \ni z \rightarrow 1$ is equivalent to that $z \in W$ and $y \rightarrow 0$.

We first show that

$$(5.2) \quad \limsup_{W \ni z \rightarrow 1} X_1(z) \leq 0.$$

Namely, for every $\epsilon > 0$ there exists $\delta > 0$ such that if $z = x + yi \in W$ and $y < \delta$, then $X_b(z) < \epsilon$.

Recall

$$X_1(z) = \sum_{n=1}^{\infty} \frac{8\pi n \sin 2\pi nx}{e^{2\pi ny} + e^{-2\pi ny} - 2 \cos 2\pi nx}$$

from (3.17). Separate this infinite sum into two parts according to whether $ny^2 < \frac{1}{2}$ or $ny^2 \geq \frac{1}{2}$. Write

$$(5.3) \quad a_n(z) = \frac{8\pi n \sin 2\pi nx}{e^{2\pi ny} + e^{-2\pi ny} - 2 \cos 2\pi nx},$$

$$(5.4) \quad A(z) = \sum_{n < 1/(2y^2)} a_n(z),$$

$$(5.5) \quad \tilde{A}(z) = \sum_{n \geq 1/(2y^2)} a_n(z),$$

so that

$$(5.6) \quad X_1(z) = A(z) + \tilde{A}(z).$$

Consider the case $n < \frac{1}{2y^2}$. Since $1 - \sqrt{1 - y^2} < y^2$ when $y \in (0, 1)$, (5.1) implies

$$(5.7) \quad 0 < 1 - x < y^2.$$

Then $0 < 2\pi n(1 - x) < 2\pi ny^2 < \pi$ and $\sin 2\pi nx = -\sin 2\pi n(1 - x) < 0$. Hence, every term $a_n(z)$ in $A(z)$ is negative and, consequently,

$$(5.8) \quad A(z) < 0.$$

When $n \geq \frac{1}{2y^2}$,

$$(5.9) \quad |a_n(z)| \leq \frac{8\pi n}{e^{2\pi ny} - 2} \leq \frac{1}{e^{\pi ny}}.$$

To see the last inequality, note

$$\frac{8\pi n}{e^{2\pi ny} - 2} \leq \frac{8\pi n}{e^{2\pi ny} - 2} (2ny^2) = \frac{16\pi (ny)^2}{e^{2\pi ny} - 1}.$$

There exists $t_0 > 0$ such that for all $t > t_0$, $\frac{16\pi t^2}{e^{2\pi t} - 1} \leq \frac{1}{e^{\pi t}}$. Since $n \geq \frac{1}{2y^2}$, $ny \geq \frac{1}{2y}$. By choosing $y < \frac{1}{2t_0}$, we have $ny > t_0$ and the last inequality of (5.9) follows. Then

$$(5.10) \quad |\tilde{A}(z)| \leq \sum_{n \geq 1/(2y^2)} \frac{1}{e^{\pi ny}} \leq \frac{e^{-\pi y \lceil \frac{1}{2y^2} \rceil}}{1 - e^{-\pi y}} \rightarrow 0 \quad \text{as } y \rightarrow 0,$$

where $\lceil \frac{1}{2y^2} \rceil$ is the integer part of $\frac{1}{2y^2}$. The claim (5.2) now follows from (5.8) and (5.10).

Next we claim that

$$(5.11) \quad \limsup_{W \ni z \rightarrow 1} X_0(z) \leq 0.$$

This is proved by a similar argument whose details are omitted. By (5.2) and (5.11) we obtain that

$$(5.12) \quad \limsup_{W \ni z \rightarrow 1} X_b(z) \leq 0.$$

From the series (3.17) and (3.19),

$$(5.13) \quad X_b(1 + yi) = 0, \quad y > 0.$$

This turns (5.12) into

$$(5.14) \quad \limsup_{W \ni z \rightarrow 1} X_b(z) = 0,$$

proving the lemma. □

Recall that for $b \in [0, B)$, the largest of the three critical points of the function $y \rightarrow f_b(yi)$, $y > 0$, is denoted q_b . This q_b is a maximum and $1 < q_b \leq \sqrt{3}$. By convention if $b \in [B, 1]$, we set $q_b = 1$ which is the unique critical point (a maximum) of $y \rightarrow f_b(yi)$, $y > 0$.

If $b \in [B, 1]$, then $1 - b \in [0, 1 - B]$ and q_{1-b} is defined as above. The transform $z = x + yi \rightarrow w = u + vi$ in (3.23) sends the point $z = q_{1-b}i$ to $w = \frac{q_{1-b}^2 - 1 + 2q_{1-b}i}{q_{1-b}^2 + 1}$. Define

$$(5.15) \quad p_b = \frac{q_{1-b}^2 - 1}{q_{1-b}^2 + 1}.$$

Then

$$(5.16) \quad \frac{q_{1-b}^2 - 1 + 2q_{1-b}i}{q_{1-b}^2 + 1} = p_b + i\sqrt{1 - p_b^2}.$$

LEMMA 5.2. *Let $b \in [0, 1 - B]$ and W be given in (3.10). Then $X_b(z) < 0$ for all $z \in W$.*

Proof. From (3.17) and (3.19) one deduces

$$(5.17) \quad X_b(yi) = 0 \quad \text{and} \quad X_b(1 + yi) = 0, \quad y > 0.$$

Also

$$(5.18) \quad \lim_{y \rightarrow \infty} X_b(z) = 0 \quad \text{uniformly with respect to } x \in \mathbb{R}.$$

On the unit circle, we know from (3.21)

$$(5.19) \quad X_b(x + i\sqrt{1 - x^2}) = \frac{\sqrt{1 - x^2}}{1 - x} Y_{1-b}\left(\frac{\sqrt{1 - x^2}}{1 - x} i\right), \quad x \in (-1, 1).$$

When $b \in [0, 1 - B]$, $1 - b \in [B, 1]$. By Lemma 4.4.2,

$$Y_{1-b}\left(\frac{\sqrt{1 - x^2}}{1 - x} i\right) < 0 \quad \text{if } x \in (0, 1).$$

This shows

$$(5.20) \quad X_b(x + i\sqrt{1 - x^2}) < 0 \quad \text{if } x \in (0, 1).$$

Since X_b is harmonic in W , the lemma follows from (5.17), (5.18), (5.20), and Lemma 5.1 by the maximum principle. $\square$

We are now ready to prove the main theorem.

Proof of Theorem 1.1.

CLAIM 1. Let $b \in [0, 1 - B]$. Then $f_b(z)$ on the upper half-plane is maximized at $q_b i$ and the points in the orbit of $q_b i$ under the group $\mathcal{G}$.

The second plot of Figure 3 demonstrates our argument. By Lemma 3.4.3 it suffices to consider f_b in $\bar{W}_{\mathbb{H}}$. In W Lemma 5.2 asserts that f_b is strictly decreasing in the horizontal direction, so it can only attain a maximum in $\bar{W}_{\mathbb{H}}$ on the part of the

unit circle in the first quadrant, i.e., $\{w \in \mathbb{C} : |w| = 1, 0 < \operatorname{Re} w < 1, \operatorname{Im} w > 0\}$, or on the part of the imaginary axis above i , i.e., $\{z \in \mathbb{C} : \operatorname{Re} z = 0, \operatorname{Im} z \geq 1\}$.

First rule out the unit circle. By Lemma 3.5

$$(5.21) \quad f_b(w) = f_{1-b}(z), \quad z \in \mathbb{H}, \quad w = \frac{z-1}{z+1} \in \mathbb{H}.$$

Take $z = yi$ to be on the imaginary axis. Then

$$w = \frac{y^2 - 1}{y^2 + 1} + \frac{2y}{y^2 + 1}i$$

is on the unit circle. As z moves from i to ∞ upward along the imaginary axis, w moves from i to 1 clockwise along the unit circle. When $b \in [0, 1-B]$, $1-b \in [B, 1]$. Since $y \rightarrow f_{1-b}(yi)$ is strictly decreasing for $y \in (1, \infty)$ by Lemma 4.4.2, $f_b(w)$ is strictly decreasing when w moves from i to 1 clockwise along the unit circle. Then f_b cannot attain a maximum on $\{z \in \mathbb{C} : |w| = 1, 0 < \operatorname{Re} w < 1, \operatorname{Im} w > 0\}$.

Therefore in $\overline{W}_{\mathbb{H}}$, f_b can only achieve a maximum on

$$\{z \in \mathbb{C} : \operatorname{Re} z = 0, \operatorname{Im} z \geq 1\}.$$

By Lemma 4.4.1, it does so at $q_b i$. This proves Claim 1.

By Lemma 4.4.1 and the convention that $q_b = 1$ if $b \in [B, 1]$, three possibilities exist for q_b when $b \in [0, 1-B]$. When $b = 0$, $q_b = \sqrt{3}$, which proves part 1 of the theorem. When $b \in (0, B)$, $q_b \in (1, \sqrt{3})$, which proves part 2 of the theorem. When $b \in [B, 1-B]$, $q_b = 1$, which proves part 3 of the theorem.

Now consider the case $B \in (1-B, 1]$.

CLAIM 2. If $b \in (1-B, 1]$, then f_b on the upper half-plane is maximized at $p_b + i\sqrt{1-p_b^2}$ and the points in its orbit under the group $\mathcal{G}$.

By Lemma 3.5, the duality property, we have

$$f_b(w) = f_{1-b}(z), \quad z \in \mathbb{H}, \quad \text{and} \quad w = \frac{z-1}{z+1} \in \mathbb{H}.$$

If w_* maximizes f_b , then $z_* = \frac{w_*+1}{-w_*+1}$ maximizes f_{1-b} . Since $b \in (1-B, 1]$, $1-b \in [0, B]$. By Claim 1, $z_* = q_{1-b}i$ or a point in the orbit of $q_{1-b}i$ under $\mathcal{G}$. Under the transform $w = \frac{z-1}{z+1}$, $z_* = q_{1-b}i$ corresponds to

$$(5.22) \quad w_* = \frac{q_{1-b}i - 1}{q_{1-b}i + 1} = \frac{q_{1-b}^2 - 1 + 2q_{1-b}i}{q_{1-b}^2 + 1} = p_b + i\sqrt{1-p_b^2}$$

by (5.16). This proves Claim 2.

When $b \in (1-B, 1)$, $q_{1-b} \in (1, \sqrt{3})$ by Lemma 4.4.1. Then by (5.16), $p_b + i\sqrt{1-p_b^2}$, identified as a maximum of f_b in Claim 2, is in $\{z \in \mathbb{C} : |z| = 1, \frac{\pi}{3} < \arg z < \frac{\pi}{2}\}$. This proves part 4 of the theorem. Finally when $b = 1$, $q_0 = \sqrt{3}$ and

$$(5.23) \quad p_1 + i\sqrt{1-p_1^2} = \frac{1 + \sqrt{3}i}{2}$$

by (5.16). This proves part 5 of the theorem. $\square$

Appendix.

Proof of (4.48). Here we prove the monotonicity of the function $y \rightarrow \frac{Y_0(yi)}{Y_1(yi)}$, $y \in (1, \sqrt{3})$, by showing

$$(A.1) \quad \frac{\partial}{\partial y} \left(\frac{Y_0(yi)}{Y_1(yi)} \right) > 0, \quad y \in (1, \sqrt{3}).$$

Our proof uses the following L'Hôpital's rule like criterion for monotonicity. See [2] for more information on this trick.

CLAIM 3.

$$(A.2) \quad \frac{\partial}{\partial y} \left(\frac{Y_0(yi)}{Y_1(yi)} \right) > 0 \text{ on } (1, \sqrt{3}) \text{ if } y \rightarrow \frac{\frac{\partial Y_0(yi)}{\partial y}}{\frac{\partial Y_1(yi)}{\partial y}} \text{ is strictly increasing on } (1, \sqrt{3}).$$

Let $y \in (1, \sqrt{3})$. There exist $y_1 \in (1, y)$ and $y_2 \in (1, y)$ such that

$$\begin{aligned} \frac{\partial}{\partial y} \left(\frac{Y_0(yi)}{Y_1(yi)} \right) &= \frac{\frac{\partial Y_0(yi)}{\partial y} Y_1(yi) - Y_0(yi) \frac{\partial Y_1(yi)}{\partial y}}{Y_1^2(yi)} \\ &= \frac{\frac{\partial Y_1(yi)}{\partial y}}{Y_1(yi)} \left(\frac{\frac{\partial Y_0(yi)}{\partial y}}{\frac{\partial Y_1(yi)}{\partial y}} - \frac{Y_0(yi)}{Y_1(yi)} \right) \\ &= \frac{\frac{\partial Y_1(yi)}{\partial y}}{Y_1(yi) - Y_1(i)} \left(\frac{\frac{\partial Y_0(yi)}{\partial y}}{\frac{\partial Y_1(yi)}{\partial y}} - \frac{Y_0(yi) - Y_0(i)}{Y_1(yi) - Y_1(i)} \right) \\ &= \frac{\frac{\partial Y_1(yi)}{\partial y}}{\frac{\partial Y_1(y_1i)}{\partial y} (y - 1)} \left(\frac{\frac{\partial Y_0(yi)}{\partial y}}{\frac{\partial Y_1(yi)}{\partial y}} - \frac{\frac{\partial Y_0(y_2i)}{\partial y}}{\frac{\partial Y_1(y_2i)}{\partial y}} \right) \end{aligned}$$

since $Y_1(i) = Y_0(i) = 0$. Because $\frac{\partial Y_1(yi)}{\partial y}$ does not change sign in $(1, \sqrt{3})$,

$$(A.3) \quad \frac{\frac{\partial Y_1(yi)}{\partial y}}{\frac{\partial Y_1(y_1i)}{\partial y} (y - 1)} > 0.$$

Moreover, since

$$y \rightarrow \frac{\frac{\partial Y_0(yi)}{\partial y}}{\frac{\partial Y_1(yi)}{\partial y}}$$

is strictly increasing,

$$(A.4) \quad \frac{\frac{\partial Y_0(yi)}{\partial y}}{\frac{\partial Y_1(yi)}{\partial y}} - \frac{\frac{\partial Y_0(y_2i)}{\partial y}}{\frac{\partial Y_1(y_2i)}{\partial y}} > 0.$$

Claim 3 then follows from (A.3) and (A.4).

We proceed to show that

$$(A.5) \quad \frac{\partial}{\partial y} \left(\frac{\frac{\partial Y_0(yi)}{\partial y}}{\frac{\partial Y_1(yi)}{\partial y}} \right) = \frac{\frac{\partial^2 Y_0(yi)}{\partial y^2} \frac{\partial Y_1(yi)}{\partial y} - \frac{\partial Y_0(yi)}{\partial y} \frac{\partial^2 Y_1(yi)}{\partial y^2}}{\left(\frac{\partial Y_1(yi)}{\partial y} \right)^2} > 0, \quad y \in (1, \sqrt{3}).$$

Define

$$(A.6) \quad T(y) = \frac{\partial^2 Y_0(yi)}{\partial y^2} \frac{\partial Y_1(yi)}{\partial y} - \frac{\partial Y_0(yi)}{\partial y} \frac{\partial^2 Y_1(yi)}{\partial y^2}.$$

By (4.3), $\frac{\partial Y_1(yi)}{\partial y} < 0$ on $(0, \infty)$. Therefore to prove (A.5), it suffices to show

$$(A.7) \quad T(y) > 0, \quad y \in (1, \sqrt{3}).$$

We divide $(1, \sqrt{3})$ into two intervals: $(1, \beta)$ and $[\beta, \sqrt{3})$, where $\beta \in (1, \sqrt{3})$ is to be determined. First consider $T(y)$ on $(1, \beta)$. Lemma 4.1 asserts that for $j = 0, 1$,

$$Y_j(yi) = \left(-\frac{1}{y^2}\right) Y_j\left(\frac{i}{y}\right).$$

Differentiation shows that

$$(A.8) \quad \frac{\partial Y_j(yi)}{\partial y} = 2y^{-3} Y_j\left(\frac{i}{y}\right) + y^{-4} \frac{\partial Y_j\left(\frac{i}{y}\right)}{\partial y},$$

$$(A.9) \quad \frac{\partial^2 Y_j(yi)}{\partial y^2} = -6y^{-4} Y_j\left(\frac{i}{y}\right) - 6y^{-5} \frac{\partial Y_j\left(\frac{i}{y}\right)}{\partial y} - y^{-6} \frac{\partial^2 Y_j\left(\frac{i}{y}\right)}{\partial y^2}.$$

Taking $y = 1$ in (A.9) and using $Y_j(i) = 0$, one obtains

$$(A.10) \quad \frac{\partial^2 Y_j(i)}{\partial y^2} = -3 \frac{\partial Y_j(i)}{\partial y}, \quad j = 1, 0.$$

In particular (A.10) implies that

$$(A.11) \quad T(1) = 0.$$

Next consider the derivative of T ,

$$(A.12) \quad T'(y) = \frac{\partial^3 Y_0(yi)}{\partial y^3} \frac{\partial Y_1(yi)}{\partial y} - \frac{\partial Y_0(yi)}{\partial y} \frac{\partial^3 Y_1(yi)}{\partial y^3}.$$

It is clear from (4.3) and (4.5) that

$$(A.13) \quad \frac{\partial Y_1(yi)}{\partial y} < 0, \quad \frac{\partial^3 Y_1(yi)}{\partial y^3} < 0, \quad y > 0.$$

Similarly to the argument following (4.30), one finds the series in (4.7) to be alternating when $y > 1$. Therefore,

$$(A.14) \quad \frac{\partial Y_0(yi)}{\partial y} > -\frac{1}{y^2} + \frac{4\pi^2 e^{-\pi y}}{(1 + e^{-\pi y})^2} - \frac{16\pi^2 e^{-2\pi y}}{(1 - e^{-2\pi y})^2}.$$

We will later choose $\beta \in (1, \sqrt{3})$ so that when $y = \beta$, the right side of (A.14) is positive, namely, choose β to make

$$(A.15) \quad -\frac{1}{\beta^2} + \frac{4\pi^2 e^{-\pi\beta}}{(1 + e^{-\pi\beta})^2} - \frac{16\pi^2 e^{-2\pi\beta}}{(1 - e^{-2\pi\beta})^2} > 0.$$

Since

$$(A.16) \quad \frac{\partial^2 Y_0(yi)}{\partial y^2} < 0, \quad y \in (1, \sqrt{3})$$

by (4.37), the condition (A.15) implies that

$$(A.17) \quad \frac{\partial Y_0(yi)}{\partial y} > 0, \quad y \in (1, \beta).$$

Regarding $\frac{\partial^3 Y_0(yi)}{\partial y^3}$, write it as

$$(A.18) \quad \frac{\partial^3 Y_0(yi)}{\partial y^3} = -\frac{6}{y^4} - 4\pi^4 \sum_{n=1}^{\infty} \frac{n^4 (-r)^n}{(1 - (-r)^n)^4} - 16\pi^4 \sum_{n=1}^{\infty} \frac{n^4 r^{2n}}{(1 - (-r)^n)^4} - 4\pi^4 \sum_{n=1}^{\infty} \frac{n^4 (-r)^{3n}}{(1 - (-r)^n)^4},$$

where

$$(A.19) \quad r = e^{-\pi y}$$

as before. Clearly, the second series in (A.18) is positive for all $y > 0$. One can also show as before that when $y > 1$, the first and the third series in (A.18) are both alternating. Pick three leading terms from the first series and one term from each of the second and the third series to form an upper bound:

$$(A.20) \quad \begin{aligned} \frac{1}{4\pi^4} \frac{\partial^3 Y_0(yi)}{\partial y^3} &< -\frac{6}{4\pi^4 y^4} + \frac{r}{(1+r)^4} - \frac{16r^2}{(1-r^2)^4} + \frac{81r^3}{(1+r^3)^4} - \frac{4r^2}{(1+r)^4} + \frac{r^3}{(1+r)^4} \\ &< -\frac{6}{4\pi^4 y^4} + \frac{r}{(1+r)^4} - \frac{16r^2}{(1-r^2)^4} + 82r^3 - \frac{4r^2}{(1+r)^4} \\ &= -\frac{6}{4\pi^4 y^4} + \frac{r}{(1-r^2)^4} \\ &\quad \cdot \left(1 - 24r + 104r^2 - 28r^3 - 311r^4 - 4r^5 + 497r^6 - 328r^8 + 82r^{10}\right) \\ &< -\frac{6}{4\pi^4 y^4} + \frac{r}{(1-r^2)^4} (1 - 24r + 104r^2) \\ &\leq -\frac{6}{4\pi^4 y^4} + \frac{1}{(1-e^{-2\pi})^4} r(1 - 24r + 104r^2). \end{aligned}$$

Denote the last line by

$$(A.21) \quad \sigma(y) = -\frac{6}{4\pi^4 y^4} + \frac{1}{(1-e^{-2\pi})^4} (e^{-\pi y} - 24e^{-2\pi y} + 104e^{-3\pi y}).$$

Compute

$$(A.22) \quad \sigma'(y) = \frac{24}{4\pi^4 y^5} + \frac{\pi e^{-\pi y}}{(1-e^{-2\pi})^4} (-1 + 48e^{-\pi y} - 312e^{-2\pi y})$$

and consider the quantity in parentheses,

$$(A.23) \quad \phi(y) = -1 + 48e^{-\pi y} - 312e^{-2\pi y}.$$

Since $\phi'(y) = \pi e^{-\pi y}(-48 + 624e^{-\pi y}) < 0$ if $y > 1$,

$$(A.24) \quad \phi(y) > \phi(\beta) \quad \text{if } y \in (1, \beta).$$

If one can make $\phi(\beta) > 0$, namely, if

$$(A.25) \quad -1 + 48e^{-\pi\beta} - 312e^{-2\pi\beta} > 0,$$

then

$$(A.26) \quad \phi(y) > 0, \quad y \in (1, \beta).$$

Consequently, by (A.22)

$$(A.27) \quad \sigma'(y) > 0, \quad y \in (1, \beta),$$

and

$$(A.28) \quad \sigma(y) < \sigma(\beta), \quad y \in (1, \beta).$$

This shows that

$$(A.29) \quad \frac{\partial^3 Y_0(yi)}{\partial y^3} \leq 4\pi^4 \sigma(\beta), \quad y \in (1, \beta).$$

If one can choose β so that $\sigma(\beta) < 0$, namely,

$$(A.30) \quad -\frac{6}{4\pi^4 \beta^4} + \frac{1}{(1 - e^{-2\pi})^4} (e^{-\pi\beta} - 24e^{-2\pi\beta} + 104e^{-3\pi\beta}) < 0,$$

then

$$(A.31) \quad \frac{\partial^3 Y_0(yi)}{\partial y^3} < 0, \quad y \in (1, \beta).$$

Following (A.13), (A.17), and (A.31), one has that

$$(A.32) \quad T'(y) > 0, \quad y \in (1, \beta).$$

By (A.11), (A.32) implies that

$$(A.33) \quad T(y) > 0, \quad y \in (1, \beta)$$

provided (A.15), (A.25), and (A.30) hold.

Next consider $T(y)$ for $y \in [\beta, \infty)$. Introduce

$$(A.34) \quad d(y) = Y_0(yi) - Y_1(yi) = \frac{\pi}{6} - \sum_{k=1}^{\infty} \frac{4\pi(2k-1)r^{2k-1}}{1 + r^{2k-1}},$$

$$(A.35) \quad d'(y) = \sum_{k=1}^{\infty} \frac{4\pi^2(2k-1)^2 r^{2k-1}}{(1 + r^{2k-1})^2},$$

$$(A.36) \quad d''(y) = \sum_{k=1}^{\infty} \frac{4\pi^3(2k-1)^3 (-r^{2k-1} + r^{2(2k-1)})}{(1 + r^{2k-1})^3}.$$

Then by (4.3), (4.4), (A.35), and (A.36),

$$\begin{aligned} T(y) &= d''(y) \frac{\partial Y_1(yi)}{\partial y} - d'(y) \frac{\partial^2 Y_1(yi)}{\partial y^2} \\ &= \sum_{k=1}^{\infty} \frac{4\pi^2(2k-1)^2 r^{2k-1}}{y^2(1+r^{2k-1})^2} \left(\frac{\pi(2k-1)(1-r^{2k-1})}{1+r^{2k-1}} - \frac{2}{y} \right) \\ (A.37) \quad &+ \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} \frac{16\pi^5(2k-1)^2(2n)^2 r^{2n+2k-1}}{(1+r^{2k-1})^2(1-r^{2n})^2} \left(\frac{(2k-1)(1-r^{2k-1})}{1+r^{2k-1}} - \frac{2n(1+r^{2n})}{1-r^{2n}} \right) \end{aligned}$$

$$(A.38) \quad = \sum_{k=1}^{\infty} c_k + \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} d_{kn},$$

where c_k and d_{kn} are defined by the terms in (A.37).

Regarding c_k , because, with $y > 1$,

$$\frac{\pi(2k-1)(1-r^{2k-1})}{1+r^{2k-1}} - \frac{2}{y} \geq \frac{\pi(1-r)}{1+r} - 2 > \frac{\pi(1-e^{-\pi})}{1+e^{-\pi}} - 2 > 0,$$

one has

$$(A.39) \quad c_k > 0 \quad \text{for all } k.$$

The terms d_{kn} have the following property:

$$(A.40) \quad d_{kn} > 0 \quad \text{if } k > n, \quad d_{kn} < 0 \quad \text{if } k \leq n.$$

To see (A.40), define

$$(A.41) \quad \rho_j(r) = \frac{j(1+(-r)^j)}{1-(-r)^j}$$

so that the quantity in the second parenthesis pair of (A.37) is $\rho_{2k-1}(r) - \rho_{2n}(r)$. The claim (A.40) follows if $\rho_j(r)$ is increasing with respect to j . To this end consider

$$(A.42) \quad \rho_{j+1}(r) - \rho_j(r) = \frac{1 - (2j+1)(r+1)(-r)^j + r^{2j+1}}{(1-(-r)^{j+1})(1-(-r)^j)}.$$

This is clearly positive when j is odd since $0 < r < 1$. When j is even, denote the numerator in (A.42) by

$$\mu_j(r) = 1 - (2j+1)(r+1)r^j + r^{2j+1}.$$

Then

$$\mu'_j(r) = (2j+1)r^{j-1}(-j - (j+1)r + r^{j+1}).$$

As $0 < r < 1$,

$$-j - (j+1)r + r^{j+1} < -j - (j+1)r + r = -j - jr < 0.$$

Hence $\mu'_j(r) < 0$ for $r \in (0, 1)$. Moreover

$$\mu_j(r) > 1 - 2(2j+1)r^j,$$

so

$$\mu_j(e^{-\pi}) > 1 - 2(2j+1)e^{-\pi j} > 0$$

for all $j \geq 1$. Therefore $\mu_j(r) > 0$ since $r \in (0, e^{-\pi})$ and (A.40) is proved.

By (A.40) drop the positive d_{kn} 's to bound the double sum in (A.34) from below by

$$(A.43) \quad \sum_{l=1}^{\infty} \sum_{n=1}^{\infty} d_{kn} > \sum_{k=1}^{\infty} d_{kk} + \sum_{k=1}^{\infty} \sum_{n=k+1}^{\infty} d_{kn}.$$

First consider $\sum_{k=1}^{\infty} d_{kk}$:

$$(A.44) \quad d_{kk} = \frac{16\pi^5(2k-1)^2(2k)^2r^{4k-1}}{(1+r^{2k-1})^2(1-r^{2k})^2} \left(\frac{(2k-1)(1-r^{2k-1})}{1+r^{2k-1}} - \frac{2k(1+r^{2k})}{1-r^{2k}} \right).$$

For $y > 1$,

$$\begin{aligned} (1+r^{2k-1})(1-r^{2k}) &= 1+r^{2k-1}(1-r-r^{2k}) \\ &\geq 1+r^{2k-1}(1-r-r^2) \\ &> 1+r^{2k-1}(1-e^{-\pi}-e^{-2\pi}) \\ &= 1+r^{2k-1} \times 0.9549 \dots > 1. \end{aligned}$$

Also, when $y > 1$, both $\frac{(4k-2)r^{2k-1}}{1+r^{2k-1}}$ and $\frac{4kr^{2k}}{1-r^{2k}}$ are decreasing with respect to k . Hence

$$\begin{aligned} \frac{(2k-1)(1-r^{2k-1})}{1+r^{2k-1}} - \frac{2k(1+r^{2k})}{1-r^{2k}} &= -1 - \frac{(4k-2)r^{2k-1}}{1+r^{2k-1}} - \frac{4kr^{2k}}{1-r^{2k}} \\ &\geq -1 - \frac{2r}{1+r} - \frac{4r^2}{1-r^2} \\ &> -1 - \frac{2e^{-\pi}}{1+e^{-\pi}} - \frac{4e^{-2\pi}}{1-e^{-2\pi}} \\ &= -\left(\frac{1+e^{-\pi}}{1-e^{-\pi}} \right). \end{aligned}$$

One estimates

$$\begin{aligned} \sum_{k=1}^{\infty} d_{kk} &> -16\pi^5 \left(\frac{1+e^{-\pi}}{1-e^{-\pi}} \right) \sum_{k=1}^{\infty} (2k-1)^2(2k)^2r^{4k-1} \\ (A.45) \quad &= -64\pi^5 \left(\frac{1+e^{-\pi}}{1-e^{-\pi}} \right) \frac{r^3(1+31r^4+55r^8+9r^{12})}{(1-r^4)^5} \end{aligned}$$

with the help of the summation formula

$$\begin{aligned} (A.46) \quad \sum_{k=1}^{\infty} (2k)^2(2k-1)^2r^{4k-1} &= \frac{r^2}{16} \left(r \left(r^{-1} \left(r \left(\frac{1}{1-r^4} \right)_r \right)_r \right)_r \right)_r \\ &= \frac{4r^3(1+31r^4+55r^8+9r^{12})}{(1-r^4)^5}. \end{aligned}$$

Next consider the double sum on the right of (A.43). Dropping the first term in the second parenthesis pair of (A.37) one obtains

$$(A.47) \quad \sum_{k=1}^{\infty} \sum_{n=k+1}^{\infty} d_{kn} > \sum_{k=1}^{\infty} \sum_{n=k+1}^{\infty} \frac{16\pi^5(2k-1)^2(2n)^2 r^{2n+2k-1}}{(1+r^{2k-1})^2(1-r^{2n})^2} \left(-\frac{2n(1+r^{2n})}{1-r^{2n}} \right).$$

For $n \geq k+1$,

$$\begin{aligned} (1+r^{2k-1})(1-r^{2n}) &= 1+r^{2k-1}(1-r^{2n-2k+1}-r^{2n}) \\ &> 1+r^{2k+2} \geq 1+r^{2n}. \end{aligned}$$

Consequently

$$\begin{aligned} \frac{1+r^{2n}}{(1+r^{2k-1})^2(1-r^{2n})^3} &< \frac{1}{(1+r^{2k-1})(1-r^{2n})^2} \\ &= \frac{1}{1+r^{2k-1}((1-r^{2n})^2+r^{2n-2k+1}(-2+r^{2n}))} \\ &\leq \frac{1}{1+r^{2k-1}((1-r^2)^2-2r^3)} < 1. \end{aligned}$$

Return to (A.47) to deduce

$$\begin{aligned} \sum_{k=1}^{\infty} \sum_{n=k+1}^{\infty} d_{kn} &> - \sum_{k=1}^{\infty} \sum_{n=k+1}^{\infty} 16\pi^5(2k-1)^2(2n)^3 r^{2n+2k-1} \\ &\geq - \sum_{k=1}^{\infty} \sum_{n=2}^{\infty} 128\pi^5(2k-1)^2 n^3 r^{2n+2k-1} \\ &= - \sum_{k=1}^{\infty} 128\pi^5(2k-1)^2 r^{2k} \left(\frac{r^3(8-5r^2+4r^4-r^6)}{(1-r^2)^4} \right) \\ &\geq - \sum_{k=1}^{\infty} 128\pi^5(2k-1)^2 r^{2k+3} \left(\frac{8}{(1-e^{-2\pi})^4} \right) \\ (A.48) \quad &= - \frac{1024\pi^5}{(1-e^{-2\pi})^4} \left(\frac{r^5(1+6r^2+r^4)}{(1-r^2)^3} \right). \end{aligned}$$

We have used the summation formulas

$$(A.49) \quad \sum_{n=2}^{\infty} n^3 r^{2n-1} = \frac{1}{8} \left(r \left(r \left(\frac{1}{1-r^2} \right)_r \right)_r \right)_r - r = \frac{r^3(8-5r^2+4r^4-r^6)}{(1-r^2)^4},$$

$$(A.50) \quad \sum_{k=1}^{\infty} (2k-1)^2 r^{2k+3} = r^5 \left(r \left(\frac{r}{1-r^2} \right)_r \right)_r = \frac{r^5(1+6r^2+r^4)}{(1-r^2)^3}$$

to reach the third line and the last line, respectively.

By (A.38), (A.39), (A.43), (A.45), and (A.48), taking two terms from $\sum_{k=1}^{\infty} c_k$ and using $1 < y < \sqrt{3}$, we find

$$\begin{aligned}
 T(y) &\geq \frac{4\pi^2 r}{y^2(1+r)^2} \left(\frac{\pi(1-r)}{1+r} - \frac{2}{y} \right) + \frac{36\pi^2 r^3}{y^2(1+r^3)^2} \left(\frac{3\pi(1-r^3)}{1+r^3} - \frac{2}{y} \right) \\
 &\quad - 64\pi^5 \left(\frac{1+e^{-\pi}}{1-e^{-\pi}} \right) \frac{r^3(1+31r^4+55r^8+9r^{12})}{(1-r^4)^5} - \frac{1024\pi^5}{(1-e^{-2\pi})^4} \left(\frac{r^5(1+6r^2+r^4)}{(1-r^2)^3} \right) \\
 &> \frac{4\pi^2 r}{y^2(1+r)^2} \left(\frac{\pi(1-r)}{1+r} - \frac{2}{y} \right) + \frac{36\pi^2 r^3}{3(1+e^{-3\pi})^2} \left(\frac{3\pi(1-e^{-3\pi})}{1+e^{-3\pi}} - 2 \right) \\
 &\quad - 64\pi^5 \left(\frac{1+e^{-\pi}}{1-e^{-\pi}} \right) \frac{r^3(1+31e^{-4\pi}+55e^{-8\pi}+9e^{-12\pi})}{(1-e^{-4\pi})^5} \\
 &\quad - \frac{1024\pi^5}{(1-e^{-2\pi})^4} \left(\frac{r^3 e^{-2\pi}(1+6e^{-2\pi}+e^{-4\pi})}{(1-e^{-2\pi})^3} \right) \\
 (A.51) \quad &= \frac{4\pi^2 r}{y^2(1+r)^2} \left(\frac{\pi(1-r)}{1+r} - \frac{2}{y} \right) + Ar^3,
 \end{aligned}$$

where

$$\begin{aligned}
 A &= \frac{36\pi^2}{3(1+e^{-3\pi})^2} \left(\frac{3\pi(1-e^{-3\pi})}{1+e^{-3\pi}} - 2 \right) \\
 &\quad - 64\pi^5 \left(\frac{1+e^{-\pi}}{1-e^{-\pi}} \right) \frac{(1+31e^{-4\pi}+55e^{-8\pi}+9e^{-12\pi})}{(1-e^{-4\pi})^5} \\
 &\quad - \frac{1024\pi^5}{(1-e^{-2\pi})^4} \left(\frac{e^{-2\pi}(1+6e^{-2\pi}+e^{-4\pi})}{(1-e^{-2\pi})^3} \right) \\
 (A.52) \quad &= -21,077.61 \dots
 \end{aligned}$$

Continuing from (A.51), one has

$$(A.53) \quad T(y) > \frac{\pi^2 r^2}{(1+r)^3} \left(\left(\frac{4\pi}{y^2} - \frac{8}{y^3} \right) r^{-1} - \left(\frac{4\pi}{y^2} + \frac{8}{y^3} \right) + \frac{Ar(1+r)^3}{\pi^2} \right).$$

Bound the last term by

$$(A.54) \quad \frac{Ar(1+r)^3}{\pi^2} \geq \frac{Ae^{-\pi\beta}(1+e^{-\pi\beta})^3}{\pi^2}$$

and define

$$(A.55) \quad \nu(y) = \left(\frac{4\pi}{y^2} - \frac{8}{y^3} \right) e^{\pi y} - \left(\frac{4\pi}{y^2} + \frac{8}{y^3} \right) + \frac{Ae^{-\pi\beta}(1+e^{-\pi\beta})^3}{\pi^2}$$

so that

$$(A.56) \quad T(y) > \frac{\pi^2 r^2}{(1+r)^3} \nu(y).$$

Regarding $\nu(y)$, one finds

$$(A.57) \quad \nu'(y) = e^{\pi y} y^{-3} \left(4\pi^2 \left(\sqrt{y} - \frac{2}{2\sqrt{y}} \right)^2 + \frac{8}{y} \right) + \frac{8\pi}{y} + \frac{24}{y^4} > 0.$$

Then (A.56) implies

$$(A.58) \quad T(y) > \frac{\pi^2 r^2}{(1+r)^3} \nu(\beta),$$

where

$$(A.59) \quad \nu(\beta) = \left(\frac{4\pi}{\beta^2} - \frac{8}{\beta^3} \right) e^{\pi\beta} - \left(\frac{4\pi}{\beta^2} + \frac{8}{\beta^3} \right) + \frac{Ae^{-\pi\beta}(1+e^{-\pi\beta})^3}{\pi^2}.$$

Therefore if we can find β so that $\nu(\beta) > 0$, namely,

$$(A.60) \quad \left(\frac{4\pi}{\beta^2} - \frac{8}{\beta^3} \right) e^{\pi\beta} - \left(\frac{4\pi}{\beta^2} + \frac{8}{\beta^3} \right) + \frac{Ae^{-\pi\beta}(1+e^{-\pi\beta})^3}{\pi^2} > 0,$$

then

$$(A.61) \quad T(y) > 0, \quad y \in [\beta, \sqrt{3}].$$

In summary, to invoke (A.33) and (A.61) one must choose β so that (A.15), (A.25), (A.30), (A.60) all hold. Our choice is

$$(A.62) \quad \beta = 1.08$$

at last. One readily checks the four conditions:

$$(A.63) \quad -\frac{1}{\beta^2} + \frac{4\pi^2 e^{-\pi\beta}}{(1+e^{-\pi\beta})^2} - \frac{16\pi^2 e^{-2\pi\beta}}{(1-e^{-2\pi\beta})^2} \Big|_{\beta=1.08} = 0.2058 \dots > 0;$$

$$(A.64) \quad -1 + 48e^{-\pi\beta} - 312e^{-2\pi\beta} \Big|_{\beta=1.08} = 0.2608 \dots > 0;$$

$$(A.65) \quad -\frac{6}{4\pi^4 \beta^4} + \frac{1}{(1-e^{-2\pi})^4} \left(e^{-\pi\beta} - 24e^{-2\pi\beta} + 104e^{-3\pi\beta} \right) \Big|_{\beta=1.08} = -0.0007930 \dots < 0;$$

$$(A.66) \quad \left(\frac{4\pi}{\beta^2} - \frac{8}{\beta^3} \right) e^{\pi\beta} - \left(\frac{4\pi}{\beta^2} + \frac{8}{\beta^3} \right) + \frac{Ae^{-\pi\beta}(1+e^{-\pi\beta})^3}{\pi^2} \Big|_{\beta=1.08} = 35.20 \dots > 0.$$

The proof of (4.48) is complete. $\square$

REFERENCES

- [1] G. ALBERTI, R. CHOKSI, AND F. OTTO, *Uniform energy distribution for an isoperimetric problem with long-range interactions*, J. Amer. Math. Soc., 22 (2009), pp. 569–605.
- [2] G. ANDERSON, M. VAMANAMURTHY, AND M. VUORINEN, *Monotonicity rules in calculus*, Amer. Math. Monthly, 113 (2006), pp. 805–816.
- [3] T. M. APOSTOL, *Modular Functions and Dirichlet Series in Number Theory*, Springer, Berlin, 1976.
- [4] E. BENDITO, A. CARMONA, A. M. ENCINAS, J. M. GESTO, A. GÓMEZ, C. MOURIÑO, AND M. T. SÁNCHEZ, *Computational cost of the Fekete problem I: The forces method on the 2-sphere*, J. Comput. Phys., 228 (2009), pp. 3288–3306.
- [5] L. BÉTERMIN, L. DE LUCA, AND M. PETRACHE, *Crystallization to the Square Lattice for a Two Body Potential*, preprint, <https://arxiv.org/abs/1907.06105>, 2019.
- [6] X. CHEN AND Y. OSHITA, *An application of the modular function in nonlocal variational problems*, Arch. Ration. Mech. Anal., 186 (2007), pp. 109–132.

- [7] R. CHOKSI AND M. A. PELETIER, *Small volume fraction limit of the diblock copolymer problem: I. Sharp interface functional*, SIAM J. Math. Anal., 42 (2010), pp. 1334–1370.
- [8] H. COHN AND A. KUMAR, *Universally optimal distribution of points on spheres*, J. Amer. Math. Soc., 20 (2006), pp. 99–148.
- [9] H. COHN, A. KUMAR, S. MILLER, D. RADCHENKO, AND M. VIAZOVSKA, *Universal Optimality of the E_8 and Leech Lattices and Interpolation Formulas*, preprint, <https://arxiv.org/abs/1902.05438>, 2019.
- [10] R. EVANS, *A fundamental region for Hecke's modular group*, J. Number Theory, 5 (1973), pp. 108–115.
- [11] D. GOLDMAN, C. B. MURATOV, AND S. SERFATY, *The gamma-limit of the two-dimensional Ohta-Kawasaki energy. I. Droplet density*, Arch. Ration. Mech. Anal., 210 (2013), pp. 581–613.
- [12] T. C. HALES, *The honeycomb conjecture*, Discrete Comput. Geom., 25 (2001), pp. 1–22.
- [13] M. MORINI AND P. STERNBERG, *Cascade of minimizers for a nonlocal isoperimetric problem in thin domains*, SIAM J. Math. Anal., 46 (2014), pp. 2033–2051.
- [14] H. NAKAZAWA AND T. OHTA, *Microphase separation of ABC-type triblock copolymers*, Macromolecules, 26 (1993), pp. 5503–5511.
- [15] T. OHTA AND K. KAWASAKI, *Equilibrium morphology of block copolymer melts*, Macromolecules, 19 (1986), pp. 2621–2632.
- [16] M. PETRACHE AND S. SERFATY, *Crystallization for Coulomb and Riesz interactions as a consequence of the Cohn-Kumar conjecture*, Proc. Amer. Math. Soc., to appear.
- [17] X. REN AND C. WANG, *A stationary core-shell assembly in a ternary inhibitory system*, Discrete Contin. Dyn. Syst., 37 (2017), pp. 983–1012.
- [18] X. REN AND C. WANG, *Stationary disc assemblies in a ternary system with long range interaction*, Commun. Contemp. Math., 21 (2019), 1850046.
- [19] X. REN AND J. WEI, *On the multiplicity of solutions of two nonlocal variational problems*, SIAM J. Math. Anal., 31 (2000), pp. 909–924.
- [20] X. REN AND J. WEI, *Triblock copolymer theory: Free energy, disordered phase and weak segregation*, Phys. D, 178 (2003), pp. 103–117.
- [21] X. REN AND J. WEI, *Many droplet pattern in the cylindrical phase of diblock copolymer morphology*, Rev. Math. Phys., 19 (2007), pp. 879–921.
- [22] X. REN AND J. WEI, *A double bubble assembly as a new phase of a ternary inhibitory system*, Arch. Ration. Mech. Anal., 215 (2015), pp. 967–1034.
- [23] E. SANDIER AND S. SERFATY, *From the Ginzburg-Landau model to vortex lattice problems*, Comm. Math. Phys., 313 (2012), pp. 635–743.
- [24] F. THEIL, *A proof of crystallization in two dimensions*, Comm. Math. Phys., 262 (2006), pp. 209–236.
- [25] C. WANG, X. REN, AND Y. ZHAO, *Bubble assemblies in ternary systems with long range interaction*, Comm. Math. Sci., 17 (2019), pp. 2309–2324.