

Redox-controlled chalcogen-bonding at tellurium: Impact on Lewis acidity and chloride anion transport properties

Benyu Zhou and François P. Gabbaï^{*,a}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Abstract: Our interests in the chemistry of atypical main group Lewis acids have led us to devise strategies that augment the affinity of chalcogen-bond donors for anionic guests. In this study, we describe the oxidative methylation of diaryltellurides as one such strategy along with its application to the synthesis of $[\text{Mes}(\text{C}_6\text{F}_5)_2\text{TeMe}]^+$ and $[(\text{C}_6\text{F}_5)_2\text{TeMe}]^+$ starting from $\text{Mes}(\text{C}_6\text{F}_5)_2\text{Te}$ and $(\text{C}_6\text{F}_5)_2\text{Te}$, respectively. These new telluronium cations have been evaluated for their ability to complex and transport chloride anions across phospholipid bilayers. These studies show that, when compared to their neutral $\text{Te}(\text{II})$ precursors, these $\text{Te}(\text{IV})$ cations display both higher Lewis acidity and transport activity. The positive attributes of these telluronium cations, which originate from a lowering of the tellurium-centered σ^* orbitals and a deepening of the associated σ -holes, demonstrate that the redox state of the main group element provides a convenient handle over its chalcogen-bonding properties.

The field of chalcogen (Ch) bonding is in rapid expansion owing to applications in supramolecular chemistry, catalysis, and anion transport.^{1–11} The formation of chalcogen-bonds, which are defined as secondary bonding interactions between a donor (D) and a chalcogenide (ChR_2 , R = substituent),¹² can be rationalized based on the presence of regions of positive electrostatic potential on the chalcogen atom. These regions often referred to as σ -holes, sit at the termini of Ch-R polar bonds and spatially coincide with the largest lobe of the Ch-R σ^* orbitals (Figure 1). Investigations into the origin of chalcogen-bonds point to the interplay of electrostatic, dispersion, and orbital delocalization effects.^{13, 14} The dispersion term is accentuated in the case of softer, inherently polarizable, chalcogens, while the electrostatic term can be controlled by adjusting the polarity of Ch-R σ -bond. The orbital delocalization term corresponding to the $\text{D} \rightarrow \text{ChR}_2$ interaction responds favourably to a lowering of the Ch-R σ^* orbital energy, as observed with heavy chalcogens. An analogous dependence is seen with respect to the polarity of the Ch-R bond which, when

increased, leads to higher chalcogen parentage of the σ^* orbital. All three bonding terms culminate at tellurium, the softest, heaviest, and most electropositive chalcogen.^{2, 5, 15, 16}

$(\text{C}_6\text{F}_5)_2\text{Te}$ is a particularly noteworthy molecule that embodies the points discussed above. The chalcogen-bond donor properties of this derivative manifest in the short intermolecular Te-F contacts found in the crystal¹⁷ as well as in the formation of adducts with donors such as tetramethylthiourea.¹⁸ Matile brought $(\text{C}_6\text{F}_5)_2\text{Te}$ back into the spotlight by demonstrating its use in anion binding catalysis¹⁹ and chloride anion transport.²⁰ The chalcogen-bond donor properties of $(\text{C}_6\text{F}_5)_2\text{Te}$ can be correlated to the electron-withdrawing properties of the pentafluorophenyl substituents, which deepen the σ -hole and stabilize the Te-C σ^* orbital.





Figure 1. Left: Orbital and electrostatic origin of the Lewis acidity in tellurium derivatives. The intensification of the blue color illustrates an energy lowering in the case of the LUMO and a greater positive charge development at tellurium in the case of the σ -hole.

Motivated by these recent developments, we are charting new strategies to predictably enhance the chalcogen-bond donor properties of tellurium compounds, with the ultimate aim of reaching *bona fide* Lewis acidity. Based on the knowledge that telluronium cations form partially covalent halide complexes^{21–23} and encouraged by recent contributions on the effect of oxidation on the Lewis acidity of phosphorus,^{24–26} antimony,^{27–29} sulfur,^{30–32} and iodine species,^{33, 34} we have now decided to probe the effect of oxidative alkylation of simple diaryltellurides as a way to stabilize the accepting Te-C

^a Department of Chemistry, Texas A&M University, College Station, Texas 77843-3255, United States of America

[†] Electronic Supplementary Information (ESI) available: Experimental and computational details. NMR spectra of novel compounds. Anion binding and transport data. Crystallographic data in CIF format. See DOI: 10.1039/x0xx00000x

σ^* orbital while at the same time deepening the coincident σ -hole as illustrated in Figure 1.




Scheme 1. Synthesis of the telluronium salts investigated in this study.

Considering stability, ease of synthesis, and synthetic modularity, we decided to target a series of telluronium cations obtained by direct methylation of the diaryltellurides shown in Scheme 1. While (C₆F₅)₂Te and Mes₂Te are known,^{17, 35} we were able to obtain Mes(C₆F₅)Te (**1**) by the reaction of a C₆F₅MgBr solution with a solution of MesTeBr generated *in situ* by the combination of bromine and dimesitylditelluride. This new telluride has been characterized by traditional means, including multinuclear NMR spectroscopy and elemental analysis. These neutral tellurides were next converted into the corresponding methyltelluronium species. While the more electron-rich derivatives Mes₂Te and **1** could be efficiently methylated using MeI/AgBF₄ in CH₂Cl₂, methylation of (C₆F₅)₂Te only proceeded with Me₃O⁺BF₄[−] upon heating to 90°C in a mixture of 1,2-dichloroethane and toluene (vol. 1:2). The harsher conditions used for this last reaction reflects the electron-deficiency of the tellurium centre induced by the two pentafluorophenyl groups. The three new telluronium cations [Mes₂TeMe]⁺ (**[2]**⁺), [Mes(C₆F₅)TeMe]⁺ (**[3]**⁺), and [(C₆F₅)₂TeMe]⁺ (**[4]**⁺) have been

isolated as their tetrafluoroborate salts and fully characterized. All three salts give well-resolved ¹H and ¹⁹F-NMR spectra. Installation of the methyl group gives rise to a singlet in the ¹H-NMR spectra of these derivatives. This singlet, which appears in the 3.17–3.45 ppm window for all three salts, is flanked by ¹²⁵Te satellites separated by ²J_{Te-H} = 29.1–31.2 Hz. These coupling constants are consistent with those observed for known methyltelluronium species.³⁶ The ¹²⁵Te-NMR resonance of the telluronium cations appears at 563.7 ppm for **[2]**⁺, 637.4 ppm for **[3]**⁺, and 598.1 for **[4]**⁺, a set of values that are distinctly downfield from those of the telluride precursors (258.7 ppm for Mes₂Te, 280.0 ppm for **1**, and 287.1 ppm for (C₆F₅)₂Te).

The structures of these new telluronium salts have also been confirmed by X-ray crystallography (Figure 2).³⁷ In the solid-state, short Te–F contacts between the tellurium atom and the tetrafluoroborate counter anion are observed. The shortest contact, which is much smaller than the sum of the van der Waals radii of Te and F ($\Sigma_{\text{vdw}}(\text{Te-F}) = 3.45 \text{ \AA}$),³⁸ is observed in the case of **[4]**[BF₄] which displays a Te---F distance of 2.6902(11) Å (Figure 2). This distance is approximately 0.3 Å shorter than in **[2]**[BF₄] (3.045(3) Å) and 0.1 Å shorter than in **[3]**[BF₄] (2.7908(15) Å), attesting to the higher Lewis acidity of the tellurium centre in the bis(pentafluorophenyl) cation **[4]**⁺. The value of the C11–Te–F6 angle in **[3]**[BF₄] (168.56°) and C8–Te–F11 angle in **[4]**[BF₄] (167.14°) shows that the fluorine atom involved in the short contact is positioned *trans* from the pentafluorophenyl ring. This position corresponds to that defined by the lowest-lying σ^* orbital or deepest σ -hole in these two salts, underscoring the importance of the molecular design principles presented in Figure 1. The higher Lewis acidity of the tellurium atom in **[4]**⁺ is further reflected by the presence of a second Te---F short contact of 2.814 Å involving the tetrafluoroborate anion of a neighboring, symmetry equivalent unit. The resulting C2–Te–F12' angle of 171.86° shows that this second contact is along a direction *trans* from the second pentafluorophenyl substituent of **[4]**⁺. It follows that **[4]**⁺ could






Figure 3. Energy level and pictorial representation of LUMO and LUMO+1 orbitals of the telluronium cations $[2]^+$, $[3]^+$, $[4]^+$.

,in principle, support interactions with two anionic guests.

The telluronium cations have also been studied by Density Functional Theory (M062X functional; mixed basis set: 6-31G(d')/6-31G+(d')/cc-pVTZ-PP). To better compare the electrostatic potential about the tellurium centre, ESP maps were plotted on the same scale (Figure 2). All cations feature pronounced σ -holes which can be conveniently characterized by their corresponding $V_{s,max}$ values. The $V_{s,max}$ values of the two σ -holes are almost identical to one another in the case of $[2]^+$ ($V_{s,max} = 103.1$ and 104.0 kcal/mol) and $[4]^+$ ($V_{s,max} = 136.0$ and 138.3 kcal/mol) which feature two identical aryl substituents. In the case of $[3]^+$, the $V_{s,max}$ values of the two σ -holes (116.3 and 122.3 kcal/mol) show a larger differentiation, with the largest value corresponding to the hole trans from the electron-withdrawing pentafluorophenyl substituent. The electron-withdrawing properties of this substituent are also responsible for the significant deepening of the σ -holes (or increase in $V_{s,max}$) upon transitioning from $[2]^+$ to $[4]^+$. This trend is paralleled by the decrease in the energies of the lowest unoccupied molecular orbitals observed upon sequential

replacement of the mesityl substituents by pentafluorophenyl substituents. This trend is illustrated in Figure 3. The largest energy drop is observed upon the introduction of the first pentafluorophenyl group in $[3]^+$ whose LUMO energy is 1.00 eV lower than that of $[2]^+$. The substitution of the second mesityl group induces a further stabilization of the LUMO by 0.75 eV, making $[4]^+$ the most electron-deficient cation of this series. The LUMO+1 shows a similar energy ordering within this series of compounds and lay only 0.24 eV above the LUMO in the case of $[2]^+$ and $[4]^+$ and 0.6 eV in the case of $[3]^+$. The LUMO and LUMO+1 of these compounds bear strong parentage from the $\sigma^*(\text{Te}-\text{C})$ orbitals, thereby defining the site of expected maximum Lewis acidity in these compounds. The larger LUMO-LUMO+1 separation in $[3]^+$ illustrates the differing electronegativity of the pentafluorophenyl and mesityl substituents in this compound. In $[4]^+$, the presence of two closely spaced, low lying vacant orbitals indicates the presence of two accessible Lewis acidic sites, a conclusion supported by the observation of two short $\text{Te} \cdots \text{F}(\text{BF}_4^-)$ contacts in the structure of $[4][\text{BF}_4]$.

Returning to the main objective of this study which is to establish the benefits of oxidation of the tellurium center on the chalcogen-bond donor properties of these derivatives, the ESP maps of $[2]^+$, $[3]^+$, and $[4]^+$ can be compared to those of their neutral precursors (Figure 3). In all cases, we observe that the $V_{s,max}$ values in the cations are significantly higher than those of the neutral precursor, indicating that the cationic systems will indeed be better chalcogen-bond donors. Comparison of the LUMO energies provides a consistent picture, with those of the neutral derivatives Mes_2Te , $\text{Mes}(\text{C}_6\text{F}_5)\text{Te}$, and $(\text{C}_6\text{F}_5)_2\text{Te}$ being >4 eV higher than those of $[2]^+$, $[3]^+$, and $[4]^+$ (see SI for details). Because electrostatic effects are unscreened and thus magnified in the gas-phase, we have also investigated the effect of solvation on the electronic property differences observed between the telluronium cations and their neutral precursors. The results obtained with the SMD solvation model and water as a solvent are presented in Figure 4 for **1** and $[3]^+$. These





Figure 4. (a) Changes observed in the energy of the LUMO and ESP map upon oxidative methylation of **1** into $[3]^+$ ($V_{s,max}$ unit: kcal/mol). (b, c) ^{19}F NMR spectra recorded upon addition of TBACl to a MeCN solution of **1** (b) and $[3]\text{BF}_4$ (c). Cl^- equiv. ($\text{eq}(\text{Cl}^-)$) used for in the titrations: $\text{eq}(\text{Cl}^-) = 0, 0.2, 0.4, 0.6, 0.8, 1.0, 1.4, 1.8, 2.2, 2.6, 3.0, 3.8, 4.6, 5.4, 6.2, 7.0, 9.8$ for spectra a-q recorded with **1** (panel (b)); $\text{eq}(\text{Cl}^-) = 0, 0.18, 0.36, 0.54, 0.72, 0.90, 1.08, 1.25, 1.44, 1.62, 1.79, 2.15, 2.51, 2.87, 3.23$ for spectra a-o recorded with $[3]\text{BF}_4$ (panel (c)).

results show that solvation dampens the extent of LUMO stabilization upon conversion of **1** into **[3]⁺**.

The above computational results predict that the telluronium cations should be more Lewis acidic than their neutral, divalent precursors. To verify this prediction, we became eager to compare the anion binding properties of the most electron deficient cations **[3]⁺** and **[4]⁺** to those of their neutral precursors Mes(C₆F₅)Te (**1**) and (C₆F₅)₂Te, respectively. We first chose to evaluate the chloride anion affinity of **[3]⁺** and **[4]⁺** by carrying out ¹⁹F NMR titrations in MeCN as illustrated in Figure 4. These experiments reveal that **[3]⁺** binds one chloride anion with an association constant of 1.2×10^4 ($\pm 6 \times 10^3$) M⁻¹. The situation presented by **[4]⁺** is more complicated in that it displays two binding events (Scheme 2, see SI for additional details). The first binding event is essentially quantitative, suggesting an association constant $K_1 \geq 10^5$. This finding is consistent with the partially covalent nature of electron-deficient triorganyltellurium(IV) chloride derivatives such as (C₆F₅)₃TeCl.²² The first binding event is followed by a second one, which we interpret as the coordination of a second chloride anion at the remaining σ hole with an association constant $K_2 = 260$ (± 80) M⁻¹ (Scheme 2). The smaller value of K_2 against K_1 is in agreement with the expected negative cooperativity between these two Lewis acidic sites. The chloride anion affinities displayed by **[3]⁺** and **[4]⁺** are respectively larger than those measured for their neutral precursors **1** and (C₆F₅)₂Te, in the same solvent. Indeed, while **1** showed no measurable chloride anion affinity in MeCN, a low binding constant of 36 M⁻¹ was measured for (C₆F₅)₂Te. The drastically increased chloride anion affinity of the cationic derivatives highlights the benefits of the approach illustrated in Figure 1. Last but not least, we note that (C₆F₅)₂Te had been previously evaluated in THF, affording a binding constant of 2,000 M⁻¹. The lower binding constant measured in MeCN shows the negative influence of a polar solvent such as MeCN that can saturate the tellurium Lewis acid while also more effectively solvating the chloride anion and the receptor.



Scheme 2. Chloride anion binding by **[4]⁺** as inferred from ¹⁹F NMR spectroscopy.

Given our interest in the development of Lewis acidic main group cations^{39, 40} as anion transporters,^{41–47} we decided to conclude this study by comparing the chloride anion transport properties of the new telluronium cations **[2]⁺**, **[3]⁺**, and **[4]⁺** with those of their neutral precursors. The main objective of these investigations was to assess the impact of increased Lewis acidity or chalcogen-bond donor properties on the transport activity of these derivatives. As a prelude to these studies, we tested the stability of **[2]⁺**, **[3]⁺**, and **[4]⁺** in water/DMSO (9:1) and did not observe the decomposition of the compounds (see SI). Next, we prepared large unilamellar vesicles (LUVs) using




Figure 5. Left: Simplified graphical summary of the chloride anion efflux experiment using EYPC LUVs vesicles in the presence of valinomycin (0.1 mol% with respect to lipid concentration). The medium was buffered using HEPES (10mM). Transporters were added as a solution in DMSO (7 μ L) to reach a final concentration of 2 mol% with respect to lipid concentration. Right: Chloride efflux data recorded with an ion selective electrode. The transporter is added 30 s after additional of valinomycin.

EYPC (egg-yolk phosphatidylcholine) as a phospholipid and a concentrated KCl solution as a cargo (Figure 5). These freshly prepared LUVs were subjected to transporter-mediated chloride anion efflux experiments monitored with an ion-selective electrode.⁴⁸ In all cases, anion transport was coupled with the addition of valinomycin as a potassium ion transporter. We first investigated the neutral derivatives with the intent of comparing the activity of Mes₂Te and **1** to that of (C₆F₅)₂Te, a derivative reported to be a potent chloride anion transporter.²⁰ While (C₆F₅)₂Te indeed mediated fast chloride efflux, the activity of Mes₂Te and **1** could not be discerned from the background efflux induced by the addition of DMSO alone (Figure 5). The superiority of (C₆F₅)₂Te is assigned to the presence of two electron-withdrawing groups and their impact on the chalcogen-bond donor properties of the tellurium centre. It is also possible that the steric bulk of the mesityl group of Mes₂Te and **1** has a detrimental influence, especially in the case of **1**. After evaluating these neutral derivatives, we turned our attention to the telluronium cations. While **[2]⁺** showed no activity, **[3]⁺** promoted a very rapid release of the chloride anion cargo, reaching almost 100% after 4.5 minutes. The contrasting behaviour of **[2]⁺** and **[3]⁺** underscores the importance of the role played by the pentafluorophenyl group. At the same time, the greater activity of **[3]⁺** when compared to its neutral precursor **1** illustrates the benefit that results from the oxidative methylation of the tellurium center. A comparison of the initial chloride efflux rate k_{ini} obtained for (C₆F₅)₂Te (0.40 %·s⁻¹) and **[3]⁺** (0.75 %·s⁻¹) indicate that the former is distinctly more active. The superiority of **[3]⁺** is further illustrated by a comparison of the EC₅₀ values obtained from a Hill analysis, which shows that the value obtained for **[3]⁺** (0.020%) is ~five times lower than that of **1** (1.1 mol%). Finally, we also evaluated the more Lewis acidic cation **[4]⁺**. We observed that it outperforms **[3]⁺** as well as its neutral precursor, as indicated by the accelerated initial chloride efflux rate ($k_{ini} = 1.68$ %·s⁻¹) and a low EC₅₀ value of 0.13%. This last result suggests a positive correlation between the Lewis acidity and the transport activity of these tellurium derivatives. A similar correlation was established in a prior study dealing with stibonium cations as transporters.³⁹ The higher transport activity of **[3]⁺** and **[4]⁺** is

notably higher than that of $(\text{C}_6\text{F}_5)_2\text{Te}$ but not disproportionately so. It is possible that the high Lewis acidity of these two cationic compounds impedes chloride anion release, thereby dampening their transport activity.^{49,50} Experiments carried out with $[3]^+$ and $[4]^+$ with chloride-loaded DPPC (DPPC = 1,2-dipalmitoyl-*sn*-glycero-3-phosphatidylcholine) vesicles show a temperature dependence consistent with a carrier-type mechanism (see SI), as observed for other transporters based on main group cations.^{39,40} Finally, $[3]^+$ and $[4]^+$ do not induce significant leakage when administered to vesicles loaded with the self-quenching dye 5(6)-carboxyfluorescein. This last result indicates that these cationic derivatives do not act as surfactants and leave the phospholipid membrane unaffected.

In summary, this work defines the oxidative alkylation of diorganotellurides as a simple strategy for enhancing the chalcogen-bond donor properties of the tellurium center. This enhancement correlates with a lowering of the Te-C σ^* orbitals and the presence of deeper σ -holes in the resulting telluronium cations. As indicated by chloride anion binding studies, the enhanced chalcogen-bond donor properties of these derivatives lead to the emergence of classical Lewis acidity. A positive outcome is also observed in the context of transmembrane chloride anion transport, with the telluronium cations displaying a higher activity than their neutral telluride precursors. We propose that this strategy may open the door to redox switchable chalcogen-bonding/group 16-centered Lewis acidity, a phenomenon that could be exploited for the design of anion transporters that respond to changes in the redox state of the medium.³¹ Finally, we note that the concept introduced in this paper is related to previously published strategies relying on the oxidation of a remote main group center as a means to enhance the activity of anion transporters.^{51,52}

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgements

Support from the Welch Foundation (Grant A-1423), the National Science Foundation (CHE-1856453), Texas A&M University (Arthur E. Martell Chair of Chemistry), and the Laboratory for Molecular Simulation at Texas A&M University (software and computational resources) is gratefully acknowledged.

References

- N. Sudha and H. B. Singh, *Coord. Chem. Rev.*, 1994, **135**, 469-515.
- A. F. Cozzolino, P. J. W. Elder and I. Vargas-Baca, *Coord. Chem. Rev.*, 2011, **255**, 1426-1438.
- M. Breugst, D. von der Heiden and J. Schmauck, *Synthesis*, 2017, **49**, 3224-3236.
- K. T. Mahmudov, M. N. Kopylovich, M. F. C. Guedes da Silva and A. J. L. Pombeiro, *Dalton Trans.*, 2017, **46**, 10121-10138.
- R. Gleiter, G. Haberhauer, D. B. Werz, F. Rominger and C. Bleiholder, *Chem. Rev.*, 2018, **118**, 2010-2041.
- K. Selvakumar and H. B. Singh, *Chem. Sci.*, 2018, **9**, 7027-7042.
- J. Y. C. Lim and P. D. Beer, *Chem*, 2018, **4**, 731-783.
- J. Bamberger, F. Ostler and O. G. Mancheño, *ChemCatChem*, 2019, **11**, 5198-5211.
- L. Vogel, P. Wönnner and S. M. Huber, *Angew. Chem. Int. Ed.*, 2019, **58**, 1880-1891.
- K. Strakova, L. Assies, A. Goujon, F. Piazzolla, H. V. Humeniuk and S. Matile, *Chem. Rev.*, 2019, **119**, 10977-11005.
- N. Biot and D. Bonifazi, *Coord. Chem. Rev.*, 2020, **413**, 213243.
- N. W. Alcock, Secondary Bonding to Nonmetallic Elements in *Advances in Inorganic Chemistry and Radiochemistry*, eds. H. J. Emeléus and A. G. Sharpe, Academic Press, 1972, vol. 15, pp. 1-58.
- C. Bleiholder, D. B. Werz, H. Köppel and R. Gleiter, *J. Am. Chem. Soc.*, 2006, **128**, 2666-2674.
- D. J. Pascoe, K. B. Ling and S. L. Cockcroft, *J. Am. Chem. Soc.*, 2017, **139**, 15160-15167.
- P. Wönnner, T. Steinke, L. Vogel and S. M. Huber, *Chem. Eur. J.*, 2020, **26**, 1258-1262.
- P. Wönnner, A. Dreger, L. Vogel, E. Engelage and S. M. Huber, *Angew. Chem. Int. Ed.*, 2019, **58**, 16923-16927.
- T. M. Klapötke, B. Krumm, P. Mayer, K. Polborn and O. P. Ruscitti, *Inorg. Chem.*, 2001, **40**, 5169-5176.
- S. Aboulkacem, D. Naumann, W. Tyrra and I. Pantenburg, *Organometallics*, 2012, **31**, 1559-1565.
- S. Benz, A. I. Poblador-Bahamonde, N. Low-Ders and S. Matile, *Angew. Chem. Int. Ed.*, 2018, **57**, 5408-5412.
- L. M. Lee, M. Tsemperouli, A. I. Poblador-Bahamonde, S. Benz, N. Sakai, K. Sugihara and S. Matile, *J. Am. Chem. Soc.*, 2019, **141**, 810-814.
- M. J. Collins, J. A. Ripmeester and J. F. Sawyer, *J. Am. Chem. Soc.*, 1987, **109**, 4113-4115.
- T. M. Klapötke, B. Krumm, P. Mayer, K. Polborn and O. P. Ruscitti, *J. Fluorine Chem.*, 2001, **112**, 207-212.
- H. Y. Zhao and F. P. Gabbai, *Nat. Chem.*, 2010, **2**, 984-990.
- J. M. Bayne and D. W. Stephan, *Chem. Soc. Rev.*, 2016, **45**, 765-774.
- M. H. Holthausen, M. Mehta and D. W. Stephan, *Angew. Chem. Int. Ed.*, 2014, **53**, 6538-6541.
- R. J. Andrews, S. S. Chitnis and D. W. Stephan, *Chem. Commun.*, 2019, **55**, 5599-5602.
- D. Tofan and F. P. Gabbai, *Chem. Sci.*, 2016, **7**, 6768-6778.
- M. Yang, D. Tofan, C.-H. Chen, K. M. Jack and F. P. Gabbai, *Angew. Chem. Int. Ed.*, 2018, **57**, 13868-13872.
- D. You and F. P. Gabbai, *Trends in Chemistry*, 2019, **1**, 485-496.
- Y. Kim, H. Zhao and F. P. Gabbai, *Angew. Chem. Int. Ed.*, 2009, **48**, 4957-4960.
- H. Zhao and F. P. Gabbai, *Org. Lett.*, 2011, **13**, 1444-1446.
- F. A. Tsao, A. E. Waked, L. Cao, J. Hofmann, L. Liu, S. Grimme and D. W. Stephan, *Chem. Commun.*, 2016, **52**, 12418-12421.
- F. Heinen, E. Engelage, A. Dreger, R. Weiss and S. M. Huber, *Angew. Chem. Int. Ed.*, 2018, **57**, 3830-3833.
- T. L. Seidl and D. R. Stuart, *J. Org. Chem.*, 2017, **82**, 11765-11771.
- M. Oba, Y. Okada, M. Endo, K. Tanaka, K. Nishiyama, S. Shimada and W. Ando, *Inorg. Chem.*, 2010, **49**, 10680-10686.
- N. S. Dance, W. R. McWhinnie, J. Mallaki and Z. Monsef-Mirzai, *J. Organomet. Chem.*, 1980, **198**, 131-143.
- CCDC 2004978-2004980 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre

- via www.ccdc.cam.ac.uk/data_request/cif. The chiral telluronium salt **[3]**[BF₄] crystallizes in the centrosymmetrical *P*2₁/*n* space group, with the two enantiomers in the unit cell.
38. S. Alvarez, *Dalton Trans.*, 2013, **42**, 8617-8636.
 39. G. Park, D. J. Brock, J.-P. Pellois and F. P. Gabbaï, *Chem*, 2019, **5**, 2215-2227.
 40. G. Park and F. P. Gabbaï, *Angew. Chem. Int. Ed.*, 2020, **59**, 5298-5302.
 41. A. P. Davis, D. N. Sheppard and B. D. Smith, *Chem. Soc. Rev.*, 2007, **36**, 348-357.
 42. J. T. Davis, O. Okunola and R. Quesada, *Chem. Soc. Rev.*, 2010, **39**, 3843-3862.
 43. S. Matile, A. Vargas Jentzsch, J. Montenegro and A. Fin, *Chem. Soc. Rev.*, 2011, **40**, 2453-2474.
 44. P. A. Gale, R. Perez-Tomas and R. Quesada, *Acc. Chem. Res.*, 2013, **46**, 2801-2813.
 45. A. Vargas Jentzsch, A. Hennig, J. Mareda and S. Matile, *Acc. Chem. Res.*, 2013, **46**, 2791-2800.
 46. P. A. Gale, J. T. Davis and R. Quesada, *Chem. Soc. Rev.*, 2017, **46**, 2497-2519.
 47. X. Wu, E. N. W. Howe and P. A. Gale, *Acc. Chem. Res.*, 2018, **51**, 1870-1879.
 48. S. Matile, N. Sakai and A. Hennig, Transport Experiments in Membranes in *Supramol. Chem.*, eds. P. A. Gale and J. W. Steed, Wiley, Chichester, U.K., 2012, vol. 2, p. 473.
 49. J. P. Behr, M. Kirch and J. M. Lehn, *J. Am. Chem. Soc.*, 1985, **107**, 241-246.
 50. A. Vargas Jentzsch, D. Emery, J. Mareda, P. Metrangolo, G. Resnati and S. Matile, *Angew. Chem. Int. Ed.*, 2011, **50**, 11675-11678.
 51. S. Benz, M. Macchione, Q. Verolet, J. Mareda, N. Sakai and S. Matile, *J. Am. Chem. Soc.*, 2016, **138**, 9093-9096.
 52. A. J. Plajer, J. Zhu, P. Proehm, A. D. Bond, U. F. Keyser and D. S. Wright, *J. Am. Chem. Soc.*, 2019, **141**, 8807-8815.