

ARNOLD-THOM GRADIENT CONJECTURE FOR THE ARRIVAL TIME

TOBIAS HOLCK COLDING AND WILLIAM P. MINICOZZI II

ABSTRACT. We prove conjectures of René Thom and Vladimir Arnold for C^2 solutions to the degenerate elliptic equation that is the level set equation for motion by mean curvature.

We believe these results are the first instances of a general principle: Solutions of many degenerate equations behave as if they are analytic, even when they are not. If so, this would explain various conjectured phenomena.

0. INTRODUCTION

By a classical result, solutions of analytic elliptic PDEs, like the Laplace equation, are analytic. Many important equations are degenerate elliptic and solutions have much lower regularity. Still, one may hope that solutions share properties of analytic functions. On the surface, such properties seem to be purely analytic; however, they turn out to be closely connected to important open problems in geometry.

For an analytic function, Łojasiewicz, [L1], proved that any gradient flow line with a limit point has finite length and, thus, limits to a unique critical point. This result has since been known as *Łojasiewicz's theorem*. The proof relied on two *Łojasiewicz inequalities* for analytic functions that had also been used to prove two conjectures around 1960: Laurent Schwarz's division conjecture in 1959 in [L3] and a conjecture of Whitney about singularities in 1963 in [L4]. Around the same time, in 1958, Hörmander proved a special case of Schwarz's division conjecture by establishing Łojasiewicz's first inequality for polynomials, [Hö].




Figure illustrates in $\mathbf{R}^3$ a situation conjectured to be impossible. The Arnold-Thom conjecture asserts that a blue integral curve does not spiral as it approaches the critical set (illustrated in red, orthogonal to the plane where the curve spirals).

The authors were partially supported by NSF Grants DMS 1404540, DMS 1206827 and DMS 1707270.

Around 1972, Thom, [T], [L2], [Ku], [A], [G], conjectured a strengthening of Lojasiewicz's theorem, asserting that each gradient flow line of an analytic function approaches its limit from a unique limiting direction:

Conjecture 0.1. If a gradient flow line $x(t)$ for an analytic function has a limit point, then the limit of secants $\lim_{t \rightarrow \infty} \frac{x(t) - x_\infty}{|x(t) - x_\infty|}$ exists.

This conjecture arose in Thom's work on catastrophe theory and singularity theory and became known as *Thom's gradient conjecture*. The conjecture was finally proven in 2000 by Kurdyka, Mostowski, and Parusinski in [KMP], but the following stronger conjecture remains open (see page 282 in Arnold's problem list, [A]):

Conjecture 0.2. If a gradient flow line $x(t)$ for an analytic function has a limit point, then the limit of the unit tangents $\frac{x'(t)}{|x'(t)|}$ exists.

It is easy to see that if $\lim_{t \rightarrow \infty} \frac{x'(t)}{|x'(t)|}$ exists, then so does $\lim_{t \rightarrow \infty} \frac{x(t) - x_\infty}{|x(t) - x_\infty|}$. It follows that the *Arnold-Thom conjecture* 0.2 implies Thom's gradient conjecture 0.1. Easy examples show that the Lojasiewicz theorem, the Lojasiewicz inequalities, and both Conjectures 0.1 and 0.2 fail for general smooth functions; see, e.g., fig. 3.5 in [Si] or fig. 1 in [CM8].

Analytic functions play an important role in differential equations since solutions of analytic elliptic equations are themselves analytic. In many instances, the properties that come from being analytic are more important than analyticity itself. We will show that solutions of an important degenerate elliptic equation have analytic properties even though solutions are not even C^3 . Namely, we will show that Conjectures 0.1, 0.2 hold for solutions of the classical degenerate elliptic equation, known as the *arrival time equation*,

$$(0.3) \quad -1 = |\nabla u| \operatorname{div} \left(\frac{\nabla u}{|\nabla u|} \right).$$

Here u is defined on a compact connected subset of $\mathbf{R}^{n+1}$ with smooth mean convex boundary. Equation (0.3) is the prototype for a family of equations, see, e.g., [OsSe], used for tracking moving interfaces in complex situations. These equations have been instrumental in applications, including semiconductor processing, fluid mechanics, medical imaging, computer graphics, and material sciences.

Even though solutions of (0.3) are a priori only in the viscosity sense, they are always twice differentiable by [CM5], though not necessarily C^2 ; see [CM6], [H2], [I], [KS]. Even when a solution is C^2 , it still might not be C^3 , Sesum, [S], let alone analytic as in Lojasiewicz's theorem. However, solutions behave like analytic functions are expected to:

Theorem 0.4. The Arnold-Thom conjecture holds for C^2 solutions of (0.3).

The geometric meaning of (0.3) is that the level sets $u^{-1}(t)$ are mean convex and evolve by mean curvature flow. One says that u is the *arrival time* since $u(x)$ is the time the hypersurfaces $u^{-1}(t)$ arrive at x under the mean curvature flow; see Chen-Giga-Goto, [ChGG], Evans-Spruck, [ES], Osher-Sethian, [OsSe], and [CM3]. Geometrically, singular points for the flow correspond to critical points for u .

We conjecture that even for solutions that are not C^2 , but merely twice differentiable, the Arnold-Thom conjecture holds:

Conjecture 0.5. Lojasiewicz's inequalities and the Arnold-Thom conjecture hold for all solutions of (0.3).

If this conjecture holds, then the gradient Lojasiewicz inequality would imply that the flow is singular at only finitely many times as has been conjectured, [W3], [AAG], [Wa], [M].

One of the important ingredients in the proof of Theorem 0.4 is an essentially sharp rate of convergence for the rescaled mean curvature flow; this will be given in Proposition 2.4 below. This rate is not fast enough to directly show the convergence of unit tangents, which is closely related to the existence of a non-integrable kernel of the linearized operator. However, we overcome this by a careful analysis of this kernel.

We believe that the principle that solutions of degenerate equations behave as though they are analytic, even when they are not, should be quite general. For instance, there should be versions for other flows, including Ricci flow; cf. [CM9].

1. LOJASIEWICZ THEOREM FOR THE ARRIVAL TIME

A function v satisfies a gradient Lojasiewicz inequality near a point y (see, e.g., [CM8]) if there exist $p > 1$, C and a neighborhood of y (all depending on v and y) so that

$$(1.1) \quad |v - v(y)| \leq C |\nabla v|^p.$$

This is nontrivial only if y is a critical point. If $\nabla v(y) = 0$ and v satisfies (1.1), then $v(y)$ is the only critical value in this neighborhood (this applies for any $p > 0$).

In this section, we show (1.1) with $p = 2$ for a C^2 solution u of (0.3). When u is not C^2 , then (1.1) can fail for any fixed $p > 1$. Namely, for any odd integer $m \geq 3$, Angenent and Velázquez construct rotationally symmetric examples in [AV] where $|u - u(y)| \approx |\nabla u|^{\frac{m}{m-1}}$ for a sequence of points tending to y . The examples in [AV] were constructed to analyze so-called type II singularities that were previously observed by Hamilton and proven rigorously to exist by Altschuler-Angenent-Giga, [AAG]; cf. also [GK].

From now on, u will be C^2 . To prove (1.1), we first recall the properties that we will use. Namely, if $S = \{x \mid \nabla u(x) = 0\}$ denotes the critical set,¹ then [CM5] and [CM6] give:

- (S1) S is a closed embedded connected k -dimensional C^1 submanifold whose tangent space is the kernel of Hess_u . Moreover, S lies in the interior of the region where u is defined.
- (S2) If $q \in S$, then $\text{Hess}_u(q) = -\frac{1}{n-k} \Pi$ and $\Delta u(q) = -\frac{n+1-k}{n-k}$, where Π is orthogonal projection onto the orthogonal complement of the kernel.

After subtracting a constant, we can assume that $\sup u = 0$.

Using these properties, the next theorem gives the gradient Lojasiewicz inequality.

Theorem 1.2. We have that $u(S) = 0$ and

$$(1.3) \quad \frac{|\nabla u|^2}{-u} \rightarrow \frac{2}{n-k} \text{ as } u \rightarrow 0.$$

In particular, there exists $C > 0$ so that $C^{-1} |\nabla u|^2 \leq -u \leq C |\nabla u|^2$.

¹The flow is smooth away from the singular set S consisting of cylindrical singularities; see, [W1], [W2], [H1], [HS1], [HS2], [HaK], [An]; cf. [B], [CM1]. See also [CM4].

Proof. The boundary of the domain is smooth and mean convex, so $\nabla u \neq 0$ on the boundary. The normalization $\sup u = 0$ implies that $u = 0$ at any maximum. Thus, there is at least one point in S with $u = 0$. By (S1), u is constant on S and we conclude that $u(S) = 0$.

Given $\epsilon > 0$, choose $\delta > 0$ so that $|p - q| < \delta$ implies that $|u_{ij}(p) - u_{ij}(q)| < \epsilon$ and, moreover, so that the δ -tubular neighborhood of S does not intersect the boundary of the domain. Let q be any point with $\text{dist}(q, S) < \delta$ and then let p be a point in the compact set S that minimizes the distance to q (note that p might not be unique). Since S is C^1 , the minimizing property implies that the vector $q - p$ is orthogonal to the tangent space to S . In particular, (S2) implies that

$$(1.4) \quad \text{Hess}_u(p)(q - p) = -\frac{q - p}{n - k}.$$

Given $t \in (0, 1]$, the fundamental theorem of calculus gives

$$(1.5) \quad \nabla u(p + t(q - p)) = \int_0^t \text{Hess}_u(p + s(q - p))(q - p) ds.$$

Combining this with (1.4) and the continuity of the Hessian gives

$$(1.6) \quad \left| \nabla u(p + t(q - p)) + t \frac{q - p}{n - k} \right| \leq \epsilon t |q - p|.$$

Using this at $t = 1$ gives

$$(1.7) \quad \left| \nabla u(q) + \frac{q - p}{n - k} \right| \leq \epsilon |q - p|.$$

Using the fundamental theorem of calculus on u this time, (1.6) gives that

$$(1.8) \quad \left| u(q) + \frac{|p - q|^2}{2(n - k)} \right| \leq \int_0^1 \left| \left\langle \nabla u(p + t(q - p)) + t \frac{q - p}{n - k}, q - p \right\rangle \right| dt \leq \frac{\epsilon}{2} |p - q|^2.$$

Since $\epsilon > 0$ is arbitrary, combining the last two inequalities gives (1.3).

The last claim follows from (1.3) since $\{u = 0\} = \{|\nabla u| = 0\} = S$. $\square$

The next theorem shows that the gradient flow lines of u have finite length (this is the Lojasiewicz theorem for u), converge to points in S , and approach S orthogonally. The first claims follow immediately from the gradient Lojasiewicz inequality of Theorem 1.2. Let Π_{axis} denote orthogonal projection onto the kernel of Hess_u .

Theorem 1.9. Each flow line γ for ∇u has finite length and limits to a point in S . Moreover, if we parametrize γ by $s \geq 0$ with $|\gamma_s| = 1$ and $\gamma(0) \in S$, then

$$(1.10) \quad u(\gamma(s)) \approx \frac{-s^2}{2(n - k)},$$

$$(1.11) \quad |\nabla u(\gamma(s))|^2 \approx \frac{s^2}{(n - k)^2},$$

$$(1.12) \quad \Pi_{\text{axis}}(\gamma_s) \rightarrow 0.$$

In particular, for s small, we have that $\gamma(s) \subset B_{2n\sqrt{-u(\gamma(s))}}(\gamma(0))$.

Proof. Each point lies on a flow line where u is increasing and limits to 0, so γ limits to S . If we parametrize γ by time t (so that $u \circ \gamma(t) = t$ and $|\gamma_t| = \frac{1}{|\nabla u|}$), then the length is

$$(1.13) \quad \int_T^0 \frac{1}{|\nabla u|} dt \approx \sqrt{\frac{n-k}{2}} \int_T^0 \frac{1}{\sqrt{-u}} dt = \sqrt{\frac{n-k}{2}} \int_T^0 \frac{1}{\sqrt{-t}} dt = \sqrt{2(k-n)T},$$

where the approximation used (1.3). In particular, the flow lines starting from $u = T$ have finite length approximately equal to $\sqrt{2(k-n)T}$. It follows that γ has a limit $\gamma(0) \in S$ as $t \rightarrow 0$ and we get the approximation (1.10). Combining (1.10) and (1.3) gives (1.11).

For $s > 0$, the arrival time equation (0.3), continuity of Δu , and (S2) give that

$$(1.14) \quad \text{Hess}_u(\gamma_s, \gamma_s) = \frac{\text{Hess}_u(\nabla u, \nabla u)}{|\nabla u|^2} = \Delta u(\gamma(s)) + 1 \rightarrow \Delta u(\gamma(0)) + 1 = -\frac{1}{n-k}.$$

Since $\text{Hess}_u \rightarrow -\frac{1}{n-k} \Pi$, we conclude that $\Pi_{\text{axis}}(\gamma_s) \rightarrow 0$, giving the third claim. Finally, the last claim follows from (1.10) and $|\gamma_s| = 1$. $\square$

2. REDUCING THEOREM 0.4 TO AN ESTIMATE FOR RESCALED MCF

In this section, we will reduce the Arnold-Thom conjecture to an estimate for rescaled mean curvature flow.

A one-parameter family of hypersurfaces M_τ evolves by *mean curvature flow* (or *MCF*) if each point $x(\tau)$ evolves by $\partial_\tau x = -H \mathbf{n}$. Here H is the mean curvature and $\mathbf{n}$ a unit normal. The rescaled MCF $\Sigma_t = \frac{1}{\sqrt{-u}} \{x \mid u(x) = -e^{-t}\}$ is equivalent to simultaneously running MCF and rescaling space, up to reparameterizations of time and the hypersurfaces. A one-parameter family of hypersurfaces Σ_t flows by the *rescaled MCF* if

$$(2.1) \quad \partial_t x = - \left(H - \frac{1}{2} \langle x, \mathbf{n} \rangle \right) \mathbf{n}.$$

It will be convenient to set $\phi = H - \frac{1}{2} \langle x, \mathbf{n} \rangle$. The fixed points for rescaled MCF are shrinkers where $\phi = 0$; the most important examples are cylinders $\mathcal{C} = \mathbf{S}^{n-k}_{\sqrt{2(n-k)}} \times \mathbf{R}^k$ where $k = 0, \dots, n-1$. Below, $\Pi : \mathbf{R}^{n+1} \rightarrow \mathbf{R}^{n-k+1}$ is orthogonal projection on the orthogonal complement of the axis $\mathbf{R}^k$ of the cylinder $\mathcal{C}$. The rescaled MCF is the negative gradient flow for the Gaussian area

$$(2.2) \quad F(\Sigma) \equiv \int_{\Sigma} e^{-\frac{|x|^2}{4}}.$$

In particular, $F(\Sigma_t)$ is non-increasing. Define the sequence δ_j by

$$(2.3) \quad \delta_j = \sqrt{F(\Sigma_{j-1}) - F(\Sigma_{j+2})}.$$

As in [CM1], the entropy $\lambda(\Sigma)$ is $\sup_{t_0 > 0, x_0 \in \mathbf{R}^{n+1}} F(t_0 \Sigma + x_0)$. We will use the Gaussian L^p norm given by $\|g\|_{L^p(\Sigma_t)}^p \equiv \int_{\Sigma_t} |g|^p e^{-\frac{|x|^2}{4}}.$

2.1. Summability of δ_j . As we will see in (2.18) below, δ_j bounds the distance that Σ_t evolves from j to $j+1$. Existence of $\lim_{t \rightarrow \infty} \Sigma_t$ is proven in [CM2] by showing that $\sum \delta_j < \infty$. We will need that δ_j is summable even after being raised to a power less than one:

Proposition 2.4. There exists $\bar{\beta} < 1$ so that

$$(2.5) \quad \sum_{j=1}^{\infty} \delta_j^{\bar{\beta}} < \infty.$$

Proof. By (6.21) and lemma 6.9 in [CM2], there exists $\rho > 1$ and C so that

$$(2.6) \quad \sum_{k=j}^{\infty} \delta_k^2 \leq 3 \left(F(\Sigma_{j-1}) - \lim_{t \rightarrow \infty} F(\Sigma_t) \right) \leq C j^{-\rho}.$$

Moreover, lemma 6.9 in [CM2] shows that this implies that $\sum \delta_j < \infty$.

We will show next that if $0 < q < \rho$, then

$$(2.7) \quad \sum \delta_j^2 j^q < \infty.$$

To prove this, set $b_j = j^q$ and $a_j = \sum_{i=j}^{\infty} \delta_i^2$, then $a_j - a_{j+1} = \delta_j^2$ and

$$(2.8) \quad b_{j+1} - b_j = (j+1)^q - j^q \leq c j^{q-1},$$

where c depends on q and we used that $j \geq 1$. Summation by parts gives

$$(2.9) \quad \begin{aligned} \sum_{j=k}^N \delta_j^2 j^q &= \sum_{j=k}^N b_j (a_j - a_{j+1}) = b_k a_k - b_N a_{N+1} + \sum_{j=k}^{N-1} a_{j+1} (b_{j+1} - b_j) \\ &\leq k^q \sum_{j=k}^{\infty} \delta_j^2 + C \sum_{j=k}^{\infty} j^{-\rho} j^{q-1}. \end{aligned}$$

This is bounded independently of N since $q < \rho$, giving (2.7).

Suppose that $a > 0$. The Hölder inequality gives $\sum \delta_j^{\beta} = \sum \left(\delta_j^2 j^a \right) j^{-a} < \infty$ if

$$(2.10) \quad \sum \delta_j^2 j^{\frac{2a}{\beta}} + \sum j^{-\frac{2a}{2-\beta}} < \infty.$$

To get (2.5), we need $\beta < 1$ and a so that both sums in (2.10) are finite. By (2.7), the first is finite if $\frac{2a}{\beta} < \rho$. The second is finite if $2 - \beta < 2a$. To satisfy both, we must have

$$(2.11) \quad 2 - \beta < 2a < \rho \beta.$$

This is possible as long as $2 < (1 + \rho) \beta$. Since $1 < \rho$, we can choose such a $\beta < 1$. $\square$

2.2. Cylindrical approximation. The rescaled MCF Σ_t converges to a limiting cylinder $\mathcal{C}$ by [CM2]. Thus, for each large integer j , Σ_j is well-approximated by $\mathcal{C}$.

In the next proposition, we will bound the distance from Σ_t to some cylinder $\mathcal{C}_t$ that is allowed to change with t . We will let Π_t denote the projection orthogonal to axis of $\mathcal{C}_t$. The operator $\mathcal{L}$ will be the drift Laplacian on the cylinder $\mathcal{C}_t$. Property (1) collects a priori estimates for the graph function w , (2) shows that w almost satisfies the linearized equation, (3) shows the approximating cylinders converge, and (4) gives a priori bounds on higher derivatives. We will only use (3) in this section; (1), (2) and (4) will be used later.

Proposition 2.12. Given $0 < \epsilon_1$ and $\beta < 1$, there exist a constant C and a sequence of radii R_j and cylinders $\mathcal{C}_j$ satisfying:

- (1) For $t \in [j, j+1]$, Σ_t is a graph over $B_{R_j} \cap \mathcal{C}_{j+1}$ of a function w with $\|w\|_{C^4} \leq \frac{\epsilon_1}{R_j}$,

$$\begin{aligned} \|w\|_{W^{3,2}}^2 + \|\phi\|_{W^{3,2}(B_{R_j})} + e^{-\frac{R_j^2}{4}} &\leq C \delta_j^\beta, \\ w^2 + |\nabla w|^2 + |\text{Hess}_w|^2 + |\nabla \text{Hess}_w|^2 + |\nabla^2 \text{Hess}_w|^2 &\leq C \delta_j^\beta e^{\frac{|x|^2}{4}}, \\ \phi^2 + |\nabla \phi|^2 + |\text{Hess}_\phi|^2 &\leq C \delta_j^{2\beta} e^{\frac{|x|^2}{4}}. \end{aligned}$$

- (2) The function w and its Euclidean partial derivatives w_i and w_{ij} on $\mathcal{C}_t$ satisfy

$$\begin{aligned} |\phi - (\mathcal{L} + 1)w| &\leq C(1 + R_j)(w^2 + |\nabla w|^2) + C(|w| + |\nabla w|) |\text{Hess}_w|, \\ \left| \phi_i - \left(\mathcal{L} + \frac{1}{2} \right) w_i \right| &\leq C(1 + R_j) (|w| + |\nabla w| + |\text{Hess}_w| + |\nabla \text{Hess}_w|) (|w| + |\nabla w| + |\nabla w_i|), \\ |\phi_{ij} - \mathcal{L} w_{ij}| &\leq C(1 + R_j) (|w| + |\nabla w| + |\text{Hess}_w| + |\nabla \text{Hess}_w| + |\nabla^2 \text{Hess}_w|)^2. \end{aligned}$$

- (3) $|\Pi_j - \Pi_{j+1}| \leq C \delta_j^\beta$.

- (4) Given any ℓ , there exists C_ℓ with $|\nabla^\ell w| + |\nabla^\ell \phi| \leq C_\ell$.

Proof. Let $\epsilon_0 > 0$ and α be fixed as in the definition of r_ℓ on page 261 in [CM2]. We will initially find a radius R'_j so that every estimate (1), (2), (3) and (4) holds except for the C^4 bound in (1) which we replace by $\|w\|_{C^{2,\alpha}} \leq \epsilon_0$. We will then use (1) to get the C^4 bound on a slightly smaller $R_j < R'_j$ with the other bounds still holding.

As in (5.2) in [CM2], define R'_j by $e^{-\frac{(R'_j)^2}{2}} = \delta_j^2$. Since $\Sigma_t \rightarrow \mathcal{C}$, we can assume that Σ_t is fixed close to $\mathcal{C}$ on a large set. Theorem 5.3 in [CM2] gives C and $\mu > 0$ and a cylinder $\mathcal{C}_{j+1}$ so that $B_{(1+2\mu)R'_j - \mathcal{C}} \cap \Sigma_t$, for $t \in [j, j+1]$, is a graph over $\mathcal{C}_{j+1}$ of a function w with $\|w\|_{C^{2,\alpha}} \leq \epsilon_0$ and, moreover, (4) holds. Furthermore, lemma 5.32 in [CM2] gives C so that

$$(2.13) \quad \int_{B_{(1+\mu)R'_j} \cap \Sigma_t} |\phi|^2 e^{-\frac{|x|^2}{4}} \leq C \delta_j^2.$$

Using theorem 0.24 from [CM2], we get for any $\beta_1 < 1$ that

$$(2.14) \quad \|w\|_{L^2}^2 \leq C_{\beta_0, \beta_1} \delta_j^{\beta_1}.$$

Using the higher derivative bound from (4) and interpolation (e.g., lemma B.1 in [CM2]), we get for any $\beta_2 < \beta_1$ that

$$(2.15) \quad \|w\|_{W^{3,2}}^2 \leq C_{\beta_0, \beta_1, \beta_2} \delta_j^{\beta_2}.$$

We have now established the first part of (1). Similarly, the second two parts of (1) follow from the first part, (4) and interpolation again.

We turn next to property (2). Lemma 4.6 in [CM2] computes the nonlinear graph equation for shrinkers; using p for points in $\mathcal{C}_t$, this gives

$$(2.16) \quad \phi = \hat{f}(w, \nabla w) + \langle p, V(w, \nabla w) \rangle + \langle \Phi(w, \nabla w), \text{Hess}_w \rangle,$$

where $\hat{f}(s, y)$, $V(s, y)$ and $\Phi(s, y)$ are smooth functions for $|s|$ small. Moreover, since $|A|^2 = \frac{1}{2}$ on $\mathcal{C}_t$, the operator $\mathcal{L} + 1$ is the linearized operator for the shrinker equation and lemma 4.10 in [CM2] gives that

$$(2.17) \quad |\phi - (\mathcal{L} + 1)w| \leq C_1(w^2 + |\nabla w|^2) + C_2(|w| + |\nabla w|) |\text{Hess}_w| ,$$

where $C_1 \leq C(1 + |p|)$ and C_2 is bounded. This gives the first claim in (2). Differentiating (2.16) in a Euclidean direction x_i and arguing similarly gives the second claim. Finally, differentiating (2.16) again gives the remaining claim in (2).

We next prove (3) by bounding the Gaussian distance from Σ_j to Σ_{j+1} by $C\delta_j$ and showing that $\mathcal{C}_j$ is Lipschitz in Σ_j . The first part follows since $|x_t| = |\phi|$ and

$$(2.18) \quad \int_j^{j+1} \|\phi\|_{L^1} dt \leq C \int_j^{j+1} \|\phi\|_{L^2} dt \leq C \left(\int_j^{j+1} \|\phi\|_{L^2}^2 dt \right)^{\frac{1}{2}} \leq C\delta_j .$$

To see that $\mathcal{C}_j$ is Lipschitz in Σ_j , we need to slightly modify the proof of theorem 0.24 in [CM2]. The choice of the cylinder in [CM2] occurs on page 240 in step 1 of the proof of proposition 2.1 there. There, the $\mathbf{R}^k$ factor is determined to be the approximate kernel of A at any point p in a fixed ball $B_{2\sqrt{2n}}$. In [CM2], p is left arbitrary – it does not effect the bounds in (1), (2) and (4) – and the $\mathbf{R}^k$ factor given by choosing any p would work (all that is needed are (2.22)–(2.24) there). To make $\mathcal{C}_j$ Lipschitz in Σ_j , we will choose the $\mathbf{R}^k$ factor by averaging over the approximate kernel of A for each point in the ball $B_{2\sqrt{2n}}$. The resulting $\mathbf{R}^k$ factor, and thus the cylinder, is then Lipschitz in Σ_j as desired.

Finally, we will fix $R_j \leq R'_j$ where $\|w\|_{C^4} \leq \frac{\epsilon_1}{R_j}$ and we still have $e^{-\frac{R_j^2}{4}} \leq C\delta_j^\beta$ where C now also depends on ϵ_1 . This follows from the pointwise C^4 bounds in (1) and $e^{-\frac{(R'_j)^2}{4}} = \delta_j$. $\square$

2.3. Reduction. The next theorem reduces Theorem 0.4 to an estimate for rescaled MCF.

Theorem 2.19. Theorem 0.4 holds if every rescaled MCF Σ_t with $\lambda(\Sigma_t) < \infty$ that goes to a cylinder as $t \rightarrow \infty$ satisfies

$$(2.20) \quad \sum_{j=1}^{\infty} \int_j^{j+1} \left(\sup_{B_{2n} \cap \Sigma_t} |\Pi_{j+1}(\nabla H)| \right) dt < \infty .$$

We will prove Theorem 2.19 here and (2.20) in Section 5. Suppose, therefore, that the function u and reparameterized gradient flow line $\gamma(s)$ are as in Section 1. In particular, $\gamma(s)$ is defined on $[0, \ell]$ with $|\gamma_s| = 1$ and $\gamma(0) \in \mathcal{S}$. We will show that γ_s has a limit as $s \rightarrow 0$. The derivative of $\gamma_s = -\frac{\nabla u}{|\nabla u|}$ is

$$(2.21) \quad \gamma_{ss} = -\frac{1}{|\nabla u|} (\text{Hess}_u(\gamma_s) - \gamma_s \langle \text{Hess}_u(\gamma_s), \gamma_s \rangle) = -\frac{(\text{Hess}_u(\gamma_s))^T}{|\nabla u|} = \nabla^T \log |\nabla u| ,$$

where $(\cdot)^T$ is the tangential projection onto the level set of u .

The simplest way to prove that $\lim \gamma_s$ exists would be to show that $\int |\gamma_{ss}| < \infty$, which is related to the rate of convergence for an associated rescaled MCF. While this rate fails to give integrability of $|\gamma_{ss}|$, it does give the following:

Lemma 2.22. Given any $\Lambda > 1$, we have $\lim_{s \rightarrow 0} \int_s^{\Lambda s} |\gamma_{ss}| ds = 0$.

Proof. Using Theorem 1.9 and the fact that $\text{Hess}_u \rightarrow -\frac{1}{n-k} \Pi$, (2.21) implies that $s |\gamma_{ss}| \rightarrow 0$. The lemma follows immediately from this. $\square$

To get around the lack of integrability, we will decompose γ_s into two pieces - the parts tangent and orthogonal to the axis - and deal with these separately. The tangent part goes to zero by (1.12) in Theorem 1.9. We will use (2.20) to control the orthogonal part.

Proof of Theorem 2.19. Translate so that $\gamma(0) = 0$ and let $\bar{H} = \frac{1}{|\nabla u|}$ be the mean curvature of the level set of u . The mean curvature H of Σ_t at time $t = -\log(-u)$ is given by

$$(2.23) \quad \bar{\nabla} \log \bar{H} = \frac{\nabla \log H}{\sqrt{-u}} \approx \frac{\sqrt{2(n-k)}}{s} \nabla \log H.$$

Note that $u(\gamma(s))$ is decreasing and Theorem 1.9 gives $t(s) \approx -2 \log s + \log(2(n-k))$ and

$$(2.24) \quad t'(s) = -\partial_s (\log(-u(\gamma(s)))) = \frac{-\partial_s u(\gamma(s))}{u(\gamma(s))} \approx -\frac{2}{s}.$$

Given a positive integer j , define s_j so that $t(s_j) = j$. Note that $\left| \log \frac{s_{j+1}}{s_j} \right|$ is uniformly bounded. Therefore, by Lemma 2.22, it suffices to show that γ_{s_j} has a limit.

We can write $\gamma_{s_j} = \Pi_{\text{axis},j}(\gamma_{s_j}) + \Pi_j(\gamma_{s_j})$. We have $\Pi_{\text{axis},j}(\gamma_{s_j}) \rightarrow 0$ since $\Pi_{\text{axis},j} \rightarrow \Pi_{\text{axis}}$ and $\Pi_{\text{axis}}(\gamma_s) \rightarrow 0$. Thus, we need that $\lim_{j \rightarrow \infty} \Pi_j(\gamma_{s_j})$ exists; this will follow from

$$(2.25) \quad \sum_j |\Pi_j(\gamma_{s_j}) - \Pi_{j+1}(\gamma_{s_{j+1}})| < \infty.$$

Theorem 1.9 gives (for s small) that $\gamma(s) \subset B_{2n\sqrt{-u(\gamma(s))}}$ and, thus, (2.21) gives

$$(2.26) \quad \begin{aligned} |\Pi_{j+1}(\gamma_{s_j}) - \Pi_{j+1}(\gamma_{s_{j+1}})| &\leq \int_{s_{j+1}}^{s_j} |\Pi_{j+1}(\gamma_{ss})| ds = \int_{s_{j+1}}^{s_j} |\Pi_{j+1}(\bar{\nabla} \log \bar{H}(\gamma(s)))| ds \\ &\leq C \int_{s_{j+1}}^{s_j} \sup_{B_{2n\sqrt{-u(\gamma(s))}}} |\Pi_{j+1}(\nabla \log \bar{H})|(\cdot, -u) ds. \end{aligned}$$

Using (2.23) and (2.24) in (2.26) and then applying Theorem 2.19 gives

$$(2.27) \quad \sum_j |\Pi_{j+1}(\gamma_{s_j}) - \Pi_{j+1}(\gamma_{s_{j+1}})| \leq C \sum_j \int_j^{j+1} \sup_{B_{2n} \cap \Sigma_t} |\Pi_{j+1}(\nabla H)| dt < \infty.$$

On the other hand, $\sum_j |\Pi_j(\gamma_{s_j}) - \Pi_{j+1}(\gamma_{s_j})| < \infty$ by (3) in Proposition 2.12 and Proposition 2.4. Therefore, the triangle inequality gives (2.25), completing the proof. $\square$

3. APPROXIMATE EIGENFUNCTIONS ON CYLINDERS

The key remaining point is summability of $\Pi_{j+1}(\nabla H)$. The bound for w^2 in (1) from Proposition 2.12 is summable by Proposition 2.4, but the bound for w is not. In particular, (1) gives a bound for ∇H that is not summable. This bound for ∇H cannot be improved due to slowly growing Jacobi fields. However, these Jacobi fields do not contribute to $\Pi_{j+1}(\nabla H)$. We will show that the remainder of w , after we subtract these Jacobi fields, is small.

In this section, we will show that if an approximate eigenfunction w on a cylinder $\mathcal{C}$ begins to grow, then it must grow rapidly. The key tool is the frequency function for the drift

Laplacian as in [CM7]; the difficulty here is handling error terms. Let $x \in \mathbf{R}^k$ be coordinates on the Euclidean factor, $f = \frac{|x|^2}{4}$, $\mathcal{L}$ the drift Laplacian $\mathcal{L} = \Delta_C - \frac{1}{2} \nabla_x = \Delta_\theta + \mathcal{L}_{\mathbf{R}^k}$, where Δ_θ is the Laplacian on $\mathbf{S}^{n-k}_{\sqrt{2(n-k)}}$, and $\text{div}_f = \text{div} - \langle \frac{x}{2}, \cdot \rangle$ the drift divergence.

In applications, w will be given by Proposition 2.12 and, thus, will satisfy (1), (2) and (4) there. Thus, we will assume that w is a function on $\{|x| < R\} \subset \mathcal{C}$ satisfying:

$$(3.1) \quad |(\mathcal{L} + 1)w - \phi| \leq \epsilon (|w| + |\nabla w|) \text{ where } \phi \text{ is a function and } \frac{8}{9} < (1 - 3\epsilon)^3,$$

$$(3.2) \quad \sup_{|x| < 4n} |(\mathcal{L} + 1)w| \leq \mu.$$

Equation (3.1) arises from w satisfying a nonlinear equation $\mathcal{M}w = \phi$ and $\mathcal{L} + 1$ is the linearization of $\mathcal{M}$. We will also assume that $\mu > 0$ is small and

$$(3.3) \quad \|w\|_{W^{3,2}}^2 + \|\phi\|_{W^{3,2}} + e^{-\frac{R^2}{4}} \leq \mu,$$

$$(3.4) \quad w^2(x) + |\nabla w(x)|^2 + |\text{Hess}_w(x)|^2 \leq \mu e^{\frac{|x|^2}{4}}.$$

We will assume that the Euclidean first derivatives w_i and second derivatives w_{ij} satisfy

$$(3.5) \quad \left| \left(\mathcal{L} + \frac{1}{2} \right) w_i \right| \leq |\phi_i| + \epsilon (|w| + |\nabla w| + |\nabla w_i|),$$

$$(3.6) \quad \sup_{|x| < r} |\mathcal{L} w_{ij}| \leq C_r \mu \text{ where } C_r \text{ depends on } r.$$

By lemma 3.26 in [CM2], the kernel of $\mathcal{L} + 1$ on the weighted Gaussian space on $\mathcal{C}$ consists of quadratic polynomials and “infinitesimal rotations” of the form

$$(3.7) \quad \tilde{w} = \sum_i a_i (x_i^2 - 2) + \sum_{i < j} a_{ij} x_i x_j + \sum_k x_k h_k(\theta),$$

where a_i, a_{ij} are constants and each h_k is a Δ_θ -eigenfunction with eigenvalue $\frac{1}{2}$.

The next theorem quadratically approximates w in $|x| \leq 3n$ by $\tilde{w}$ as in (3.7). Namely, while (3.4) gives $|w| \leq C \mu^{\frac{1}{2}}$, the next theorem gives $|w - \tilde{w}| \leq C \mu^\nu$ with $\nu \approx 1$.

Theorem 3.8. Given $\nu < 1$, there exists $C, \bar{\epsilon}$, and $\mu_0 > 0$ so that if w satisfies (3.1)–(3.6) with $\mu_0 > \mu$ and $\bar{\epsilon} > \epsilon$, then there is a function $\tilde{w}$ as in (3.7) with

$$(3.9) \quad \sup_{|x| \leq 3n} |w - \tilde{w}| \leq C \mu^\nu.$$

3.1. First reduction.

Lemma 3.10. If w satisfies (3.1)–(3.6), then there is a function $\tilde{w}$ as in (3.7) so that $v = w - \tilde{w}$ satisfies (3.1)–(3.6) and

- (A1) Each Euclidean second derivative v_{ij} has $\int_{x=0} v_{ij} = 0$.
- (A2) Each Euclidean first derivative v_i has $\int_{x=0} v_i h = 0$ for any h with $\Delta_\theta h = -\frac{1}{2} h$.
- (A3) We have $|\int_{x=0} v| \leq \mu \text{Vol}(x=0)$.

Proof. Given $\tilde{w}$ as in (3.7), the Euclidean first and second derivatives are given at $x = 0$ by

$$(3.11) \quad \tilde{w}_i = h_i(\theta), \tilde{w}_{ii} = 2a_i, \tilde{w}_{ij} = a_{ij} \text{ for } i < j.$$

To arrange (A1), define a_i and a_{ij} by

$$(3.12) \quad 2a_i \int_{x=0} 1 = \int_{x=0} w_{ii} \text{ and } a_{ij} \int_{x=0} 1 = \int_{x=0} w_{ij} \text{ for } i < j.$$

Similarly, for (A2), let h_i be the projection of w_i onto the $\frac{1}{2}$ -eigenspace of Δ_θ at $x = 0$. Claim (A3) follows by integrating $(\mathcal{L} + 1)w$ at $x = 0$ and using (3.2) and (A1). $\square$

For a function v , we let $\text{Hess}_v^x = \frac{\partial^2 v}{\partial x_i \partial x_j}$ denote its Euclidean Hessian.

Corollary 3.13. Given $\beta > 2$, there exists C so that if v satisfies (3.1)–(3.6) and (A1)–(A3), then

$$(3.14) \quad \sup_{|x| < 3n} \left(|v|^\beta + |\nabla_{\mathbf{R}^k} v|^\beta + |\text{Hess}_v^x|^\beta \right) \leq C \mu^2 + C \int_{|x| < 4n} |\text{Hess}_v^x|^2.$$

Similarly, given $\beta > 2$ and $r > 3n$, there exists $C_{\beta,r}$ so that

$$(3.15) \quad \sup_{|x| < r} \left(|v|^\beta + |\nabla_{\mathbf{R}^k} v|^\beta + |\text{Hess}_v^x|^\beta \right) \leq C_{\beta,r} \mu^2 + C_{\beta,r} \int_{|x| < r+1} |\text{Hess}_v^x|^2.$$

Proof. We will prove (3.14); (3.15) follows similarly. Set $\delta^2 = \mu^2 + \int_{|x| < 4n} |\text{Hess}_v^x|^2$. Since we have uniform higher derivative bounds on v , interpolation implies that all norms are equivalent if we go to any worse power. Thus, given any $\beta_1 < 1$, (3.2) gives

$$(3.16) \quad \|\text{Hess}_v^x\|_{C^2} + \|(\mathcal{L} + 1)v\|_{C^2} \leq C_1 \delta^{\beta_1},$$

where C_1 depends on β_1 . It follows that $|(\Delta_\theta + 1)v(\theta, 0)| \leq C_1 \delta^{\beta_1}$. Since $\Delta_\theta + 1$ is invertible (lemma 2.5 in [CM2]), this (and interpolation again) gives for any $\beta_2 < \beta_1$ that

$$(3.17) \quad |v(\theta, 0)| \leq C_2 \delta^{\beta_2}.$$

Given a Euclidean first derivative v_i , (3.16) gives that

$$(3.18) \quad \left| \left(\Delta_\theta + \frac{1}{2} \right) v_i(\theta, 0) \right| \leq C_1 \delta^{\beta_1}.$$

The operator $(\Delta_\theta + \frac{1}{2})$ is not invertible, but (A2) implies that $v_i(\theta, 0)$ is orthogonal to the kernel so we get (using interpolation again) that $|v_i(\theta, 0)| \leq C_2 \delta^{\beta_2}$. The bound on v_i at $x = 0$ and the Hessian bound give a bound on v_i everywhere. Integrating this and using (3.17) gives the desired pointwise bound on v , completing the proof. $\square$

3.2. The frequency. Given a function u on $\mathcal{C}$, define I and D by²

$$(3.19) \quad I(r) = r^{1-k} \int_{|x|=r} u^2,$$

$$(3.20) \quad D(r) = r^{2-k} \int_{|x|=r} u u_r = e^{\frac{r^2}{4}} r^{2-k} \int_{|x| < r} (|\nabla u|^2 + u \mathcal{L} u) e^{-f}.$$

Here u_r denotes the normal derivative of u on the level set $|x| = r$. Note that f is proper. It is easy to see that $I' = \frac{2D}{r}$ and $(\log I)' = \frac{2U}{r}$, where the frequency $U = \frac{D}{I}$; cf. [Be], [CM7].

²When $k = 1$ and the sphere is disconnected, let r be signed distance and set $I(|r|) = \int_{x=r} u^2 + \int_{x=-r} u^2$.

The next theorem shows that if the growth of an approximate eigenfunction hits a certain threshold, then it grows very rapidly. The theorem is stated for eigenvalue 1, but generalizes easily to other eigenvalues. The case $(\mathcal{L} + 1)v = 0$, where $\epsilon = \phi = 0$, follows from [CM7].

Theorem 3.21. Given $r_1 > \max\{9n, 4n + 64\sqrt{2}\}$, there exist $\bar{R} = \bar{R}(n, r_1)$, $C = C(n, r_1)$ so that if v is a function on $\{|x| \leq R\}$ satisfying (3.1), where $\bar{R} \leq R$, and

$$(3.22) \quad 2 \int_{|x| < 9n} v^2 e^{-f} \leq \int_{|x| < r_1} v^2 e^{-f} \text{ and } \frac{r_1^2}{16} < U(r_1),$$

then for any $\Lambda \in (0, 1/3)$

$$(3.23) \quad \int_{|x| < 4n} v^2 e^{-f} \leq \Lambda^{-2} \|\phi\|_{L^2}^2 + C I(R) R^{2n+68} e^{-\frac{(1-3\epsilon-\Lambda)R^2}{2(1+3\epsilon+\Lambda)^2}}.$$

To prove Theorem 3.8, we will find a scale r_1 where Theorem 3.21 applies to give that w is bounded by μ^ν . To do this, we will find a long stretch where Hess_w^x must grow and, thus, w must also have grown. Note that Hess_w^x is easier to work with since each $\mathbf{R}^k$ derivative lowers the eigenvalue by $1/2$ and, thus, lowers the threshold for growth (cf. [CM7]).

The proof of Theorem 3.21 uses a modified version of the frequency. Define E and U_E by

$$(3.24) \quad E(r) = r^{2-k} e^{\frac{r^2}{4}} \int_{|x| < r} \{|\nabla v|^2 - v^2\} e^{-f} = D(r) - r^{2-k} e^{\frac{r^2}{4}} \int_{|x| < r} (v\mathcal{L}v + v^2) e^{-f},$$

$$(3.25) \quad U_E(r) = \frac{E(r)}{I(r)}.$$

Lemma 3.26. If $E(r) > 0$, then

$$(3.27) \quad (\log U_E)'(r) \geq \frac{2-k}{r} + \frac{r}{2} - \frac{r}{U_E} + \frac{U(r)}{r} \left(\frac{D(r)}{E(r)} - 2 \right).$$

Proof. The Cauchy-Schwarz inequality $(\int uu_r)^2 \leq \int u^2 \int |\nabla u|^2$ gives

$$(3.28) \quad E'(r) = \frac{2-k}{r} E + \frac{r}{2} E + r^{2-k} \int_{|x|=r} (|\nabla v|^2 - v^2) \geq \frac{2-k}{r} E + \frac{r}{2} E + \frac{UD}{r} - rI.$$

The lemma follows from this since $\frac{I'}{I} = \frac{2U}{r}$. □

The next lemma is valid for any function v .

Lemma 3.29. If $r > \bar{r} > 3n$ and

$$(3.30) \quad \int_{|x| < \bar{r}} v^2 e^{-f} \leq \int_{\bar{r} < |x| < r} v^2 e^{-f},$$

then

$$(3.31) \quad \int_{|x| < r} v^2 e^{-f} \leq \frac{32}{\bar{r}^2 - 8n} \int_{|x| < r} |\nabla v|^2 e^{-f}.$$

Proof. Since $\mathcal{L}|x|^2 = 2k - |x|^2$, we have $\operatorname{div}_f(v^2 x) = 2v\langle x, \nabla v \rangle + v^2(k - |x|^2/2)$. Using the absorbing inequality $2|v\langle x, \nabla v \rangle| \leq |x|^2 v^2/4 + 4|\nabla v|^2$, the divergence theorem gives

$$(3.32) \quad \begin{aligned} \int_{|x|<r} \left(\frac{|x|^2}{2} - k \right) v^2 e^{-f} &= -r e^{-\frac{r^2}{4}} \int_{|x|=r} v^2 + 2 \int_{|x|<r} v \langle x, \nabla v \rangle e^{-f} \\ &\leq -r e^{-\frac{r^2}{4}} \int_{|x|=r} v^2 + \int_{|x|<r} \left(v^2 \frac{|x|^2}{4} + 4|\nabla v|^2 \right) e^{-f}. \end{aligned}$$

It follows that

$$\left(\frac{\bar{r}^2}{4} - k \right) \int_{\bar{r}<|x|<r} v^2 e^{-f} - k \int_{|x|<\bar{r}} v^2 e^{-f} \leq \int_{|x|<r} \left(\frac{|x|^2}{4} - k \right) v^2 e^{-f} \leq 4 \int_{|x|<r} |\nabla v|^2 e^{-f}.$$

Bringing in the assumption (3.30) gives

$$\left(\frac{\bar{r}^2}{4} - 2k \right) \int_{\bar{r}<|x|<r} v^2 e^{-f} \leq 4 \int_{|x|<r} |\nabla v|^2 e^{-f}.$$

The lemma follows from this and using the assumption again. $\square$

Proof of Theorem 3.21. We can assume that $\|\phi\|_{L^2}^2 \leq \Lambda^2 \int_{|x|<4n} v^2 e^{-f}$ since otherwise we get (3.23) immediately. Therefore, given any $r \geq 4n$, (3.1) gives

$$(3.33) \quad |D - E|(r) \leq r^{2-k} e^{\frac{r^2}{4}} \int_{|x|<r} ((\epsilon + \Lambda)v^2 + \epsilon|\nabla v|^2) e^{-f}.$$

Suppose now that some $r \geq 4n$ satisfies

$$(\star 1) \quad \int_{|x|<r} v^2 e^{-f} \leq \frac{1}{2} \int_{|x|<r} |\nabla v|^2 e^{-f}.$$

We will use $(\star 1)$ to show that $D(r)$ and $E(r)$ are comparable, get a differential inequality for $U_E(r)$ and bound the ratio of the derivatives of quantities in $(\star 1)$. Namely, $(\star 1)$ gives

$$(3.34) \quad \frac{1}{2} r^{2-k} e^{\frac{r^2}{4}} \int_{|x|<r} |\nabla v|^2 e^{-f} \leq r^{2-k} e^{\frac{r^2}{4}} \int_{|x|<r} (|\nabla v|^2 - v^2) e^{-f} = E(r).$$

Similarly, using (3.33), $(\star 1)$ and (3.34) gives that

$$(3.35) \quad |D - E|(r) \leq \left(\frac{3\epsilon}{2} + \frac{\Lambda}{2} \right) r^{2-k} e^{\frac{r^2}{4}} \int_{|x|<r} |\nabla v|^2 e^{-f} \leq (3\epsilon + \Lambda) E(r).$$

We conclude that $D(r)$, and thus also $I'(r)$, are also positive and

$$(3.36) \quad |U - U_E|(r) \leq (3\epsilon + \Lambda) U_E(r),$$

$$(3.37) \quad (1 - 3\epsilon - \Lambda) U_E \leq U(r) \leq (1 + 3\epsilon + \Lambda) U_E.$$

Using this in Lemma 3.26 gives the differential inequality at r

$$(3.38) \quad (\log U_E)' \geq \frac{2-k}{r} + \frac{r}{2} - \frac{r}{U_E} - (1 + 3\epsilon + \Lambda)^2 \frac{U_E}{r}.$$

From (3.37), the definition of U , and the Cauchy-Schwarz inequality, we get at r that

$$(3.39) \quad (1 - 3\epsilon - \Lambda)^2 U_E^2 I^2 \leq U^2 I^2 = D^2 \leq I r^{3-k} \int_{|x|=r} |\nabla v|^2.$$

Noting that $3\epsilon + \Lambda < \frac{1}{2}$, we get

$$(3.40) \quad \frac{U_E^2(r)}{4r^2} \leq (1 - 3\epsilon - \Lambda)^2 \frac{U_E^2(r)}{r^2} \leq \frac{\int_{|x|=r} |\nabla v|^2}{\int_{|x|=r} v^2}.$$

We will also need a second property (the first part is the strict form of $(\star 1)$):

$$(\star 2) \quad \int_{|x|<r} v^2 e^{-f} < \frac{1}{2} \int_{|x|<r} |\nabla v|^2 e^{-f} \text{ and } \frac{r^2}{32} < U_E(r).$$

Set $r_0 = 4n + 64\sqrt{2}$. We will show that if $(\star 2)$ holds for some $r \geq r_0$, then it holds for all $s \geq r$. We will argue by contradiction, so suppose that $s > r$ is the first time $(\star 2)$ fails. Note that $(\star 2)$ is equivalent to $U_E(r) > \frac{r^2}{32}$ and $F(r) > 0$ where $F(r) = \int_{|x|<r} (\frac{1}{2} |\nabla v|^2 - v^2) e^{-f}$. Since s is the first time, we have $F(s) \geq 0$ (i.e., $(\star 1)$), $U_E(s) - \frac{s^2}{32} \geq 0$ and

$$(3.41) \quad F(t) > 0 \text{ and } U_E(t) - \frac{t^2}{32} > 0 \text{ for all } t \in [r, s].$$

We also have that at least one of $F(s)$ and $U_E(s) - \frac{s^2}{32}$ is zero. Suppose first that $F(s) = 0$ and, thus, $F'(s) \leq 0$. However, (3.40) and $U_E(s) - \frac{s^2}{32} \geq 0$ give that

$$(3.42) \quad \left(\frac{s}{64}\right)^2 \leq \frac{U_E^2(s)}{4s^2} \leq \frac{\int_{|x|=s} |\nabla v|^2}{\int_{|x|=s} v^2}.$$

However, this implies that $F'(s) > 0$ as long as $s > 64\sqrt{2}$, giving the desired contradiction in the first case. Suppose now that $U_E(s) = \frac{s^2}{32}$ and, thus, $U'_E(s) \leq \frac{s}{16}$ and

$$(3.43) \quad (\log U_E)'(s) \leq \frac{2}{s}.$$

On the other hand, (3.38) gives that

$$(3.44) \quad (\log U_E)'(s) \geq \frac{2-k}{s} + \frac{s}{2} - \frac{32}{s} - (1 + 3\epsilon + \Lambda)^2 \frac{s}{32} \geq \frac{3s}{8} - \frac{k+30}{s}.$$

This contradicts (3.43) since $s \geq r_0$, completing the proof of the claim.

We will now show that $(\star 2)$ holds for r_1 . Using the first part of (3.22), we can apply Lemma 3.29 (with $\bar{r} = 9n$) to get

$$(3.45) \quad \int_{|x|<r_1} v^2 e^{-f} \leq \frac{32}{81n^2 - 8n} \int_{|x|<r_1} |\nabla v|^2 e^{-f} < \frac{1}{2} \int_{|x|<r_1} |\nabla v|^2 e^{-f},$$

where the last inequality used that $81n^2 > 8n + 64$. This gives the first part of $(\star 2)$; in particular, $(\star 1)$ holds and (3.37) gives that

$$(3.46) \quad U(r_1) \leq (1 + 3\epsilon + \Lambda) U_E(r_1) \leq \frac{3}{2} U_E(r_1).$$

Since $\frac{r_1^2}{16} \leq U(r_1)$ by the second part of (3.22), the second part of $(\star 2)$ also holds.

We have established that $(\star 2)$ holds for all $r \geq r_1$, so we get the differential inequality (3.38) for U_E and the equivalence (3.36) between U and U_E . This will give the desired

growth of U and, thus also, I . We do this next. Set $\kappa = (3\epsilon + \Lambda)$. We claim that there exists $\bar{R} = \bar{R}(k, r_1) \geq r_1$ so that for all $r \geq \bar{R}$ we have

$$(3.47) \quad U_E(r) > \frac{r^2 - 2k - 68}{2(1 + \kappa)^2}.$$

The key is that if (3.47) fails for some $r \geq r_1$, then (3.38) implies that

$$(3.48) \quad (\log U_E)' \geq \frac{r}{2} - \frac{k + 30}{r} - (1 + \kappa)^2 \frac{U_E}{r} \geq \frac{4}{r}.$$

On the other hand, for $r \geq 4k$, we have

$$(3.49) \quad \left(\log \frac{r^2 - 2k - 68}{2(1 + \kappa)^2} \right)' = \frac{2r}{r^2 - 2k - 68} < \frac{3}{r},$$

where the last inequality used that $6k + 204 < r_0^2$. Integrating (3.48) and (3.49) and using that $U_E \geq \frac{r^2}{32}$, gives an upper bound for the maximal interval where (3.47) fails. The first derivative test, (3.48), and (3.49) imply that once (3.47) holds for some $R \geq r_1$, then it also holds for all $r \geq R$. This gives the claim. Using (3.36) and (3.47), we get for $r \geq \bar{R}$ that

$$(3.50) \quad U(r) \geq (1 - \kappa)U_E(r) > \frac{(1 - \kappa)}{(1 + \kappa)^2} \left(\frac{r^2}{2} - k - 34 \right).$$

Integrating this from $\bar{R}$ to R gives that

$$(3.51) \quad \log \frac{I(R)}{I(\bar{R})} = 2 \int_{\bar{R}}^R \frac{U(r)}{r} dr \geq \frac{(1 - \kappa)}{(1 + \kappa)^2} \left(\frac{R^2 - \bar{R}^2}{2} - (2k + 68) \log \frac{R}{\bar{R}} \right).$$

Since $\bar{R}$ depends only on k and r_1 , exponentiating gives $C = C(k, r_1)$ so that

$$(3.52) \quad \sup_{r_1 \leq r \leq \bar{R}} I(r) = I(\bar{R}) \leq C I(R) R^{(2k+68) \frac{1-\kappa}{(1+\kappa)^2}} e^{-\frac{(1-\kappa)}{2(1+\kappa)^2} R^2}.$$

Choose $r_2 \in [r_1, 2r_1]$ that achieves the minimum of D on $[r_1, 2r_1]$. Since $I' = \frac{2D}{r}$, it follows that $D(r_2) \leq I(2r_1)$. Therefore, since $(\star 1)$ holds for r_2 , we have

$$(3.53) \quad r_2^{2-k} e^{\frac{r_2^2}{4}} \int_{|x| < r_2} v^2 e^f \leq E(r_2) \leq \frac{D(r_2)}{(1 - \kappa)} \leq 2 I(2r_1).$$

Finally, combining (3.52) and (3.53) gives (3.23). $\square$

4. GENERAL FREQUENCY

In this section, we will prove Theorem 3.8 by showing that either we already have the bound on w or (3.22) holds and Theorem 3.21 bounds w . Throughout this section, we will assume that w satisfies (3.1)–(3.6) and (A1)–(A3).

The main task left is to prove the following proposition:

Proposition 4.1. Given $r_+ \geq 9n$, there exist $\lambda > r_+$ and ζ_2 so that if $\zeta \geq \zeta_2$ and

$$(4.2) \quad \int_{|x| < 4n} |\text{Hess}_w^x|^2 \geq \zeta \mu^2,$$

then there exists $r_1 \in (r_+, \lambda)$ satisfying (3.22).

Throughout this section, C_r will be a constant that depends on r (but not on w or μ) that will be allowed to change from line to line.

4.1. Proof of Theorem 3.8 assuming Proposition 4.1.

Lemma 4.3. Given $r > 0$, there exists C_r so that

$$(4.4) \quad \left| \int_{|x|<r} \text{Hess}_w^x \right| + \left| \int_{|x|<r} w \right| \leq C_r \mu.$$

Furthermore, given any h on $\mathbf{S}^{n-k}$ with $\Delta_\theta h = -\frac{1}{2}h$ and $\int h^2(\theta) d\theta = 1$, we have

$$(4.5) \quad \left| \int_{|x|<r} h \nabla^{\mathbf{R}^k} w \right| \leq C_r \mu.$$

Proof. Let w_{ij} be a Euclidean second derivative and define the spherical average

$$(4.6) \quad J_{ij}(r) = r^{1-k} \int_{|x|=r} w_{ij}.$$

By (A1), we have $J_{ij}(0) = 0$. Note that $|\mathcal{L} w_{ij}| \leq C_r \mu$, so we have

$$(4.7) \quad |J'_{ij}|(r) \leq r^{1-k} e^{\frac{r^2}{4}} \int_{|x|<r} |\mathcal{L} w_{ij}| e^{-f} \leq C_r \mu.$$

Thus, we get that $|J_{ij}(r)| \leq C_r \mu$. Integrating this gives the integral bound on Hess_w in (4.4) and, thus, the same bound on $\left| \int_{|x|<r} \Delta_{\mathbf{R}^k} w \right|$. The bound on $\left| \int_{|x|<r} w \right|$ follows similarly by setting $J(r) = r^{1-k} \int_{|x|=r} w$. Namely, (A3) bounds $J(0)$ and we bound $J'(r)$ by using that $\Delta_\theta w$ integrates to zero over each sphere and $\left| \int_{|x|<r} \Delta_{\mathbf{R}^k} w \right| \leq C_r \mu$.

To get the last claim, define a vector-valued function $J_h(r)$ by

$$(4.8) \quad J_h(r) = r^{1-k} \int_{|x|=r} h \nabla^{\mathbf{R}^k} w,$$

so that $J_h(0) = 0$ by (A2). Arguing as above and using the integral bound on the Euclidean Hessian bounds $J_h(r)$ and integrating this gives the last claim. $\square$

Corollary 4.9. Given $\bar{r} > 4n$, there exist $C_{\bar{r}}$ so that

$$(4.10) \quad \int_{|x|<\bar{r}} |\text{Hess}_w^x|^2 \leq C_{\bar{r}} \mu^2 + C_{\bar{r}} \int_{\bar{r}<|x|<\bar{r}+1} |\text{Hess}_w^x|^2.$$

Proof. Set $\mathcal{A} = \{\bar{r} < |x| < \bar{r} + 1\}$. Let w_{ij} be a Euclidean second derivative and η a cutoff function that is one for $|x| < \bar{r}$, zero for $\bar{r} + 1 < |x|$, and $|\nabla \eta| \leq 2$. Given $\delta > 0$, we get

$$(4.11) \quad \begin{aligned} \text{div}_f (\eta^2 w_{ij} \nabla w_{ij}) &= \eta^2 (|\nabla w_{ij}|^2 + w_{ij} \mathcal{L} w_{ij}) + 2\eta w_{ij} \langle \nabla w_{ij}, \nabla \eta \rangle \\ &\geq \eta^2 \left(\frac{1}{2} |\nabla w_{ij}|^2 - \delta w_{ij}^2 - \frac{1}{4\delta} |\mathcal{L} w_{ij}|^2 \right) - 2|\nabla \eta|^2 w_{ij}^2. \end{aligned}$$

We get that

$$\begin{aligned}
 \int_{|x|<\bar{r}} |\nabla w_{ij}|^2 e^{-f} &\leq 2\delta \int_{|x|<\bar{r}} w_{ij}^2 e^{-f} + \frac{1}{2\delta} \int_{|x|<\bar{r}+1} |\mathcal{L} w_{ij}|^2 e^{-f} + (8+2\delta) \int_{\mathcal{A}} w_{ij}^2 e^{-f} \\
 (4.12) \quad &\leq 2\delta \int_{|x|<\bar{r}} w_{ij}^2 e^{-f} + C_{\bar{r}} \frac{\mu^2}{\delta} + (8+2\delta) \int_{\mathcal{A}} w_{ij}^2 e^{-f}.
 \end{aligned}$$

On the other hand, Lemma 4.3 and the Neumann Poincaré inequality give $C_{\bar{r}}$ so that

$$(4.13) \quad \int_{|x|<\bar{r}} w_{ij}^2 e^{-f} \leq C_{\bar{r}} \left(\mu^2 + \int_{|x|<\bar{r}} |\nabla w_{ij}|^2 e^{-f} \right).$$

Using this to bound the first term on the right in (4.12) and taking $\delta > 0$ small enough (depending on $\bar{r}$), this can be absorbed. Finally, summing over i, j gives the corollary. $\square$

Proof of Theorem 3.8. Lemma 3.10 gives $\tilde{w}$ as in (3.7) so that $v = w - \tilde{w}$ satisfies (3.1)–(3.6) and (A1), (A2) and (A3). By Corollary 3.13, it suffices to get $\int_{|x|<4n} v^2 \leq C \mu^\beta$ with $\nu < \beta$.

Proposition 4.1 gives λ and ζ_2 (depending just on n) so that if (4.2) holds with $\zeta \geq \zeta_2$, then there exists r_1 satisfying (3.22) with

$$(4.14) \quad r_1 \in (\max\{9n, \sqrt{8n+256}\}, \lambda).$$

We can assume that (4.2) holds with $\zeta \geq \zeta_2$ since the theorem otherwise follows from Corollary 3.13. Therefore, Theorem 3.21 applies and we get $\bar{R} = \bar{R}(n, r_1)$ and $C = C(n, r_1)$ so that for any $\Lambda \in (0, 1/3)$

$$\begin{aligned}
 \int_{|x|<4n} v^2 e^{-f} &\leq \Lambda^{-2} \|\phi\|_{L^2}^2 + C I(R) R^{2n+68} e^{-\frac{(1-3\epsilon-\Lambda) R^2}{2(1+3\epsilon+\Lambda)^2}} \\
 (4.15) \quad &\leq \Lambda^{-2} \mu^2 + C R^{2n+68} \left(e^{-\frac{R^2}{2}} \right)^{\frac{(1-3\epsilon-\Lambda)}{(1+3\epsilon+\Lambda)^2}}.
 \end{aligned}$$

This required $R \geq \bar{R}$; if $\bar{R} > R$, then there is a positive lower bound for μ and the theorem holds trivially. Since $e^{-\frac{R^2}{2}} \leq \mu^2$, the theorem follows by taking $\epsilon, \Lambda > 0$ small enough that

$$(4.16) \quad (1 - 3\epsilon - \Lambda) > \nu (1 + 3\epsilon + \Lambda)^2.$$

$\square$

4.2. Proof of Proposition 4.1. We will get a positive lower bound for the frequency U_2 for Hess_w^x that will force Hess_w^x to grow very rapidly. We will then combine Poincaré and reverse Poincaré inequalities to show that w itself grows rapidly as claimed. To do this, define quantities I_2 , D_2 and U_2 for Hess_w^x by

$$(4.17) \quad I_2(r) = r^{1-k} \int_{|x|=r} |\text{Hess}_w^x|^2,$$

$D_2 = \frac{r}{2} I_2'$, and $U_2 = \frac{D_2}{I_2}$ so that $(\log I_2)' = \frac{2U_2}{r}$. Define $\psi = (\mathcal{L} + 1)w$ so that $\mathcal{L} w_{ij} = \psi_{ij}$. Differentiating I_2 , we see that

$$(4.18) \quad D_2(r) = r^{2-k} \sum_{i,j} \int_{|x|=r} w_{ij} \partial_r w_{ij} = r^{2-k} e^{\frac{r^2}{4}} \int_{|x|<r} \left(|\nabla \text{Hess}_w^x|^2 + \sum_{i,j} w_{ij} \psi_{ij} \right) e^{-f}.$$

The next two lemmas give a differential inequality for U_2 when $U_2 > 0$ and then establish that $U_2(r)$ is positive on an interval.

Lemma 4.19. If $U_2(r) > 0$, then

$$(4.20) \quad (\log U_2)'(r) \geq \frac{2-k}{r} + \frac{r}{2} - \frac{U_2}{r} - \frac{r}{U_2} \left(\frac{r^{1-k} \int_{|x|=r} \sum_{i,j} \psi_{ij}^2}{I_2} \right)^{\frac{1}{2}}.$$

Proof. Differentiating D_2 gives that

$$(4.21) \quad D_2'(r) = \frac{2-k}{r} D_2 + \frac{r}{2} D_2 + r^{2-k} \int_{|x|=r} \left(|\nabla \text{Hess}_w^x|^2 + \sum_{i,j} w_{ij} \psi_{ij} \right).$$

The first equality in (4.18) and the Cauchy-Schwarz inequality give that

$$(4.22) \quad D_2^2(r) \leq I_2(r) r^{3-k} \int_{|x|=r} \sum_{i,j} (\partial_r w_{ij})^2 \leq I_2(r) r^{3-k} \int_{|x|=r} |\nabla \text{Hess}_w^x|^2.$$

Since $D_2(r) > 0$ (by assumption), using (4.22) in (4.21) and dividing by $D_2(r)$ gives

$$(4.23) \quad (\log D_2)'(r) \geq \frac{2-k}{r} + \frac{r}{2} + \frac{U_2}{r} - \frac{r^{2-k}}{D_2} \int_{|x|=r} \left| \sum_{i,j} w_{ij} \psi_{ij} \right|.$$

The lemma follows from this and the Cauchy-Schwarz inequality since $(\log I_2)' = \frac{2U_2}{r}$. $\square$

Lemma 4.24. Given $\lambda > 4n$, there exists ζ_0 so that if (4.2) holds for $\zeta \geq \zeta_0$, then for each $r \in (4n, 2\lambda)$ we have $U_2(r) \geq 0$ and, moreover, for $r \in (4n, 2\lambda - 1)$ there exists $c_r > 0$ so

$$(4.25) \quad \max \{U_2(s) \mid s \in [r, r+1]\} > c_r.$$

Proof. Given $r \in (4n, 2\lambda)$, the Neumann Poincaré inequality, (4.4) and (4.2) give

$$\int_{|x|<r} |\text{Hess}_w^x|^2 \leq C_r \mu^2 + C_r \int_{|x|<r} |\nabla \text{Hess}_w^x|^2 \leq \frac{C_r}{\zeta} \int_{|x|<4n} |\text{Hess}_w^x|^2 + C_r \int_{|x|<r} |\nabla \text{Hess}_w^x|^2.$$

If ζ is large enough (depending on λ), we can absorb the first term on the right to get

$$(4.26) \quad \int_{|x|<r} |\text{Hess}_w^x|^2 \leq C_r \int_{|x|<r} |\nabla \text{Hess}_w^x|^2.$$

To bound the error term in (4.18), use the absorbing inequality to get for any $\delta > 0$

$$(4.27) \quad \sum_{i,j} \int_{|x|<r} |\psi_{ij} w_{ij}| e^{-f} \leq \delta \int_{|x|<r} |\text{Hess}_w^x|^2 e^{-f} + \frac{C_r \mu^2}{\delta}.$$

Taking $\delta > 0$ small and then ζ even larger, the last two inequalities give that

$$(4.28) \quad \sum_{i,j} \int_{|x|<r} |\psi_{ij} w_{ij}| e^{-f} \leq \frac{1}{2} \int_{|x|<r} |\nabla \text{Hess}_w^x|^2 e^{-f}.$$

Using this in (4.18), we conclude that

$$(4.29) \quad D_2(r) \geq \frac{r^{2-k}}{2} e^{\frac{r^2}{4}} \int_{|x|<r} |\nabla \text{Hess}_w^x|^2 e^{-f} \geq C_r \int_{|x|<r} |\text{Hess}_w^x|^2 = C_r' \int_0^r s^{k-1} I_2(s) ds.$$

In particular, $U_2(r) \geq 0$. Moreover, we get for $r \in (4n, 2\lambda - 1)$ that

$$(4.30) \quad D_2(r+1) \geq C_r I_2(r).$$

Note that Corollary 4.9 implies that $I_2(r) > 0$. We have either $I_2(r+1) \leq 2I_2(r)$ or $2I_2(r) < I_2(r+1)$; the claim (4.25) follows from (4.30) in each case, completing the proof. $\square$

The next lemma gives r_n so that $U_2(r) \geq \frac{r^2}{3}$ when $r \geq r_n$ as long as (4.2) holds for a large ζ that depends on λ . It will be crucial that r_n does not depend on λ .

Lemma 4.31. Given $\lambda > 4n$, there exists ζ_1 so that if (4.2) holds for $\zeta \geq \zeta_1$, then for each $r \in (4n, 2\lambda)$ we have $U_2(r) > 0$ and, moreover,

$$(4.32) \quad U_2(r) \geq \frac{r^2}{3} \text{ for } r \in (r_n, 2\lambda), \text{ where } r_n \text{ depends only on } n.$$

Proof. We will choose ζ_1 even greater than the ζ_0 given by Lemma 4.24. Thus, Lemma 4.24 gives that $I_2'(r) \geq 0$ for $r \in (4n, 2\lambda)$ and (4.25) holds. Let c_{4n} be the constant from (4.25) with $r = 4n$, so that there exists $s \in (4n, 4n+1)$ with

$$(4.33) \quad U_2(s) \geq c_{4n} > 0.$$

Corollary 4.9 and the monotonicity of I_2 give C_0 so that $C_0 \zeta \mu^2 \leq I_2(r)$ and, thus,

$$(4.34) \quad \left(\frac{r^{1-k} \int_{|x|=r} \sum_{i,j} \psi_{ij}^2}{I_2} \right)^{\frac{1}{2}} \leq \frac{C_\lambda}{\zeta}.$$

Using this in Lemma 4.19 gives for $r \in (4n, 2\lambda)$ that

$$(4.35) \quad (\log U_2)'(r) \geq \frac{2-k}{r} + \frac{r}{2} - \frac{U_2}{r} - \frac{C_\lambda}{\zeta U_2}.$$

Now choose $\zeta_1 > \zeta_0$ so that $\frac{C_\lambda}{\zeta_1 c_{4n}} \leq \frac{1}{4}$. Thus, if $c_{4n} \leq U_2(r) \leq \frac{11r^2}{30}$ and $r \in (4n, 2\lambda)$, then

$$(4.36) \quad (\log U_2)'(r) \geq \frac{r^2 - 2U_2(r)}{2r} - \frac{1}{2} \geq \frac{4r - 15}{30} \geq \frac{r}{24} > 0.$$

Combining this with (4.33), we see that $U_2 \geq c_{4n}$ for $r \in (4n+1, 2\lambda)$. Arguing as in the proof of (3.47), (4.36) gives that

- There exists r_n depending on n so that there is $r_1 \in [4n+1, r_n]$ with $U_2(r_1) > \frac{r_1^2}{3}$.
- There cannot be a first $r \in (r_1, 2\lambda)$ with $U_2(r) = \frac{r^2}{3}$.

$\square$

The next lemma uses reverse Poincaré inequalities to bound the Euclidean Hessian in terms of the L^2 norm of the function on a larger set.

Lemma 4.37. If $r > 4n$ and (4.2) holds for $\zeta \geq 84$, then

$$(4.38) \quad \int_{|x|<r} |\text{Hess}_w^x|^2 e^{-f} \leq 204 \int_{|x|<r+2} w^2 e^{-f}.$$

Proof. Given a compactly supported function $\eta(x)$ and a Euclidean partial derivative w_i , we have $(\mathcal{L} + \frac{1}{2}) w_i = \psi_i$ and, thus,

$$(4.39) \quad \operatorname{div}_f (\eta^2 w_i \nabla w_i) = \eta^2 \left(|\nabla w_i|^2 - \frac{1}{2} w_i^2 + w_i \psi_i \right) + 2\eta w_i \langle \nabla \eta, \nabla w_i \rangle.$$

The divergence theorem and Cauchy-Schwarz and absorbing inequalities give

$$(4.40) \quad \int \eta^2 |\nabla w_i|^2 e^{-f} \leq \int ((4|\nabla \eta|^2 + 2\eta^2) w_i^2 + \eta^2 \psi_i^2) e^{-f}.$$

Taking $\eta \leq 1$ identically one for $|x| < r$ and cutting off linearly for $r < |x| < r+1$, we get

$$(4.41) \quad \int_{|x| < r} |\nabla w_i|^2 e^{-f} \leq \int_{|x| < r+1} |\psi_i|^2 e^{-f} + 6 \int_{|x| < r+1} |w_i|^2 e^{-f}.$$

Since (3.5) gives that $|\psi_i| \leq |\nabla \phi| + \epsilon (|w| + |\nabla w| + |\nabla w_i|)$ and $\|\nabla \phi\|_{L^2} \leq \mu$, we get

$$(4.42) \quad \int_{|x| < r} |\operatorname{Hess}_w^x|^2 e^{-f} \leq 2\mu^2 + \int_{|x| < r+1} (2w^2 + 10|\nabla w|^2) e^{-f}.$$

We will now argue similarly to bound the right-hand side of (4.42) in terms of w itself. We will again let η be a cutoff function (on a different set). We have

$$(4.43) \quad \operatorname{div}_f (\eta^2 w \nabla w) = \eta^2 (|\nabla w|^2 - w^2 + w \psi) + 2\eta w \langle \nabla w, \nabla \eta \rangle.$$

Using the absorbing inequality $|2\eta w \langle \nabla w, \nabla \eta \rangle| \leq \eta^2 |\nabla w|^2 / 2 + 2|\nabla \eta|^2 w^2$ and the Cauchy-Schwarz inequality on the $w\psi$ term, the divergence theorem gives that

$$(4.44) \quad \int \eta^2 |\nabla w|^2 e^{-f} \leq \int (3\eta^2 + 4|\nabla \eta|^2) w^2 e^{-f} + \|\eta \psi\|_{L^2}^2.$$

Equation (3.1) gives that $\psi^2 \leq 2\phi^2 + 2\epsilon(w^2 + |\nabla w|^2)$, so we get

$$(4.45) \quad \int \eta^2 |\nabla w|^2 e^{-f} \leq 2\mu^2 + 2\epsilon \int \eta^2 |\nabla w|^2 e^{-f} + \int ((3 + 2\epsilon)w^2 + |\nabla \eta|^2) w^2 e^{-f}.$$

Since $\epsilon < \frac{1}{9}$, we can absorb the $|\nabla w|^2$ term. Thus, taking $\eta \leq 1$ identically one for $|x| < r+1$ and cutting off linearly for $r+1 < |x| < r+2$, we get

$$(4.46) \quad \int_{|x| < r+1} |\nabla w|^2 e^{-f} \leq 4\mu^2 + 10 \int_{|x| < r+2} w^2 e^{-f}.$$

Combining this with (4.42) gives that

$$(4.47) \quad \int_{|x| < r} |\operatorname{Hess}_w^x|^2 e^{-f} \leq 42\mu^2 + 102 \int_{|x| < r+2} w^2 e^{-f},$$

The lemma follows since (4.2) implies that $42\mu^2 \leq \frac{1}{2} \int_{|x| < 4n} |\operatorname{Hess}_w^x|^2 e^{-f}$. $\square$

Given a function u on the cylinder, $y \in \mathbf{R}^k$, and $\lambda \in \mathbf{R}$, let $\Psi_{\lambda, u, y}$ be the norm squared of the projection of u on the λ eigenspace of Δ_θ on the sphere $x = y$. Let B_R^k be the ball in $\mathbf{R}^k$.

Lemma 4.48. Given $\lambda \in \mathbf{R}$, there exists C depending on λ, k, n so that

$$(4.49) \quad \int_{|x| < R} u^2 \leq \int_{B_R^k} \Psi_{\lambda, u, x} + C R^2 \int_{|x| < R} |\nabla^{\mathbf{R}^k} u|^2 + C \int_{|x| < R} \{((\mathcal{L} + \lambda)u)^2 + |\operatorname{Hess}_u^x|^2\}$$

Proof. Since $\mathcal{L} + \lambda = (\Delta_\theta + \lambda) + \mathcal{L}_{\mathbf{R}^k}$, we get for each $y \in B_R^k$ that

$$\begin{aligned} \int_{x=y} u^2 &\leq \Psi_{\lambda,u,y} + c \int_{x=y} ((\Delta_\theta + \lambda)u)^2 \leq \Psi_{\lambda,u,y} + 2c \int_{x=y} ((\mathcal{L} + \lambda)u)^2 + 2c \int_{x=y} (\mathcal{L}_{\mathbf{R}^k}u)^2 \\ (4.50) \quad &\leq \Psi_{\lambda,u,y} + 2c \int_{x=y} ((\mathcal{L} + \lambda)u)^2 + 4kc \int_{x=y} |\text{Hess}_u^x|^2 + 4cR^2 \int_{x=y} |\nabla^{\mathbf{R}^k} u|^2. \end{aligned}$$

Integrating this over B_R^k gives the lemma. $\square$

The next lemma is a Poincaré inequality bounding w by Hess_w^x .

Lemma 4.51. We have

$$(4.52) \quad \int_{|x|<R} w^2 \leq C_R \int_{|x|<R+1} |\text{Hess}_w^x|^2 + C_R \mu^2$$

Proof. Lemma 4.48 with $u = w$ and $\lambda = 1$, so that $\Psi_{\lambda,u,x} \equiv 0$ and $|(\mathcal{L} + 1)w| \leq C_r \mu$, gives

$$(4.53) \quad \int_{|x|<R} w^2 \leq C R^2 \int_{|x|<R} |\nabla^{\mathbf{R}^k} w|^2 + C \int_{|x|<R} |\text{Hess}_w^x|^2 + C_R \mu^2.$$

We need to absorb the first term on the right side. Let w_i be a Euclidean derivative of w . Applying Lemma 4.48 with $u = w_i$ and $\lambda = \frac{1}{2}$ gives

$$(4.54) \quad \int_{|x|<R} w_i^2 \leq \int_{B_R^k} \Psi_{\frac{1}{2},w_i,x} + C R^2 \int_{|x|<R} |\text{Hess}_w^x|^2 + C \int_{|x|<R} |\text{Hess}_{w_i}^x|^2 + C_R \mu^2.$$

Let $\{h_j\}$ be an L^2 -orthonormal basis of $\frac{1}{2}$ -eigenfunctions for Δ_θ and define $\Psi_{ij}(x) = \int h_j(\theta) w_i(x, \theta) d\theta$. It follows that

$$(4.55) \quad \Psi_{\frac{1}{2},w_i,x} = \sum_j (\Psi_{ij}(x))^2.$$

By Lemma 4.3, $\left(\int_{|x|<r} \Psi_{ij}(x)\right)^2 \leq C_r \mu^2$. The Poincaré inequality on B_R^k gives

$$(4.56) \quad \int_{B_R^k} (\Psi_{ij}(x))^2 \leq C_r \mu^2 + C_R \int_{B_R^k} |\nabla^{\mathbf{R}^k} \Psi_{ij}|^2 \leq C_r \mu^2 + C_R \int_{|x|<R} |\text{Hess}_w^x|^2.$$

Putting this together gives

$$(4.57) \quad \int_{|x|<R} w^2 \leq C_R \int_{|x|<R} |\nabla^{\mathbf{R}^k} \text{Hess}_w^x|^2 + C_R \int_{|x|<R} |\text{Hess}_w^x|^2 + C_R \mu^2.$$

Finally, to complete the proof, we use $|\mathcal{L} \text{Hess}_w^x| \leq C_r \mu$ and the reverse Poincaré inequality to bound the $|\nabla \text{Hess}_w^x|$ term. $\square$

Proof of Proposition 4.1. We will fix λ at the end depending just on n and r_+ and then choose ζ_2 . Given $r > 4n$, Lemma 4.51 and (4.2) give

$$(4.58) \quad \int_{|x|<r} w^2 \leq C'_r \int_{|x|<r+1} |\text{Hess}_w^x|^2 + C_r \mu^2 \leq C_r \int_{4n<|x|<r+1} |\text{Hess}_w^x|^2.$$

Let r_n be given by Lemma 4.31, so that $U_2(r) \geq \frac{r^2}{3}$ for $r \geq r_n$. If $r_n \leq r$ and $r+1 \leq s \leq \lambda-2$, then (4.58) and Lemma 4.31 give that

$$(4.59) \quad \int_{|x|<r} w^2 \leq C_r I_2(r+1) \leq C_r I_2(s) e^{-\frac{s^2}{3}}.$$

On the other hand, Lemma 4.37 gives that

$$(4.60) \quad \int_0^s t^{k-1} I_2(t) e^{-\frac{t^2}{4}} dt = \int_{|x|<s} |\text{Hess}_w^x|^2 e^{-f} \leq 204 \int_{|x|<s+2} w^2 e^{-f}.$$

It follows that we can choose r_a , depending just on n , so that $2 \int_{|x|<9n} w^2 e^{-f} \leq \int_{|x|<r_a} w^2 e^{-f}$. In particular, the first part of (3.22) holds for any $r_1 \geq r_a$.

Let I_0 and U_0 be the quantities I and U for w . We repeat the argument starting from $r = \max\{r_a, r_+\}$ using $U_2 \geq \frac{r^2}{3}$ to force I_0 to grow. For λ large, depending on n and r_+ , this gives $r_1 \in (\max\{r_a, r_+\}, \lambda)$ with $U_0(r_1) \geq \frac{r^2}{16}$ (we could do this for any rate below $\frac{r^2}{3}$). Finally, choose $\zeta_2 > 84$ larger than the ζ_1 from Lemma 4.31 with this λ . $\square$

5. PROVING THE ESTIMATE FOR RESCALED MCF

We will now prove (2.20) and, thus, complete the proof of Theorem 0.4. From now on, $\Sigma_t \subset \mathbf{R}^{n+1}$ is a rescaled MCF with $\lambda(\Sigma_t) < \infty$ and Σ_t converges as $t \rightarrow \infty$ to a cylinder $C = \frac{\mathbf{S}^{n-k}}{\sqrt{2(n-k)}} \times \mathbf{R}^k$. The sequence δ_j is defined in (2.3).

Proposition 5.1. (2.20) holds.

Proof. We will assume that $k \geq 1$ as the case $k = 0$ follows similarly, but much more easily. Let $\bar{\beta} < 1$ be given by Proposition 2.4. The proposition will follow once we show

$$(5.2) \quad \sup_{t \in [j, j+1]} \sup_{B_{2n} \cap \Sigma_t} |\Pi_{j+1}(\nabla \bar{H})| \leq C \delta_j^{\bar{\beta}},$$

where C does not depend on j .

We next explain how the parameters will be chosen. First, since $\bar{\beta} < 1$, we can choose $\nu, \beta < 1$ so that $\bar{\beta} < \nu\beta$. Next, given this ν , Theorem 3.8 gives $\bar{\epsilon} > 0$. Finally, we choose the constant $\epsilon_1 > 0$ in Proposition 2.12 to ensure that (3.1) holds with $\bar{\epsilon}$.

Proposition 2.12 with ϵ_1 and $\beta \in (\bar{\beta}, 1)$ as above gives constants R_j, C and cylinders $\mathcal{C}_j$ so that $B_{R_j} \cap \Sigma_t$ is a graph over $\mathcal{C}_{j+1}$ of a function w for each $t \in [j, j+1]$. Moreover, (1), (2) and (4) in Proposition 2.12 give (3.1)–(3.6) with $\epsilon < \bar{\epsilon}$ and $\mu = C \delta_j^{\bar{\beta}}$. Theorem 3.8 now applies with our choice of $\nu < 1$ above. Thus, we get a constant C_ν and function

$$(5.3) \quad \tilde{w} = \sum_i a_i (x_i^2 - 2) + \sum_{i < k} a_{ik} x_i x_k + \sum_k x_k h_k(\theta),$$

where a_i, a_{ik} are constants and each $h_k(\theta)$ is a $\frac{1}{2}$ -eigenfunction for Δ_θ , and we have

$$(5.4) \quad \sup_{|x| \leq 3n} |w - \tilde{w}| \leq C_\nu \mu^\nu = C \delta_j^{\beta\nu}.$$

The $\mathbf{R}^k$ unit vector fields ∂_{x_i} on $\mathcal{C}_{j+1}$ push forward to vector fields on Σ_t that we still denote ∂_{x_i} . Since $\{|x| \leq 3n\} \cap \Sigma_t$ is the graph over $\mathcal{C}_{j+1}$ of w with $\|w\|_{C^2(|x| \leq 3n)}^2 \leq C \delta_j^{\bar{\beta}}$, it follows

that $|\nabla \bar{H}| \leq C \delta_j^{\frac{\beta}{2}}$ and $|\Pi_{j+1}(\partial_{x_i})| \leq C \delta_j^{\frac{\beta}{2}}$ on $|x| \leq 3n$. Therefore, (5.2) follows from

$$(5.5) \quad \sup_{|x| \leq 3n} |\nabla_{\theta} \bar{H}| \leq C \delta_j^{\frac{\beta}{2}},$$

where $\bar{H}$ is now regarded as a function on $\mathcal{C}_{j+1}$ itself. It remains to establish (5.5).

The mean curvature $\bar{H}$ of the graph of w is given at each point explicitly as a function of w , ∇w and Hess_w ; see corollary A.30 in [CM2]. We can write this as the first order part (in $w, \nabla w, \text{Hess}_w$) plus a quadratic remainder

$$(5.6) \quad \bar{H} = H_{\mathcal{C}} + \left(\Delta_{\theta} + \Delta_x + \frac{1}{2} \right) w + O(w^2).$$

Here $O(w^2)$ is a term that depends at least quadratically on $w, \nabla w, \text{Hess}_w$ and the constant $H_{\mathcal{C}} = \frac{\sqrt{n-k}}{\sqrt{2}}$ is the mean curvature of $\mathcal{C}$. We will show that the θ derivative of each of the terms in (5.6) is bounded by $C \delta_j^{\frac{\beta}{2}}$. This is obvious for the constant term. It is also obvious for the quadratic term $O(w^2)$ using interpolation and $\|w\|_{C^2(|x| \leq 3n)}^2 \leq C \delta_j^{\beta}$. Similarly, since $\beta \nu > \bar{\beta}$, the estimate (5.4) and interpolation give that

$$(5.7) \quad \left| \nabla \left(\Delta_{\theta} + \Delta_x + \frac{1}{2} \right) (w - \tilde{w}) \right| \leq C \delta_j^{\frac{\beta}{2}}.$$

The proposition follows from the above since applying the linearized operator to $\tilde{w}$ gives

$$(5.8) \quad \nabla_{\theta} \left(\Delta_{\theta} + \Delta_x + \frac{1}{2} \right) \tilde{w} = \nabla_{\theta} \sum_k \left(\Delta_{\theta} + \Delta_x + \frac{1}{2} \right) x_k h_k = 0.$$

□

REFERENCES

- [AAG] S. Altschuler, S. Angenent, and Y. Giga, *Mean curvature flow through singularities for surfaces of rotation*. J. Geom. Anal. 5(3), 293–358 (1995).
- [An] B. Andrews, *Noncollapsing in mean-convex mean curvature flow*. G&T 16 (2012), no. 3, 1413–1418.
- [AV] S. Angenent and J. J. L. Velázquez, *Degenerate neckpinches in mean curvature flow*. J. Reine Angew. Math. 482 (1997), 15–66.
- [A] V. I. Arnold, *Arnold's problems*. Translated and revised edition of the 2000 Russian original. Springer-Verlag, Berlin; PHASIS, Moscow, 2004.
- [Be] J. Bernstein, *Asymptotic structure of almost eigenfunctions of drift Laplacians on conical ends*, preprint.
- [B] S. Brendle, *Embedded self-similar shrinkers of genus 0*, Annals of Math. 183, 715–728 (2016).
- [ChGG] Y.G. Chen, Y. Giga, and S. Goto, *Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations*. J. Differential Geom. 33 (1991), no. 3, 749–786.
- [CM1] T.H. Colding and W.P. Minicozzi II, *Generic mean curvature flow I; generic singularities*, Annals of Math., 175 (2012), 755–833.
- [CM2] ———, *Uniqueness of blowups and Łojasiewicz inequalities*, Annals of Math., 182 (2015), 221–285.
- [CM3] ———, *Level set method for motion by mean curvature*, Notices of the AMS, (2016) Vol. 63, No. 10, 1148–1153.
- [CM4] ———, *The singular set of mean curvature flow with generic singularities*, Inventiones Math., 204 (2) (2016), 443–471.
- [CM5] ———, *Differentiability of the arrival time*, CPAM, Vol. LXIX, (2016) 2349–2363.
- [CM6] ———, *Regularity of the level set flow*, CPAM, to appear.
- [CM7] ———, *Sharp frequency bounds for eigenfunctions of the Ornstein-Uhlenbeck operator*, preprint.

- [CM8] ———, *Łojasiewicz inequalities and applications*. Surveys in differential geometry 2014. Regularity and evolution of nonlinear equations, 6382, Surv. Differ. Geom., 19, Int. Press, Somerville, MA, 2015.
- [CM9] ———, *Rigidity of cylinders for Ricci flow*, preprint.
- [ES] L.C. Evans and J. Spruck, *Motion of level sets by mean curvature I*, JDG 33 (1991) 635–681.
- [GK] Z. Gang and D. Knopf, *Universality in mean curvature flow neckpinches*. Duke Math. J. 164 (2015), no. 12, 2341–2406.
- [G] P. Goldstein, *Gradient flow of a harmonic function in $\mathbf{R}^3$* . J. Diff. Eq. 247 (2009), no. 9, 2517–2557.
- [HaK] R. Haslhofer and B. Kleiner, *Mean curvature flow of mean convex hypersurfaces*, CPAM, (2017), 511–546.
- [Hö] L. Hörmander, *On the division of distributions by polynomials*, Ark. Mat. 3 (1958) 555–568.
- [H1] G. Huisken, *Asymptotic behavior for singularities of the mean curvature flow*. JDG 31 (1990), 285–299.
- [H2] ———, *Local and global behaviour of hypersurfaces moving by mean curvature*. Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990), 175–191, Proc. Sympos. Pure Math., 54, Part 1, AMS, Providence, RI, 1993.
- [HS1] G. Huisken and C. Sinestrari, *Mean curvature flow singularities for mean convex surfaces*, Calc. Var. Partial Differ. Equ. 8 (1999), 1–14.
- [HS2] ———, *Convexity estimates for mean curvature flow and singularities of mean convex surfaces*, Acta Math. 183 (1999), no. 1, 45–70.
- [I] T. Ilmanen, *Generalized Flow of Sets by Mean Curvature on a Manifold*, Indiana J. Math. 41 (1992) 671–705.
- [Ku] N. Kuiper, *On gradient curves of an analytic function near a critical point*, preprint IHES, 1991.
- [KMP] K. Kurdyka, T. Mostowski, and A. Parusinski, *Proof of the gradient conjecture of R. Thom*, Annals of Math., 152 (2000), 763–792.
- [KS] R.V. Kohn and S. Serfaty, *A deterministic control based approach to motion by curvature*. CPAM. 59 (2006), no. 3, 344–407.
- [L1] S. Łojasiewicz, *Ensembles semi-analytiques*, IHES notes (1965).
- [L2] ———, *Sur la geometrie semi- et sous-analytique*, Ann. Inst. Fourier 43 (1993), 1575–1595.
- [L3] ———, *Sur le problème de la division*, Studia Math. 18 (1959) 87–136.
- [L4] ———, *Une propriété topologique des sous-ensembles analytiques réels*, Les Équations aux Dérivées Partielles (1963) 87–89, CNRS, Paris.
- [M] T. Miura, *An example of a mean-convex mean curvature flow developing infinitely many singular epochs*. J. Geom. Anal. 26 (2016), no. 4, 3019–3026.
- [OsSe] S. Osher and J.A. Sethian, *Fronts propagating with curvature-dependent speed: algorithms based on Hamilton-Jacobi formulations*. J. Comput. Phys. 79 (1988), no. 1, 12–49.
- [S] N. Sesum, *Rate of convergence of the mean curvature flow*. CPAM 61 (2008), no. 4, 464–485.
- [Si] L. Simon, *Theorems on Regularity and Singularity of Energy Minimizing Maps*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 1996.
- [T] R. Thom, *Problemes rencontres dans mon parcours mathematique: un bilan*, Publ. Math. IHES 70 (1989), 200–214.
- [Wa] X.-J. Wang, *Convex solutions to the mean curvature flow*. Ann. of Math. (2) 173 (2011), 1185–1239.
- [W1] B. White, *The size of the singular set in mean curvature flow of mean-convex sets*. JAMS 13 (2000), 665–695.
- [W2] ———, *The nature of singularities in mean curvature flow of mean-convex sets*. JAMS 16 (2003), 123–138.
- [W3] ———, *Evolution of curves and surfaces by mean curvature*, in Proceedings of the International Congress of Mathematicians, Vol. I (Beijing, 2002), Higher Ed. Press, Beijing, 2002, 525–538.

MIT, DEPT. OF MATH., 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139-4307.

E-mail address: colding@math.mit.edu and minicozz@math.mit.edu