

Within-host priority effects and epidemic timing determine outbreak severity in coinfecting populations

Authors: Patrick A. Clay^{1,2*}, Meghan A. Duffy¹, Volker H.W. Rudolf²

¹University of Michigan, Department of Ecology and Evolutionary Biology

²Rice University, Biosciences Department

pclay@umich.edu, duffymeg@umich.edu, volker.rudolf@rice.edu

*Corresponding Author

Running Title: priority effects and outbreak severity

Key words: Coinfection, Within-Host, Between-Host, Priority effects, Epidemic, Daphnia, Pathogen, Community Ecology

ABSTRACT-

Coinfections of hosts by multiple pathogen species are ubiquitous, but predicting their impact on disease remains challenging. Interactions between coinfecting pathogens within hosts can alter pathogen transmission, with the impact on transmission typically dependent on the relative arrival order of pathogens within hosts (within-host priority effects). However, it is unclear how these within-host priority effects influence multi-pathogen epidemics, particularly when the arrival order of pathogens at the host population scale varies. Here we combined models and experiments with zooplankton and their naturally co-occurring fungal and bacterial pathogens to examine how within-host priority effects influence multi-pathogen epidemics. Epidemiological models parameterized with within-host priority effects measured at the single host scale predicted that advancing the start date of bacterial epidemics relative to fungal epidemics would decrease mean bacterial prevalence in a multi-pathogen setting, while models without within-host priority effects predicted the opposite effect. We tested these predictions with experimental multi-pathogen epidemics. Empirical dynamics matched predictions from the model including within-host priority effects, providing evidence that within-host priority effects influenced epidemic dynamics. Overall, within-host priority effects may be a key element of predicting multi-pathogen epidemic dynamics in the future, particularly as shifting disease phenology alters the order of infection within hosts.

Introduction

Epidemics of infectious diseases can strongly influence natural, agricultural and human populations. They can result in rapid degradation of host health, regulate host population dynamics, and can even put species at risk of extinction [1–3]. Predicting the dynamics and severity of epidemics ahead of time is therefore imperative for timely public health interventions, such as limiting trade of infected livestock, or immunizing at-risk human and wildlife populations. However, the vast majority of host populations are coinfecting by multiple pathogen species [4–7], and epidemics of different pathogens often alter one another’s trajectories. For instance, influenza epidemics historically increased population susceptibility to pneumonia [8]. Thus, understanding how coinfecting pathogens interact at the individual and population scales is essential to predicting epidemic dynamics and mitigating epidemic severity.

Co-occurring pathogens can increase or decrease one another’s epidemic severity by interacting at the individual host scale. Pathogens can interact by competing for resources such as nutrients or body tissue, by directly interfering with or facilitating one another (e.g. by producing bacteriocins), and/or by indirectly interacting via the immune system [9–12]. These within-host interactions alter host susceptibility, pathogen transmission rates, and the duration of infections, thus determining the rate at which pathogens transmit through a host population [13,14]. Therefore, the severity of an epidemic may be difficult to predict without knowing the identities of coinfecting pathogens and how they interact within hosts.

While the many ways that pathogens interact within hosts are well documented [15], scaling these interactions up to predict epidemic severity has proven challenging because within-host interactions are influenced by the order of pathogen arrival at multiple spatial scales [16]. Within-host interactions have a deterministic component based on fitness asymmetry between

pathogens, but can be modified by the order in which pathogens infect hosts, both in multi-strain coinfections [17] and in multi-species coinfections [18]. These within-host priority effects can alter both the strength and direction of within-host interactions [19], and can alter disease risk [20]. Furthermore, the first pathogen to spread through a host population during a multi-pathogen epidemic is likely to be the first pathogen to infect individual hosts [16,21]. Thus, differences in the arrival order of co-occurring epidemics can alter within-host interactions, and thereby could change epidemic dynamics. However, large enough inter-specific asymmetry in pathogen fitness within and across hosts could override context dependent within-host priority effects. In this scenario, infection order may not be important and will simply add “noise” to the dynamics of epidemics. Given that variation in timing of epidemics is common because pathogens typically respond differentially to seasonal forcings [22], it is important to understand whether within-host priority effects can alter epidemic severity, and whether measuring them will allow us to better predict future epidemic dynamics.

One challenge to understanding the role of within-host priority effects in determining epidemic severity is that within-host priority effects may themselves be context dependent. Many studies have documented within-host priority effects experimentally [18,23–26], but these interactions are often measured in isolated hosts under ideal conditions that do not reflect the complex, stressful, and ever-changing environmental conditions hosts and pathogens experience during epidemics. For instance, host and pathogen populations at the start, peak, or end of an epidemic experience very different resource conditions [27], population densities [2], and age/stage structures [28] that could all modify within-host interactions and thus alter within-host priority effects. Consequently, measuring within-host interactions and priority effects

experimentally in isolated hosts may not allow us to understand their role in determining epidemic dynamics at the host population scale.

To help fill these conceptual gaps, we asked (1) how does relative epidemic start date interact with within-host priority effects to alter epidemic dynamics? And (2) does measuring within-host priority effects at the individual host scale improve our ability to predict multi-pathogen epidemics? To answer these questions, we used individuals and populations of zooplankton coinfecting with naturally co-occurring fungal and bacterial pathogens as a model system. We ran predictive epidemic models parameterized with or without within-host priority effects, and then tested the predictions of these models with experimental multi-pathogen epidemics. Predictive models with or without within-host priority effects showed clear qualitative differences in the impact of epidemic arrival order on epidemic dynamics, and our experimental epidemics qualitatively matched patterns predicted only by our within-host priority effect model. Together, these results indicate that within-host priority effects interact with relative epidemic start date to alter multi-pathogen epidemic dynamics in our system, and that we might better predict multi-pathogen epidemics by taking within-host priority effects into account.

Methods

We predict that epidemic timing will alter epidemic dynamics by creating feedback loops mediated by within-host priority effects. The first pathogen to enter a host population will most likely be the first pathogen to enter individual hosts in the first host generation of a multi-pathogen epidemic. In later generations, epidemic arrival order will alter the order of infection at the single host scale indirectly. The arrival order in the first generation of hosts will determine the transmission of pathogens from those hosts, and thus the force of infection of both pathogens. When two pathogens circulate in a host population, the pathogen with the higher force of

infection will be more likely to infect hosts first [19]. Thus, the relative start dates of co-occurring epidemics can indirectly alter the arrival order of pathogens at the single host scale several generations after the start of the epidemics.

It is intractable to measure these priority effect mediated feedback loops directly, as we would need to measure both the infection order of pathogens in each individual host, and the transmission of pathogens from those hosts, during an epidemic. Instead, we built predictive epidemic models parameterized with experiments conducted at the single host scale and ran these models with and without within-host priority mediated feedback loops. This approach allowed us to understand how these feedback loops influence multi-pathogen dynamics and isolate within-host priority effects from effects arising simply from co-infections. We then tested the qualitative predictions made by our epidemic models by running experimental multi-pathogen epidemics in the lab.

Study system

We built models to predict epidemic dynamics in the zooplankton species *Daphnia dentifera* (henceforth *zooplankton*). This zooplankton species is common in lakes in the midwestern United States [29]. In late summer, zooplankton populations commonly host pathogens, including the fungus *Metschnikowia bicuspidata* and the bacterium *Pasteuria ramosa* (henceforth *fungus* and *bacterium*). While lakes differ in their pathogen communities, the fungus and bacterium do co-occur in some lakes. Zooplankton ingest infectious spores of both pathogens while filtering for algae. After ingestion, spores replicate until host death, at which point they escape into the water column until they are ingested by a new host [30]. For this study, we used a clonal zooplankton line (“Mid37”) and pathogen isolates which had been grown in lab for several years (“Standard” fungus and “G/18” bacterium).

132 To predict the impact of within-host interactions and within-host priority effects on
133 epidemic severity, and to test whether we can predict epidemic severity by measuring within-
134 host interactions experimentally, we ran multi-pathogen epidemic simulations parameterized
135 with experimental measurements of within-host priority effects. We compared models which (i)
136 included within-host interactions, (ii) included within-host interactions but not within-host
137 priority effects, and (iii) did not include any within-host interactions (Figure 1). We include a no
138 within-host interaction model to determine whether coinfection and epidemic arrival order alter
139 epidemic dynamics purely by altering host population dynamics.

140 We parameterized our model with within-host priority effects measured and fully
141 described in [31]. Age-controlled hosts were singly infected, simultaneously coinfecting, or
142 sequentially infected, with either the bacterium or the fungus arriving four days after the other
143 pathogen. Comparing glms with heteroscedasticity using the glht function in R [32], we
144 determined that fungal spore yield (number of infectious spores in the host at death) was
145 significantly lower in coinfecting hosts the fungus arrived first in compared to other coinfecting
146 host classes. Bacterial spore load was significantly lower in coinfecting hosts than singly infected
147 hosts in all cases. To parameterize the within-host priority effects for this study, we used the
148 R2jags package in R [33] to find Bayesian estimates of the mean values of fungal spores
149 (normally distributed) and bacterial spores (poisson distributed) from each infected host class
150 (Figure 1). We used a Bayesian approach to more efficiently carry forward uncertainty through
151 nested layers of modelling (parameters estimated from single hosts were used to estimate
152 parameters in single-pathogen epidemics, which were used to make predictions about multi-
153 pathogen epidemics).





Figure 1: Pathogen fitness in all models: those that contained within-host interactions/priority effects, corresponding to empirical data (“full model”), those that did not contain within-host priority effects (“no priority effects”), and those that did not contain any within-host interactions (“no interactions”). The fitness of pathogens was measured as average number of infectious spores released from hosts at death [31]. Points indicate relative spore yield of pathogens in coinfecting hosts compared to mean spore yield of singly infected hosts when the fungus arrived first (Fun), when pathogens arrived at the same time (Simul), or when the bacterium arrived first (Bac). Blue points represent the bacterium, yellow points represent the fungus. The gray shading indicates the likelihood that a pathogen will be competitively excluded, ranging from 0 in the white to 1 in the darkest gray (see supplement for calculation). Bars indicate 95% of posterior estimate for each point. Raw data and relevant methods can be found in Clay et al. 2019.

Here, we present our full model that includes both within-host interactions and within-host priority effects (Figure 2). See methods supplement for equations describing all models, and for discussion of model assumptions. We model epidemics as a discrete time SI system of

170 environmentally transmitted bacterial and fungal pathogens, with environmental spore pools B
 171 and F . Hosts may be susceptible (S), singly infected (I_B or I_F), coinfecting simultaneously (C_{sim}),
 172 sequentially infected with the fungus first (C_{FB}), or sequentially infected with the bacterium first
 173 (C_{BF}). Further, upon coinfection, pathogens may competitively exclude one another. Hosts in
 174 class $X_{i,sim}$, $X_{i,FB}$, or $X_{i,BF}$ were previously coinfecting, but pathogen i competitively excluded its
 175 competitor within the host. Once a pathogen has been competitively excluded, it cannot re-infect
 176 a host.



177

178 **Figure 2:** Model flowchart, highlighting most likely transmission pathways for each multi-
 179 pathogen experimental treatment, e.g. in our “Bacterium First” treatment, we predict susceptible
 180 individuals will first become singly infected by the bacterium (I_B), and then coinfecting (C_{BF}).

(Note, however, that all highlighted transmission pathways can occur in all treatments). Blue is used to indicate state variables and parameters related to the bacterium, including hosts singly infected by the bacterium (I_B), bacterial infection (solid blue arrow), bacterial spore release (dashed blue arrow), and bacterium competitively excluding the fungus within hosts (dotted blue arrow). Similarly, yellow represents state variables and parameters related to the fungus, including hosts singly infected by the fungus, fungal infection, fungal spore release, and the fungus competitively excluding the fungus within hosts. Green represents coinfecting hosts and simultaneous coinfection. Multi-colored lines mean that hosts can move back and forth between two classes. Line thickness represents relative rate, e.g. bacterial spore release is less from coinfecting individuals than from singly infected individuals, and thus the bacterial spore release line is thinner from coinfecting hosts. Note that competitive exclusion happens instantaneously upon coinfection, thus individuals move directly from singly infected individuals to cleared individuals in model equations.

To differentiate between simultaneous coinfections (infections occur on the same day) and sequential coinfections (infections happen on different days), we model dynamics in discrete time, with a time step of one day. Dynamics of the susceptible class are given by

$$S_{t+1} = S_t + \overbrace{bS_{t-j} \left(1 - \frac{N_{t-j}}{K}\right)}^{\text{Births}} - \overbrace{(\theta_{B,t} + \theta_{F,t} - \theta_{B,t}\theta_{F,t})S_t}^{\text{Infection}} - \overbrace{\widehat{d_S}S_t}^{\text{Deaths}} \quad eq. 1$$

Where b is the number of offspring/host/day at low population density, K is host carrying capacity, and N_t is total population size at time t . In experimental, single-pathogen epidemics (described below), we observed no developing offspring in the brood chamber of any infected zooplankton. Thus, infected hosts do not give birth in our model. Hosts must go through a juvenile stage, lasting j days, before they are counted as part of the susceptible, adult population.

Susceptible hosts become singly infected by feeding on spores in the water column. Thus, $\theta_{i,t}$ is the proportion of hosts to become infected by pathogen i at time t , given by

$$\theta_{B,t} = f_t \mu_B B_t \quad eq. 2$$

$$\theta_{F,t} = f_t \mu_F F_t \quad eq. 3$$

Where f_t is the total proportion of spores in the environment eaten by each host at time step t , and μ_i is the per spore infectivity of pathogen i (eq. 2 and 3 were established to be mass action processes based on previous experiments, see supplement). We assume that pathogens do not alter host susceptibility to one another, as shown in [31]. In these prior experiments, we found that the order of infection of the fungal and bacterial pathogens only altered spore yield. We found no evidence of protection or immune suppression (prior exposure by one pathogen did not reduce or increase the likelihood of infection by the other pathogen). Thus, we assume that infection likelihood is not altered by the infection status of the host. The proportion of susceptible hosts to become simultaneously coinfectd at time t is $\theta_{B,t} \theta_{F,t}$, and the proportion of susceptible hosts to become singly infected by pathogen i is the proportion of susceptible hosts to become infected by pathogen i , minus the proportion of those hosts to also become simultaneously coinfectd. At no point in our simulations does $\theta_{B,t}$ or $\theta_{F,t}$ become greater than one. Finally d_S is the rate of intrinsic host mortality.

Hosts singly infected by the fungus may become coinfectd upon ingesting bacterial spores, and vice versa. Upon coinfection, the probability of remaining coinfectd until death with no competitive exclusion is ω_j , where j is the coinfection class. $\lambda_{B,j}$ is the probability that the fungus will competitively exclude the bacterium, and $\lambda_{F,j}$ is the probability that the bacterium will competitively exclude the fungus. See supplement for calculation of competitive exclusion

probabilities. After the fungus is cleared from infected hosts, the hosts can become reinfected by the fungus, returning to their original coinfecting class. Upon reinfection, the bacterium and the fungus may once again competitively exclude one another, with the original probabilities associated with that coinfecting class. We assume that once the bacterium is excluded by the fungus, it cannot reinfect hosts, as the time between fungal infection and host death is relatively short. Finally, since fungal infection reduces host lifespan [31], all hosts infected by the fungus die at an increased rate (d_F), with their death rate reverting to d_S if the bacterium excludes the fungus. Thus, changes in the numbers of infected and coinfecting hosts are then given by

$$I_{B,t+1} = I_{B,t} + \overbrace{(\theta_{B,t}(1 - \theta_{F,t}))S_t}^{\text{Infection}} - \overbrace{\theta_{F,t}I_{B,t}}^{\text{Coinfection}} - \overbrace{d_S I_{B,t}}^{\text{Deaths}} \quad eq. 4$$

$$I_{F,t+1} = I_{F,t} + \overbrace{(\theta_{F,t}(1 - \theta_{B,t}))S_t}^{\text{Infection}} - \overbrace{\theta_{B,t}I_{F,t}}^{\text{Coinfection}} - \overbrace{d_F I_{F,t}}^{\text{Deaths}} \quad eq. 5$$

$$C_{sim,t+1} = C_{sim,t} + \overbrace{\theta_{B,t}\theta_{F,t}\omega_{sim}S_t}^{\text{Coinfection}} + \overbrace{\theta_{F,t}\omega_{sim}X_{B,sim,t}}^{\text{Reinfection}} - \overbrace{d_F C_{sim,t}}^{\text{Deaths}} \quad eq. 6$$

$$X_{B,sim,t+1} = X_{B,sim,t} + \overbrace{\theta_{B,t}\theta_{F,t}\lambda_{F,sim}S_t}^{\text{Competitive Exclusion}} - \overbrace{\theta_{F,t}X_{B,sim,t}(1 - \lambda_{F,sim})}^{\text{Reinfection}} - \overbrace{d_S X_{B,sim,t}}^{\text{Deaths}} \quad eq. 7$$

$$X_{F,sim,t+1} = X_{F,sim,t} + \overbrace{\theta_{B,t}\theta_{F,t}\lambda_{B,sim}S_t}^{\text{Competitive Exclusion}} + \overbrace{\theta_{F,t}\lambda_{B,sim}X_{B,sim,t}}^{\text{Reinfection}} - \overbrace{d_F X_{F,sim,t}}^{\text{Deaths}} \quad eq. 8$$

$$C_{BF,t+1} = C_{F,t} + \overbrace{\theta_{F,t}\omega_{BF}I_{B,t}}^{\text{Coinfection}} + \overbrace{\theta_{F,t}\omega_{BF}X_{B,BF,t}}^{\text{Reinfection}} - \overbrace{d_F C_{F,t}}^{\text{Deaths}} \quad eq. 9$$

$$X_{B,BF,t+1} = X_{B,BF,t} + \overbrace{\theta_{F,t}\lambda_{F,BF}I_{B,t}}^{\text{Competitive Exclusion}} - \overbrace{\theta_{F,t}X_{B,BF,t}(1 - \lambda_{F,BF})}^{\text{Reinfection}} - \overbrace{d_S X_{B,BF,t}}^{\text{Deaths}} \quad eq. 10$$

$$X_{F,BF,t+1} = X_{F,BF,t} + \overbrace{\theta_{F,t}\lambda_{B,BF}I_{B,t}}^{\text{Competitive Exclusion}} + \overbrace{\theta_{F,t}\lambda_{B,BF}X_{B,BF,t}}^{\text{Reinfection}} - \overbrace{d_F X_{F,BF,t}}^{\text{Deaths}} \quad eq. 11$$

$$241 \quad C_{FB,t+1} = C_{FB,t} + \overbrace{\theta_{B,t}\omega_{FB}I_{F,t}}^{\text{Coinfection}} + \overbrace{\theta_{F,t}\omega_{FB}X_{B,FB,t}}^{\text{Reinfection}} - \overbrace{d_F C_{FB,t}}^{\text{Deaths}} \quad eq. 12$$

$$242 \quad X_{B,FB,t+1} = X_{B,FB,t} + \overbrace{\theta_{B,t}\lambda_{F,FB}I_{F,t}}^{\text{Competitive Exclusion}} - \overbrace{\theta_{F,t}X_{B,FB,t}(1 - \lambda_{F,FB})}^{\text{Reinfection}} - \overbrace{d_S X_{B,FB,t}}^{\text{Deaths}} \quad eq. 13$$

$$243 \quad X_{F,FB,t+1} = X_{F,FB,t} + \overbrace{\theta_{B,t}\lambda_{B,FB}I_{F,t}}^{\text{Competitive Exclusion}} + \overbrace{\theta_{F,t}\lambda_{B,FB}X_{B,FB,t}}^{\text{Reinfection}} - \overbrace{d_F X_{F,FB,t}}^{\text{Deaths}} \quad eq. 14$$

244

245 Infected hosts transmit spores into environmental spore pools upon death, whose
 246 dynamics are given by

$$247 \quad B_{t+1} = B_t +$$

$$248 \quad \overbrace{d_S \beta_{B(B)}(I_{B,t} + X_{B,BF,t} + X_{B,FB,t} + X_{B,sim,t}) + d_F \beta_{B(BF)}C_{BF,t} + d_F \beta_{B(FB)}C_{FB,t} + d_F \beta_{B(sim)}C_{sim,t}}^{\text{Spore Release}}$$

$$249 \quad - \overbrace{\alpha_B B_t}^{\text{Loss}} - \overbrace{f_t \mu_B B_t N_t}^{\text{Uptake}} \quad eq. 15$$

250 And

$$251 \quad F_{t+1} = F_t +$$

$$252 \quad \overbrace{d_F \beta_{F(F)}(I_{F,t} + X_{F,BF,t} + X_{F,FB,t} + X_{F,sim,t}) + d_F \beta_{F(BF)}C_{BF,t} + d_F \beta_{F(FB)}C_{FB,t} + d_F \beta_{F(sim)}C_{sim,t}}^{\text{Spore Release}}$$

$$253 \quad - \overbrace{\alpha_F F_t}^{\text{Loss}} - \overbrace{f_t F_t N_t}^{\text{Uptake}} \quad eq. 16$$

254 Where $\beta_{i(j)}$ represents of the number of spores i released from host class j . Thus, all hosts that
 255 are infected by a given pathogen add spores of that pathogen to the environment upon death. We
 256 assume that if a pathogen is competitively excluded from a host, the host's spore yield reverts to

that of a singly infected host. Spores have a loss rate (α_i), which represents spore degradation and spores moving out of the system (e.g., due to settling), and fungal spores are removed from the environment by host feeding (f) [34]. Bacterial spores, alternatively, can survive passage through the host gut, and thus are not removed by host feeding unless they actively infect the host [35].

Models were parameterized with a combination of three experiments. First, we measured lifespan and birthrates of uninfected hosts with a life table experiment (supplement). We then measured pathogen fitness (infectious spores released upon host death) in singly infected, sequentially infected, and simultaneously coinfecting hosts [31]. Finally, we estimated carrying capacity and spore degradation from the uninfected and singly infected treatments of our experimental epidemics (supplement). All parameters were estimated with the R2jags package in R [33], in order to create Bayesian posterior distributions of parameter value probabilities. See supplement for full methods and all parameter values (Table S1).

To predict epidemic dynamics under shifting epidemic arrival order, we ran our model under conditions where fungal epidemics started a week before bacterial epidemics, where epidemics started simultaneously, and where bacterial epidemics started a week before fungal epidemics, matching our experimental treatments (Table 1). We simulated each scenario 1,000 times, each time drawing parameters from posterior distributions of Bayesian parameter estimates (supplement). Within-host interactions and priority effects in this system primarily impact infectious spore production [31]. Thus, for our model with no within-host interactions, we set the spore release rate from all coinfecting hosts equal to the spore release rate from singly infected hosts. For our model which included within-host interactions but no within-host priority

effects, we set the spore release rate of sequentially coinfecting hosts equal to that of simultaneously coinfecting hosts (Figure 1, see supplement for exact equations for each model).

Testing model predictions

To test our epidemic model predictions empirically, we ran experimental multi-pathogen epidemics. We seeded 1 liter beakers of filtered pond water with 100 adult zooplankton each on day 1 of the experiment. On day 2, we homogenized singly infected zooplankton to create infectious spore slurries, and seeded beakers with either 190 fungal spores/ml, 1000 bacterial spores/ml, 190 fungal spores/ml and 1000 bacterial spores/ml, or a control dosage of ground up uninfected *Daphnia*. We repeated this procedure on day 9 to create populations with no epidemics, single-pathogen epidemics, or multi-pathogen epidemics with varying epidemic arrival orders (Table 1), with 8 replicate populations per treatment.

Treatment No.	Treatment Day 2	Treatment day 9
1	Control	Control
2	Fungus	Control
3	Bacterium	Control
4	Fungus/Bacterium	Control
5	Fungus	Bacterium
6	Bacterium	Fungus

Table 1: Treatments in host population scale experiments.

To monitor host-pathogen dynamics, we subsampled 1/10th of each zooplankton population every 5 days, filtering out *Daphnia* and returning water to each replicate after sampling. Juveniles were counted and discarded. Adult individuals were homogenized and

infectious spores inside each adult were identified. We then categorized adult zooplankton as uninfected, singly infected by the fungus, singly infected by the bacterium, or coinfecting. We ran the experiment for 67 days, ending when the majority of infected populations had gone extinct. We fed each replicate 20,000 *Ankistrodesmus sp.* cells per ml 4 days a week (Mon, Tue, Thu, Fri).

Comparing models predictions and experimental epidemics

To compare model predictions and empirical epidemics, we focused on treatment differences in three key metrics that characterize different aspect of epidemics: mean prevalence, epidemic length (time between first observed and last observed infected host), and integrated prevalence (the summed prevalence across all sampling dates). For each metric and pathogen, we first calculated significant pairwise differences between single pathogen epidemics and each multi-pathogen epidemic treatment. Second, for each pathogen we calculated the relationship between a metric and the relative timing of infection (day of fungal spore introduction – day of bacterial spore introduction) among our coinfecting treatments. Residuals of the mean fell under gamma or normal distributions for all variables in each treatment. Thus, all comparisons were performed with the generalized linear model (glm) function in R [36].

Our models predicted that within-host priority effects resulted in testable differences in key aspect of epidemics (see results) across arrival treatments. Thus, if our no interaction model (Figure 1, “no interactions”) correctly predicts epidemic dynamics, then within-host interactions are not important drivers of epidemic dynamics in our system. If our no within-host priority effects model (Figure 1, “no priority effects”) correctly predicts epidemic dynamics, then within-host interactions, but not within-host priority effects are important drivers of epidemic dynamics in our system. If only our full model (Figure 1, “full model”) predicts epidemic

dynamics, then we can conclude that within-host priority effects alter epidemic dynamics, and that by measuring them we can accurately predict epidemic dynamics a priori. If no models correctly predict epidemic dynamics, then measurements of within-host interactions made experimentally do not necessarily represent within-host interactions that occur during an epidemic.

Results

Model Predictions

Adding within-host priority effects to our predictive models altered the predicted impact of epidemic arrival order on bacterial epidemic dynamics. In all models, the presence of a fungal epidemic reduced the integrated prevalence, mean prevalence, and epidemic length of bacterial epidemics (Figure S2). All models also predicted the same impact of epidemic arrival order on bacterial epidemic length, with bacterial epidemics lasting longer when bacterial epidemics start before fungal epidemics. However, models disagreed on the impact of epidemic arrival order on mean prevalence. Models that did not include within-host interactions predicted no effect of epidemic arrival order on mean prevalence. Models that accounted for within-host interactions but not within-host priority effects predicted that mean prevalence would be higher when the bacterial epidemic started before the fungal epidemic than in other treatments. In sharp contrast, our full model (including within-host priority effects) predicted the opposite result: as bacterial epidemic start date moved later, and fungal epidemic start date moved earlier, mean bacterial prevalence should decrease (Figures 3, S2).




Figure 3: Empirical estimates of the impact of epidemic arrival order on mean bacterial prevalence and epidemic length match predictions made by our full model. The Y-axis shows the mean prevalence (A-D) and length (E-H) of bacterial epidemics in experiments (A,E) and in models when the fungal epidemic precedes the bacterial epidemic (“Fungus”), and when the bacterial epidemic precedes the fungal epidemic (“Bacterium”). Points for empirical data represent means with standard error across replicates, while model data is represented by boxplots of 1,000 model runs.

All models predicted no change in fungal epidemic dynamics due to coinfection or epidemic arrival order (Figure S3).

Empirical single pathogen epidemic dynamics

Both pathogens spread readily through host populations. In single pathogen epidemics the fungus and the bacterium both reached 100% prevalence in all replicates, and maintained themselves in host populations until host populations went extinct (Figure 4,S3). Bacterial

epidemics reached 100% prevalence roughly 40 days after epidemic start date and stayed close to 100% prevalence until populations went extinct (Figure 4E). Fungal epidemics spread much faster and first peaked near 100% prevalence at day 11 of our experiment, then proceeded to cycle through multiple epidemic peaks of varying intensity (Figure 4F).

How do coinfection and epidemic timing alter empirical epidemic severity?

Coinfection and relative epidemic start date interacted to change epidemic dynamics. We saw coinfecting individuals at a high rate in all coinfecting treatments (Figure S4). Coinfection reduced the total size of all bacterial epidemics (integrated prevalence of infected individuals over time, hereafter referred to as integrated prevalence: only bacterium vs. fungus first $p=0.017$, only bacterium vs. simultaneous epidemics $p=0.046$, only bacterium vs. bacterium first $p=0.048$, Figure 3). The integrated prevalence did not significantly change with shifting epidemic start date ($p=1.0$). However, as bacterial epidemics shifted earlier and fungal epidemics shifted later, bacterial epidemics became longer ($p=0.036$), and had lower average prevalence ($p=0.0015$, Figure 3A,E, Figure 4B-D). Overall these results indicate that relative epidemic start date in our system shaped how epidemics were distributed over time, though not their overall size. Fungal epidemics were not significantly different in integrated prevalence, mean prevalence, or length across any treatments (Table S2; fungal epidemics appear longer in singly infected populations in Figure 3F due to a single replicate).




Figure 4: Empirical bacterial epidemics (blue) changed with coinfection and epidemic arrival order, while fungal epidemics (yellow) did not. Y-axis shows prevalence (proportion) of hosts infected over the course of the epidemic. Prevalence includes both singly infected and coinfecting hosts. Lines represent mean prevalence across non-extinct replicates, with ribbons representing 95% confidence intervals at each time point. Coinfection reduced the integrated prevalence of all bacterial epidemics (B vs. E $p=0.017$, C vs. E $p=0.046$, D vs. E $p=0.048$). The integrated prevalence did not significantly change with shifting epidemic start date ($p=1.0$). As bacterial epidemics shifted earlier and fungal epidemics shifted later, bacterial epidemics became longer

($p=0.036$), and had lower average prevalence ($p=0.0015$). Fungal epidemics were not significantly different between treatments.

Testing model predictions with empirical dynamics

The dynamics of our experimental epidemics best matched predictions made by our full model. In cases where models agreed with one another (predicting that coinfection would reduce the integrated size of bacterial epidemics, and that bacterial epidemics would become shorter as the bacterial epidemic shifted later and the fungal epidemic shifted earlier), empirical epidemics matched the predictions of all models (Figure 3, E-H, Figure 4, B-E). However, as the bacterial epidemic shifted later and the fungal epidemic shifted earlier, our empirical data rejected the predictions of our no priority effect models that the mean prevalence of bacterial epidemics would increase or stay the same (Figure 3, C-D, Figure 4, B-E), instead supporting the prediction of our full model that the mean prevalence of bacterial epidemics should decrease.

Discussion

In natural systems, the arrival order of pathogens within a host can determine the fitness and transmission of those pathogens. However, it is still unclear how these within-host priority effects can alter disease dynamics during multi-pathogen epidemics. Our study shows that in our system pathogen arrival order at the within-host scale interacts with pathogen arrival order at the host population scale to determine multi-pathogen epidemic dynamics. Specifically, we show that coinfection reduced epidemic size, while the relative start dates of co-occurring epidemics determined how epidemics were distributed over time. Predictive epidemic models only accurately captured qualitative multi-pathogen dynamics if they included within-host priority effects measured at the single host scale, indicating that within-host priority effects can be a

mechanism through which disease phenology alters multi-pathogen epidemics. Taken together, this evidence suggests that measuring within-host priority effects may increase our ability to predict multi-pathogen epidemics in the future, particular under shifting disease phenology.

Mechanisms of single pathogen epidemics dynamics

The two pathogens in our study both created large epidemics, but exhibited different temporal dynamics. In our single pathogen treatments, the sustained high prevalence of the bacterium (Figure 3E) is likely driven by its low probability of degradation (Table S1), and because this bacterium can remain infectious after passing through host digestive tracts [35]. Therefore, the bacterial spore load in the environment, and thus the probability of infection, could remain high throughout our experiment. The cycling pattern in our fungal epidemics (Figure 3F) is due to the synchronized infection and mortality of infected host cohorts. After fungal spores are introduced to the environment, hosts will remove most of them via ingestion. A proportion of those hosts will become infected, and then all die roughly 10 to 12 days later due to fungus induced mortality, releasing spores back into the environment, and restarting the cycle.

Mechanisms of multi-pathogen epidemics dynamics

Entering host populations before the fungus could have increased mean bacterial prevalence as the bacterium had time to spread without fungal interference, or could have decreased mean bacterial prevalence as the bacterium had the lowest fitness within coinfecting hosts when it arrived first (Figure 1). Ultimately, we found that bacterial prevalence was lowest when bacterial epidemics had a head start (Figure 3,4). Our epidemic models tease apart the mechanics of how relative epidemic start date alters mean prevalence of the bacterium. Mean bacterial prevalence shrinks when the bacterium arrives first compared to when the fungus

arrives first because there is a high probability of competitive exclusion for the bacterium when it infects hosts before the fungus (Figure 1). Thus, as competitive exclusion of the bacterium increases due to within-host priority effects, the mean prevalence of the bacterium during the epidemic decreases. Therefore, in this system, within-host priority effects are strong enough for epidemic start date to change epidemic severity.

Fungal epidemic dynamics showed no change between treatments, despite our fungal pathogen experiencing stronger within-host priority effects than the bacterium (Figure 1). However, this pattern may be driven by epidemic saturation: fungal pathogens reached 100% prevalence very quickly in all single pathogen treatments. This high prevalence makes it difficult to detect if epidemics facilitate one another, as epidemic prevalence cannot exceed 100%. Furthermore, it suggests that the spore load of the fungus in the environment may have been far greater than necessary to infect all hosts. Under such conditions, fluctuations in infectious spore yields from co-infected hosts would have no effect on epidemic dynamics. This indicates that for pathogens with very high transmission and prevalence, such as some gut helminths in wildlife populations [37], their dynamics may not be impacted by interactions with other pathogens, or by relative epidemic timing.

Ultimately, how epidemic timing and within-host priority effects altered experimental epidemics depended on the type of within-host priority effects we observed. In our system, pathogens had lowest fitness when they infect coinfecting hosts first (Figure 1). For the fungus, we hypothesize coinfection with the bacterium facilitates the fungus, as the bacterium causes gigantism via castration [38], allowing for hosts to contain more spores. However, if the fungus arrives first, it may disrupt bacterial castration, removing the facilitative effect. Further experiments are needed to confirm this hypothesis. For a system where within-host priority

effects take a different form, e.g. where pathogens have a fitness advantage from prior residency in hosts [17], the impact of shifting epidemic phenology will be different than that seen in our system. However, our results provide evidence that within-host priority effects can scale up by interacting with disease phenology, and may do so in other systems.

Role of within host interactions

Overall, this study experimentally demonstrates that within-host priority effects can alter epidemic dynamics in our system by shifting pathogen fitness in coinfecting hosts. Previous studies have demonstrated that sequential infections can alter epidemic severity if early arriving pathogens increase or decrease host susceptibility to further infection [11,16,20,39]. For instance, endemic infection by helminths makes tuberculosis invasion easier as helminths increase host susceptibility to microparasites [11]. Or, in certain grass hosts, the first arriving pathogen changes the likelihood of coinfection by all other pathogens in the system [16]. Previous theory further shows that within-host priority effects may alter disease dynamics not only by changing the likelihood of infection, but also by altering competitive outcomes within hosts, altering the ability of pathogens to spread through a previously infected population [40,41]. Our study adds to this body of work by providing experimental evidence that within-host competitive outcomes driven by infection order scale up to alter epidemic dynamics. These results may apply even when epidemics do not have separate start dates. This is because hosts will usually be infected first by the pathogen with the highest infection risk in a given system [16,21]. Thus, our results emphasize that we should examine the role of within-host priority effects in driving disease dynamics whenever a host population is infected by multiple pathogens.

Though within-host priority effects are important, they are not the only mechanism through which epidemic timing impacts epidemic dynamics in our system. We found that

between-host interactions, rather than within-host interactions, determined variation in epidemic length in our experiments. Between-host interactions occur when pathogens decrease the population size of susceptible hosts, decreasing the transmission of density-dependent pathogens [42]. In our experiments, bacterial epidemics were longer when the bacterium entered the population first than when the fungus entered the population first. This is because coinfection greatly decreased population density and drove host populations to extinction (Figure S5C-E). In total, the length of a bacterial epidemic was proportional to the time between when bacterium entered the host population, and when the host population went extinct. Similarly, in all of our models, bacterial epidemics are longest when the bacterium enters the population first (Figure 2,4), because epidemics start when the bacterium enters the host population, and end when the host population goes extinct. Thus, within-host interactions are not needed to predict some aspects of how epidemic timing alters epidemic patterns. However, the strength of between-host interactions increases with pathogen prevalence [39], which depends in part on within-host interactions. Therefore, within-host interactions and between host interactions likely feed back on one another to determine epidemic severity.

Making Predictions

Our results demonstrate that we can predict qualitative changes in multi-pathogen epidemic dynamics in our system. Specifically, our models parameterized with within-host priority effects correctly predicted that epidemic start date would alter the length and mean prevalence of bacterial epidemics (Figure 3,S2) but would not alter fungal epidemics (Figure S3). Our models did not predict quantitative epidemic dynamics (e.g. our full model could not predict pathogen prevalence at a specific time point in the epidemic, figure S6) because *Daphnia* *sp.* population dynamics are highly stochastic, follow boom and bust cycles, and depend on

asymmetric competition between different daphnia age classes [43,44]. We chose a simpler model that would not capture *Daphnia* population dynamics, but would better show the impact of within-host priority effects at the host-population scale. Further, qualitative disease predictions are highly valuable in identifying areas of concern, such as identifying where shifts in disease phenology will lead to an increase in epidemic severity.

Importantly, our model predicted the qualitative impact of within-host interactions *a priori*. Previous multi-pathogen epidemic models have accurately predicted epidemic patterns, but these models were parameterized by either observing or inferring within-host interactions during the epidemic the models were predicting [8,45,46]. The value of models with post-hoc parameterization lies in predicting multi-pathogen epidemics that occur in common spatial and temporal combinations. For instance, Abu-Raddad et al. (2006) predicted the dynamics of endemic strains of HIV and malaria in Kenya by observing endemic HIV/malaria interactions in Malawi. However, for novel pathogen combinations or arrival orders, experimentally measured within-host interactions must be used to predict how pathogen interactions will alter epidemic dynamics *a priori*. Our results indicate that this approach can accurately predict qualitative metrics, but also that we must take into account how within-host interactions will translate from a lab setting to a field setting, where pathogens face fundamentally different conditions [47]. For instance, for our models to make accurate predictions, we needed to assume that within-host competition between pathogens was higher during experimental epidemics than during the experiments performed on isolated hosts used to parameterize our predictive models (an assumption with empirical support, see supplement). How within-host interactions will change from field to lab settings will depend on within-host mechanisms. For instance, hosts rarely compete for resources during experimental measurements of within-host interactions

[25,26,31,48,49]. Within-host competition for resources may then increase when hosts compete for host resources in the field [50], but apparent competition via the immune system may decrease when hosts compete for resources [51]. Thus, to predict how changing ecology will qualitatively change multi-pathogen epidemics before they occur, researchers should examine the mechanisms underlying within-host priority effects, rather than simply measuring the impact of sequential infection on pathogen fitness in a lab setting.

Conclusion

Infectious disease research now acknowledges that many pathogens usually circulate within host populations [52]. Epidemics of these pathogens occur in response to environmental conditions (temperature, rainfall), host population dynamics (aggregation, dispersal, foraging), or within-host conditions (seasonal weakening of immune systems)[22]. These variables are constantly changing, and as long as pathogens respond differentially to environmental or host cues, then epidemics will continue to fluctuate in their relative timing, and thus severity. Further, we expect to see shifts in environmental variables and host population dynamics under future climate conditions, potentially creating novel timing conditions of co-occurring epidemics. If we do not take both within-host interactions and within-host priority effects into account, we may become increasingly unable to predict epidemic severity in the future.

Author Contributions: PAC and VHWR conceptualized and designed this experiment. MAD provided pathogens and hosts. PAC analyzed empirical data and built epidemiological models. All authors contributed to writing of manuscript.

Data Accessibility: When this manuscript is accepted, all data and code will be uploaded to dryad.

Acknowledgments: We thank Rebecca Bilich, Katie Hunsberger, Laura Lopez, Clara Shaw, Karana Wickens, and Khadijah Payne for lab support. We further thank Tom E.X. Miller and Amy Dunham for helpful discussions on experimental planning and data analysis. Work was supported by NSF grants DEB-1748729 and DEB-1601353.

References

1. Roelke-Parker, M. E. et al. 1996 A canine distemper virus epidemic in Serengeti lions (*Panthera leo*). *Nature*; Feb 1 **379**.
2. Hudson, P. J. 1998 Prevention of Population Cycles by Parasite Removal. *Science* (80-.). **282**, 2256–2258. (doi:10.1126/science.282.5397.2256)
3. Ezenwa, V. O. & Jolles, A. E. 2015 Opposite effects of anthelmintic treatment on microbial infection at individual versus population scales. *Science* (80-.). **347**, 175–177. (doi:10.1126/science.1261714)
4. Petney, T. N. & Andrews, R. H. 1998 Multiparasite communities in animals and humans: frequency, structure and pathogenic significance. *Int. J. Parasitol.* **28**, 377–393. (doi:10.1016/S0020-7519(97)00189-6)
5. Brogden, K. A., Guthmiller, J. M. & Taylor, C. E. 2005 Human polymicrobial infections. *Lancet* **365**, 253–255. (doi:10.1016/S0140-6736(05)17745-9)
6. Balmer, O. & Tanner, M. 2011 Prevalence and implications of multiple-strain infections. *Lancet Infect. Dis.* **11**, 868–878. (doi:10.1016/S1473-3099(11)70241-9)
7. Cox, F. E. G. 2011 Concomitant infections, parasites and immune responses. *Parasitology* **122**, S23–S38. (doi:10.1017/S003118200001698X)

- 557 8. Shrestha, S., Foxman, B., Weinberger, D. M., Steiner, C., Viboud, C. & Rohani, P. 2013
558 Identifying the interaction between influenza and pneumococcal pneumonia using
559 incidence data. *Sci. Transl. Med.* **5**, 191ra84. (doi:10.1126/scitranslmed.3005982)
- 560 9. Graham, A. L. 2008 Ecological rules governing helminth-microparasite coinfection. *Proc.*
561 *Natl. Acad. Sci. U. S. A.* **105**, 566–70. (doi:10.1073/pnas.0707221105)
- 562 10. Inglis, R. F., Gardner, A., Cornelis, P. & Buckling, A. 2009 Spite and virulence in the
563 bacterium *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 5703–7.
564 (doi:10.1073/pnas.0810850106)
- 565 11. Ezenwa, V. O., Etienne, R. S., Luikart, G., Beja-Pereira, A. & Jolles, A. E. 2010 Hidden
566 consequences of living in a wormy world: nematode-induced immune suppression
567 facilitates tuberculosis invasion in African buffalo. *Am. Nat.* **176**, 613–24.
568 (doi:10.1086/656496)
- 569 12. Wale, N., Sim, D. G. & Read, A. F. 2017 A nutrient mediates intraspecific competition
570 between rodent malaria parasites in vivo. *Proc. R. Soc. London B Biol. Sci.* **284**.
- 571 13. Ezenwa, V. O. & Jolles, A. E. 2011 From Host Immunity to Pathogen Invasion: The
572 Effects of Helminth Coinfection on the Dynamics of Microparasites. *Integr. Comp. Biol.*
573 **51**, 540–551. (doi:10.1093/icb/ucr058)
- 574 14. Clark, P. R., Ward, W. T., Lang, S. A., Saghbini, A. & Kristan, D. M. 2013 Order of
575 Inoculation during *Heligmosomoides bakeri* and *Hymenolepis microstoma* Coinfection
576 Alters Parasite Life History and Host Responses. *Pathog. (Basel, Switzerland)* **2**, 130–52.
577 (doi:10.3390/pathogens2010130)

- 578 15. Fenton, A., Lamb, T. & Graham, A. L. 2008 Optimality analysis of Th1/Th2 immune
579 responses during microparasite-macroparasite co-infection, with epidemiological
580 feedbacks. *Parasitology* **135**, 841–853. (doi:10.1017/S0031182008000310)
- 581 16. Halliday, F. W., Umbanhowar, J. & Mitchell, C. E. 2017 Interactions among symbionts
582 operate across scales to influence parasite epidemics. *Ecol. Lett.* **20**, 1285–1294.
583 (doi:10.1111/ele.12825)
- 584 17. de Roode, J. C., Helinski, M. E. H., Anwar, M. A. & Read, A. F. 2005 Dynamics of
585 multiple infection and within-host competition in genetically diverse malaria infections.
586 *Am. Nat.* **166**, 531–42. (doi:10.1086/491659)
- 587 18. Hoverman, J. T., Hoyer, B. J. & Johnson, P. T. J. 2013 Does timing matter? How priority
588 effects influence the outcome of parasite interactions within hosts. *Oecologia* **173**, 1471–
589 80. (doi:10.1007/s00442-013-2692-x)
- 590 19. Clay, P. A., Cortez, M. H., Duffy, M. A. & Rudolf, V. H. W. 2019 Priority effects within
591 coinfecting hosts can drive unexpected population-scale patterns of parasite prevalence.
592 *Oikos* **128**, 571–583. (doi:10.1111/oik.05937)
- 593 20. Halliday, F. W., Umbanhowar, J. & Mitchell, C. E. 2018 A host immune hormone
594 modifies parasite species interactions and epidemics: insights from a field manipulation.
595 *Proc. R. Soc. B Biol. Sci.* **285**, 20182075. (doi:10.1098/rspb.2018.2075)
- 596 21. Fukami, T. 2015 Historical Contingency in Community Assembly: Integrating Niches,
597 Species Pools, and Priority Effects. *Annu. Rev. Ecol. Evol. Syst.* **46**, 1–23.
598 (doi:10.1146/annurev-ecolsys-110411-160340)

- 599 22. Altizer, S., Dobson, A., Hosseini, P., Hudson, P., Pascual, M. & Rohani, P. 2006
600 Seasonality and the dynamics of infectious diseases. *Ecol. Lett.* **9**, 467–484.
601 (doi:10.1111/j.1461-0248.2005.00879.x)
- 602 23. Al-Naimi, F. A., Garrett, K. A. & Bockus, W. W. 2005 Competition, facilitation, and
603 niche differentiation in two foliar pathogens. *Oecologia* **143**, 449–457.
604 (doi:10.1007/s00442-004-1814-x)
- 605 24. Lohr, J. N., Yin, M. & Wolinska, J. 2010 Prior residency does not always pay off--co-
606 infections in *Daphnia*. *Parasitology* **137**, 1493–500. (doi:10.1017/S0031182010000296)
- 607 25. Doublet, V., Natsopoulou, M. E., Zschiesche, L. & Paxton, R. J. 2015 Within-host
608 competition among the honey bees pathogens *Nosema ceranae* and Deformed wing virus
609 is asymmetric and to the disadvantage of the virus. *J. Invertebr. Pathol.* **124**, 31–34.
610 (doi:10.1016/J.JIP.2014.10.007)
- 611 26. Rynkiewicz, E. C., Brown, J., Tufts, D. M., Huang, C.-I., Kampen, H., Bent, S. J., Fish, D.
612 & Diuk-Wasser, M. A. 2017 Closely-related *Borrelia burgdorferi* (sensu stricto) strains
613 exhibit similar fitness in single infections and asymmetric competition in multiple
614 infections. *Parasit. Vectors* **10**, 64. (doi:10.1186/s13071-016-1964-9)
- 615 27. Pedersen, A. B. & Greives, T. J. 2008 The interaction of parasites and resources cause
616 crashes in a wild mouse population. *J. Anim. Ecol.* **77**, 370–377. (doi:10.1111/j.1365-
617 2656.2007.01321.x)
- 618 28. Hite, J. L., Penczykowski, R. M., Shocket, M. S., Strauss, A. T., Orlando, P. A., Duffy, M.
619 A., Cáceres, C. E. & Hall, S. R. 2016 Parasites destabilize host populations by shifting
620 stage-structured interactions. *Ecology*. **97**.

- 621 29. Tessier, A. J. & Woodruff, P. 2002 Trading off the ability to exploit rich versus poor food
622 quality. *Ecol. Lett.* **5**, 685–692. (doi:10.1046/j.1461-0248.2002.00373.x)
- 623 30. Auld, S. K. J. R., Hall, S. R., Housley Ochs, J., Sebastian, M. & Duffy, M. A. 2014
624 Predators and patterns of within-host growth can mediate both among-host competition
625 and evolution of transmission potential of parasites. *Am. Nat.* , S77-90.
626 (doi:10.1086/676927)
- 627 31. Clay, P. A., Dhir, K., Rudolf, V. H. W. & Duffy, M. A. 2019 Within-host priority effects
628 systematically alter pathogen coexistence. *Am. Nat.* **193**, 187–199. (doi:10.1086/701126)
- 629 32. Hothorn, T., Bretz, F. & Westfall, P. 2008 Simultaneous Inference in General Parametric
630 Models. *Biometrical J.* **50**, 346–363. (doi:10.1002/bimj.200810425)
- 631 33. Su, Y.-S. & Yajima, M. 2015 R2jags: Using R to Run ‘JAGS’. R package version 0.5-7.
632 <https://CRAN.R-project.org/package=R2jags>.
- 633 34. Civitello, D. J., Pearsall, S., Duffy, M. A. & Hall, S. R. 2013 Parasite consumption and
634 host interference can inhibit disease spread in dense populations. *Ecol. Lett.* **16**, 626–634.
635 (doi:10.1111/ele.12089)
- 636 35. King, K. C., Auld, S. K. J. R., Wilson, P. J., James, J. & Little, T. J. 2013 The bacterial
637 parasite *Pasteuria ramosa* is not killed if it fails to infect: implications for coevolution.
638 *Ecol. Evol.* **3**, 197–203. (doi:10.1002/ece3.438)
- 639 36. R Core Team (2017) 2017 R: A language and environment for statistical computing. *R*
640 *Found. Stat. Comput.*
- 641 37. Lotz, J. M. & Font, W. 1994 Excess Positive Associations in Communities of Intestinal

- 642 Helminths of Bats: A Refined Null Hypothesis and a Test of the Facilitation Hypothesis.
643 *J. Parasitology* **80**, 398–413.
- 644 38. Ebert, D. 1996 Development, life cycle, ultrastructure and phylogenetic position of
645 *Pasteuria ramosa* Metchnikoff 1888: Rediscovery of an obligate endoparasite of *Daphnia*
646 *magna* Straus. *Philos. Trans. R. Soc. B Biol. Sci.* **351**, 1689–1701.
647 (doi:10.1098/rstb.1996.0151)
- 648 39. Vasco, D. A., Wearing, H. J. & Rohani, P. 2007 Tracking the dynamics of pathogen
649 interactions: modeling ecological and immune-mediated processes in a two-pathogen
650 single-host system. *J. Theor. Biol.* **245**, 9–25. (doi:10.1016/j.jtbi.2006.08.015)
- 651 40. Clay, P. A., Dhir, K., Rudolf, V. H. W. & Duffy, M. A. 2018 Within-Host Priority Effects
652 Systematically Alter Pathogen Coexistence. *Am. Nat.* , 000–000. (doi:10.1086/701126)
- 653 41. Bushman, M., Antia, R., Udhayakumar, V. & de Roode, J. C. 2018 Within-host
654 competition can delay evolution of drug resistance in malaria. *PLOS Biol.* **16**, e2005712.
655 (doi:10.1371/journal.pbio.2005712)
- 656 42. Rohani, P., Green, C. J., Mantilla-Beniers, N. B. & Grenfell, B. T. 2003 Ecological
657 interference between fatal diseases. *Nature* **422**, 885–8. (doi:10.1038/nature01542)
- 658 43. Ananthasubramaniam, B., Nisbet, R. M., Nelson, W. A., McCauley, E. & Gurney, W. S.
659 C. 2011 Stochastic growth reduces population fluctuations in *Daphnia* –algal systems.
660 *Ecology* **92**, 362–372. (doi:10.1890/09-2346.1)
- 661 44. Grover, J. P., McKee, D., Young, S., Godfray, H. C. J. & Turchin, P. 2000 Periodic
662 dynamics in daphnia populations: Biological interactions and external forcings. *Ecology*

- 663 **81**, 2781–2798. (doi:10.1890/0012-9658(2000)081[2781:PDIDPB]2.0.CO;2)
- 664 45. Abu-Raddad, L. J., Patnaik, P. & Kublin, J. G. 2006 Dual infection with HIV and malaria
665 fuels the spread of both diseases in sub-Saharan Africa. *Science* **314**, 1603–6.
666 (doi:10.1126/science.1132338)
- 667 46. Abu-Raddad, L. J., van der Ventel, B. I. S. & Ferguson, N. M. 2008 Interactions of
668 Multiple Strain Pathogen Diseases in the Presence of Coinfection, Cross Immunity, and
669 Arbitrary Strain Diversity. *Phys. Rev. Lett.* **100**, 168102.
670 (doi:10.1103/PhysRevLett.100.168102)
- 671 47. Laine, A.-L. 2011 Context-dependent effects of induced resistance under co-infection in a
672 plant-pathogen interaction. *Evol. Appl.* **4**, 696–707. (doi:10.1111/j.1752-
673 4571.2011.00194.x)
- 674 48. Natsopoulou, M. E., McMahon, D. P., Doublet, V., Bryden, J. & Paxton, R. J. 2015
675 Interspecific competition in honeybee intracellular gut parasites is asymmetric and favours
676 the spread of an emerging infectious disease. *Proc. Biol. Sci.* **282**, 20141896.
677 (doi:10.1098/rspb.2014.1896)
- 678 49. Marchetto, K. M. & Power, A. G. 2017 Coinfection Timing Drives Host Population
679 Dynamics through Changes in Virulence. *Am. Nat.* , 000–000. (doi:10.1086/695316)
- 680 50. Budischak, S. A., Sakamoto, K., Megow, L. C., Cummings, K. R., Urban, J. F. & Ezenwa,
681 V. O. 2015 Resource limitation alters the consequences of co-infection for both hosts and
682 parasites. *Int. J. Parasitol.* **45**, 455–463. (doi:10.1016/j.ijpara.2015.02.005)
- 683 51. Hite, J. L. & Cressler, C. E. 2018 Resource-driven changes to host population stability

684 alter the evolution of virulence and transmission. *Philos. Trans. R. Soc. B Biol. Sci.* **373**,
685 20170087. (doi:10.1098/rstb.2017.0087)

686 52. Johnson, P. T. J., de Roode, J. C. & Fenton, A. 2015 Why infectious disease research
687 needs community ecology. *Science* (80-.). **349**, 1259504–1259504.
688 (doi:10.1126/science.1259504)

689