



# Generalized minimum distance functions and algebraic invariants of Geramita ideals



Susan M. Cooper<sup>a</sup>, Alexandra Seceleanu<sup>b,\*</sup>,  
Ştefan O. Tohăneanu<sup>c</sup>, Maria Vaz Pinto<sup>d</sup>, Rafael H. Villarreal<sup>e</sup>

<sup>a</sup> Department of Mathematics, 420 Machray Hall, 186 Dysart Road University of Manitoba, Winnipeg, MB R3T 2N2 Canada

<sup>b</sup> Department of Mathematics, 203 Avery Hall, Lincoln, NE 68588, United States of America

<sup>c</sup> Department of Mathematics, University of Idaho, 875 Perimeter Drive, MS 1103, Moscow, ID 83844-1103, United States of America

<sup>d</sup> Departamento de Matemática, Instituto Superior Técnico, Universidade de Lisboa, Avenida Roiviso País, 1, 1049-001 Lisboa, Portugal

<sup>e</sup> Departamento de Matemáticas, Centro de Investigación y de Estudios Avanzados del IPN, Apartado Postal 14-740, 07000 Mexico City, CDMX, Mexico

## ARTICLE INFO

### Article history:

Received 13 January 2019

Received in revised form 19 July 2019

Accepted 10 September 2019

Available online 23 September 2019

### MSC:

primary 13P25

secondary 14G50, 94B27, 11T71

### Keywords:

Generalized minimum distance

Fat points

Linear codes

Basic parameters

Degree

Hilbert function

## ABSTRACT

Motivated by notions from coding theory, we study the generalized minimum distance (GMD) function  $\delta_I(d, r)$  of a graded ideal  $I$  in a polynomial ring over an arbitrary field using commutative algebraic methods. It is shown that  $\delta_I$  is non-decreasing as a function of  $r$  and non-increasing as a function of  $d$ . For vanishing ideals over finite fields, we show that  $\delta_I$  is strictly decreasing as a function of  $d$  until it stabilizes. We also study algebraic invariants of Geramita ideals. Those ideals are graded, unmixed, 1-dimensional and their associated primes are generated by linear forms. We also examine GMD functions of complete intersections and show some special cases of two conjectures of Tohăneanu–Van Tuyl and Eisenbud–Green–Harris.

Published by Elsevier Inc.

\* Corresponding author.

E-mail addresses: [susan.cooper@umanitoba.ca](mailto:susan.cooper@umanitoba.ca) (S.M. Cooper), [aseceleanu@unl.edu](mailto:aseceleanu@unl.edu) (A. Seceleanu), [tohaneanu@uidaho.edu](mailto:tohaneanu@uidaho.edu) (Ş.O. Tohăneanu), [vazpinto@math.ist.utl.pt](mailto:vazpinto@math.ist.utl.pt) (M.V. Pinto), [vila@math.cinvestav.mx](mailto:vila@math.cinvestav.mx) (R.H. Villarreal).

## 1. Introduction

Let  $K$  be any field, and let  $C$  be a linear code that is the image of some  $K$ -linear map  $K^s \rightarrow K^n$ . Suppose  $G$  is the  $s \times n$  matrix representing this map with respect to some chosen bases and assume that  $G$  has no zero columns. By definition, the *minimum (Hamming) distance* of  $C$  is

$$\delta(C) := \min\{\text{wt}(\mathbf{v}) \mid \mathbf{v} \in C \setminus \{\mathbf{0}\}\},$$

where for any vector  $\mathbf{w} \in K^n$ , the *weight* of  $\mathbf{w}$ , denoted  $\text{wt}(\mathbf{w})$ , is the number of nonzero entries in  $\mathbf{w}$ . More generally, for  $1 \leq r \leq \dim_K(C)$ , the  $r$ -th *generalized Hamming distance*, denoted  $\delta_r(C)$ , is defined as follows. For any subcode, i.e., linear subspace,  $D \subseteq C$  define the support of  $D$  to be

$$\chi(D) := \{i \mid \text{there exists } (x_1, \dots, x_n) \in D \text{ with } x_i \neq 0\}.$$

Then the  $r$ -th generalized Hamming distance of  $C$  is

$$\delta_r(C) := \min_{D \subseteq C, \dim D = r} |\chi(D)|.$$

The *weight hierarchy* of  $C$  is the sequence  $(\delta_1(C), \dots, \delta_k(C))$ , where  $k = \dim(C)$ . Observe that  $\delta_1(C)$  equals the minimum distance  $\delta(C)$ . The study of these weights is related to trellis coding,  $t$ -resilient functions, and was motivated by some applications from cryptography [35]. It is the study of the generalized Hamming weight of a linear code that motivates our definition of a generalized minimum distance function for any graded ideal in a polynomial ring [18,20].

If the rank of  $G$  is  $s$ , then it turns out (see [35]) that

$$\delta_r(\mathcal{C}) = n - \text{hyp}_r(\mathcal{C}), \quad (1.1)$$

where  $\text{hyp}_r(\mathcal{C})$ , is the maximum number of columns of  $G$  that span an  $(s-r)$ -dimensional vector subspace of  $K^s$ . Moreover, if  $G$  also has no proportional columns then the columns of  $G$  determine the coordinates of  $n$  (projective) points in  $\mathbb{P}^{s-1}$ , not all contained in a hyperplane. Denote this set  $\mathbb{X} = \{P_1, \dots, P_n\}$  and let  $I := I(\mathbb{X}) \subset S := K[t_1, \dots, t_s]$  be the defining ideal of  $\mathbb{X}$ . We have:

- the (Krull) dimension of  $S/I$  is  $\dim(S/I) = 1$ , and the degree is  $\deg(S/I) = n$ ;

- the ideal  $I$  is given by  $I = \mathfrak{p}_1 \cap \cdots \cap \mathfrak{p}_n$ , where  $\mathfrak{p}_i$  is the vanishing ideal of the point  $P_i$ , so  $I$  is unmixed, each associated prime ideal  $\mathfrak{p}_i$  is generated by linear forms, and  $I = \sqrt{I}$ ;
- $\text{hyp}_r(\mathcal{C}) = \max_{F \in \mathcal{F}_r} \{\deg(S/(I, F))\}$ , where  $\mathcal{F}_r$  is the set of  $r$ -tuples of linear forms of  $S$  that are linearly independent. With this, we can conclude that

$$\delta_r(\mathcal{C}) = \deg(S/I) - \max_{F \in \mathcal{F}_r} \{\deg(S/(I, F))\}.$$

A similar approach can be taken for projective Reed–Muller-type codes. Let  $\mathbb{X} = \{P_1, \dots, P_n\}$  be a finite subset of  $\mathbb{P}^{s-1}$ . Let  $I := I(\mathbb{X}) \subset S = K[t_1, \dots, t_s]$ , be the defining ideal of  $\mathbb{X}$ . Via a rescaling of the homogeneous coordinates of the points  $P_i$ , we can assume that the first non-zero coordinate of each  $P_i$  is 1. Fix a degree  $d \geq 1$ . Because of the assumption on the coordinates of the  $P_i$ , there is a well-defined  $K$ -linear map given by the evaluation of the homogeneous polynomials of degree  $d$  at each point in  $\mathbb{X}$ . This map is given by

$$\text{ev}_d: S_d \rightarrow K^n, \quad f \mapsto (f(P_1), \dots, f(P_n)),$$

where  $S_d$  denotes the  $K$ -vector space of homogeneous polynomials of  $S$  of degree  $d$ . The image of  $S_d$  under  $\text{ev}_d$ , denoted by  $C_{\mathbb{X}}(d)$ , is called a *projective Reed–Muller-type code* of degree  $d$  on  $\mathbb{X}$  [5,11,15]. The *parameters* of the linear code  $C_{\mathbb{X}}(d)$  are:

- *length*:  $|\mathbb{X}| = \deg(S/I)$ ;
- *dimension*:  $\dim_K C_{\mathbb{X}}(d) = H_{\mathbb{X}}(d)$ , the Hilbert function of  $S/I$  in degree  $d$ ;
- *r-th generalized Hamming weight*:  $\delta_{\mathbb{X}}(d, r) := \delta_r(C_{\mathbb{X}}(d))$ .

By [13, Theorem 4.5] the  $r$ -th generalized Hamming weight of a projective Reed–Muller code is given by

$$\delta_{\mathbb{X}}(d, r) = \deg(S/I) - \max_{F \in \mathcal{F}_{d,r}} \{\deg(S/(I, F))\},$$

where  $\mathcal{F}_{d,r}$  the set of  $r$ -tuples of forms of degree  $d$  in  $S$  which are linearly independent over  $K$  modulo the ideal  $I$  and the maximum is taken to be 0 if  $\mathcal{F}_{d,r} = \emptyset$ .

As we can see above, the generalized Hamming weights for any linear code can be interpreted using the language of commutative algebra. Motivated by the notion of generalized Hamming weight described above and following [13] we define *generalized minimum distance (GMD) functions* for any homogeneous ideal in a polynomial ring. This allows us to extend the notion of generalized Hamming weights to codes arising from algebraic schemes, rather than just from reduced sets of points. Another advantage to formulating the notion of generalized minimum distance in the language of commutative algebra is that it allows the use of various homological invariants of graded ideals to study the possible values for these GMD functions.

Let  $S = K[t_1, \dots, t_s] = \bigoplus_{d=0}^{\infty} S_d$  be a polynomial ring over a field  $K$  with the standard grading and let  $I \neq (0)$  be a graded ideal of  $S$ . Given  $d, r \in \mathbb{N}_+$ , let  $\mathcal{F}_{d,r}$  be the set:

$$\mathcal{F}_{d,r} := \{ \{f_1, \dots, f_r\} \subset S_d \mid \bar{f}_1, \dots, \bar{f}_r \text{ are linearly independent over } K, \\ (I: (f_1, \dots, f_r)) \neq I \},$$

where  $\bar{f} = f + I$  is the class of  $f$  modulo  $I$ , and  $(I: (f_1, \dots, f_r)) = \{g \in S \mid gf_i \in I, \text{ for all } i\}$ . If necessary we denote  $\mathcal{F}_{d,r}$  by  $\mathcal{F}_{d,r}(I)$ . We denote the *degree* of  $S/I$  by  $\deg(S/I)$ .

**Definition 1.1.** Let  $I \neq (0)$  be a graded ideal of  $S$ . The function  $\delta_I: \mathbb{N}_+ \times \mathbb{N}_+ \rightarrow \mathbb{Z}$  given by

$$\delta_I(d, r) := \begin{cases} \deg(S/I) - \max\{\deg(S/(I, F)) \mid F \in \mathcal{F}_{d,r}\} & \text{if } \mathcal{F}_{d,r} \neq \emptyset, \\ \deg(S/I) & \text{if } \mathcal{F}_{d,r} = \emptyset, \end{cases}$$

is called the *generalized minimum distance function* of  $I$ , or simply the GMD function of  $I$ .

This notion recovers (Proposition 3.14) and refines the algebraic-geometric notion of degree. If  $r = 1$  one obtains the minimum distance function of  $I$  [23]. In this case we denote  $\delta_I(d, 1)$  simply by  $\delta_I(d)$  and  $\mathcal{F}_{d,r}$  by  $\mathcal{F}_d$ .

The aims of this paper are to study the behavior of  $\delta_I$ , to introduce algebraic methods to estimate this function, and to study the algebraic invariants (minimum distance function, v-number, regularity, socle degrees) of special ideals that we call Geramita ideals. Recall that an ideal  $I \subset S$  is called *unmixed* if all its associated primes have the same height; this notion is sometimes called height unmixed in the literature. We call an ideal  $I \subset S$  a *Geramita ideal* if  $I$  is an unmixed graded ideal of dimension 1 whose associated primes are generated by linear forms. Defining ideals of schemes of finite sets of points in projective space and unmixed monomial ideals of codimension 1 are examples of Geramita ideals.

The following function is closely related to  $\delta_I$  as illustrated in Eq. (1.1).

**Definition 1.2.** Let  $I$  be a graded ideal of  $S$ . The function  $\text{hyp}_I: \mathbb{N}_+ \times \mathbb{N}_+ \rightarrow \mathbb{N}$ , given by

$$\text{hyp}_I(d, r) := \begin{cases} \max\{\deg(S/(I, F)) \mid F \in \mathcal{F}_{d,r}\} & \text{if } \mathcal{F}_{d,r} \neq \emptyset, \\ 0 & \text{if } \mathcal{F}_{d,r} = \emptyset, \end{cases}$$

is called the *hyp function* of  $I$ .

If  $r = 1$ , we denote  $\text{hyp}_I(d, 1)$  by  $\text{hyp}_I(d)$ . Finding upper bounds for  $\text{hyp}_I(d, r)$  is equivalent to finding lower bounds for  $\delta_I(d, r)$ . If  $I(\mathbb{X})$  is the vanishing ideal of a finite

set  $\mathbb{X}$  of reduced projective points, then  $\text{hyp}_{I(\mathbb{X})}(d, 1)$  is the maximum number of points of  $\mathbb{X}$  contained in a hypersurface of degree  $d$  (see [32, Remarks 2.7 and 3.4]). There is a similar geometric interpretation for  $\text{hyp}_{I(\mathbb{X})}(d, r)$  [13, Lemma 3.4].

To compute  $\delta_I(d, r)$  is a difficult problem even when  $K$  is a finite field and  $r = 1$ . However, we show that a generalized footprint function, which is more computationally tractable, gives lower bounds for  $\delta_I(d, r)$ . Fix a monomial order  $\prec$  on  $S$ . Let  $\text{in}_\prec(I)$  be the initial ideal of  $I$  and let  $\Delta_\prec(I)$  be the *footprint* of  $S/I$ , consisting of all the *standard monomials* of  $S/I$  with respect to  $\prec$ . The footprint of  $S/I$  is also called the *Gröbner escalier* of  $I$ . Given integers  $d, r \geq 1$ , let  $\mathcal{M}_{\prec, d, r}$  be the set of all subsets  $M$  of  $\Delta_\prec(I)_d := \Delta_\prec(I) \cap S_d$  with  $r$  distinct elements such that  $(\text{in}_\prec(I) : (M)) \neq \text{in}_\prec(I)$ .

**Definition 1.3.** The *generalized footprint function* of  $I$ , denoted  $\text{fp}_I$ , is the function  $\text{fp}_I: \mathbb{N}_+ \times \mathbb{N}_+ \rightarrow \mathbb{Z}$  given by

$$\text{fp}_I(d, r) := \begin{cases} \deg(S/I) - \max\{\deg(S/(\text{in}_\prec(I), M)) \mid M \in \mathcal{M}_{\prec, d, r}\} & \text{if } \mathcal{M}_{\prec, d, r} \neq \emptyset, \\ \deg(S/I) & \text{if } \mathcal{M}_{\prec, d, r} = \emptyset. \end{cases}$$

If  $r = 1$  one obtains the footprint function of  $I$  that was studied in [27] from a theoretical point of view (see [23,24] for some applications). In this case we denote  $\text{fp}_I(d, 1)$  simply by  $\text{fp}_I(d)$  and  $\mathcal{M}_{\prec, d, r}$  by  $\mathcal{M}_{\prec, d}$ . The importance of the footprint function is that it gives a lower bound on the generalized minimum degree function (Theorem 3.9) and it is computationally much easier to determine than the generalized minimum degree function. See the Appendix for scripts that implement these computations.

The content of this paper is as follows. In Section 2 we present some of the results and terminology that will be needed throughout the paper. In some of our results we will assume that there exists a linear form  $h$  that is regular on  $S/I$ , that is,  $(I : h) = I$ . There are wide families of ideals over finite fields that satisfy this hypothesis, e.g., vanishing ideals of parameterized codes [29]. Thus our results can be applied to a variety of Reed–Muller type codes [15], to monomial ideals, and to ideals that satisfy  $|K| > \deg(S/\sqrt{I})$ .

In Section 3 we study GMD functions of unmixed graded ideals. The *footprint matrix* ( $\text{fp}_I(d, r)$ ) and the *weight matrix* ( $\delta_I(d, r)$ ) of  $I$  are the matrices whose  $(d, r)$ -entries are  $\text{fp}_I(d, r)$  and  $\delta_I(d, r)$ , respectively. We show that the entries of each row of the weight matrix form a non-decreasing sequence and that the entries of each column of the weight matrix form a non-increasing sequence (Theorem 3.9). We also show that  $\text{fp}_I(d, r)$  is a lower bound for  $\delta_I(d, r)$  (Theorem 3.9). This was known when  $I$  is the vanishing ideal of a finite set of projective points [13, Theorem 4.9].

Let  $I \subset S$  be an unmixed graded ideal whose associated primes are generated by linear forms. In Section 4 we study the minimum distance functions of these ideals. For  $\delta_I(d) = \delta_I(d, 1)$ , the *regularity index* of  $\delta_I$ , denoted  $\text{reg}(\delta_I)$ , is the smallest  $d \geq 1$  such that  $\delta_I(d) = 1$ . If  $I$  is prime, we set  $\text{reg}(\delta_I) = 1$ . The regularity index of  $\delta_I$  is the index where the value of this numerical function stabilizes (Remark 3.10), named by analogy

with the regularity index for the Hilbert function of a fat point scheme  $Z$  which is the index where the Hilbert function  $H_Z$  of  $Z$  stabilizes.

In order to study the behavior of  $\delta_I$  we introduce a numerical invariant called the *v-number* (Definition 4.1). We give a description for this invariant in Proposition 4.2 that will allow us to compute it using computer algebra systems, e.g. *Macaulay2* [16] (Example 4.3).

**Proposition 4.6.** *Let  $I \subsetneq \mathfrak{m} \subset S$  be an unmixed graded ideal whose associated primes are generated by linear forms. Then  $\text{reg}(\delta_I) = \text{v}(I)$ .*

From the viewpoint of algebraic coding theory it is important to determine  $\text{reg}(\delta_I)$ . Indeed let  $\mathbb{X}$  be a set of projective points over a finite field  $K$ , let  $C_{\mathbb{X}}(d)$  be its corresponding Reed-Muller type code, and let  $\delta_{\mathbb{X}}(d)$  be the minimum distance of  $C_{\mathbb{X}}(d)$  (see Section 5), then  $\delta_{\mathbb{X}}(d) \geq 2$  if and only if  $1 \leq d < \text{reg}(\delta_{I(\mathbb{X})})$ . Our results give an effective method—that can be applied to any Reed-Muller type code—to compute the regularity index of the minimum distance (Corollary 5.6, Example 4.5).

The minimum socle degree  $s(I)$  of  $S/I$  (Definition 2.7) was used in [32] to obtain homological lower bounds for the minimum distance of a fat point scheme  $Z$  in  $\mathbb{P}^{s-1}$ . We relate the minimum socle degree, the v-number and the Castelnuovo-Mumford regularity for Geramita ideals in Theorem 4.10. For radical ideals it is an open problem whether or not  $\text{reg}(\delta_I) \leq \text{reg}(S/I)$  [27, Conjecture 4.2]. In dimension 1, the conjecture is true because of Proposition 4.6 and Theorem 4.10. Moreover, via Theorem 4.10, we can extend the notion of a Cayley–Bacharach scheme [10] by defining the notion of a Cayley–Bacharach ideal (Definition 4.14). It turns out that Cayley–Bacharach ideals are connected to Reed–Muller type codes and to minimum distance functions.

Letting  $H_I$  be the Hilbert function of  $I$ , we have  $\delta_I(d) > \deg(S/I) - H_I(d) + 1$  for some  $d \geq 1$  when  $I$  is unmixed of dimension at least 2 (Proposition 4.21). One of our main results is:

**Theorem 4.19.** *If  $I \subset S$  is a Geramita ideal and there exists  $h \in S_1$  regular on  $S/I$ , then*

$$\delta_I(d) \leq \deg(S/I) - H_I(d) + 1$$

for  $d \geq 1$  or equivalently  $H_I(d) - 1 \leq \text{hyp}_I(d)$  for  $d \geq 1$ .

This inequality is well known when  $I$  is the vanishing ideal of a finite set of projective points [29, p. 82]. In this case the inequality is called the *Singleton bound* [33, Corollary 1.1.65].

Projective Reed–Muller-type codes are studied in Section 5.

The main result of Section 5 shows that the entries of each column of the weight matrix  $(\delta_{\mathbb{X}}(d, r))$  form a decreasing sequence until they stabilize.

In particular one recovers the case when  $\mathbb{X}$  is a set, lying on a projective torus, parameterized by a finite set of monomials [12, Theorem 12]. Then we show that  $\delta_{\mathbb{X}}(d, H_{\mathbb{X}}(d))$  is equal to  $|\mathbb{X}|$  for  $d \geq 1$  (Corollary 5.7).

In Section 6 we examine minimum distance functions of complete intersection ideals and show some special cases of the following two conjectures.

**Conjecture 6.3.** *Let  $\mathbb{X}$  be a finite in  $\mathbb{P}^{s-1}$  and suppose that  $I = I(\mathbb{X})$  is a complete intersection generated by  $f_1, \dots, f_c$ ,  $c = s - 1$ , with  $d_i = \deg(f_i)$ , and  $2 \leq d_i \leq d_{i+1}$  for all  $i$ .*

- (a) (Tohăneanu–Van Tuyl [32, Conjecture 4.9])  $\delta_I(1) \geq (d_1 - 1)d_2 \cdots d_c$ .
- (b) (Eisenbud–Green–Harris [8, Conjecture CB10]) *If  $f_1, \dots, f_c$  are quadratic forms, then  $\text{hyp}_I(d) \leq 2^c - 2^{c-d}$  for  $1 \leq d \leq c$  or equivalently  $\delta_I(d) \geq 2^{c-d}$  for  $1 \leq d \leq c$ .*

We prove part (a) of this conjecture, in a more general setting, when  $I$  is equigenerated, that is, all minimal homogeneous generators have the same degree (Proposition 6.4, Remark 6.6). The conjecture also holds for  $\mathbb{P}^2$  [32, Theorem 4.10] (Corollary 6.5). According to [8], part (b) of this conjecture is true for the following values of  $d$ :  $1, c - 1, c$ .

For all unexplained terminology and additional information we refer to [4,6,26] (for the theory of Gröbner bases, commutative algebra, and Hilbert functions), and [22,33] (for the theory of error-correcting codes and linear codes).

## 2. Preliminaries

In this section we present some of the results that will be needed throughout the paper and introduce some more notation. All results of this section are well-known. To avoid repetitions, we continue to employ the notations and definitions used in Section 1.

**Commutative algebra.** Let  $I \neq (0)$  be a graded ideal of  $S$  of Krull dimension  $k$ . The *Hilbert function* of  $S/I$  is:  $H_I(d) := \dim_K(S_d/I_d)$  for  $d = 0, 1, 2, \dots$ , where  $I_d = I \cap S_d$ . By a theorem of Hilbert [31, p. 58], there is a unique polynomial  $P_I(x) \in \mathbb{Q}[x]$  of degree  $k-1$  such that  $H_I(d) = P_I(d)$  for  $d \gg 0$ . By convention the degree of the zero polynomial is  $-1$ .

The *degree* or *multiplicity* of  $S/I$  is the positive integer

$$\deg(S/I) := \begin{cases} (k-1)! \lim_{d \rightarrow \infty} H_I(d)/d^{k-1} & \text{if } k \geq 1, \\ \dim_K(S/I) & \text{if } k = 0. \end{cases}$$

As usual  $\text{ht}(I)$  will denote the height of the ideal  $I$ . By the dimension of  $I$  (resp.  $S/I$ ) we mean the Krull dimension of  $S/I$  denoted by  $\dim(S/I)$ .

One of the most useful and well-known facts about the degree is its additivity:

**Proposition 2.1.** (Additivity of the degree [28, Proposition 2.5]) *If  $I$  is an ideal of  $S$  and  $I = \mathfrak{q}_1 \cap \cdots \cap \mathfrak{q}_m$  is an irredundant primary decomposition, then*

$$\deg(S/I) = \sum_{\text{ht}(\mathfrak{q}_i) = \text{ht}(I)} \deg(S/\mathfrak{q}_i).$$

If  $F \subset S$ , the *ideal quotient* of  $I$  with respect to  $(F)$  is given by  $(I : (F)) = \{h \in S \mid hF \subset I\}$ . An element  $f$  of  $S$  is called a *zero-divisor* of  $S/I$ —as an  $S$ -module—if there is  $\bar{0} \neq \bar{a} \in S/I$  such that  $f\bar{a} = \bar{0}$ , and  $f$  is called *regular* on  $S/I$  if  $f$  is not a zero-divisor. Thus  $f$  is a zero-divisor if and only if  $(I : f) \neq I$ . An associated prime of  $I$  is a prime ideal  $\mathfrak{p}$  of  $S$  of the form  $\mathfrak{p} = (I : f)$  for some  $f$  in  $S$ .

**Theorem 2.2.** [34, Lemma 2.1.19, Corollary 2.1.30] *If  $I$  is an ideal of  $S$  and  $I = \mathfrak{q}_1 \cap \cdots \cap \mathfrak{q}_m$  is an irredundant primary decomposition with  $\text{rad}(\mathfrak{q}_i) = \mathfrak{p}_i$ , then the set of zero-divisors  $\mathcal{Z}(S/I)$  of  $S/I$  is equal to  $\bigcup_{i=1}^m \mathfrak{p}_i$ , and  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  are the associated primes of  $I$ .*

**Definition 2.3.** If  $I$  is a graded ideal of  $S$ , the *Hilbert series* of  $S/I$ , denoted  $F_I(x)$ , is given by

$$F_I(x) = \sum_{d=0}^{\infty} H_I(d)x^d, \text{ where } x \text{ is a variable.}$$

**Theorem 2.4.** (Hilbert–Serre [31, p. 58]) *Let  $I \subset S$  be a graded ideal of dimension  $k$ . Then there is a unique polynomial  $h(x) \in \mathbb{Z}[x]$  such that*

$$F_I(x) = \frac{h(x)}{(1-x)^k} \text{ and } h(1) > 0.$$

**Remark 2.5.** The leading coefficient of the Hilbert polynomial  $P_I(x)$  is equal to  $h(1)/(k-1)!$ . Thus  $h(1)$  is equal to  $\deg(S/I)$ .

**Definition 2.6.** Let  $I \subset S$  be a graded ideal. The  $a$ -invariant of  $S/I$ , denoted  $a(S/I)$ , is the degree of  $F_I(x)$  as a rational function, that is,  $a(S/I) = \deg(h(x)) - k$ . If  $h(x) = \sum_{i=0}^r h_i x^i$ ,  $h_i \in \mathbb{Z}$ ,  $h_r \neq 0$ , the vector  $(h_0, \dots, h_r)$  is called the  $h$ -vector of  $S/I$ .

**Definition 2.7.** Let  $I \subset S$  be a graded ideal and let  $\mathbf{F}$  be the minimal graded free resolution of  $S/I$  as an  $S$ -module:

$$\mathbf{F} : 0 \rightarrow \bigoplus_j S(-j)^{b_{g,j}} \rightarrow \cdots \rightarrow \bigoplus_j S(-j)^{b_{1,j}} \rightarrow S \rightarrow S/I \rightarrow 0.$$

The *Castelnuovo–Mumford regularity* of  $S/I$  (*regularity* of  $S/I$  for short) and the *minimum socle degree* (*s-number* for short) of  $S/I$  are defined as

$$\text{reg}(S/I) = \max\{j - i \mid b_{i,j} \neq 0\} \quad \text{and} \quad \text{s}(I) = \min\{j - g \mid b_{g,j} \neq 0\}.$$

If  $S/I$  is Cohen-Macaulay (i.e.  $g = \dim(S) - \dim(S/I)$ ) and there is a unique  $j$  such that  $b_{g,j} \neq 0$ , then the ring  $S/I$  is called *level*. In particular, a level ring for which the unique  $j$  such that  $b_{g,j} \neq 0$  is  $b_{g,j} = 1$  is called *Gorenstein*.

An excellent reference for the regularity of graded ideals is the book of Eisenbud [7].

**Definition 2.8.** The *regularity index* of the Hilbert function of  $S/I$ , or simply the *regularity index* of  $S/I$ , denoted  $\text{ri}(S/I)$ , is the least integer  $n \geq 0$  such that  $H_I(d) = P_I(d)$  for  $d \geq n$ .

The next result is valid over any field; see for instance [34, Theorem 5.6.4].

**Theorem 2.9.** [10] Let  $I$  be a graded ideal with  $\text{depth}(S/I) > 0$ . The following hold.

- (i) If  $\dim(S/I) \geq 2$ , then  $H_I(i) < H_I(i+1)$  for  $i \geq 0$ .
- (ii) If  $\dim(S/I) = 1$ , then there is an integer  $r$  and a constant  $c$  such that

$$1 = H_I(0) < H_I(1) < \cdots < H_I(r-1) < H_I(i) = c \quad \text{for } i \geq r.$$

**Lemma 2.10.** Let  $I \subset J \subset S$  be graded ideals of the same height. The following hold.

- (a) [9, Lemma 8] If  $I$  and  $J$  are unmixed, then  $I = J$  if and only if  $\deg(S/I) = \deg(S/J)$ .
- (b) If  $I \subsetneq J$ , then  $\deg(S/I) > \deg(S/J)$ .

**Proof.** (b) Since any associated prime of  $J/I$  is an associated prime of  $S/I$ ,  $\dim(J/I) = \dim(S/I)$ . From the short exact sequence

$$0 \rightarrow J/I \rightarrow S/I \rightarrow S/J \rightarrow 0$$

we obtain  $\deg(S/I) = \deg(J/I) + \deg(S/J)$ . As  $J/I$  is not zero, one has  $\deg(S/I) > \deg(S/J)$ .  $\square$

**Lemma 2.11.** [34, p. 122] Let  $I \subset S$  a graded ideal of height  $r$ . If  $K$  is infinite and  $I$  is minimally generated by forms of degree  $p \geq 1$ , then there are forms  $f_1, \dots, f_m$  of degree  $p$  in  $I$  such that  $f_1, \dots, f_r$  is a regular sequence and  $I$  is minimally generated by  $f_1, \dots, f_m$ .

**The footprint of an ideal.** Let  $\prec$  be a monomial order on  $S$  and let  $(0) \neq I \subset S$  be an ideal. If  $f$  is a non-zero polynomial in  $S$ , the *leading monomial* of  $f$  is denoted by  $\text{in}_\prec(f)$ . The *initial ideal* of  $I$ , denoted by  $\text{in}_\prec(I)$ , is the monomial ideal given by  $\text{in}_\prec(I) = (\{\text{in}_\prec(f) \mid f \in I\})$ .

We will use the following multi-index notation: for  $a = (a_1, \dots, a_s) \in \mathbb{N}^s$ , set  $t^a := t_1^{a_1} \cdots t_s^{a_s}$ . A monomial  $t^a$  is called a *standard monomial* of  $S/I$ , with respect to  $\prec$ , if  $t^a$  is not in the ideal  $\text{in}_\prec(I)$ . A polynomial  $f$  is called *standard* if  $f \neq 0$  and  $f$  is a  $K$ -linear combination of standard monomials. The set of standard monomials, denoted  $\Delta_\prec(I)$ , is called the *footprint* of  $S/I$ . The image of the standard polynomials of degree  $d$ , under the canonical map  $S \mapsto S/I$ ,  $x \mapsto \bar{x}$ , is equal to  $S_d/I_d$ , and the image of  $\Delta_\prec(I)$  is a basis of  $S/I$  as a  $K$ -vector space. This is a classical result of Macaulay (for a modern approach see [4, Chapter 5]). In particular, if  $I$  is graded, then  $H_I(d)$  is the number of standard monomials of degree  $d$ .

**Lemma 2.12.** [3, p. 3] *Let  $I \subset S$  be an ideal generated by  $\mathcal{G} = \{g_1, \dots, g_r\}$ , then*

$$\Delta_\prec(I) \subset \Delta_\prec(\text{in}_\prec(g_1), \dots, \text{in}_\prec(g_r)).$$

**Lemma 2.13.** [13, Lemma 4.7] *Let  $\prec$  be a monomial order, let  $I \subset S$  be an ideal, let  $F = \{f_1, \dots, f_r\}$  be a set of polynomial of  $S$  of positive degree, and let  $\text{in}_\prec(F) = \{\text{in}_\prec(f_1), \dots, \text{in}_\prec(f_r)\}$  be the set of initial terms of  $F$ . If  $(\text{in}_\prec(I) : (\text{in}_\prec(F))) = \text{in}_\prec(I)$ , then  $(I : (F)) = I$ .*

Let  $\prec$  be a monomial order and let  $\mathcal{F}_{\prec, d, r}$  be the set of all subsets  $F = \{f_1, \dots, f_r\}$  of  $S_d$  such that  $(I : (F)) \neq I$ ,  $f_i$  is a standard polynomial for all  $i$ ,  $\bar{f}_1, \dots, \bar{f}_r$  are linearly independent over the field  $K$ , and  $\text{in}_\prec(f_1), \dots, \text{in}_\prec(f_r)$  are distinct monomials.

The next result is useful for computations with *Macaulay2* [16] (see Procedure A.2).

**Proposition 2.14.** [13, Proposition 4.8] *The generalized minimum distance function of  $I$  is given by the following formula*

$$\delta_I(d, r) = \begin{cases} \deg(S/I) - \max\{\deg(S/(I, F)) \mid F \in \mathcal{F}_{\prec, d, r}\} & \text{if } \mathcal{F}_{\prec, d, r} \neq \emptyset, \\ \deg(S/I) & \text{if } \mathcal{F}_{\prec, d, r} = \emptyset. \end{cases}$$

An ideal  $I \subset S$  is called *radical* if  $I$  is equal to its radical. The radical of  $I$  is denoted by  $\sqrt{I}$ .

**Lemma 2.15.** [13, Lemma 3.3] *Let  $I \subset S$  be a radical unmixed graded ideal. If  $F = \{f_1, \dots, f_r\}$  is a set of homogeneous polynomials of  $S \setminus \{0\}$ ,  $(I : (F)) \neq I$ , and  $\mathcal{A}$  is the set of all associated primes of  $S/I$  that contain  $F$ , then  $\text{ht}(I) = \text{ht}(I, F)$ ,  $\mathcal{A} \neq \emptyset$ , and*

$$\deg(S/(I, F)) = \sum_{\mathfrak{p} \in \mathcal{A}} \deg(S/\mathfrak{p}).$$

### 3. Generalized minimum distance function of a graded ideal

In this section we study the generalized minimum distance function of a graded ideal.

Part (c) of the next lemma was known for vanishing ideals and part (b) for unmixed radical ideals [13, Proposition 3.5, Lemma 4.1].

**Lemma 3.1.** *Let  $I \subset S$  be an unmixed graded ideal, let  $\prec$  be a monomial order, and let  $F$  be a finite set of homogeneous polynomials of  $S$  such that  $(I: (F)) \neq I$ . The following hold.*

- (a)  $\text{ht}(I) = \text{ht}(I, F)$ .
- (b)  $\deg(S/(I, F)) < \deg(S/I)$  if  $I$  is an unmixed ideal and  $(F) \not\subset I$ .
- (c)  $\deg(S/I) = \deg(S/(I: (F))) + \deg(S/(I, F))$  if  $I$  is an unmixed radical ideal.
- (d) [13, Lemma 4.1]  $\deg(S/(I, F)) \leq \deg(S/(in_{\prec}(I), in_{\prec}(F))) \leq \deg(S/I)$ .

**Proof.** (a) As  $I \subsetneq (I: (F))$ , there is  $g \in S \setminus I$  such that  $g(F) \subset I$ . Hence the ideal  $(F)$  is contained in the set of zero-divisors of  $S/I$ . Thus, by Theorem 2.2 and since  $I$  is unmixed,  $(F)$  is contained in an associated prime ideal  $\mathfrak{p}$  of  $S/I$  of height  $\text{ht}(I)$ . Thus  $I \subset (I, F) \subset \mathfrak{p}$ , and consequently  $\text{ht}(I) = \text{ht}(I, F)$ . Therefore the set of associated primes of  $(I, F)$  of height equal to  $\text{ht}(I)$  is not empty and is equal to the set of associated primes of  $S/I$  that contain  $(F)$ .

(b) The inequality follows from part (a) and Lemma 2.10 (b).

(c) Let  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  be the associated primes of  $S/I$ . As  $I$  is a radical ideal, one has the decompositions

$$I = \bigcap_{i=1}^m \mathfrak{p}_i \quad \text{and} \quad (I: (F)) = \bigcap_{i=1}^m (\mathfrak{p}_i: (F)).$$

Note that  $(\mathfrak{p}_i: (F)) = S$  if  $F \subset \mathfrak{p}_i$  and  $(\mathfrak{p}_i: (F)) = \mathfrak{p}_i$  if  $F \not\subset \mathfrak{p}_i$ . Therefore, using the additivity of the degree of Proposition 2.1 and Lemma 2.15, we get

$$\deg(S/(I: (F))) = \sum_{F \not\subset \mathfrak{p}_i} \deg(S/\mathfrak{p}_i) \quad \text{and} \quad \deg(S/(I, F)) = \sum_{F \subset \mathfrak{p}_i} \deg(S/\mathfrak{p}_i).$$

Thus  $\deg(S/I) = \sum_{i=1}^m \deg(S/\mathfrak{p}_i) = \deg(S/(I: (F))) + \deg(S/(I, F))$ .  $\square$

**Definition 3.2.** Let  $I \subset S$  be a graded ideal. A sequence  $f_1, \dots, f_r$  of elements of  $S$  is called a  $(d, r)$ -sequence of  $S/I$  if the set  $F = \{f_1, \dots, f_r\}$  is in  $\mathcal{F}_{d, r}$ .

**Lemma 3.3.** *Let  $I \subset S$  be a graded ideal. A sequence  $f_1, \dots, f_r$  is a  $(d, r)$ -sequence of  $S/I$  if and only if the following conditions hold*

- (a)  $f_1, \dots, f_r$  are homogeneous polynomials of  $S$  of degree  $d \geq 1$ ,
- (b)  $(I: (f_1, \dots, f_r)) \neq I$ , and
- (c)  $f_i \notin (I, f_1, \dots, f_{i-1})$  for  $i = 1, \dots, r$ , where we set  $f_0 = 0$ .

**Proof.** The proof is straightforward.  $\square$

**Definition 3.4.** If  $I \subset S$  is a graded ideal, the *Vasconcelos function* of  $I$  is the function  $\vartheta_I: \mathbb{N}_+ \times \mathbb{N}_+ \rightarrow \mathbb{N}$  given by

$$\vartheta_I(d, r) := \begin{cases} \min\{\deg(S/(I:(F))) \mid F \in \mathcal{F}_{d,r}\} & \text{if } \mathcal{F}_{d,r} \neq \emptyset, \\ \deg(S/I) & \text{if } \mathcal{F}_{d,r} = \emptyset. \end{cases}$$

The next result was shown in [13, Theorem 4.5] for vanishing ideals over finite fields.

**Theorem 3.5.** Let  $I \subset S$  be a graded unmixed radical ideal. Then

$$\vartheta_I(d, r) = \delta_I(d, r) \text{ for } d \geq 1 \text{ and } 1 \leq r \leq H_I(d).$$

**Proof.** If  $\mathcal{F}_{d,r} = \emptyset$ , then  $\delta_I(d, r)$  and  $\vartheta_I(d, r)$  are equal to  $\deg(S/I)$ . Now assume that  $\mathcal{F}_{d,r} \neq \emptyset$ . Using Lemma 3.1(c), we obtain

$$\begin{aligned} \vartheta_I(d, r) &= \min\{\deg(S/(I:(F))) \mid F \in \mathcal{F}_{d,r}\} \\ &= \min\{\deg(S/I) - \deg(S/(I, F)) \mid F \in \mathcal{F}_{d,r}\} \\ &= \deg(S/I) - \max\{\deg(S/(I, F)) \mid F \in \mathcal{F}_{d,r}\} = \delta_I(d, r). \quad \square \end{aligned}$$

As the next result shows for  $r = 1$  we do not need the assumption that  $I$  is a radical ideal. For  $r \geq 2$  this assumption is essential, as shown in the next Example 3.6.

**Example 3.6.** Let  $I$  be the ideal  $(t_1^2, t_1 t_2, t_2^2)$  of the polynomial ring  $S = K[t_1, t_2]$  over a field  $K$  and let  $F = \{t_1, t_2\}$ . Then  $(I:(F)) = (I, F) = (t_1, t_2)$  and

$$3 = \deg(S/I) \neq \deg(S/(I:(F))) + \deg(S/(I, F)) = 2.$$

**Theorem 3.7.** [23, Theorem 4.4] Let  $I \subset S$  be an unmixed graded ideal. If  $\mathfrak{m} = (t_1, \dots, t_s)$  and  $d \geq 1$  is an integer such that  $\mathfrak{m}^d \not\subset I$ , then

$$\delta_I(d) = \min\{\deg(S/(I:f)) \mid f \in S_d \setminus I\}.$$

Recall from the introduction that the definition of  $\delta_I(d, r)$  was motivated by the notion of generalized Hamming weight of a linear code [18,35]. The following compilation of facts reflects the monotonicity of the generalized minimum distance function with respect of its two input values for the case of linear codes corresponding to reduced sets of points.

**Theorem 3.8.** Let  $C$  be a linear code of length  $m$  and dimension  $k$ . The following hold.

- (a) [35, Theorem 1, Corollary 1]  $1 \leq \delta_1(C) < \dots < \delta_k(C) \leq m$ .
- (b) [33, Corollary 1.1.65]  $r \leq \delta_r(C) \leq m - k + r$  for  $r = 1, \dots, k$ .

(c) If  $\delta_1(C) = m - k + 1$ , then  $\delta_r(C) = m - k + r$  for  $r = 1, \dots, k$ .

**Proof.** (c): By (a), one has  $m - k + 1 = \delta_1(C) \leq \delta_i(C) - (r - 1)$ . Thus  $m - k + r \leq \delta_i(C)$  and, by (b), equality holds.  $\square$

Below we consider more generally the behavior of the generalized minimum distance function and the footprint function for arbitrary graded ideals. The next result shows that the entries of any row (resp. column) of the weight matrix of  $I$  form a non-decreasing (resp. non-increasing) sequence. Parts (a)-(c) of the next result are broad generalizations of [13, Theorem 4.9] and [27, Theorem 3.6].

**Theorem 3.9.** *Let  $I \subset S$  be an unmixed graded ideal, let  $\prec$  be a monomial order on  $S$ , and let  $d \geq 1$ ,  $r \geq 1$  be integers. The following hold.*

- (a)  $\text{fp}_I(d, r) \leq \delta_I(d, r)$  for  $1 \leq r \leq H_I(d)$ .
- (b)  $\delta_I(d, r) \geq 1$ .
- (c)  $\text{fp}_I(d, r) \geq 1$  if  $\text{in}_\prec(I)$  is unmixed.
- (d)  $\delta_I(d, r) \leq \delta_I(d, r + 1)$ .
- (e) If there is  $h \in S_1$  regular on  $S/I$ , then  $\delta_I(d, r) \geq \delta_I(d + 1, r) \geq 1$ .

**Proof.** (a) If  $\mathcal{F}_{d,r} = \emptyset$ , then  $\delta_I(d, r) = \deg(S/I) \geq \text{fp}_I(d, r)$ . Now assume  $\mathcal{F}_{d,r} \neq \emptyset$ . Let  $F$  be any set in  $\mathcal{F}_{\prec, d, r}$ . By Lemma 2.13,  $\text{in}_\prec(F)$  is in  $\mathcal{M}_{\prec, d, r}$ , and by Lemma 3.1,  $\deg(S/(I, F)) \leq \deg(S/(\text{in}_\prec(I), \text{in}_\prec(F)))$ . Hence, by Proposition 2.14 and Lemma 3.1(b),  $\text{fp}_I(d, r) \leq \delta_I(d, r)$ .

(b) If  $\mathcal{F}_{d,r} = \emptyset$ , then  $\delta_I(d, r) = \deg(S/I) \geq 1$ , and if  $\mathcal{F}_{d,r} \neq \emptyset$ , then using Lemma 3.1(b) it follows that  $\delta_I(d, r) \geq 1$ .

(c) If  $\mathcal{M}_{\prec, d, r} = \emptyset$ , then  $\text{fp}_I(d, r) = \deg(S/I) \geq 1$ . Next assume that  $\mathcal{M}_{\prec, d, r}$  is not empty and pick  $M$  in  $\mathcal{M}_{\prec, d, r}$  such that

$$\text{fp}_I(d, r) = \deg(S/I) - \deg(S/(\text{in}_\prec(I), M)).$$

As  $\text{in}_\prec(I)$  is unmixed, by Lemma 3.1(b),  $\text{fp}_I(d, r) \geq 1$ .

(d) If  $\mathcal{F}_{d,r+1}$  is empty, then  $\delta_I(d, r) \leq \deg(S/I) = \delta_I(d, r + 1)$ . We may then assume  $\mathcal{F}_{d,r+1}$  is not empty and pick  $F = \{f_1, \dots, f_{r+1}\}$  in  $\mathcal{F}_{d,r+1}$  such that  $\text{hyp}_I(d, r + 1) = \deg(S/(I, F))$ . Setting  $F' = \{f_1, \dots, f_r\}$  and noticing that  $I \subsetneq (I : (F)) \subset (I : (F'))$ , we get  $F' \in \mathcal{F}_{d,r}$ . By the proof of Lemma 3.1, one has  $\text{ht}(I) = \text{ht}(I, F) = \text{ht}(I, F')$ . Taking Hilbert functions in the exact sequence

$$0 \longrightarrow (I, F)/(I, F') \longrightarrow S/(I, F') \longrightarrow S/(I, F) \longrightarrow 0$$

it follows that  $\deg(S/(I, F')) \geq \deg(S/(I, F))$ . Therefore

$$\text{hyp}_I(d, r) \geq \deg(S/(I, F')) \geq \deg(S/(I, F)) = \text{hyp}_I(d, r + 1) \Rightarrow \delta_I(d, r) \leq \delta_I(d, r + 1).$$

(e) By part (b),  $\delta_I(d, r) \geq 1$  for  $d \geq 1$ . Assume  $\mathcal{F}_{d,r} = \emptyset$ . Then  $\delta_I(d, r) = \deg(S/I)$ . If the set  $\mathcal{F}_{d+1,r}$  is empty, one has

$$\delta_I(d, r) = \delta_I(d+1, r) = \deg(S/I).$$

If the set  $\mathcal{F}_{d+1,r}$  is not empty, there is  $F \in \mathcal{F}_{d+1,r}$  such that

$$\delta_I(d+1, r) = \deg(S/I) - \deg(S/(I, F)) \leq \deg(S/I) = \delta_I(d, r).$$

Thus we may now assume  $\mathcal{F}_{d,r} \neq \emptyset$ . Pick  $F = \{f_1, \dots, f_r\}$  in  $\mathcal{F}_{d,r}$  such that

$$\delta_I(d, r) = \deg(S/I) - \deg(S/(I, F)).$$

By assumption there exists  $h \in S_1$  such that  $(I : h) = I$ . Hence the set  $h\overline{F} = \{h\bar{f}_i\}_{i=1}^r$  is linearly independent over  $K$ ,  $hF \subset S_{d+1}$ , and

$$I \subsetneq (I : F) \subset (I : hF),$$

that is,  $hF$  is in  $\mathcal{F}_{d+1,r}$ . Note that there exists  $\mathfrak{p} \in \text{Ass}(S/I)$  that contains  $(I, F)$  (see Lemma 3.1(a)). Hence the ideals  $(I, F)$  and  $(I, hF)$  have the same height because a prime ideal  $\mathfrak{p} \in \text{Ass}(S/I)$  contains  $(I, F)$  if and only if  $\mathfrak{p}$  contains  $(I, hF)$ . Therefore taking Hilbert functions in the exact sequence

$$0 \longrightarrow (I, F)/(I, hF) \longrightarrow S/(I, hF) \longrightarrow S/(I, F) \longrightarrow 0$$

it follows that  $\deg(S/(I, hF)) \geq \deg(S/(I, F))$ . As a consequence we get

$$\begin{aligned} \delta_I(d, r) &= \deg(S/I) - \deg(S/(I, F)) \geq \deg(S/I) - \deg(S/(I, hF)) \\ &\geq \deg(S/I) - \max\{\deg(S/(I, F')) \mid F' \in \mathcal{F}_{d+1,r}\} = \delta_I(d+1, r). \quad \square \end{aligned}$$

**Remark 3.10.** (a) Let  $I$  be a non-prime ideal and let  $\mathfrak{p}$  be an associated prime of  $I$ . There is  $f \in S_d$ ,  $d \geq 1$ , such that  $(I : f) = \mathfrak{p}$ . Note that  $f \in \mathcal{F}_d$ . By Theorem 3.7 one has  $\delta_I(d) = 1$ .

(b) If  $\dim(S/I) \geq 1$ , then  $\text{reg}(\delta_I)$  is the smallest  $n \geq 1$  such that  $\delta_I(d) = 1$  for  $d \geq n$ . This follows from Theorems 3.7 and 3.9.

**Example 3.11.** Let  $S = K[t_1, \dots, t_6]$  be a polynomial ring over the finite field  $K = \mathbb{F}_3$  and let  $I$  be the ideal  $(t_1t_6 - t_3t_4, t_2t_6 - t_3t_5)$ . The regularity and the degree of  $S/I$  are 2 and 4, respectively, and  $H_I(1) = 6$ ,  $H_I(2) = 19$ . Using Procedure A.2 and Theorem 3.9(a) we obtain:

$$(\text{fp}_I(d, r)) = \begin{bmatrix} 1 & 3 & 4 & 4 & 4 & 4 & \infty \\ 1 & 1 & 1 & 1 & 2 & 3 & 3 \end{bmatrix}, \quad d = 1, 2 \text{ and } r = 1, \dots, 7,$$

and  $(\delta_I(1, 1), \dots, \delta_I(1, 5)) = (3, 3, 4, 4, 4)$ .

**Definition 3.12.** If  $\text{fp}_I(d) = \delta_I(d)$  for  $d \geq 1$ , we say that  $I$  is a *Geil–Carvalho ideal*. If  $\text{fp}_I(d, r) = \delta_I(d, r)$  for  $d \geq 1$  and  $r \geq 1$ , we say that  $I$  is a *strongly Geil–Carvalho ideal*.

The next result generalizes [24, Proposition 3.11].

**Proposition 3.13.** *If  $I$  is an unmixed monomial ideal and  $\prec$  is any monomial order, then  $\delta_I(d, r) = \text{fp}_I(d, r)$  for  $d \geq 1$  and  $r \geq 1$ , that is,  $I$  is a strongly Geil–Carvalho ideal.*

**Proof.** The inequality  $\delta_I(d, r) \geq \text{fp}_I(d, r)$  follows from Theorem 3.9(a). To show the reverse inequality notice that  $\mathcal{M}_{\prec, d, r} \subset \mathcal{F}_{\prec, d, r}$  because one has  $I = \text{in}_{\prec}(I)$ . Also notice that  $\mathcal{M}_{\prec, d, r} = \emptyset$  if and only if  $\mathcal{F}_{\prec, d, r} = \emptyset$ , this follows from the proof of [13, Proposition 4.8]. Therefore one has  $\text{fp}_I(d, r) \geq \delta_I(d, r)$ .  $\square$

**Proposition 3.14.** *If  $I \subset S$  is an unmixed graded ideal and  $\dim(S/I) \geq 1$ , then*

$$\delta_I(d, H_I(d)) = \deg(S/I) \text{ for } d \geq 1.$$

**Proof.** We set  $r = H_I(d)$ . It suffices to show that  $\mathcal{F}_{d, r} = \emptyset$ . We proceed by contradiction. Assume that  $\mathcal{F}_{d, r}$  is not empty and let  $F = \{f_1, \dots, f_r\}$  be an element of  $\mathcal{F}_{d, r}$ . Let  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  be the associated primes of  $I$ . As  $I \subsetneq (I : (F))$ , we can pick  $g \in S$  such that  $g(F) \subset I$  and  $g \notin I$ . Then  $(F)$  is contained  $\cup_{i=1}^m \mathfrak{p}_i$ , and consequently  $(F) \subset \mathfrak{p}_i$  for some  $i$ . Since  $r = H_I(d)$ , one has

$$S_d/I_d = K\bar{f}_1 \oplus \dots \oplus K\bar{f}_r \Rightarrow S_d = Kf_1 + \dots + Kf_r + I_d.$$

Hence  $S_d \subset \mathfrak{p}_i$ , that is,  $\mathfrak{m}^d \subset \mathfrak{p}_i$ , where  $\mathfrak{m} = (t_1, \dots, t_s)$ . Therefore  $\mathfrak{p}_i = \mathfrak{m}$ , a contradiction because  $I$  is unmixed and  $\dim(S/I) \geq 1$ .  $\square$

**Example 3.15.** Let  $S = K[t_1, t_2, t_3]$  be a polynomial ring over a field  $K$  and let  $(\text{fp}_I(d, r))$  and  $(\delta_I(d, r))$  be the footprint matrix and the weight matrix of the ideal  $I = (t_1^3, t_2 t_3)$ . The regularity and the degree of  $S/I$  are 3 and 6. Using Procedure A.1 we obtain:

$$(\text{fp}_I(d, r)) = \begin{bmatrix} 3 & 5 & 6 & \infty & \infty & \infty \\ 2 & 3 & 4 & 5 & 6 & \infty \\ 1 & 2 & 3 & 4 & 5 & 6 \end{bmatrix}.$$

If  $r > H_I(d)$ , then  $\mathcal{M}_{\prec, d, r} = \emptyset$  and the  $(d, r)$ -entry of this matrix is equal to 6, but in this case we write  $\infty$  for computational reasons. Therefore, by Proposition 3.13,  $(\text{fp}_I(d, r))$  is equal to  $(\delta_I(d, r))$ . Setting  $F = \{t_1^2 t_2, t_1 t_2^2, t_1 t_3^2, t_1^2 t_3\}$  and  $F' = \{t_1^2 t_2, t_1 t_2^2, t_1 t_3^2 + t_2^3, t_1^2 t_3\}$ , we get

$$\delta_I(3, 4) = \deg(S/I) - \deg(S/(I, F)) = 4 \text{ and } \deg(S/I) - \deg(S/(I, F')) = 5.$$

Thus  $\delta_I(3, 4)$  is attained at  $F$ .

#### 4. Minimum distance function of a graded ideal

In this section we study minimum distance functions of unmixed graded ideals whose associated primes are generated by linear forms and the algebraic invariants of Geramita ideals.

##### 4.1. Minimum distance function for unmixed ideals

We begin by introducing the following numerical invariant which will be used to express the regularity index of the minimum distance function (Proposition 4.6).

**Definition 4.1.** The *v-number* of a graded ideal  $I$ , denoted  $v(I)$ , is given by

$$v(I) := \begin{cases} \min\{d \geq 1 \mid \text{there exists } f \in S_d \text{ and } \mathfrak{p} \in \text{Ass}(I) \text{ with } (I: f) = \mathfrak{p}\} & \text{if } I \subsetneq \mathfrak{m}, \\ 0 & \text{if } I = \mathfrak{m}, \end{cases}$$

where  $\text{Ass}(I)$  is the set of associated primes of  $S/I$  and  $\mathfrak{m} = (t_1, \dots, t_s)$  is the irrelevant maximal ideal of  $S$ .

The *v*-number is finite for any graded ideal by the definition of associated primes. If  $\mathfrak{p}$  is a prime ideal and  $\mathfrak{p} \neq \mathfrak{m}$ , then  $v(\mathfrak{p}) = 1$ .

Let  $I \subsetneq \mathfrak{m} \subset S$  be a graded ideal and let  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  be its associated primes. One can define the *v*-number of  $I$  locally at each  $\mathfrak{p}_i$  by

$$v_{\mathfrak{p}_i}(I) := \min\{d \geq 1 \mid \exists f \in S_d \text{ with } (I: f) = \mathfrak{p}_i\}.$$

The *v*-number of  $I$  is equal to  $\min\{v_{\mathfrak{p}_1}(I), \dots, v_{\mathfrak{p}_m}(I)\}$ . If  $I = I(\mathbb{X})$  is the vanishing ideal of a finite set  $\mathbb{X} = \{P_1, \dots, P_m\}$  of reduced projective points and  $\mathfrak{p}_i$  is the vanishing ideal of  $P_i$ , then  $v_{\mathfrak{p}_i}(I)$  is the degree of  $P_i$  in  $\mathbb{X}$  in the sense of [10, Definition 2.1].

We give an alternate description for the *v*-number using initial degrees of certain modules. This will allow us to compute the *v*-number using *Macaulay2* [16] (see Example 4.3). For a graded module  $M \neq 0$  we denote  $\alpha(M) = \min\{\deg(f) \mid f \in M, f \neq 0\}$ . By convention, for  $M = 0$  we set  $\alpha(0) = 0$ .

**Proposition 4.2.** Let  $I \subset S$  be an unmixed graded ideal. Then  $I \subsetneq (I: \mathfrak{p})$  for  $\mathfrak{p} \in \text{Ass}(I)$ ,

$$v(I) = \min\{\alpha((I: \mathfrak{p})/I) \mid \mathfrak{p} \in \text{Ass}(I)\},$$

and  $\alpha((I: \mathfrak{p})/I) = v_{\mathfrak{p}}(I)$  for  $\mathfrak{p} \in \text{Ass}(I)$ .

**Proof.** The strict inclusion  $I \subsetneq (I: \mathfrak{p})$  follows from the equivalence of Eq. (4.1) below. As a preliminary step of the proof of the equality we establish that for a prime  $\mathfrak{p} \in \text{Ass}(I)$  we have

$$(I : f) = \mathfrak{p} \text{ if and only if } f \in (I : \mathfrak{p}) \setminus I. \quad (4.1)$$

If  $(I : f) = \mathfrak{p}$ , it is clear that we have  $f \in (I : \mathfrak{p})$  and since  $(I : f) \neq S$  it follows that  $f \notin I$ . Conversely, if  $f \in (I : \mathfrak{p}) \setminus I$ , then  $\mathfrak{p} \subset (I : f)$ . Let  $\mathfrak{q} \in \text{Ass}(I : f)$ , which is a nonempty set since  $f \notin I$ . Since  $\text{Ass}(I : f) \subset \text{Ass}(I)$  and  $I$  is height unmixed, we have  $\text{ht}(\mathfrak{q}) = \text{ht}(\mathfrak{p})$  and  $\mathfrak{p} \subset (I : f) \subset \mathfrak{q}$ . It follows that  $\mathfrak{p} = (I : f) = \mathfrak{q}$ .

The equivalence of Eq. (4.1) implies that  $\alpha((I : \mathfrak{p})/I) = v_{\mathfrak{p}}(I)$ , and shows the equality

$$\{f \mid (I : f) = \mathfrak{p} \text{ for some } \mathfrak{p} \in \text{Ass}(I)\} = \bigcup_{\mathfrak{p} \in \text{Ass}(I)} (I : \mathfrak{p}) \setminus I.$$

The claim now follows by considering the minimum degree of a homogeneous element in the above sets.  $\square$

**Example 4.3.** Let  $S = \mathbb{Q}[t_1, t_2, t_3, t_4]$  be a polynomial ring over the rational numbers and let  $I$  be the ideal of  $S$  given by

$$I = (t_2^{10}, t_3^9, t_4^4, t_2 t_3 t_4^3) \cap (t_1^4, t_3^4, t_4^3, t_1 t_3 t_4^2) \cap (t_1^4, t_2^5, t_4^3) \cap (t_1^3, t_2^5, t_3^{10}).$$

The associated primes of  $I$  are  $\mathfrak{p}_1 = (t_2, t_3, t_4)$ ,  $\mathfrak{p}_2 = (t_1, t_3, t_4)$ ,  $\mathfrak{p}_3 = (t_1, t_2, t_4)$ ,  $\mathfrak{p}_4 = (t_1, t_2, t_3)$ . Using Proposition 4.2 together with Procedure A.3 we get  $s(I) = 10$ ,  $v(I) = 12$ ,  $\text{reg}(S/I) = 19$ ,  $v_{\mathfrak{p}_1}(I) = 12$ ,  $v_{\mathfrak{p}_2}(I) = 15$ ,  $v_{\mathfrak{p}_i}(I) = 18$  for  $i = 3, 4$ . Thus the minimum socle degree  $s(I)$  can be smaller than the  $v$ -number  $v(I)$ .

**Corollary 4.4.** *If  $I \subsetneq \mathfrak{m}$  is a graded ideal of  $S$  and  $\dim(S/I) = 0$ , then the minimum socle degree  $s(I) := \alpha((I : \mathfrak{m})/I)$  of  $S/I$  is equal to  $v(I)$ .*

**Proof.** The socle of  $S/I$  is given by  $\text{Soc}(S/I) = (I : \mathfrak{m})/I$ . Thus, by Proposition 4.2, one has the equality  $s(I) = v(I)$ .  $\square$

This corollary does not hold in dimension 1. There are examples of Geramita monomial ideals satisfying the strict inequality  $s(I) < v(I)$  (see Example 4.3). If  $S/I$  is a Cohen–Macaulay ring, the socle is understood to be the socle of some Artinian reduction of  $S/I$  by linear forms.

**Example 4.5.** Let  $K$  be the finite field  $\mathbb{F}_3$  and let  $\mathbb{X}$  be the following set of points in  $\mathbb{P}^2$ :

$$\begin{aligned} &[(1, 0, 1)], \quad [(1, 0, 0)], \quad [(1, 0, 2)], \quad [(1, 1, 0)], \quad [(1, 1, 1)], \\ &[(1, 1, 2)], \quad [(0, 0, 1)], \quad [(0, 1, 0)], \quad [(0, 1, 1)], \quad [(0, 1, 2)]. \end{aligned}$$

Using Propositions 4.2 and 4.6, together with Procedure A.4, we get  $v(I) = \text{reg}(\delta_{\mathbb{X}}) = 3$ ,  $\text{reg}(S/I) = 4$ ,  $\delta_{\mathbb{X}}(1) = 6$ ,  $\delta_{\mathbb{X}}(2) = 3$ , and  $\delta_{\mathbb{X}}(d) = 1$  for  $d \geq 3$ . The vanishing ideal of  $\mathbb{X}$  is generated by  $t_1 t_2^2 - t_1^2 t_2$ ,  $t_1 t_3^3 - t_1^3 t_3$ , and  $t_2 t_3^3 - t_2^3 t_3$ .

**Proposition 4.6.** *Let  $I \subsetneq \mathfrak{m} \subset S$  be an unmixed graded ideal whose associated primes are generated by linear forms. Then  $\text{reg}(\delta_I) = \text{v}(I)$ .*

**Proof.** Let  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  be the associated primes of  $I$ . We may assume that  $I$  not a prime ideal, otherwise  $\text{reg}(\delta_I) = \text{v}(I) = 1$ . If  $d_1 = \text{v}(I)$ , there are  $f \in S_{d_1}$  and  $\mathfrak{p}_i$  such that  $(I: f) = \mathfrak{p}_i$ . Then, by Theorem 3.7, one has  $\delta_I(d_1) = 1$ . Thus  $\text{reg}(\delta_I) \leq \text{v}(I)$ .

To show the reverse inequality set we set  $d_0 = \text{reg}(\delta_I)$ . Then  $\delta_I(d_0) = 1$ . Note that  $\mathfrak{m}^{d_0} \not\subset I$ ; otherwise  $\mathcal{F}_{d_0}(I) = \emptyset$  and by definition  $\delta_I(d_0)$  is equal to  $\deg(S/I)$ , a contradiction because  $I \subsetneq \mathfrak{m}$  and by Lemma 2.10  $\deg(S/I) > 1$ . Then, by Theorem 3.7, there is  $f \in S_{d_0} \setminus I$  such that  $\delta_I(d_0) = \deg(S/(I: f)) = 1$ . Let  $I = \cap_{i=1}^m \mathfrak{q}_i$  be the minimal primary decomposition of  $I$ , where  $\mathfrak{q}_i$  is a  $\mathfrak{p}_i$ -primary ideal. Note that  $(\mathfrak{q}_i: f)$  is a primary ideal if  $f \notin \mathfrak{q}_i$  because  $S/(\mathfrak{q}_i: f)$  is embedded in  $S/\mathfrak{q}_i$ . Thus the primary decomposition of  $(I: f)$  is  $\cap_{f \notin \mathfrak{q}_i} (\mathfrak{q}_i: f)$ . Therefore, by the additivity of the degree of Proposition 2.1, we get that  $(I: f) = (\mathfrak{q}_k: f)$  for some  $k$  such that  $f \notin \mathfrak{q}_k$  and  $\deg(S/(\mathfrak{q}_k: f)) = 1$ . Since  $S/\mathfrak{p}_k$  has also degree 1 and  $(\mathfrak{q}_k: f) \subset \mathfrak{p}_k$ , by Lemma 2.10, we get  $(I: f) = (\mathfrak{q}_k: f) = \mathfrak{p}_k$ , and consequently  $\text{v}(I) \leq \text{reg}(\delta_I)$ .  $\square$

**Corollary 4.7.** *Let  $I \subset S$  be an unmixed radical graded ideal. If all the associated primes of  $I$  are generated by linear forms and  $v = \text{v}(I)$  is its v-number, then*

$$\delta_I(1) > \dots > \delta_I(v-1) > \delta_I(v) = \delta_I(d) = 1 \quad \text{for } d \geq v.$$

**Proof.** It follows from [27, Theorem 3.8] and Proposition 4.6.  $\square$

The minimum distance function behaves well asymptotically.

**Corollary 4.8.** *Let  $I \subsetneq \mathfrak{m} \subset S$  be an unmixed graded ideal of dimension  $\geq 1$  whose associated primes are generated by linear forms. Then  $\delta_I(d) = 1$  for  $d \geq \text{v}(I)$ .*

**Proof.** This follows from Remark 3.10(b) and Proposition 4.6.  $\square$

The next result relates the minimum socle degree and the v-number.

**Proposition 4.9.** *Let  $I \subset S$  be an unmixed non-prime graded ideal whose associated primes are generated by linear forms and let  $h \in S_1$  be a regular element on  $S/I$ . The following hold:*

- (a) *If  $\delta_I(d) = \deg(S/(I: f))$ ,  $f \in \mathcal{F}_d \cap (I, h)$ , then  $d \geq 2$  and  $\delta_I(d) = \delta_I(d-1)$ .*
- (b) *If  $S/I$  is Cohen–Macaulay, then  $\text{v}(I, h) \leq \text{v}(I)$ .*
- (c) *If  $K$  is infinite and  $S/I$  is Cohen–Macaulay, then  $\text{s}(I) \leq \text{v}(I)$ .*

**Proof.** (a) Writing  $f = g + f_1 h$ , for some  $g \in I_d$  and  $f_1 \in S_{d-1}$ , one has  $(I: f) = (I: f_1)$ . Note that  $d \geq 2$ , otherwise if  $d = 1$ , then  $(I: f) = I$ , a contradiction because  $f \in \mathcal{F}_d$ . Therefore noticing that  $f_1 \in \mathcal{F}_{d-1}$ , by Theorems 3.7 and 3.9, we obtain

$$\delta_I(d) = \deg(S/(I:f)) = \deg(S/(I:f_1)) \geq \delta_I(d-1) \geq \delta_I(d) \Rightarrow \delta_I(d) = \delta_I(d-1).$$

(b) We set  $v = v(I)$ . By Proposition 4.2 there is an associated prime  $\mathfrak{p}$  of  $I$  and  $f \in (I:\mathfrak{p}) \setminus I$  such that  $f \in S_v$ . Then  $(I:f) = \mathfrak{p}$ ,  $f \in \mathcal{F}_d$ , and  $\delta_I(d) = \deg(S/(I:f)) = 1$ . We claim that  $f$  is not in  $(I,h)$ . If  $f \in (I,h)$ , then by part (a) one has  $v \geq 2$  and  $\delta_I(v-1) = 1$ , a contradiction because  $v$  is the regularity index of  $\delta_I$  (see Proposition 4.6). Thus  $f \notin (I,h)$ . Next we show the equality  $(\mathfrak{p},h) = ((I,h):f)$ . The inclusion “ $\subset$ ” is clear because  $(I:f) = \mathfrak{p}$ . Take an associated prime  $\mathfrak{p}'$  of  $((I,h):f)$ . The height of  $\mathfrak{p}'$  is  $\text{ht}(I) + 1$  because  $(I,h)$  is Cohen–Macaulay. Then  $\mathfrak{p}' = (\mathfrak{p}'',h)$  for some  $\mathfrak{p}''$  in  $\text{Ass}(I)$ . Taking into account that  $\mathfrak{p}$  and  $\mathfrak{p}''$  are generated by linear forms, we get the equality  $(\mathfrak{p},h) = (\mathfrak{p}'',h)$ . Thus  $(\mathfrak{p},h)$  is equal to  $((I,h):f)$ . Hence  $\delta_{(I,h)}(v) = 1$ , and consequently  $v(I,h) = \text{reg}(\delta_{(I,h)}) \leq \text{reg}(\delta_I) = v(I) = v$ .

(c) There exists a system of parameters  $\underline{h} = h_1, \dots, h_t$  of  $S/I$  consisting of linear forms, where  $t = \dim(S/I)$ . As  $S/I$  is Cohen–Macaulay,  $\underline{h}$  is a regular sequence on  $S/I$ . Hence, by part (b), we obtain

$$v(I, \underline{h}) = v(I, h_1, \dots, h_t) \leq \dots \leq v(I, h_1) \leq v(I).$$

Thus, by Corollary 4.4, we get  $s(I) = s(I, \underline{h}) = \alpha(((I, \underline{h}):\mathfrak{m})/(I, \underline{h})) = v(I, \underline{h}) \leq v(I)$ .  $\square$

#### 4.2. Minimum distance function for Geramita ideals and Cayley–Bacharach ideals

The minimum socle degree  $s(I)$ , the local  $v$ -number  $v_{\mathfrak{p}}(I)$ , and the regularity  $\text{reg}(S/I)$ , are related below. For complete intersections of dimension 1 they are all equal. In particular in this case one has  $\delta_I(d) \geq 2$  for  $1 \leq d < \text{reg}(S/I)$ .

**Theorem 4.10.** *Let  $I \subset S$  be a Geramita ideal and  $\mathfrak{p} \in \text{Ass}(I)$ . If  $I$  is not prime, then*

$$s(I) \leq v_{\mathfrak{p}}(I) \leq \text{reg}(S/I),$$

*with equality everywhere if  $S/I$  is a level ring.*

**Proof.** We set  $M = S/I$ ,  $r_0 = \text{reg}(S/I)$ ,  $n = v_{\mathfrak{p}}(I)$ , and  $I' = (I:\mathfrak{p})$ . To show the inequality  $n \leq r_0$  we proceed by contradiction. Assume that  $n > r_0$ . The  $S$ -modules in the exact sequence

$$0 \longrightarrow I'/I \longrightarrow S/I \longrightarrow S/I' \longrightarrow 0$$

are nonzero Cohen–Macaulay modules of dimension 1. Indeed, that  $I'/I \neq 0$  (resp.  $S/I' \neq 0$ ) follows from Proposition 4.2 (resp.  $I$  is not prime). That the modules are Cohen–Macaulay follows observing that  $I$  and  $I'$  are unmixed ideals of dimension 1. Since  $n$  is  $v_{\mathfrak{p}}(I)$  and  $r_0 < n$ , one has  $(I'/I)_{r_0} = 0$  (see the equivalence of Eq. (4.1) in the

proof of Proposition 4.2). Hence taking Hilbert functions in the above exact sequence in degree  $d = r_0$  (resp. for  $d \gg 0$ ), by Theorem 2.9, we get

$$\deg(S/I) = H_I(r_0) = H_{I'}(r_0) \leq \deg(S/I') \quad (\text{resp. } \deg(S/I) = \deg(I'/I) + \deg(S/I')).$$

As  $I'/I \neq 0$ ,  $\deg(I'/I) > 0$ . Hence  $\deg(S/I) > \deg(S/I')$ , a contradiction. Thus  $n \leq r_0$ .

To show the inequality  $s(I) \leq v_p(I)$  we make a change of coefficients. Consider the algebraic closure  $\overline{K}$  of  $K$ . We set

$$\overline{S} = S \otimes_K \overline{K} = \overline{K}[t_1, \dots, t_s] \quad \text{and} \quad \overline{I} = I\overline{S}.$$

Note that  $K \hookrightarrow \overline{K}$  is a faithfully flat extension. Apply the functor  $S \otimes_K (-)$ . By base change, it follows that  $S \hookrightarrow \overline{S}$  is a faithfully flat extension. Therefore  $H_I(d) = H_{\overline{I}}(d)$  for  $d \geq 0$  and  $\deg(S/I) = \deg(\overline{S}/\overline{I})$ . Furthermore the minimal graded free resolutions and the Hilbert series of  $S/I$  and  $\overline{S}/\overline{I}$  are identical. Thus  $S/I$  and  $\overline{S}/\overline{I}$  have the same regularity,  $s(I) = s(\overline{I})$ , and  $\overline{I}$  is Cohen–Macaulay of dimension 1. The ideal  $\overline{p} = p\overline{S}$  is a prime ideal of  $\overline{S}$  because  $p$  is generated by linear forms, and so is  $\overline{p}$ . The ideal  $\overline{I}$  is Geramita. To show this, let  $I = \cap_{i=1}^m q_i$  be the minimal primary decomposition of  $I$ , where  $q_i$  is a  $p_i$ -primary ideal. Since  $p_i\overline{S}$  is prime, the ideal  $q_i\overline{S}$  is a  $p_i\overline{S}$ -primary ideal of  $\overline{S}$ , and the minimal primary decomposition of  $\overline{I}$  is

$$\overline{I} = \left( \bigcap_{i=1}^m q_i \right) \overline{S} = \bigcap_{i=1}^m (q_i \overline{S}),$$

see [25, Sections 3.H, 5.D and 9.C]. Thus  $\overline{I}$  is a Geramita ideal. Recall that  $n \geq 1$  is the smallest integer such that there is  $f \in S_n$  with  $(I : f) = p$ . Fix  $f$  with these two properties. Then  $f \in (I : p) \setminus I$  and since  $\overline{I} \cap S = I$  and  $(I : p)\overline{S} = (I\overline{S} : p\overline{S})$ , one has  $f \in (I\overline{S} : p\overline{S}) \setminus I\overline{S}$ . Therefore, setting  $\overline{p} = p\overline{S}$ , we obtain  $v_{\overline{p}}(\overline{I}) \leq v_p(I)$ . Altogether using Proposition 4.9(c), we obtain

$$s(I) = s(\overline{I}) \leq v(\overline{I}) \leq v_{\overline{p}}(\overline{I}) \leq v_p(I) \leq \text{reg}(S/I) = \text{reg}(\overline{S}/\overline{I}).$$

If  $S/I$  is level then so is  $\overline{S}/\overline{I}$ , because the Betti numbers ( $b_{i,j}$  in Definition 2.7) for  $S/I$  and  $\overline{S}/\overline{I}$  agree [6, 6.10]. Furthermore, since the ring  $\overline{S}/\overline{I}$  is level, we have  $s(\overline{I}) = \text{reg}(\overline{S}/\overline{I})$  by [7, 4.13, 4.14] and which gives equality everywhere.  $\square$

**Definition 4.11.** [17,32] Let  $Z = a_1P_1 + \dots + a_mP_m \subset \mathbb{P}^{s-1}$  be a set of fat points, and suppose that  $Z' = a_1P_1 + \dots + (a_i - 1)P_i + \dots + a_mP_m$  for some  $i = 1, \dots, m$ . We call  $f \in S_d$  a *separator of  $P_i$  of multiplicity  $a_i$*  if  $f \in I(Z') \setminus I(Z)$ . The *vanishing ideal*  $I(Z)$  of  $Z$  is  $\cap_{i=1}^m \mathfrak{p}_i^{a_i}$ , where  $\mathfrak{p}_i$  is the vanishing ideal of  $P_i$ . If  $Z$  is a set of reduced points (i.e.,  $a_1 = \dots = a_m = 1$ ), the *degree* of  $P_i$ , denoted  $\deg_Z(P_i)$ , is the least degree of a separator of  $P_i$  of multiplicity 1.

**Remark 4.12.** If  $f$  is a separator of  $P_i$  of multiplicity  $a_i$  and  $\mathfrak{p}_i$  is the vanishing ideal of  $P_i$ , then  $f \in (I : \mathfrak{p}_i) \setminus I$ . The converse hold if  $a_i = 1$ .

**Corollary 4.13.** [32, Theorem 3.3] Let  $Z = a_1P_1 + \cdots + a_mP_m \subset \mathbb{P}^{s-1}$  be a set of fat points, and suppose that  $Z' = a_1P_1 + \cdots + (a_i - 1)P_i + \cdots + a_mP_m$  for some  $i = 1, \dots, m$ . If  $f$  is a separator of  $P_i$  of multiplicity  $a_i$ , then  $\deg(f) \geq v(I) \geq s(I)$ .

**Proof.** If  $f$  is a separator of  $P_i$  of multiplicity  $a_i$  and  $\mathfrak{p}_i$  be the vanishing ideal of  $P_i$ , then  $f \in (I : \mathfrak{p}_i) \setminus I$ . Hence, by Proposition 4.2 and Theorem 4.10, one has  $\deg(f) \geq v(I) \geq s(I)$ .  $\square$

A finite set  $\mathbb{X} = \{P_1, \dots, P_m\}$  of reduced points in  $\mathbb{P}^{s-1}$  is *Cayley-Bacharach* if every hypersurface of degree less than  $\text{reg}(S/I(\mathbb{X}))$  which contains all but one point of  $\mathbb{X}$  must contain all the points of  $\mathbb{X}$  or equivalently if  $\deg_{\mathbb{X}}(P_i) = \text{reg}(S/I(\mathbb{X}))$  for all  $i = 1, \dots, m$  [10, Definition 2.7]. Since  $\deg_{\mathbb{X}}(P_i) = v_{\mathfrak{p}_i}(I)$ , where  $\mathfrak{p}_i$  is the vanishing ideal of  $P_i$ , by Theorem 4.10 one can extend this notion to Geramita ideals.

**Definition 4.14.** A Geramita ideal  $I \subset S$  is called *Cayley-Bacharach* if  $v_{\mathfrak{p}}(I)$  is equal to  $\text{reg}(S/I)$  for all  $\mathfrak{p} \in \text{Ass}(I)$ .

As the next result shows Cayley-Bacharach ideals are connected to Reed–Muller type codes and to minimum distance functions.

**Corollary 4.15.** A Geramita ideal  $I \subset S$  is Cayley–Bacharach if and only if

$$\text{reg}(\delta_I) = v(I) = \text{reg}(S/I).$$

**Proof.** It follows from Proposition 4.6 and Theorem 4.10.  $\square$

There are some families of Reed–Muller type codes where the minimum distance and its index of regularity are known [21,30]. In these cases one can determine whether or not the corresponding sets of points are Cayley–Bacharach.

**Corollary 4.16.** If  $K = \mathbb{F}_q$  is a finite field and  $\mathbb{X} = \mathbb{P}^{s-1}$ , then  $I(\mathbb{X})$  is Cayley–Bacharach.

**Proof.** It follows from Corollary 4.15 because according to [30] the regularity index of  $\delta_{I(\mathbb{X})}$  is equal to  $\text{reg}(S/I(\mathbb{X}))$ .  $\square$

Next we give a lemma that allows comparisons between the generalized minimum distances of ideals related by containment.

**Lemma 4.17.** If  $I, I'$  are unmixed graded ideals of the same height and  $J$  is a graded ideal such that  $I' = (I : J)$ , then  $\mathcal{F}_d(I') \subset \mathcal{F}_d(I)$  and

$$\deg(S/I') - \delta_{I'}(d) \leq \deg(S/I) - \delta_I(d).$$

**Proof.** Let  $f \in \mathcal{F}_d(I')$ . Then  $f \notin I'$  and  $(I': f) \neq I'$ , and since we have the following relations

$$I' \subsetneq (I': f) = ((I: J): f) = (I: (fJ)) = ((I: f): J)$$

we deduce that  $(I: f) \neq I$  (otherwise the last ideal displayed above would be  $I'$ ). Note that  $I \subset I'$ , so  $f \notin I$ . The second statement follows from the inequality

$$\begin{aligned} \deg(S/I') - \delta_{I'}(d) &= \max\{\deg(S/(I', f)) \mid f \in \mathcal{F}_d(I')\} \\ &\leq \max\{\deg(S/(I, g)) \mid g \in \mathcal{F}_d(I)\} = \deg(S/I) - \delta_I(d). \end{aligned}$$

This inequality is a consequence of the observation that if  $f \in \mathcal{F}_d(I')$ , then  $\text{ht}(I', f) = \text{ht}(I')$ , and since  $f \in \mathcal{F}_d(I)$  one also has  $\text{ht}(I, f) = \text{ht}(I)$  by Lemma 3.1(a). Thus  $\deg(S/(I', f)) \leq \deg(S/(I, f))$ .  $\square$

One of our main results shows that the function  $\eta: \mathbb{N}_+ \rightarrow \mathbb{Z}$  given by

$$\eta(d) := (\deg(S/I) - H_I(d) + 1) - \delta_I(d)$$

non-negative for Geramita ideals (see Theorem 4.19).

**Lemma 4.18.** *Let  $I \subset S$  be a Geramita ideal. If  $\mathcal{F}_{d_0} = \emptyset$  for some  $d_0 \geq 1$ , then  $\eta(d_0) = 0$  and  $\eta(d) \geq 0$  for all  $d \geq 1$ .*

**Proof.** Let  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  be the associated primes of  $I$ . As  $\mathfrak{p}_k$  is generated by linear forms, the initial ideal of  $\mathfrak{p}_k$ , w.r.t. the lexicographical order  $\prec$ , is generated by  $s - 1$  variables. Hence, as  $\mathfrak{p}_k$  and  $\text{in}_{\prec}(\mathfrak{p}_k)$  have the same Hilbert function,  $\deg(S/\mathfrak{p}_k) = 1$  and  $H_{\mathfrak{p}_k}(d) = 1$  for  $d \geq 1$ . Assume that  $\mathcal{F}_{d_0} = \emptyset$ . Then  $\delta_I(d_0) = \deg(S/I)$  and  $(I: f) = I$  for any  $f \in S_{d_0} \setminus I$ . Hence, by Theorem 2.2, we get

$$(\mathfrak{p}_1)_{d_0} \subset \left( \bigcup_{i=1}^m \mathfrak{p}_i \right) \cap S_{d_0} \subset I_{d_0} \subset (\mathfrak{p}_1)_{d_0}.$$

Thus  $I_{d_0} = (\mathfrak{p}_1)_{d_0}$ ,  $H_I(d_0) = H_{\mathfrak{p}_1}(d_0) = 1$ ,  $H_I(0) = 1$ , and  $\eta(d_0) = 0$ . Using Theorem 2.9(ii), one has  $H_I(d) = 1$  for  $d \geq 1$ . Therefore  $\eta(d) \geq 0$  for  $d \geq 1$ .  $\square$

#### 4.3. Singleton bound

We come to one of our main results. The inequality in the following theorem is well known when  $I$  is the vanishing ideal of a finite set of projective points [29, p. 82]. In this case the inequality is called the *Singleton bound* [33, Corollary 1.1.65].

**Theorem 4.19.** *Let  $I \subset S$  be an unmixed graded ideal whose associated primes are generated by linear forms and such that there exists  $h \in S_1$  regular on  $S/I$ . If  $\dim(S/I) = 1$ , then*

$$\delta_I(d) \leq \deg(S/I) - H_I(d) + 1 \quad \text{for } d \geq 1$$

or equivalently  $H_I(d) - 1 \leq \text{hyp}_I(d)$  for  $d \geq 1$ .

**Proof.** The proof is by induction on  $\deg(S/I)$ . If  $\deg(S/I) = 1$ , then  $I = \mathfrak{p}$  is a prime generated by linear forms,  $H_I(d) = 1$  for all  $d \geq 0$  and  $\mathcal{F}_d(I) = \emptyset$  for all  $d \geq 1$ . The latter follows since for any prime  $\mathfrak{p}$ ,  $(\mathfrak{p}: f) \neq \mathfrak{p}$  implies  $f \in \mathfrak{p}$ . So the result is verified in this case. Let  $v = v(I)$  be the v-number of  $I$ . By Proposition 4.2,  $v = \alpha((I: \mathfrak{p})/I)$  for some  $\mathfrak{p} \in \text{Ass}(I)$ . Set  $I' = (I: \mathfrak{p})$ . The short exact sequence

$$0 \longrightarrow I'/I \longrightarrow S/I \longrightarrow S/I' \longrightarrow 0$$

together with the unmixed property of  $S/I$  show that  $\dim(I'/I) = 1$  and  $\text{depth}(I'/I) = 1$ . Therefore,  $H_{I'/I}(d) = 0$  for  $d < \alpha((I: \mathfrak{p})/I) = v$  and  $H_{I'/I}(d) > 0$  for  $d \geq \alpha((I: \mathfrak{p})/I) = v$ , and consequently  $H_{I'}(d) = H_I(d)$  for  $d < v$  and  $H_{I'}(d) > H_I(d)$  for  $d \geq v$ . The last statement yields that  $\deg(S/I) > \deg(S/I')$ . This also follows from Lemma 2.10(b).

If  $d < v$  we deduce from Lemma 4.17, the inductive hypothesis and  $H_{I'}(d) = H_I(d)$  that

$$\deg(S/I) - \delta_I(d) \geq \deg(S/I') - \delta_{I'}(d) \geq H_{I'}(d) - 1 = H_I(d) - 1,$$

which is the desired inequality. If  $d \geq v$  we know that there exists  $f \in S_v$  such that  $(I: f) = \mathfrak{p}$  and thus  $(I: h^{d-v}f) = \mathfrak{p}$ . Therefore  $\delta_I(d) = 1$  and since  $\deg(S/I) \geq H_I(d)$  for any  $d$  the desired inequality follows.  $\square$

The next result is known for complete intersection vanishing ideals over finite fields [14, Lemma 3]. As an application we extend this result to Geramita Gorenstein ideals.

**Corollary 4.20.** *Let  $I \subset S$  be a Geramita ideal. If  $I$  is Gorenstein and  $r_0 = \text{reg}(S/I) \geq 2$ , then  $\delta_I(r_0 - 1)$  is equal to 2.*

**Proof.** By Proposition 4.6 and Theorem 4.10,  $r_0$  is the regularity index of  $\delta_I$ . Thus  $\delta_I(r_0 - 1) \geq 2$ .

We show that  $\deg(S/I) = 1 + H_I(r_0 - 1)$ . For this, we may assume that  $K$  is infinite. Indeed, consider the algebraic closure  $\overline{K}$  of  $K$ . We set  $\overline{S} = S \otimes_K \overline{K}$  and  $\overline{I} = I\overline{S}$ . From [31, Lemma 1.1], we have  $\overline{I}$  is Gorenstein,  $H_I(d) = H_{\overline{I}}(d)$  for  $d \geq 0$  and  $\deg(S/I) = \deg(\overline{S}/\overline{I})$ . Since  $\overline{K}$  is infinite, there is  $h \in \overline{S}$  that is regular on  $\overline{S}/\overline{I}$ . Then by [2, 3.1.19](b) the quotient ring  $A = \overline{S}/(\overline{I}, h)$  is Gorenstein of dimension 0, by [7, 4.13, 4.14] it has

$r_0 = \text{reg}(\overline{S}/\overline{I}) = \text{reg}(A)$ , and by [2, 4.7.11(b)]  $\deg(\overline{S}/\overline{I}) = \deg(A) = \sum_{i=0}^{r_0} H_A(i)$ . By [2, proof of 4.1.10],  $F_A(x) = (1-x)F_I(x)$  hence  $H_{\overline{I}}(n) = \sum_{i=0}^n H_A(i)$  for any  $n \geq 0$ . From here, using that  $H_A(r_0) = 1$ , since  $A$  is Gorenstein, we deduce that

$$\begin{aligned} \deg(S/I) &= \deg(\overline{S}/\overline{I}) = \deg(A) = \sum_{i=0}^{r_0} H_A(i) = 1 + \sum_{i=0}^{r_0-1} H_A(i) = 1 + H_{\overline{I}}(r_0-1) \\ &= 1 + H_I(r_0-1). \end{aligned}$$

Finally, making  $d = r_0 - 1$  in Theorem 4.19, we get  $\delta_I(r_0 - 1) \leq 2$ . Thus equality holds.  $\square$

Note that the situation is quite different from the conclusion of Theorem 4.19 if  $\dim(S/I) \geq 2$ .

**Proposition 4.21.** *Let  $I \subset S$  be an unmixed graded ideal. If  $\dim(S/I) \geq 2$ , then*

$$\delta_I(d) > \deg(S/I) - H_I(d) + 1 \text{ for some } d \geq 1.$$

**Proof.** Note that  $\mathfrak{m} = (t_1, \dots, t_s)$  is not an associated prime of  $I$ , that is,  $\text{depth}(S/I) \geq 1$ . Assume that  $\mathcal{F}_d = \emptyset$  for some  $d \geq 2$ . As  $H_I(0) = 1$  and  $\delta_I(d)$  is equal to  $\deg(S/I)$ , by Theorem 2.9(i), one has  $H_I(d) > 1$  and the inequality holds. Now assume that  $\mathcal{F}_d \neq \emptyset$  for  $d \geq 2$ . For each  $d \geq 2$  pick  $f_d \in \mathcal{F}_d$  such that

$$\delta_I(d) = \deg(S/I) - \deg(S/(I, f_d)).$$

As  $H_I$  is strictly increasing by Theorem 2.9(i), using Lemma 3.1(b), we get

$$\deg(S/(I, f_d)) < \deg(S/I) < H_I(d) - 1$$

for  $d \gg 0$ . Thus the required inequality holds for  $d \gg 0$ .  $\square$

## 5. Reed-Muller type codes

In this section we give refined information on the minimum distance function for the Reed–Muller codes defined in the Introduction. The key insight is that, in the case of the projective Reed–Muller codes, this minimum distance function can be realized as a generalized minimum distance function for a finite set of points in projective space, often called evaluation points in the algebraic coding context.

**Theorem 5.1.** [13, Theorem 4.5] *Let  $\mathbb{X}$  be a finite set of points in a projective space  $\mathbb{P}^{s-1}$  over a field  $K$  and let  $I(\mathbb{X})$  be its vanishing ideal. If  $d \geq 1$  and  $1 \leq r \leq H_{\mathbb{X}}(d)$ , then*

$$\delta_r(C_{\mathbb{X}}(d)) = \delta_{I(\mathbb{X})}(d, r).$$

By Theorem 3.8 (a) and [35, Theorem 1, Corollary 1], the entries of each row of the weight matrix  $(\delta_{\mathbb{X}}(d, r))$  form an increasing sequence until they stabilize. We show in Theorem 5.3 below that the entries of each column of the weight matrix  $(\delta_{\mathbb{X}}(d, r))$  form a decreasing sequence.

Before we can prove this result we need an additional lemma. Recall that the *support*  $\chi(\beta)$  of a vector  $\beta \in K^m$  is  $\chi(K\beta)$ , that is,  $\chi(\beta)$  is the set of non-zero entries of  $\beta$ .

**Lemma 5.2.** *Let  $D$  be a subcode of  $C$  of dimension  $r \geq 1$ . If  $\beta_1, \dots, \beta_r$  is a  $K$ -basis for  $D$  with  $\beta_i = (\beta_{i,1}, \dots, \beta_{i,m})$  for  $i = 1, \dots, r$ , then  $\chi(D) = \cup_{i=1}^r \chi(\beta_i)$  and the number of elements of  $\chi(D)$  is the number of non-zero columns of the matrix:*

$$\begin{bmatrix} \beta_{1,1} & \cdots & \beta_{1,i} & \cdots & \beta_{1,m} \\ \beta_{2,1} & \cdots & \beta_{2,i} & \cdots & \beta_{2,m} \\ \vdots & \cdots & \vdots & \cdots & \vdots \\ \beta_{r,1} & \cdots & \beta_{r,i} & \cdots & \beta_{r,m} \end{bmatrix}.$$

**Theorem 5.3.** *Let  $\mathbb{X}$  be a finite set of points in  $\mathbb{P}^{s-1}$ , let  $I = I(\mathbb{X})$  be its vanishing ideal, and let  $1 \leq r \leq |\mathbb{X}|$  be a fixed integer. Then there is an integer  $d_0 \geq 1$  such that*

$$\delta_I(1, r) > \delta_I(2, r) > \cdots > \delta_I(d_0, r) = \delta_I(d, r) = r \text{ for } d \geq d_0.$$

**Proof.** Let  $[P_1], \dots, [P_m]$  be the points of  $\mathbb{X}$ . By Theorem 5.1 there exists a linear subcode  $D$  of  $C_{\mathbb{X}}(d)$  of dimension  $r$  such that  $\delta_I(d, r) = \delta_{\mathbb{X}}(d, r) = |\chi(D)|$ . Pick a  $K$ -basis  $\beta_1, \dots, \beta_r$  of  $D$ . Each  $\beta_i$  can be written as

$$\beta_i = (\beta_{i,1}, \dots, \beta_{i,k}, \dots, \beta_{i,m}) = (f_i(P_1), \dots, f_i(P_k), \dots, f_i(P_m))$$

for some  $f_i \in S_d$ . Consider the matrix  $B$  whose rows are  $\beta_1, \dots, \beta_m$ :

$$B = \begin{bmatrix} f_1(P_1) & \cdots & f_1(P_k) & \cdots & f_1(P_m) \\ f_2(P_1) & \cdots & f_2(P_k) & \cdots & f_2(P_m) \\ \vdots & \cdots & \vdots & \cdots & \vdots \\ f_r(P_1) & \cdots & f_r(P_k) & \cdots & f_r(P_m) \end{bmatrix}.$$

As  $B$  has rank  $r$ , by permuting columns and applying elementary row operations, the matrix  $B$  can be brought to the form:

$$B' = \begin{bmatrix} g_1(Q_1) & & & g_1(Q_{r+1}) & \cdots & g_1(Q_m) \\ & g_2(Q_2) & & g_2(Q_{r+1}) & \cdots & g_2(Q_m) \\ & & \mathbf{0} & & & \\ & & & \ddots & & \vdots \\ & & & & g_r(Q_r) & g_r(Q_{r+1}) & \cdots & g_r(Q_m) \end{bmatrix},$$

where  $g_1, \dots, g_r$  are linearly independent polynomials over the field  $K$  modulo  $I$  of degree  $d$ ,  $Q_1, \dots, Q_m$  are a permutation of  $P_1, \dots, P_m$ , the first  $r$  columns of  $B'$  form a diagonal

matrix such that  $g_i(Q_i) \neq 0$  for  $i = 1, \dots, r$ , and the ideals  $(f_1, \dots, f_r)$  and  $(g_1, \dots, g_r)$  are equal. Let  $D'$  be the linear space generated by the rows of  $B'$ . The operations applied to  $B$  did not affect the size of the support of  $D$  (Lemma 5.2), that is,  $|\chi(D)| = |\chi(D')|$ .

Note that  $\delta_r(C_{\mathbb{X}}(d))$  depends only on  $\mathbb{X}$ , that is,  $\delta_r(C_{\mathbb{X}}(d))$  is independent of how we order the points in  $\mathbb{X}$  (cf. Theorem 5.1). Let  $\text{ev}'_d: S_d \rightarrow K^m$  be the evaluation map,  $f \mapsto (f(Q_1), \dots, f(Q_m))$ , relative to the points  $[Q_1], \dots, [Q_m]$ . By Theorem 3.8,  $\delta_{\mathbb{X}}(d, r) \geq r$ .

First we assume that  $\delta_{\mathbb{X}}(d, r) = r$  for some  $d \geq 1$  and  $r \geq 1$ . Then the  $i$ -th column of  $B'$  is zero for  $i > r$ . For each  $1 \leq i \leq r$  pick  $h_i \in S_1$  such that  $h_i(Q_i) \neq 0$ . The polynomials  $h_1g_1, \dots, h_rg_r$  are linearly independent modulo  $I$  because  $(h_ig_i)(Q_j)$  is not 0 if  $i = j$  and is 0 if  $i \neq j$ . The image of  $Kh_1g_1 \oplus \dots \oplus Kh_rg_r$ , under the map  $\text{ev}'_{d+1}$ , is a subcode  $D''$  of  $C_{\mathbb{X}}(d+1)$  of dimension  $r$  and  $|\chi(D'')| = r$ . Thus  $\delta_{\mathbb{X}}(d+1, r) \leq r$ , and consequently  $\delta_{\mathbb{X}}(d+1, r) = r$ .

Next we assume that  $\delta_{\mathbb{X}}(d, r) > r$ . Then  $B'$  has a nonzero column  $(g_1(Q_k), \dots, g_r(Q_k))^{\top}$  for some  $k > r$ . It suffices to show that  $\delta_{\mathbb{X}}(d, r) > \delta_{\mathbb{X}}(d+1, r)$ . According to [23, Lemma 2.14(ii)] for each  $1 \leq i \leq r$  there is  $h_i$  in  $S_1$  such that  $h_i(Q_i) \neq 0$  and  $h_i(Q_k) = 0$ . Let  $B''$  be the matrix:

$$B'' = \begin{bmatrix} h_1g_1(Q_1) & & & h_1g_1(Q_{r+1}) & \cdots & h_1g_1(Q_m) \\ & h_2g_2(Q_2) & & h_2g_2(Q_{r+1}) & \cdots & h_2g_2(Q_m) \\ \mathbf{0} & & \ddots & & & \vdots \\ & & & h_rg_r(Q_r) & h_rg_r(Q_{r+1}) & \cdots & h_rg_r(Q_m) \end{bmatrix}.$$

The image of  $Kh_1g_1 \oplus \dots \oplus Kh_rg_r$ , under the map  $\text{ev}'_{d+1}$ , is a subcode  $V$  of  $C_{\mathbb{X}}(d+1)$  of dimension  $r$  because the rank of  $B''$  is  $r$ , and since the  $k$ -column of  $B''$  is zero, we get

$$\delta_{\mathbb{X}}(d, r) = |\chi(D)| = |\chi(D')| > |\chi(V)| \geq \delta_{\mathbb{X}}(d+1, r).$$

Thus  $\delta_{\mathbb{X}}(d, r) > \delta_{\mathbb{X}}(d+1, r)$ .  $\square$

**Corollary 5.4.** *Let  $\mathbb{X}$  be a finite set of points in  $\mathbb{P}^{s-1}$  and let  $I = I(\mathbb{X})$  be its vanishing ideal. If  $I$  is a complete intersection, then  $\delta_I(d) \geq \text{reg}(S/I) - d + 1$  for  $1 \leq d < \text{reg}(S/I)$ .*

**Proof.** If  $r_0$  denotes the regularity of  $S/I$ , by Theorem 4.10, one has  $\text{v}(I) = r_0$ . Thus  $\delta_I(r_0 - 1) \geq 2$  and the result follows from Theorem 5.3 by setting  $r = 1$ .  $\square$

**Corollary 5.5.** [12, Theorem 12] *If  $\mathbb{X}$  is a set parameterized by monomials lying on a projective torus and  $1 \leq r \leq |\mathbb{X}|$  be a fixed integer, then there is an integer  $d_0 \geq 1$  such that*

$$\delta_r(C_{\mathbb{X}}(1)) > \delta_r(C_{\mathbb{X}}(2)) > \dots > \delta_r(C_{\mathbb{X}}(d_0)) = \delta_r(C_{\mathbb{X}}(d)) = r \text{ for } d \geq d_0.$$

**Proof.** It follows at once from Theorems 5.1 and 5.3.  $\square$

**Corollary 5.6.** *Let  $\mathbb{X}$  be a finite set of points of  $\mathbb{P}^{s-1}$  and let  $\delta_{\mathbb{X}}(d)$  be the minimum distance of  $C_{\mathbb{X}}(d)$ . Then  $\delta_{\mathbb{X}}(d) = 1$  if and only if  $d \geq v(I)$ .*

**Proof.** It follows from Proposition 4.6, and Theorems 5.1 and 5.3.  $\square$

**Corollary 5.7.** *If  $\mathbb{X}$  is a finite set of  $\mathbb{P}^{s-1}$  over a field  $K$ , then  $\delta_{\mathbb{X}}(d, H_{\mathbb{X}}(d)) = |\mathbb{X}|$  for  $d \geq 1$ .*

**Proof.** It follows at once from Proposition 3.14 and Theorem 5.1.  $\square$

## 6. Complete intersections

In this section we examine minimum distance functions of complete intersection ideals.

**Definition 6.1.** An ideal  $I \subset S$  is called a *complete intersection* if there exist  $g_1, \dots, g_r$  in  $S$  such that  $I = (g_1, \dots, g_r)$ , where  $r = \text{ht}(I)$  is the height of  $I$ .

There are a number of interesting open problems regarding the minimum distance of complete intersection functions. We discuss one such problem in Conjecture 6.2 below and relate this problem to [8, Conjecture CB12]) in the second part of this section.

**Conjecture 6.2.** [24] *Let  $I \subset S := K[t_1, \dots, t_s]$  be a complete intersection graded ideal of dimension 1 generated by forms  $f_1, \dots, f_c$ ,  $c = s-1$ , with  $d_i = \deg(f_i)$  and  $2 \leq d_i \leq d_{i+1}$  for  $i \geq 1$ . If the associated primes of  $I$  are generated by linear forms, then*

$$\delta_I(d) \geq (d_{k+1} - \ell)d_{k+2} \cdots d_c \quad \text{if } 1 \leq d \leq \sum_{i=1}^c (d_i - 1) - 1,$$

where  $0 \leq k \leq c-1$  and  $\ell$  are integers such that  $d = \sum_{i=1}^k (d_i - 1) + \ell$  and  $1 \leq \ell \leq d_{k+1} - 1$ .

This conjecture holds if the initial ideal of  $I$  with respect to some monomial order is a complete intersection [24, Theorem 3.14]. Our results show that for complete intersections  $v(I) = \text{reg}(S/I)$  (Theorem 4.10) and  $\delta_I(d) \geq \text{reg}(S/I) - d + 1$  for  $1 \leq d < \text{reg}(S/I)$  if  $I$  is a vanishing ideal (Corollary 5.4). Thus the conjecture is best possible for vanishing ideals in the sense that it covers all cases where  $\delta_I(d) > 1$  because the regularity of  $S/I$  is equal to  $\sum_{i=1}^c (d_i - 1)$ .

Two special cases of the conjecture that are still open are the following.

**Conjecture 6.3.** *Let  $\mathbb{X}$  be a finite set of reduced points in  $\mathbb{P}^{s-1}$  and suppose that  $I = I(\mathbb{X})$  is a complete intersection generated by  $f_1, \dots, f_c$ ,  $c = s-1$ , with  $d_i = \deg(f_i)$  for  $i = 1, \dots, c$ , and  $2 \leq d_i \leq d_{i+1}$  for all  $i$ . Then*

- (a) [32, Conjecture 4.9]  $\delta_I(1) \geq (d_1 - 1)d_2 \cdots d_c$ .
- (b) If  $f_1, \dots, f_c$  are quadratic forms, then  $\delta_I(d) \geq 2^{c-d}$  for  $1 \leq d \leq c$  or equivalently  $\text{hyp}_I(d) \leq 2^c - 2^{c-d}$  for  $1 \leq d \leq c$ .

We prove part (a) of this conjecture, in a more general setting, when  $I$  is equigenerated.

**Proposition 6.4.** *Let  $I \subset S$  be an unmixed graded ideal of height  $c$ , minimally generated by forms of degree  $e \geq 2$ , whose associated primes are generated by linear forms. Then*

$$\text{hyp}_I(1) \leq e^{c-1}$$

and  $\delta_I(1) \geq \deg(S/I) - e^{c-1}$ . Furthermore  $\delta_I(1) \geq e^c - e^{c-1}$  if  $I$  is a complete intersection.

**Proof.** Since the associated primes of  $I$  are generated by linear forms and  $e \geq 2$ , one has  $\mathcal{F}_1(I) \neq \emptyset$ . Take any linear form  $h = t_k - \sum_{j \neq i} \lambda_j t_j$  in  $\mathcal{F}_1(I)$ ,  $\lambda_j \in K$ . For simplicity of notation assume  $k = 1$ . It suffices to show that  $\deg(S/(I, h)) \leq e^{c-1}$ . Let  $\{f_1, \dots, f_n\}$  be a minimal set of generators of  $I$  consisting of homogeneous polynomials with  $\deg(f_i) = e$  for all  $i$ . Setting  $f'_i = f_i(\sum_{j \neq 1} \lambda_j t_j, t_2, \dots, t_s)$  for  $i = 1, \dots, n$ ,  $S' = K[t_2, \dots, t_s]$ , and  $I' = (f'_1, \dots, f'_n)$ , there is an isomorphism  $\varphi$  of graded  $K$ -algebras

$$S/(I, h) \xrightarrow{\varphi} S'/I', \quad t_1 \mapsto \lambda_2 t_2 + \cdots + \lambda_s t_s, \quad t_i \mapsto t_i, \quad i = 2, \dots, s.$$

Note that  $\varphi(f + (I, h)) = f(\lambda_2 t_2 + \cdots + \lambda_s t_s, t_2, \dots, t_s) + I'$  for  $f$  in  $S$  and that  $\varphi$  has degree 0, that is,  $\varphi$  is degree preserving. Hence  $S/(I, h)$  and  $\deg(S'/I')$  have the same degree and the same dimension. Since  $\text{ht}(I, h) = \text{ht}(I)$ , we get  $\text{ht}(I') = \text{ht}(I) - 1$ , that is,  $\text{ht}(I') = c - 1$ . By definition  $f'_i$  is either 0 or has degree  $e$ , that is,  $I'$  is generated by forms of degree  $e$ . As  $K$  is infinite, there exists a minimal set of generators of  $I'$ ,  $\{g_1, \dots, g_t\}$ , such that  $\deg(g_i) = e$  for all  $i$  and  $g_1, \dots, g_{c-1}$  form a regular sequence (see Lemma 2.11). From the exact sequence

$$0 \longrightarrow I'/(g_1, \dots, g_{c-1}) \longrightarrow S'/(g_1, \dots, g_{c-1}) \longrightarrow S'/I' \longrightarrow 0,$$

we get  $e^{c-1} = \deg(S/(g_1, \dots, g_{c-1})) \geq \deg(S'/I') = \deg(S/(I, h))$ . This proves that  $\text{hyp}_I(1)$  is less than or equal to  $e^{c-1}$ . Hence  $\delta_I(1) \geq \deg(S/I) - e^{c-1}$ . Therefore, if  $I$  is a complete intersection,  $\deg(S/I) = e^c$  and we obtain the inequality  $\delta_I(1) \geq e^c - e^{c-1}$ .  $\square$

As a consequence, we recover the fact that Conjecture 6.3(a) holds for  $\mathbb{P}^2$  [32, Theorem 4.10].

**Corollary 6.5.** *Let  $I \subset S$  be a graded ideal of height 2, minimally generated by two forms  $f_1, f_2$  of degrees  $e_1, e_2$ , with  $2 \leq e_1 \leq e_2$ , whose associated primes are generated by linear forms. Then  $\text{hyp}_I(1) \leq e_2$  and  $\delta_I(1) \geq e_1 e_2 - e_2$ .*

**Proof.** It follows adapting the proof of Proposition 6.4.  $\square$

**Corollary 6.6.** *Let  $I \subset S$  be an unmixed graded ideal minimally generated by forms of degree  $e \geq 2$  whose associated primes are generated by linear forms. If  $1 \leq r \leq \text{ht}(I)$ , then*

$$\text{hyp}_I(1, r) \leq e^{\text{ht}(I)-r},$$

$\delta_I(1, r) \geq \deg(S/I) - e^{\text{ht}(I)-r}$ , and  $\delta_I(1, r) \geq e^{\text{ht}(I)} - e^{\text{ht}(I)-r}$  if  $I$  is a complete intersection.

**Proof.** This follows by adapting the proof of Proposition 6.4 and observing the following. If  $f_1, \dots, f_r$  are linearly independent linear forms and  $t_1 \succ \dots \succ t_s$  is the lexicographical order, we can find linear forms  $h_1, \dots, h_r$  such that  $\text{in}_\prec(h_1) \succ \dots \succ \text{in}_\prec(h_r)$  and  $(f_1, \dots, f_r)$  is equal to  $(h_1, \dots, h_r)$ .  $\square$

**Cayley-Bacharach conjectures.** In the following we explore the connections between a modified form of Conjecture 6.2 and a conjecture of Eisenbud-Green-Harris [8, Conjecture CB12].

**Conjecture 6.7** (Strong form of [8, Conjecture CB12]). *Let  $\Gamma$  be any subscheme of a zero-dimensional complete intersection of hypersurfaces of degrees  $d_1 \leq \dots \leq d_c$  in a projective space  $P^c$ . If  $\Gamma$  fails to impose independent conditions on hypersurfaces of degree  $m$ , then*

$$\deg(\Gamma) \geq (e+1)d_{k+2}d_{k+3} \cdots d_c$$

where  $e$  and  $k$  are defined by the relations

$$\sum_{i=k+2}^c (d_i - 1) \leq m + 1 < \sum_{i=k+1}^c (d_i - 1) \quad \text{and} \quad e = m + 1 - \sum_{i=k+2}^c (d_i - 1).$$

**Proposition 6.8.** *Conjectures 6.2 and 6.7 are equivalent for radical complete intersections.*

**Proof.** We first prove that Conjecture 6.7 for  $m = \sum_{i=k+1}^c (d_i - 1) - \ell - 1$  and  $e = d_{k+1} - \ell - 1$  implies Conjecture 6.2. Let  $I$  be a radical complete intersection ideal minimally generated by forms of degrees  $d_1 \leq \dots \leq d_c$ . Let  $H$  be any hypersurface defined by a form  $F$  of degree  $d$ . Let  $\mathbb{X}$  be the scheme defined by  $I(\mathbb{X}) = (I, F)$  and let  $\Gamma$  be the residual scheme defined by  $I(\Gamma) = I : F$ . By the Cayley-Bacharach Theorem [8, CB7],  $\Gamma$  must fail to impose independent conditions on hypersurfaces of degree  $\sum_{i=1}^c (d_i - 1) - d - 1 = m$ . Now Conjecture 6.7 implies  $\deg(S/I : F) = \deg(\Gamma) \geq ed_{k+2}d_{k+3} \cdots d_c$ , which in view of Theorem 3.5 gives

$$\delta_I(d) \geq (e+1)d_{k+2}d_{k+3} \cdots d_c = (d_{k+1} - \ell)d_{k+2}d_{k+3} \cdots d_c.$$

For the converse, we prove that Conjecture 6.2 with  $d = \sum_{i=1}^c (d_i - 1) - m - 1$  and  $\ell = d_{k+1} - e - 1$  recovers Conjecture 6.7. Let  $\Gamma$  be any subscheme of a complete intersection, and suppose that  $\Gamma$  fails to impose independent conditions on hypersurfaces of degree  $m$ . Assuming that  $\Gamma$  spans a projective space  $\mathbb{P}^c$ , take a radical complete intersection ideal  $I$  contained in  $I_\Gamma$ , and let  $\mathbb{X}$  be the scheme defined by  $I(\mathbb{X}) = I : I(\Gamma)$ . By [8, CB7],  $\mathbb{X}$  lies on a hypersurface of degree  $\sum_{i=1}^c (d_i - 1) - m - 1 = d$ . Then Conjecture 6.2 and Theorem 3.5 give

$$\deg(\Gamma) = \deg(S/I : F) \geq (d_{k+1} - \ell)d_{k+2}d_{k+3} \cdots d_c = (e + 1)d_{k+2}d_{k+3} \cdots d_c. \quad \square$$

Conjecture 6.7 has been recently proven in [19, Theorem 5.1] for  $k = 1$  under additional assumptions on the Picard group of the complete intersection. We now consider the case when  $d_1 = \cdots = d_c = 2$ . In this case Conjecture 6.2 specializes to Conjecture 6.3(b) and Conjecture 6.7 is related to [8, Conjecture CB10].

**Proposition 6.9.** *The following statements are equivalent:*

- (1) [Conjecture 6.3(b)] Let  $I$  be a complete intersection generated by  $c$  quadratic forms. Then  $\delta_I(d) \geq 2^{c-d}$  for  $1 \leq d \leq c$  or equivalently  $\text{hyp}_I(d) \leq 2^c - 2^{c-d}$  for  $1 \leq d \leq c$ .
- (2) [8, Conjecture CB10] If  $\mathbb{X}$  is an ideal-theoretic complete intersection of  $c = s - 1$  quadrics in  $\mathbb{P}^{s-1}$  and  $f \in S := K[t_1, \dots, t_s]$  is a homogeneous polynomial of degree  $d$  such that  $\deg(S/(I(\mathbb{X}), f)) > 2^c - 2^{c-d}$ , then  $f \in I_{\mathbb{X}}$ .

**Proof.** Let  $I = I(\mathbb{X})$  be a complete intersection ideal of  $c$  quadratic homogeneous polynomials. Then  $\deg(S/I) = 2^c$  and (2) is equivalent to the statement for any  $f \in \mathcal{F}_d$ ,  $\deg(S/(I(\mathbb{X}), f)) \leq 2^c - 2^{c-d}$ . Using Definition 1.1, this is in turn equivalent to  $\delta_I(d) \geq 2^{c-d}$ , for  $d = 1, \dots, c$ , which is precisely the statement of (1).  $\square$

As an application of our earlier results we recover the following cases of Conjecture 6.3(b) under the more general hypothesis that  $I(\mathbb{X})$  is a not necessarily a radical complete intersection.

**Corollary 6.10.** [8] *If  $I$  is a complete intersection ideal generated by  $c$  quadratic forms, then  $\delta_I(d) \geq 2^{c-d}$ , for  $d = 1, c - 1$ , and  $c$ .*

**Proof.** Case  $d = 1$  follows by taking  $e = 2$  in Proposition 6.4. Case  $d = c - 1$  follows from Corollary 4.20 because complete intersections are Gorenstein and vanishing ideals of finite sets of points are Geramita. For Case  $d = c$  recall that By Proposition 4.6 and Theorem 4.10,  $\text{reg}(S/I)$  is the regularity index of  $\delta_I$ . Thus Case  $d = c$  follows since  $\text{reg}(S/I) = c$ .  $\square$

If  $\mathbb{X} \subset \mathbb{P}^c$  is a reduced set of points such that  $I(\mathbb{X})$  is a complete intersection ideal and the points of  $\mathbb{X}$  are in linearly general position, i.e., any  $c + 1$  points of  $\mathbb{X}$  span  $\mathbb{P}^c$ ,

then [1, Theorem 1] adds more cases when [8, Conjecture CB10] holds. If we assume that the quadrics that cut out  $\mathbb{X}$  are generic, then the assumptions of that result are satisfied, and we can conclude by Proposition 6.9 that if  $1 \leq d \leq c-1$ , then  $\delta_I(d) \geq c(c-1-d)+2$ . Suppose  $d := c-k$ , for some  $1 \leq k \leq c-1$ . Then this inequality becomes  $\delta_I(c-k) \geq c(k-1)+2$ . We observe that this inequality is sufficiently strong to establish [Conjecture 6.3(b)] *asymptotically* for  $c$  sufficiently large:

- $k = 2$ . If  $c \geq 2$ , then  $\delta_I(c-2) \geq c+2 \geq 2^2$ .
- $k = 3$ . If  $c \geq 3$ , then  $\delta_I(c-3) \geq 2c+2 \geq 2^3$ .
- $k = 4$ . If  $c \geq 5$ , then  $\delta_I(c-4) \geq 3c+2 \geq 2^4$ .
- $k \geq 5$ . If  $c \geq (2^k-2)/(k-1)$ , then  $\delta_I(c-k) \geq 2^k$ .

## Acknowledgments

We thank the Banff International Research Station (BIRS) where this project was started during under the auspices of their focused research groups program. Computations with *Macaulay2* [16] were crucial to verifying many of the examples given in this paper. The authors acknowledge the support of the following funding agencies: Cooper was partially funded by NSERC grant RGPIN-2018-05004, Seceleanu was partially supported by NSF grant DMS-1601024 and EPSCoR grant OIA-1557417.

## Appendix A. Procedures for *Macaulay2*

**Procedure A.1.** Computing the footprint matrix with *Macaulay2* [16]. This procedure corresponds to Example 3.15. It can be applied to any vanishing ideal  $I$  to obtain the entries of the matrix  $(\text{fp}_I(d, r))$  and is reasonably fast.

```
S=QQ[t1,t2,t3], I=ideal(t1^3,t2*t3)
M=coker gens gb I
regularity M, degree M, init=ideal(leadTerm gens gb I)
er=(x)-> if not quotient(init,x)==init then degree ideal(init,x) else 0
fpr=(d,r)->degree M - max apply(apply(apply(
subsets(flatten entries basis(d,M),r),toSequence),ideal),er)
hilbertFunction(1,M),fpr(1,1),fpr(1,2),fpr(1,3)
--gives the first row of the footprint matrix
```

**Procedure A.2.** Computing the GMD function with *Macaulay2* [16] over a finite field and computing an upper bound over any field using products of linear forms. This procedure corresponds to Example 3.11.

```
q=3,S=ZZ/3[t1,t2,t3,t4,t5,t6],I=ideal(t1*t6-t3*t4,t2*t6-t3*t5)
G=gb I, M=coker gens gb I
regularity M, degree M, init=ideal(leadTerm gens gb I)
```

```

genmd=(d,r)->degree M-max apply(apply(subsets(apply(apply(apply(
toList (set(0..q-1))^**hilbertFunction(d,M))-(
set{0})^**hilbertFunction(d,M)),toList),x->basis(d,M)*vector x),
z->ideal(flatten entries z)),r),ideal),
x-> if #set flatten entries mingens ideal(leadTerm gens x)==r
and not quotient(I,x)==I then degree(I+x) else 0)
hilbertFunction(1,M),fpr(1,1),fpr(1,2),fpr(1,3),fpr(1,4),fpr(1,5),
fpr(1,6)
genmd(1,1), L={t1,t2,t3,t4,t5,t6}
linearforms=(d,r)->degree M - max
apply(apply(apply((subsets(apply(apply(
(subsets(L,d)),product),x-> x % G),r)),toList),ideal),
x-> if #set flatten entries mingens ideal(leadTerm gens x)==r
and not quotient(I,x)==I then degree(I+x) else 0)
linearforms(1,2),linearforms(1,3),linearforms(1,4),linearforms(1,5)
--gives upper bound for genmd

```

**Procedure A.3.** Computing the minimum socle degree and the v-number of an ideal  $I$  with Macaulay2 [16]. This procedure corresponds to Example 4.3.

```

S=QQ[t1,t2,t3,t4]
p1=ideal(t2,t3,t4),p2=ideal(t1,t3,t4),p3=ideal(t1,t2,t4),
p4=ideal(t1,t2,t3)
I=intersect(ideal(t2^10,t3^9,t4^4,t2*t3*t4^3),
ideal(t1^4,t3^4,t4^3,t1*t3*t4^2),ideal(t1^4,t2^5,t4^3),
ideal(t1^3,t2^5,t3^10))
h=ideal(t1+t2+t3+t4)--regular element on S/I
J=quotient(I+h,m), regularity coker gens gb I
soc=J/(I+h), degrees mingens soc
J1=quotient(I,p1), soc1=J1/I, degrees mingens soc1

```

**Procedure A.4.** Computing the v-number of a vanishing ideal  $I(\mathbb{X})$ , the regularity index of  $\delta_{\mathbb{X}}$ , and the minimum distance  $\delta_{\mathbb{X}}(d)$  of the Reed–Muller-type code  $C_{\mathbb{X}}(d)$  with Macaulay2 [16]. This procedure corresponds to Example 4.5.

```

q=3, G=ZZ/q, S=G[t3,t2,t1,MonomialOrder=>Lex]
p1=ideal(t2,t1-t3),p2=ideal(t2,t3),p3=ideal(t2,2*t1-t3)
p4=ideal(t1-t2,t3),p5=ideal(t1-t3,t2-t3), p6=ideal(2*t1-t3,2*t2-t3)
p7=ideal(t1,t2),p8=ideal(t1,t3),p9=ideal(t1,t2-t3),p10=ideal(t1,2*t2-t3)
I=intersect(p1,p2,p3,p4,p5,p6,p7,p8,p9,p10)
M=coker gens gb I, regularity M, degree M
init=ideal(leadTerm gens gb I)
genmd=(d,r)->degree M-max apply(apply(subsets(apply(apply(apply(

```

```

toList (set(0..q-1))^**(hilbertFunction(d,M))-
(set{0})^**(hilbertFunction(d,M)),toList),x->basis(d,M)*vector x),
z->ideal(flatten entries z)),r),ideal),
x-> if #set flatten entries mingens ideal(leadTerm gens x)==r
and not quotient(I,x)==I then degree(I+x) else 0)
genmd(1,1), genmd(2,1)
J1=quotient(I,p1), soc1=J1/I, degrees mingens soc1--gives 4
J7=quotient(I,p7), soc7=J7/I, degrees mingens soc7--gives 3

```

## References

- [1] E. Ballico, C. Fontanari, The Horace method for error-correcting codes, *Appl. Algebra Engrg. Comm. Comput.* 17 (2006) 135–139.
- [2] W. Bruns, J. Herzog, *Cohen-Macaulay Rings*, revised edition, Cambridge University Press, 1997.
- [3] C. Carvalho, On the second Hamming weight of some Reed-Muller type codes, *Finite Fields Appl.* 24 (2013) 88–94.
- [4] D. Cox, J. Little, D. O’Shea, *Ideals, Varieties, and Algorithms*, Springer-Verlag, 1992.
- [5] I.M. Duursma, C. Rentería, H. Tapia-Recillas, Reed-Muller codes on complete intersections, *Appl. Algebra Engrg. Comm. Comput.* 11 (6) (2001) 455–462.
- [6] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, 1995.
- [7] D. Eisenbud, *The Geometry of Syzygies: A Second Course in Commutative Algebra and Algebraic Geometry*, Graduate Texts in Mathematics, vol. 229, Springer-Verlag, New York, 2005.
- [8] D. Eisenbud, M. Green, J. Harris, Cayley–Bacharach theorems and conjectures, *Bull. Amer. Math. Soc. (N.S.)* 33 (3) (1996) 295–324.
- [9] B. Engheta, On the projective dimension and the unmixed part of three cubics, *J. Algebra* 316 (2) (2007) 715–734.
- [10] A.V. Geramita, M. Kreuzer, L. Robbiano, Cayley–Bacharach schemes and their canonical modules, *Trans. Amer. Math. Soc.* 339 (1) (1993) 163–189.
- [11] L. Gold, J. Little, H. Schenck, Cayley–Bacharach and evaluation codes on complete intersections, *J. Pure Appl. Algebra* 196 (1) (2005) 91–99.
- [12] M. González-Sarabia, E. Camps, E. Sarmiento, R.H. Villarreal, The second generalized Hamming weight of some evaluation codes arising from a projective torus, *Finite Fields Appl.* 52 (2018) 370–394.
- [13] M. González-Sarabia, J. Martínez-Bernal, R.H. Villarreal, C.E. Vivares, Generalized minimum distance functions, *J. Algebraic Combin.* (2019), in press.
- [14] M. González-Sarabia, C. Rentería, The dual code of some Reed–Muller type codes, *Appl. Algebra Engrg. Comm. Comput.* 14 (2004) 329–333.
- [15] M. González-Sarabia, C. Rentería, H. Tapia-Recillas, Reed-Muller-type codes over the Segre variety, *Finite Fields Appl.* 8 (4) (2002) 511–518.
- [16] D. Grayson, M. Stillman, Macaulay2, a software system for research in algebraic geometry, available at <http://www.math.uiuc.edu/Macaulay2/>.
- [17] E. Guardo, L. Marino, A. Van Tuyl, Separators of fat points in  $\mathbb{P}^n$ , *J. Algebra* 324 (7) (2010) 1492–1512.
- [18] T. Helleseth, T. Kløve, J. Mykkelheit, The weight distribution of irreducible cyclic codes with block lengths  $n_1((q^l - 1)/N)$ , *Discrete Math.* 18 (1977) 179–211.
- [19] J. Hotchkiss, B. Ullery, The gonality of complete intersection curves, preprint, arXiv:1706.08169, 2018.
- [20] T. Kløve, The weight distribution of linear codes over  $GF(q^l)$  having generator matrix over  $GF(q)$ , *Discrete Math.* 23 (2) (1978) 159–168.
- [21] H.H. López, C. Rentería, R.H. Villarreal, Affine cartesian codes, *Des. Codes Cryptogr.* 71 (1) (2014) 5–19.
- [22] F.J. MacWilliams, N.J.A. Sloane, *The Theory of Error-Correcting Codes*, North-Holland, 1977.
- [23] J. Martínez-Bernal, Y. Pitones, R.H. Villarreal, Minimum distance functions of graded ideals and Reed-Muller-type codes, *J. Pure Appl. Algebra* 221 (2017) 251–275.

- [24] J. Martínez-Bernal, Y. Pitones, R.H. Villarreal, Minimum distance functions of complete intersections, *J. Algebra Appl.* 17 (11) (2018) 1850204.
- [25] H. Matsumura, *Commutative Algebra*, Benjamin-Cummings, Reading, MA, 1980.
- [26] H. Matsumura, *Commutative Ring Theory*, Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, 1986.
- [27] L. Núñez-Betancourt, Y. Pitones, R.H. Villarreal, Footprint and minimum distance functions, *Commun. Korean Math. Soc.* 33 (1) (2018) 85–101.
- [28] L. O’Carroll, F. Planas-Vilanova, R.H. Villarreal, Degree and algebraic properties of lattice and matrix ideals, *SIAM J. Discrete Math.* 28 (1) (2014) 394–427.
- [29] C. Rentería, A. Simis, R.H. Villarreal, Algebraic methods for parameterized codes and invariants of vanishing ideals over finite fields, *Finite Fields Appl.* 17 (1) (2011) 81–104.
- [30] A. Sørensen, Projective Reed-Muller codes, *IEEE Trans. Inform. Theory* 37 (6) (1991) 1567–1576.
- [31] R. Stanley, Hilbert functions of graded algebras, *Adv. Math.* 28 (1978) 57–83.
- [32] S. Tohăneanu, A. Van Tuyl, Bounding invariants of fat points using a coding theory construction, *J. Pure Appl. Algebra* 217 (2) (2013) 269–279.
- [33] M. Tsfasman, S. Vladut, D. Nogin, *Algebraic Geometric Codes: Basic Notions*, Mathematical Surveys and Monographs, vol. 139, American Mathematical Society, Providence, RI, 2007.
- [34] R.H. Villarreal, *Monomial Algebras*, second edition, Monographs and Research Notes in Mathematics, Chapman and Hall/CRC, 2015.
- [35] V.K. Wei, Generalized Hamming weights for linear codes, *IEEE Trans. Inform. Theory* 37 (5) (1991) 1412–1418.