

Big Torelli groups: generation and commensuration

Javier Aramayona, Tyrone Ghaswala, Autumn E. Kent,
Alan McLeay, Jing Tao, and Rebecca R. Winarski

Abstract. For any surface Σ of infinite topological type, we study the *Torelli subgroup* $\mathcal{I}(\Sigma)$ of the mapping class group $\mathrm{MCG}(\Sigma)$, whose elements are those mapping classes that act trivially on the homology of Σ . Our first result asserts that $\mathcal{I}(\Sigma)$ is topologically generated by the subgroup of $\mathrm{MCG}(\Sigma)$ consisting of those elements in the Torelli group which have compact support. Next, we prove the abstract commensurator group of $\mathcal{I}(\Sigma)$ coincides with $\mathrm{MCG}(\Sigma)$. This extends the results for finite-type surfaces [9, 6, 7, 16] to the setting of infinite-type surfaces.

Mathematics Subject Classification (2010). Primary: 20F65; Secondary: 57M50.

Keywords. Infinite-type surfaces, mapping class groups, Torelli group.

1. Introduction

Let Σ be a connected oriented surface of infinite topological type – that is, a surface with fundamental group that is not finitely generated. We will also assume the boundary components of Σ are compact. The *mapping class group* of Σ is the group:

$$\mathrm{MCG}(\Sigma) = \mathrm{Homeo}(\Sigma, \partial\Sigma) / \mathrm{Homeo}_0(\Sigma, \partial\Sigma),$$

where $\mathrm{Homeo}(\Sigma, \partial\Sigma)$ is the group of self-homeomorphisms of Σ which fix $\partial\Sigma$ pointwise, equipped with the compact-open topology, and $\mathrm{Homeo}_0(\Sigma, \partial\Sigma)$ is the connected component of the identity in $\mathrm{Homeo}(\Sigma, \partial\Sigma)$. We equip $\mathrm{MCG}(\Sigma)$ with the quotient topology.

There is a natural homomorphism $\mathrm{MCG}(\Sigma) \rightarrow \mathrm{Aut}(H_1(\Sigma, \mathbb{Z}))$, whose kernel is commonly referred to as the *Torelli group* $\mathcal{I}(\Sigma) < \mathrm{MCG}(\Sigma)$. While Torelli groups of finite-type surfaces have been the object of intense study (see for example [3, 4, 12, 15, 17, 20, 22, 24, 26]) not much is known about them in the case of surfaces of infinite type. The present article aims to be a first step in this direction.

Generation. In a recent article, Patel and Vlamis [23] give a topological generating set for the *pure* mapping class group $\text{PMCG}(\Sigma)$ (i.e. the subgroup of $\text{MCG}(\Sigma)$ consisting of those mapping classes which fix every *end* of Σ ; see Section 2). More concretely, they show that $\text{PMCG}(\Sigma)$ is topologically generated by the subgroup of elements with compact support if Σ has at most one end *accumulated by genus*; otherwise, $\text{PMCG}(\Sigma)$ is topologically generated by the union of the set of compactly-supported elements and the set of *handle shifts*; see Section 2.

Observe that $\mathcal{I}(\Sigma) < \text{PMCG}(\Sigma)$. Denote by $\mathcal{I}_c(\Sigma)$ the subgroup of $\mathcal{I}(\Sigma)$ consisting of those elements with compact support, and let $\overline{\mathcal{I}_c(\Sigma)}$ be the closure of $\mathcal{I}_c(\Sigma)$ in $\text{PMCG}(\Sigma)$. Our first result asserts that, for any infinite-type surface Σ , the set of compactly-supported mapping classes contained in the Torelli group topologically generates the Torelli group:

Theorem 1.1. *For any connected oriented surface Σ of infinite type, we have $\mathcal{I}(\Sigma) = \overline{\mathcal{I}_c(\Sigma)}$.*

Birman [4] and Powell [24] showed that the Torelli group of a finite-type surface with at most one boundary or puncture is generated by *separating twists* (i.e. Dehn twists about separating curves), plus *bounding pair maps* (that is, products of twists of the form $T_\gamma T_\delta^{-1}$, where γ and δ are disjoint non-separating curves but their union separates and γ and δ are homologous). We will show the same is true for finite-type surfaces with arbitrary number of boundary components and punctures. By exhausting Σ by an appropriate sequence of finite-type subsurfaces, we obtain:

Theorem 1.2. *Let Σ be a connected oriented surface of infinite topological type. Then $\mathcal{I}(\Sigma)$ is topologically generated by separating twists and bounding-pair maps.*

Theorem 1.1 implies $\mathcal{I}(\Sigma)$ is a closed subgroup of $\text{MCG}(\Sigma)$. Since $\text{MCG}(\Sigma)$ is a Polish group [1] and closed subgroups of Polish groups are Polish, we have the following corollary.

Corollary 1.3. *Let Σ be a connected oriented surface of infinite topological type. Then $\mathcal{I}(\Sigma)$ is a Polish group.*

Commensurations. Recall that, given a group G , its abstract commensurator $\text{Comm}(G)$ is the group of equivalence classes of isomorphisms between finite-index subgroups of G ; here, two isomorphisms are equivalent if they agree on a finite-index subgroup. Observe that there is a natural homomorphism

$$\text{Aut}(G) \longrightarrow \text{Comm}(G).$$

We will prove the following result.

Theorem 1.4. *For any connected oriented surface Σ of infinite topological type and without boundary we have*

$$\mathrm{Comm}\mathcal{J}(\Sigma) \cong \mathrm{Aut}\mathcal{J}(\Sigma) \cong \mathrm{MCG}(\Sigma).$$

Before continuing we stress that to the best of our knowledge, it is not known whether the Torelli group of an infinite-type surface has any finite-index subgroups, and so in particular it is possible that $\mathrm{Comm}\mathcal{J}(\Sigma)$ may in fact coincide *a priori* with $\mathrm{Aut}\mathcal{J}(\Sigma)$.

Historical context and idea of proof. Theorem 1.4 was previously known to hold for finite-type surfaces. Indeed, Farb and Ivanov [9] proved it for closed surfaces of genus at least five, which was then extended (and generalized to the Johnson Kernel) by Brendle and Margalit to all closed surfaces of genus at least three [6, 7]. Kida extended the result of Brendle and Margalit to all finite-type surfaces of genus at least four [16]. Finally, recent work of Brendle and Margalit and McLeay has further generalized the result to apply to a large class of normal subgroups of finite-type surfaces [5, 21].

In order to prove the theorem, we closely follow Brendle and Margalit’s strategy. First, we adapt ideas of Bavard, Dowdall, and Rafi [2] to show that every commensuration of the Torelli group respects the property of being a separating twist or a bounding pair map. From this we deduce that every commensuration induces an automorphism of a combinatorial object called the *Torelli complex*. This complex was originally introduced, for closed surfaces, by Brendle and Margalit [6], who proved that its automorphism group coincides with the mapping class group; this was later extended by Kida [16] to finite-type surfaces with punctures. Using this, plus an inductive argument due to Ivanov [14], we will show that every automorphism of the Torelli complex of an infinite-type surface is induced by a surface homeomorphism. At this point, Theorem 1.4 will follow easily using a well-known argument of Ivanov [14].

Acknowledgments. This note is the result of a conversation during the conference “Geometry of Teichmüller spaces and mapping class groups” at the University of Warwick in the Spring of 2018. We are grateful to the organizers, Tara Brendle, Mladen Bestvina, and Chris Leininger for the opportunity to participate. We also thank Dan Margalit for several helpful comments.

We are very thankful to the referee for comments and suggestions that helped improve the paper, especially for an alternate argument establishing one of the containments in Theorem 1.1; see the end of Section 3. We also thank the referee for telling us about Lemma 2.2, whose proof is due to Federica Fanoni. Finally, we thank Justin Malestein for providing the proofs in Section 3.1

Kent, Tao, and Winarski were partially supported by NSF grants DMS-1107452, DMS-1107263, DMS-1107367 “RNMS: Geometric Structures and Representation Varieties” (the GEAR Network). Aramayona was partially supported by grants RYC-2013-13008 and MTM-2015-67781. Kent acknowledges support from NSF grant DMS-1350075 and the Vilas Trustees at the University of Wisconsin. Tao acknowledges partial support from NSF grant DMS-1440140. Winarski acknowledges partial support from the AMS-Simons Travel Grants which are administered by the American Mathematical Society, with support from the Simons Foundation.

2. Definitions

In this section we introduce the main objects needed for the proofs of our results.

2.1. Surfaces. Throughout, by a *surface* we mean a connected, oriented, second-countable topological surface with (possibly empty) compact boundary. We say that Σ has *finite type* if its fundamental group is finitely generated; otherwise, we say that Σ has *infinite type*. In the finite type case, we will sometimes use the notation $\Sigma = \Sigma_{g,p}^b$, where g , p , and b are, respectively, the genus, the number of punctures, and the number of boundary components of Σ . In this case, we define the *complexity* of Σ to be the integer $\xi(\Sigma) = 3g - 3 + p + b$.

The *space of ends* of Σ is the set

$$\text{Ends}(\Sigma) = \varprojlim \pi_0(\Sigma \setminus K),$$

where the inverse limit is taken over the set of compact subsets $K \subset \Sigma$, directed with respect to inclusion. Here, the topology on $\text{Ends}(\Sigma)$ is given by the limit topology obtained by equipping each $\pi_0(\Sigma \setminus K)$ with the discrete topology. See [27] for further details.

We say that e in $\text{Ends}(\Sigma)$ is *accumulated by genus* if every neighborhood of e has infinite genus; otherwise, we say that e is *planar*. In particular, observe that every puncture of Σ is a planar end. We denote by $\text{Ends}_g(\Sigma)$ the subset of $\text{Ends}(\Sigma)$ consisting of ends accumulated by genus. It is a classical theorem (see [27] for a discussion and proof) that the homeomorphism type of Σ is determined by the tuple

$$(g(\Sigma), b(\Sigma), \text{Ends}(\Sigma), \text{Ends}_g(\Sigma)),$$

where $g(\Sigma)$ and $b(\Sigma)$ denote the genus and the number of boundary components of Σ .

2.2. Curves and domains. By a *curve* on Σ we mean the free homotopy class of a simple closed curve that does not bound a disk or a disk containing a single planar end of Σ . Abusing notation, we will not make any distinction between a curve and any of its representatives.

We say that a curve γ is *separating* if $\Sigma \setminus \gamma$ has two connected components; otherwise, we say that γ is *non-separating*. We say that two curves are *disjoint* if they have disjoint representatives in Σ . A *multicurve* is a set of pairwise disjoint curves. Given two curves α and β , we denote by $i(\alpha, \beta)$ their geometric intersection number. The intersection number between two multicurves is defined additively.

A *domain* Y in Σ is a closed subset which is itself a surface and the inclusion map is a proper, π_1 -injective embedding. Note that domains are only defined up to isotopy. A *subsurface* of Σ is a disjoint union of domains. The following definition appears in [2].

Definition 2.1. A domain Y of Σ is called *principal* if Y has finite-type and every component of $\Sigma \setminus Y$ has infinite-type.

The following lemma was communicated to us by the referee. The statement and the proof are due to Federica Fanoni.

Lemma 2.2. *Let σ be multicurve. Then there are curves realizing the homotopy classes in σ which do not accumulate in any compact set of Σ if and only if for every $\alpha \notin \sigma$, the set $\{\beta \in \sigma \mid i(\alpha, \beta) \neq 0\}$ is finite.*

Proof. The forward direction is clear so we will focus on the other direction. Choose a complete hyperbolic metric on Σ without half planes and realize all curves as geodesics. The assumption on the metric implies that we can decompose Σ into pairs of pants with geodesic boundary, funnels and/or cusps (as shown in [13]). If the geodesic representative of σ has an accumulation point, then we can find a pair of pants P of Σ containing such a point. However, by assumption, there are only finitely many curves of σ that intersect the boundary of P . Thus, only finitely many curves of σ intersect P . In particular, they cannot accumulate in P , a contradiction. $\square$

Given an element $f \in \text{MCG}(\Sigma)$, there is a canonical (possibly empty) multicurve ∂f in Σ for which $f(\partial f) = \partial f$, defined as follows. Let $\mathcal{O}(f)$ be the set of curves α such that $\{f^k(\alpha) \mid k \in \mathbb{Z}\}$ is finite. Then ∂f is the set of curves in $\mathcal{O}(f)$ that are disjoint from all other elements of $\mathcal{O}(f)$. See [11, Section 2] and also [2] for further details.

Given $f \in \text{MCG}(\Sigma)$ and a subsurface Y , we say Y *supports* f or f is *supported* on Y if f can be realized by a homeomorphism which is the identity outside of Y . An element $f \in \text{MCG}(\Sigma)$ is said to have *compact support* if there is a compact subsurface that supports f . Similarly, f has *finite support* if there is

a subsurface of finite-type that supports f . If f is a product of Dehn twists about a multicurve α , then we will also say α supports f .

The following statement follows from [2]. The first assertion is Lemma 2.6, while the second follows from its proof.

Lemma 2.3. *If $f \in \text{MCG}(\Sigma)$ is nontrivial and has finite support, then f has infinite order and ∂f is a nonempty multicurve in Σ . Furthermore, let Y be the (finite) union of all finite-type domains in $\Sigma \setminus \partial f$, then Y supports f .*

2.3. Pure mapping classes. The *pure mapping class group* $\text{PMCG}(\Sigma)$ is the normal subgroup of $\text{MCG}(\Sigma)$ whose elements fix every end of Σ .

The set of compactly-supported mapping classes $\text{PMCG}_c(\Sigma)$ forms a subgroup of $\text{PMCG}(\Sigma)$ which is normal in $\text{MCG}(\Sigma)$. Note that if f lies in $\text{PMCG}(\Sigma)$ and f has finite support, then it necessarily has compact support. Since $\text{PMCG}_c(\Sigma)$ is a direct limit of pure mapping class groups of compact surfaces, a classical result due to Dehn and Lickorish (see [10, Section 4], for instance) implies that $\text{PMCG}_c(\Sigma)$ is generated by *Dehn twists*.

2.4. Handle Shifts. For any subgroup $\Gamma < \text{MCG}(\Sigma)$, we denote by $\bar{\Gamma}$ its topological closure in $\text{MCG}(\Sigma)$.

Patel and Vlamis introduced *handle shifts* and showed that handle shifts and Dehn twists topologically generate $\text{PMCG}(\Sigma)$, see [23]. Subsequently, in [1] it was shown that $\text{PMCG}(\Sigma) = \overline{\text{PMCG}_c(\Sigma)} \rtimes H$, where H is a particular subgroup isomorphic to a direct product of pairwise commuting *handle shifts*. We now recall the definition of a handle shift.

Let Λ be the surface obtained from $\mathbb{R} \times [-1, 1]$ by removing disks of radius $\frac{1}{4}$ centered at $(t, 0)$ for $t \in \mathbb{Z}$ and gluing in a torus with one boundary component, identifying the boundary of the torus with the boundary of the removed disk. Let $\sigma: \Lambda \rightarrow \Lambda$ be the homeomorphism that shifts the handle at $(t, 0)$ to the handle at $(t + 1, 0)$, and is the identity on $\mathbb{R} \times \{-1, 1\}$ (see [1] or [23] for an image of such a homeomorphism). The isotopy class of σ is called a *handle shift* of Λ .

An element h in $\text{MCG}(\Sigma)$ is a *handle shift* if there exists a proper embedding $\iota: \Lambda \rightarrow \Sigma$ which induces an injective map on ends, and such that $[h] = [\delta]$ where $\delta|_{\iota(\Lambda)} = \sigma$ and δ is the identity outside $\iota(\Lambda)$. As a consequence of our definition, we must have $|\text{Ends}_g(\Sigma)| \geq 2$; also, for each handle shift there is an attracting end ϵ_+ and a repelling end ϵ_- in $\text{Ends}_g(\Sigma)$, and they are distinct.

We say a handle shift h with attracting end ϵ_+ and repelling end ϵ_- is *dual* to a separating curve γ if each component of $\Sigma \setminus \gamma$ contains exactly one of ϵ_+ and ϵ_- .

2.5. Principal exhaustions. We now introduce a minor modification of the notion of *principal exhaustion* from [1, 2].

Definition 2.4. A *principal exhaustion* of Σ is an infinite sequence of principal domains $\{P_1, P_2, \dots\}$ such that, for every $i \geq 1$,

- (1) $P_i \subset P_{i+1}$;
- (2) every component of ∂P_i is separating
- (3) no component of ∂P_i is isotopic to a component of ∂P_{i+1} ;
- (4) $\Sigma = \bigcup P_i$.

Lemma 2.5. Let Σ be a connected infinite-type surface and let $\{P_i\}$ be a principal exhaustion of Σ . Then for all i ,

- for all $j > i$, $H_1(P_j) \cong H_1(P_i) \oplus M$ for some $M < H_1(P_j \setminus P_i)$;
- $H_1(\Sigma) \cong H_1(P_i) \oplus M'$ for some $M' < H_1(\Sigma \setminus P_i)$.

Proof of Lemma 2.5. We will let W be either P_j or Σ to prove both cases simultaneously.

Let $\partial_1 P_i, \dots, \partial_m P_i$ be the boundary components of P_i .

Since every component of $\Sigma - P_i$ is of infinite type, every component of $\overline{W - P_i}$ either contains an end of Σ or a boundary component of W . So there is a collection of pairwise disjoint rays and arcs $\gamma_1, \dots, \gamma_m$ properly embedded in $\overline{W - P_i}$ such that $\gamma_k \cap \partial_k P_i$ is a single point for all k .

By the Regular Neighborhood Theorem, we may deformation retract W along the γ_k , fixing P_i throughout, to obtain a subsurface Δ homotopy equivalent to W that contains P_i and such that $P_i \cap \overline{\Delta - P_i}$ is a disjoint union of arcs $\alpha_1, \dots, \alpha_m$, as pictured in Figure 1.




Figure 1. Pictured at the top is the surface W , the subsurface P_i , and the arcs $\gamma_1, \dots, \gamma_m$. Below is the surface Δ , obtained by deleting open neighborhoods of the interiors of the γ_k . The α_k are the dotted arcs.

Consideration of the Mayer–Vietoris sequence gives us an exact sequence

$$0 \longrightarrow H_1(P_i) \oplus H_1(\Delta - P_i) \longrightarrow H_1(W) \xrightarrow{\partial} H_0(\alpha_1 \sqcup \cdots \sqcup \alpha_m).$$

This gives us the direct sum decomposition of $H_1(W)$. Since $\partial_\ell P_i$ are separating, then so are the α_ℓ . This implies that the boundary map ∂ is zero, and since $H_1(\Delta - P_i)$ is naturally a subgroup of $H_1(W - P_i)$, the proof is complete. $\square$

3. Compactly generating the Torelli group

Let Σ be an infinite-type surface. We define the *compactly supported Torelli group*

$$\mathcal{I}_c(\Sigma) := \{f \in \mathcal{I}(\Sigma) \mid f \text{ has compact support}\}.$$

The aim of this section is to prove the first main result of the introduction, whose statement we now recall:

Theorem 3.1. *For any connected oriented surface Σ of infinite type, we have $\mathcal{I}(\Sigma) = \overline{\mathcal{I}_c(\Sigma)}$.*

Remark 3.2. The referee suggested an alternate way of proving that every element of $\mathcal{I}(\Sigma)$ is a limit of mapping classes with compact support; we give this argument at the end of this section. However, we have decided to keep our original argument, phrased in terms of a new type of mapping class (*pseudo handle shifts*, see below), which may be of independent interest.

We will need to know that certain, possibly infinite, products of handle shifts are inaccessible by compactly supported mapping classes. For a general product of handle shifts, this is too much to hope for. For example, in a surface with two ends, the product of two commuting handle shifts with opposite dynamics is a limit of compactly supported classes.

More generally, there are products of infinitely many commuting handle shift that are limits of compactly supported classes. For example, there is the “boundary leaf shift,” which we now explain.

Example 3.3 (boundary leaf shift). Start with an infinite regular tree $\mathcal{T}$ properly embedded in the hyperbolic plane $\mathbb{H}^2$ with boundary a Cantor set in $\partial\mathbb{H}^2$. Orient $\partial\mathbb{H}^2$ counterclockwise. Build a surface by taking the boundary of a regular neighborhood of $\mathcal{T}$ in $\mathbb{H}^2 \times \mathbb{R}$ and attach handles periodically (in the hyperbolic metric) along each side of $\mathcal{T}$, see Figure 2. The orientation on $\partial\mathbb{H}^2$ defines a product $\mathcal{H}$ of handle shifts by shifting the handles in each region of $\mathbb{H}^2 - \mathcal{T}$ in the clockwise direction.

To see the boundary leaf shift is in $\overline{\text{PMCG}_c(\Sigma)}$, pick a basepoint $*$ in $\mathcal{T}$ and consider the n -neighborhood $B(n)$ of $*$ in $\mathcal{T}$. Then we may move the handles incident to $B(n)$ around in a counterclockwise fashion to get a compactly supported class f_n in $\text{PMCG}_c(\Sigma)$. The sequence $\{f_n\}$ converges to the boundary leaf shift.




Figure 2. The boundary leaf shift.

Let γ be a separating curve in Σ whose complementary components are both noncompact. Let Σ_- and Σ_+ be the closures of the two components of $\Sigma - \gamma$. By the same argument as in Lemma 2.5, Σ deformation retracts to a subspace homeomorphic to $X \vee \gamma \vee Y$, where X and Y are subspaces of Σ_- and Σ_+ , respectively. It follows that $H_1(\Sigma)$ splits as $A \oplus \langle \gamma \rangle \oplus B$, where $A = H_1(X)$ and $B = H_1(Y)$.

Similarly, if h is a handle shift dual to γ , then

$$H_1(\Sigma) \cong L \oplus \langle \gamma \rangle \oplus H_1(\text{supp}(h)) \oplus R,$$

where L and R are subgroups of A and B .

Definition 3.4 (pseudo handle shift). We say that a mapping class $\mathcal{H}$ is a *pseudo handle shift dual to a separating curve γ with associated handle shift h* if the following hold:

- (1) h is a handle shift dual to γ ;
- (2) $\mathcal{H}_*$ agrees with h_* on $H_1(\text{supp}(h))$;
- (3) $\mathcal{H}_*([\gamma]) = [\gamma]$;
- (4) $\mathcal{H}_*(L) < A$;
- (5) $\mathcal{H}_*(R) < B$.

In what follows, we always assume that the repelling end of h is on the “ A -side.”

Examples 3.5. Let h be a handle shift dual to a separating curve γ . Then h is itself a pseudo handle shift dual to γ (with associated handle shift h).

For a less trivial example, take a connected component X of $\Sigma \setminus (\gamma \cup \text{supp}(h))$. Then the composition h with any element $g \in \text{MCG}(\Sigma)$ supported on X is a pseudo handle shift associated to h . In particular, if $h' = \prod h_i$ is a product of commuting handle shifts h_i with dual curves γ_i , such that h and h' commute, $\text{supp}(h')$ is disjoint from γ and $\text{supp}(h)$ is disjoint from all γ_i , then the composition $k = h \circ h'$ is a pseudo handle shift associated to h .

Note that since pseudo handle shifts are defined by their action on homology, if $f_* = \mathcal{H}_*$ for a pseudo handle shift $\mathcal{H}$ associated to h , then f is also a pseudo handle shift associate to h .

Theorem 3.6 (pseudo handle shifts are unapproachable). *A pseudo handle shift $\mathcal{H}$ dual to a separating curve γ is not a limit of compactly supported mapping classes.*

Proof. Let h be the associated handle shift dual to γ . Let ϵ_- and ϵ_+ be the ends of Σ corresponding to the repelling and attracting ends of h , respectively, and let Σ_- and Σ_+ be the complementary components of $\Sigma - \gamma$ containing ϵ_- and ϵ_+ , respectively. Choose some principal exhaustion $\{P_i\}$ of Σ , and let $\Sigma_-^i = (\Sigma - P_i) \cap \Sigma_-$ and $\Sigma_+^i = (\Sigma - P_i) \cap \Sigma_+$.

The curve γ partitions the space of ends into two closed subspaces E_- and E_+ of which Σ_- and Σ_+ are neighborhoods, respectively. The subsurfaces Σ_-^i and Σ_+^i are also neighborhoods of E_- and E_+ . Since $\mathcal{H}$ is pure, $\mathcal{H}(\Sigma_-^i)$ and $\mathcal{H}(\Sigma_+^i)$ are also neighborhoods of E_- and E_+ . Since E_- and E_+ are disjoint, the intersection of the closures of $\mathcal{H}(\Sigma_\pm^i)$ and $\Sigma_\mp$ is compact, and since the $\Sigma_\pm^i$ are nested and have empty intersection, we may take i large enough so that the term P_i in our principal exhaustion contains γ and satisfies $\mathcal{H}(\Sigma_\pm^i) \cap \Sigma_\mp$ is empty.

The handle shift h is supported on a strip $\mathcal{S}$ with equally spaced handles and standard basis $\{\alpha_p, \beta_p\}_{k \in \mathbb{Z}}$ of $H_1(\mathcal{S})$ so that $h_*(\alpha_p) = \alpha_{p+1}$ and $h_*(\beta_p) = \beta_{p+1}$. We choose once and for all curves in $\mathcal{S}$ representing these classes. After reindexing the α_p and β_p by translating p , we assume that α_1 and β_1 lie in Σ_-^i . Since α_p and β_p tend to ϵ_+ , there is some $j > 1$ such that α_j and β_j lie in Σ_+^i .

Suppose that $\mathcal{H}$ is a limit of compactly supported $\mathcal{H}_n$. Pick n large enough so that $\mathcal{H}_n$ agrees with $\mathcal{H}$ on P_i and so that $\mathcal{H}_{n*}$ agrees with h_* on both $H_1(P_i)$ and $\langle \alpha_1, \beta_1, \dots, \alpha_j, \beta_j \rangle$. Let P_k be some term in the exhaustion with $k \geq i$ that contains the support of $\mathcal{H}_n$.

We have a direct sum decomposition

$$H_1(P_k) \cong \mathbb{Z}^\ell \oplus \mathbb{Z}^{2j} \oplus \mathbb{Z}^r$$

where $\mathbb{Z}^\ell$ is a subgroup of $H_1(\Sigma_-) \oplus \langle \gamma \rangle$, $\mathbb{Z}^{2j} = \langle \alpha_1, \beta_1, \dots, \alpha_j, \beta_j \rangle$, and $\mathbb{Z}^r$ is a subgroup of $H_1(\Sigma_+)$. Picking a basis $\langle x_1, \dots, x_\ell, \alpha_1, \beta_1, \dots, \alpha_j, \beta_j, y_1, \dots, y_r \rangle$ for $H_1(P_k)$ compatible with this decomposition, we see that $\mathcal{H}_{n*}$ has a block decomposition:

$$\mathcal{H}_{n*} = \begin{matrix} & \ell & 2j-2 & 2 & r \\ \begin{matrix} \ell \\ 2 \\ 2j-2 \\ r \end{matrix} & \begin{pmatrix} * & \mathbf{0} & \mathbf{0} & Y \\ * & \mathbf{0} & \mathbf{0} & Z \\ * & I & \mathbf{0} & * \\ X & \mathbf{0} & A & B \end{pmatrix} \end{matrix}$$

By properties (4) and (5) of a pseudo handle shift, and our choice of i , the blocks X , Y , and Z are all zero. So the matrix is

$$\mathcal{H}_{n*} = \begin{matrix} & \ell & 2j-2 & 2 & r \\ \begin{matrix} \ell \\ 2 \\ 2j-2 \\ r \end{matrix} & \begin{pmatrix} * & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ * & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ * & I & \mathbf{0} & * \\ \mathbf{0} & \mathbf{0} & A & B \end{pmatrix} \end{matrix}$$

This matrix is column equivalent to

$$\mathcal{H}_{n*} = \begin{matrix} & \ell & 2j-2 & 2 & r \\ \begin{matrix} \ell \\ 2 \\ 2j-2 \\ r \end{matrix} & \begin{pmatrix} * & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ * & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ * & I & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & A & B \end{pmatrix} \end{matrix}$$

But the matrix $[A \ B]$ is an $r \times (r + 2)$ matrix, and so its Jordan form cannot have a pivot in every column. So the matrix for $\mathcal{H}_{n*}$ is equivalent to a matrix with a zero column. But $\mathcal{H}_{n*}$ is an isomorphism, and this contradiction completes the proof. $\square$

We are now ready to prove Theorem 3.1.

Proof of Theorem 3.1. We will first show that $\mathcal{J}(\Sigma) < \overline{\text{PMCG}_c(\Sigma)}$. By [23, Theorem 1], we only need to consider the case when Σ has at least two ends accumulated by genus. We observe that $\mathcal{J}(\Sigma) < \text{PMCG}(\Sigma)$. Let g be in $\text{PMCG}(\Sigma)$ so that g is not a limit of compactly supported mapping classes. We show that g is not in $\mathcal{J}(\Sigma)$.

By Theorem 3 and Corollary 4 from [1], g can be written $g = fk^{-1}$ where f is a limit of compactly supported classes and k is a product of pairwise commuting handle shifts h_i . The handle shift h_i has the property that the support of h_i is disjoint from the dual curve γ_j for h_j whenever $i \neq j$. Thus, for any i , k is a pseudo-handle shift dual to γ_i associated to h_i (see Examples after the definition of pseudo handle shifts). If g were in Torelli, then $f_* = g_*k_* = k_*$, which implies f is a pseudo handle shift, but this violates Theorem 3.6. This shows

$$\mathcal{J}(\Sigma) < \overline{\text{PMCG}_c(\Sigma)}.$$

If ϕ_n is a sequence in $\mathcal{J}_c(\Sigma)$ that converges to ϕ , then ϕ lies in $\mathcal{J}(\Sigma)$, since $\phi_n(\alpha)$ eventually agrees with $\phi(\alpha)$ for any given simple closed curve α . So

$$\overline{\mathcal{J}_c(\Sigma)} < \mathcal{J}(\Sigma).$$

For the other containment, let ϕ be an element of $\mathcal{J}(\Sigma)$ and let $\{\psi_n\}$ be a sequence in $\text{PMCG}_c(\Sigma)$ converging to ϕ . We would like to convert ψ_n into a sequence of compactly supported ϕ_n in $\mathcal{J}(\Sigma)$ converging to ϕ . The idea is that the homology classes affected by ψ_n must move further and further away from a given basepoint, and so we can precompose the ψ_n with a mapping class supported far from the basepoint to produce the desired ϕ_n .

Fix a principal exhaustion $\{P_i\}$ of Σ . For each i , pick a $j > i$ such that P_j contains $\phi^{-1}(P_i)$. Pick an N large enough so that, for all $n \geq N$, the map ψ_n has a representative that agrees with a fixed representative of ϕ on P_j . Note that ψ_{n*} agrees with ϕ_* on $H_1(P_j)$. Pick a $k > j$ such that P_k contains the support of ψ_n .

By Lemma 2.5, we have $H_1(P_k) \cong H_1(P_i) \oplus Q \oplus R$ for some Q a subgroup of $H_1(P_j - P_i)$ and R a subgroup of $H_1(P_k - P_j)$. Let α be element of $H_1(P_k)$ and write $\alpha = \gamma + \mu + \nu$ where γ , μ , and ν are in $H_1(P_i)$, Q , and R , respectively. So $\psi_{n*}(\alpha) = \gamma + \mu + \psi_{n*}(\nu)$.

The class ν is represented by a 1-manifold $\mathcal{N}$ in $P_k - P_j$. By our choice of j and n , the 1-manifold $\psi_n(\mathcal{N})$ is disjoint from P_i . So $\psi_{n*}(\nu)$ is in $Q \oplus R$. Therefore $\psi_{n*}: H_1(P_k) \rightarrow H_1(P_k)$ may be represented by a square matrix

$$A = \begin{bmatrix} I & \mathbf{0} \\ \mathbf{0} & B \end{bmatrix},$$

where I is the identity on $H_1(P_i)$ and B is a square matrix. Since A is the induced map on homology associated to a homeomorphism of P_k , it is invertible and respects the intersection form, and so the same is true of B .

We claim that the matrix B is represented by a homeomorphism $F': \overline{P_k - P_i} \rightarrow \overline{P_k - P_i}$ that is the identity on $\partial P_i \cap \overline{P_k - P_i}$. To see this, note that B preserves the intersection form and the homology classes of the boundary components of each component of $\overline{P_k - P_i}$. Let X be the surface obtained from $\overline{P_k - P_i}$ by capping off all the boundary components with disks. The homology of X is a quotient of that of $\overline{P_k - P_i}$, and B induces an automorphism of the homology of X that preserves the intersection form. There is a homeomorphism $X \rightarrow X$ inducing this automorphism that preserves each component of X , by Burkhardt's theorem [8]. By an isotopy, we may assume that Burkhardt's homeomorphism of X fixes, point-wise, small disks around the centers of the disks we added to construct X . Restricting this to $\overline{P_k - P_i}$ is the desired F' .

We extend F' by the identity to all of Σ and call the result F .

Now consider the homeomorphism $\phi_n = \psi_n \circ F^{-1}$. By the construction of F , this homeomorphism acts trivially on the homology of Σ , and agrees with ψ_n on P_i .

This completes the proof. $\square$

We finish this section with the alternate proof of $\mathcal{I}(\Sigma) < \overline{\text{PMCG}_c(\Sigma)}$, as suggested by the referee. Suppose, for contradiction, that there is $f \in \mathcal{I}(\Sigma)$ which may not be expressed as the limit of a sequence of mapping classes with compact support. A consequence of [1, Theorem 4.5] is that there is a separating curve $\gamma \subset \Sigma$ such that γ and $f(\gamma)$ have different topological types in every compact subsurface of Σ containing them. In particular, they induce different splittings of $H_1(\Sigma, \mathbb{Z})$, which contradicts the fact that f acts trivially on homology.

3.1. Generation by separating twists and bounding pairs. The proofs in this section are due to Justin Malestein.

Proposition 3.7. *For any finite-type surface $\Sigma = \Sigma_{g,p}^b$, $\mathcal{I}(\Sigma)$ is generated by separating twists and bounding pair maps.*

Proof. The Dehn twist about each boundary component of Σ is in $\mathcal{J}(\Sigma)$ and counts as a separating twist. By capping off each boundary component of Σ with a once-punctured disk, we get an exact sequence

$$1 \longrightarrow \prod_{\alpha \in \partial \Sigma} \langle T_\alpha \rangle \longrightarrow \mathcal{J}(\Sigma_{g,p}^b) \longrightarrow \mathcal{J}(\Sigma_{g,p+b}) \longrightarrow 1.$$

Since a separating twist (resp. bounding pair map) from $\mathcal{J}(\Sigma_{g,p+b})$ can be lifted to a separating twist (resp. bounding pair map) in $\mathcal{J}(\Sigma_{g,p}^b)$, we may assume $b = 0$.

Recall that $\mathcal{J}(\Sigma) \leq \text{PMCG}(\Sigma)$ and $\text{PMCG}(\Sigma)$ is generated by Dehn twists. When $g = 0$, then all curves are separating, so $\text{PMCG}(\Sigma) = \mathcal{J}(\Sigma)$. When $g = 1$ and $p \leq 1$, $\mathcal{J}(\Sigma)$ is the trivial group. When $g \geq 2$ and $p \leq 1$, then the statement of the proposition holds by Birman [4] and Powell [24]. We now assume $g \geq 1$ and induct on p using the Birman exact sequence.

Set $\Sigma = \Sigma_{g,p}$, $g \geq 1$ and $p \geq 1$. Pick a point $x \in \Sigma$ and let $\Sigma - x$ denote the surface obtained from puncturing Σ at x . The inclusion map $\Sigma - x \hookrightarrow \Sigma$ gives rise to the Birman exact sequence:

$$1 \longrightarrow \pi_1(\Sigma, x) \xrightarrow{\rho} \text{PMCG}(\Sigma - x) \longrightarrow \text{PMCG}(\Sigma) \longrightarrow 1,$$

where $\pi_1(\Sigma, x) \xrightarrow{\rho} \text{PMCG}(\Sigma - x)$ is the point-pushing map. Let

$$\Delta = \rho^{-1}(\mathcal{J}(\Sigma - x)) \cap \pi_1(\Sigma, x).$$

The Birman sequence induces the exact sequence

$$1 \longrightarrow \Delta \longrightarrow \mathcal{J}(\Sigma - x) \longrightarrow \mathcal{J}(\Sigma) \longrightarrow 1.$$

By the induction hypothesis, $\mathcal{J}(\Sigma)$ is generated by separating twists and bounding pair maps. Every bounding pair (resp. separating curve) in Σ has a lift to $\Sigma - x$ which is also a bounding pair (resp. separating curve). Thus, separating twists and bounding pair maps in $\mathcal{J}(\Sigma)$ can be lifted to such maps in $\mathcal{J}(\Sigma - x)$. We now claim $\rho(\Delta)$ is contained in the group generated by separating twists.

Given a simple curve $\alpha \in \pi_1(\Sigma, x)$, the point-pushing map ρ_α along α is equal to a product of two Dehn twists; namely $\rho_\alpha = T_{\alpha'} T_{\alpha''}^{-1}$, where α' and α'' are freely homotopic to α in Σ but not in $\Sigma - x$. Let $\Delta_0 = [\pi_1(\Sigma, x), \pi_1(\Sigma, x)]$. By [25, Lemma A.1], $\rho(\Delta_0)$ is generated by point-pushing maps along simple (null-homologous) separating curves. In particular, if α is separating, then ρ_α is a product of two separating twists. This shows $\rho(\Delta_0)$ is contained in the group generated by separating twists. Now let $\{x_1, \dots, x_p\}$ be the set of punctures of Σ . For each x_i , let $\alpha_i \in \pi_1(\Sigma, x)$ be a simple curve which is freely homotopic to x_i . In this case, ρ_{α_i} is a single separating twist along a curve which is freely homotopic to x_i in Σ but not in $\Sigma - x$.

We now claim $\Delta = \langle \Delta_0, \alpha_1, \dots, \alpha_p \rangle$. Let $\gamma \in \pi_1(\Sigma, x) - \langle \Delta_0, \alpha_1, \dots, \alpha_p \rangle$. Then $[\gamma] \in H_1(\Sigma)/\langle \alpha_1, \dots, \alpha_p \rangle$ is nontrivial, so there is a simple non-separating curve α on Σ such that $[\gamma] = [\alpha^k]$ in $H_1(\Sigma)/\langle \alpha_1, \dots, \alpha_p \rangle$. But then ρ_α is a product of Dehn twists along two disjoint non-separating, non-homologous curves, so cannot be in $\mathcal{J}(\Sigma - x)$. This shows $\Delta = \langle \Delta_0, \alpha_1, \dots, \alpha_p \rangle$, and in particular $\rho(\Delta)$ is contained in the group generated by separating twists. This finishes the proof that $\mathcal{J}(\Sigma - x)$ is generated by separating twists and bounding pair maps. $\square$

Lemma 3.8. *Let Σ be a connected oriented surface of infinite type. If P is a principal domain of Σ such that each boundary component of P is separating, then the inclusion map $P \hookrightarrow \Sigma$ induces an embedding $i_*: \text{MCG}(P) \rightarrow \text{MCG}(\Sigma)$ and $i_*^{-1}(\mathcal{J}(\Sigma)) \cap \text{MCG}(P) = \mathcal{J}(P)$.*

Proof. By our assumption, all boundary curves of P are essential and distinct in Σ , hence Dehn twists along the boundary curves induce non-trivial mapping classes in $\text{MCG}(\Sigma)$. This shows i_* is injective. For the second statement, by Lemma 2.5, we have a direct sum decomposition $H_1(\Sigma) = H_1(P) \oplus M$ for some $M < H_1(\Sigma \setminus P)$, thus $i_*^{-1}(\mathcal{J}(\Sigma)) \cap \text{MCG}(P) = \mathcal{J}(P)$. $\square$

We now prove the second main result of the introduction; namely:

Theorem 3.9. *Let Σ be a connected oriented surface of infinite type. Then $\mathcal{J}(\Sigma)$ is topologically generated by separating twists and bounding-pair maps.*

Proof. Let $\{P_1, P_2, \dots\}$ be a principal exhaustion of Σ . By the previous lemma, elements in $\mathcal{J}(\Sigma)$ with support on P_i is $\mathcal{J}(P_i)$. By Theorem 3.1, $\mathcal{J}(\Sigma) = \overline{\bigcup_i \mathcal{J}(P_i)}$. Finally, by Proposition 3.7, $\mathcal{J}(P_i)$ is generated by separating twists and bounding pair maps. $\square$

4. Abstract commensurators of the Torelli group

In this section we prove Theorem 1.4. As in the statement of the theorem, throughout this section we will assume that the surface Σ has empty boundary. As mentioned in the introduction, the first step of the argument consists of proving that an element of $\text{Comm}\mathcal{J}(\Sigma)$ induces a simplicial automorphism of a combinatorial object associated to Σ , called the *Torelli complex*, introduced by Brendle and Margalit in [6].

4.1. Torelli complex. Recall that the *curve complex* of Σ is the (infinite-dimensional) simplicial complex whose vertex set is the set of isotopy classes of curves in Σ , and where a collection of vertices spans a simplex if and only if the corresponding curves are pairwise disjoint. The curve complex was used by Ivanov [14],

Korkmaz [18], and Luo [19] to prove that, for all but a few finite-type surfaces Σ ,

$$\text{Comm MCG}(\Sigma) \cong \text{Aut MCG}(\Sigma) \cong \text{MCG}(\Sigma).$$

Subsequently, Bavard, Dowdall, and Rafi [2] established the analogous result for *every* infinite-type surface. In a similar fashion, Farb and Ivanov [9], Brendle and Margalit [6, 7, 5], and Kida [16] proved that, for all but a few finite-type surfaces,

$$\text{Comm } \mathcal{J}(\Sigma) \cong \text{Aut } \mathcal{J}(\Sigma) \cong \text{MCG}(\Sigma).$$

Here, we will adapt the ideas of Brendle and Margalit [6] to the infinite-type setting. Given an infinite-type surface Σ , we define its Torelli complex to be the (infinite-dimensional) simplicial complex whose vertex set is the set of isotopy classes of separating curves and bounding pairs in Σ , and where a collection of vertices spans a simplex if and only if the corresponding curves are pairwise disjoint. In order to relax notation, we will blur the distinction between vertices of $\mathcal{J}(\Sigma)$ and the curves (or multicurves) they represent.

The Torelli complex of a finite-type surface is connected [9]. As a consequence, the same holds for the Torelli complex of an infinite-type surface also. We record the following observation as a separate lemma, as we will need to make use of it later:

Lemma 4.1. *The Torelli complex $\mathcal{J}(\Sigma)$ has infinite diameter if and only if Σ has finite type.*

Proof. If Σ has finite type, a slick limiting argument due to Feng Luo (see the comment after Proposition 4.6 of [20]) shows that the curve complex has infinite diameter. The obvious adaptation of this method to the case of the Torelli complex also implies that $\mathcal{J}(\Sigma)$ has infinite diameter.

For the other direction, suppose Σ has infinite type. Since curves are compact, given multicurves $\gamma, \delta \subset \Sigma$, we can find a separating curve $\eta \subset \Sigma$ which is disjoint from both γ and δ . In particular, $\mathcal{J}(\Sigma)$ has diameter two. $\square$

4.2. Automorphisms of the Torelli complex. Denote by $\text{Aut } \mathcal{J}(\Sigma)$ the group of simplicial automorphisms of $\mathcal{J}(\Sigma)$, and observe that there is a natural homomorphism $\text{MCG}(\Sigma) \rightarrow \text{Aut } \mathcal{J}(\Sigma)$. We want to prove:

Theorem 4.2. *Let Σ be an infinite-type surface without boundary. The natural homomorphism $\text{MCG}(\Sigma) \rightarrow \text{Aut } \mathcal{J}(\Sigma)$ is an isomorphism.*

As noted above, the finite-type case is due to Brendle and Margalit [6, 7, 5] and Kida [16]. Indeed, the notion of *sides* which is used in this section is adapted from arguments that may be found in Brendle and Margalit [6], and which find their way back to ideas of Ivanov [14].

Sides. Recall that the *link* of a vertex v of a simplicial complex X is the set of all vertices of X that span an edge with v . In particular, v is not an element of its link. For any finite-dimensional simplex σ let $\text{Link}(\sigma)$ be the intersection of the links of each of the vertices in σ . We say that two vertices α, β in $\text{Link}(\sigma)$ lie on the same *side* of σ if there exists a vertex γ in $\text{Link}(\sigma)$ that fails to span an edge with both α and β , that is, if there exists a curve in $\text{Link}(\sigma)$ that intersects both α and β . Observe that “being on the same side” defines an equivalence relation $\sim_\sigma$ on $\text{Link}(\sigma)$, that is, the *sides* of σ are the equivalence classes of $\sim_\sigma$ in $\text{Link}(\sigma)$.

In particular, we may consider the sides of a vertex of $\mathcal{T}(\Sigma)$. We say that γ in $\mathcal{T}(\Sigma)$ is k -sided if there are k equivalence classes with respect to $\sim_\gamma$. As we shall see, k is in $\{1, 2\}$.

For any vertex γ of $\mathcal{T}(\Sigma)$ there exist two subsurfaces $R, L \subset \Sigma$ obtained by cutting Σ along γ such that γ is isotopic to the boundary components of both R and L . Suppose R is of finite type. We call γ a *pants curve* if γ is a separating curve and $R \cong \Sigma_{0,2}^1$, a sphere with two punctures and one boundary component. We call γ a *genus curve* if γ is a separating curve and $R \cong \Sigma_{1,0}^1$, a torus with one boundary component. If γ is any other type of separating curve then we say it is type X .

If γ is a bounding pair and one of the associated subsurfaces of Σ is homeomorphic to $\Sigma_{1,0}^2$ then we call it a *genus bounding pair*.

Lemma 4.3. *A vertex γ in $\mathcal{T}(\Sigma)$ is 2-sided if and only if it is type X or it is a genus bounding pair. Otherwise, γ is 1-sided.*

Proof. We first prove that if γ has type X then it has exactly two sides. Let R and L be the two subsurfaces of Σ obtained by cutting along γ . Let $\alpha, \beta \in \text{Link}(\gamma)$. If $\alpha \subset R$ and $\beta \subset L$, then any vertex of $\mathcal{T}(\Sigma)$ that intersects both α and β must also intersect γ . This implies that γ has at least two sides. If $\alpha, \beta \subset R$ then there exists an element of the $\text{MCG}(\Sigma)$ -orbit of α that intersects both α and β and is contained in R . An identical argument holds for two vertices contained in L and so it follows that γ has exactly two sides.

Now let γ be a genus one separating curve or a pants curve. Define $L, R \subset \Sigma$ as above. Observe that neither $\Sigma_{0,2}^1$ nor $\Sigma_{1,0}^1$ contains any non-peripheral separating curves or bounding pairs. Therefore $\text{Link}(\gamma)$ does not contain any curves in R . As above, all vertices contained in L are on the same side and so γ is 1-sided.

We now move on to the case where γ is a bounding pair. We define R and L as above. Assume that neither R nor L is homeomorphic to $\Sigma_{1,0}^2$. Let $\alpha, \beta \in \text{Link}(\gamma)$ be such that $\alpha \subset R$ and $\beta \subset L$. As shown in Figure 3, there exists a bounding pair $\gamma' = \{\delta_R, \delta_L\}$ such that

- any pair of curves in γ or γ' forms a bounding pair,
- $\delta_R \subset R$ and $\delta_L \subset L$, and
- $\delta_R \cap \alpha \neq \emptyset$ and $\delta_L \cap \beta \neq \emptyset$.




Figure 3. A general bounding pair γ is 1-sided. For any two vertices α, β in $\mathcal{T}(\Sigma)$ adjacent to γ , we can find a bounding pair not adjacent to α and β but adjacent to γ . Informally, bounding pairs can “pass through” each other.

That is, γ' is in $\text{Link}(\gamma)$ and there is no edge between γ' and α or between γ' and β . It follows that γ has exactly one side.

If γ is a genus bounding pair then no such γ' exists. Indeed, every non-separating curve in R that forms a bounding pair with a curve in γ is also isotopic to a curve in γ . By the same argument as for type X vertices, we conclude that γ is 2-sided. $\square$

Let σ be a finite-dimensional simplex of $\mathcal{T}(\Sigma)$ consisting entirely of curves of type X . Using similar methods to the above proofs it is straightforward to show that the set of sides of σ is in bijective correspondence with the subsurfaces of Σ obtained by cutting Σ along σ .

The following lemma follows immediately from Lemma 2.2.

Lemma 4.4. *Let σ be a simplex in $\mathcal{T}(\Sigma)$. Then there are curves realizing the homotopy classes in σ which do not accumulate in any compact set of Σ if and only if for every vertex $v \notin \sigma$, the set $\{w \in \sigma \mid i(v, w) \neq 0\}$ is finite.*

Proof of Theorem 4.2. Let

$$\Phi: \text{MCG}(\Sigma) \longrightarrow \text{Aut}(\mathcal{T}(\Sigma))$$

be the natural homomorphism; that is, for f in $\text{MCG}(\Sigma)$, $\Phi(f)$ is the automorphism of $\mathcal{T}(\Sigma)$ determined by the rule

$$\Phi(f)(\gamma) = f(\gamma)$$

for every separating curve or bounding pair γ .

First, we show that Φ is injective. To this end, suppose $\Phi(f) = \text{Id}$. Then we argue that $f(\gamma) = \gamma$ for every curve γ . Indeed, if γ is separating, then γ is a vertex of $\mathcal{T}(\Sigma)$, so $\Phi(f)(\gamma) = \gamma$ and we are done. If γ is non-separating, there is some curve γ' such that γ and γ' form a bounding pair. Because $\Phi(f)$ fixes the vertex corresponding to $\gamma \cup \gamma'$, it must be the case that either $f(\gamma) = \gamma$ and $f(\gamma') = \gamma'$ or $f(\gamma) = \gamma'$ and $f(\gamma') = \gamma$. But there exists a separating curve η that intersects γ but not γ' . Because $f(\eta) = \eta$, it cannot be the case that $f(\gamma) = \gamma'$. Therefore $f(\gamma) = \gamma$ as desired.

By the Alexander method for infinite-type surfaces, due to Hernández, Moralez, and Valdez [13], we deduce that f is the identity in $\text{MCG}(\Sigma)$.

We now show that Φ is surjective. Let $\phi: \mathcal{T}(\Sigma) \rightarrow \mathcal{T}(\Sigma)$ be an automorphism. Fix a principal exhaustion $\{P_1, P_2, \dots\}$ of Σ such that P_1 has complexity at least six. Define σ_i to be the simplex of $\mathcal{T}(\Sigma)$ corresponding to the multicurve ∂P_i . Denote by $\mathcal{P}_i$ the subcomplex of $\mathcal{T}(\Sigma)$ spanned by the curves and bounding pairs contained in P_i . Denote by P_i° the surface obtained by gluing once-punctured disks to each boundary component of P_i . By construction, σ_i contains only type X vertices and therefore $\mathcal{T}(\mathcal{P}_i)$ is isomorphic to $\mathcal{T}(P_i^\circ)$. As the complexity of P_i is at least six, $\mathcal{P}_i$ is connected for all i . By Lemma 4.1, we know that $\mathcal{P}_i$ is the unique side of σ_i whose diameter is infinite.

Since ϕ is a simplicial automorphism, it induces a bijection between the sides of σ_i and the sides of $\phi(\sigma_i)$. Because all simplicial automorphisms of $\mathcal{T}(\Sigma)$ are isometries, $\phi(\sigma_i)$ has a unique side of infinite diameter. From Lemma 4.3 we have that every vertex of $\phi(\sigma_i)$ is of type X or it is a genus bounding pair. However, since each side of $\phi(\sigma_i)$ is connected, it cannot contain a genus bounding pair by Lemma 4.3.

We write $\mathcal{Q}_i \subset \text{Link}(\phi(\sigma_i))$ for the unique side of $\phi(\sigma_i)$ with infinite diameter, and $Q_i \subset \Sigma$ for the finite-type subsurface which it defines. By Lemma 4.4, we can realize the curves $\bigcup \mathcal{Q}_i$ by non-accumulating curves in Σ . The $\mathcal{Q}_i$'s form a sequence of nested subsurfaces, thus their union is an open set. The nonaccumulation property implies that $\bigcup \mathcal{Q}_i$ is the full surface Σ .

Since each vertex of $\phi(\sigma_i)$ has type X we have that $\mathcal{Q}_i \cong \mathcal{T}(Q_i^\circ)$. Furthermore, ϕ restricts to an isomorphism

$$\phi_i: \mathcal{P}_i \longrightarrow \mathcal{Q}_i.$$

Since each P_i is assumed to have complexity at least six, the combination of results of Kida [16] and Korkmaz [18] implies that ϕ_i is induced by a homeomorphism $f_i: P_i \rightarrow Q_i$. Moreover, the homeomorphism f_{i+1} may be chosen so that it restricts to f_i on the subsurface P_i . Hence their direct limit is a homeomorphism of $\bigcup P_i$ to $\bigcup Q_i$ inducing ϕ . Since $\Sigma = \bigcup P_i = \bigcup Q_i$, this completes the proof. $\square$

We now end this subsection with the following two observations which will be useful later.

Lemma 4.5. *For any simplex σ in $\mathcal{T}(\Sigma)$ and any compactly-supported $f \in \mathcal{I}(\Sigma)$, if f preserves σ then f fixes σ pointwise.*

Proof. Consider two vertices $v, w \in \sigma$, and assume $f(v) = w$. Then v and w are either both separating curves or they are both bounding pairs.

Suppose v separates Σ into two infinite-type subsurfaces. Since v and w are disjoint, we can find a compact domain $Y \subset \Sigma$ such that v and w are of different topological type. By choosing Y large enough, we can assume that Y supports f . But then it is impossible for $f|_Y(v) = f(v) = w$.

Suppose now that v (and w) separates Σ into a finite-type subsurface and an infinite-type subsurface. Call the finite-type subsurfaces V and W , corresponding to v and w respectively. Since $v \in \Sigma \setminus W$ and $w \in \Sigma \setminus V$, the subsurfaces V and W are disjoint. By assumption $f(v) = w$, therefore V and W must have the same topological type and $f(V) \subset W$. But this is impossible if $f \in \mathcal{I}(\Sigma)$. $\square$

Lemma 4.6. *For any bounding pair $v \in \mathcal{T}(\Sigma)$ and any $f \in \mathcal{I}(\Sigma)$, if f preserves v , then f must fix the curves in v .*

Proof. Suppose $v = \{\alpha, \beta\}$ and $f(\alpha) = \beta$. The complement of v in Σ has two components. Since f is pure it preserves the components. Let Y be one of the components. The orientation of Y induces an orientation on α and β . These oriented curves (as homology classes) satisfy $\alpha = -\beta$. The map f is orientation preserving, so it must preserve the orientation of α and β . That is, $f(\alpha) = \beta$ as oriented curves. But then α and $f(\alpha)$ cannot be homologous, contradicting that $f \in \mathcal{I}(\Sigma)$. $\square$

4.3. Algebraic characterization of twists and bounding pair maps. Before proving Theorem 1.4 we will need one more ingredient. Notice that the vertices of $\mathcal{T}(\Sigma)$ define supports of elements in $\mathcal{I}(\Sigma)$. We must now show that commensurations of $\mathcal{I}(\Sigma)$ preserve such elements and therefore define a permutation of the vertices of the complex. We will adapt the algebraic characterization of Dehn twists of Bavard, Dowdall, and Rafi [2] to our setting.

We first introduce some terminology to facilitate the characterization of twists and bounding pairs. Let $G < \text{MCG}(\Sigma)$. We denote by $\mathcal{F}_G$ the set of elements of G whose conjugacy class (in G) is countable. Bavard, Dowdall, and Rafi proved that if G is finite-index in $\text{MCG}(\Sigma)$ then f is in $\mathcal{F}_G$ if and only if it has compact support [2, Proposition 4.2]. Using similar methods, we will show:

Proposition 4.7. *Let $G < \mathcal{I}(\Sigma)$ be a finite-index subgroup. An element f in G has compact support if and only if f is in $\mathcal{F}_G$.*

Proof. It is clear that compactly-supported mapping classes have countable conjugacy classes. For the opposite direction, the argument in [2, Proposition 4.2] exhibits a infinite sequence of pairwise-disjoint curves a_i such that the Dehn twists about the a_i give rise to uncountably many conjugates of f . Since Σ has infinite type, the curves a_i may be chosen to be separating, so that the corresponding twists belong to $\mathcal{I}(\Sigma)$. Hence the result follows. $\square$

We denote by $Z(H)$ the center of H . If h is in H , we write $C_H(h)$ for the centralizer of h in H . Given a finite-index subgroup $G < \mathcal{I}(\Sigma)$ we write $\mathcal{M}_G$ for the set of elements f in G which satisfy the following three conditions:

- (1) $f \in \mathcal{F}_G$,
- (2) $Z(\mathcal{F}_G \cap C_G(f))$ is an infinite cyclic group, and
- (3) $C_G(f) = C_G(f^k)$ for every $k > 0$.

We now prove that, for any finite-index subgroup G of $\mathcal{I}(\Sigma)$, powers of Dehn twists and bounding pair maps belong to the set $\mathcal{M}_G$.

Lemma 4.8. *Let $G < \mathcal{I}(\Sigma)$ be a finite-index subgroup. If $f \in G$ is a power of a Dehn twist about a separating curve or a bounding pair map then f belongs to $\mathcal{M}_G$.*

Proof. Since f has compact support, $f \in \mathcal{F}_G$. Suppose first f is a power of a Dehn twist about the separating curve γ . We have that

$$C_{\mathcal{I}(\Sigma)}(T_\gamma^k) = \{g \in \mathcal{I}(\Sigma) \mid g(\gamma) = \gamma\},$$

for $k \neq 0$. It follows that all powers of T_γ have the same centralizer in $\mathcal{I}(\Sigma)$ and hence, in any subgroup. A similar argument holds if f is a power of bounding pair map. This implies the third condition in the definition of $\mathcal{M}_G$.

To see that f satisfies the second condition, once again assume first that f is a power of the Dehn twist about a separating curve γ . Let g be a nontrivial element of $\mathcal{F}_G \cap C_G(f)$ and assume that g is not a power of T_γ . Then there exists a curve δ disjoint from γ such that $g(\delta) \neq \delta$. If δ is a separating curve then T_δ^k is in $C_G(f)$, for some $k > 0$, but $gT_\delta^k \neq T_\delta^k g$, so g is not in $Z(\mathcal{F}_G \cap C_G(f))$. On the other hand, suppose δ is a non-separating curve; since g acts trivially on homology we have that δ and $g(\delta)$ are homologous curves. Because $T_\gamma g(\delta) = gT_\gamma(\delta) = g(\delta)$, we have that $g(\delta)$ is disjoint from γ . It follows that some power of $T_\delta T_{g(\delta)}^{-1}$ belongs to $C_G(f)$, but does not commute with g . We have therefore shown that g is not central in $\mathcal{F}_G \cap C_G(f)$ as desired.

A similar argument also shows that, if f is a power of a bounding pair map, then g belongs to $Z(\mathcal{F}_G \cap C_G(f))$ if and only if it is a power of the same bounding pair map. $\square$

When G is a finite-index subgroup of $\text{MCG}(\Sigma)$, all elements of $\mathcal{M}_G$ are powers of multi Dehn twists, see [2, Lemma 4.5]. In stark contrast, this is no longer true in our setting. When G is a finite-index subgroup of $\mathcal{I}(\Sigma)$, then $\mathcal{M}_G$ may contain elements which are not supported on a disjoint union of annuli: for example, we may take a pure braid on a nonseparating planar subsurface with at least three boundary components. The following proposition characterizes those elements that lie in $\mathcal{M}_G$, when G is a finite-index subgroup of $\mathcal{I}(\Sigma)$. Recall the definition of ∂f for a mapping class f , which is non-empty when f is compactly-supported. We have the following statements.

Lemma 4.9. *Let $G < \mathcal{I}(\Sigma)$ be a finite-index subgroup. Given a nontrivial $f \in \mathcal{M}_G$, let $\sigma \subset \partial f$ be the set of all separating curves or bounding pairs. The following statements hold.*

- (1) *For all $g \in \mathcal{F}_G \cap C_G(f)$, g fixes every component of $\Sigma \setminus \partial f$.*
- (2) *If σ is empty, then for each finite-type domain Y of $\Sigma \setminus \partial f$, $f|_Y$ is either the identity map or pseudo-Anosov. Furthermore, there is at least one pseudo-Anosov component of f .*
- (3) *If σ is non-empty then $\sigma = \partial f$ is either a separating or a bounding pair, and $f = T_\sigma^j$ for some non-zero j , where T_σ is either a separating twist or a bounding pair map.*

Proof. For any $g \in \mathcal{F}_G \cap C_G(f)$, $g(\partial f) = \partial(gfg^{-1}) = \partial f$. To show that g fixes every component of $\Sigma \setminus \partial f$, it is enough to show g fixes each component of ∂f . To see this, suppose there exists a domain Y of $\Sigma \setminus \partial f$ such that $g(Y)$ is not Y . Since g fixes the components of ∂f , Y and $g(Y)$ share the same boundary. This means $\Sigma = Y \cup g(Y) \cup \partial Y$, so Y must have infinite type. But then g cannot be a pure mapping class, contradicting the fact that $g \in \mathcal{I}(\Sigma)$. In the following, we will consider the two cases that σ is empty or σ is not empty. For each case, we show g fixes the components of ∂f along with the statements of (2) and (3).

First suppose that σ is empty. In this case, every curve of ∂f is non-separating and no two form a bounding pair. In particular, no two curves of ∂f are homologous, so every element of $\mathcal{F}_G \cap C_G(f)$ must fix the components of ∂f . Now let Y be a component of $\Sigma \setminus \partial f$ of finite-type. The map $f|_Y$ is irreducible (otherwise the reducing curves would also be in ∂f), so it is either of finite order or pseudo-Anosov. We now show it is not possible for $f|_Y$ to be non-trivial and have finite order. Since ∂f does not contain any separating curves or bounding pairs, it has cardinality at least 3. This means $\chi(Y) < 0$, in which case $\text{MCG}(Y)$ is torsion-free ([10, Corollary 7.3]). Hence, if $f|_Y$ is not pseudo-Anosov, then $f|_Y$ is the identity. Finally, if there is no pseudo-Anosov component of the support of f , then there exists a power $k \geq 1$ such that $f^k|_Y$ is the identity for any component Y in $\Sigma \setminus \partial f$. But f has infinite order by Lemma 2.3, so f^k must be a product of powers of Dehn twists about curves in ∂f . This is impossible as $f \in \mathcal{I}(\Sigma)$, see [28].

Now suppose σ is non-empty. Regard σ as a simplex in $\mathcal{I}(\Sigma)$. Every element $g \in \mathcal{F}_G \cap C_G(f)$ preserves ∂f , and hence also σ . By Lemma 4.5, g fixes each vertex of σ . Then, by Lemma 4.6, we can further conclude that g also fixes each curve in σ . Let v be a vertex of σ , and let T_v be the twist about v , which belongs to $\mathcal{I}(\Sigma)$.

We now show $f = T_v^j$ for some non-zero j . In particular, $v = \sigma = \partial f$. Since every $g \in \mathcal{F}_G \cap C_G(f)$ preserves v , g commutes with T_v , showing $T_v^k \in Z(\mathcal{F}_G \cap C_G(f))$ for some $k \geq 1$. Since f also lies in $Z(\mathcal{F}_G \cap C_G(f))$ which is infinite cyclic by assumption, we must have $f^m = T_v^n$ for some non-zero integers

m and n . Let w a vertex of $\mathcal{T}(\Sigma)$ disjoint from v and choose $k \geq 1$ so that $T_w^k \in G$. Since v and w are disjoint, we have

$$T_w^k \in C_G(T_v^n) = C_G(f^m) = C_G(f).$$

This is only possible if $f(w) = w$. Thus, f fixes every separating curve and bounding pair disjoint from v . Applying Lemma 4.6, we obtain that f fixes every curve disjoint from v . This shows $f = T_v^j$ for some non-zero integer j , concluding the proof. $\square$

Definition 4.10 (essential support). For each element $f \in \mathcal{M}_G$, we define the *essential support* $\text{supp}(f)$ of f as follows. Let $\sigma \subset \partial f$ be the set of all separating curves or bounding pairs.

- If σ is nonempty, then $\text{supp}(f) = \sigma$.
- If σ is empty, then $\text{supp}(f)$ is the union over all domains Y in $\Sigma \setminus \partial(f)$ such that $f|_Y$ is a pseudo-Anosov. Note that by Lemma 4.9, $\text{supp}(f)$ is nonempty.

Let

$$C_{\mathcal{M}_G}(f) = \{g \in \mathcal{M}_G \mid fg = gf\}.$$

We also define a further subset:

$$(\mathcal{P}_G)_f = \{g \in C_{\mathcal{M}_G}(f) \mid g \text{ is supported in } \Sigma \setminus \text{supp}(f)\}.$$

The next lemma tells us that the elements supported in $\text{supp}(f)$ are precisely those that are central.

Lemma 4.11. *Let $G < \mathcal{I}(\Sigma)$ be a finite-index subgroup. For any element f in $\mathcal{M}_G$ we have that*

$$C_{\mathcal{M}_G}(f) = Z(C_{\mathcal{M}_G}(f)) \oplus (\mathcal{P}_G)_f.$$

Proof. Let $\sigma \subset \partial f$ be the set of all separating curves or bounding pairs. If σ is nonempty, then by Lemma 4.9, $\sigma = \partial f$ is a separating curve or a bounding pair, and $f = T_\sigma^j$, where T_σ is a separating twist or a bounding pair map. In this case, any element $g \in C_{\mathcal{M}_G}(f)$ fixes every curve of σ . Thus, if g has support in $\text{supp}(f)$, then g is itself a power of T_σ . This shows $g \in Z(C_{\mathcal{M}_G}(f))$.

Now suppose σ is empty. By Lemma 4.9, $\text{supp}(f)$ is non-empty. Let $g \in C_{\mathcal{M}_G}(f) \subset \mathcal{F} \cap C_G(f)$ be an element with support in $\text{supp}(f)$. We want to show g lies in the center of $C_{\mathcal{M}_G}(f)$. If $h \in C_{\mathcal{M}_G}(f)$ has support disjoint from $\text{supp}(f)$, then h and g clearly commute. Henceforth, we may assume this is not the case.

By Lemma 4.9, f , g , and h all preserve every component of $\text{supp}(f)$. Thus, we can assume without a loss of generality that h is also supported in $\text{supp}(f)$.

Let $Y_1, \dots, Y_n$ be the components of $\text{supp}(f)$. For each i , let f_i , (resp. g_i and h_i) denote the restriction of f (resp. g and h) to Y_i . Since the components of ∂f are non-separating and no two form a bounding pair, we can write

$$f = f_1 f_2 \cdots f_n \quad g = g_1 g_2 \cdots g_n, \quad h = h_1 h_2 \cdots h_n,$$

where each f_i is pseudo-Anosov on Y_i . Since $\text{MCG}(Y_i)$ is torsion-free and $f \in \mathcal{M}_G$, the centralizer of f_i in $\text{MCG}(Y_i)$ is cyclic. As f commutes with g and h , each f_i commutes with g_i and h_i . Thus, each g_i or h_i is contained in the centralizer of f_i , yielding g_i and h_i commute for all i . This shows g and h commute as required. $\square$

Finally, we can prove the characterization of Dehn twists and bounding pair maps.

Proposition 4.12. *Let $G < \mathcal{I}(\Sigma)$ be a finite-index subgroup, and let f lie in G . Then f is a power of a Dehn twist or of a bounding pair map if and only if f is in $\mathcal{M}_G$, and for all g in $\mathcal{M}_G$ such that $(\mathcal{P}_G)_g = (\mathcal{P}_G)_f$ we have that there exist integers $i, j \neq 0$ such that $f^i = g^j$.*

Proof. The forward direction is Lemma 4.8 and the definition of $(\mathcal{P}_G)_f$.

For the other direction, we prove the contrapositive. Assume that f is not a power of a Dehn twist or bounding pair map. Our goal is to find an element $g \in \mathcal{M}_G$ with $\text{supp}(g) = \text{supp}(f)$ such that no powers of f and g are equal, but $(\mathcal{P}_G)_f = (\mathcal{P}_G)_g$.

By Lemma 4.9, there exists a domain Y in $\Sigma \setminus \partial f$ on which $f|_Y$ is a pseudo-Anosov. Since Y supports a pseudo-Anosov, we may choose an element $h \in \text{MCG}(\Sigma)$ such that h preserves the components of $\Sigma \setminus \partial f$, is identity outside of Y , and the restriction to Y of f and $g = h f h^{-1}$ are two independent pseudo-Anosovs. Since $\mathcal{I}(\Sigma)$ is normal in $\text{MCG}(\Sigma)$, $g \in \mathcal{I}(\Sigma)$, and by construction, $\text{supp}(g) = \text{supp}(f)$. Thus $(\mathcal{P}_G)_g = (\mathcal{P}_G)_f$, but f and g do not have a common power. $\square$

4.4. Abstract commensurators of the Torelli group. We can now finally prove Theorem 1.4. For a bounding pair $\gamma = \{\gamma_1, \gamma_2\}$ we use the shorthand T_γ for the bounding pair map $T_{\gamma_1} T_{\gamma_2}^{-1}$.

Proof of Theorem 1.4. Let $[\psi]$ be an element of $\text{Comm } \mathcal{I}(\Sigma)$ representing the isomorphism of finite index subgroups

$$\psi: G_1 \longrightarrow G_2.$$

Let γ be a separating curve or a bounding pair and choose n in $\mathbb{N}$ so that T_γ^n is in G_1 . By Proposition 4.12, T_γ^n is in $\mathcal{M}_{G_1}$ and for all g in $\mathcal{M}_{G_1}$ such that $(\mathcal{P}_{G_1})_g = (\mathcal{P}_{G_1})_{T_\gamma^n}$, there exist integers i, j so that $(T_\gamma^n)^i = g^j$. Since these conditions

are preserved by isomorphism, we have that $\psi(T_\gamma^n)$ lies in $\mathcal{M}_{G_2}$, Proposition 4.12 implies there exists a separating curve or bounding pair δ and a nonzero integer m such that $\psi(T_\gamma^n) = T_\delta^m$.

At this point, and again with respect to the above notation, we obtain that ψ induces a map

$$\begin{aligned}\psi_*: \mathcal{J}(\Sigma) &\longrightarrow \mathcal{J}(\Sigma), \\ \gamma &\longmapsto \delta.\end{aligned}$$

We observe that ψ_* is a simplicial map, since powers of Dehn twists and bounding pair maps commute if and only if the underlying curves are disjoint. Moreover, the map is also bijective, with inverse the simplicial map associated to the inverse of ψ^{-1} .

By Theorem 4.2, there exists an $f \in \text{MCG}(\Sigma)$ such that $\psi_*(\gamma) = f(\gamma)$ for every separating curve or bounding pair γ . Now, for any g in G_1 we have

$$\psi(gT_\gamma^n g^{-1}) = \psi(g)\psi(T_\gamma^n)\psi(g^{-1}) = \psi(g)T_{f(\gamma)}^n\psi(g^{-1}) = T_{\psi(g)f(\gamma)}^n,$$

and therefore

$$T_{\psi(g)f(\gamma)}^n = \psi(gT_\gamma^n g^{-1}) = \psi(T_{g(\gamma)}^n) = T_{fg(\gamma)}^n.$$

Therefore $\psi(g)f(\gamma) = fg(\gamma)$. By use of the Alexander method [13] we conclude that $\psi(g) = fgf^{-1}$. This shows that every abstract commensurator of $\mathcal{J}(\Sigma)$ is defined by conjugation by a mapping class of Σ , and in particular, so is every automorphism of $\mathcal{J}(\Sigma)$.

On the other hand, suppose there exists an f in $\text{MCG}(\Sigma)$ and a finite-index subgroup $H < \mathcal{J}(\Sigma)$ such that conjugation by f induces the identity map on H . For any separating curve or bounding pair γ , there exists some $m \geq 1$ such that T_γ^m lies in H . Thus

$$T_\gamma^m = fT_\gamma^m f^{-1} = T_{f\gamma}^m.$$

By [2, Lemma 2.5], $f\gamma = \gamma$, and thus f is the identity by Theorem 4.2. This completes the proof. $\square$

References

- [1] J. Aramayona, P. Patel, and N. G. Vlamis, The first integral cohomology of pure mapping class groups. Preprint, 2017. [arXiv:1711.03132](https://arxiv.org/abs/1711.03132) [math.GT]
- [2] J. Bavard, S. Dowdall, and K. Rafi, Isomorphisms between big mapping class groups. To appear in *Internat. Math. Res. Notices*. Preprint, 2017. [arXiv:1708.08383](https://arxiv.org/abs/1708.08383) [math.GR]
- [3] M. Bestvina, K.-U. Bux, and D. Margalit, The dimension of the Torelli group. *J. Amer. Math. Soc.* **23** (2010), no. 1, 61–105. [Zbl 1233.20033 MR 2552249](https://doi.org/10.2307/2552249)

1398J. Aramayona, T. Ghaswala, A. E. Kent, A. McLeay, J. Tao, and R. R. Winarski

- [4] J. S. Birman, On Siegel's modular group. *Math. Ann.* **191** (1971), 59–68. [Zbl 0208.10601](#) [MR 0280606](#)
- [5] T. E. Brendle and D. Margalit, Normal subgroups of mapping class groups and the metaconjecture of Ivanov. Preprint, 2017. [arXiv:1710.08929](#) [math.GT]
- [6] T. E. Brendle and D. Margalit, Commensurations of the Johnson kernel. *Geom. Topol.* **8** (2004), 1361–1384. [Zbl 1079.57017](#) [MR 2119299](#)
- [7] T. E. Brendle and D. Margalit, Addendum to: “Commensurations of the Johnson kernel.” *Geom. Topol.* **12** (2008), no. 1, 97–101. [Zbl 1128.57303](#) [MR 2377246](#)
- [8] H. Burkhardt, Grundzüge einer allgemeinen Systematik der hyperelliptischen Functionen I. Ordnung. Nach Vorlesungen von F. Klein. *Math. Ann.* **35** (1889), no. 1-2, 198–296. [JFM 1510603](#) [MR 1510603](#)
- [9] B. Farb and N. V. Ivanov, The Torelli geometry and its applications. research announcement. *Math. Res. Lett.* **12** (2005), no. 2-3, 293–301. [Zbl 1073.57012](#) [MR 2150885](#)
- [10] B. Farb and D. Margalit, *A primer on mapping class groups*. Princeton Mathematical Series, 49. Princeton University Press, Princeton, N.J., 2012. [Zbl 1245.57002](#) [MR 2850125](#)
- [11] M. Handel and W. P. Thurston, New proofs of some results of Nielsen. *Adv. in Math.* **56** (1985), no. 2, 173–191. [Zbl 0584.57007](#) [MR 0788938](#)
- [12] A. Hatcher and D. Margalit, Generating the Torelli group. *Enseign. Math.* (2) **58** (2012), no. 1-2, 165–188. [Zbl 1273.57011](#) [MR 2985015](#)
- [13] J. H. Hernández, I. Morales, and F. Valdez, The Alexander method for infinite-type surfaces. Preprint, 2017. [arXiv:1703.00407](#) [math.GT]
- [14] N. V. Ivanov, Automorphisms of complexes of curves and of Teichmüller spaces. *Internat. Math. Res. Notices* **1997**, no. 14, 651–666. [Zbl 0890.57018](#) [MR 1460387](#)
- [15] D. L. Johnson, Homeomorphisms of a surface which act trivially on homology. *Proc. Amer. Math. Soc.* **75** (1979), no. 1, 119–125. [Zbl 0407.57003](#) [MR 0529227](#)
- [16] Y. Kida, Automorphisms of the Torelli complex and the complex of separating curves. *J. Math. Soc. Japan* **63** (2011), no. 2, 363–417. [Zbl 1378.57027](#) [MR 2793105](#)
- [17] Y. Kida, The co-Hopfian property of the Johnson kernel and the Torelli group. *Osaka J. Math.* **50** (2013), no. 2, 309–337. [Zbl 1282.20033](#) [MR 3080802](#)
- [18] M. Korkmaz, Automorphisms of complexes of curves on punctured spheres and on punctured tori. *Topology Appl.* **95** (1999), no. 2, 85–111. [Zbl 0926.57012](#) [MR 1696431](#)
- [19] F. Luo, Automorphisms of the complex of curves. *Topology* **39** (2000), no. 2, 283–298. [Zbl 0951.32012](#) [MR 1722024](#)
- [20] D. McCullough and A. Miller, The genus 2 Torelli group is not finitely generated. *Topology Appl.* **22** (1986), no. 1, 43–49. [Zbl 0579.57007](#) [MR 0831180](#)
- [21] A. McLeay, Geometric normal subgroups in mapping class groups of punctured surfaces. Preprint, 2018. [arXiv:1810.00742](#) [math.GT]
- [22] G. Mess, The Torelli groups for genus 2 and 3 surfaces. *Topology* **31** (1992), no. 4, 775–790. [Zbl 0772.57025](#) [MR 1191379](#)

- [23] P. Patel and N. G. Vlamis, Algebraic and topological properties of big mapping class groups. *Algebr. Geom. Topol.* **18** (2018), no. 7, 4109–4142. [Zbl 07006387](#) [MR 3892241](#)
- [24] Jerome Powell. Two theorems on the mapping class group of a surface. *Proc. Amer. Math. Soc.* **68** (1978), no. 3, 347–350. [Zbl 0391.57009](#) [MR 0494115](#)
- [25] A. Putman, Cutting and pasting in the Torelli group. *Geom. Topol.* **11** (2007), 829–865. [Zbl 1157.57010](#) [MR 2302503](#)
- [26] A. Putman, Small generating sets for the Torelli group. *Geom. Topol.* **16** (2012), no. 1, 111–125. [Zbl 1296.57008](#) [MR 2872579](#)
- [27] I. Richards, On the classification of noncompact surfaces. *Trans. Amer. Math. Soc.* **106** (1963), 259–269. [Zbl 0156.22203](#) [MR 0143186](#)
- [28] W. R. Vautaw, Abelian subgroups of the Torelli group. *Algebr. Geom. Topol.* **2** (2002), 157–170. [Zbl 0997.57035](#) [MR 1917048](#)

Received October 19, 2018

Javier Aramayona, Departamento de Matematicas,
Universidad Autónoma de Madrid & ICMAT, Ciudad Universitaria de Cantoblanco,
s/n, 28049 Madrid, Spain

e-mail: aramayona@gmail.com

Tyrone Ghaswala, Department of Mathematics, University of Manitoba, Machray Hall,
420, 186 Dysart Rd, Winnipeg, MB R3T 2M8, Canada

e-mail: ty.ghaswala@gmail.com

Autumn E. Kent, Department of Mathematics, University of Wisconsin, 480 Lincoln Dr,
Madison, WI 53706, USA

e-mail: kent@math.wisc.edu

Alan McLeay, Mathematics Research Unit, Université du Luxembourg, Esch-sur-Alzette,
Luxembourg, 2, avenue de l'Université, 4365 Esch-sur-Alzette, Luxembourg

e-mail: mcleay.math@gmail.com

Jing Tao, Department of Mathematics, University of Oklahoma, 73069, 601 Elm Ave,
Norman, OK 73019, USA

e-mail: jing@ou.edu

Rebecca R. Winarski, Department of Mathematics, University of Michigan,
530 Church St, Ann Arbor, MI 48109, USA

e-mail: rebecca.winarski@gmail.com